1   **Diagnosing and predicting mixed culture fermentations with unicellular and**

2   **guild-based metabolic models**

3   *Running Title:* MCFA Metabolic Models
4
5

6   *Matthew J. Scarborough,[1,2,3]* Joshua J. Hamilton[4] Elizabeth A. Erb,[5] Timothy J. Donohue[1,6]*

7   *and Daniel R. Noguera[1,2]*

8
9

1.   The Great Lakes Bioenergy Research Center, UW-Madison, Madison, WI

2.   Department of Civil and Environmental Engineering, UW-Madison, Madison, WI

3.   Department of Civil and Environmental Engineering, University of Vermont, Burlington,

VT

4.   Department of Biochemistry, UW-Madison, Madison, WI

5.   Department of Chemical and Biological Engineering, UW-Madison, Madison, WI

6.   Department of Bacteriology, UW-Madison, Madison, WI

*corresponding author e-mail: mscarbor@uvm.edu

ABSTRACT

Multi-species microbial communities determine the fate of materials in the environment and can be harnessed to produce beneficial products from renewable resources. In a recent example, fermentations by microbial communities have produced medium-chain fatty acids (MCFAs). Tools to predict, assess, and improve the performance of these communities, however, are limited. To provide such tools, we constructed two metabolic models of MCFA-producing microbial communities based on available genomic, transcriptomic and metabolomic data. The first model is a unicellular model (iFermCell215), while the second model (iFermGuilds789) separates fermentation activities into functional guilds. Ethanol and lactate are fermentation products known to serve as substrates for MCFA production, while acetate is another common co-metabolite during MCFA production. Simulations with iFermCell215 predict that low molar ratios of acetate-to-ethanol favor MCFA production, whereas the products of lactate and acetate co-utilization are less dependent on the acetate-to-lactate ratio. In simulations of an MCFA-producing community fed a complex organic mixture derived from lignocellulose, iFermGuilds789 predicted that lactate was an extracellular co-metabolite that served as a substrate for butyrate (C4) production. Extracellular hexanoic (C6) and octanoic acids (C8) were predicted by iFermGuilds789 to be from community members that directly metabolize sugars. Modeling results provide several hypotheses that can improve our understanding of microbial roles in a MCFA-producing microbiome and inform strategies to increase MCFA production. Further, these models represent novel tools for exploring the role of mixed microbial communities in carbon recycling in the environment, as well as on beneficial reuse of organic residues.

IMPORTANCE

Microbiomes are vital to human health, agriculture, and protecting the environment.

Predicting behavior of self-assembled or synthetic microbiomes, however, remains a challenge. In

this work, we used unicellular and guild-based metabolic models to investigate production of

medium-chain fatty acids by a mixed microbial community that is fed multiple organic substrates.

Modeling results provided insights into metabolic pathways of three medium-chain fatty acid-

producing guilds and identified potential strategies to increase production of medium-chain fatty

acids. This work demonstrates the role of metabolic models in augmenting multi-omic studies to

gain greater insights into microbiome behavior.

INTRODUCTION

Mixed microbial fermentations have benefited humanity throughout history.[1] Since the

mid-18[th] century, mixed culture fermentations have produced valuable chemicals and recovered

energy from wastes.[2] Many mixed microbial fermentations involve the synergistic activities of

several functional groups to convert organic substrates into valuable products. In anaerobic

digestion, fermentation products like acetic acid (C2) and H_2 are used to support microbial

production of methane, thus enabling the recovery of a large fraction of the chemical energy

originally present in the complex organic substrates. While anaerobic digestion technologies are

well established, other strategies to expand the range of potential products of mixed culture

fermentations are only beginning to emerge. One example of such a mixed culture fermentation is

the carboxylate platform, which produces medium-chain fatty acids (MCFAs) as a potentially

valuable set of bio-based products.[3]

MCFAs are attractive carboxylate platform products because of their many industrial uses.[4]

Octanoic acid (C8), an 8-carbon linear monocarboxylic acid, is a particularly valuable MCFA

because of its high market value and relative ease of recovery. Further, when C8 is reduced to its

corresponding alkane, it can substitute for octane in liquid transportation fuels. Recent research

has demonstrated the feasibility of using mixed culture fermentations to transform renewable

resources into MCFAs.[5-7] While C8 is produced by some self-assembled bioreactor microbial

communities, hexanoic acid (C6) is often the primary MCFA, along with two short-chain fatty

acids (SCFAs), acetic (C2) and butyric (C4) acids.[5-10]

The microbial metabolic networks involved in converting complex organic substrates to

MCFAs by mixed culture fermentations are not fully elucidated. Consequently, there is a lack of

knowledge on how to obtain a carboxylate platform community that produces C8 as the main

MCFA, or how to minimize the accumulation of SCFAs. Initial models of mixed culture

fermentations to produce products other than methane have focused on simulating metabolic

networks for production of C2, C4, propionic acid, ethanol, lactic acid and H_2 from glucose.[11]

Subsequent models took into account that multiple electron carriers (e.g., ferredoxin, NADH) play

different roles in these metabolic networks, enabling the simulation of energy-conserving electron-

bifurcating reactions during production of C4 by reverse β-oxidation.[12] While a recent model

considered the role of homoacetogens as a H_2-consuming member of mixed culture fermentation,

it was limited to producing C4 as the major product of reverse β-oxidation and did not include

MCFA production.[13] Additionally, recent metabolic models assessed the potential for producing

C2, C3, C4 and valerate (C5) from amino acids.[14] While all of these models have enabled

predictions of mixed-culture fermentations from a single organic substrate, they lack the ability to

simulate production of MCFAs from complex feedstocks.

In this study, we constructed a unicellular model (iFermCell215) and a guild-based

microbial community model (iFermGuilds789) that offer expanded substrates and metabolic

pathways compared to existing models. We used iFermCell215 and iFermGuilds789 to analyze

pathways that lead to MCFA production. Simulations were performed with single organic

substrates, including sugars, glycerol, lactate, and acetate, with ethanol and acetate or lactate and

acetate as co-substrates, and with complex organic mixtures. The iFermGuilds789 was also used

to evaluate predictions of observed bioreactor behavior and of the flow of carbon and electrons

through different guilds within the mixed microbial community. In total, results of individual

model simulations generated new hypotheses on how to increase MCFA production by mixed

microbial communities.

MATERIALS AND METHODS

Metabolic networks of mixed culture fermentation that contain reactions that occur during

anaerobic metabolism of a previously described lignocellulosic biorefinery residue[15] were

assembled. The first network, iFermCell215, contains 215 reactions and 154 metabolites, models

mixed culture fermentation as a single unit, and expands the metabolic pathways and substrate

range of previously published mixed culture fermentation networks.[11-13] iFermCell215 describes

metabolism of exogenous substrates including glucans, glucose, xylans, xylose, and glycerol,

along with lactate, ethanol, acetate, CO_2 and H_2, which are potential intermediates in this mixed

culture fermentation (**Fig. 1**).

The second network, iFermGuilds789, represents activities of different guilds (i.e., groups

of organisms with similar functions) within the microbial community. The iFermGuilds789

network describes the same reactions and metabolites as in the iFermCell215 model but these are

separated into six functional guilds with each guild representing a subset of compartmentalized

reactions (**Fig. 2**). iFermGuilds789 also includes additional transport reactions that simulate

metabolite exchange among guilds. Combining the six guilds, the iFermGuilds789 network

includes 789 reactions and 579 metabolites. The guilds in the iFermGuilds789 model represent

previously described organisms that perform fatty acyl chain elongation from sugars (SEOs), sugar

fermenting organisms that produce lactate, acetate, and ethanol as fermentation products (SFO),

hydrogen-producing sugar fermenters (HSFs), organisms that perform fatty acyl chain elongation

from lactate (LEOs) or ethanol (EEOs), and homoacetogenic organisms (HAOs). Examples of

organisms falling within each functional guild are described in **Supplementary Text 1**.

When constructing each metabolic network, reactions were added to capture the exchange

of substrates and products with the extracellular space.[16] In addition to intracellular (denoted as

"_c" in the metabolic models) and extracellular ("_e") compartments, a compartment for ion-

motive force generation was included (denoted as "_i" in the metabolic models). Where possible,

standard metabolite and reaction abbreviations from the Biochemical, Genetic, and Genomic

(BiGG) knowledgebase were used.[17] All reactions were balanced for mass and charge assuming

an intracellular pH of 7.0 with the dominant ionic form of all metabolites obtained from the BiGG

database.[17]

Since many members of this mixed microbial community are Firmicutes, simplified

biomass equations for both models were created based on the biomass composition of the well-

studied gram-positive bacterium *Bacillus subtilis*.[18-20] We simplified all of the biosynthetic

pathways for production of biomass components to capture 16 precursor compounds produced by

the models. The precursor compounds included α-ketoglutarate, oxaloacetate, glucose-6-

phosphate, glucose-3-phosphate, 3-phosphoglycerate, pyruvate, phosphoenolpyruvate, acetyl-

CoA, erythrose-4-phosphate, ribulose-5-phosphate, ATP, NADPH, NAD+, ammonium, sulfate,

and H_2O. Additional information and calculations for the biomass equation are provided in

Supplementary Text 2, Supplementary Data File 1, Figures S1-S6, and Table S1. Both

metabolic models (iFermCell215 and iFermGuilds789) and instructions for their use are available

at https://github.com/mscarbor/Mixed-Culture-Fermentation-Models.

The production capabilities of both metabolic networks were simulated using constraint-

based methods.[21] These methods predict the steady-state flux distribution through a metabolic

network that is subject to physiochemical constraints. In this work, we used flux sampling[22] and

parsimonious flux balance analysis (pFBA).[23] Flux sampling was performed using artificial

centering hit and run (ACHR)[22] using 101 simulations and randomly generated seeds. A solution

with pFBA provides a flux distribution by minimizing the sum of all fluxes predicted by the

model.[23] When modeling bioreactors with the presence of solute gradients across the cell

membrane, we considered the ATP required for transport as described in **Supplementary Text 3.**

The objective function for all modeling scenarios was to maximize biomass production. When

simulating the iFermGuilds789 network for a steady-state condition, we required all individual

guilds to achieve the same growth rate (hr^{-1}). All simulations were performed in Python 3.7 using

the cobrapy package (v 0.18.1).[24]

Throughout this work, we present results in terms of electron equivalents (eeq), which

report the number of electrons potentially transferred to an electron acceptor if the compound is

completely oxidized (using factors provided in **Table S2** to convert from molar concentrations to

eeq concentrations). For each scenario, we calculated the overall reaction thermodynamics using

the predicted stoichiometry of the model and the standard free energy of formation (ΔG_f^0) of

metabolites (**Table S2**) obtained from KBase.[25] Unless otherwise noted, we ensured that at least

50 kJ was available per ATP produced under standard conditions at neutral pH.[26, 27]

RESULTS

Simulation of mixed culture fermentations with iFermCell215

By constraining the iFermCell215 model to produce defined products, we initially tested

its predictive ability with well-established fermentation conditions (See **Supplementary Text 4**

and **Table S3**). We then investigated predictions derived from an iFermCell215 model under

conditions where single substrates were consumed within this unit. The iFermCell215 model

depicts a simplified representation of fermentation in a mixed microbial community where

interspecies transfer of metabolites is assumed to have no energetic penalty, and is similar to

previously published models of mixed culture fermentations.[11-13] For this analysis, we required

that all possible solutions released $\geq$ 50 kJ per mol ATP produced.[26]

Using flux sampling, iFermCell215 provided 101 solutions that maximized biomass

production depending on the substrates provided to the simulation (**Fig. 3**). For instance, when we

modeled the utilization of glucose or xylose as the sole substrate, iFermCell215 predicted multiple

solutions that allowed for maximum biomass production (40% of eeq are incorporated into

biomass) with H_2, C2, C4, C6, and C8 as predicted products (**Fig. 3A**). None of the solutions in

this flux sampling analysis predicted ethanol or lactate as significant products. The predicted

biomass yield (eeq incorporated into biomass per eeq consumed) decreased with glycerol (37%),

lactate (20%), and ethanol (5.9%) as sole substrates. Consumption of ethanol as a sole substrate

was predicted to produce C8 and H_2 as the most energetically favorable solution (**Fig. 3D**), whereas

consumption of lactate as a sole substrate produced optimal solutions through heterofermentation

(**Fig. 3C**) and consumption of H_2 and CO_2 was predicted to produce C2 for optimal biomass

growth, which is consistent with the activity of homoacetogens (**Fig 3E**). These results suggest

that sugars, glycerol, and lactate can produce significant amounts of multiple fermentation

products, including C2, C4, C6, and C8, while supporting optimal biomass production, and that

production of ethanol or lactate, the two common fermentation products described as intermediates

in MCFA production,[28] are not as energetically favorable as producing other fermentation

products. It further suggests that consumption of ethanol as a sole substrate should favor

production of C8, and therefore, that ethanol may be a better intermediate product for optimizing

C8 production than lactate. However, since the fermentation of complex organic substrates by

mixed communities often results in accumulation of multiple fermentation products, we also

simulated conditions in which ethanol and acetate or lactate and acetate were co-substrates for

MCFA production.

187 Simulation of ethanol and acetate utilization with iFermCell215

Ethanol is a substrate that has often been used for chain elongation,[29] and early work on

MCFA production was performed using the ethanol- and acetate-consuming *Clositridium*

kluyveri.[30, 31] During initial work with *C. kluyveri*, it was proposed that C4 could be produced from

two moles of ethanol, without using acetate as an electron acceptor.[31] Recent work has confirmed

that *C. kluyveri* produces MCFAs from ethanol as a sole substrate in batch experiments,[32] in

agreement with the iFermCell215 prediction (**Fig. 3E**). Production of MCFAs from C2 as a sole

substrate has not been observed and is not predicted by iFermCell215. Past studies of ethanol

elongation have investigated optimal ratios of ethanol-to-acetate for MCFA production.[29, 33, 34]

Here, we use iFermCell215 to assess impacts of different molar ratios of acetate-to-ethanol

(C2:EtOH) on predicted product formation.

When supplied with both ethanol and C2, the iFermCell215 predicted maximum growth

rate of 0.11 hr^{-1} is more than double the predicted growth rate with ethanol as the sole substrate

(**Fig. 4A**), suggesting an ecological advantage for microorganisms that can co-utilize ethanol and

acetate. This growth rate can be achieved at C2:EtOH ratios from 0.56 to 1.28 mol:mol. At ratios

of 0.56 mol:mol and lower, C8 and H_2 are the only major predicted products (**Fig. 4**, left panels).

At ratios between 0.56 and 1.20, a mixture of C8, C6, and C4 are predicted by iFermCell215, and

at ratios of 1.28 and above, C4 is predicted as the sole product. These model predictions are

consistent with previous experimental work that shows that low ratios of C2:EtOH improve

specificity of C8 production.[33] The overall production rate of MCFAs, however, has been shown

to improve at higher ratios of C2:EtOH.[33] The highest overall MCFA production rate was achieved

by Spiritio et. al at a ratio of 0.53 mol:mol,[33] which is very close to the model predicted ratio of

0.56 that optimizes biomass growth and C8 production. The production of H_2 at low ratios of

C2:EtOH raises the question as to the impact of H_2 partial pressure on the thermodynamic

feasibility of the model results. We used iFermCell215 to calculate the maximum H_2 partial

pressure for all 101 scenarios and found that the lowest allowable H_2 partial pressure for growth

is greater than 10^6 atm (**Fig S7**), indicating that the proposed transformations are

thermodynamically favorable under all bioreactor H_2 partial pressure conditions.

In total, iFermCell215 predicts that the C2:EtOH ratio could either be used to improve

production of C8 as a desired product, or could be an indication that the environmental conditions

in a bioreactor would lead to a mixture of fermentation products. At low C2:EtOH ratios (<0.56),

iFermCell215 predicts that C8 should be the abundant product. At medium C2:EtOH ratios (0.56-

1.20) iFermCell215 predicts a mixture of C4, C6, and C8 would be produced, while at high ratios

(>1.20) of these substrates, iFermCell215 predicts C4 would be the predominant product.

Therefore, in a microbial community fermenting a mixture of substrates, iFermCell215 predicts

that different ratios of ethanol and C2 will impact the composition and abundance of products.

Additionally, iFermCell215 predicts that, if feeding a reactor with ethanol and C2, attention should

be paid to creating a feeding regime that optimizes production of the target product.

Simulation of lactate and acetate utilization with iFermCell215

In addition to ethanol, lactate is often used as a substrate for microbial chain elongation.

The well-studied *Megasphaera elsdenii* has been shown to produce MCFAs from lactate,[35] and

recent chain elongation studies with microbial communities have explored lactate as both a

substrate[9, 36] and a proposed key intermediate that supports chain elongation from carbohydrate

fermentation.[15, 37] While ethanol and lactate are both potential fermentation products and substrates

for chain elongation, iFermCell215 predicts that the pathways for utilization of these substrates

have key differences that impact the terminal fermentation products, as shown in the flux sampling

analyses (**Fig. 3D-E**), and may impact the products when C2 is a co-substrate. Therefore, similar

to the analysis shown above for C2:EtOH ratios, we used iFermCell215 to assess impacts of

different C2-to-lactate ratios.

In the case of lactate, the predicted maximum growth rate does not change when C2 is a

co-substrate (**Fig. 4E**). This suggests that, unlike ethanol and C2 co-utilization, there is no apparent

energetic benefit (reflected in growth rate) for a microorganism to co-utilize lactate and C2. Low

ratios of C2:lactate are predicted by iFermCell215 to produce a variety of end products, including

C8, C6, C4, and H_2 (**Fig. 4**, right panels). As the C2:lactate ratio increases, the product distribution

is predicted by iFermCell215 to change, with a decrease in MCFA and H_2 production accompanied

by an increase in C4 production, and at ratios of 0.96 and above, C4 is predicted as the sole product.

Like ethanol and C2 co-utilization, lactate and C2 co-utilization is not predicted by iFermCell215

to be impacted by bioreactor H_2 partial pressure (**Fig S7**). These results suggest that when lactate

is a key intermediate in an MCFA-producing community, controlling the relative amounts of C2

and lactate may not be an effective strategy for increasing C6 and C8 production. However, like

with ethanol, an excess amount of acetate is predicted by iFermCell215 to increase C4 production.

The predicted differences in product formation when acetate is co-metabolized with

ethanol or lactate suggests important differences in the metabolism of these substrates. Therefore,

to understand these differences, we also used iFermCell215 to assess the predicted metabolic

pathways of ethanol and lactate utilization.

Predicted metabolic pathways for ethanol and lactate utilization

The flux sampling analysis obtained with iFermCell215 consuming ethanol as a sole

substrate converged to a solution in which C8 was the only significant fermentation product (**Fig.**

3E). The flux of electron equivalents in this simulation predicted that reactions catalyzed by

ferredoxin hydrogenase (Hyd1), proton-translocating ferredoxin:NAD+ oxidoreductase (aka,

Rhodobacter nitrogen fixation complex; RNF), a coenzyme A transferase (CoAT), and acetate

kinase (AckR) were key enzymes used in the network (**Fig 5A**). The genome of the well-studied

ethanol chain-elongating organism, *C. kluyveri,* is predicted to contain genes that encode the

proteins for catalyzing all of these reactions.[38] Further, Hyd1 is the only hydrogen-producing gene

annotated in the genome of *C. kluyveri*.

The flux through the above enzymes predicts extensive use of energy conserving

mechanisms that rely on the generation of an ion-motive force (IMF). When ethanol is the sole

substrate (**Fig. 5A**), iFermCell215 predicted that the IMF is generated by the ATP synthase while

hydrolyzing ATP. The model predicts that the RNF enzyme uses the IMF to generate reduced

ferredoxin and that ATP is provided by acetate kinase. The reverse β–oxidation cycle uses an

electron bifurcating acyl-CoA dehydrogenase that also generates reduced ferredoxin, and redox

balance is maintained by Hyd1 that uses the reduced ferredoxin to generate H_2 **(Fig. 5A)**. When

C2 is available in addition to ethanol, iFermCell215 predicts that no net H_2 production is required

for redox balance **(Fig. 5B)**. Instead, iFermCell215 predicts that the activities of several enzymes

are reversed, with the RNF complex generating IMF, ATP synthase producing ATP, and acetate

kinase consuming ATP to generate acetyl-CoA. Thus, the same set of reactions is predicted to be

important by iFermCell215 when using ethanol as a sole substrate or ethanol and C2 as co-

substrates, but the predicted direction of key reactions change (**Fig 5A-B**). These predictions

suggest that organisms that elongate ethanol may readily transition between utilizing ethanol as a

sole substrate and co-utilizing ethanol and C2 since the same enzymes are predicted to be used,

and the co-utilization of C2 and ethanol improves growth rates (**Fig 4A**).

Unlike ethanol utilization, lactate conversion to acetyl-CoA is predicted by iFermCell215

to produce CO_2, formate, and reduced ferredoxin (**Fig. 5C-D**). The Wood-Ljungdahl pathway is

also predicted by iFermCell215 to use this CO_2, formate and reduced ferredoxin to produce C2

(**Fig. 5C-D**). The predicted ability to generate reduced ferredoxin during lactate consumption

(compared to only NADH from ethanol consumption) enables iFermCell215 to utilize a much

wider range of pathways for generating energy and maintaining redox balance. The reverse

β−oxidation cycle also uses an electron bifurcating acyl-CoA dehydrogenase under these

conditions, generating reduced ferredoxin, with RNF also predicted to be active in redox

balancing. These predictions of iFermCell215 explain the wide range of products that are possible

from utilization of lactate as sole substrate and co-utilization of C2 and lactate. Further, this

suggests that limiting the available metabolic pathways can impact the predicted products from

lactate. Indeed, removing specific reactions in iFermCell215 impacts the predicted product

spectrum (**Supplementary Text 5**, **Fig S7**). This demonstrates that strategies to improve MCFA

production may depend on the available set of microbial transformations and may differ between

bioreactor microbial communities.

Simulating community behavior with iFermCell215 and iFermGuilds789

The iFermCell215 model predicted that multiple substrates could lead to C8 production

(**Fig. 3**), but that biomass could be maximized by co-production of other fermentation products.

We wanted to assess if this single-unit model could simulate the observed product yields of a

mixed microbial community fed lignocellulosic biorefinery residues, a complex organic

substrate.[15] The previously described bioreactor was fed lignocellulosic biorefinery residues for

96 days, at which time its microbial community was evaluated using metagenomic and

metatranscriptomic analyses along with end-product sampling. To represent organic substrate

metabolism, the substrate uptake fluxes were set according to the observed consumption of

compounds in the feedstock (**Table S6**). We constrained iFermCell215 by setting the growth rate

of the model to 0.0069 hr^{-1}, corresponding to the bioreactor retention time of 6 d and we again

constrained the model only by requiring a minimum of 50 kJ of free energy release per mol of

ATP produced. Flux sampling results from this scenario predicted a range of potential products

(**Fig. 6A**), including ethanol, C2, H_2, even chain carboxylic acids, and odd-chain carboxylic acids.

Overall, the amount of MCFAs predicted by iFermCell215 exceeded observed values, exposing a

limitation of the single unit model.

In an attempt to better represent the compartmentalized nature of activities within mixed

microbial communities we developed the functional guild model iFermGuilds789. In

iFermGuilds789, the metabolic activities of a mixed microbial community are partitioned among

a number of functional guilds. This biological division of labor negates the free exchange of

reducing equivalents (e.g., NADH, ferredoxin) and of many pathway intermediates (e.g., acetyl-

CoA) between community members, ultimately constraining how individual organisms maintain

redox balance, which can potentially impact ATP and product yields. For instance, the single unit

model (iFermCell215) cannot predict the production of lactate via a NADH-dependent

dehydrogenase and re-consumption of lactate via an electron-confurcating dehydrogenase, activities that may happen in a mixed microbial community that includes lactate producers and lactate consumers. The iFermGuilds789 model contains fermentation pathways in six guilds (**Fig. 2**), with each guild including predictions on transport reactions to simulate uptake and release of intermediate metabolites by the most abundant members of the community as predicted from previous metagenomics analyses.[15]

To simulate the activities in the MCFA-producing bioreactor we used a version of iFermGuilds789 with only four guilds, which represented the most abundant community members in the bioreactor as determined by metagenome sequencing.[15] These guilds included the sugar elongating Cand. *Weimeria bifida* (SEOs), lactate-elongating Cand. *Pseudoramibacter fermentans* (LEOs),[39] five sugar fermenting *Lactobacillus* species (SFOs), and three hydrogenic sugar fermenting *Coriobacteriaceae* species (HSFs).[15] We constrained the abundance of each guild in iFermGuilds789 according to the relative abundance of DNA for each guild in the bioreactor. Guilds were assumed to have all pathways associated with their guild (**Fig. 2**), and we required each guild to have the same growth rate of 0.0069 hr[-1]. The results of this iFermGuilds789 simulation (**Fig. 6B**) predicted a narrower range of products than iFermCell215 (**Fig. 6A**), with even- and odd-chain fatty acid products predicted. With these parameters, the iFermGuilds789 model overpredicted the production of C2 and C8 while underpredicting C4 and C6 production.

In an additional iFermGuilds789 simulation, we restricted pathways in guild members based on previous metatranscriptomic analyses.[15] SEOs were constrained to use glucose, xylose and glycerol; SFOs were restricted to consuming glucose, xylose, xylans, glucans, and glycerol; HSFs were constrained to only consume glucans, glucose and glycerol; and LEOs were restricted to consuming only lactate and glycerol (**Table S7**). Further, SEOs and LEOs were constrained to

not produce odd-chain volatile fatty acids (propionate, pentanoate, and heptanoate). LEOs were

further constrained to not use acetate kinase, which is absent from the genome of Cand.

Pseudoramibacter fermentans.[39] This iFermGuilds789 simulation predicted ranges of ethanol,

lactate, C2, and C6 that contained the experimentally observed values (**Fig 6C**). With these

parameters, C8 and H_2 production were overpredicted by iFermGuilds789, while C4 production

was underpredicted.

One potential explanation for the differences in the various iFermGuilds789 predictions

may be the specificity of chain elongating enzymes for producing products of defined chain

lengths. This could include any of the enzymes catalyzing the four steps of reverse β-oxidation

itself or the terminal enzymes that cleave the final products of reverse β-oxidation from CoA. We

tested this by constraining the flux through CoAT and acyl-CoA hydrolase reactions for both SEOs

and LEOs according to carbon-chain length. When constraining both CoAT and acyl-CoA

hydrolase in both SEOs and LEOs to 0.02 mmol gDCW^{-1} hr^{-1} for C8 and 1 mmol gDCW^{-1} hr^{-1} for

C6, iFermGuilds789 modeling predictions aligned much better with community performance (**Fig.**

6D). This suggests that reverse β-oxidation and/or terminal enzymes in this pathway have higher

affinity for C4 and C6 than for C8 in this specific microbial community.

Taken together, initial predictions from iFermCell215 (**Fig. 6A**) and iFermGuilds789 (**Fig.**

6B) did not align with observed bioreactor performance. Predictions of iFermGuilds789 improved

by incorporating knowledge of specific reactions within each guild (**Fig. 6C**). Further, the best

alignment between observations and model predictions resulted from constraining the fluxes

related to individual products with iFermGuilds789 (**Fig. 6D**). This suggests that accurate

predictions of MCFA production may require incorporation of enzyme affinity and additional

characterization of enzymes from chain-elongating species to improve modeling predictions.

Assessing interspecies metabolite flux using iFermGuilds789

Our previous analyses predicted that microbes within the MCFA-producing microbial community perform different functions, compete for resources, and participate in interspecies transfer of metabolites that allows for conversion of lignocellulosic biorefinery residues to SCFAs and MCFAs.[15] We used iFermGuilds789 to predict the function of each guild in the transformation of substrates as well as the production and transfer of intermediate metabolites between microbes. In this case, unlike previous simulations with iFermGuilds789, we did not constrain the growth rate (**Fig. 6B-D**) or fluxes for MCFA production (**Fig. 6D**). Instead, we constrained iFermGuilds789 to accumulate extracellular products at the rates observed in the MCFA-producing bioreactor (**Table S6**), constrained the reactions within each guild as described above, and predicted fluxes using pFBA, which calculates a single flux distribution when the model is constrained to the observed production rates. Under these conditions, iFermGuilds789 predicted a growth rate of 0.0051 hr^{-1}, which is 74% of the expected growth rate (0.0069 hr^{-1}). This growth rate corresponds to a retention time of 8 days (compared to the actual retention time of 6 days) and these differences may reflect difficulty in estimating the amount of metabolically active cells in the reactor or absence of unknown metabolic pathways that may increase the predicted growth rate.

By constraining the substrates and products according to observed community behavior, iFermGuilds789 predicted that SFOs and HSFs consumed xylans and glucans, while SEOs consumed xylose, glucose, and 99% of the glycerol (**Fig 7**). SFOs were predicted to produce ethanol, lactate and C2, while HSFs were predicted to produce a small amount of the lactate and C2. SEOs are also predicted to produce lactate and C2. In total, 22% of the eeq in the feedstock are predicted by this instance of iFermGuilds789 to be converted to lactate and 27% are predicted

to be converted to C2. All of the lactate and 52% of the C2 is predicted to be consumed by LEOs.

LEOs are predicted to produce all of the C4. SEOs are predicted to produce all of the C6 and C8

along with all of the H_2. SEOs are further predicted to have a flux for C6 production of 0.46 mmol

gDCW^{-1} hr^{-1} and C8 production of 0.033 mmol gDCW^{-1} hr^{-1} (**Supplementary Data File 5**), both

of which are lower than the maximum flux constraints applied previously (**Fig. 6D).**

Predicted routes of sugar and lactate elongation

The above prediction that essentially all MCFAs were produced by SEOs consuming

xylose, glycerol, and glucose, while LEOs produced C4 from lactate and C2 were different than

the predictions made earlier based on transcriptomic analyses alone.[15] To gain further insights into

SEO and LEO metabolism, we investigated the specific reactions predicted for both guilds by

iFermGuilds789. SEOs are predicted to produce pyruvate via glycolysis. Some pyruvate is

predicted to be converted to lactate, while most is converted to acetyl-CoA via pyruvate flavodoxin

oxidoreductase (PFOR, **Fig 8A**). iFermGuilds789 also predicted that acetyl-CoA is derived from

the phosphoketolase pathway (PKETX, **Fig 8A**). Acetyl-CoA is predicted to be directed to reverse

β-oxidation, but a portion is also predicted to be hydrolyzed by an acyl-CoA hydrolase. Acyl-CoA

hydrolase enzymes are also predicted as terminal enzymes of reverse β-oxidation rather than

energy-conserving CoA transferase enzymes (CoAT). SEOs are further predicted to use Hyd1 for

hydrogen production, to generate IMF via RNF1, and produce ATP via ATP synthase (ATPS4r)

(**Fig. 8A**).

LEOs are predicted to convert lactate to pyruvate via an electron-confurcating lactate

dehydrogenase (**Fig. 8B**, ECLDH), and pyruvate is predicted to form acetyl-CoA via PFOR. All

of the acetyl-CoA is predicted to undergo reverse β-oxidation and C2 is incorporated by a CoAT

that transfers the CoA from butyryl-CoA to acetyl-CoA (**Fig. 8B**). LEOs are predicted to consume

equimolar concentrations of C2 and lactate, which was previously predicted to favor C4 production

(**Fig. 4**). Further, LEOs are not predicted to produce H_2. iFermGuilds789 predicted that H_2

produced by Hyd1 is re-consumed via a proton-translocating energy conserving hydrogenase that

produces reduced ferredoxin (ECH, **Fig. 8B**). This ferredoxin augments that produced by PFOR

and is predicted to be used for lactate consumption by ECLDH and the RNF complex. The use of

an energy conserving hydrogenase for ferredoxin production from H_2 has been demonstrated

previously.[40]

While both SEOs and LEOs are predicted by iFermGuilds789 to use reverse β-oxidation

to elongate acetyl-CoA, the energy-conserving and redox-balancing reactions needed to support

this process are predicted to differ. While SEOs are predicted to use an acyl-CoA hydrolase as the

terminal enzyme of reverse β-oxidation, LEOs are predicted to use a CoAT which conserves

energy by activating C2 to acetyl-CoA, which would otherwise require ATP. In the previous

metatranscriptomic analyses of these community members,[15, 39] we found that both SEOs and

LEOs expressed genes for both acyl-CoA hydrolases and CoAT. The specificity of these predicted

chain elongating enzymes for different substrates, however, is not known. The iFermGuilds789

model predictions reinforce the potential importance of terminal enzymes in reverse β-oxidation

in determining the final product length; specifically, it predicts that all C4 is produced using CoAT

and all MCFAs are using acyl-CoA hydrolase.

Furthermore, SEOs are predicted by iFermGuilds789 to maintain redox balance by

producing H_2 with Hyd1, while LEOs are predicted to recycle all H_2 internally using both Hyd1

and ECH. Previous metatranscriptomic analyses showed that both LEOs and SEOs expressed

genes for both Hyd1 and ECH.[15] This agrees with iFermGuilds789 predictions for LEOs, but not

with SEOs that are predicted to only use Hyd1. When constraining iFermGuilds789 to require

SEOs to use ECH, there are no changes in interspecies metabolite transfer (**Supplementary Data**

File 5), but SEOs are predicted to generate additional ATP via ATP synthase while reducing flux

through other energy-conserving pathways to achieve the same growth rate (**Supplementary Data**

File 5). These findings suggest that SEOs and LEOs have multiple methods for conserving energy

and maintaining redox balance and this metabolic flexibility can maintain the same overall

transformation of complex substrates to products. This metabolic flexibility may reflect ecological

advantages for fermenting organisms, but are also challenges for the accurate predictions of MCFA

production, which may strongly depend on the specific organisms present in the microbial

community.

DISCUSSION

Under anaerobic conditions, a diverse set of mixed microbial communities catalyze the

transformation of organic substrates to a variety of products. For over a century, anaerobic

microbial communities have been harnessed to recover energy, in the form of methane-rich biogas,

from complex organic matter in anaerobic digestion.[2] Some fermentation metabolisms are well

understood, and are used by genetically tractable pure cultures that can be analyzed, modeled and

improved for use in industrial applications.[41, 42] In contrast, the activities of mixed microbial

communities have been harder to analyze, predict or redirect to improve substrate utilization or

product yields. The ability to harness the metabolic activity of mixed microbial communities to

produce alternative fermentation products is limited since metabolic and kinetic models that

accurately predict the distribution of fermentation products are only starting to emerge.[11-13] To

address this knowledge gap, we constructed metabolic models of mixed culture fermentations with

the objective of gaining greater insights into how to increase MCFA production by mixed

microbial communities. We focused modeling the production of C6 and C8, linear medium-length

MCFAs of industrial value that can be produced from a variety of organic residues.[3] To build and

test these models we took advantage of prior metagenomic, metatranscripotomic, and metabolomic

analyses of a mixed community that produces C6 and C8 fatty acids from a lignocellulosic refinery

residue.[15, 39]

The predictions of our models generated several hypotheses that should be explored with

further analyses. First, the single unit iFermCell215 predicted production of MCFAs from ethanol

as a sole source of electrons and carbon (**Fig. 4A-D**), and this is supported by recent experiments

with mixed cultures.[33] More work is needed to determine bioreactor conditions that promote

elongation of ethanol as a sole substrate or with low C2:EtOH ratios. The iFermCell215

simulations predict a threshold of 0.56 mol C2 per mol ethanol below which C8 production benefits

from C2 co-utilization, and above which C8 production starts to decrease and to be replaced by

production of shorter acids. This shift in product profile was predicted by iFermCell215 to be

associated with reversal in the direction of enzymatic activities related to energy conserving

metabolism, an observation that still lacks experimental verification.

Second, even though lactate is a known substrate for chain elongation,[9] iFermGuilds789

simulation results suggest that, in this mixed community, lactate may be converted to C4 instead

of C6 or C8 **(Fig. 7)**. Although at lower C2 to lactate ratios iFermCells215 simulations predicted

that this ratio was not as influential on determining the product distribution as the C2:EtOH ratio,

it also predicted that excess C2 could result in C4 production instead of C6 or C8 from both

intermediate metabolites (**Fig. 4**). The iFermGuilds789 simulation predicted co-utilization of C2

and lactate at a molar ratio of 1.0 (**Fig. 8B**), which according to the iFermCell215 simulations (**Fig.**

4) this is already an excess C2 condition predicted to yield C4 as the main fermentation product.

These results have the important implication that efforts directed to construct synthetic

communities or control self-assembled communities for production of MCFA should focus on

minimizing C2 production and accumulation.

Third, iFermCell215 modeling results suggest that controlling product formation from

lactate may not be possible by controlling C2-to-lactate ratios (**Fig. 4E-G**), whereas C2:EtOH

ratios suggests very specific targets to maximize C8 production (**Fig. 4A-D**). This suggests that

for MCFA production from complex organic substrates, in addition to targeting the reduction of

C2 production, efforts should also be directed to creating conditions in which ethanol instead of

lactate is the key intermediate metabolite for MCFA production. However, this may not be

achievable in self-assembled communities, and since simulations predict that the products of

lactate fermentation are dependent on specific metabolic features of LEOs (**Fig S8**), it is important

to continue the discovery of the natural diversity of lactate fermentation metabolisms, which

eventually could help identify preferred LEOs for the development of novel strategies to promote

MCFA production when lactate is the key intermediate metabolite.

Fourth, iFermGuilds789 predicts that enzyme affinity plays a major role in determining the

final products of reverse β-oxidation and mixed culture fermentation (**Fig 6D**). Additional work is

needed to assess the affinity of reverse β-oxidation, acyl-CoA hydrolase, and CoA transferase

enzymes for different chain lengths to identify ideal microorganisms or engineer microorganisms

to improve MCFA production. Past studies have identified and engineered acyl-ACP thioesterase

enzymes (the terminal enzyme of fatty acid synthesis via the fatty acid biosynthesis pathway) with

high affinity for C8.[43-45] Similar studies should be performed with acyl-CoA hydrolase and acyl-

CoA transferase enzymes to characterize and improve the terminal enzymes of reverse β-

oxidation.

Since many microbes in natural and self-assembled fermentation communities remain

uncultured, our knowledge and ability to predict important metabolic networks relies largely on

information from metagenomics, metatranscriptomics, and metabolomics analyses to build the

functional activities in each guild.[15, 46] We show in this work that metabolic modeling offers

additional insight that is not gained by these analyses, but that is critical in the evaluation of the

microbial communities and in developing hypothesis of how to direct metabolism towards a

specific set of fermentation products. From prior analyses,[15] we anticipated that both SEOs and

LEOs participated in MCFA production, but the iFermGuilds789 simulation of interspecies

metabolite flux (**Fig. 7**) predicted that to achieve the observed growth rate of the microbial

community, C6 and C8 were produced solely by SEOs, while LEOs were responsible for

production of C4. This difference between predictions based on metatranscriptomic and simulation

analyses can be tracked to the importance of co-utilization of fermentation products (i.e., C2 and

lactate), which is easier to explore using metabolic models.

iFermCell215 and iFermGuilds789 are examples of first-generation predictive models of a

mixed culture fermentation. Their utility can be improved by obtaining additional knowledge of

metabolic pathways involved in fermentation through studying individual microbes within each

functional guild. Further effort is needed to simulate poorly appreciated conditions that shape

community structure, such as the toxicity of fermentation products on the specific guilds or the

contribution(s) of low abundance microbes to the overall activity of the mixed culture

fermentation. Additionally, computational methods to predict the abundance of specific guilds,

rather than relying on relative abundances as a model input, are needed. Efforts to assemble such

models have begun for the human gut microbiome,[47, 48] and as these approaches are applied to

mixed microbial fermentations, their benefits to the production of MCFA and other chemicals of

societal importance will continue to improve.

ACKNOWLEDGMENTS

This work was funded by the DOE Great Lakes Bioenergy Research Center (DOE BER

Office of Science DE-SC0018409). M.J.S. was supported by the National Science Foundation

Graduate Research Fellowship Program under grant No. DGE-1256259 and by the University of

Vermont College of Engineering and Mathematical Sciences. J.J.H. was supported by the National

Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2016-

67012-24709. The authors extend our gratitude to Robert Aidan Campbell for assistance with

Python coding.

REFERENCES

1.   McGovern, P. E.; Zhang, J.; Tang, J.; Zhang, Z.; Hall, G. R.; Moreau, R. A.; Nunez, A.;
Butrym, E. D.; Richards, M. P.; Wang, C. S.; Cheng, G.; Zhao, Z.; Wang, C., Fermented beverages
of pre- and proto-historic China. *Proc. Natl. Acad. Sci. U. S. A.* **2004,** *101*, (51), 17593-8.

2.   Bond, T.; Templeton, M. R., History and future of domestic biogas plants in the developing
world. *Energy for Sustainable Development* **2011,** *15*, (4), 347-354.

3.   Angenent, L. T.; Richter, H.; Buckel, W.; Spirito, C. M.; Steinbusch, K. J. J.; Plugge, C.
M.; Strik, D.; Grootscholten, T. I. M.; Buisman, C. J. N.; Hamelers, H. V. M., Chain Elongation
with Reactor Microbiomes: Open-Culture Biotechnology To Produce Biochemicals. *Environ. Sci.*
Technol. **2016,** *50*, (6), 2796-2810.

4.   Sarria, S.; Kruyer, N. S.; Peralta-Yahya, P., Microbial synthesis of medium-chain
chemicals from renewables. *Nat. Biotechnol.* **2017,** *35*, (12), 1158-1166.

5.   Scarborough, M. J.; Lynch, G.; Dickson, M.; McGee, M.; Donohue, T. J.; Noguera, D. R.,
Increasing the economic value of lignocellulosic stillage through medium-chain fatty acid
production. *Biotechnology for Biofuels* **2018,** *11*, (1), 200.

6.   Nelson, R.; Peterson, D.; Karp, E.; Beckham, G.; Salvachúa, D., Mixed Carboxylic Acid
Production by Megasphaera elsdenii from Glucose and Lignocellulosic Hydrolysate. *Fermentation*
2017, *3*, (1), 10.

7.   Kucek, L. A.; Xu, J. J.; Nguyen, M.; Angenent, L. T., Waste Conversion into n-Caprylate
and n-Caproate: Resource Recovery from Wine Lees Using Anaerobic Reactor Microbiomes and
In-line Extraction. *Front. Microbiol.* **2016,** *7*.

8.   Zhu, X.; Zhou, Y.; Wang, Y.; Wu, T.; Li, X.; Li, D.; Tao, Y., Production of high-
concentration n-caproic acid from lactate through fermentation using a newly isolated
Ruminococcaceae bacterium CPB6. *Biotechnol Biofuels* **2017,** *10*, 102.

9.   Kucek, L. A.; Nguyen, M.; Angenent, L. T., Conversion of L-lactate into n-caproate by a
continuously fed reactor microbiome. *Water Research* **2016,** *93*, 163-171.

10.   Coma, M.; Vilchez-Vargas, R.; Roume, H.; Jauregui, R.; Pieper, D. H.; Rabaey, K.,
Product Diversity Linked to Substrate Usage in Chain Elongation by Mixed-Culture Fermentation.
Environ. Sci. Technol. **2016,** *50*, (12), 6467-76.

11.   Rodriguez, J.; Kleerebezem, R.; Lema, J. M.; van Loosdrecht, M. C. M., Modeling product
formation in anaerobic mixed culture fermentations. *Biotechnol. Bioeng.* **2006,** *93*, (3), 592-606.

12.   Zhang, F.; Zhang, Y.; Chen, M.; van Loosdrecht, M. C. M.; Zeng, R. J., A modified
metabolic model for mixed culture fermentation with energy conserving electron bifurcation
reaction and metabolite transport energy. *Biotechnol. Bioeng.* **2013,** *110*, (7), 1884-1894.

13.   Regueira, A.; Gonzalez-Cabaleiro, R.; Ofiteru, I. D.; Rodriguez, J.; Lema, J. M., Electron
bifurcation mechanism and homoacetogenesis explain products yields in mixed culture anaerobic
fermentations. *Water Res.* **2018,** *141*, 349-356.

14.   Regueira, A.; Lema, J. M.; Carballa, M.; Mauricio-Iglesias, M., Metabolic modeling for
predicting VFA production from protein-rich substrates by mixed-culture fermentation. *Biotechnol*
Bioeng **2020,** *117*, (1), 73-84.

15.   Scarborough, M. J.; Lawson, C. E.; Hamilton, J. J.; Donohue, T. J.; Noguera, D. R.,
Metatranscriptomic and Thermodynamic Insights into Medium-Chain Fatty Acid Production
Using an Anaerobic Microbiome. *mSystems* **2018,** *3*, (6).

16.   Covert, M. W.; Schilling, C. H.; Famili, I.; Edwards, J. S.; Goryanin, I. I.; Selkov, E.;
Palsson, B. O., Metabolic modeling of microbial strains in silico. *Trends Biochem. Sci.* **2001,** *26*,
(3), 179-186.

17.   King, Z. A.; Lu, J.; Drager, A.; Miller, P.; Federowicz, S.; Lerman, J. A.; Ebrahim, A.;
Palsson, B. O.; Lewis, N. E., BiGG Models: A platform for integrating, standardizing and sharing
genome-scale models. *Nucleic Acids Res.* **2016,** *44*, (D1), D515-22.

18.   Henry, C. S.; Zinner, J. F.; Cohoon, M. P.; Stevens, R. L., iBsu1103: a new genome-scale
metabolic model of Bacillus subtilis based on SEED annotations. *Genome Biol* **2009,** *10*, (6), R69.

19.   Oh, Y. K.; Palsson, B. O.; Park, S. M.; Schilling, C. H.; Mahadevan, R., Genome-scale
reconstruction of metabolic network in Bacillus subtilis based on high-throughput phenotyping
and gene essentiality data. *J Biol Chem* **2007,** *282*, (39), 28791-9.

20.   Dauner, M.; Sauer, U., Stoichiometric growth model for riboflavin-producing Bacillus
subtilis. *Biotechnol Bioeng* **2001,** *76*, (2), 132-43.

21.   Bordbar, A.; Monk, J. M.; King, Z. A.; Palsson, B. O., Constraint-based models predict
metabolic and associated cellular functions. *Nat. Rev. Genet.* **2014,** *15*, (2), 107-20.

22.   Megchelenbrink, W.; Huynen, M.; Marchiori, E., optGpSampler: an improved tool for
uniformly sampling the solution-space of genome-scale metabolic networks. *PLoS One* **2014,** *9*,
(2), e86587.

23.   Lewis, N. E.; Hixson, K. K.; Conrad, T. M.; Lerman, J. A.; Charusanti, P.; Polpitiya, A.
D.; Adkins, J. N.; Schramm, G.; Purvine, S. O.; Lopez-Ferrer, D.; Weitz, K. K.; Eils, R.; Konig,
R.; Smith, R. D.; Palsson, B. O., Omic data from evolved E. coli are consistent with computed
optimal growth from genome-scale models. *Mol Syst Biol* **2010,** *6*, 390.

24.   Ebrahim, A.; Lerman, J. A.; Palsson, B. O.; Hyduke, D. R., COBRApy: COnstraints-Based
Reconstruction and Analysis for Python. *BMC Syst. Biol.* **2013,** *7*, 74.

25.   Arkin, A. P.; Cottingham, R. W.; Henry, C. S.; Harris, N. L.; Stevens, R. L.; Maslov, S.;
Dehal, P.; Ware, D.; Perez, F.; Canon, S.; Sneddon, M. W.; Henderson, M. L.; Riehl, W. J.;
Murphy-Olson, D.; Chan, S. Y.; Kamimura, R. T.; Kumari, S.; Drake, M. M.; Brettin, T. S.; Glass,
E. M.; Chivian, D.; Gunter, D.; Weston, D. J.; Allen, B. H.; Baumohl, J.; Best, A. A.; Bowen, B.;
Brenner, S. E.; Bun, C. C.; Chandonia, J. M.; Chia, J. M.; Colasanti, R.; Conrad, N.; Davis, J. J.;
Davison, B. H.; DeJongh, M.; Devoid, S.; Dietrich, E.; Dubchak, I.; Edirisinghe, J. N.; Fang, G.;
Faria, J. P.; Frybarger, P. M.; Gerlach, W.; Gerstein, M.; Greiner, A.; Gurtowski, J.; Haun, H. L.;

He, F.; Jain, R.; Joachimiak, M. P.; Keegan, K. P.; Kondo, S.; Kumar, V.; Land, M. L.; Meyer, F.; Mills, M.; Novichkov, P. S.; Oh, T.; Olsen, G. J.; Olson, R.; Parrello, B.; Pasternak, S.; Pearson, E.; Poon, S. S.; Price, G. A.; Ramakrishnan, S.; Ranjan, P.; Ronald, P. C.; Schatz, M. C.; Seaver, S. M. D.; Shukla, M.; Sutormin, R. A.; Syed, M. H.; Thomason, J.; Tintle, N. L.; Wang, D.; Xia, F.; Yoo, H.; Yoo, S.; Yu, D., KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat. Biotechnol.* **2018,** *36*, (7), 566-569.

26.   Thauer, R. K.; Jungermann, K.; Decker, K., Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol. Rev.* **1977,** *41*, (1), 100-80.

27.   Alberty, R. A., *Thermodynamics of biochemical reactions*. John Wiley & Sons, Inc. : Hoboken, New Jersey, 2003.

28.   Han, W. H.; He, P. J.; Shao, L. M.; Lu, F., Road to full bioconversion of biowaste to biochemicals centering on chain elongation: A mini review. *J Environ Sci-China* **2019,** *86*, 50-64.

29.   Spirito, C. M.; Richter, H.; Rabaey, K.; Stams, A. J.; Angenent, L. T., Chain elongation in anaerobic reactor microbiomes to recover resources from waste. *Curr Opin Biotechnol* **2014,** *27*, 115-22.

30.   Schoberth, S.; Gottschalk, G., Considerations on the energy metabolism of Clostridium kluyveri. *Arch Mikrobiol* **1969,** *65*, (4), 318-28.

31.   Thauer, R. K.; Jungermann, K.; Henninger, H.; Wenning, J.; Decker, K., The energy metabolism of Clostridium kluyveri. *Eur J Biochem* **1968,** *4*, (2), 173-80.

32.   Yin, Y.; Zhang, Y.; Karakashev, D. B.; Wang, J.; Angelidaki, I., Biological caproate production by Clostridium kluyveri from ethanol and acetate as carbon sources. *Bioresour Technol* **2017,** *241*, 638-644.

33.   Spirito, C. M.; Marzilli, A. M.; Angenent, L. T., Higher Substrate Ratios of Ethanol to Acetate Steered Chain Elongation toward n-Caprylate in a Bioreactor with Product Extraction. *Environ Sci Technol* **2018,** *52*, (22), 13438-13447.

34.   Roghair, M.; Hoogstad, T.; Strik, D.; Plugge, C. M.; Timmers, P. H. A.; Weusthuis, R. A.; Bruins, M. E.; Buisman, C. J. N., Controlling Ethanol Use in Chain Elongation by CO2 Loading Rate. *Environ Sci Technol* **2018,** *52*, (3), 1496-1505.

35.   Weimer, P. J.; Moen, G. N., Quantitative analysis of growth and volatile fatty acid
production by the anaerobic ruminal bacterium Megasphaera elsdenii T81. *Appl Microbiol*
Biotechnol **2013,** *97*, (9), 4075-81.

36.   Zhu, X.; Tao, Y.; Liang, C.; Li, X.; Wei, N.; Zhang, W.; Zhou, Y.; Yang, Y.; Bo, T., The
synthesis of n-caproate from lactate: a new efficient process for medium-chain carboxylates
production. *Sci Rep* **2015,** *5*, 14360.

37.   Andersen, S. J.; De Groof, V.; Khor, W. C.; Roume, H.; Props, R.; Coma, M.; Rabaey, K.,
A Clostridium Group IV Species Dominates and Suppresses a Mixed Culture Fermentation by
Tolerance to Medium Chain Fatty Acids Products. *Front Bioeng Biotechnol* **2017,** *5*, 8.

38.   Seedorf, H.; Fricke, W. F.; Veith, B.; Bruggemann, H.; Liesegang, H.; Strittmatter, A.;
Miethke, M.; Buckel, W.; Hinderberger, J.; Li, F.; Hagemeier, C.; Thauer, R. K.; Gottschalk, G.,
The genome of Clostridium kluyveri, a strict anaerobe with unique metabolic features. *Proc Natl*
Acad Sci U S A **2008,** *105*, (6), 2128-33.

39.   Scarborough, M. J.; Myers, K. S.; Donohue, T. J.; Noguera, D. R., Medium-Chain Fatty
Acid Synthesis by "Candidatus Weimeria bifida" gen. nov., sp. nov., and "Candidatus
Pseudoramibacter fermentans" sp. nov. *Appl Environ Microbiol* **2020,** *86*, (3).

40.   Hedderich, R.; Forzi, L., Energy-converting [NiFe] hydrogenases: More than just H-2
activation. *J Mol Microb Biotech* **2005,** *10*, (2-4), 92-104.

41.   Juturu, V.; Wu, J. C., Microbial production of lactic acid: the latest development. *Crit. Rev.*
Biotechnol. **2016,** *36*, (6), 967-977.

42.   Buschke, N.; Schafer, R.; Becker, J.; Wittmann, C., Metabolic engineering of industrial
platform microorganisms for biorefinery applications--optimization of substrate spectrum and
process robustness by rational and evolutive strategies. *Bioresour. Technol.* **2013,** *135*, 544-54.

43.   Sarria, S.; Bartholow, T. G.; Verga, A.; Burkart, M. D.; Peralta-Yahya, P., Matching
Protein Interfaces for Improved Medium-Chain Fatty Acid Production. *ACS Synth Biol* **2018,** *7*,
(5), 1179-1187.

44.   Hernandez Lozada, N. J.; Lai, R. Y.; Simmons, T. R.; Thomas, K. A.; Chowdhury, R.;
Maranas, C. D.; Pfleger, B. F., Highly Active C8-Acyl-ACP Thioesterase Variant Isolated by a
Synthetic Selection Strategy. *ACS Synth Biol* **2018,** *7*, (9), 2205-2215.

45.   Deng, X.; Chen, L.; Hei, M.; Liu, T.; Feng, Y.; Yang, G. Y., Structure-guided reshaping of
the acyl binding pocket of 'TesA thioesterase enhances octanoic acid production in E. coli. *Metab*
Eng **2020,** *61*, 24-32.

46.   Han, W.; He, P.; Shao, L.; Lu, F., Metabolic Interactions of a Chain Elongation
Microbiome. *Appl. Environ. Microbiol.* **2018**.

47.   Zomorrodi, A. R.; Maranas, C. D., OptCom: a multi-level optimization framework for the
metabolic modeling and analysis of microbial communities. *PLoS Comput. Biol.* **2012,** *8*, (2),
e1002363.

48.   Kumar, M.; Ji, B.; Zengler, K.; Nielsen, J., Modelling approaches for studying the
microbiome. *Nat Microbiol* **2019,** *4*, (8), 1253-1267.
674

Figure 1. Summary of metabolic networks in iFermCell215. Metabolites with an asterisk are required for biomass production. Open hexagons indicate consumption of ATP (orange), NADH (green), NADPH (grey), and reduced ferredoxin (red). Closed hexagons indicate production of these compounds and consumption of ADP (orange), NAD+ (green), NADP+ (grey), and oxidized

ferredoxin (red). For reversible reactions, larger arrowheads indicate the reaction directionality

corresponding with consumption and/ or production of ATP, NADH, NADPH and reduced

ferredoxin.. Dashed lines indicate multiple reactions. Key reactions are indicated according to the

abbreviation used in iFermCell215 (**Supplementary File 3**). Pathways are identified by number:

(1) upper glycolysis; (2) phosphoketolase metabolism of fructose; (3) non-oxidative pentose

phosphate pathway; (4) phosphoketolase metabolism of xylulose; (5) glycerol utilization; (6) lower

glycolysis; (7) gluconeogenesis; (8) pyruvate formate lyase; (9) lactate metabolism; (10) pyruvate

flavodoxin oxidoreductase; (11) pyruvate dehydrogenase; (12) acetate metabolism; (13) ethanol

metabolism; (14) incomplete citric acid cycle with glyoxylate shunt; (15) anaplerotic metabolism

of phosphoenolpyruvate, pyruvate, and oxaloacetate; (16) ATP citrate lyase; (17) propionate

production via the acryloyl-CoA pathway; (18) propionate production via the methylmalonyl-CoA

pathway; (19) even-chain reverse β-oxidation; (20) odd-chain reverse β-oxidation; (21)

homoacetogenesis via the Wood Ljungdahl pathway; (22) ferredoxin-dependent hydrogenase; (23)

proton-translocating hydrogenase; (24) electron confurcating hydrogenase; (25) proton-

translocating ferredoxin: NAD^+ oxidoreductase; (26) NAD(P) transhydrogenase. For clarity, not

all reactions or metabolites are shown. More information on the reactions and metabolites

contained in iFermCell215 is provided in **Supplementary File 3**.

Figure 2. Summary of metabolic networks for six functional guilds contained in iFermGuilds789.

Refer to **Fig. 1** for details on pathways included in the iFermGuilds789 model. More information

on the reactions and metabolites contained in the iFermGuilds789 model is provided in

Supplementary File 4.

Figure 3. Flux sampling results for iFermCell215 with objective function to optimize biomass

production when fed substrates indicated within the plots: (A) glucose, (B) xylose, (C) glycerol,

(D) lactate, (E) ethanol, (F) H_2 and CO_2 (with H_2 as the limiting substrate). Box plots show the

results of 101 simulations. Median values are indicated by a horizontal line within the box, and the

top and bottom of the box indicate the 25^{th} and 75^{th} percentile, respectively. Whiskers indicate

values within 1.5 times the interquartile range and closed circles indicate values outside of this

range.

Figure 4. Flux sampling results for iFermCell215 when consuming ethanol and acetate (A-D) or

lactate and acetate (E-H) at different molar ratios of acetate-to-ethanol (C2:EtOH) or acetate-to-

lactate (C2:Lactate). Results are shown for fluxes of octanoate (C8), hexanoate (C4), butyrate (C4)

and H_2. These results are shown as box plots, as described previously (Figure 3), representing the

results of 101 simulations for each modeled ratio. The predicted maximum growth rate (hr^{-1}) is

shown for ethanol (A) and lactate (E) as red triangles plotted on the secondary y-axis.

Figure 5. Predicted pathways of ethanol utilization by iFermCell215 using pFBA when (A) ethanol is the sole substrate; (B) ethanol and acetate are co-substrates; (C) lactate is a sole substrate; and (D) lactate and acetate are co-substrates. Flux values (shown as white text) have units of mmol gDCW⁻¹ hr⁻¹. For both scenarios, the model was constrained to uptake 1 mmol gDCW⁻¹ hr⁻¹ of ethanol or 1 mmol gDCW⁻¹ hr⁻¹ of lactate. For scenarios with acetate co-utilization, acetate was provided in excess. Key reactions indicated by red text include a ferredoxin hydrogenase (Hyd1), a CoA transferase (CoAT), a proton-translocating ferredoxin: NAD+ oxidoreductase (RNF1), ATP synthase (ATPS4r), NADH:NADP transhydrogenase (THD2), and energy-conserving hydrogenase (ECH). Metabolite abbreviations are and are defined in **Supplementary Data File 2**.

Figure 6. Predicted and observed product formation for a mixed culture fermentation bioreactor fed lignocellulosic biorefinery residues. (A) predictions with iFermCell215; (B) predictions with iFermGuilds789; (C) predictions with iFermGuilds789 with reactions knocked-out based on metagenomic analyses of the microbial community; (D) predictions with iFermGuilds789 with additional constraints for maximum fluxes through acyl-CoA and CoAT reactions for C6 and C8 production in SEOs and LEOs

Figure 7. Predicted substrate flow through a mixed culture fermentation producing a mixture of MCFA from a complex feedstock using iFermGuilds789. Values were obtained using pFBA while constraining the products according to observed bioreactor performance. The relative abundance of each functional guild was assumed based on reactor DNA abundance. The relative amount of eeq in the substrates, products, and intermediates are also provided. Line width indicates the relative amount of each substrate consumed and product created by each of the functional guilds. H_2 was not measured, so for these simulations it was assumed that the difference between eeq in the substrate and eeq in the measured soluble and insoluble products represented H_2 production. In this simulation, 22% of the eeq in conversion residue was predicted to be directed to lactate as an intermediate and 27% to acetate.

Figure 8. Predicted chain elongation metabolism by SEOs and LEOs using iFermGuilds789.

Predicted fluxes values are shown as white numbers and have units of mmol gDCW^{-1} hr^{-1}. (A)

SEOs are predicted to produce all of the hexanoate (hxa) and octanoate (octa) in the community

and use an acyl-CoA hydrolase (ACH) reaction as the terminal step of reverse β-oxidation. (B)

LEOs are predicted to produce all of the butyrate in the community while using a CoA transferase

758   (CoAT). LEOs are predicted to recycle all of the H_2 that they produce via an energy conserving

759   hydrogenase.