Enhancing Myocardial Repair with CardioClusters

Megan M. Monsanto, BS[1], Bingyan J. Wang, MS[1], Zach R. Ehrenberg, BS[1], Oscar Echeagaray, BS[1], Kevin S. White, BS[1], Roberto Alvarez Jr., PhD[1], Kristina Fisher, BS[1], Sharon Sengphanith, BS[1], Alvin Muliono, BS[1], Natalie A. Gude, PhD[1], Mark A. Sussman, PhD[1]*

[1]San Diego Heart Research Institute, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182, USA

*To whom correspondence should be addressed

SDSU Heart Institute and Department of Biology
San Diego State University
5500 Campanile Drive, San Diego, CA 92182
(619) 594-2983 (voice)
(619) 594-2610 (fax)
heartman4ever@icloud.com (e-mail)

Running Title: Enhancing Myocardial Repair with CardioClusters
Abstract word count: 390
Main text word count: 5,957

Abstract

Background: Cellular therapy to treat heart failure is an ongoing focus of intense

research and development, but progress has been frustratingly slow due to limitations of

current approaches. Engineered augmentation of established cellular effectors

overcomes impediments, enhancing reparative activity with improved outcomes relative

to conventional techniques. Such 'next generation' implementation includes delivery of

combinatorial cell populations exerting synergistic effects. Concurrent isolation and

expansion of three distinct cardiac-derived interstitial cell types from human heart

tissue, as previously reported by our group, prompted design of a three-dimensional

(3D) structure that maximizes cellular interaction, allows for defined cell ratios, controls

size, enables injectability, and minimizes cell losses upon delivery.

Methods: Three distinct populations of human cardiac interstitial cells including

mesenchymal stem cells (MSCs), endothelial progenitor cells (EPCs), and c-Kit$^+$ cardiac

interstitial cells (cCICs) when cultured together spontaneously form scaffold-free 3D

microenvironments termed CardioClusters. Biological consequences of CardioCluster

formation were assessed by multiple assays including single cells RNA-Seq

transcriptional profiling. Protective effects of CardioClusters in vitro were measured

using cell culture models for oxidative stress and myocardial ischemia in combination

with freshly isolated neonatal rat ventricular myocytes. Long-term impact of adoptively

transferred CardioClusters upon myocardial structure and function in a xenogenic model

of acute infarction using NOD^{scid} mice was assessed over a longitudinal time course of

20-weeks.

Results: CardioCluster design enables control over composite cell types, cell ratios, size, and preservation of structural integrity during delivery. Profound changes for biological properties of CardioClusters relative to constituent parental cell populations include enhanced expression of stem cell-relevant factors, adhesion/extracellular-matrix molecules, and cytokines. The CardioCluster 3D microenvironment maximizes cellular interaction while maintaining a more native transcriptome similar to endogenous cardiac cells. CardioCluster delivery improves cell retention following intramyocardial injection with preservation of long-term cardiac function relative to monolayer-cultured cells when tested in an experimental murine infarction model followed for up to 20 weeks post-challenge. CardioCluster-treated hearts show increases in capillary density, preservation of cardiomyocyte size, and reduced scar size indicative of blunting pathologic infarction injury.

Conclusions: CardioClusters are a novel 'next generation' development and delivery approach for cellular therapeutics that potentiate beneficial activity and enhance protective effects of human cardiac interstitial cell mixed populations. CardioClusters utilization in this preclinical setting establishes fundamental methodologic and biologic insights, laying the framework for optimization of CardioCluster design to provide greater efficacy in cell-based therapeutic interventions intended to mitigate cardiomyopathic damage.

Introduction

Cellular therapy continues to be pursued as an experimental approach to treat cardiomyopathic damage following infarction in both preclinical animal models and human clinical trials[1-9]. While promising, cellular therapy has been hindered by marginal improvement in cardiac function, in part due to low engraftment and persistence of the transplanted cells[10-12]. As such, improving cell retention and survival has been a major area of research using a plethora of techniques, including biomaterials[13, 14], cytokines and growth factors[15], repeated administration of cells[16], or genetic enhancement with pro-survival and anti-apoptotic genes[17, 18]. Combinatorial cell therapies are an additional conceptual approach intended to promote additive or synergistic effects in myocardial repair[19-22]; however, cell retention and survival remain poor whether single or combinatorial cell delivery is attempted. Furthermore, coincident delivery of multiple cell populations does not ensure effective cell interaction and cross talk following injection. Therefore, novel strategic approaches to prolong cell retention and survival are highly sought after, preferably also favoring cell phenotypic characteristics that promote short-term mitigation of injury and long-term recovery of myocardial structure and function.

Traditional monolayer culture, while both prevalent and convenient, promotes loss of cell identity and spatial organization, thereby impacting many cellular aspects, including cell morphology, proliferation, differentiation, viability, and transcriptome profile[23-25]. Approaches intended to mitigate deleterious effects of artificial *in vitro* environments through recreating native three dimensional (3D) architecture include engineered heart tissue grafts[26, 27], organoids[28, 29], and 3D bioprinting.[30, 31] When grown in 3D cells aggregate and form complex cell-cell and cell-matrix interactions that mimic

the natural environment found *in vivo,* allowing for cell differentiation and tissue

organization not possible in conventional two-dimensional (2D) culture systems[32]. For

example, cardiosphere formation[33, 34] enhances cellular communication through gap

junction protein connexin 43 fostering signal propagation and cell differentiation*[35-37]*.

However, cardiospheres are formed by random cell aggregation leading to inconsistent

cell characterization markers[33, 34], variable sphere size, and muddled communication

within the aggregate microenvironment[33, 34, 38]. A superior preferable methodology would

utilize combinations of defined cell populations aggregated in 3D under controlled

conditions that allow for intact delivery without dissociation of the spherical

microenvironment to promote preservation of cellular phenotype and optimize cellular

retention. This combination of desired features formed the rational basis for our novel

methodological approach distinct in several critical aspects from cardiosphere-derived

cell therapy.

A 'next generation' conceptual approach for a deliverable cell therapeutic based

upon the aforementioned literature precedents would involve 1) combinations of multiple

cardiac-derived cell types, 2) *ex vivo* culture adaptation to a 3D microenvironment to

maximize cellular interaction, 3) control over cell composition, cell number, and

aggregate cell volume, and 4) capability to deliver *ex vivo* formed cell aggregate 'niches'

intact to facilitate preservation of microenvironment biological properties and retention.

Guided by these stipulated parameters to create a novel artificially engineered cell

aggregate, inception began with essential methodological studies to isolate and expand

three distinct cardiac-derived interstitial cell types from a single human heart biopsy as

previously reported by our group[39]. Human cardiac tissue of varying age and gender

was used as source material to isolate c-Kit[+] cardiac interstitial cells (cCIC),

mesenchymal stem cells (MSCs), and endothelial progenitor cells (EPCs). Having

isolated and characterized three distinct cardiac interstitial cell types each known to

possess beneficial properties to blunt cardiomyopathic damage, a protocol was

designed for spontaneous self-assembly of mixed cell populations into an optimal

conceptual engineered solution that we have named "CardioClusters."

CardioClusters harness the distinct phenotypic attributes of three well defined

cardiac cell populations. MSCs support myocardial reparative activity through secretion

of paracrine factors that activate endogenous cells, promote angiogenesis, protect

cardiomyocytes, and reduce scar formation[40, 41]. MSCs also secrete cell adhesion

molecules such as integrins and cadherins integral to cellular aggregation[42]. EPCs

promote paracrine dependent vasculogenesis and angiogenesis and differentiate into

mature endothelial cells[43, 44]. Prior studies demonstrate EPCs transplanted *in vivo* are

capable of forming microvessels, but regress without MSCs to support vessel maturity[45,

46]. cCICs distributed in the CardioCluster contribute to support of myocardial

homeostasis, response to injury, and remodeling. CardioCluster characteristics are

particularly well suited to mediating myocardial repair in the wake of acute pathological

damage by providing a more natural niche microenvironment for augmented delivery

and extended functional activity to improve the outcome of cellular therapy.

CardioClusters advance the application of combinatorial cell therapy by

integrating complementary and synergistic properties from multiple cardiac-resident cell

types into a single injectable product. CardioCluster formation is a rapid, reproducible,

and controllable process demonstrated in preclinical testing to mediate significant

improvements in myocardial structure and function following infarction injury.

Enhancement of cell therapy with CardioClusters represents an important advance to

improve clinical application of cell therapy for treatment of cardiomyopathic damage and

disease.

Methods

Full materials and methods are available in the Online Data Supplements.

Human Cardiac Interstitial Cell Isolation

NIH guidelines for human research were followed as approved by IRB review (Protocol #120686). Neonatal heart tissue procured from post-mortem infants provided by a commercial source (Novogenix Laboratories) was used for isolation of human cardiac cells. The protocol detailing cardiac cell isolation can be found in our previous publication[39] and Online Data Supplement. All cells used in this study were mid-passage (passages 5–10).

CardioCluster Formation

CardioClusters are formed using 96 well, ultra-low attachment multiwell round-bottom plates (Corning, catalog #CLS7007) in a two-step process. Step 1 generates the inner core composed of cCICs and MSCs in a 1:2 ratio. The inner core of cCICs and MSCs is seeded in 100 µL/well MSC media for 24 hours at 37°C in 5% CO_2 incubator. Step 2 forms the outer EPC layer using a cell number equal to the number of cells used to create the central core. The EPCs are added in an additional 50 µL/well MSC media and incubated at 37°C in 5% CO_2 for an additional 24 hours until CardioCluster 3D structure has formed. The radius of a CardioCluster is approximately 150 µm, composed of a total of 400±100 cells.

Single-cell RNA-seq Preparation, Data Analysis, and Data Availability

1   Size distribution was quantified for single cell suspensions of cCIC, MSC, EPC

2 and dissociated CardioClusters to verify cell size met droplet platform specifications

3 (Supplemental Figure 6A). Cells were loaded on a Chromium™ Controller (10x

4 Genomics) and single-cell RNA-Seq libraries were prepared using Chromium™ Single

5 Cell 3' Library & Gel Bead Kit v2 (10x Genomics) following manufacturer's

6 protocol. Each library was tested with Bioanalyzer (average library size: 450-490 bp).

7 The sequencing libraries were quantified by quantitative PCR (KAPA Biosystems

8 Library Quantification Kit for Illumina platforms P/N KK4824) and Qubit 3.0 with dsDNA

9 HS Assay Kit (Thermo Fisher Scientific). Sequencing libraries were loaded at 2 pM on

10 an Illumina HiSeq2500 with 2X75 paired-end kits using the following read length: 98 bp

11 Read1, 8 bp i7 Index, and 26 bp Read2. Batch effects concerns were obviated in our

12 analyses by mitigating strategies in our experimental design. Specifically, the following

13 steps were taken: 1) all samples were processed in the same microfluidics chip, and 2)

14 libraries were prepared in parallel and sequenced in the same lane in the HiSeq 4000.

15 Libraries were aggregated and normalized using CellRanger. Final removal of unwanted

16 sources of variation and batch effect corrections was performed using Seurat R

17 Package (v2.3.4)[47].

18   Raw sequencing data was processed with the Cell Ranger pipeline (10X

19 Genomics; version 2.0). Sequencing reads were aligned to the human genome hg19.

20 Cells with fewer than 1,000 genes or more than 10% of mitochondrial gene UMI count

21 were filtered out and genes detected in fewer than three cells were filtered out using

22 Seurat R Package (v2.3.4)[47]. The first 20 principal components were found to be

23 significant to perform dimensionality reduction. Preparations derived from *in vitro*

studies yielded 5659 barcoded cells for analysis, from which 1125, 1717, 1403 and

1414 corresponded to CardioClusters, cCICs, EPCs and MSCs, respectively.

Approximately 2,029 variable genes were selected based on their expression and

dispersion. The first 20 principal components were used for the t-SNE projection and

unsupervised clustering[47]. Differential expression analysis was done using Wilcoxon

rank sum test and selecting for an adjusted p-value ≤0.05 and a log (FC)>0.25. Global

differential expression analysis was done using Loupe Cell Browser 2.0.0. Gene

ontology analysis was performed using R package clusterProfiler[48]. scRNA-Seq data

generated in this study have been uploaded to the Gene Expression Omnibus (GEO)

database (accession number GSE133832). Datasets for freshly isolated CICs have

been previously published by our group[25] and are available at the GEO database

(accession number GSE114280).

Myocardial Infarction and Intramyocardial Injection

Animal protocols and experimental procedures are approved by the Institutional

Animal Care and Use Committee at San Diego State University. Animals were

randomized for treatments that were blinded to personnel carrying out the surgical

procedures, injections, and physiological function analysis. Appropriate animal sample

size was determined using sample size calculator (http://www.lasec.cuhk.edu.hk/sample-size-

calculation.html). A total of 60 mice were used in this study. Inclusion criteria were set prior

to study commencement and required a drop in ejection fraction between 10-50% 1-

week following infarction injury. Myocardial infarctions were carried out on 8-week old

NOD.CB17-Prkdc^{scid}/J female mice (The Jackson Laboratory, catalog #001303) under

2% isoflurane (Victor Medical, catalog #NDC 57319-474-06) as previously described[49].

Briefly, the 3rd and 4th ribs were separated enough to get adequate exposure of the

operating region, but the ribs were kept intact. The heart was squeezed out by pressing

the thorax lightly and the left anterior descending artery (LAD) was ligated at the distal

diagonal branch with a 7-0 suture. Infarction was confirmed by blanching of anterior left

myocardium wall. Following ligation, either CardioClusters (n=17), 2:3:1 ratio of

cCIC+EPC+MSC (C+E+M; n=15), or a vehicle control (PBS plus 0.5% sodium alginate

[NovaMatrix, catalog #4209001]; n=16) were delivered intramyocardially at 3 separate

sites in the vicinity bordering the blanched area. A total of 90,000 cells per heart were

introduced into CardioCluster and C+E+M animals. The hearts were immediately placed

back into the intrathoracic space followed by muscle and skin closure. Animals in sham

group (n=12) received a comparable surgical procedure without LAD ligation or

injection. Each animal received 20 µL analgesic treatment with 0.3 mg/mL Buprenex

(Victor Medical, catalog #12496-0757) at time of surgery and 12 hours post-surgery. No

animals required exclusion from study.

Statistical Analysis

Data are expressed as mean±SEM. Statistical analyses of multiple groups were

assessed by 1-way ANOVA with Bonferroni post hoc test (comparison among all

groups) or Dunnett's post hoc test (versus single group). Multiple groups over time were

analyzed by 2-way ANOVA. Statistical analysis was performed using GraphPad Prism

version 5.0 software. Experiments were performed in triplicate unless stated otherwise.

A p-value of less than 0.05 was considered statistically significant.

Results

Three distinct cardiac nonmyocyte cell types are used for CardioCluster

formation

CardioClusters are comprised of three cardiac-resident cell populations concurrently isolated as previously described by our group with full phenotypic characterization[39]. Cell surface marker profiling of cell lines used to produce CardioClusters were similar to previous findings (cCIC express c-Kit^{high}, CD90^{high}, CD105^{low}, CD133^{low}, CD45^{neg}; EPC express CD133^{high}, CD105^{high}, c-Kit^{low}, CD90^{neg}, CD45^{neg}; and MSC express CD90^{high}, CD105^{high}, c-Kit^{low}, CD133^{low}, CD45^{neg}; data not shown). Commitment toward angiogenic and smooth muscle fates was assessed for the three cardiac-derived cell populations relative to control cell lines human umbilical vein endothelial cells (HUVECs) and bone marrow-derived MSCs (BM MSCs). Tube formation assays demonstrated robust angiogenic responses from both EPCs and HUVECs *in vitro* using growth factor-reduced Matrigel (Supplemental Figure 1A). Transcript levels of endothelial-related genes *CD31* and *von Willebrand factor* (*vWF*) were elevated in HUVECs and EPCs (p<0.001 versus cCICs; Supplemental Figure 1B and 3C). *Smooth muscle actin* (*SMA*) transcripts were highly expressed in BM MSCs and cardiac MSCs (p<0.001 versus cCICs), with both endothelial populations (EPC or HUVEC) expressing near undetectable levels (p<0.05 versus cCICs; Supplemental Figure 1D). *GATA4* was expressed by cCICs (1.0±0.05) and to a lesser extent by EPCs (0.87±0.03) and MSCs (0.33±0.01), with non-cardiac controls expressing undetectable levels (Supplemental Figure 1E). Collectively, these three cardiac-derived cell populations recapitulate and validate previous results of phenotypic characterization for

cell types obtained using our published protocol[39]. Distinct phenotypic properties of

these three cardiac-derived cell populations fulfills the conceptual design of combining

multiple cell types for CardioClusters formation.

The three distinct cardiac derived cell populations were modified with lentiviral

vectors to introduce fluorescent proteins for tracking purposes (eGFP tagged cCICs

[green], mOrange tagged EPCs [blue], and Neptune tagged MSCs [red]; tagging

efficiency 99.1±0.2%; Supplemental Figure 2A and 2B). Distinct morphology for each

cell population is evident in representative brightfield images with companion

immunofluorescent images demonstrating corresponding fluorophore expression in

cCICs (Figure 1A), EPCs (Figure 1B), and MSCs (Figure 1C). Cell morphology

measurement of area, roundness, and L/W ratio for each cell type confirmed distinct

phenotypes (Figure 1, D-F). MSCs were significantly larger (18,563±1,021) relative to

both cCIC (3383±121) and EPC (3272±102) (Figure 1D). EPCs were significantly

rounder (EPC, 0.55±0.012; cCIC, 0.19±0.0097; MSC, 0.36±0.015) (Figure 1E), while

cCICs show increased L/W ratio (cCIC, 5.2±0.19; EPC, 2.1±0.063; MSC, 2.8±0.11)

(Figure 1F). Morphometric parameters clustered by cell type (Supplemental Figure 3),

with minor variation between heart samples. EPCs exhibited a proliferative rate similar

to cCICs, with both populations showing increased proliferation over MSCs based on

CyQuant proliferation assays (Figure 1G). EPCs were significantly more resistant to cell

death and retained 92±0.76% cell viability, versus only 54±5.6% for cCIC and 79±1.5%

for MSCs after 4 hours H_2O_2 treatment (Figure 1, H-J). Cumulatively, characterization

showed phenotypic and biological distinctions between cardiac interstitial cell

populations fundamental to CardioCluster design and utility, such as elevated

resistance to oxidative stress-induced cell death, high proliferative activity, and pro-

angiogenic nature of EPCs.

Generation of CardioClusters

CardioClusters are formed in a two-step process (Figure 1K). cCICs and MSCs

are seeded to form the inner core, with EPCs added 24 hours later to provide an

endothelial cell-enriched outer layer for the CardioCluster. The outer EPC layer provides

enhanced resistance to oxidative stress relative to the more sensitive cCICs and MSCs

within the CardioCluster core (Figure 1J). Individual cells within an assembled

CardioCluster were readily visualized with their cognate fluorophore tags, obviating the

need for antibody-mediated detection (Figure 1L and 1M). CardioCluster size

reproducibly and predictably corresponds to cell number seeded per microwell (Figure

1N). CardioClusters ranging from 100-1000 cells were examined to determine changes

in size and morphology over a 7 day time period (Supplemental Figure 2D).

CardioCluster diameter and area increased over 7 days, except for CardioClusters

seeded with 1000 cells, whose diameter plateaued after day 3. This finding, also

previously observed with 3D aggregated cells[50], is consistent with reduced oxygen and

nutrient diffusion within dense cellular structures >200 µm[51, 52].

Preserving CardioCluster 3D structural integrity for intramyocardial delivery to

promote intercellular contact and enhance retention is essential to improve upon typical

approaches involving dissociated single cell suspensions. Intramyocardial injection for

cell delivery in murine hearts uses a standard 30-gauge needle with a 159 µm internal

diameter ([I.D.]), so CardioClusters were engineered for a diameter allowing for injection

to preserve 3D structure. The maximum number of cells that could comprise a

CardioCluster and pass through a 30-gauge needle is 400 cells based on morphometric

quantitative analysis (Figure 1O). EPC, cCICs and MSCs were combined in a 3:2:1

ratio, based upon the consideration of larger MSC size occupying relatively more

volume relative to cCIC or EPC (Figure 1D). CardioCluster spontaneous self-assembly

as revealed using time-lapse video microscopy shows the MSC population immediately

migrating towards the central core (Supplemental Video 1). cCIC/MSC interaction was

allowed to progress for 24 hours, at which point EPCs were added to interact with

established cCIC/MSC cores. Interestingly, EPCs initially form their own clusters instead

of adhering to cCIC/MSC cores and then subsequently envelop the cCIC/MSC core

(Supplemental video 1 and Supplemental Figure 2). Architecture of an MSC-enriched

core was invariant regardless of seeding sequence, as plating of cCIC+EPCs prior to

adding MSCs consistently resulted in MSCs migrating and localizing within the

CardioCluster core rather than surface (Supplemental Video 2) consistent with the

preferential localization of MSC to hypoxic environments.[53, 54]. CardioCluster formation

consistently occurs with MSCs in the core and cCIC/EPCs on the outer layers.

      CardioClusters possess a high percentage of live cells maintained within the 3D

structure (93.9-98% of cells alive; Supplemental Figure 4). Robust vitality of

CardioClusters was confirmed by recovery from long term liquid nitrogen storage, where

the percentage of live cells was comparable to that of control non-frozen CardioClusters

(Supplemental Figure 4A and 4B). When cultured on standard tissue culture-treated

plastic, cells adhered and migrated out from the CardioCluster whether frozen or not

non-frozen with comparable cell morphology (Supplemental Figure 4C). These findings

support "off-the-shelf" feasibility of using frozen banked CardioClusters for therapeutic

purposes rather than necessitating *de novo* creation prior to use.

CardioCluster cells undergo reprogramming toward the transcriptome profile of

freshly isolated cardiac interstitial cells

Transcriptional profiling of CardioClusters and their monolayer cultured parental

counterparts reveals significant reprogramming consequential to 3D aggregation. Since

CardioClusters are heterogeneous cell populations by design, single-cell RNA

sequencing (scRNA-seq) was employed to reveal cellular transcriptome heterogeneity

within the CardioCluster (Supplemental Figure 5) at the single cell level with a level of

resolution not achievable with bulk population analysis. Quality control testing validated

parameters of cell size distribution, sequence alignment and filtering of multiplets and

dying cells (Supplemental Figure 6). Dimensionality reduction by t-SNE reveals

segregation of CardioClusters (orange cluster) distinguished by a unique transcriptome

profile separating them from their constituent parental populations, which form their own

clusters (red, green, blue clusters representing MSCs, cCICs and EPCs respectively;

Figure 2A). Differentially expressed genes (DEGs) are increased in the CardioCluster

environment (620) relative to cCIC (296), EPC (167), or MSC (211) (Figure 2B,

Supplemental Table 4). Taken together, these results demonstrate CardioClusters

distinguishing themselves as a transcriptionally unique population diverged from

parental cell lines.

Cellular identities of the three constituent parental lines comprising a

CardioCluster are consistent with detected transcripts in each cell type. EPCs highly

express ECSCR, ESM1, EGFL7 and RAC2, which are endothelial–related genes

important for neovasculature and the angiogenic response (Figure 2B and 2C). The

highly specific marker vascular endothelial statin (VE-statin) referred to as EGFL7,

exhibits near-exclusive expression and action upon endothelial cells[55] and is highly

expressed in EPCs (Figure 2B). In contrast, many secreted angiogenic signaling

molecules including vascular endothelial growth factor (VEGF) are expressed by

nonendothelial cell types such as fibroblasts. MSC-enriched transcripts include the cell

surface marker THY1 (also referred to as CD90) as well as Smooth muscle α-2 actin

(ACTA2) (Figure 2B and 2C). Gene ontology analysis of MSCs reveals expression of

extracellular matrix and adhesion molecules such as COL1A2, TIMP3 and FN1 (Figure

2B and 2D). Consistent with preference for hypoxia, MSCs are enriched for HIF1A, a

transcription factor that plays a key role in response to hypoxic stimuli (Figure 2B).

Lastly, transcripts associated with cell proliferation and anti-apoptotic activity such as

BIRC5 and HMGB1 are differentially expressed in cCICs, as well as the chemotactic

signaling molecule CXCL12, commonly referred to as stromal derived factor-1 (SDF-1),

and developmental genes DKK1 and FGF5 (Figure 2B). Collectively, these data

highlight that the 3 parental populations are distinctly different from one another, with

EPCs and MSCs expressing endothelial and stromal-associated genes.

Transcriptome profiling of CardioClusters by scRNA-seq reveals several features

distinct from the three parental cell lines. CardioClusters are enriched for transcripts in

multiple categories including stem cell-relevant factors (KLF4, LIF, JAK1, SMAD7,

BAMBI, NOTCH3), adhesion/extracellular-matrix molecules (integrin-α2, laminin-γ1,

type 1 collagen-α1, BMP1, MMP2), and cytokines (SOD2, SDF-1, FGF2). These

aforementioned DEGs were similarly enriched in freshly isolated cardiac interstitial cells

(Figure 2B-D, 2F, and Supplemental Table 5), suggesting that CardioClusters adopt a

transcriptome profile with features reminiscent of cardiac interstitial cells present in the

myocardium rather than cultured cells. Indeed, 89 out of the 448 DEGs present in

CardioClusters are also found in freshly isolated cardiac interstitial cells (Figure 2E and

2F, Supplemental Table 5), in stark contrast to the overlap with the 2D-cultured parental

cell lines where only 7 DEGs are shared with freshly isolated cardiac interstitial cells

(Figure 2E and 2F, Supplemental Table 5). This finding supports that standard tissue

culture causes expanded cells to lose their identity, unlike cells within a 3D

microenvironment. Consistent with this observation, the size of cells grown within a

CardioCluster were smaller relative to 2D cultured parental counterparts (p<0.01;

Supplemental Figure 6A) resembling a size more similar to freshly isolated cells,

consistent with cellular remodeling that occurs alongside topological changes[56]. Thus,

the 3D microenvironment of a CardioCluster promotes a more native phenotype similar

to endogenous or freshly isolated cardiac cells.

CardioClusters exert protective effects with serum starvation *in vitro* assay

Protective effects mediated by cCICs and MSCs are conferred upon serum

starved neonatal rat cardiomyocytes (NRCMs) in co-culture[57]. Similarly, beneficial

effects mediated by CardioClusters were assessed by co-culture with NRCMs in serum

depleted conditions relative to effects conferred by cCIC, EPCs, MSCs, and a combined

mixture of cCIC+EPC+MSC (C+E+M) (Figure 3A). NRCMs maintained in low serum

(0.5%) were smaller relative to NRCMs maintained in high serum condition (10%)

(Figure 3B and 3C, Supplemental Figure 7). CardioCluster co-culture with low serum

treated NRCMs restored cardiomyocyte size within 24 hours relative to all other

treatments (p<0.05; Figure 3B and 3C, Supplemental Figure 7) and also increased

mRNA expression for *Desmin,* a muscle-specific type III intermediate filament protein

(p<0.001; Figure 3D). Furthermore, CardioCluster co-culture increased mRNA for *Sdf-1*

(p<0.05; Figure 3E), a cardioprotective cytokine and chemotactic factor for MSCs that

plays an additional role in recruitment of EPCs important for angiogenesis[58, 59].

Importantly, CardioClusters offered significantly greater protection upon NRCM than

actions exerted by any individual parental population (cCICs, EPCs, MSCs) or the

combined C+E+M mixed population. Collectively, these results demonstrate superior

protective effects of CardioClusters for NRCM in response to serum starvation

challenge.

Paracrine gene expression is increased in CardioClusters after NRCM *in vitro* co-

culture

16   Paracrine factor action is considered a primary mechanism for cardioprotective[60],

so mRNA transcript level expression for growth and immunomodulatory factors was

assessed after CardioCluster co-culture for 5 days with serum depleted NRCM

(Supplemental Figure 8A). mRNA levels for CardioClusters, parental cells, and the

C+E+M mixed population were measured by separating fluorescently tagged cells away

from the NRCM population using flow cytometric sorting (Supplemental Figure 7).

mRNA transcript levels for *insulin-like growth factor* (*IGF*) and *interleukin-6* (*IL-6*) were

highly elevated in CardioClusters co-cultured with NRCMS relative to any of the

individual parental population (cCICs, EPCs, MSCs) or the combined C+E+M mixed

population (p<0.001 and p<0.05 respectively, versus cCIC; Supplemental Figure 8B and

7C). *IGF* exerts chemotactic and growth-stimulatory effects[60] in addition to anti-

apoptotic properties[61-63]. Early release of anti-inflammatory cytokines such as *IL-6* after

acute cardiac damage has been shown to be beneficial by signaling protective

responses in local tissue and initiating wound healing[64]. Additionally, the

cardioprotective cytokines *SDF-1* and *hepatocyte growth factor* (*HGF*), both trended

towards increased expression in CardioClusters following co-culture experiments

(Supplemental Figure 8D and 8E). *HGF* stimulates cell proliferation, motility,

morphogenesis, angiogenesis and importantly tissue regeneration[63, 65]. Collectively

these results show that at the transcript level CardioClusters induction of paracrine

factors *IGF* and *IL-6* exceeds that of parental cell populations or C+E+M group when

co-cultured with serum depleted NRCMs.

Several mRNAs associated with lineage specification were analyzed following

co-culture of CardioClusters or parental cell populations with NRCMs. *GATA4* showed

the highest expression in cCIC co-culture. Predictably, EPCs displayed the largest

induction of endothelial marker *CD31*, whereas MSCs induced *SMA* gene expression

after 5 days of co-culture with NRCMs (Supplemental Figure 8, F-H). Neither *CD31* nor

SMA were significantly upregulated in CardioCluster group (Supplemental Figure 8G

and 8H).

CardioClusters are resistant to oxidative stress by *in vitro* assay

CardioClusters were substantially more resistant to cell death induced by 4 hours of

H_2O_2 treatment after overnight low serum culture (Figure 4). Dying cells were divided

into groups of early apoptosis, late apoptosis, and necrosis based on Annexin V and

Sytox Blue staining (Supplemental Figure 9). CardioClusters showed significantly fewer

cells in necrosis relative to any individual parental population (cCICs, EPCs, MSCs)

(p<0.05 versus cCIC; Figure 4B and Supplemental Figure 9F) as visually evidenced by

fewer cells rounding up and detaching from the tissue culture dish (transparent arrows;

Figure 4C and 4D). These results demonstrate superiority of CardioClusters to survive

oxidative stress challenge *in vitro*.

CardioClusters improve myocardial structure and function following infarction

injury

Therapeutic efficacy of CardioClusters was assessed in a murine experimental

myocardial infarction injury model of permanent coronary artery occlusion. Xenogenic

human cell treatment into NOD^{SCID} recipient mice was performed at the time of

infarction with direct comparison between CardioClusters and the C+E+M combined

population group administered as a single cell suspension mixture. Myocardial structure

and function were assessed by parasternal long axis echocardiography for four

experimental groups: non-injured sham, CardioCluster, C+E+M, and vehicle-treated

(Figure 5A). All groups had comparable reduction in cardiac function at 1 week post

injection (wpi) demonstrating consistency of infarction injury (Figure 5, B-D, F-G and

Supplemental Table 6) with average EF for all infarcted groups of approximately 30%

(CardioCluster, 27±2.9%; C+E+M, 32±2.2%; Vehicle, 29±2.0%; Supplemental Figure

10). The CardioCluster-treated group showed significant cardiac functional improvement

starting 4 wpi, which was sustained throughout the 20-week time course, with increased

fractional shortening (FS; Figure 5B) and ejection fraction (EF; Figure 5C) versus

C+E+M treatment 4 and 8 wpi and was significant at study completion for EF. In

comparison, EF and FS improvements in the C+E+M treated group only began to

appear at 12 and 16 wpi, respectively (Figure 5B and 5C). Terminal EF measurements

at 20 wpi show EF value is highest in the CardioCluster group relative to vehicle only or

C+E+M groups (40±1.9% versus 16±1.0% or 30±4.5%, respectively; Supplemental

Figure 10, B-D). CardioCluster treatment shows a 45±7% improvement in EF at study

completion relative to the starting value at week 1. In contrast, EF at study completion

relative to week 1 for the C+E+M or vehicle only-treated groups decreased by 5±14%

and 46±3%, respectively (Figure 5E). Furthermore, CardioCluster treatment group

exhibits significantly smaller left ventricular internal diameter both in systole (LVID;s)

and diastole (LVID;d), as well as reduction in LV end systolic and diastolic volumes (LV

Vol;s and LV Vol;d). Heart rate was not significantly different among treatment groups

(Figure 5, F-G and Supplemental Figure 11, A-C). Structural and functional data are

detailed in Supplemental Table 6.

Speckle-tracking based strain analysis is a highly sensitive echocardiographic

technique for assessing left ventricular (LV) function[66, 67]. LV function was similarly

reduced in all infarcted mice at 1 wpi (Figure 6A). Progressive changes consistent with

adverse ventricular remodeling occur in vehicle-treated animals in agreement with

conventional echocardiographic measures of function (Figure 5). LV systolic

deformation in the CardioCluster-treated group showed significant improvement starting

at 8 wpi and progressing through 20 wpi compared to vehicle-treated animals (p<0.001;

Figure 6, B-D). Radial strain measurements at 8 and 20 wpi confirmed significant

functional benefit provided by CardioCluster treatment versus 2D cultured parental

C+E+M mixed population (p<0.01; Figure 6C). Peak longitudinal strain was improved in

the C+E+M treated group versus vehicle alone (p<0.05; Figure 6D). Regional strain

measurements assessing the area of injury further demonstrate significant improvement

in LV function for CardioCluster-treated animals (Figure 6E and 6F). In the area of

injury, the absolute difference in radial strain (week 1-to-week 20) for CardioCluster-

treated group was 7.59±1.28% which was significantly improved relative to C+E+M and

vehicle-treated groups (-1.56±1.96% and -1.88±1.61% respectively, p<0.01 versus

CardioCluster group, Figure 6F). Similar improvement was seen for absolute difference

in longitudinal strain 20 wpi.

　　　　CardioCluster superiority for restoring myocardial structure and function relative

to the mixed population C+E+M is further reinforced by tissue morphometry and

hemodynamic measurements. Cardiac hypertrophy was not a contributing factor to

increasing anterior wall thickness (AWT) at 20 weeks in the CardioCluster-treated group

(Figure 5D) as heart weight to tibia length ratios did not increase (HW/TL; Figure 5H)

relative to the sham-operated control. In contrast, a significant increase in HW/TL is

present in both vehicle control as well as C+E+M treatment groups. Fibrotic area is

significantly smaller in CardioCluster-treated mice compared to vehicle at 20 wpi

(38.4±4.5% of CardioCluster LV versus 55.3±4.3% of vehicle LV; Figure 5I-M) although

infarct size is not significantly different between hearts receiving C+E+M or

CardioClusters. Invasive hemodynamic measurement validates functional superiority of

the CardioCluster-treated group showing significantly improved developed pressure

1 over time (dP/dT) versus vehicle (Supplemental Figure 11D), in addition to increasing

2 left ventricular developed pressure (LVDP) and P_{max}-P_{min} (Supplemental Figure 11E).

3 Collectively, these findings are evidence that CardioClusters offer significantly greater

4 benefit for restoration of myocardial performance in this murine myocardial infarction

5 injury model.

7 CardioClusters engraft and persist in the myocardial wall following

8 intramyocardial injection

Characteristics of CardioClusters including multicellular 3D architecture and

enhanced survival are attractive features to mediate increased persistence following

delivery compared to dissociated single cell suspensions such as the C+E+M mixed

population. CardioCluster persistence *in vivo* was longitudinally assessed over a 20-

week period by confocal microscopy (Figure 7). CardioCluster localization was tracked

with co-injection of FluoSpheres tracking beads in pilot studies to confirm delivery

location in tissue sections (Figure 7A). Cryosectioned hearts allowed for direct

visualization of fluorophore tags without antibody labeling. All three constituent cell

types were readily visualized in the myocardial wall, with MSCs at the center of the

CardioCluster surrounded by a layer of cCICs and EPCs, similar to architecture

observed *in vitro* (Figure 1). With CardioCluster localization confirmed coincident with

the injection site, subsequent injections and imaging were performed without

FluoSpheres tracking beads for long-term functional studies. CardioClusters were

clearly visible within the myocardium at serial time points: 1 day, 3 days, 1-, 4-, 12-, and

20- weeks post injection (Figure 7, B-H). Antibody labeling confirmed CardioCluster

persistence up to 20 weeks post injection (Figure 7E-H).

CardioClusters increase capillary density in the infarct area

Capillary density was measured in the infarct, border zone, and remote regions

at 20 wpi. Non-injured controls (sham) serve as the control group compared to injured

hearts (Figure 8). Notably, the CardioCluster group exhibited significantly more isolectin

labeled vessels in the infarct region at 20 wpi versus both vehicle and C+E+M-treated

groups (Figure 8, A and E–G). CardioCluster group capillary density in the infarct region

increased 62% or 83% versus the C+E+M or vehicle only control, respectively. Within

the infarct border zone at 20 wpi, both CardioCluster and C+E+M-treated groups

trended toward increased capillary density versus vehicle control, but not achieving

significance (Figure 8B). The remote region did not significantly increase capillary

density in either CardioCluster or C+E+M groups relative to vehicle or sham (Figure 8C

and 8D). Taken together, these data demonstrate a superior level of

microvascularization prompted by CardioCluster treatment relative to dissociated mixed

cell preparation or vehicle only control groups.

CardioCluster treatment antagonizes cardiomyocyte hypertrophy in the border

and remote regions and preserves cardiomyocyte size in the infarct region

Cell therapy reduces hypertrophic remodeling following pathologic injury to blunt

progression of heart failure after MI. Individual cardiomyocyte cross-sectional area was

traced (Supplemental Figure 12, A-C) along with average cardiomyocyte cross-sectional

area for infarct, border zone, and remote regions (Supplemental Figure 12, D-F).

Treatment groups receiving either CardioClusters or the C+E+M mixed population both

exhibited normalized cardiomyocyte size in the infarct region nearly identical to

uninjured sham control hearts at 20 wpi (Figure 9A). Cardiomyocytes within the border

zone proximal to infarction or in remote regions from the injury site were significantly

smaller in the CardioCluster treatment group compared to either C+E+M or vehicle only

control groups (p<0.001; Figure 9, B-C and E-G). Indeed, cardiomyocyte size in remote

regions was normalized in the CardioCluster group to values similar with non-injured

controls (Figure 9C and 9D). Collectively these data validate the action of CardioCluster

treatment to blunt hypertrophic cellular enlargement better than dissociated mixed cell

preparation or vehicle only control groups.

Discussion

The preceding decade of cardiac cell therapy has produced substantial

knowledge regarding optimization as well as limitations of current therapeutic

interventions. In a field sometimes overshadowed by contentious debate[68-72], it is

important to remember that there are also many points of consensus. Specifically, all

parties agree that inefficient cell delivery to the site of injury, low cell retention and

modest efficacy of cells that do remain within the tissue are factors hampering

advancement. Technical reinvention building upon prior success by incorporating 'next

generation' approaches to surmount these established barriers represent the frontier of

cell therapy research. CardioClusters introduced in this report are a novel and effective

solution that integrates multiple cardiac-resident cell types into a single injectable

product. The evolutionary advance offered by CardioClusters is mitigation of single cell

delivery challenges through spherical self-assembly of a larger 3D structure, which

provides enhanced retention to mediate repair after delivery. Inspiration for

CardioClusters is drawn from prior studies showing superiority of combinatorial cell

therapy[19-22] and prolonged survival and functional engraftment of cells grown in 3D

environments[73-76] or tissue-like constructs [27]. The multicellular structure and composition

of CardioClusters represents a distinctly unique *in vitro* engineered platform to enhance

the outcome of cellular therapeutics as demonstrated in preclinical testing using an

established murine infarction injury model treated with a xenograft of human cardiac-

derived cells.

CardioClusters were deliberately designed with multiple features anticipated to

enhance efficacy. Among these properties, the essential combination of multiple cardiac

cell types was enabled by our prior methodological studies to isolate and expand three

distinct cardiac-derived interstitial cell types from patients with end-stage heart failure

undergoing implantation of a left ventricular assist device (LVAD)[39]. End stage heart

failure patients such as LVAD recipients represent likely candidates for interventional

autologous therapy using cells derived from their own cardiac tissue. Combinatorial

approaches harnessing beneficial attributes of multiple adult cell types are gaining

acceptance as a method to enhance biological properties and efficacy based upon the

tenets that: 1) no single cell population possesses all the requisite attributes for effective

repair, and 2) both cardiomyogenic and non-cardiomyogenic cells contribute to

myocardial repair and regeneration. Combining multiple cell types with complementary

roles more efficiently mediates repair in preclinical experimental animal models of heart

failure[19-22] and is currently being assessed in the CONCERT clinical trial with patients

receiving mixtures of MSCs and cardiac progenitor cells (cCIC)[77]. Similarly, induced

pluripotent stem cell-derived cardiomyocytes combined with vascular cells[78] or MSCs[79]

potentiates myocardial repair, likely due to enhanced stimulation of endogenous repair

mechanisms. Efficient isolation and expansion of three distinct cardiac-resident non-

myocyte populations brought together *ex vivo* to form CardioClusters is now technically

feasible (Figure 1, Supplemental Figures 1 and 3) [39]. CardioCluster biological variability

depending upon the source, condition, and pathologic state of donor tissue is an

important and intriguing unresolved issue to be addressed in follow-up studies based

upon the proof-of-principle provided in the present report.

Another enabling feature of CardioClusters is the profound influence of

aggregation upon phenotypic and biological properties of the constituent cell

populations. Specifically, CardioClusters foster a transcriptional profile more consistent

with freshly isolated cardiac interstitial cells compared to their monolayer counterparts

(Figure 2). Even relatively short-term *in vitro* expansion of cCIC in 2D monolayer culture

results in loss of identity marker gene expression and decreased population

heterogeneity by single cell RNA-Seq transcriptome profiling[25]. And although cells

derived from typical 2D monolayer cultures are used to seed CardioClusters, the

transcriptome signature of CardioCluster cells collectively resemble each other far more

than original parental cells. The CardioCluster microenvironment promotes intercellular

coordination initiated within a 3D environment, unlike traditional monolayer expansion[23-

25]. Increased expression of collagen type I and III, integrins (ITGA2, ITGB1, ITGA11,

ITGA1, ITGAV), fibronectin, and matrix remodeling enzymes (MMP-1, MMP-2, MMP-14,

TIMP-1, TIMP-2) in CardioClusters is consistent with enhanced matrix remodeling

capacity of cells maintained in a 3D microenvironment[80]. Transcript data for

CardioClusters also showed elevated expression of Notch3 that exerts an important

regulatory role in the contexts of development and tissue regeneration for maintenance

of a progenitor pool and tissue homeostasis[81]. Elevated Notch expression and

superiority of 3D aggregation culture relative to conventional 2D conditions is consistent

with results using pediatric cCICs cultured in 3D spheres of approximately 1500 cells,

wherein 3D aggregated cCICs exert enhanced repair with increased notch signaling

compared with their 2D counterparts in a right ventricular heart failure model.[50]

Restoring fresh and/or youthful characteristics to isolated cells expanded *in vitro*[82-85]

may be one way that CardioClusters provide functional benefits to the collective

population.

A third enabling feature of CardioClusters is cardioprotective action, particularly

under conditions of environmental stress. *In vitro* testing in co-culture assays is an

established protocol to assess the potential of candidate cell types to inhibit

cardiomyocyte death from pro-apoptotic challenge[57, 86]. Superior ability of

CardioClusters to blunt NRCM death relative to single parental cell types supports the

rationale for culturing the mixed cell population together in a 3D configuration (Figure 3).

Animal studies have demonstrated that transplantation of exogenous cells exert

cardioprotection through release of paracrine factors and, to a lesser extent, trans-

differentiation into cardiac resident cells[79, 87-93]. This idea is supported by our

transcriptomic and quantitative PCR analysis revealing the ability of cardiac interstitial

cells to express a wide array of paracrine cytokines and factors that promote

angiogenesis (such as VE-statin and RAC2) and cell survival (such as FGF, SDF-1,

IGF, IL-6 and HGF). In vitro functional assays revealed that cardiac interstitial cells

exerted prosurvival and proangiogenic effects. Furthermore, the secretome of

CardioClusters also exerted protective effects on neonatal rat cardiomyocytes.

Importantly, the paracrine effects of CardioClusters were more powerful than those

shown with the single parental populations. These findings establish the justification for

subsequent *in vivo* testing and a potential paracrine mechanistic basis for CardioCluster

action.

Superior restoration of structure and function following cardiomyopathic injury is

evident from comparative testing with either CardioClusters or the dissociated cell

mixture of C+E+M in xenogenic treatment of NOD^{SCID} mice (Figure 5). Empirical control

of CardioCluster size to <160 μm (Figure 1N and 1O) allowed for injection through a 30-

gauge needle without dissociation into single cells. Improvement in FS and EF was

observed starting at week 4 and maintained during the entirety of the 20-week study,

with CardioCluster-treated animals showing significantly improved myocardial wall

structure compared to C+E+M-treated animals concomitant with increased capillary

density (Figure 8) and preserved cardiomyocyte size (Figure 9). CardioClusters

persisted within the myocardial wall, with cells clearly visible up to 5 months post-

injection (Figure 7). The use of human cells necessitated the use of NOD^{SCID} mice in our

study. Although having a fully functioning immune system may lead to a different

outcome, in view of what has been seen in previous studies the results should improve

when the cells have been introduced into a model that has a working immune system.

Future studies will be needed to address this point.

Meta-analysis examining cardiac stem cell (CSC) and MSC ability to treat MI in

animal models found that treatment culminated in an absolute difference in EF ranging

from 8-10.7% compared to control animals[94, 95]. In comparison, by 20 weeks

CardioCluster treatment showed a 24.2% increase in EF compared with vehicle

treatment. Our data shows the significant 2-fold improvement possible with

CardioClusters versus traditional single cell therapy approaches. "Off-the-shelf"

potential of CardioClusters preserved in liquid nitrogen demonstrated high viability and

structural integrity indistinguishable from non-frozen counterparts (Supplemental Figure

4). Frozen/thawed CardioCluster efficacy remains to be tested *in vivo*, but the ability to

mass-produce and preserve CardioClusters in frozen storage is attractive for clinical

implementation planning.

The conceptual framework of CardioClusters offers almost infinite possibilities for

modification and optimization for therapeutic use, as well as basic investigation of

cellular interactions. For example, parameters including cell ratios, cell types, cluster

size, and number of CardioClusters to inject are all worthy of further consideration.

CardioCluster contain approximately 300 cells crucial for injectability through the inner

diameter of a 30-gauge needle, which constrained diameter to <160 μm (Figure 1O) in

murine studies. However, a larger animal could tolerate a larger gauge needle and

concomitantly scaled up CardioCluster size for a greater total number of cells to be

injected. With the advent of ultra-low attachment microcavity plates with between 79 –

15,000+ wells per plate, the feasibility of scaling up production is possible for future

clinical application and warrants further investigation. Alternatively, 'mini-CardioClusters'

of 50-100 cells with smaller diameter would allow a greater number of individual clusters

to be injected. Ability to fine-tune CardioCluster size is a benefit distinct from traditional

3D cell aggregates such as cardiospheres where diameter is not controllable,

necessitating dissociation into single cell suspensions of cardiosphere-derived cells for

clinical use. Additional strategies to further enhance the CardioCluster concept could

involve incorporating genetically modified cells with pro-survival factors such as Pim-1[83,]

[96] or overexpression of chemokine receptor (CCR1) to enhance migration, survival and

engraftment[97]. Likewise, culture condition modification using hypoxia to favor cell growth

and blunt senescence-associated characteristics[98] could dramatically alter

CardioCluster biological properties. Cellular interactions occurring within 3D

environments can also be tested, such as seeding the three C+E+M founder cell types

together rather than sequentially, which appears to create a hollow CardioCluster (data

not shown, Sussman lab). This may be attributable to EPC/MSC interaction allowing for

internal cavity formation as seen during organ and tissue development[99, 100] or may be

more similar to pericyte/MSC-like endothelial cell interactions[101]. With knowledge

regarding combinatorial cell therapy at a rudimentary level [102-104] the 'next generation'

CardioCluster approach will benefit from further investigation given the multiple

possibilities for tweaking the system to enhance the outcome.

As with any novel technological approach, there are unknowns and limitations

that need to be resolved for CardioCluster development. A benefit of CardioCluster

design is the quick formation time of only 48 hours from start to finish, however creation

of CardioClusters necessitates that the multiple composite founder cell types must be

ready for utilization in sequence within a short time frame. Since CardioClusters could

be conceived using a plethora of possible cells, the time required for expansion of the

parental cells may differ depending on the cell types chosen, particularly if using cells

isolated from aged patients suffering from cardiomyopathic disease. Allogeneic

implementation for CardioClusters might incorporate immunosuppressive agents or

assembly using 'universal' donor cells engineered by CRISPR-mediated genome

editing[105] . Prospective preparation and freeze storage of either parental cells or

CardioClusters will help ease issues with timing for assembly and delivery. With respect

to delivery, CardioClusters may present a safety concerns as microemboli if

administered intravenously, so direct intramyocardial injection will be used, which would

be the preferred approach regardless to enhance efficacy[106, 107].

This study presents the debut of CardioClusters as a novel technical 'next

generation' approach to improve upon established protocols using dissociated single

cell preparations proven to be safe for administration to patients but of limited efficacy.

Unlike other tissue engineering microfabrication approaches, the spontaneously formed

CardioCluster 3D structure maximizes cellular interaction and allows for defined cell

ratios, controlled size, and facilitates injectability without dissociation. This combination

of features makes CardioClusters unique among current cell therapeutic approaches

with demonstrated superiority over single cell mixed suspensions in mitigation of

myocardial infarction damage. This initial step toward enhanced cell therapy provides a

readily manipulatable platform that will benefit from further research development with

the goal of potentiating cell-based therapeutic efficacy to mediate myocardial repair.

Acknowledgments

M. Monsanto and M. Sussman designed experiments. M. Monsanto, K. White, B.

Wang, Z. Ehrenberg, R. Alvarez, O. Echeagaray, S. Sengphanith, A. Muliono, N. Gude,

and K. Fisher performed experiments and analyzed data. M. Monsanto and M.

Sussman wrote the article. All authors read and approved the final article.

Sources of Funding

M. Monsanto is supported by NIH grant R01HL122525, Rees-Stealy Research

Foundation Phillips Gausewitz, M.D., Scholars of the SDSU Heart Institute,

Achievement Rewards for College Scientists (ARCS), and Inamori Fellowship. M.A.

Sussman is supported by NIH grants: R01HL067245, R37HL091102, R01HL105759,

R01HL113647, R01HL117163, P01HL085577, and R01HL122525, as well as an award

from the Fondation Leducq.

Disclosures

M. Sussman is Chief Scientific Officer and a founding member of CardioCreate

Inc.

References

1. Assmus, B. *et al.* Transplantation of Progenitor Cells and Regeneration Enhancement in Acute Myocardial Infarction (TOPCARE-AMI). *Circulation* **106**, 3009-3017 (2002).
2. Hare, J.M. *et al.* Comparison of allogeneic vs autologous bone marrow-derived mesenchymal stem cells delivered by transendocardial injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA* **308**, 2369-2379 (2012).
3. Makkar, R.R. *et al.* Intracoronary cardiosphere-derived cells for heart regeneration after myocardial infarction (CADUCEUS): a prospective, randomised phase 1 trial. *Lancet* **379**, 895-904 (2012).
4. Menasche, P. *et al.* The Myoblast Autologous Grafting in Ischemic Cardiomyopathy (MAGIC) trial: first randomized placebo-controlled study of myoblast transplantation. *Circulation* **117**, 1189-1200 (2008).
5. Mills, J.S. & Rao, S.V. REPAIR-AMI: stem cells for acute myocardial infarction. *Future Cardiol* **3**, 137-140 (2007).
6. Traverse, J.H. *et al.* Effect of intracoronary delivery of autologous bone marrow mononuclear cells 2 to 3 weeks following acute myocardial infarction on left ventricular function: the LateTIME randomized trial. *JAMA* **306**, 2110-2119 (2011).
7. Perin, E.C. *et al.* Transendocardial, autologous bone marrow cell transplantation for severe, chronic ischemic heart failure. *Circulation* **107**, 2294-2302 (2003).
8. Bolli, R. *et al.* Intracoronary delivery of autologous cardiac stem cells improves cardiac function in a porcine model of chronic ischemic cardiomyopathy. *Circulation* **128**, 122-131 (2013).
9. Lee, S.T. *et al.* Intramyocardial injection of autologous cardiospheres or cardiosphere-derived cells preserves function and minimizes adverse ventricular remodeling in pigs with heart failure post-myocardial infarction. *J Am Coll Cardiol* **57**, 455-465 (2011).
10. Sanganalmath, S.K. & Bolli, R. Cell therapy for heart failure: a comprehensive overview of experimental and clinical studies, current challenges, and future directions. *Circ Res* **113**, 810-834 (2013).
11. Higuchi, A. *et al.* Stem cell therapies for myocardial infarction in clinical trials: bioengineering and biomaterial aspects. *Lab Invest* **97**, 1167-1179 (2017).
12. Zhu, K. *et al.* Lack of Remuscularization Following Transplantation of Human Embryonic Stem Cell-Derived Cardiovascular Progenitor Cells in Infarcted Nonhuman Primates. *Circ Res* **122**, 958-969 (2018).
13. Fujita, J. *et al.* Myocardial cell sheet therapy and cardiac function. *Am J Physiol Heart Circ Physiol* **303**, H1169-1182.
14. Cheng, K. *et al.* Functional performance of human cardiosphere-derived cells delivered in an in situ polymerizable hyaluronan-gelatin hydrogel. *Biomaterials* **33**, 5317-5324.
15. Beohar, N., Rapp, J., Pandya, S. & Losordo, D.W. Rebuilding the damaged heart: the potential of cytokines and growth factors in the treatment of ischemic heart disease. *J Am Coll Cardiol* **56**, 1287-1297.
16. Tokita, Y. *et al.* Repeated Administrations of Cardiac Progenitor Cells Are Markedly More Effective Than a Single Administration: A New Paradigm in Cell Therapy. *Circ Res* **119**, 635-651 (2016).
17. Fischer, K.M. *et al.* Enhancement of myocardial regeneration through genetic engineering of cardiac progenitor cells expressing Pim-1 kinase. *Circulation* **120**, 2077-2087 (2009).
18. Mohsin, S. *et al.* Human Cardiac Progenitor Cells Engineered With Pim-I Kinase Enhance Myocardial Repair. *J Am Coll Cardiol*.
19. Williams, A.R. *et al.* Enhanced effect of combining human cardiac stem cells and bone marrow mesenchymal stem cells to reduce infarct size and to restore cardiac function after myocardial infarction. *Circulation* **127**, 213-223.

20. Karantalis, V. *et al.* Synergistic Effects of Combined Cell Therapy for Chronic Ischemic Cardiomyopathy. *J Am Coll Cardiol* **66**, 1990-1999 (2015).

21. Latham, N. *et al.* Human blood and cardiac stem cells synergize to enhance cardiac repair when cotransplanted into ischemic myocardium. *Circulation* **128**, S105-112 (2013).

22. Avolio, E. *et al.* Combined intramyocardial delivery of human pericytes and cardiac stem cells additively improves the healing of mouse infarcted hearts through stimulation of vascular and muscular repair. *Circ Res* **116**, e81-94 (2015).

23. Tibbitt, M.W. & Anseth, K.S. Hydrogels as extracellular matrix mimics for 3D cell culture. *Biotechnol Bioeng* **103**, 655-663 (2009).

24. Gupta, N. *et al.* Microfluidics-based 3D cell culture models: Utility in novel drug discovery and delivery research. *Bioeng Transl Med* **1**, 63-81 (2016).

25. Kim, T. *et al.* In situ transcriptome characteristics are lost following culture adaptation of adult cardiac stem cells. *Sci Rep* **8**, 12060 (2018).

26. Zimmermann, W.H. *et al.* Engineered heart tissue grafts improve systolic and diastolic function in infarcted rat hearts. *Nat Med* **12**, 452-458 (2006).

27. Li, J. *et al.* Human Pluripotent Stem Cell-Derived Cardiac Tissue-like Constructs for Repairing the Infarcted Myocardium. *Stem Cell Reports* **9**, 1546-1559 (2017).

28. Voges, H.K. *et al.* Development of a human cardiac organoid injury model reveals innate regenerative potential. *Development* **144**, 1118-1127 (2017).

29. Nugraha, B., Buono, M.F., von Boehmer, L., Hoerstrup, S.P. & Emmert, M.Y. Human Cardiac Organoids for Disease Modeling. *Clin Pharmacol Ther* **105**, 79-85 (2019).

30. Ong, C.S. *et al.* 3D and 4D Bioprinting of the Myocardium: Current Approaches, Challenges, and Future Prospects. *Biomed Res Int* **2018**, 6497242 (2018).

31. Noor, N. *et al.* 3D Printing of Personalized Thick and Perfusable Cardiac Patches and Hearts. *Adv Sci (Weinh)* **6**, 1900344 (2019).

32. Huh, D., Hamilton, G.A. & Ingber, D.E. From 3D cell culture to organs-on-chips. *Trends Cell Biol* **21**, 745-754 (2011).

33. Messina, E. *et al.* Isolation and expansion of adult cardiac stem cells from human and murine heart. *Circ Res* **95**, 911-921 (2004).

34. Chimenti, I. *et al.* Relative roles of direct regeneration versus paracrine effects of human cardiosphere-derived cells transplanted into infarcted mice. *Circ Res* **106**, 971-980.

35. Cho, H.J. *et al.* Secondary sphere formation enhances the functionality of cardiac progenitor cells. *Mol Ther* **20**, 1750-1766.

36. Smith, R.R. *et al.* Regenerative potential of cardiosphere-derived cells expanded from percutaneous endomyocardial biopsy specimens. *Circulation* **115**, 896-908 (2007).

37. Leri, A., Kajstura, J. & Anversa, P. Role of cardiac stem cells in cardiac pathophysiology: a paradigm shift in human myocardial biology. *Circ Res* **109**, 941-961.

38. Lee, S.T. *et al.* Intramyocardial injection of autologous cardiospheres or cardiosphere-derived cells preserves function and minimizes adverse ventricular remodeling in pigs with heart failure post-myocardial infarction. *J Am Coll Cardiol* **57**, 455-465.

39. Monsanto, M.M. *et al.* Concurrent Isolation of 3 Distinct Cardiac Stem Cell Populations From a Single Human Heart Biopsy. *Circ Res* **121**, 113-124 (2017).

40. Bernstein, H.S. & Srivastava, D. Stem cell therapy for cardiac disease. *Pediatr Res* **71**, 491-499.

41. Gnecchi, M., Zhang, Z., Ni, A. & Dzau, V.J. Paracrine mechanisms in adult stem cell signaling and therapy. *Circ Res* **103**, 1204-1219 (2008).

42. Wagner, W. *et al.* Molecular and secretory profiles of human mesenchymal stromal cells and their abilities to maintain primitive hematopoietic progenitors. *Stem Cells* **25**, 2638-2647 (2007).

43. Jujo, K., Ii, M. & Losordo, D.W. Endothelial progenitor cells in neovascularization of infarcted myocardium. *J Mol Cell Cardiol* **45**, 530-544 (2008).

44. Thal, M.A. *et al.* Enhanced angiogenic and cardiomyocyte differentiation capacity of epigenetically reprogrammed mouse and human endothelial progenitor cells augments their efficacy for ischemic myocardial repair. *Circ Res* **111**, 180-190.

45. Lee, W.Y. *et al.* Vascularization and restoration of heart function in rat myocardial infarction using transplantation of human cbMSC/HUVEC core-shell bodies. *Biomaterials* **33**, 2127-2136.

46. Saleh, F.A., Whyte, M., Ashton, P. & Genever, P.G. Regulation of mesenchymal stem cell activity by endothelial cells. *Stem Cells Dev* **20**, 391-403 (2011).

47. Macosko, E.Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

48. Yu, G., Wang, L.G., Han, Y. & He, Q.Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284-287 (2012).

49. Gao, E. *et al.* A novel and efficient model of coronary artery ligation and myocardial infarction in the mouse. *Circ Res* **107**, 1445-1453 (2010).

50. Trac, D., Maxwell, J.T., Brown, M.E., Xu, C. & Davis, M.E. Aggregation of Child Cardiac Progenitor Cells Into Spheres Activates Notch Signaling and Improves Treatment of Right Ventricular Heart Failure. *Circ Res* **124**, 526-538 (2019).

51. Nichols, M.G. & Foster, T.H. Oxygen diffusion and reaction kinetics in the photodynamic therapy of multicell tumour spheroids. *Phys Med Biol* **39**, 2161-2181 (1994).

52. Wu, J., Rostami, M.R., Cadavid Olaya, D.P. & Tzanakakis, E.S. Oxygen transport and stem cell aggregation in stirred-suspension bioreactor cultures. *PLoS One* **9**, e102486 (2014).

53. Tang, Y.L. *et al.* Hypoxic preconditioning enhances the benefit of cardiac progenitor cell therapy for treatment of myocardial infarction by inducing CXCR4 expression. *Circ Res* **104**, 1209-1216 (2009).

54. Yan, F. *et al.* Hypoxic preconditioning improves survival of cardiac progenitor cells: role of stromal cell derived factor-1alpha-CXCR4 axis. *PLoS One* **7**, e37948.

55. Nichol, D. & Stuhlmann, H. EGFL7: a unique angiogenic signaling factor in vascular development and disease. *Blood* **119**, 1345-1352 (2012).

56. Yu, J. *et al.* Topological Arrangement of Cardiac Fibroblasts Regulates Cellular Plasticity. *Circ Res* **123**, 73-85 (2018).

57. Quijada, P. *et al.* Cardiac Stem Cell Hybrids Enhance Myocardial Repair. *Circ Res* **117**, 695-706 (2015).

58. Zheng, H., Fu, G., Dai, T. & Huang, H. Migration of endothelial progenitor cells mediated by stromal cell-derived factor-1alpha/CXCR4 via PI3K/Akt/eNOS signal transduction pathway. *J Cardiovasc Pharmacol* **50**, 274-280 (2007).

59. Kanzler, I. *et al.* Differential roles of angiogenic chemokines in endothelial progenitor cell-induced angiogenesis. *Basic Res Cardiol* **108**, 310 (2013).

60. Hodgkinson, C.P., Bareja, A., Gomez, J.A. & Dzau, V.J. Emerging Concepts in Paracrine Mechanisms in Regenerative Cardiovascular Medicine and Biology. *Circ Res* **118**, 95-107 (2016).

61. Wang, M., Tsai, B., Brown, J.W. & Meldrum, D.R. Insulin-like growth factor-1 in myocardial tissue: interaction with tumor necrosis factor. *Crit Care* **7**, 417-419 (2003).

62. Kawaguchi, N. *et al.* c-kitpos GATA-4 high rat cardiac stem cells foster adult cardiomyocyte survival through IGF-1 paracrine signalling. *PLoS One* **5**, e14297 (2010).

63. Ellison, G.M. *et al.* Endogenous cardiac stem cell activation by insulin-like growth factor-1/hepatocyte growth factor intracoronary injection fosters survival and regeneration of the infarcted pig heart. *J Am Coll Cardiol* **58**, 977-986 (2011).

64. Fontes, J.A., Rose, N.R. & Cihakova, D. The varying faces of IL-6: From cardiac protection to cardiac failure. *Cytokine* **74**, 62-68 (2015).

65. Nakamura, T. & Mizuno, S. The discovery of hepatocyte growth factor (HGF) and its significance for cell biology, life sciences and clinical medicine. *Proc Jpn Acad Ser B Phys Biol Sci* **86**, 588-610 (2010).

66. Bauer, M. *et al.* Echocardiographic speckle-tracking based strain imaging for rapid cardiovascular phenotyping in mice. *Circ Res* **108**, 908-916 (2011).

67. de Lucia, C. *et al.* Echocardiographic Strain Analysis for the Early Detection of Left Ventricular Systolic/Diastolic Dysfunction and Dyssynchrony in a Mouse Model of Physiological Aging. *J Gerontol A Biol Sci Med Sci* **74**, 455-461 (2019).

68. Sussman, M.A. Cardiac nonmyocyte subpopulations: a secular congregation. *Regen Med* (2019).

69. Gude, N.A. & Sussman, M.A. Chasing c-Kit through the heart: Taking a broader view. *Pharmacol Res* **127**, 110-115 (2018).

70. Khan, M. & Koch, W.J. c-kit+ Cardiac Stem Cells: Spontaneous Creation or a Perplexing Reality. *Circ Res* **118**, 783-785 (2016).

71. Sultana, N. *et al.* Resident c-kit(+) cells in the heart are not cardiac stem cells. *Nat Commun* **6**, 8701 (2015).

72. van Berlo, J.H. *et al.* c-kit+ cells minimally contribute cardiomyocytes to the heart. *Nature* **509**, 337-341 (2014).

73. Le, L.V., Mkrtschjan, M.A., Russell, B. & Desai, T.A. Hang on tight: reprogramming the cell with microstructural cues. *Biomed Microdevices* **21**, 43 (2019).

74. Kobel, S. & Lutolf, M. High-throughput methods to define complex stem cell niches. *Biotechniques* **48**, ix-xxii.

75. Zuppinger, C. 3D culture for cardiac cells. *Biochim Biophys Acta* **1863**, 1873-1881 (2016).

76. Moon, S.H. *et al.* The use of aggregates of purified cardiomyocytes derived from human ESCs for functional engraftment after myocardial infarction. *Biomaterials* **34**, 4013-4026 (2013).

77. Bolli, R. *et al.* Rationale and Design of the CONCERT-HF Trial (Combination of Mesenchymal and c-kit(+) Cardiac Stem Cells As Regenerative Therapy for Heart Failure). *Circ Res* **122**, 1703-1715 (2018).

78. Masumoto, H. *et al.* Human iPS cell-engineered cardiac tissue sheets with cardiomyocytes and vascular cells for cardiac regeneration. *Sci Rep* **4**, 6716 (2014).

79. Yoshida, S. *et al.* Maturation of Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes by Soluble Factors from Human Mesenchymal Stem Cells. *Mol Ther* **26**, 2681-2695 (2018).

80. van Marion, M.H. *et al.* Behavior of CMPCs in unidirectional constrained and stress-free 3D hydrogels. *J Mol Cell Cardiol* **87**, 79-91 (2015).

81. Gude, N. *et al.* Notch activation enhances lineage commitment and protective signaling in cardiac progenitor cells. *Basic Res Cardiol* **110**, 29 (2015).

82. Hariharan, N. *et al.* Nucleostemin rejuvenates cardiac progenitor cells and antagonizes myocardial aging. *J Am Coll Cardiol* **65**, 133-147 (2015).

83. Mohsin, S. *et al.* Rejuvenation of human cardiac progenitor cells with Pim-1 kinase. *Circ Res* **113**, 1169-1179 (2013).

84. Li, S.H. *et al.* Young Bone-Marrow Sca-1(+) Stem Cells Rejuvenate the Aged Heart and Improve Function after Injury through PDGFRbeta-Akt pathway. *Sci Rep* **7**, 41756 (2017).

85. Li, S.H. *et al.* Reconstitution of aged bone marrow with young cells repopulates cardiac-resident bone marrow-derived progenitor cells and prevents cardiac dysfunction after a myocardial infarction. *Eur Heart J* **34**, 1157-1167 (2013).

86. Arisi, M.F. *et al.* Myocardial apoptosis and mesenchymal stem cells with acute exercise. *Physiol Rep* **5** (2017).

87. Ferrini, A., Stevens, M.M., Sattler, S. & Rosenthal, N. Toward Regeneration of the Heart: Bioengineering Strategies for Immunomodulation. *Front Cardiovasc Med* **6**, 26 (2019).

88.	Gomez, I., Duval, V. & Silvestre, J.S. Cardiomyocytes and Macrophages Discourse on the Method to Govern Cardiac Repair. *Front Cardiovasc Med* **5**, 134 (2018).

89.	Haider, H., Jiang, S., Idris, N.M. & Ashraf, M. IGF-1-overexpressing mesenchymal stem cells accelerate bone marrow stem cell mobilization via paracrine activation of SDF-1alpha/CXCR4 signaling to promote myocardial repair. *Circ Res* **103**, 1300-1308 (2008).

90.	Jackson, R. *et al.* Paracrine Engineering of Human Cardiac Stem Cells With Insulin-Like Growth Factor 1 Enhances Myocardial Repair. *J Am Heart Assoc* **4**, e002104 (2015).

91.	Mayfield, A.E. *et al.* Interleukin-6 Mediates Post-Infarct Repair by Cardiac Explant-Derived Stem Cells. *Theranostics* **7**, 4850-4861 (2017).

92.	Forte, G. *et al.* Human cardiac progenitor cell grafts as unrestricted source of supernumerary cardiac cells in healthy murine hearts. *Stem Cells* **29**, 2051-2061 (2011).

93.	Oskouei, B.N. *et al.* Increased potency of cardiac stem cells compared with bone marrow mesenchymal stem cells in cardiac repair. *Stem Cells Transl Med* **1**, 116-124 (2012).

94.	Jansen Of Lorkeers, S.J. *et al.* Similar effect of autologous and allogeneic cell therapy for ischemic heart disease: systematic review and meta-analysis of large animal studies. *Circ Res* **116**, 80-86 (2015).

95.	Zwetsloot, P.P. *et al.* Cardiac Stem Cell Treatment in Myocardial Infarction: A Systematic Review and Meta-Analysis of Preclinical Studies. *Circ Res* **118**, 1223-1232 (2016).

96.	Quijada, P. *et al.* Preservation of myocardial structure is enhanced by pim-1 engineering of bone marrow cells. *Circ Res* **111**, 77-86 (2012).

97.	Huang, J. *et al.* Genetic modification of mesenchymal stem cells overexpressing CCR1 increases cell viability, migration, engraftment, and capillary density in the injured myocardium. *Circ Res* **106**, 1753-1762 (2010).

98.	Korski, K.I. *et al.* Hypoxia Prevents Mitochondrial Dysfunction and Senescence in Human c-Kit(+) Cardiac Progenitor Cells. *Stem Cells* **37**, 555-567 (2019).

99.	Durdu, S. *et al.* Luminal signalling links cell communication to tissue architecture during organogenesis. *Nature* **515**, 120-124 (2014).

100.	Bedzhov, I. & Zernicka-Goetz, M. Self-organizing properties of mouse pluripotent cells initiate morphogenesis upon implantation. *Cell* **156**, 1032-1044 (2014).

101.	Chen, W.C. *et al.* Human myocardial pericytes: multipotent mesodermal precursors exhibiting cardiac specificity. *Stem Cells* **33**, 557-573 (2015).

102.	Lemcke, H., Voronina, N., Steinhoff, G. & David, R. Recent Progress in Stem Cell Modification for Cardiac Regeneration. *Stem Cells Int* **2018**, 1909346 (2018).

103.	Quijada, P. & Sussman, M.A. Making it stick: chasing the optimal stem cells for cardiac regeneration. *Expert Rev Cardiovasc Ther* **12**, 1275-1288 (2014).

104.	Hatzistergos, K.E. & Vedenko, A. Cardiac Cell Therapy 3.0: The Beginning of the End or the End of the Beginning? *Circ Res* **121**, 95-97 (2017).

105.	Hawksworth, J. *et al.* Enhancement of red blood cell transfusion compatibility using CRISPR-mediated erythroblast gene editing. *EMBO Mol Med* **10** (2018).

106.	Hayashi, M., Li, T.S., Ito, H., Mikamo, A. & Hamano, K. Comparison of intramyocardial and intravenous routes of delivering bone marrow cells for the treatment of ischemic heart disease: an experimental study. *Cell Transplant* **13**, 639-647 (2004).

107.	Dib, N., Khawaja, H., Varner, S., McCarthy, M. & Campbell, A. Cell therapy for cardiovascular disease: a comparison of methods of delivery. *J Cardiovasc Transl Res* **4**, 177-181 (2011).

A cCIC — BF — eGFP — 100 μm — 0 μm 50
B EPC — BF — mOrange — 100 μm — 0 μm 50
C MSC — BF — Neptune — 100 μm — 0 μm 50

D Area (a.u.): cCIC, EPC, MSC — *** ***
E Roundness: cCIC, EPC, MSC — *** *** ***
F L/W Ratio: cCIC, EPC, MSC — *** *** ***

G Proliferation — Fold Change — cCIC, EPC, MSC — Day 0, Day 1, Day 3, Day 5
H Apoptosis — Annexin V⁺ cells — cCIC, EPC, MSC — *** ***
I Necrosis — Sytox Blue⁺ cells — cCIC, EPC, MSC — **
J Healthy — Live cells — cCIC, EPC, MSC — *** *** *

K Parental Cell Populations — EPC, cCIC, MSC — Day 1 cCIC/MSC Inner Core — Day 2 Addition of EPC — Day 3 CardioCluster Formed
L CardioCluster — cCIC, EPC, MSC — 0 μm 75
M 3D Cross Section — cCIC, EPC, MSC — 0 μm 75

N 100 cells, 200 cells, 400 cells, 600 cells, 800 cells, 1000 cells — Day 1, Day 3, Day 7
O CardioCluster Average Size — Diameter (um) — Cell Number — I.D. of 30 Gauge Needle

Figure 1. Three distinct cardiac cell types used to generate CardioClusters

A-C, Representative brightfield (BF) and immunofluorescent images for cCIC (eGFP$^+$) **(A)**, EPC (mOrange$^+$) **(B)** and MSC (Neptune$^+$) **(C)**. Scale bars: brightfield, 100 µm; immunofluorescent, 50 µm. DAPI to visualize nuclei (white). **D-F**, Cell morphometric parameters measuring area (a.u.: arbitrary units; **D**), roundness **(E)**, and length-to-width (L/W) ratio (**F;** n=3 human heart isolations; minimum of 30 cells traced per cell type, per heart). **G**, Cardiac cell proliferation measured using CyQuant assays at day 0, day 1, day 3, and day 5 (n=3). **H-J**, Annexin V/Sytox Blue labeling for apoptotic **(H)**, necrotic **(I)** and healthy **(J)** cells following cell death assay: 24 hours of low serum (75% serum reduction) followed by 4 hours treatment with 30 µM H_2O_2 in low serum medium (n=4). Data are presented as 1-way ANOVA, *p<0.05, **p<0.01, ***p<0.001. **K**, Schematic showing CardioCluster formation using three cell populations isolated from human heart tissue: MSC (red), cCIC (green), EPC (blue). **L**, Live CardioCluster visualized by endogenous fluorescent tags showing cCIC (eGFP; FITC channel; green), MSC (Neptune; APC channel; red) and EPC (mOrange; PE channel; blue). Scale bar, 75 µm. **M**, 3D cross section of a CardioCluster lightly fixed in 4% paraformaldehyde and stained with 4′,6-diamidino-2-phenylindole (DAPI; white) to visualize nuclei and cells exhibiting fluorescent tags without the need for antibody labeling. Scale bar, 75 µm. **N**, Representative brightfield images of CardioClusters ranging in size from 100-1000 cells cultured over a 7-day period. Scale bar, 100 µm. **O**, Determination of cell number enabling CardioClusters to pass through a 30-gauge needle, which has an inner diameter (I.D.) of 159 µm. Graph plot of CardioCluster diameters averaged over 3 days (n=4-7).

Figure 2. CardioCluster single cell RNA-Seq reveals transcriptional profile acquisition with increased similarity to freshly isolated cardiac interstitial cells

A-F, Transcriptional profiling on CardioClusters and parental cells using scRNA-seq. **A**, t-SNE map showing cells grown within a 3D CardioCluster (orange) predominately cluster together, while cCICs (green), EPCs (blue), and MSCs (red) primarily cluster into their own individual groups. **B**, Heatmap of differentially expressed genes (DEGs) among CardioClusters and parental cells. Selected DEGs for each group are color-coded and shown on the right. **C**, Violin plots of expression distribution for selected DEGs. **D**, Gene Ontology (GO) term analysis of molecular functions that are enriched based on DEGs. **E-F**, DEGs compared to genes expressed by freshly isolated mouse cardiac interstitial cells (mFreshCIC) represented in a Venn diagram **(E)** and heatmap of the gene set coexpressed by freshly isolated cells in comparison to CardioCluster and parental cell populations **(F)**. hParental indicates human parental cells; hCardioCluster, human CardioCluster cells; BAMBI, BMP and activin membrane bound inhibitor; LIF, leukemia inhibitory factor; CXCL12, C-X-C motif chemokine 12 or stromal cell-derived factor 1 (SDF1); ACTA2, smooth muscle alpha (α)-2 actin; C. Cluster, CardioCluster.

Figure 3. CardioClusters protect cardiomyocytes against low serum conditions *in vitro*
A, Timeline for neonatal rat cardiomyocyte (NRCM) low serum assay over a 4-day (D) period. **B**, Representative images of NRCM conditions: serum starved (48 hours of 0.5% serum), rescued (24 hours of 0.5% serum with 10% serum added for additional 24 hours), and experimental groups (24 hours of 0.5% serum with addition of either cCIC, EPC, MSC, C+E+M, or CardioCluster for additional 24 hours). Cardiomyocytes visualized by staining with sarcomeric actinin (α-actinin; red). TO-PRO-3 iodide (Topro; white) used to visualize nuclei. Scale bar, 50 μm. **C**, Quantitation of cardiomyocyte size relative to serum starved control. **D-E**, Gene expression of *desmin* **(D)** and *sdf-1* **(E)** in cardiomyocytes with and without the addition of cells. Data are presented as 1-way ANOVA, *p<0.05 and ***p<0.001, versus serum starved control (n=3).

Figure 4. CardioClusters exhibit increased protection from oxidative stress
A-D, Cell death assay performed on cardiac cell populations under 24 hours of low serum (75% serum reduction), followed by 4 hours of treatment with 30 µM H_2O_2 in low serum medium. **A-B**, Apoptosis visualized by Annexin V **(A)** and necrosis visualized by Sytox Blue **(B)**. Data are presented as 1-way ANOVA, *p<0.05, **p<0.01, ***p<0.001, versus cCIC (n=3-5). **C-D**, Brightfield images of single cells **(C)** and CardioClusters **(D)** 4 hours after H_2O_2 treatment. Transparent arrows highlight adherent (healthy) cells and black arrows highlight cells that have rounded up and detached from the tissue culture surface, presumably undergoing cell death.

Figure 5. CardioCluster treatment improves left ventricular wall structure and cardiac function after myocardial injury

A, Representative 2D echocardiography images (M-mode) at study completion (week 20). Para-sternal short-axis view showing LV anterior wall and posterior wall movement. **B-D**, Longitudinal assessment of LV fractional shortening (FS, %) **(B),** ejection fraction (EF, %) **(C)**, and anterior wall thickness in systole (LVAW;s, mm) **(D)** over 20 weeks. Data are presented as 1-way ANOVA, analyzed per timepoint, **p<0.01 and ***p<0.001, CardioCluster versus vehicle. +p<0.05 and +++p<0.001, C+E+M versus vehicle. $p<0.05, CardioCluster versus C+E+M. Sample size specified in Supplemental Table 6. **E**, Percent change in EF from week 1 to week 20. Data are presented as 1-way ANOVA, ***p<0.001, versus CardioCluster. ++p<0.01, versus C+E+M (n=7–8 mice per group). **F-G**, Bar graph showing LV volume in systole (Vol;s, µl; **(F)** and in diastole (Vol;d, µl; **(G)**. Data are presented as 1 Way ANOVA, *p<0.05, **p<0.01, ***p<0.001, versus CardioCluster. Sample size specified in Supplemental Table 6. **H**, Heart weight to tibia length ratio (HW/TL; mg/mm) at week 20. Data are presented as 1-way ANOVA, *p<0.05, **p<0.01, versus sham (n=4–5 mice per group). **I-M**, Masson's Trichrome staining used to evaluate LV fibrotic area. Percentage of fibrotic LV in CardioCluster, C+E+M and vehicle-treated hearts **(I)**. Data are presented as 1-way ANOVA, *p<0.05, versus CardioCluster (n=4–5 mice per group). Representative histology sections of sham **(J)**, vehicle **(K)**, CardioCluster **(L)**, and C+E+M **(M)** treated hearts 20 weeks after MI. Collagen-rich areas (scar tissue) are colored in blue and healthy myocardium in red.

Figure 6. Impact of CardioClusters on cardiac function measured by cardiac strain
A-B, Representative images of long-axis echocardiography recording (left panel), with cross-sectional segment synchronicity map (middle panels), and radial and longitudinal strain curves (right panels) at week 1 **(A)** and week 20 **(B)**. Strain curves (representing strain measures over time) are generated for the 6 standard myocardial regions, with a 7th line (shown in black) denoting the average strain. **C-D**, Speckle tracking echocardiography analysis used to determine global peak radial strain (%) **(C)** and global peak longitudinal strain (%) **(D)** at week 1, week 8, and week 20. Data are presented as 2-way ANOVA, *p<0.05, **p<0.01, ***p<0.001, versus sham. +p<0.01, ++p<0.05, +++p<0.001, versus CardioCluster. $p<0.05, versus C+E+M (n=6–8 mice per group). **E**, Speckle tracking echocardiography analysis used to determine regional peak radial and longitudinal strain (%) in the area of injury at week 1 and week 20. Data are presented as 2-way ANOVA, **p<0.01, versus vehicle. +++p<0.001, versus CardioCluster. $p<0.05, versus C+E+M (n=6–9 mice per group). **F**, Total change in regional radial and longitudinal strain from week 1 to week 20 in the area of injury. Data are presented as 1-way ANOVA, *p<0.05, **p<0.01, ***p<0.001, versus CardioCluster (n=6–9 mice per group).

Figure 7. CardioCluster show enhanced engraftment and persistence in the myocardial wall

A, Immunofluorescent tile scan of a cryosectioned heart from an uninjured animal injected with CardioClusters day 3 post-injection tracked by FluoSpheres with no antibody labeling required for visualization of cells. **A'-A'''**, Higher magnification of areas with white dotted boxes. A', Illustrates the removal of the 555nm channel in order to better visualize cCIC (green) and MSC (blue), without FluoSpheres. **A''- A'''**, 555nm channel restored **(A'')** and field of view magnified **(A''')**. **B-D**, Immunofluorescent images from cryosectioned MI hearts injected with CardioClusters at day 1 **(B)**, day 3 **(C)**, and week 1 **(D)** . **E-H**, Antibody labeled immunofluorescent images from MI hearts injected with CardioClusters at week 1 **(E)**, week 4 **(F)**, week 12 **(E)**, and week 20 **(H)**. **E'**, Higher magnification of areas with white dotted boxes. **G'**, Higher magnification of cCICs shown within white dotted box. **H'**, Higher magnification of cCICs and EPCs shown within white dotted boxes. Antibody labeling: anti-GFP labels cCIC, anti-mCherry labels EPC and MSC **(E)**; anti-GFP labels cCIC, anti-myc labels EPC and anti-HA labels MSC (F-H). Scale bars: 1mm (A); 100µm (B-H).

Figure 8. CardioCluster treatment increases capillary density in the infarct region
A-C, Quantitative analysis represents measurement of capillary density in the infarct **(A)**, border zone **(B)**, and remote **(C)** heart regions (n= 4-6 mice per group). Dashed line: mean capillary density of sham group. **D-G**, Representative image of isolectin+ vessels in sham **(D)** and infarct regions of Vehicle **(E)**, CardioCluster **(F)**, and C+E+M **(G)** used to quantitate capillaries (isolectin B4; green), cardiac troponin T (cTNT; red), and nuclei (DAPI; white). Right panels show higher magnification of isolectin+ vessels from areas highlighted by yellow boxes. Scale bar, 200 μm. Data are presented as 1-way ANOVA, ***p<0.001, versus CardioCluster.

Figure 9. CardioCluster treatment antagonizes cardiomyocyte hypertrophy in the border and remote region and preserves cardiomyocyte size in the infarct region

A-C, Quantitative analysis represents measurement of cross-sectional area of cardiomyocytes from infarct **(A)**, border zone **(B)**, and remote **(C)** heart regions (n=4-5 mice per group). Dashed line: mean cross-sectional area of sham group. **D-G**, Representative image from sham **(D)** and border zone regions of vehicle **(E)**, CardioCluster **(F)**, and C+E+M **(G)** used to quantitate cardiomyocyte cross-sectional area (wheat germ agglutinin [WGA]; cyan), cardiac troponin T (cTNT; red), and nuclei (DAPI; blue). Right panels show higher magnification of areas highlighted by yellow boxes. The area of three traced cardiomyocytes per region are shown within magnified view. Scale bar, 200 µm. Data are presented as 1-way ANOVA, ***p<0.001, versus CardioCluster. +++p<0.001, versus C+E+M. $$$p<0.001, versus sham.