1 **Original Article**

2 # Platelet proteolytic machinery assessment in Alzheimer's Diseases

3 **Roy G. Muriu [1], Jessica M. Sage [1], Abdulbaki Agbas [1]**

4 *[1]Department of Basic Sciences, Kansas City University of Medicine and Biosciences, Kansas City, MO 64106 ,*
5 *USA*
6

Correspondence to: Dr. Abdulbaki Agbas, Department of Basic Sciences, Kansas City University of Medicine and
Biosciences, 1750 Independence Avenue, Kansas City, MO 64106 , USA, aagbas@kcumb.edu; ORCID:
0000-0003-3854-7784

Abstract

Aim: Platelets provide substantial information about the proteolytic system profile in neurodegenerative
diseases. Assessment of autophagy and proteasome activities in platelets may reflect tissue proteolytic
system profile in central nervous system in Alzheimer's diseases (AD). We aimed to demonstrate the
optimum assay conditions and identify target proteins in platelet proteolytic system.

Methods: Autophagosome forming proteins were identified by Western blotting analysis. Standard gel
electrophoresis and electro transfer apparatus were used for protein transfer. Several vendors' products
were tested to identify the best working antibodies and their concentrations were optimized. An ELISA
kit was used for platelet proteasome protein determination. Infrared imaging technology was used for
visualizing the proteins on the membrane. Platelet samples were obtained from clinically verified AD
patients and age-matched non-demented control subjects that were recruited by University of Kansas
Alzheimer's disease Center.

Results: Autophagosome forming proteins showed elevated levels in AD patient platelet cytosol. Only
LC3-I autophagosome protein levels were significantly elevated. The platelet lysate proteasome
concentrations were assessed. AD patient's proteasome levels were elevated but they were statistically
not important as compared to controls.

Conclusions: Platelets can be used for assessing diseased protein levels and whether proteolytic system
is functional. Blood-based sampling from human donors is less-invasive and analyzing platelet
proteolytic system profile may help to develop pharmaceutical intervention approaches for
neurodegenerative diseases in general.

Keywords: Autophagy, Proteasome, platelets, Alzheimer's disease, protein aggregation, TDP-43,
neurodegeneration

INTRODUCTION

The purpose of this study is to demonstrate that human blood-derived platelets may provide critical
information about malfunctioned proteolytic machinery leading to diseased protein aggregation in a
neurodegenerative disease. Alzheimer's Disease (AD) is mainly characterized by protein aggregations
and deposition that are toxic and lethal to cellular structures in the central nervous system [1] . The
proteolytic system that includes autophagy and proteasome pathways, degrade and allow recyclization of
targeted molecules and organelles in all eukaryotic cells (Fig.1). The reason certain proteins are

allowed to aggregate is the inhibition of proteolysis and down regulation of effector proteins that
stimulate the pathogenesis of the degradation [2]. A dysfunctional proteolytic system, including autophagy
and proteasome, is implicated in the pathogenesis of AD. Autophagy is a lysosomal degradative process
by which cellular residents are inducted into homeostasis through quality control by clearance of
pathogenic proteins, recycling of macromolecules, and response to energy requirements [3] . In
neurodegeneration, this system has been proven to be dysfunctional and therefore associated with lack of
proper protein disposal

Figure-1 Schematic diagram of the proteolytic system in autophagic and
Proteasomal pathways.

and thereby aggregation [4] . A two-stage autophagy impairment (i.e., induction and lysosomal
acidification) leads to pathogenesis of AD [2] . During the induction process, autophagy requires the
release and presence of beclin-1 protein from endoplasmic reticulum and formation of a multimeric
complex [2] .This is the site of autophagy initiation through vesicle nucleation then formation of isolation
membrane [5]. Beclin-1 is a crucial regulator of autophagy and whose expression in the hippocampus is
decreased at the RNA and protein level in AD especially with advanced age. This protein is necessary for
nucleation of a phagophore membrane before the autophagosome vesicle is fully formed [6]. This concept
produced comparative results against control subjects that demonstrated an increase in protein
aggregation [7].
Proteasomes are molecular machines that degrade aberrant proteins through polyubiquitination process
followed by recognition by regulatory particle, deubiquitylation by deubiquitinase (DUB), and finally
degradation of proteins into fragments [8] . In AD, the components that are linked to this pathology are
abnormal formation of ubiquitin and activity inhibition [9, 10]. For example, paired helical filaments of tau
(PHF-tau) bind to proteasomes and thereby reduce its activity [9], resulting less polyubiquitination
Without this step, a protein without proper ubiquitination cannot be recognized by the regulatory particle
of the proteasomes [11]. Therefore, the autophagy and proteasome activities are a critical component and
can be measured in platelets. Although the functions of autophagy within platelets are largely unclear but
thus far, it is known that, its impairment leads to a lack of platelet aggregation and adhesion [12].
We have demonstrated that a TAR-DNA/RNA binding protein (TDP-43) and its phosphorylated
derivative (pTDP-43) levels were elevated in platelets obtained from AD patients as part of the

blood-based biomarker development studies [13]. In this study, we hypothesize that platelets are known to

have an active autophagic (specific aim-1) and proteasome (specific aim-2) machinery; therefore,

autophagic and proteasome protein expression will be expected to be different in platelets from AD and

non-demented individuals. We predict that previously observed TDP-43 protein accumulation [13] might

be relevant to dysfunctional proteolytic system.

METHODS

1. *Human Platelets* : AD patients and age-matched non-demented control subject human platelets were

obtained from the biorepository of University of Kansas Alzheimer's Disease Center under the

approved IRB protocol (KUADC#11132). 5-6 days old platelets were obtained from a local

community blood center (CBC) for initial studies before analyzing AD patients and control human

platelet samples.

2. *Materials*:

2.1. Antibodies:

2.1.1. MBL antibodies (MBL International, 15A Constitution way, Woburn, MA 01801, USA):

Anti-LC3(pAb, PM036Y), Anti-LC3 (8E10, mAb, M186-3Y), Anti-LC3 (4E12, mAb,

M152-3Y), Anti-Beclin-1 (pAb, PD017Y), Anti-Atg (pAb, PM040Y), Anti-p62

(SQSTM1, pAb, PM045Y), Anti-Atg5 (pAb, PM050Y), Positive control for anti-LC3

(PM036-PNY).

2.1.2. Cell signaling antibodies (Cell Signaling Technology, 3 Trask Lane, Danvers, MA 01923,

USA): Anti-Beclin-1 (pAb, 3738S), LC3A (mAb, 4599S), LC3B (pAb, 2775S).

2.1.3. Abcam antibody (Abcam, Inc., 1 Kendall Square Suite, B2304, Cambridge, MA

02139-1517, USA): p62 (SQSTM1, AB56416).

2.1.4. Secondary antibodies (LI-COR Inc., 4308 Progressive Ave., Lincoln, Nebraska 68504,

USA): Goat anti-Mouse (green wavelength) antibody (Li-Cor, C40213-01), Goat

anti-Rabbit (green wavelength) antibody (Li-Cor, C30829-02)

2.2. SDS/PAGE   and Western Blotting reagents

Butanol, 1.5M Tris-HCl (pH 8.8), 1.0M Tris-HCl (pH 6.8), 10% Sodium Dodecyl Sulfate

(SDS), N,N,N',N'-Tetra-methyl ethylenediamine (TEMED, Bio-Rad, 161-0801), 10%

ammonium persulfate (APS) (Bio-Rad # 161-0700), 30% acrylamide and bis-acrylamide

solution 29:1 (Bio-Rad, 1610156), urea (VWR, BDH4214-500G). Pre-Cast 4-20% gradient

gel (Bio-Rad #456-1096 PVDF membrane (Millipore, Immobilon-FL Transfer Membranes

IPFL00010) Membrane blocking agent, SeaBlock (ThermoFisher, # 37527); Total protein

staining solution (REVERT Total Protein Stain kit, LI-COR Inc., # 926-11010; Pyronin Y

(Sigma-Aldrich # P9172)

2.3. Protein concentration determination : Pierce^{TM} bicinchoninic acid (BCA) protein assay kit

(Thermo Scientific) (UB276872),

2.4. ELISA commercial kit: For quantifying proteasome (Enzo 20S/26S Proteosome ELISA Kit,

Catalog # BML-PW0575-0001)

2.5. *Software:* Image Studio Lite (Ver.4.0) for image analyses. This software is part of Odyssey

(LI-COR) image analyzer. Online-based free statistics calculator was used for statistical

analysis (

125   (https://www.danielsoper.com/statcalc/calculator.aspx?id=47)

126   *2.6. Experimental apparatus:*

127   2.6.1. Sonic dismembrator (Fisher Scientific, Model: XL2000-350)

128   2.6.2. Table top centrifuge (Eppendorf, Model: 5418)

129   2.6.3. Odyssey Infrared Imager (Model : 9120, LI-COR Inc., 4308 Progressive Ave. Lincoln,

130   Nebraska 68504)

131   2.6.4. Mini Protean III Electrophoresis system (BioRad   165-3301

132   2.6.5. Electro transfer system (Mini Trans-Blot Electrophoretic Transfer Cell Bio-Rad 170-3930

133   2.6.6. Multi-well plate reader (Bio-Tek Cytation 5 or Bio-Tek Synergy HT)

134   3.   Procedures:

135   3.1. *SDS/PAGE and Western Blotting*: The cytosolic proteins of platelet lysates form AD patients

136   and control subjects were separated on a homemade 4-12% gradient gel with 1.5mm width 20

137   well, MINI Protean II casted gel, using sodium dodecyl sulfate polyacrylamide gel

138   electrophoresis (SDS-PAGE). The apparatus was filled with 1x electrophoresis buffer. With

139   the gel inside the buffer containing cell, each lane was loaded with pyronin Y lane marker and

140   30 µg total protein (1 mg/mL) Electrophoresis was performed at 75 volts for an average of 100

141   min. until front dye (Pyronin Y) was at the bottom of the gel. The gel was removed from the

142   sandwiched plates and processed for protein transferring to a methanol-activated

143   polyvinylidene difluoride (PVDF) membrane. Electro-transfer process was carried out at 75

144   volts for 30 minutes. Transfer buffer did not include methanol. The low-temperature of transfer

145   unit was maintained by either inserting an ice-block or placing the transfer unit in an ice-filled

146   container. The PVDF membrane was then stained for total protein visualization by total protein

147   staining kit (REVERT, Total Protein Stain Kit LI-COR Inc., 926-11010). Transferred proteins

148   were imaged by Odyssey (LI-COR) imaging system at the 700 nm set channel. The staining

149   was optionally removed by incubating with REVERT reversal solution for a maximum of 10

150   minutes as per protocol, then finally rinsed twice with nanopure water. The membrane was

151   blocked in 2 mL of 1:1 Seablock /TBS buffer for 1 hour at room temperature (RT) on an orbital

152   shaker. Afterwards, the membrane is directly transferred to a container with 10 mL of 1:1

153   Seablock/TBST and primary antibody dilution in1:500 or 1:1000 while incubating on an orbital

154   shaker for overnight (at least 12 hours) at 4℃. Next day, the membrane was washed with 1X

155   TBST for   20-minutes then incubated horseradish peroxidase conjugated secondary antibody,

156   goat anti-mouse (LC3) and goat anti-rabbit (P62, Beclin-1, and Atg5-12) at 1:10,000 dilution

157   for 1 hour at RT on an orbital shaker. Additional 20-minutes wash with 1x TBST was

158   performed to remove unbound antibodies prior to imaging was completed. The membrane was

159   scanned by Odyssey (LI-COR) imager and analyzed by densitometric quantification using

160   Image Studio Lite (Ver.4.0).

161   3.2. *26s/20s Proteasome analysis by ELISA method*:   We initially utilized platelets obtained

162   from CBC to optimize the assay conditions for the proteasome quantification. Three sample

163   fractions (i.e., whole platelet lysate, clear supernatant, and membranous pellet) were

164   analyzed by proteasome ELISA kit. Whole platelet lysate (40 µg per assay well) for

165   proteasome protein content determination in AD patients   and in non-demented control

subjects. Assay procedure was performed according to the manufacturer provided protocol.

Absorbance was read at 450 nm wavelength using a multi-well plate reader (Bio-Tek,

Synergy HT). Standard curve was established and slope of the curve was used to determine

the concentration of unknown sample proteasome 26s/20s.

4. *Statistical analyses :* The quantitative analysis for Western blots were performed using Image Studio

Lite software program (V 4.0.) The quantitative value of autophagy protein markers represent

arbitrary units (a.u.) based on the intensity of the bands. Statistical results were calculated by a

two-tailed unpaired student t-test and Mann-Whitney U test. This was coupled with calculating the

value of Cohen's d and the effect-size correlation, R, using the means and standard deviations of two

groups (AD and control). Error bars on all data represents standard error of the mean ($\pm$S.E.M)

RESULTS

We have tested a battery of antibodies for probing autophagy protein purchased from three different

vendors and compared to each other for selecting the best working antibodies. Individual platelet lysate

samples from AD and control group (n=9) were analyzed by Western blotting technique using selected

antibodies. The results from

three replicates averaged and independently tested as an interval type of data. We only found an increase

of LC3B-I ($p \leq 0.02$) in AD when compared to Control (Fig.2). The select autophagosome proteins

showed an elevated profile; however, there was no statistical difference between AD patients and control

group.

Figure-2 Autophagy protein profiles in Human AD and in non-demented (control) platelet cytosols. Comparative analysis of autophagy proteins in platelet cytosols. The LC3B-I/II and Atg5-12 proteins were identified using antibodies from MBL vendor. The Beclin-1 was detected using Cell-Signaling antibody, and the p62 protein was detected by using Sigma-Aldrich antibody. Each samples were analyzed three times and Mean±SEM was presented. The protein band intensities were normalized to total protein staining. Student t-test was employed and only LC3B-1 showed the significance ($p \leq 0.02$).

187
188
189

190 The proteasome concentrations of platelet lysates from AD and age-matched control subjects (n=12)
191 were
192 assessed by ELISA method. Although AD patient platelet lysate proteasome levels were elevated, no
193 statistical difference between patient and control group was obtained (Fig.3)

Figure-3 Proteasome concentration measurements in whole platelet lysate. The bar graph represents the proteasome concentration in whole platelet lysate obtained from AD and control subjects (Mean±SEM; n=12). Although an increased proteasome protein profile observed in AD samples, no statistical importance was obtained. The Inset figure shows a typical standard curve for quantifying proteasome 20s concentrations.

DISCUSSION

$A\beta_{1-42}$ is widely understood to accumulate in the early stages of the AD pathology. This malfunctioning
protein recruits and triggers microglial, astrocyte facilitated clearance, but soon $A\beta_{1-42}$ depositions
overwhelm the response leading to microglia-mediated neuroinflammation [14-16]$A\beta_{1-42}$ also stimulates the
catalysis by NADPH oxidase. This enzyme activity increases reactive oxygen species production. The
presence of these harmful chemicals activates Atg4, which cleaves and lipidates LC3-I with
phosphoethanolamine (PE), as part of the induction of the autophagic pathway. Since AD is a multi-faceted
disease, somehow the proteolytic system is suppressed due to downregulation[2, 6], allowing for the
characteristic extracellular amyloid plaques and neurofibrillary tangles to develop.

Some reports also recently discovered that platelets contain a proteolytic system by which hemostasis and
thrombosis is acquired [17, 18]. Since it was also discovered that an elevation of pathological pTDP43 within
the platelet cytosol is correlated to hippocampal cortex of AD patients [13] , it was imperative that we
analyze these platelet proteolytic systems. We attempted to establish any change of the protein
concentrations of proteolytic system, since platelet proteasome concentrations have not been reported prior
to our studies. It was initially believed and hypothesized, based on aforementioned reports, the proteolytic
system is somehow altered and that we would expect to observe some degree of variations in proteasome
concentrations.

In specific aim-1, certain autophagy marker antibodies (i.e. Beclin-1, Atg5, p62, and LC3) were analyzed
for autophagy forming proteins in platelets. This was in reference to previous autophagy activity
measurements by Gupta eta al[17] .The antibodies from multiple manufacturers have different affinities for
a given sample. A series of antibody dilutions and platelet sample concentrations were optimized in order
to match the best antibody /protein combination to be used in the assays. Those results suggested 30 μg
whole platelet lysate proteins paired with a primary antibody dilution of 1:1000 was ideal for future
analysis.

Next, we evaluated only LC3A and LC3B antibodies in Western blotting.These findings distinguished that

human platelet samples are not sensitive to the detection of LC3B isoform. Other researchers have stated
that out of the three possible isoforms (LC3A, LC3B, and LC3C). LC3A and LC3B autophagosome
components exhibit distinct expression patterns in different human tissues [19]. After the final
measurement of three replicates, the only reasonable results found were the proteins levels of LC3-I (a
cleaved form of LC3). LC3-II (lipidated LC3-I) is a conjugated protein marker for platelet autophagy
system [20] .Since there was no change in levels of LC3-II, the difference in LC3-I between the groups (p ≤
0.021) could be a result of autophagic pathway being blocked in its initial steps in AD [21]. The overarching
implication we believe is that in AD, probably to alleviate aggregate stress, platelet allocated autophagy
may be functionally upregulated and induced but inhibited in early or late steps of the pathway[20] . By only
immunoblotting the for LC3, our results are more likely inconclusive. The immunoreactivity of LC3-I and
LC3-II are different. LC3-II tends to be more sensitive to antibodies[21]. A stimulated autophagic pathway
is represented by reduction of autophagy marker LC3II. So, when we analyzed that there was no change in
LC3-II, its more indicative of no detection of flux than the changes of LC3-I. Therefore, an autophagy
activity analysis assessment should be performed in freshly isolated intact platelets.

In specific aim-2, we assessed the proteasome concentration in platelet lysates. Poteasome 20S protein
levels were measured by enzyme-linked immunosorbent assay (ELISA) techniques in cell lysates,
supernatants, and pellets in CBC obtained healthy samples. We relied on supernatant isolations of platelet
lysates to measure AD profiled proteasome concentrations. We found no statistical difference (p= 0.373)
between AD patients   and age-matched Control subjects in these platelet samples based on a two-tailed,
student t-test and Mann-Whitney U test.

To assess the relationship of platelet proteolytic machinery protein levels in AD more effectively than the
use of only p values we incorporated effect sizes [22, 23]. In addition to selecting a student t-test (interval,
parametric), we also stepped down to a Mann-Whitney (ordinal, non-parametric) level of measurement.
This was justified by the expression of a large variation in protein concentrations between individual
samples within each group. Which means that our sample did not meet the parametric assumption of equal
variances. This adjustment ascribed the data into ranked categories before comparing the median values
between the groups.

Despite the statistical insignificance between the two groups means of Atg5-12, p62, Beclin-1, a measure
of strength of an effect points to a promising conclusion in future analysis. A moderate to large standard
deviations (d = 0.41–1.0) and correlations (R = 0.36-0.44) between AD and control were detected.
Denoting a probability that there is a difference in autophagic pathway protein levels, but our sample is not
large enough to produce statistical significance. On the other hand, the insignificance found between AD
and control regarding LC3-II protein bands was coincided with a small magnitude of effect between
groups. (d = 0.27, R = 0.13, p = 0.58) (Fig.1B) About proteasome testing (p = 0.373) its statistical
insignificance was also matched with a small group effect (d = 0.34, R = 0.17).

In retrospect, there are some caveats and setbacks, largely rendering our results with a low difference. First,
our continuous and averaged data comparison was statistically insignificant for all except one based on our
sample cohort (n=9 for autophagosome and n=12 proteasome analyses).As mentioned before, human
sample do have a tremendous variation in concentration and effect between each sample and group
measurements. Especially with a small effect size for the main autophagy monitor, LC3-II, increasing the n
value to ~220 per group (n=220) should produce a significant t-test at a probability level of 0.05 for all
samples [24]. Secondly, as it has been reported that LC3-I is unstable and sensitive to freezing-thawing
cycles or SDS buffered cocktail [25]. Future samples should be freshly assessed and not exposed to a
repeated freeze-thaw cycle. Thirdly, immunoblotting has its limitation of determining autophagic
fluctuations. As we understand that LC3-II is the go-to marker, an increase in its presence could be due to
several reasons. It could probably correlate with autophagosome accumulation but autophagic flux is not
guaranteed [26]. In more relevance to AD neuropathology, beclin-1 is downregulated in this disease;
however, LC3-II can still form without a phagophore and on ectopic membranes [21, 25].

It is known that a neurodegenerative condition such as AD effects the periphery by increase in thrombin and von Willebrand factor (vWF) protein. Thrombin and vWF are platelet activators [18] . In AD, the patients are known to express an increased amount of vWF, which is partially used to convert quiescent platelets into activated platelets [27] . Hence, activated platelets of these patients produce and release more Aβ into circulation compared to the controls [28]. Aβ usually interacts with fibrin thereby promoting coagulation and fibrin aggregation. Therefore, an increase of Aβ could mean an increase in platelet aggregation that implies an alternate flux in hemostasis. We did not analyze Aβ peptide levels in platelet lysates of AD patients and age-matched control subjects in this study; however, we plan to include the platelet lysate $Aβ_{1-42}$ measurements in a follow-up study

Since our results are supposed to be a testament of basal autophagy in quiescent platelets, AD derived samples might have already been activated when obtained from the patients. Activated platelets have increased autophagic pathway activity [18]. Therefore, the significant measurements might be due to that previous condition. The best way to adjust for that variable would be activate the control platelets before any protein concentration measurements. An alternative approach would be to inhibit the platelet activation factor by including an inhibitor (PGI2) during the platelet isolation from whole blood.   Platelet proteolytic system analysis has allowed us to differentiate probable proteostasis between probable AD and control in cross-sectional fashion. It is clear that the elevated presence of TDP-43 is equally represented in AD brain homogenate segments of the hippocampus and isolated platelet lysates [13]. One of the possibilities that can be discerned from our study is that there may be elevated induction of autophagy, beyond basal amount. With a higher probability, somewhere along the autophagic pathway, there is suppression of progression into the pathway.

In light of this information, platelet autophagic profile may be similar to neurons of this disease profile. This study was meant to extrapolate the extent to which AD can influence the state of these systems during an elevated presence of disease-related proteins. From now on, we would need to use freshly obtained human samples, to include PGI2 inhibitor during the platelet isolation, to increase our sample size to about 220. Autophagy flux and proteasome assessment in intact platelet are necessary for obtaining more evidences that are conclusive. A similar cross-sectional study with a few adjustments should be considered before testing other neurodegenerative disease group in a similar fashion.

DECLARATIONS

Authors' contributions
Muriu, RG made substantial contributions to conception and design of the study and performed data analysis and interpretation
Sage, JM performed data acquisition, as well as provided administrative, technical, and material support
Agbas, A conceptualized the study, analyzed the data, and wrote the manuscript

Availability of data and materials

Not applicable

Financial support and sponsorship

Research reported in this publication was supported by several pilot project funds from QS85523J, University of Kansas Medical Center Research Institute, Inc.(QS85523J), FONTIERS-Trail Blazer Award (01–2429–001) and KCU intramural grants. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed. No writing assistance was utilized in the production of this manuscript.

Conflicts of interest

All authors declared that there are no conflicts of interest

Ethical approval and consent to participate

The authors state that they have obtained appropriate institutional review board approval (KUADC#11132) or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

Consent for publication

Not applicable

REFERENCES

1. Tuite, M.F. and R. Melki, *Protein misfolding and aggregation in ageing and disease: molecular processes and therapeutic perspectives.* Prion, 2007. **1**(2): p. 116-20.
2. Ghavami, S., et al., *Autophagy and apoptosis dysfunction in neurodegenerative disorders.* Prog Neurobiol, 2014. **112**: p. 24-49.
3. Klionsky, D.J., *Autophagy: from phenomenology to molecular understanding in less than a decade.* Nat Rev Mol Cell Biol, 2007. **8**(11): p. 931-7.
4. Meijer, A.J. and P. Codogno, *Autophagy: regulation and role in disease.* Crit Rev Clin Lab Sci, 2009. **46**(4): p. 210-40.
5. Virgin, H.W. and B. Levine, *Autophagy genes in immunity.* Nat Immunol, 2009. **10**(5): p. 461-70.
6. Pickford, F., et al., *The autophagy-related protein beclin 1 shows reduced expression in early Alzheimer disease and regulates amyloid beta accumulation in mice.* J Clin Invest, 2008. **118**(6): p. 2190-9.
7. He, C., et al., *Exercise-induced BCL2-regulated autophagy is required for muscle glucose homeostasis.* Nature, 2012. **481**(7382): p. 511-5.
8. Gadhave, K., et al., *The ubiquitin proteasomal system: a potential target for the management of Alzheimer's disease.* J Cell Mol Med, 2016. **20**(7): p. 1392-407.
9. Keck, S., et al., *Proteasome inhibition by paired helical filament-tau in brains of patients with Alzheimer's disease.* J Neurochem, 2003. **85**(1): p. 115-22.
10. van Leeuwen, F.W., et al., *Frameshift mutants of beta amyloid precursor protein and ubiquitin-B in Alzheimer's and Down patients.* Science, 1998. **279**(5348): p. 242-7.
11. Chen, X. and D. Petranovic, *Role of frameshift ubiquitin B protein in Alzheimer's disease.* Wiley Interdiscip Rev Syst Biol Med, 2016. **8**(4): p. 300-13.
12. Feng, W., et al., *Dissection of autophagy in human platelets.* Autophagy, 2014. **10**(4): p. 642-51.
13. Wilhite, R., et al., *Platelet phosphorylated TDP-43: an exploratory study for a peripheral surrogate biomarker development for Alzheimer's disease.* Future Sci OA, 2017. **3**(4): p. FSO238.
14. Kinney, J.W., et al., *Inflammation as a central mechanism in Alzheimer's disease.* Alzheimers Dement (N Y), 2018. **4**: p. 575-590.
15. Hickman, S.E., E.K. Allison, and J. El Khoury, *Microglial dysfunction and defective beta-amyloid clearance pathways in aging Alzheimer's disease mice.* J Neurosci, 2008. **28**(33): p. 8354-60.

16. Rothhammer, V. and F.J. Quintana, *Control of autoimmune CNS inflammation by*
astrocytes. Semin Immunopathol, 2015. **37**(6): p. 625-38.

17. Gupta, N., et al., *Proteasome proteolysis supports stimulated platelet function and*
thrombosis. Arterioscler Thromb Vasc Biol, 2014. **34**(1): p. 160-8.

18. Ouseph, M.M., et al., *Autophagy is induced upon platelet activation and is essential*
for hemostasis and thrombosis. Blood, 2015. **126**(10): p. 1224-33.

19. He, H., et al., *Post-translational modifications of three members of the human*
MAP1LC3 family and detection of a novel type of modification for MAP1LC3B. J Biol
Chem, 2003. **278**(31): p. 29278-87.

20. Zhang, X.J., et al., *Why should autophagic flux be assessed?* Acta Pharmacol Sin, 2013.
34(5): p. 595-9.

21. Mizushima, N. and T. Yoshimori, *How to interpret LC3 immunoblotting.* Autophagy,
2007. **3**(6): p. 542-5.

22. Lakens, D., *Calculating and reporting effect sizes to facilitate cumulative science: a*
practical primer for t-tests and ANOVAs. Front Psychol, 2013. **4**: p. 863.

23. Nakagawa, S. and I.C. Cuthill, *Effect size, confidence interval and statistical*
significance: a practical guide for biologists. Biol Rev Camb Philos Soc, 2007. **82**(4): p.
591-605.

24. Sullivan, G.M. and R. Feinn, *Using Effect Size-or Why the P Value Is Not Enough.* J Grad
Med Educ, 2012. **4**(3): p. 279-82.

25. Klionsky, D.J., et al., *Guidelines for the use and interpretation of assays for monitoring*
autophagy in higher eukaryotes. Autophagy, 2008. **4**(2): p. 151-75.

26. Gimenez-Xavier, P., et al., *LC3-I conversion to LC3-II does not necessarily result in*
complete autophagy. Int J Mol Med, 2008. **22**(6): p. 781-5.

27. Peyvandi, F., I. Garagiola, and L. Baronciani, *Role of von Willebrand factor in the*
haemostasis. Blood Transfus, 2011. **9 Suppl 2**: p. s3-8.

28. Tang, K., et al., *Platelet amyloid precursor protein processing: a bio-marker for*
Alzheimer's disease. J Neurol Sci, 2006. **240**(1-2): p. 53-8.

396