Hybridizing salamanders experience accelerated diversification

Austin H. Patton[1,*], Mark J. Margres[1,2], Brendan Epstein[1,3], Jon Eastman[1], Luke J. Harmon[4], and Andrew Storfer[1]

[1]Washington State University, School of Biological Sciences, Pullman, WA, 99164, USA
[2]Harvard University, Department of Organismic and Evolutionary Biology, Cambridge, MA, 02138, USA
[3]University of Minnesota, College of Biological Sciences, Saint Paul, MN, 55108, USA
[4]University of Idaho, Department of Biological Sciences and IBEST, Moscow, ID, 83844, USA
[*]austin.patton@wsu.edu

ABSTRACT

Whether hybridization generates or erodes species diversity has long been debated, but to date most studies have been conducted at small taxonomic scales. Salamanders (order Caudata) represent a taxonomic order in which hybridization plays a prevalent ecological and evolutionary role. We employed a recently developed model of trait-dependent diversification to test the hypothesis that hybridization impacts the diversification dynamics of species that are currently hybridizing. We find strong evidence supporting this hypothesis, showing that hybridizing salamander lineages have significantly greater net-diversification rates than non-hybridizing lineages. This pattern is driven by concurrently increased speciation rates and decreased extinction rates in hybridizing lineages. Our results support the hypothesis that hybridization can act as a generative force in macroevolutionary diversification.

Introduction

A leading unresolved question in evolutionary biology is whether hybridization, defined as the interbreeding between two genetically distinct lineages[1], acts as a creative or destructive evolutionary force[2,3,4,5,6]. The prevailing view in the animal literature is that hybridization constrains lineage diversification because hybrid lineages are often documented to be less fit

than parentals[4,5,7]. Under this scenario, hybridization is predicted to increase extinction rates.

Further, introgressive hybridization has the potential to "wash away" accumulating divergence

among incompletely isolated lineages[8,9,10], leading to a prediction of decreased speciation rates.

In contrast, the prevailing view in the plant literature is that hybridization enhances adaptive

potential by introducing novel genetic and phenotypic variation[11,12,13]. Reinforcement, or the

accumulation of post-zygotic reproductive isolation through selection against hybrids[14], has long

been considered to expedite the speciation of diverging lineages[9]. Additionally, hybridization-

mediated shuffling of old genetic variants may fuel rapid diversification, as outlined by the

combinatorial view on speciation[15]. Accordingly, hybridization is predicted to increase

speciation rates and/or decrease extinction rates.

Whereas evidence of hybrid speciation in plants has long been abundant, evidence for

widespread hybrid speciation in animals is relatively scarce[6,10,16,17,18,19]. In allopolypoid hybrid

speciation, a mode of speciation common in plants but rare in animals, nearly complete

reproductive isolation may evolve in a single generation due to a change in ploidy[11]. In contrast,

homoploid hybrid speciation typically has to occur in the face of continued gene flow, which acts

to homogenize the diverging hybrid lineages[10].

Interest in hybrid-mediated speciation has recently burgeoned, but studies have typically

been limited in taxonomic scope. Studies of the effect of hybridization on diversification have

most commonly been conducted among closely related pairs or small clades of taxa, and results

have been equivocal[1,20,21,22,23,24,25,26]. Additionally, studies of hybridization often occur along

different stages of the speciation continuum[27], whereby hybridization can appear as a force that

either facilitates or impedes speciation. Recent work highlights this uncertainty by demonstrating

that the outcomes of hybridization depend on the underlying nature of selection pressures and

demography[28]. We suggest that studies at broad phylogenetic and macroevolutionary timescales

can help overcome these limitations by providing a phylogenetic context in which to view

repeated hybridization events over evolutionary timescales and the consequent impact on lineage

diversification rates.

Here, we conduct a taxonomically-broad test of the relationship of hybridization with

macroevolutionary diversification rates. We study salamanders (order Caudata, ca. 716 spp as of

October 2018[29]), which are particularly suitable because hybridization is pervasive and has been

studied extensively (i.e., nearly 1/3 of N. American species hybridize: Supplementary Fig. S1,

Supplementary Table S1). Additionally, sufficient sequence data are available to resolve the

phylogenetic relationships among most (~63%) taxa within this group[30]. If hybridization plays a

meaningful role in the diversification process, differences in diversification rates among

hybridizing and non-hybridizing taxa are expected. We thus test the hypothesis that there is a

difference in diversification rates (speciation and extinction rates) between contemporaneously

hybridizing and non-hybridizing salamander lineages. Note that we are simply testing whether

contemporary hybridization influences diversification rates, not whether ancient hybridization

facilitated the present radiation as postulated by the hybrid swarm hypothesis[6], because our

experimental design cannot address this (See supplement). We replicate this test across four

datasets to investigate the robustness of our results: 1) including all available data; 2) exclusion

of species that do not exhibit sympatry (defined as <10% geographic range overlap) thereby

lacking the opportunity to hybridize; 3) only the family Plethodontidae, which are the most

widely hybridizing and diverse of the 10 salamander families; 4) all (nine) salamander families

except the Plethodontidae.

Materials and Methods

Data Collection

We used the time-calibrated phylogeny of Amphibia[3] as the source for downstream analyses. This tree[30] was constructed using nine nuclear genes and three mitochondrial genes as data using RAxMLv7.2.8[31], and time-calibrated using treePL[32]. Using the APE package (v.3.4[33]) in R, we extracted the subtree containing salamanders for subsequent analyses. This approach yielded a tree containing 469 of 716 extant species of salamanders (Supplementary Fig. S1[30]) representing approximately two-thirds of the known diversity.

Each species was scored as 'non-hybridizable' (NH) or 'hybridizable' (H), based on an extensive literature review of hybridization using the search engines Google Scholar and ISI Web of Knowledge. To do so we paired each species with the terms 'hybrid' and 'introgress' as well as dialectical and structural variants thereof (e.g., 'hybridization', 'introgressed') to search for cases of hybridization. Criteria for hybridizability under the first, "narrow" definition was the documentation of hybridization among natively distributed species, specifically the observation of heterospecific mating or hybrid offspring in the wild as detected by morphological and molecular intermediacy. Criteria for hybridizability under the second, "broad" definition included those criteria described previously, as well as: 1) the observation of hybridization occurring in laboratory settings or among introduced and native species, and; 2) inference of historical introgression as determined using molecular lines of evidence (e.g. substantial and replicated genealogical discordance among molecular markers or detection via Approximate Bayesian Computation methods). This latter definition is less conservative than the former. Although inference of hybridization through the detection of genealogical discordance warrants caution, our narrow definition of hybridization does not recognize these species as hybridizable.

In total, we retrieved 56 papers (date-range: 1957-2017). We confirmed hybridization for

roughly 11 and 13 percent of extant Caudates for the narrow and broad datasets, respectively (78

and 92 of 716 species: 17 and 20 percent of sampled taxa; Supplementary Table S1).

Documented hybridization was absent from four families (Cryptobranchidae, Sirenidae,

Proteidae, Rhyacotritonidae) in the narrow dataset and two families (Rhyacotritonidae,

Sirenidae) in the broad dataset.

In addition to analyzing the entire salamander subclade from the Pyron[30] amphibian

phylogeny (469 total species), we compared diversification rates using three additional datasets,

produced using both the narrow and broad datasets (i.e., for a total of eight datasets). In the first

dataset, we required that "non-hybridizable" taxa were sympatric (>10% range overlap) with

another species, thus possessing sufficient opportunity to hybridize. Consequently, species that

were not sympatric with any other salamander taxa were excluded from the dataset. Therefore,

species classified as non-hybridizing in this analysis may have limited opportunity to hybridize

but have not been observed to do so. Although we cannot account for historical species

distributions (e.g. species' ranges may have previously overlapped), our primary (narrow)

definition of hybridizability necessitates contemporary hybridization. Thus, our designation of

sympatry occurs at the same time scale as our designation of hybridizability.

In the second and third datasets, we tested for a family-specific effect of hybridization on

diversification rates. As plethodontids are the most diverse family of salamanders (Fig. 1B) and

also the most well studied, the greatest number of instances of hybridization have been

documented in this group. Additionally, many plethodontids of the genus *Plethodon* in the

Eastern United States have been described as species on the basis of a threshold genetic distance

(e.g. [34, 35,36,37]). Coincidentally, nearly half of these species have been observed to hybridize in

nature. Thus, we produced datasets including either only members of family Plethodontidae, or

the nine families that exclude plethodontid salamanders. In total, eight analyses were conducted,

using four trees for each of the narrow and broad datasets, respectively. Information on the total

number of species included in each tree, as well as the number of species in each state (i.e., H or

NH) may be found in Supplementary Table S1.

Assessment of Trait Dependent Diversification

We applied the HiSSE (Hidden State Speciation & Extinction[38]) trait-dependent diversification

model to test for differences in speciation and extinction rates between hybridizing and non-

hybridizing lineages. HiSSE infers speciation and extinction rates for a binary character while

allowing for heterogeneity in diversification rate to exist within each character state. In the

HiSSE model, hidden states are co-distributed with the trait of interest and account for

unsampled traits that may simultaneously contribute to the diversification process. The hidden

state need not be associated with any single trait but may instead be associated with a set of traits

or suite of traits. Inclusion of the hidden state thus ameliorates the confounding effects of

unsampled traits on diversification rate estimations by allowing for greater rate heterogeneity in

the tree than in previous SSE models[38]. Thus, we are in essence measuring the impact of

contemporary hybridization while controlling for other, correlated traits on diversification rate.

Finally, by accounting for increased rate heterogeneity in character-dependent and character-

independent (null) models through the inclusion of hidden states, model rejection properties are

greatly improved[38] relative to previous SSE models.

We evaluated a total of 14 competing models using HiSSE, half of which represent a

model of character-dependent diversification and the remaining half represented models of

character independence (Supplementary Tables S2-S9). Models of trait dependence varied in the

number of hidden states included and in the number of free transitions among states. In

character-dependent models, our four states where Hybridizing (H) and Non-hybridizing (NH),

each being associated with one of the two hidden state (A & B) for four total character

states/diversification regimes (H-**A**, H-**B**, NH-**A**, NH-**B**).

Additionally, to account for the fact that diversification rates may be biased by

incomplete sampling of extant diversity within a phylogeny[39,40,41], we assume that 20% of extant

species of salamanders hybridize in nature. This value was chosen because it approximately

equals the mean frequency of hybridization across our datasets (19.75). To explore the effect this

assumption had on our results, we repeated these analyses assuming 1) we have sampled all

extant hybridizing species (Supplementary Fig. S2), 2) our sampling of character states is

proportional to their prevalence in nature (Supplementary Fig. S3), and 2) that 30% of extant

species hybridize (Supplementary Fig. S4). Thus, while unable to designate all hybridizing taxa

as such in our phylogeny due to their not being represented in the literature, we have explicitly

addressed this uncertainty in our analysis. We elaborate upon our choice of sampling fractions in

the Supplementary Materials.

To assess whether the inclusion of extremely young species impacted our diversification

rate estimates, we ran a single analysis excluding species younger than 1MY, as per Beaulieu &

O'Meara[38] using the narrow dataset and all species. This in turn led to the removal of 14 species.

Results were qualitatively identical to those obtained including these young species, so all

subsequent analyses were conducted including them. To improve the performance of the

Maximum Likelihood optimization procedure implemented in HiSSE, we used simulated

annealing to first traverse the likelihood surface to identify optimal starting values for subsequent

ML-optimization. Rather than reporting the results of individual model fits, we instead take the

approach of investigating model-averaged parameter estimates for each sampled character state[42]

(see Supplementary Information for further justification). That is, parameter estimates obtained

from each fitted model are averaged together such that their contribution to the average is

proportional to their relative support (Akaike weights) among the set of candidate models

(Supplementary Table S13). This leads the best supported models to have the greatest impact on

the final model averaged parameter estimates. Diversification rates are returned for each sampled

state respectively, as not all models include hidden states.

To test for significance among diversification rates inferred for each state, we calculated

all possible ratios between non-hybridizing and hybridizing species' model-averaged parameter

estimates and calculating the proportion of comparisons in which the value for the non-

hybridizing lineage is greater than that of hybridizing lineage. Thus, we obtain empirical P-

values (reported in Supplementary Table S14) in which a value of 0 means in every comparison,

hybridizing lineages were inferred to have rates greater than those of non-hybridizing lineages,

and vice versa. This test is extremely conservative and tests the null hypothesis that non-

hybridizing species always experience diversification rates greater than hybridizing species.

In summary, four phylogenies (all species, sympatric species, plethodontids, & non-

plethodontids) were analyzed using two datasets (narrow & broad definitions of hybridization).

Each data/tree-set combination was tested assuming the three aforementioned differences in

prevalence of hybridization in nature. In all, a total of 32 rounds of model testing were

performed, comparing seven models of trait-independent diversification and seven models of

trait-dependent diversification for each sampling fraction. Further information on how these

data/tree-sets were produced and analyzed may be found in the Supplementary Materials. Lastly,

a description of our test of sensitivity to phylogenetic uncertainty may also be found in the

Supplement (Supplementary Fig. S6 & S7; Supplementary Table S15)

To complement our HiSSE analyses in a manner that is largely insensitive to the

potentially confounding relationship between branch lengths and propensity to hybridize (see

discussion), we conducted sister clade comparisons. Specifically, using all comparisons of three

or more taxa (i.e. two sister species hybridize, and the sister lineage does not), we used the

method of Barraclough, Harvey and Nee[43] to test the hypothesis that hybridizing clades had

greater richness than non-hybridizing clades. To assess confidence, 1000 permutations of

contrast signs were conducted. This test was repeated for each of the eight datasets described

above (Supplementary Table S16).

To test for possible circularity of causality between diversification rates, species richness

and opportunity to hybridize, we quantified the relationships between the three. Specifically, we

tested for a relationship between 1) mean diversification rates and the proportion family

hybridizing, 2) mean diversification rates and species richness, and 3) proportion family

hybridizing and species richness (Supplementary Fig. S8). All analyses were conducted at the

family level, and mean diversification rates were obtained using lineage-specific model-averaged

diversification (speciation, extinction, net-diversification) rates. Simple linear regressions were

conducted in R[44].

Results

Across both datasets (narrow and broad) assuming 20% of species hybridize, three of four

analyzed phylogenies consistently found that hybridizing species experience increased speciation

rates, decreased extinction rates, and therefore increased net-diversification rates (all significant:

Figs. 1, 2 & 3; Tables 1, 2, Supplementary Tables S6 & S7). Net-diversification of hybridizing

lineages in these trees were on average 4X greater than that of non-hybridizing lineages. Our

results were insensitive to phylogenetic uncertainty (Supplementary Fig. S7; Supplementary

Table S15).

215
216   **Figure 1. Inferred speciation rate along the salamander phylogeny for both hybridizing and**
217   **non-hybridizing species.** Branch outlines depict speciation rates corresponding to the inset
218   density plot. Branch interiors depict the probability that a lineage is hybridizable (white) or non-
219   hybridizable (black). The arrow labeled **A** denotes family Plethodontidae, and the arrow labeled **B**
220   denotes the *Plethodon glutinosus* group.
221

Dataset	Tree	Speciation λ		Extinction μ		Net Diversification $r = \lambda - \mu$		Extinction Fraction $\varepsilon = \mu \div \lambda$		Turnover $\tau = \lambda + \mu$	
		H	NH	H	NH	H	NH	H	NH	H	NH
Narrow	All Species	**0.124 ± 5.90e-4**	0.055 ± 1.69e-3	9.39e-3 ± 8.62e-5	**0.038 ± 8.36e-4**	**0.115 ± 5.04e-4**	0.017 ± 2.54e-3	0.076 ± 3.20e-4	**0.794 ± 0.036**	**0.134 ± 6.76e-4**	0.094 ± 8.60e-4
	Sympatric Species	**0.111 ± 8.140e-3**	0.042 ± 1.73e-3	8.57e-4 ± 8.68e-5	**9.45e-3 ± 3.5e-4**	**0.110 ± 8.04e-3**	0.032 ± 2.08e-3	7.54e-3 ± 1.53e-4	**0.283 ± 0.020**	**0.112 ± 8.22e-3**	0.051 ± 1.38e-3
	Plethodontids	**0.098 ± 5.50e-6**	0.041 ± 1.48e-3	4.26e-4 ± 1.20e-8	**0.014 ± 7.66e-4**	**0.098 ± 5.48e-6**	0.028 ± 2.24e-3	4.35e-3 ± 1.63e-7	**0.435 ± 0.051**	**0.098 ± 5.50e-6**	0.055 ± 7.12e-4
	Non-Plethodontids	**0.078 ± 3.38e-3**	0.041 ± 1.73e-3	1.89e-3 ± 4.92e-5	**0.039 ± 7.88e-4**	**0.076 ± 3.42e-3**	2.2e-3 ± 2.5e-3	0.025 ± 1.19e-3	**1.040 ± 0.066**	**8.00e-2 ± 3.32e-3**	0.080 ± 9.80e-4
Broad	All Species	**0.117 ± 2.72e-3**	0.053 ± 2.04e-3	2.39e-3 ± 1.55e-4	**0.027 ± 7.24e-4**	**0.115 ± 2.88e-3**	0.026 ± 2.74e-3	0.021 ± 1.95e-3	**0.647 ± 0.039**	**0.120 ± 2.58e-3**	0.080 ± 1.37e-3
	Sympatric Species	**0.106 ± 4.60e-3**	0.047 ± 1.85e-3	3.02e-3 ± 1.64e-4	**0.027 ± 1.10e-3**	**0.103 ± 4.74e-3**	0.019 ± 2.96e-3	0.031 ± 2.04e-3	**0.732 ± 0.057**	**0.109 ± 4.46e-3**	0.074 ± 7.58e-4
	Plethodontids	**0.143 ± 1.79e-4**	0.049 ± 1.64e-3	3.48e-6 ± 1.33e-6	**1.40e-3 ± 6.66e-5**	**0.143 ± 1.78e-4**	0.048 ± 1.70e-3	2.42e-5 ± 9.24e-6	**0.033 ± 2.26e-3**	**0.143 ± 1.80e-4**	0.051 ± 1.57e-3
	Non-Plethodontids	0.071 ± 5.42e-3	0.072 ± 2.88e-3	1.44e-4 ± 1.04e-5	**1.66e-4 ± 3.72e-7**	0.071 ± 5.42e-3	0.072 ± 2.88e-3	2.38e-3 ± 4.80e-4	**2.60e-3 ± 2.52e-4**	0.071 ± 5.42e-3	0.072 ± 2.88e-3

Table 1. Model-averaged diversification rate estimates at the tips of the phylogeny ± 2 SE. Results assume 20% of extant species hybridize. Bold indicate parameter estimates that differ significantly among character states, with the boldened values as the larger rate estimate. Significance was determined by calculating all possible ratios between non-hybridizing and hybridizing species' model-averaged parameter estimates and calculating the proportion of comparisons in which the value for the non-hybridizing lineage is greater than that of hybridizing lineage. This comparison thus produced an empirical P-value with which significance could be determined.

In all cases of inferred trait dependent diversification, a HiSSE model was the best supported (Supplementary Tables S2-S9). Of these HiSSE models, two groups emerged: one in which hybridizing species did not harbor a second hidden state (All Species – Narrow, Plethodontids – Narrow & Broad), and one in which both hybridizing and non-hybridizing taxa had two hidden states (Table 2). The former of these (without a second hidden state for hybridizing taxa) is interpretable as meaning that there is less diversification rate heterogeneity experienced by hybridizing taxa than by non-hybridizing taxa. Interestingly, non-hybridizing taxa were sometimes inferred to have slightly negative net-diversification rates (Table 2). In the case of the best-fit model using all taxa assuming 20% of species hybridize, this leads to an expected waiting time of 128 million years before the next net-loss in diversity. These negative rates are not persistent however; examination of ancestral state reconstructions indicates that the

hidden state responsible (NH-B) for these rates is distributed primarily along the tips

(Supplementary Figs S9 – S12).

As the average magnitude of increase in speciation rate across all 32 analyses (95% CI:

0.0557 ± 0.0059 species/MY) is significantly greater than the average decrease in extinction rate

(95% CI: 0.0145 ± 0.0037 species/MY), we conclude it is primarily differential speciation that is

driving the increase in net-diversification in hybridizing lineages. Interpreted as a waiting time,

this means that hybridizing species, on average, speciate every 9.2 (95% CI: 8.84 - 9.62) million

years and go extinct every 160.7 (118.97 – 247.96) million years.

Analysis of the tree containing all species recovered strong signal of trait dependent

diversification, with significant differences between hybridization or non-hybridizing lineages

(as measured by empirical P-values) found between states for all five parameter estimates (Figs.

2 & 3; Tables 1, 2, Supplementary Tables S6 – S7), regardless of whether the broad or narrow

criterion was used. Speciation (λ), net-diversification ($r = \lambda - \mu$), and turnover rate ($\tau = \lambda + \mu$)

were greater in hybridizing than non-hybridizing lineages, whereas extinction rate (μ) and

extinction fraction ($\mathcal{E} = \mu \div \lambda$) were lower. Increases were on average 125% (95% CI: 117 -

133%) for speciation, 576% (486 - 699%) for net diversification and 43% (41 - 45%) for

turnover. In contrast, decreases were on average 305% (292 - 317%) for extinction and 945%

(893 - 997%) for extinction fraction (Fig. 2). Additional analyses that assume different

percentages of extant species hybridize produced qualitatively similar results and are discussed

in the supplement (Supplementary Figs. S2-S4, Supplementary Tables S2-14).

Analysis of the tree including only sympatric taxa regardless of the hybridization

criterion (narrow versus broad) datasets recovered patterns identical to those obtained using the

complete phylogeny (Figs. 2 & 3; Tables 1, 2, Supplementary Tables S6-S7). Speciation rate,

net-diversification rate, and turnover increased by 164% (135 - 196%; λ), 244% (199 - 295%; *r*),

and 120% (98 - 142%; τ) respectively in hybridizing versus non-hybridizing lineages. In

contrast, extinction rate and the extinction fraction decreased in hybridizing lineages by 1003%

(864 - 1172%; µ) and 3653% (3319 - 4002%; ε) respectively.

271
Figure 2. Model-averaged lineage-specific diversification rate estimates at the tips of the
phylogeny assuming 20% of species hybridize. Results using different trees are displayed by
column, whereas results for different parameters are displayed by row. Hybridizing lineages (H)
are displayed in red, whereas non-hybridizing (NH) lineages are displayed in blue. Results for both
the narrow and broad datasets are shown; the narrow dataset includes only instances of
contemporary hybridization in nature among natively distributed species, whereas the broad
datasets includes instances of historical introgression and non-natural hybridization.
279
Similarly, plethodontids exhibited the same patterns described above (Figs. 2 & 3; Tables

1, 2, Supplementary Tables S6-S7). Hybridizing lineages experienced rates of speciation, net-

diversification and turnover that were on average 139% (130 - 148%; λ), 250% (224 - 281%; *r*),

and 78% (76 - 81%; τ) greater than those experienced by non-hybridizing lineages. Extinction

rate and extinction fraction were reduced by 3186% (3007 - 3366%; µ) and 9000% (8727 -

11073%; ε) in hybridizing lineages relative to non-hybridizing lineages.

Definition of Hybridization	Dataset	Model	Akaike Weight	Speciation				Extinction				Net Diversification			
				NH-A	*NH-B*	*H-A*	*H-B*	*NH-A*	*NH-B*	*H-A*	*H-B*	*NH-A*	*NH-B*	*H-A*	*H-B*
Narrow	**All Species**	**HiSSE: No H-B, all transitions**	**0.715**	**0.114**	**0.033**	**0.120**	**NA**	**4.24E-7**	**0.041**	**0.010**	**NA**	**0.114**	**-0.008**	**0.110**	**NA**
		HiSSE: No double transitions	0.263	0.025	0.148	0.131	0.181	0.076	0.019	0.007	0.015	-0.051	0.129	0.124	0.166
	Sympatric Species	**HiSSE: All parameters free**	**0.980**	**0.086**	**0.019**	**0.255**	**0.078**	**1.77E-10**	**0.014**	**0.002**	**0.001**	**0.086**	**0.005**	**0.253**	**0.077**
	Plethodontids	**HiSSE: No H-B, all transitions**	**0.523**	**0.063**	**0.009**	**0.099**	**NA**	**1.31E-10**	**0.027**	**2.04E-10**	**NA**	**0.063**	**-0.018**	**0.099**	**NA**
		HiSSE: No H-B, no double transitions	0.475	0.073	0.011	0.097	NA	1.50E-10	0.001	0.033	NA	0.073	0.010	0.064	NA
	Non-plethodontids	**HiSSE: No double transitions**	**0.648**	**0.022**	**0.070**	**0.073**	**0.204**	**0.067**	**0.003**	**0.026**	**0.001**	**-0.044**	**0.068**	**0.047**	**0.203**
		HiSSE: No H-B, no double transitions	0.206	0.089	0.014	0.067	NA	1.83E-10	0.010	1.37E-10	NA	0.089	0.004	0.067	NA
		HiSSE: No H-B, all transitions	0.085	0.017	0.092	0.071	NA	0.031	0.058	0.002	NA	-0.015	0.035	0.069	NA
Broad	**All Species**	**HiSSE: No H-B, all transitions**	**0.631**	**0.106**	**0.026**	**0.065**	**0.126**	**2.19E-10**	**0.025**	**0.003**	**2.59E-10**	**0.106**	**1.00E-4**	**0.062**	**0.126**
		HiSSE: No H-B, no double transitions	0.229	0.089	0.014	0.067	NA	1.83E-10	0.010	1.37E-10	NA	0.089	0.004	0.067	NA
		HiSSE: No H-B, all transitions	0.130	0.028	0.152	0.111	NA	0.024	0.069	0.006	NA	0.004	0.083	0.106	NA
	Sympatric Species	**HiSSE: No double transitions**	**0.637**	**0.027**	**0.075**	**0.099**	**0.194**	**0.044**	**0.013**	**0.005**	**4.00E-10**	**-0.018**	**0.063**	**0.094**	**0.194**
		HiSSE: All parameters free	0.361	0.096	0.024	0.055	0.146	1.98E-10	0.032	1.13E-10	3.01E-10	0.096	-0.008	0.055	0.146
	Plethodontids	**HiSSE: No H-B, no double transitions**	**0.970**	**0.036**	**0.085**	**0.143**	**NA**	**0.002**	**1.74E-10**	**2.95E-10**	**NA**	**0.035**	**0.085**	**0.143**	**NA**
	Non-plethodontids	**CID-2: Three transitions, no double**	**0.949**	**0.014**	**0.082**	**0.014**	**0.082**	**2.86E-11**	**1.69E-10**	**2.86E-11**	**1.69E-10**	**0.014**	**0.082**	**0.014**	**0.082**

Table 2. Best-fit models assuming 20% of species hybridize. Included are models that received >5% Akaike weights for their respective analyses. For each dataset, the best fit model is bold. Maximum-likelihood parameter estimates for speciation, extinction, and net diversification are reported. Non-hybridizing is abbreviated as NH, Hybridizing as H; A and B indicate the two hidden states.

In contrast, analysis of non-plethodontids revealed greater ambiguity as to the impact of hybridization on diversification rates. Whereas the narrow dataset inferred trait-dependent diversification in which hybridization drove increased diversification, the broad dataset did not (Figs. 2 & 3; Tables 1, 2, Supplementary Tables S6-S7). Using the narrow dataset, all rates except turnover were found to differ significantly (Supplementary Tables S6-S7, S13-S14). Speciation rate and net diversification were 90% (75 - 107%; λ) and 3355% (1444 - 26573%; r) greater in hybridizing lineages, whereas extinction rate and extinction fraction were reduced by 1964% (1871 - 2062%; μ) and 4060% (3619 - 4545%; ϵ) in hybridizing lineages relative to non-hybridizing lineages. Analysis of the broad dataset revealed no significant differences and parameter estimates between hybridizing and non-hybridizing species.

Sister clade contrasts broadly supported results of the HiSSE analysis. That is,

305   hybridizing clades were found to have significantly greater species richness than non-hybridizing

306   clades for all datasets except for plethodontid salamanders. Details on significance of these tests

307   may be found in Supplemental Table S16.

308
309
Figure 3. Comparison of model averaged parameter estimates among character states.
Results reported here are those assuming 20% of species hybridize. A, B and C illustrate the
distributions of non-hybridizing to hybridizing lineages diversification rates as estimated at the
tips of the phylogeny using the narrow dataset, whereas D, E and F are those using the broad
dataset. A value > 1 corresponds to a comparison in which non-hybridizing lineages experience
rates greater than those of hybridizing lineages and vice-versa. Dotted vertical lines are placed at
1, at which rates are equal among states.
317

Rapidly diversifying species may have greater opportunity to hybridize due to an

associated increase in species richness. We did not recover evidence supporting this

interpretation. Both mean diversification rate (Adjusted R^2: λ = -0.095; μ = -0.084; r = -0.092)

and proportion of family hybridizing (Adjusted R^2 = -0.112) are decoupled from family species

richness, despite a significant correlation (P: λ = 0.0002; μ = 0.0004; r = 0.0002) between mean

diversification rate and proportion family hybridizing (Adjusted R^2: $\lambda = 0.822$; $\mu = 0.786$; $r =$

0.811; Supplementary Fig. S8).

Discussion

Here, we show strong support that contemporary hybridization is correlated with elevated

diversification rates in the order Caudata. Net-diversification of hybridizing species tends to be

significantly greater than that of non-hybridizing species, driven primarily by a coincident

increase in speciation and decrease in extinction rates (Figs. 2, 3; Tables 1, 2). The accelerated

diversification of hybridizing salamanders appears ephemeral, however; rate differences rapidly

become less pronounced deeper in the tree due to turnover of the hidden states (Fig. 1,

Supplementary Figs. S9-S12). Possible mechanisms leading to this result include frequent range

expansions and contractions (i.e. ref [45]) that have been documented in salamanders (e.g. refs

[46,47,48]) and the process of reinforcement which has long been recognized to contribute to the

diversification process[14,49,50,51]. We outline the potential contribution of each below.

Salamanders often exhibit substantial genetic differentiation at small geographic

distances (e.g., 200m[52]) owing to limited dispersal abilities and low rates of gene flow[53], thus

leading to an abundance of opportunities to evolve in allopatry. Additionally, terrestrial species

such as the *Plethodontid* salamanders of the southeastern United States experience elevational

range expansions and contractions associated with climatic change[48]. Perhaps this combination

of the primarily sessile nature of many salamander species and frequent repeated secondary

contact leads to hybridization occurring regularly across evolutionary timescales. Under these

scenarios, hybridization may then play a creative evolutionary role in the diversification process

similar to that observed in haplochromine cichlids[22,23]. Allopatric speciation of haplochromine

cichlids has occurred in lakes that frequently have dried, split, and reformed, whereas sympatric

speciation has occurred within lakes in which lineages exhibited extreme habitat specificity and

have been reproductive isolated at fine spatial scales[54]. Under these circumstances, hybridization

may have afforded genetic rescue from the consequences of small population size by providing

increased standing genetic variation and thus expedited adaptation to novel stressors, as in Lake

Victorian cichlids post-colonization[23].

There is now clear evidence for latitudinal and elevational range shifts mediated by

climate change[55,56,57,58,59] and a consequent increase in frequency of hybridization among

previously isolated taxa[60,61,62,63]. An informed understanding of the influence of hybridization on

macroevolutionary diversification may thus provide invaluable context for contemporary

processes. This possibility of climate-change mediated hybridization has already been

demonstrated in plethodontid salamanders (*P. shermani* & *P. teyahalee*[48]), as well as in

ecologically divergent subspecies of salamandrid salamanders (*S. salamandra*[64]). Thus, it seems

likely that salamander species worldwide, particularly those found at high elevations due to their

more limited potential geographic distributions, may experience a heightened frequency of

hybridization as climate change advances. While generalizations regarding the outcome of

hybridization should be made with caution[28], our study indicates that perhaps speciation

reversal[65,66,67] need not be the expectation. Rather, our study implies that hybridization may

facilitate adaptation to novel conditions under climate change, leading to diversification of new

salamander lineages.

Here, we show the novel result of a strong correlation of contemporary hybridization with

elevated speciation and net diversification at a large taxonomic scale. However, reinforcement,

defined as the strengthening of prezygotic reproductive isolation in sympatry[14], is intrinsically

intertwined with hybridization. Reinforcement has been documented both experimentally[49] and

observationally[51] to accelerate the initiation and/or completion of the speciation process[50,68,69].

For instance, reinforcement is likely to play an important role in the speciation process due to

strong interspecific sexual selection and mate choice in plethodontids[70]. Indeed, patterns of

sexual isolation among populations of *Plethodon jordani* and *P. teyahalee* match expectations of

reinforcement[14], with sexual selection being stronger in sympatry than in allopatry[71]. Although

we cannot currently quantify the contribution of reinforcement to diversification rate differences

using our data, we urge further research measuring the degree of association between

contemporaneous hybridization and reinforcement among taxa. Nonetheless, were reinforcement

to play a role in the production of the patterns observed in this study, the very occurrence/process

of hybridization would be the ultimate driver (i.e., cannot have reinforcement without

hybridization). Under such a scenario, our study design is well-suited to identify such a signal.

Although a generative role of hybridization is robustly supported across three of our four

datasets, evidence for such a role outside of the Plethodontidae is more limited (Table 1,

Supplementary Tables S2-S14). We find two possible explanations for this finding. Firstly, the

positive association between diversification rates and hybridization may be unique to

Plethodontid salamanders. However, family Plethodontidae is the largest extant family of

salamanders, comprising approximately 2/3rds of the present diversity (471 of 716 species:

amphibiaweb.org). Thus, our observation of hybridization facilitating the diversification process

applies to the majority of salamanders and implies that, at a minimum, contemporaneous

hybridization does not impede the diversification process of extant salamanders.

Secondly, it is highly probable that our analysis of non-plethodontid salamanders is

lacking in power. SSE models have long been known to lose much of their power when dealing

with small number of OTUs (trees < 300 taxa[72]). For example, for trees of 300 species, BiSSE

attains a power of at most 50%, with power dropping below 15% of trees of 100 taxa[72]. Our

phylogeny of non-plethodontids includes only 167 species; that we detected a positive

relationship between hybridization and speciation rates using our narrow (most conservative)

dataset despite such reduced power is a testament to the strength of the signal in our data.

Whereas our larger datasets [complete (469 spp), sympatric (368 spp) and plethodontids (306

spp)] have greater power, our lack of detection of a relationship between hybridization and

diversification rates in non-plethodontids using our broad definition of hybridization is perhaps

unsurprising, given the low power of the analysis (also see Supplementary Materials for an

elaboration of power). Although the power of HiSSE under such scenarios has not been

specifically established, accuracy of parameter estimation does decay with decreasing tree size[38].

Consequently, we cautiously interpret the results of the analysis of non-plethodontids.

Interestingly, sister clade comparisons consistently supported a positive relationship between

hybridization and species richness in non-plethodontids, despite not supporting such a

relationship in plethodontids (Supplementary Table S16). These results are insensitive to branch-

lengths, thereby ameliorating potential concerns related to the relationship between

hybridization, branch-lengths, and diversification rates[21].

Importantly, parameter estimates are largely reasonable. For instance, the greatest

speciation rate inferred by any analysis (Plethodontids assuming 30% of taxa hybridize using the

full tree: Supplementary Table S13), of 0.159 species/million years (MY) can be interpreted as a

waiting time, such that on average, hybridizing species speciate every 6.29 MY. Extinction rates

appear less reliably estimated however; some estimates functionally equal zero, leading to the

large percent decrease in extinction rates observed for hybridizing relative to non-hybridizing

species. In some cases, extinction rates in non-hybridizing taxa leads to negative net-

diversification rates. That being said, averaged extinction rates across all analyses for hybridizing

and non-hybridizing taxa led to more reasonable waiting times of 127.9 and 54.2 MY

respectively. Further, recent studies have documented even more negative net-diversification

rates than inferred herein[73]. Taken together, it appears that extinction plays an important role in

the diversification of salamanders, leading to a reduction in net-diversification rates towards the

present relative to hybridizing species.

422   An important question regarding the interpretation of our results is the relationship

between lineage diversification rates, species richness, and opportunity to hybridize. Because the

relationship between lineage diversification rate and opportunity to hybridize are not necessarily

independent, rapidly diversifying lineages may simply have greater opportunity to hybridize due

to increased diversification rates. Although a legitimate concern, we did not find evidence that

the increased diversification rates we observe are due to increased family-level species richness

leading to increased opportunity to hybridize (Supplementary Fig. S8). Further, it is unlikely that

non-random taxonomic sampling has biased our results, as there is no relationship between clade

specific sampling fraction and frequency of hybridization (Supplementary Fig. S5).

431   Although the ability of methods to accurately infer extinction rates has been debated

432   recently[40,74,75], we emphasize that our results are robust to this concern. Our central result, that

433   hybridizing lineages experience increased net diversification, is driven by both increased

434   speciation rates and decreased extinction rates. Further, in nearly all cases, the magnitude of

435   increase of speciation rate is greater than that of the decrease in extinction. Thus, our results are

436   likely robust even to inaccuracies in the estimation of extinction rate.

437   An important distinction between our study and most previous studies investigating the

influence that hybridization exerts on the diversification process is that of the time-scale at which

hybridization is being assessed. Following Seehausen's[6] landmark paper "Hybridization and

Adaptive Radiation," tests and discussion of his hypothesis, that ancient, widespread

hybridization facilitates adaptive radiation became abundant in the literature (e.g. refs

[16,20,21,76,77]). Whereas much of the subsequent studies focused on *ancient* hybridization, our study

instead focuses on the effects of *contemporary* hybridization.

In a pertinent study, Wiens et al.[21], tested the hybrid swarm hypothesis in the *Plethodon*

glutinosus group (indicated in Fig. 1B) using two nuclear and two mitochondrial genes. They did

not recover strongly supported evidence of genealogical discordance at the base of this group;

these results were interpreted as not being supportive of Seehausen's hypothesis. Further, they

identified a positive relationship between age of species and reproductive isolation. They argue

that the observed relationship between diversification rate and hybridization in this group was a

consequence of this relationship. Although a legitimate concern, we argue that this hybridization

is likely to still have biologically relevant consequences on diversification rates. Specifically,

hybridization may either 1) facilitate the divergence of these young species i.e. through

reinforcement/strengthening of prezygotic isolation, or 2) erode their divergence leading to

species collapse. Whereas the former hypothesis predicts increased speciation rates, the latter

predicts increased extinction rates. We find strong, consistent evidence in favor of the former.

We explicitly tested the hypothesis that contemporary hybridization plays a creative role

in the diversification process in the broadest taxonomic and temporal scale study to date, and our

observations strongly supported the predictions of this hypothesis. Specifically, hybridization

was found to be correlated with both increased speciation rates and decreased extinction rates,

resulting in increased net diversification rates relative to non-hybridizing lineages. Although

other factors certainly contribute to the observed diversification dynamics, we have shown that

hybridization plays a significant role, while accounting for hidden, correlated states in our

analysis. Nearly all studies of hybridization have focused on individual case studies in which

hybridization results in species collapse[66] or promotes diversification in a single species

group[12,13,20,25]. Such studies are necessarily limited in the extent to which their results may be

generalized[28], particularly because results were equivocal across studies. Consequently, we

advocate that our approach can be applied at broad taxonomic and evolutionary timescales to

facilitate robust tests of the role of hybridization in the lineage diversification process. We

anticipate our results are broadly generalizable to animal groups in which homoploid

hybridization occurs because only 17 species of salamanders are known to be polyploid[78], and

our dataset includes only seven (*Ambystoma mexicanum, A. barbouri, A. jeffersonium, A.*

laterale, *A. texanum*, *A. tigrinum*, and *Lissotriton vulgaris*) hybridizing polyploid taxa (none of

which are plethodontids). Our study adds to the growing evidence that hybridization may fuel

rapid diversification and is a compliment to speciation genomics studies characterizing the

genomic basis of this process (e.g. ref [15]). Herein we have shown that hybridization may act as a

generative force across a phylogenetic order, and additional studies at such macroevolutionary

scales are needed to determine if this pattern holds more generally across the tree of life.

Acknowledgements

We thank Adam Leaché, David Weisrock & Maria Servedio for comments on earlier versions of

the manuscript. We also thank Daniel Caetano, Josef Uyeda, Rosana Zenil-Ferguson and Jeremy

Beaulieu for insightful discussions regarding the study and for technical advice. The authors

have no conflicts of interest to declare.

Funding

This work was supported by NSF DEB-1316549 AS, NSF DEB 1208912 to LJH, and the

WSU School of Biological Sciences Elling Endowment Fund to AHP.

Author Contributions

A.H.P. and J.E. conceived of and designed the study, A.H.P. conducted all analyses, and wrote

the manuscript. A.H.P, M.J.M, B.E., J.E., L.J.H, and A.S. contributed to revisions of the

manuscript. A.H.P., J.E., & M.J.M. conducted the literature review and B.E. contributed to early

analyses.

References

1. Barton, N. H. & Hewitt, G. M. Analysis of hybrid zones. *Annu. Rev. Ecol. Syst.* **16,** 113–148 (1985).

2. Darwin, C. *On the Origin of Species by Means of Natural Selection. (Murray.* (1859).

3. Fisher, R. *The Genetical Theory of Natural Selection. (Clarendon Press.* (1930).

4. Mayr, E. *Animal species and evolution. Harvard University Press.* (1963).

5. Futuyma, D. J. On the role of species in anagenesis. *Am. Nat.* **130,** 465–473 (1987).

6. Seehausen, O. Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19,** 198–207 (2004).

7. Grant, P. R. & Grant, B. R. Hybridization of bird species. *Science* **256,** 193–197 (1992).

8. Seehausen, O., Van Alphen, J. J. & Witte, F. Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* **277,** 1808–1811 (1997).

9. Turelli, M., Barton, N. H. & Coyne, J. A. Theory and speciation. *Trends Ecol. Evol.* **16,** 330–343 (2001).

10. Buerkle, C. A., Morris, R. J., Asmussen, M. A. & Rieseberg, L. H. The likelihood of homoploid hybrid speciation. *Heredity.* **84,** 441–451 (2000).

11. Dowling, T. E. & Secor, C. L. The role of hybridization and introgression in the

diversification of animals. *Annu. Rev. Ecol. Syst.* **28,** 593–619 (1997).

12.   Rieseberg, L. H., Archer, M. A. & Wayne, R. K. Transgressive segregation, adaptation and speciation. *Heredity.* **83,** 363 (1999).

13.   Soltis, P. S. & Soltis, D. E. The role of hybridization in plant speciation. *Annu. Rev. Plant Biol.* **60,** 561–588 (2009).

14.   Coyne, J. A. & Orr, H. A. *Speciation. (Sinauer Associates Sunderland.* (MA, 2004).

15.   Marques, D. A., Meier, J. I. & Seehausen, O. A combinatorial view on speciation and adaptive radiation. *Trends in Ecol. & Evol.* 2019.

16.   Mallet, J. Hybrid speciation. *Nature* **446,** 279–83 (2007).

17.   Mavárez, J. & Linares, M. Homoploid hybrid speciation in animals. *Mol. Ecol.* **17,** 4181–4185 (2008).

18.   Gross, B. & Rieseberg, L. H. The ecological genetics of homoploid hybrid speciation. *J. Hered.* **96,** 241–252 (2005).

19.   Schumer, M., Rosenthal, G. G. & Andolfatto, P. How common is homoploid hybrid speciation? *Evolution.* **68,** 1553–1560 (2014).

20.   Mavárez, J. *et al.* Speciation by hybridization in Heliconius butterflies. *Nature* **441,** 868–871 (2006).

21.   Wiens, J. J., Engstrom, T. N. & Chippindale, P. T. Rapid diversification, incomplete isolation, and the "speciation clock" in North American salamanders (genus *Plethodon*): testing the hybrid swarm hypothesis of rapid radiation. *Evolution.* **60,** 2585–2603 (2006).

22.   Keller, I. *et al.* Population genomic signatures of divergent adaptation, gene flow and hybrid speciation in the rapid radiation of Lake Victoria cichlid fishes. *Mol. Ecol.* **22,** 2848–2863 (2013).

23.   Meier, J. I. *et al.* Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat. Commun.* **8,** 14363 (2017).

24.   Grant, P. R. & Grant, B. R. *Darwin's finches on Daphne Major island.* (Princeton University Press, 2014).

25.   MacLeod, A. *et al.* Hybridization masks speciation in the evolutionary history of the Galápagos marine iguana. *Proc. R. Soc. B* **282,** (2015).

26.   Yakimowski, S. B. & Rieseberg, L. H. The role of homoploid hybridization in evolution: a century of studies synthesizing genetics and ecology. *Am. J. Bot.* **101,** 1247–1258 (2014).

27.   Seehausen, O. *et al.* Genomics and the origin of species. *Nat. Rev. Genet.* **15,** 176–192 (2014).

28. Gompert, Z. & Buerkle, C. A. What, if anything, are hybrids: enduring truths and challenges associated with population structure and gene flow. *Evol. Appl.* **9,** 909–923 (2016).

29. Amphibiaweb.org. *Species by the numbers.* (2018). at <http://www.amphibiaweb.org/amphibian/speciesnums.html>

30. Pyron, R. A. Biogeographic analysis reveals ancient continental vicariance and recent oceanic dispersal in amphibians. *Syst. Biol.* **63,** 779–797 (2014).

31. Stamatakis, A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22,** 2688–2690 (2006).

32. Smith, S. A. & O'meara, B. C. treePL: divergence time estimation using penalized likelihood for large phylogenies. *Bioinformatics* **28,** 2689–2690 (2012).

33. Paradis, E., Claude, J. & Strimmer, K. APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* **20,** 289–290 (2004).

34. Highton, R. Speciation in eastern North American salamanders of the genus *Plethodon*. *Annu. Rev. Ecol. Syst.* **26,** 579–600 (1995).

35. Highton, R. Geographic protein variation and speciation in the *Plethodon* dorsalis complex. *Herpetologica* **1997,** 345–356 (1997).

36. Highton, R. Geographic protein variation and speciation in the salamanders of the *Plethodon cinereus* group with the description of two new species. *Herpetologica* **1999,** 43–90 (1999).

37. Highton, R. & Peabody, R. in *The Biology of Plethodontid Salamanders.* (eds. Bruce, R. C., Jaeger, R. G. & Houck, L. D.) 31–93 (Kluwer Academic/Plenum Publishers, 2000).

38. Beaulieu, J. M. & O'Meara, B. C. Detecting hidden diversification shifts in models of trait dependent speciation and extinction. *Syst. Biol.* **65,** 583–601 (2016).

39. Stadler, T. How can we improve accuracy of macroevolutionary rate estimates? *Syst. Biol.* **62,** 321–329 (2013).

40. Beaulieu, J. M. & O'Meara, B. C. Extinction can be estimated from moderately sized molecular phylogenies. **69,** 1036–1043 (2015).

41. Rabosky, D. L. & Goldberg, E. E. Model inadequacy and mistaken inferences of trait-dependent speciation. *Syst. Biol.* **64,** 340–55 (2015).

42. Caetano, D. S., O'Meara, B. C. & Beaulieu, J. M. Hidden state models improve state-dependent diversification approaches, including biogeographical models. *Evolution.* (2018).

43. Barraclough, T. G., Harvey, P. H. & Nee, S. Rate of rbcL gene sequence evolution and species diversification in flowering plants (angiosperms). *Proc. R. Soc. B* **263,** 589–591 (1996).

44. R Core Team *R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing.* (Vienna, 2017).

45. Hewitt, G. M. Some genetic consequences of ice ages, and their role in divergence and speciation. *Biol. J. Linn. Soc.* **58,** 247–276 (1996).

46. Kozak, K. H., Blaine, R. A. & Larson, A. Gene lineages and eastern North American palaeodrainage basins: phylogeography and speciation in salamanders of the *Eurycea bislineata* species complex. *Mol. Ecol.* **15,** 191–207 (2006).

47. Shepard, D. B. & Burbrink, F. T. Phylogeographic and demographic effects of Pleistocene climatic fluctuations in a montane salamander, *Plethodon fourchensis*. *Mol. Ecol.* **18,** 2243–2262 (2009).

48. Walls, S. C. The role of climate in the dynamics of a hybrid zone in Appalachian salamanders. *Glob. Chang. Biol.* **15,** 1903–1910 (2009).

49. Rice, W. R. & Hostert, E. E. Laboratory experiments on speciation: what have we learned in 40 years? **47,** 1637–1653 (1993).

50. Servedio, M. R. & Noor, M. A. The role of reinforcement in speciation: theory and data. *Annu. Rev. Ecol. Evol. Syst.* **34,** 339–364 (2003).

51. Matute, D. R. & Ortiz-Barrientos, D. Speciation: The strength of natural selection driving reinforcement. *Curr. Biol.* **24,** 955–957 (2014).

52. Zamudio, K. R. & Wieczorek, A. M. Fine-scale spatial genetic structure and dispersal among spotted salamander (*Ambystoma maculatum*) breeding populations. *Mol. Ecol.* **16,** 257–274 (2007).

53. Emel, S. L. & Storfer, A. A decade of amphibian population genetic studies: synthesis and recommendations. *Conserv. Genet.* **13,** 1685–1689 (2012).

54. Rico, C. & Turner, G. F. Extreme microallopatric divergence in a cichlid species from Lake Malawi. *Mol. Ecol.* **11,** 1585–1590 (2002).

55. Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421,** 37–42 (2003).

56. Root, T. L., *et al.* Fingerprints of global warming on wild animals and plants. *Nature* **421,** 57–60 (2003).

57. Kelly, A. E. & Goulden, M. L. Rapid shifts in plant distribution with recent climate change. *Proc. Natl. Acad. Sci.* **105,** 11823–11826 (2008).

58. Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid range shifts of species associated with high levels of climate warming. *Science.* **333,**

1024–1026 (2011).

59. Elsen, P. R. & Tingley, M. W. Global mountain topography and the fate of montane species under climate change. *Nat. Clim. Chang.* **5,** 772–776 (2015).

60. Garroway C.J., *et al.* Climate change induced hybridization in flying squirrels. *Glob. Chang. Biol.* **16,** 113–121 (2010).

61. Hoffmann, A. A. & Sgrò, C. M. Climate change and evolutionary adaptation. *Nature* **470,** 479–85 (2011).

62. Chunco, A. J. Hybridization in a warmer world. *Ecol. Evol.* **4,** 2019–2031 (2014).

63. McQuillan, M. A. & Rice, A. M. Differential effects of climate and species interactions on range limits at a hybrid zone: potential direct and indirect impacts of climate change. *Ecol. Evol.* **5,** 5120–5137 (2015).

64. Pereira, R. J., Martínez-Solano, I. & Buckley, D. Hybridization during altitudinal range shifts: nuclear introgression leads to extensive cyto-nuclear discordance in the fire salamander. *Mol. Ecol.* **25,** 1551–1565 (2016).

65. Seehausen, O. Conservation: losing biodiversity by reverse speciation. *Curr. Biol.* **16,** 334-337 (2006).

66. Seehausen, O., Takimoto, G., Roy, D. & Jokela, J. Speciation reversal and biodiversity dynamics with hybridization in changing environments. *Mol. Ecol.* **17,** 30–44 (2008).

67. Vonlanthen, P. *et al*. Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature* **482,** 357–362 (2012).

68. Hoskin, C. J., Higgie, M., McDonald, K. R. & Moritz, C. Reinforcement drives rapid allopatric speciation. *Nature* **437,** 1353–1356 (2005).

69. Pfennig, K. S. Reinforcement as an initiator of population divergence and speciation. *Curr. Zool.* **62,** 145–154 (2016).

70. Arnold, S. J., Reagan, N. L. & Verrell., P. A. Reproductive isolation and speciation in plethodontid salamanders. *Herpetologica* **49,** 216–228 (1993).

71. Reagan, N. L. *Evolution of sexual isolation in salamanders of the genus* Plethodon *(Doctoral dissertation*. (University of Chicago, 1992).

72. Davis, W. C. & Twitty, V. C. Courtship behavior and reproductive isolation in the species of *Taricha* (Amphibia, Caudata). *Copeia* **1964,** 601–610 (1964).

73. Freyman, W. A. & Höhna, S. Stochastic character mapping of state-dependent Diversification reveals the tempo of evolutionary decline in self-compatible *Onagraceae* lineages. *Syst. Biol.* **68,** 505–519 (2019).

74. Rabosky, D. L. Extinction rates should not be estimated from molecular phylogenies. *Evolution.* **64,** 1816–1824 (2010).

75. Rabosky, D. L. Challenges in the estimation of extinction from molecular phylogenies: a response to Beaulieu and O'Meara. *Evolution.* **70,** 218–228 (2016).

76. Kearney, M. Hybridization, glaciation and geographical parthenogenesis. *Trends Ecol. Evol.* **20,** 495–502 (2005).

77. Willis, B. L., van Oppen, M. J., Miller, D. J., Vollmer, S. V & Ayre, D. J. The role of hybridization in the evolution of reef corals. *Annu. Rev. Ecol. Evol. Syst* **37,** 489–517 (2006).

78. Schmid, M., Evans, B.J. & Bogart, J. P. Polyploidy in amphibia. *Cytog. and Gen. Res.*, *145*, 315–330 (2015).

494