Specialization for cell-free or cell-to-cell spread of BAC-cloned HCMV strains is determined by factors beyond the UL128-131 and RL13 loci.

Eric P. Schultz [1,4], Jean-Marc Lanchy [1], Le Zhang Day [1,3,4], Qin Yu [1], Christopher Peterson [1,2,4], Jessica Preece [1], and Brent J. Ryckman[1,2,3,4*]

Division of Biological Sciences[1], Cellular, Molecular and Microbial Biology Program[2], Biochemistry and Biophysics Program[3], Center for Biomolecular Structure and Dynamics[4], University of Montana, Missoula, Montana, U.S.A.

*Corresponding author: Dr. Brent J. Ryckman
Division of Biological Sciences
Interdisciplinary Science Building Rm. 215
The University of Montana
Missoula, MT 59812
Tel: 406-243-6948
Fax: 406-243-4304
Email: brent.ryckman@mso.umt.edu

Running title: Cell-free or cell-to-cell spread of HCMV strains.

ABSTRACT

It is widely held that clinical isolates of human cytomegalovirus (HCMV) are highly cell-associated,

and mutations affecting the UL128-131 and RL13 loci that arise in culture lead to the appearance of a cell-

free spread phenotype. The BAC-clone Merlin (ME), expresses abundant UL128-131, is RL13-impaired and

produces low infectivity virions in fibroblasts, whereas TB40/e (TB) and TR are low in UL128-131, RL13-

intact and produce virions of much higher infectivity. Despite these differences, quantification of spread by

flow cytometry revealed remarkably similar spread efficiencies in fibroblasts. In epithelial cells, ME spread

more efficiently, consistent with robust UL128-131 expression. Strikingly, ME spread far better than TB or

TR in the presence of neutralizing antibodies on both cell types, indicating that ME is not simply deficient at

cell-free spread, but is particularly efficient at cell-to-cell spread, whereas TB and TR cell-to-cell spread is

poor. Sonically disrupted ME-infected cells contained scant infectivity, suggesting that the efficient cell-to-

cell spread mechanism of ME depends on features of the intact cells such as junctions or intracellular

trafficking processes. Even when UL128-131 was transcriptional repressed, cell-to-cell spread of ME was

still more efficient than TB or TR. Moreover, RL13 expression comparably reduced both cell-free and cell-to-

cell spread of all three strains, suggesting that it acts at a stage of assembly and/or egress common to both

routes of spread. Thus, HCMV strains can be highly specialized for either for cell-free or cell-to-cell spread

and these phenotypes are determined by factors beyond the UL128-131 or RL13 loci.

IMPORTANCE

Both cell-free and cell-to-cell spread are likely important for the natural biology of HCMV. In culture, strains

clearly differ in their capacity for cell-free spread as a result of differences in the quantity and infectivity of

extracellular released progeny. However, it has been unclear whether "cell-associated" phenotypes are

simply the result of poor cell-free spread, or are indicative of particularly efficient cell-to-cell spread

mechanisms. By measuring the kinetics of spread at early time points, we were able to show that HCMV

strains can be highly specialized to either cell-free or cell-to-cell mechanisms, and this was not strictly linked

the efficiency of cell-free spread. Our results provide a conceptual approach to evaluating intervention

strategies for their ability to limit cell-free or cell-to-cell spread as independent processes.

INTRODUCTION

Intervention approaches to control human cytomegalovirus (HCMV) infection, including DNA replication inhibitors such as ganciclovir, and vaccines designed to elicit neutralizing antibodies have had limited success (1, 2), and this may be due in part to the complex genetic diversity of HCMV circulating within human populations (3–9). Basic annotations of the 235kbp HCMV genome identify 165 canonical open reading frames (ORFs), although there is evidence of extensive transcription and translation beyond these loci (10, 11). While nucleotide polymorphisms are found throughout the genome, most sequences are well conserved. 21 of the 165 canonical ORFs show high nucleotide diversity and are distributed as islands throughout the genome. Several groups have reported evidence of frequent recombination within the more conserved regions, which may essentially mix and match the more diverse loci into many different genotype combinations (3, 4, 7, 8). Adding to this complexity, some studies have shown evidence of gene inactivating mutations (pseudogenes), and gene deletions (7, 8). While some infected individuals may harbor relatively pure populations of HCMV genotypes, complex, multiple genotype infections, and sequential infections by genetically distinct populations also occur. Cloning of HCMV from clinical samples on bacterial artificial chromosomes (BAC) has resulted in numerous, genetically distinct strains for use in laboratory studies. It is not clear how the genetic and phenotypic differences among BAC-cloned HCMVs might reflect their natural diversity.

The glycoprotein (g) H/L complex is functionally conserved among the herpesvirus family and is involved in the initial receptor engagement and the regulation of the membrane fusion protein gB (reviewed in (12)). The HCMV gH/gL is present in the virion envelope as a gH/gL/gO complex (13, 14), a gH/gL/UL128-131 complex (15–18), and in complex with gB (19). Transient expression of HCMV gH/gL and gB is sufficient to drive cell-cell fusion, but is unclear which forms of gH/gL contribute to membrane fusion during HCMV infection (20). The gH/gL/gO and gH/gL/UL128-131 complexes bind to various cell surface receptors through the gO and UL128-131 domains (21–23) and these interactions are important for entry into a broad range of cell types. gH/gL/gO is likely critical for infection of most, or all cell types, with fibroblasts, epithelial, and endothelial cells being the most extensively studied (24–27). Mutants of HCMV lacking gO are deficient at adsorption to cells and soluble gH/gL/gO blocks such adsorption (25, 28). In

contrast, gH/gL/UL128-131 is dispensable for infection of fibroblasts and neuronal cells but critical for epithelial and endothelial cells, monocyte-macrophages, and dendritic cells (16, 18, 29–31). Soluble gH/gL/UL128-131 did not block virus adsorption, but there is evidence that engagement of receptors can elicit signal transduction pathways that influence the nature of the resulting infection pathway (21, 28, 30). Murine and guinea pig CMVs contain homologous gH/gL complexes that play similar roles in cell tropism, although there are some differences in the reported requirements for the complexes for infection of different cell types (32–37).

　　　　We, and others have noted striking phenotypic variation among the HCMV strains TB40/e (TB), TR, and Merlin (ME) in terms of content of the gH/gL complexes in the virion envelope and the corollary effects on entry and tropism. TB and TR contain gH/gL mostly in the form of gH/gL/gO, and very little gH/gL/UL128-131, whereas ME virions contain overall lower amounts of gH/gL and this is mostly in the form of gH/gL/UL128-131 (27). The genetic polymorphism(s) responsible for these differences are unclear. Murrell et al. described a G>T mutation in the UL128 locus of TB that when engineered into ME, reduced the assembly of gH/gL/UL128-131 through effects on mRNA splicing (38). However, this did not fully explain the observed strain differences, as TR is also low in gH/gL/UL128-131 but is congenic to ME at this locus (27). Zhang et al. showed that expression of gO during replication is lower in ME-infected cells as compared to TR, but again the genetic correlates of this difference were not clearly identified (39). The cell-free infectivity of HCMV strains on both fibroblasts and epithelial cells correlates with the amounts of gH/gL/gO in the virion envelope; TB is by far the most infectious, followed by TR, while ME virions are poorly infectious (27). Repression of the UL131 promoter in ME resulted in virions with dramatically reduced amounts of gH/gL/UL128-131, somewhat higher levels of gH/gL/gO, and dramatically improved cell-free infectivity (27, 40).

　　　　HCMV can spread through monolayer cell cultures by diffusion of cell-free virus in the culture supernatant or by a more direct cell-to-cell mode, but the mechanistic distinctions between these types of spread are not well characterized. Moreover, the pathway of cell-to-cell spread for HCMV has not been extensively studied. There have been suggestions of limited fusion between infected and uninfected cells allowing the transfer of subviral components, but the efficiency of these processes to facilitate the spread of

HCMV infection is not clear (45–47). Syncytium formation has been observed in HCMV infected cultures,

but it is not clear whether these cell-cell fusions contribute to viral spread, or merely represent the

coalescence of late-stage infected cells, long after progeny virus has exited and spread to adjacent cells (48,

49). Different HCMV strains can be characterized as cell-free or cell-associated based on the appearance of

the foci formed in a culture monolayer. For example, ME is considered cell-associated due to formation of

tightly localized foci whereas TB forms more diffuse focal pattern characteristic of cell-free spread (40, 41).

However, it is not clear whether the apparent cell-associated nature of ME simply reflects the lack of efficient

cell-free spread, or a specific mechanism to facilitate efficient cell-to-cell spread. Nor is it clear how strains

like ME and TB compare in their ability to spread cell-to-cell. Finally, the genetic loci responsible for the

apparent cell-free and cell-associated natures of strains are not clear. Some reports have monitored foci

size in the presence of neutralizing antibodies and attributed the cell-associated nature of ME to the

expression of the UL128-131 and RL13 loci (24, 42). It seems clear that either gH/gL/gO or gH/gL/UL128-

131 is required for cell-to-cell spread on fibroblasts, whereas gH/gL/UL128-131 seems to be required for

cell-to-cell spread in epithelial or endothelial cells (24, 26, 43, 44). Here, we report the use of flow cytometry

to compare the spread of HCMV TB, TR, and ME in fibroblasts and epithelial cells at early time points. The

results showed that these strains differ not only in their ability to spread cell-free, but also cell-to-cell and that

neither UL128-131 nor the RL13 are the determinate loci for the mode of spread.

RESULTS

Representative HCMV strains differ not only in their efficiencies of cell-free spread, but also in

their ability to spread cell-to-cell. Focal sizes and patterns can vary considerably among strains of HCMV

and likely reflect differences in their proclivity for cell-free versus cell-to-cell modes of spread.

Representative focal spread patterns of GFP-expressing HCMV strains TB, TR, and ME are depicted in

Figure 1. Cultures of fibroblasts and epithelial cells were infected at low multiplicities and spread was

monitored by fluorescence microscopy over 18 days. In fibroblasts, the spread of all three strains was

localized to small, tight foci for the first 6 days. By day 12, TB and TR began to show signs of more diffuse

spread that continued to increase through day 18, whereas ME foci remained generally smaller and more

localized over the entire experiment (Fig 1A). In contrast, foci in epithelial cell cultures for all three strains

were smaller and more tightly localized throughout the experiment (Fig. 1B). The more diffuse focal

patterns of TB and TR in fibroblasts were likely indicative of efficient cell-free spread, whereas the localized

spread of ME was suggestive of less efficient cell-free spread. This was consistent with our previous

observations that progeny virus released to culture supernatants by TB and TR were far more infectious than

extracellular ME progeny (27). While similar focal spread characterizations have been reported by other

groups (40) (24) (38) (50), such comparisons of focal size and patterns are not easily quantified and do not

clearly distinguish whether smaller, more localized foci form as a passive result of inefficient cell-free spread,

or rather reflect an active preference for a specialized cell-to-cell mechanism. Moreover, while it is clear that

these strains of HCMV can differ in their ability to spread via cell-free progeny, whether they also differ in

their ability to spread cell-to-cell has not been addressed.

For quantitative comparisons of spread among HCMV strains, experiments similar to those described

for Figure 1 were performed, but instead of microscopy, flow cytometry was used to measure the increasing

number of infected (GFP-expressing) cells over the first 12 days of replication.   In fibroblasts, the number of

infected cells for both TB and TR increased exponentially over 12 days, and fit well to a log-linear

regression, indicating a constant rate of spread over the course of the experiment (Fig 2A, B). In contrast,

ME spread considerably faster between days 3 and 6, and then spread slower between days 9 and 12. The

average spread rates (LN GFP+ cells/day as indicated by the regression slopes in Fig. 2B) of all three

strains were within 1.2-fold of each other, with only the difference between TB and TR reaching statistical

significance (Fig. 2C). This was somewhat unexpected, as ME has been described to replicate poorly

compared to other strains (38). Thus, the growth of the smaller, more tightly localized foci of ME were

characterized by initially rapid expansion followed by progressive slowing, whereas the more diffuse foci of

TB and TR were characterized by a constant rate of expansion. These results highlight how visual analyses

of focal size and pattern fail to detect the rapid, early focal expansion of ME. In epithelial cells, spread rates

for all three stains were lower than in fibroblasts, and there were greater differences between strains. ME

was approximately 2-fold faster than TR and 1.5 fold faster than TB, and all differences were statistically

significant (Fig. 2D-F). The more efficient spread by ME on epithelial cells was consistent with the higher

levels of gH/gL/UL128-131 as compared to TB and TR (38) (51) (27), but the tightly localized focal patterns

suggested inefficient cell-free spread by all three strains on this cell type.

Neutralizing antibodies were used to distinguish the contributions of cell-free and cell-to-cell

mechanisms to the rate of spread for each strain.   Antibodies chosen for these experiments were mAb 14-

4b, an anti-gH antibody that likely targets a discontinuous epitope at the membrane proximal region (52,

53), as well as a mixture of rabbit anti-peptide sera that target the epithelial tropism factors UL130 and

UL131 (17). The relative potency of these antibodies to neutralize cell-free TB, TR, and ME was verified in

neutralization experiments shown in Figure 3. Note that on fibroblasts, 14-4b was approximately 10-fold

more potent against ME than for TB and TR, and there was a residual 20% of TR infectivity that was

resistant even at very high antibody concentrations (Fig 3A). On epithelial cells, potency of neutralization by

anti-UL130/131 antibodies was more similar among the strains, and complete neutralization of each was

achieved (Fig. 3B).

Spread rates of each strain in either fibroblast or epithelial cell cultures were then determined in the

presence or absence of anti-gH mAb 14-4b or anti-UL130/131 sera at concentrations sufficient for maximal

neutralization of cell-free virions. In fibroblast cultures, anti-gH 14-4b reduced the spread rate of TB and TR

by 70% and 55%, respectively, whereas ME spread was reduced by only 25% (Fig. 4 A-C). The apparent

resistance of ME spread was especially noteworthy since cell-free ME was more sensitive to neutralization

than either TB or TR (Fig. 3A). In all cases, the apparent antibody resistant spread was greater than the

spread in the presence of ganciclovir, even for TR, which is known to harbor resistance mutations in the

UL97 kinase (54). This demonstrated that the apparent antibody resistant spread was not simply the

transfer of cytoplasmic contents from infected to uninfected cells, but rather represented *bona fide* viral

spread dependent on the production of new viral genomes. These results supported the hypothesis that

spread of TB in fibroblasts predominately involved cell-free virions that were highly sensitive to neutralizing

antibodies, whereas ME spread predominately by a distinct, antibody resistant cell-to-cell mechanism. The

intermediate antibody sensitivity of TR spread might indicate intermediate contributions of cell-free and cell-

to-cell spread mechanisms. However, it may also reflect the lack of complete neutralization of cell-free TR

as described in Fig 3A. In epithelial cell cultures, the presence of anti-UL130/131 antibodies resulted in 20-

30% reductions in spread rates for all three strains, indicating predominantly cell-to-cell modes of spread in

this cell type (Fig 4 D-F). Again, ME spread was most efficient in the presence of neutralizing antibodies,

despite TB and TR displaying similar resistance over 12 days. While virions derived from fibroblasts are well

characterized (51) (27), less is known about virions produced in epithelial cells. To this end, virion quantity

and infectivity generated from epithelial cells infected with TB, TR, or ME was determined using subcellular

fractionation, and compared to fibroblasts.

HCMV ME is uniquely efficient at cell-to-cell spread via a mechanism that depends on the

intact cell monolayer, and overcomes the low infectivity of progeny virus. The efficient cell-free spread

of TB in fibroblasts was likely a function of the quantity and infectivity of progeny virus release to culture

supernatants. Similarly, it was expected that highly efficient cell-to-cell spread of ME would correlate with

the quantity and/or infectivity of progeny virus within the infected cells, or "cell-associated virus". To test

these hypotheses, infected cultures were separated into supernatants and cells, and the cells further

fractionated into nuclear, and cytoplasmic portions. This fractionation allowed for more detailed comparisons

of DNA replication and nuclear egress between strains. The quantity and infectivity of progeny virus within

the infected cells and in the supernatants were then determined, and results shown in Figure 5.

In fibroblasts, total intracellular viral genome accumulation was comparable for all three strains (Fig

5A). The vast majority was in the nuclear fraction, likely in the form of non-infectious, unpackaged genomes

and non-tegumented, non-enveloped nucleocapsids. The number of viral genomes in cytoplasmic and

supernatant fractions was slightly lower for TB-infected fibroblasts than for either TR or ME. At least a

portion of these cytoplasmic and supernatant genomes would be expected to represent the infectious

progeny that could contribute to either cell-to-cell or cell-free spread, respectively. Despite the lower

numbers of progeny genomes for TB in cytoplasmic and supernatant fractions, these corresponded to log-

folds greater cell-associated (i.e., sonically disrupted cells) and cell-free (i.e., supernatant) infectivity per cell

as compared to TR or ME (Fig. 5B). The greater infectivity of TB, despite lower virion numbers was in

agreement with our previous measures of lower specific infectivity ratios (genome/PFU) for TB compared to

TR and ME (27), and was positively correlated with the relative efficiencies of cell-free spread for these

strains. Surprisingly, cell-to-cell spread was inversely correlated to the levels of cell-associated infectivity,

with cell-to-cell spread of ME being highly efficient despite producing the lowest detectable cell-associated

infectivity.

In epithelial cells, accumulation of whole cell, cytoplasmic, and nuclear progeny genomes were

similar among all three strains (Fig. 5C). In contrast, ME released nearly 10-fold more progeny genomes to

culture supernatants than did TB, and TR was by far the least efficient at releasing progeny to supernatants.

These results suggest major differences among strains in the later assembly and release stages of the

replication cycle in epithelial cells. As was observed in fibroblasts, ME had the lowest cell-associated

infectivity in epithelial cells, despite the more efficient cell-to-cell spread in this cell type (Fig 5D and 4F).

Infectivity in epithelial cell culture supernatants mirrored the quantities of genomes, indicating that the

epithelial-derived virions of all three strains are of comparable specific infectivity. This was in stark contrast

to the drastic differences in specific infectivity among these strains when produced in fibroblasts (Fig 5 A and

B) (27).

Taken together, these results confirmed that the robust cell-free spread of TB and TR in fibroblasts

was associated with the release of highly infectious progeny virus to the culture supernatant and that the

poor cell-free spread of ME was due to the low infectivity of the cell-free progeny. Surprisingly, ME was

much more efficient at cell-to-cell spread than TB despite having far less intracellular infectious progeny than

TB or TR. In epithelial cell cultures, the cell-free characteristics for all of the strains were more similar to one

another than they were on fibroblasts, and this was consistent with a comparably small contribution of cell-

free spread for all three strains in this cell type. Thus, spread by all three strains in epithelial cells was

predominantly cell-to-cell, and ME was by far the most efficient whereas TR was notably inefficient. The

cell-to-cell spread of ME thus depends on efficient trafficking of progeny to adjacent cells, and this seems to

overcome the poor infectivity of the intracellular progeny.

The highly efficient cell-to-cell spread mechanism of HCMV ME is not determined by the high

expression of UL128-131. The cell-associated nature of ME has been linked to the high expression levels

of UL128-131 and the corollary poor infectivity of cell-free ME virions (24, 27, 40, 42). To address this in our

quantitative spread assay, we made use of the previously described tetracycline (Tet) repression system

developed by Stanton et al. (40). Briefly, the BAC clone of ME used in these studies contained tetracycline-

operator (TetO) sequences in the promoter of UL131. We previously showed that replication of this

recombinant ME in HFFFtet fibroblasts, which express the tetracycline-repressor (TetR) protein, produced

extracellular virus with dramatically reduced gH/gL/UL128-131, slightly higher gH/gL/gO and as a result,

greatly improved cell-free infectivity (27). We refer to ME propagated under conditions of UL131 repression

as "Merlin-T" (MT).

Direct comparisons of ME spread rates in nHDF the HFFFtet cells were complicated by the fact that

in addition to expressing TetR, HFFFtet were also telomerase immortalized and this could have independent

effects on the HCMV replication cycle. Moreover, since we also wished to address the role of UL128-131

expression on spread in epithelial cells, two new TetR-expressing cell lines were generated; nHDFtet and

ARPEtet.   A luciferase reporter assay demonstrated efficient repression of TetO-containing promoters in

each of these cell lines (Fig 6A). Consistent with our previous reports, ME virions produced from nHDF

contained high levels of gH/gL/UL128-131 and low amounts of gH/gL/gO, whereas MT virions produced in

HFFFtet contained far less gH/gL/UL128-131 but more gH/gL/gO (Fig 6B) (27). MT produced in the new

nHDFtet cell line showed a similar reduction of gH/gL/UL128-131 and increase in gH/gL/gO, and MT virions

derived from either tetR-expressing fibroblasts cell line were dramatically more infectious than ME derived

from nHDF cells (Figs 6 B and C). ME virions produced in control ARPE cells were gH/gL/UL128-131-rich

and, as in fibroblasts, Tet-repression dramatically reduced amounts of gH/gL/UL128-131 in epithelial-derived

MT virions (Fig 6B). However, unlike fibroblast-derived MT, epithelial-derived MT did not contain increased

gH/gL/gO and the infectivity was actually reduced (Figs 6B and C).

In nHDF fibroblasts, MT spread slightly faster than ME, consistent with a gain in the ability to spread

cell-free as a result of the improved virion infectivity (Fig 7A). However, in the presence of neutralizing

antibody 14-4b, the spread of ME and MT was indistinguishable. Similar experiments were performed with

HFFFtet cells, which generate MT. The high-gH/gL/UL128-131 condition of ME was generated by addition

of doxycycline to the cultures to block the TetR repression of UL131 (Fig 6A). Spread rates were generally

lower in HFFFtet cells compared to nHDF cells, but as expected, MT spread more efficiently than ME, even

in the presence of neutralizing antibodies (Figs 7 A and B).   Thus, in both fibroblast systems, the efficiency

of ME cell-to-cell spread was not dependent on the high levels of gH/gL/UL128-131. Nor was the apparent

cell-to-cell spread a default mechanism of poor cell-free spread, and these results demonstrate a previously

unappreciated distinction of the mechanisms of cell-free and cell-to-cell spread.

In epithelial cells, repression of gH/gL/UL128-131 resulted in a decrease in average spread rate

(from 0.38 to 0.29), and small increase in the sensitivity of that spread to neutralizing anti-UL130/131

antibodies (from 10% to 30% inhibition) (Fig 7B). The reduction in spread rate was consistent with the

important role of gH/gL/UL128-131 for HCMV in these cells. One interpretation of the small increase in

antibody sensitivity was a shift towards increased cell-free spread. However, this seemed unlikely since

epithelial-derived MT were actually less infectious than epithelial-derived ME (Fig 6C). Thus, it seems more

likely that the observed spread of MT was still predominantly cell-to-cell, but that the reduced gH/gL/UL128-

131 allowed for more efficient blocking of the cell-to-cell spread by these antibodies. It was notable that the

spread rates of MT in epithelial cells were comparable to those of TB both in the presence and absence of

neutralizing antibodies (compare Figs 4D and 7B). This was consistent with the notion that both strains

spread in epithelial cells predominantly via cell-to-cell spread over the first 12 days. Thus, while the high-

levels of gH/gL/UL128-131 of ME clearly enhance the efficiency of the spread in epithelial cells, this does not

appear to determine the mechanism, and these results suggest other intrastrain variable factors can affect

cell-to-cell spread in this cell type.

The RL13 glycoprotein tempers both cell-free and cell-to-cell spread. The RL13 ORF encodes

an envelope glycoprotein that has been described as a selective, or preferential inhibitor of cell-free spread

over cell-to-cell spread inasmuch as genotypes with inactivating mutations in RL13 arise during serial

propagation of HCMV in both fibroblasts and epithelial cells, and this has been correlated with the

appearance of more cell-free spread characteristics (40, 50, 55). While the ME recombinant used in the

present studies harbors such an inactivating RL13 mutation, the RL13 expression status of the TB and TR

BAC-clones was unclear. DNA sequencing confirmed that the RL13 ORF of both TB and TR was intact

(data not shown), but lack of quality antibodies precluded confirmation of RL13 glycoprotein expression.

Thus, to directly and fairly compare the effects of RL13 on the spread of all three strains, and to avoid

potential selection of RL13 mutants in TB and TR during spread experiments, fibroblast and epithelial cell

lines that express RL13 were engineered. Immunoblot analysis demonstrated that these cell lines

expressed of both mature and immature proteoforms, and flow cytometry confirmed similar expression levels

(Fig 8).

In fibroblasts, RL13 expression resulted in a comparable 40-50% reduction in spread rate for all three

strains (Fig 9A). The relative effect of neutralizing antibodies on spread by TB or TR was unaltered by RL13

expression, but spread of ME was 2-fold more sensitive to neutralizing antibodies in the RL13-expressing

fibroblasts. In epithelial cells, RL13 expression had no effect on the spread of TR but resulted in a modest

20% reduction for both TB and ME, although only for ME was this statistically reproducible (Fig 9B). Effects

of RL13 expression on antibody sensitivity were also different in epithelial cells than in fibroblasts. RL13

expression increased the sensitivity of TB spread to antibody inhibition, but had little effect on TR or ME (Fig

9B). The notion that RL13 acts selectively or preferentially to restrict cell-free spread would have predicted

that a greater fraction of the observed spread in RL13-expressing cells would be cell-to-cell and therefore

less sensitive to neutralizing antibodies. On the contrary, results showed that sensitivity of spread to

neutralizing antibodies was either unaffected or enhanced in RL13-expressing cells compared to the control

cells. Thus, RL13 appears to exert its effect on HCMV spread in a manner that affects both the cell-free and

the cell-to-cell mechanisms, not preferentially cell-free spread.

RL13 exerts strain-dependent and cell type-dependent effects on the quantity and infectivity

of progeny virus released to culture supernatants. Two plausible mechanisms by which RL13 expression

could negatively affect the cell-free mode of spread include 1) reducing the numbers of cell-free progeny

released to culture supernatants, and 2) reducing the infectivity of the cell-free progeny released. Both

possibilities were tested using the RL13-expressing fibroblasts and epithelial cells. In fibroblasts, there were

modest reductions in the numbers of progeny released to culture supernatants for TB and ME (Fig 10A).

There were also modest reductions in detectable infectivity in culture supernatants of the RL-13 expressing

cells (Fig 10B.) Taken together, the correlation between reduced numbers and reduced infectivity suggest

that the specific infectivity of cell-free virions was not substantially affected by RL13-expression. For the

efficient cell-free spreaders TB and TR, these reductions in cell-free progeny quantity and/or infectivity may

help explain the reductions in spread rates described above. In contrast, the reduction of cell-to-cell spread

for ME could not be attributed to reduced cell-free progeny quantity or infectivity.

In epithelial cells, there were no effects on the numbers of supernatant progeny released for TB or

TR, but there were actually more cell-free ME progeny produced from RL13-expressing cells (Fig. 10C).

Despite the greater numbers of ME progeny in RL13-expressing epithelial cell supernatants, there were

drastically lower levels of infectivity, indicating a substantial reduction in the per virion infectivity (Fig. 10D).

However, as was noted above for fibroblasts, since the spread of ME in epithelial cells was determined to be

predominately cell-to-cell, these effects on cell-free virus may not fully explain how RL13 tempers spread in

this cell type.

The drastic reduction of ME cell-free infectivity due to RL13 expression in epithelial cells suggested

effects on the envelope glycoproteins. In particular, gH/gL/UL128-131 is the only envelope glycoprotein

currently known to be specifically required for efficient infection of epithelial cells. Thus, ME virions produced

by RL13-expressing or control fibroblasts and epithelial cells were analyzed by immunoblot for the levels of

gH/gL complexes. Consistent with our previous reports (Fig. 6B, and (51)), ME virions produced by

fibroblasts had gH/gL predominately in the form of gH/gL/UL128-131 (Fig 11A). Strikingly, epithelial-derived

ME contained substantially greater amounts of gH/gL/UL128-131 than fibroblast-derived ME (Fig 11A). In

both cell types, RL13 expression resulted in a slight reduction in gH/gL/UL128-131, but the reductions were

too small to provide a compelling explanation for the effects on virion infectivity observed. Levels of gB were

also analyzed as a control, but surprising results were obtained. Antibodies used to detect gB were specific

for both the 55 kDa portion of the furin-cleaved proteoform, and the the full length, uncleaved 165-170 kDa

form and (56–58). In fibroblasts, ME virions contained almost entirely furin-cleaved gB, as evidenced by

detection of only a 55kDa band (Fig 11B). This was not affected by RL13-expression. In contrast, epithelial-

derived ME contained mostly uncleaved gB, and RL13-expression drastically reduced the amount of this

proteoform (Fig 11B). Together, these results indicated that RL13 can affect the processing and

incorporation of gH/gL complexes and gB into cell-free progeny, albeit in a cell type-dependent manner .

DISCUSSION

Distinctions between cell-free and cell-associated phenotypes of different HCMV strains and isolates

have been noted, but the importance of different genetic loci involved in these spread modes remains to be

understood. Sinzger et al. showed that during the initial rounds of cell-associated subculturing in fibroblasts,

some clinical HCMV isolates formed small and tightly packed foci, but tended towards larger and more

diffuse foci during later rounds of passage (59). The change in foci morphology coincided with the

appearance of infectious virus in culture supernatants and the basic interpretation was that the adaptation of

clinical HCMV isolates to the cell culture environment involved a shift from predominately cell-to-cell spread

to cell-free spread. A more recent report by Galitska et al. noted considerable variation among clinical

HCMV samples, with some displaying the characteristic small foci, but others showing larger diffuse foci

even in early rounds of subculturing (60). This variation fits well with the growing evidence of genotypic

diversity of HCMV populations within natural populations of HCMV (3, 4, 6, 8, 9) and with the earlier

suggestion by Sinzger that the observed culture adaptation of HCMV clinical isolates might represent

random sampling or purifying selection of preexisting genotypic variants, rather than the acquisition of *de*

novo mutations (59). In line with this view, Subramanian et al. used a flow cytometry method to determine

the increase in HCMV infected cells over time to better quantitate spread kinetics and found that some low

passage clinical isolates spread more rapidly than others (61). They interpreted slow kinetics to represent a

cell-associated-only phenotype and fast kinetics to represent a phenotype of both cell-associated and cell-

free spread. Indeed, the faster spreading isolates were more sensitive to the presence of neutralizing

antibodies. Together, these and other studies suggest considerable variation in the modes of spread

exhibited by genetically distinct HCMV, but the exact genetic variations that determine these phenotypes

remain unclear.

Here we applied a flow cytometry-based approach to compare the spread characteristics of

three commonly studied and genetically distinct BAC clones of HCMV (TB, TR, and ME) in both fibroblast

and epithelial cell cultures. It has been widely appreciated that TB and TR exhibit characteristics of cell-free

spread, whereas ME is essentially restricted to cell-to-cell spread. Previous studies suggested that the

distinction between these strains in the ability to spread cell-free in fibroblast cultures is due largely, if not

entirely to the fact that cell-free TB and TR virions are log-folds more infectious than ME virions and that cell-

free spread of ME in fibroblast cultures can be enhanced by transcriptional repression of

epithelial/endothelial cell tropism factor gH/gL/UL128-131, which in turn leads to improved infectivity of cell-

free ME virions (40) (38) (27). However, it has remained unclear whether the apparent cell-associated

nature of ME is due simply to the poor cell-free infectivity, or reflects a specific ability to facilitate more

efficient cell-to-cell spread than viruses like TB and TR that can spread efficiently cell-free. An important

corollary consideration for our studies was that the UL128-131 and RL13 loci have been implicated as

important for determining the cell-free or cell-associated nature of HCMV strains (24, 27, 40, 42) (40, 50,

55). The results presented here demonstrate that the cell-associated characteristic of ME is not simply due

to the poor cell-free infectivity. Rather, ME is capable of a specifically efficient cell-to-cell mechanism of

spread, TB is specifically inefficient at cell-to-cell spread, and TR may be of intermediate cell-to-cell spread

capacity. Moreover, we demonstrate that cell-free and cell-associated spread characteristics are not

determined by either the expression levels of UL128-131 or by RL13 expression, although these loci do

influence the efficiency in strain-dependent and cell type-dependent manners. Thus, we suggest the

involvement of other variable loci in distinct mechanisms of cell-free and cell-to-cell spread, and that in

epistatic relationships among these loci govern the efficiency of these processes.

One candidate for a variable factor that might influence cell-free and cell-to-cell spread in an epistatic

manner is gO, which is one of the more diverse proteins encoded by HCMV. There are 8 alleles of the

UL74(gO) ORF that vary between 10-30% of predicted amino acids (62, 63) (64) (51). Mutational inactivation

of UL74 in TR and TB dramatically reduced cell-free infectivity, but allowed cell-to-cell spread, albeit reduced

(26) (65). In contrast, spread of a gO_null ME was indistinguishable from the parental (24). Together these

observations suggested that gH/gL/gO is critical for cell-free spread, but cell-to-cell spread can be mediated

by gH/gL/UL128-131 alone. However, it has more recently been reported that viruses lacking gH/gL/UL128-

131 can spread cell-to-cell, but is dependent on the gH/gL/gO receptor, PDGFRa (43). Furthermore, a

recent unpublished study by our group demonstrated that switching the UL74 allele in both TR and ME

backgrounds could influence the efficiency of cell-to-cell spread, demonstrating that there were epistatic

phenomena involved inasmuch as the influence of some UL74 alleles were observed in only one of the two

genetic backgrounds (66).

Our finding that the level of gH/gL/UL128-131 expression did not affect the spread rate of ME in

fibroblasts conflicts with a report by Stanton et al., which showed UL128-null ME plaques significantly larger

than UL128-intact ME plaques (40). The apparent contradiction may be explained by the fact that our flow

cytometry assays measured spread rates averaged over the first 12 days, as compared to the

measurements of 21-day plaque sizes by Stanton et al. Linear regression analyses of the data collected by

flow cytometry indicated that ME spread at a faster rate between days 3 and 6, and then began to slow after

day 9, whereas the constant rate of cell-free spread demonstrated by TB and TR (Fig. 2). The slowdown

displayed by ME may simply reflect a limiting number of potential uninfected target cells available via cell-to-

cell spread during focal expansion. Alternatively, it could be that the cells at the periphery of an expanding

ME foci rapidly become less permissive, for example via activation of the antiviral state. In any case, the

increase in 21 day plaque size of a UL128-null ME is consistent with enhanced cell-free infectivity, but our

analysis of spread rates over 12 days indicates this does not come at the expense of a distinct and

comparably efficient mode of cell-to-cell spread. This interpretation is supported by the observation that the

12 day spread rate of ME was highly resistant to neutralizing antibodies regardless of gH/gL/UL128-131

expression level (Fig. 7) and is consistent with Laib Sampaio et al., who showed much greater cell-free

dispersion of UL128-null ME foci, but more similar antibody resistant, cell-associated spread with or without

UL128 (24).

In epithelial cells, repression of gH/gL/UL128-131 did reduce the 12 day spread rate for ME (Fig 7).

This was consistent with reduced plaques sizes of UL128-null ME documented by Murrell et al., and likely

reflects the important role of gH/gL/UL128-UL131 in epithelial cells (38). However, as observed in

fibroblasts, spread in epithelial cells was still highly resistant to neutralizing antibodies when gH/gL/UL128-

131 was repressed. This was different than Murrell et al., who concluded that spread of ME in epithelial cells

was more sensitive to antibodies when gH/gL/UL128-131 was reduced (42). Again the discrepancy may

indicate that analyses of 21-day plaque size likely accentuate the influence of cell-free infectivity and masks

the contribution of cell-to-cell spread. Thus, by measuring average spread rates over 12 days, we find that

gH/gL/UL128-131 levels are not the sole determinate of the cell-free versus cell-associated spread

phenotype for ME. Rather, the increase in cell-free infectivity due to reduction in gH/gL/UL128-131 and the

concomitant increase in gH/gL/gO, is independent of the highly efficient cell-to-cell mode of spread.

The RL13 glycoprotein was another candidate for a factor influencing the cell-free and cell-

associated phenotypes. Previous reports have suggested that expression of RL13 limits HCMV replication

in fibroblast, epithelial, and endothelial cell cultures through preferential effects on cell-free spread (24, 40,

50, 55). On the contrary, we find that RL13 tempers HCMV spread by either cell-free or cell-to-cell

mechanisms and that this temperance was more pronounced in fibroblasts compared to epithelial cells (Fig.

9). Interestingly, RL13 expression dramatically reduced the infectivity of cell-free virions released by ME

from epithelial cells, and this was consistent with the previous report showing low cell-free titers for RL13-

intact ME in epithelial cell cultures compared to RL13 mutant ME (40). The physiological basis of this

infectivity loss remains unclear, but it correlated with the observations that ME virions produced in RL13-

expressing epithelial cells contained less full length, non-furin cleaved gB (Fig. 11). Strive et al. showed that

furin cleavage was not required for the function of HCMV gB (67). Therefore, while the change in gB may

not directly explain the change in infectivity, it is conceivable that RL13 could affect the processing and

incorporation of other important envelope glycoproteins.

Our results may help to explain inconsistencies regarding the selection of UL128-131 and RL13

mutations during laboratory culturing of HCMV isolates, phenomenon that has been universally ascribed to

HCMV, despite clear strain variability. While ME is highly sensitive to the selection of *de novo* UL128-131 in

fibroblast cultures, TB and TR are remarkably stable in this respect (40, 50, 59). This disparity might simply

reflect that many subculturing methods that including supernatant transfers, cell sonication, and high split-

ratio transfer of intact infected cells, would be expected to heavily favor cell-free spread. Thus, given the

highly efficient cell-to-cell spread of ME reported in this study, propagation by low split-ratio passage of

intact, infected cells might reduce the selective bottleneck on loss of UL128-131 function to favor cell-free

spread. Likewise, RL13 inactivating mutations have also been reported for many strains and clinical isolates

in studies involving fibroblasts, epithelial, and endothelial cells (10, 55). The BAC clones of TB and TR have

intact RL13 ORFs, but since expression of RL13 from transduced cells reduced replication by these strains,

it may be that both strains carry mutations or polymorphisms in promoter or other regulatory sequences that

impact RL13 expression. Moreover, our results suggest that the selection against RL13 may be more

extensive in fibroblasts but less pronounced in epithelial cells.

Notably, a recent report showed that the presence of neutralizing antibodies during subculturing of

clinical urine-derived HCMV in fibroblasts stabilized the UL128-131 and RL13 loci in the resultant virus

populations (68). If these clinical isolates were similar to ME in their proclivity for cell-to-cell spread, it would

seem logical that neutralizing antibodies would reduce the selective advantage of cell-free infectivity

conferred by UL128-131 mutation. Indeed, it is interesting that ME was also isolated from a clinical urine

sample (10). However, the preservation of RL13 seems less intuitive since RL13 impacts both cell-free and

cell-to-cell modes of spread. Given the variable effects we observed of RL13 on different strains, these

observations again seem to point to epistasis phenomena, where the relative effect of any given perturbation

at one variable locus is influenced by variable physiology related to other polymorphic loci.

Our data also shed new light on the notion that the cell type might influence phenotypic properties in

a non-genetic manner through changes to the protein composition of the progeny virus. Among the

herpesviruses, these so called "producer cell effects" have been particularly well characterized for EBV,

where the mechanism alternates the tropism of progeny virus between epithelial and B-cells (reviewed in

(69)). Scrivano et al. suggested a similar phenomenon for HCMV based in part on data showing that

progeny virus produced from endothelial cells had less UL128 protein in the envelope compared to fibroblast

derived progeny (35). This is in stark contrast to our results that epithelial-derived HCMV contained greater

amounts of gH/gL/UL128-131 than fibroblast-derived virus (Fig 9). There are several possible, non-mutually

exclusive explanations for this discrepancy. This could reflect differences between strains as the prior study

used TB, whereas our analyses used ME. There could also be differences between endothelial and

epithelial cells. Finally, the Scrivano et al. analysis normalized virion immunoblots to gB, whereas we used

major capsid protein (MCP). It is possible that incorporation of gB into progeny virus is affected by

glycoprotein processing/secretory pathway differences between cell type, whereas plasticity in the number of

subunits per capsid subunits seems less likely. Indeed, we observed differences in the proteolytic

processing and virion incorporation of gB when ME was produced in fibroblast and epithelial cells.

In conclusion, we have characterized very different cell-free and cell-associated spread phenotypes

among three commonly studied BAC clones of HCMV. Our data suggest that these phenotypic variances

are manifest through the combined influenced of diversity at multiple genetic loci other than UL128-131 and

RL13. How this phenotypic variation is represented *in vivo* remains unclear. There is little clear evidence to

support the notion that HCMV is predominately cell-associated *in vivo*. The endemic nature, and the

pleomorphic disseminated disease presentations suggest that HCMV is able to thrive in many different

bodily environments as it spreads within and among individual human hosts. Copious amounts of cell-free

virus are shed in urine, saliva, and breast milk, and this is likely a major route of transmission between

individuals, whereas HCMV in the blood is likely highly associated with leukocytes (70-72). Thus, it seems

likely that both cell-free and cell-associated modes of spread are important to the natural history of HCMV.

MATERIALS AND METHODS

Cell lines. Primary neonatal human dermal fibroblasts (nHDF; Thermo Fisher Scientific), RL13-nHDFs

(nHDFs transduced with lentiviral vectors encoding RL13 of HCMV strain Merlin, selected with puromycin

resistance), nHDF-tet (nHDFs transduced with retroviral vectors encoding the tetracycline repressor protein,

selected with puromycin resistance, MRC-5 fibroblasts (American Type Culture Collection; CCL-171), and

HFFF-tet (40) (provided by Richard Stanton, Cardiff University, United Kingdom) were grown in Dulbecco's

modified Eagle's medium (DMEM)(Sigma) supplemented with 6% heat-inactivated fetal bovine serum (FBS)

(Rocky Mountain Biologicals, Missoula, MT, USA) and 6% Fetalgro® (Rocky Mountain Biologicals, Missoula,

MT, USA). Retinal pigment epithelial cells (ARPE19)(American Type Culture Collection, Manassas, VA,

USA), ARPE-tet (ARPE19 cells transduced with retroviral vectors encoding the tetracycline repressor

protein, and selected by puromycin resistance), and RL13-ARPE (transduced with lentiviral vectors encoding

RL13 of HCMV strain Merlin, selected with puromycin resistance) were grown in a 1:1 mixture of DMEM and

Ham's F-12 medium (DMEM:F-12)(Gibco) and supplemented with 10% FBS.

Retroviral vectors. The Tet repressor protein bearing a nuclear localization signal was extracted by PCR

from the integrated sequence in the HFFF-tet chromosomal DNA and cloned into the same pMXs retrovirus

vector used to construct HFFF-tet (40). The pMXs retrovirus vector plasmid was a gift from Dr. Toshio

Kitamura at the Institute of Medical Science, University of Tokyo (73). The tet-containing vector plasmid

was transformed in 293T cells together with the pUMVC and pMD2.G helper plasmids. The pUMVC helper

plasmid was a gift from Bob Weinberg (Addgene plasmid # 8449) (74). Two days after transformation, the

retroviral particles in the supernatant were purified from cell debris through syringe filtration and

centrifugation. After titration, the particles were used to transduce low passage nHDF or ARPE-19 cells.

After a week of puromycin selection, cells aliquots were tested for tetR expression after transfection of a

firefly luciferase tetR-reporter system and aliquots were stored in liquid nitrogen until further usage. The

codon-optimized RL13 with an intact ORF from Merlin HCMV strain (NCBI ref YP_081461) was constructed

by Gibson Assembly and used to replace the eGFP ORF in the pLJM1-EGFP lentiviral transfer vector

plasmid. The pLJM1-EGFP plasmid was a gift from David Sabatini (Addgene plasmid # 19319) (75). The

RL13-containing vector plasmid was transformed in 293T cells together with three lentiviral helper plasmids.

The pMDLg/pRRE, pRSV-Rev, and pMD2.G helper plasmids were a gift from Didier Trono (Addgene

plasmids # 12251, # 12253, # 12259, respectively) (76). Two days after transformation, the lentiviral

particles in the supernatant were purified from cell debris thru syringe filtration and centrifugation. After

titration, the particles were used to transduce either low passage nHDF or ARPE-19 cells. After a week of

puromycin selection, cells were tested for RL13 expression and aliquots were stored in liquid nitrogen until

further usage.

HCMV. All human cytomegalovirus (HCMV) strains were derived from bacterial artificial chromosome (BAC)

clones. BAC clone TB40/e (BAC4) was provided by Christian Sinzger (University of Ulm, Germany) (41).

BAC clone TR was provided by Jay Nelson (Oregon Health and Sciences University, Portland, OR, USA)

(77). BAC clone Merlin (ME)(pAL1393), which contains tetracycline operator sequences within the transcriptional promoter of UL130 and UL131, was provided by Richard Stanton (Cardiff University, Cardiff, United Kingdom) (40). All BAC clones were modified to express green fluorescent protein (GFP) with En passant recombineering (78) by replacing US11 with the eGFP gene. The constitutive expression of eGFP allows the monitoring of HCMV infection early and is strain-independent. Infectious HCMV was recovered by electroporation of BAC-DNA into HFFFs, as previously described (26). For infectious unit (IU) determination, viruses were serial diluted and infectivity was determined on fibroblasts or epithelial cells using flow cytometry 48 hours post infection.

Antibodies. Monoclonal antibodies specific to HCMV major capsid protein (MCP), gH (14-4b), and gB (27-156) were provided by Bill Britt (University of Alabama, Birmingham, AL) (52, 57, 79). 14-4b was purified by FPLC prior to use. Rabbit polyclonal sera against HCMV gL, UL130, and UL131 were provided by David Johnson (Oregon Health and Sciences University, Portland, OR) (17).

Vrial spread assays. Approx. 1×10^5 (3×10^5) nHDFs (ARPEs) were seeded into 6-well culture plates and allowed to grow to confluence. Confluent monolayers of nHDFs or ARPE cells were inoculated with 100-1000 IUs of strains TB, TR, or ME at for 4hrs at 37°C. Cells were then washed extensively with PBS and cultured in the appropriate growth medium supplemented with 2% FBS. Viral spread in the presence or absence of neutralizing antibodies or ganciclovir was monitored over 12 days by flow cytometry. For fibroblasts, 100ug/mL anti-gH 14-4b was used; for epithelial cells, a 1:1000 dilution of anti-UL130/131 rabit sera. 50uM ganciclovir was used on both cell types. All experiments were performed in triplicate and a minimum of 3 experiments was conducted for each condition. Spread rates were determined by plotting LN GFP+ cells over time and fitting the data to the log-linear rate expression $LN(I)_t = m(t) + LN(I)_0$ where (I) is the number of GFP+ cells, (t) is the time in days, and m is the spread rate.

Flow cytometry. Recombinant GFP-expressing HCMV-infected cells were washed twice with PBS and lifted with trypsin. Trypsin was quenched with DMEM containing 10% FBS and cells were spun at 500xg for 5 min at RT. Cells were fixed in PBS containing 2% paraformaldehyde for 10 min at RT, then washed and resuspended in PBS. Samples were analyzed using an AttuneNxT flow cytometer. Cells were identified using FSC-A and SSC-A, and single cells were gated using FSC-W and FSC-H. BL-1 laser (488nm) was used to identify GFP+ cells, and only cells with median GFP intensities 10-fold above background were

considered positive. RL13 expression was measured using an intracellular staining kit (Thermo) and an anti-

6x His antibody conjugated to AlexaFluor-647 (Thermo) using the RL-2 laser (647nm).

qPCR. The real-time quantitative PCR (qPCR) assay used to quantify viral or cellular DNA molecules was

performed as previously described (27). Briefly, HCMV DNA inside cell-free particles was purified using the

PureLink Viral RNA/DNA mini kit (Thermo Scientific). A region within UL83 conserved among ME, TR, and

TB was chosen as the HCMV-specific amplicon, and viral genomes were quantified by SYBR green qPCR

as previously described. Standard curves were performed using serial dilutions of a single PCR DNA band

containing the sequences of the viral UL83 and cellular beta2-microglobulin (see below) amplicons and the

corresponding set of primers. Finally, the concentration of HCMV DNA genomes in the supernatant was

extrapolated from the UL83 standard curve and expressed as genome molecules per ml of media

(genome/ml). Total intracellular or cytoplasmic HCMV DNA was quantified with the UL83-based qPCR after

extracting viral and cellular DNA from infected cells with the PureLink Genomic DNA mini kit (Thermo

Scientific). We also measured the chromosomal DNA molecules present in these samples with an amplicon

located in the human beta-2 microglobulin gene (80). These chromosomes numbers were used to correct

for the number of cells when measuring the whole cell, nuclear, and cytoplasmic genomes from different

samples. To obtain these fractions, infected cells were subjected to a Subcellular Fractionation Kit for

cultured cells (Thermo Scientific) to purify the cytoplasmic HCMV DNA from the excess of nuclear HCMV

DNA. Lastly, genomes present in the cytoplasmic fraction were purified using the Genomic DNA mini kit and

their number was measured with the UL83-based qPCR assay.

Particle production. Production of intracellular and extracellular HCMV genomes was determined over a

time course of 4 days. Briefly, nHDFs and ARPEs were infected with HCMV strains TB, TR, or ME at an

MOI of 1, and the cells were extensively washed 3 days post infection. At 3, 5, and 7 dpi culture

supernatants were collected and spun at 500xg for 10 minutes at RT, while cells were lifted with trypsin and

then quenched with culture media containing 10% FBS. Cells were spun at 500xg and resuspended into

aliquots. Cells were either resuspended in PBS for whole cell analysis, or fractionated for nuclear and

cytoplasmic analysis using the Subcellular Fractionation Kit for Cultured Cells (Thermo). Cells were also

sonicated to release cell-associated virus for infectivity analysis, or processed for flow cytometry to

determine the number of GFP+ cells. The number of HMCV genomes in the whole cell, nuclear,

cytoplasmic, and supernatant fractions was measured by qPCR, normalized to load, and divided by the number of GFP+ cells. The total infectivity of supernatant and sonicated fractions was determined by flow cytometry 48 hours after inoculation of the producer cell type. The accumulation at day 7 from nHDFs and ARPEs is shown in Fig 5.

Immunoblot analyses. Cell-free virions were solubilized in a buffer containing 20mM Tris-buffered saline (TBS) (pH 6.8) and 2% SDS. Protein samples were separated by SDS-polyacrylamide gel electrophoreses (SDS-PAGE) and electrophoretically transferred to polyvinylidene difluoride membranes in a buffer containing 10mM $NaHCO_3$ and 3mM Na_2CO_3 (pH 9.9) and 10% methanol. Transferred proteins were first probed with MAbs or rabbit sera, then anti-mouse or anti-rabbit secondary antibodies conjugated to horseradish peroxidase (Sigma), and Pierce ECL-Western Blotting substrate (Sigma). Chemiluminescence was detected using a Bio-Rad ChemiDoc MP imaging system.

ACKNOWLEDGEMENTS

We are grateful to Richard Stanton, Bill Britt, Jay Nelson, and David Johnson, for generously supplying HCMV BAC clones, antibodies, and cell lines as indicated in the Materials and Methods. Additionally, we are grateful to the Center for Biomolecular Structure and Dynamics (CBSD), University of Montana, Missoula, MT, for purification of monoclonal antibodies, and well as the Flow Cytometry Core of the Center for Environmental Heath Sciences (CEHS), University of Montana, Missoula, MT, for guidance on experimental design, acquisition, and analysis of the flow cytometry-based approaches used for this study.

This work was supported by grant from the National Institutes of Health to B.J.R (R01AI097274), a fellowship from American Heart Association to E.P.S (17POST33350043) and a National Institutes of Health CoBRE award to Center for Biomolecular Structure and Dynamics at University of Montana (PG20GM103546).

Experiments were designed by E.P.S, B.J.R, and J.L., and performed by E.P.S, J.L, L.Z.D., C.P., J.P, and Q.Y., and the manuscript was prepared by B.J.R., E.P.S., and J.L.

REFERENCES

1. Cui, X., Z. Cao, S. Wang, M. Flora, S. P. Adler, M. A. McVoy, and C. M. Snapper. 2019. Immunization of Rabbits with Recombinant Human Cytomegalovirus Trimeric versus Monomeric gH/gL Protein Elicits Markedly Higher Titers of Antibody and Neutralization Activity. *Int J Mol Sci* 20:
2. Razonable, R. R. 2018. Drug-resistant cytomegalovirus: clinical implications of specific mutations. *Curr Opin Organ Transplant* 23: 388-394.
3. Cudini, J., S. Roy, C. J. Houldcroft, J. M. Bryant, D. P. Depledge, H. Tutill, P. Veys, R. Williams, A. J. J. Worth, A. U. Tamuri, R. A. Goldstein, and J. Breuer. 2019. Human cytomegalovirus haplotype reconstruction reveals high diversity due to superinfection and evidence of within-host recombination. *Proc Natl Acad Sci U S A* 116: 5693-5698.
4. Lassalle, F., D. P. Depledge, M. B. Reeves, A. C. Brown, M. T. Christiansen, H. J. Tutill, R. J. Williams, K. Einer-Jensen, J. Holdstock, C. Atkinson, J. R. Brown, F. B. van Loenen, D. A. Clark, P. D. Griffiths, G. M. G. M. Verjans, M. Schutten, R. S. B. Milne, F. Balloux, and J. Breuer. 2016. Islands of linkage in an ocean of pervasive recombination reveals two-speed evolution of human cytomegalovirus genomes. *Virus Evol* 2: vew017.
5. Renzette, N., B. Bhattacharjee, J. D. Jensen, L. Gibson, and T. F. Kowalik. 2011. Extensive genome-wide variability of human cytomegalovirus in congenitally infected infants. *PLoS Pathog* 7: e1001344.
6. Renzette, N., L. Gibson, B. Bhattacharjee, D. Fisher, M. R. Schleiss, J. D. Jensen, and T. F. Kowalik. 2013. Rapid Intrahost Evolution of Human Cytomegalovirus Is Shaped by Demography and Positive Selection. *PLoS Genet* 9: e1003735.
7. Sijmons, S., K. Thys, M. Mbong Ngwese, E. Van Damme, J. Dvorak, M. Van Loock, G. Li, R. Tachezy, L. Busson, J. Aerssens, M. Van Ranst, and P. Maes. 2015. High-throughput analysis of human cytomegalovirus genome diversity highlights the widespread occurrence of gene-disrupting mutations and pervasive recombination. *J Virol* 89: 7673-7695.
8. Suárez, N. M., G. S. Wilkie, E. Hage, S. Camiolo, M. Holton, J. Hughes, M. D. Maabar, V. B. Sreenu, A. Dhingra, U. A. Gompels, G. W. G. Wilkinson, F. Baldanti, M. Furione, D. Lilleri, A. Arossa, T. Ganzenmueller, G. Gerna, P. Hubáček, T. F. Schulz, D. Wolf, M. Zavattoni, and A. J. Davison. 2019. Human Cytomegalovirus Genomes Sequenced Directly from Clinical Material: Variation, Multiple-Strain Infection, Recombination and Gene Loss. *J Infect Dis*
9. Suárez, N. M., K. G. Musonda, E. Escriva, M. Njenga, A. Agbueze, S. Camiolo, A. J. Davison, and U. A. Gompels. 2019. Multiple-Strain Infections of Human Cytomegalovirus with High Genomic Diversity are Common In Breast Milk from HIV-Positive Women in Zambia. *J Infect Dis*
10. Dolan, A., C. Cunningham, R. D. Hector, A. F. Hassan-Walker, L. Lee, C. Addison, D. J. Dargan, D. J. McGeoch, D. Gatherer, V. C. Emery, P. D. Griffiths, C. Sinzger, B. P. McSharry, G. W. Wilkinson, and A. J. Davison. 2004. Genetic content of wild-type human cytomegalovirus. *J Gen Virol* 85: 1301-1312.
11. Stern-Ginossar, N., B. Weisburd, A. Michalski, V. T. Le, M. Y. Hein, S. X. Huang, M. Ma, B. Shen, S. B. Qian, H. Hengel, M. Mann, N. T. Ingolia, and J. S. Weissman. 2012. Decoding human cytomegalovirus. *Science* 338: 1088-1093.
12. Connolly, S. A., J. O. Jackson, T. S. Jardetzky, and R. Longnecker. 2011. Fusing structure and function: a structural view of the herpesvirus entry machinery. *Nat Rev Microbiol* 9: 369-381.
13. Huber, M. T., and T. Compton. 1997. Characterization of a novel third member of the human cytomegalovirus glycoprotein H-glycoprotein L complex. *J Virol* 71: 5391-5398.
14. Li, L., J. A. Nelson, and W. J. Britt. 1997. Glycoprotein H-related complexes of human cytomegalovirus: identification of a third protein in the gCIII complex. *J Virol* 71: 3090-3097.
15. Adler, B., L. Scrivano, Z. Ruzcics, B. Rupp, C. Sinzger, and U. Koszinowski. 2006. Role of human cytomegalovirus UL131A in cell type-specific virus entry and release. *J Gen Virol* 87: 2451-2460.
16. Hahn, G., M. G. Revello, M. Patrone, E. Percivalle, G. Campanini, A. Sarasini, M. Wagner, A. Gallina, G. Milanesi, U. Koszinowski, F. Baldanti, and G. Gerna. 2004. Human cytomegalovirus UL131-128 genes are indispensable for virus growth in endothelial cells and virus transfer to leukocytes. *J Virol* 78: 10023-10033.
17. Ryckman, B. J., B. L. Rainish, M. C. Chase, J. A. Borton, J. A. Nelson, M. A. Jarvis, and D. C.

631   Johnson. 2008. Characterization of the human cytomegalovirus gH/gL/UL128-131 complex that
632   mediates entry into epithelial and endothelial cells. *J Virol* 82: 60-70.

18. Wang, D., and T. Shenk. 2005. Human cytomegalovirus UL131 open reading frame is required for epithelial cell tropism. *J Virol* 79: 10330-10338.

19. Vanarsdall, A. L., P. W. Howard, T. W. Wisner, and D. C. Johnson. 2016. Human Cytomegalovirus gH/gL Forms a Stable Complex with the Fusion Protein gB in Virions. *PLoS Pathog* 12: e1005564.

20. Vanarsdall, A. L., B. J. Ryckman, M. C. Chase, and D. C. Johnson. 2008. Human cytomegalovirus glycoproteins gB and gH/gL mediate epithelial cell-cell fusion when expressed either in cis or in trans. *J Virol* 82: 11837-11850.

21. E, X., P. Meraner, P. Lu, J. M. Perreira, A. M. Aker, W. M. McDougall, R. Zhuge, G. C. Chan, R. M. Gerstein, P. Caposio, A. D. Yurochko, A. L. Brass, and T. F. Kowalik. 2019. OR14I1 is a receptor for the human cytomegalovirus pentameric complex and defines viral epithelial cell tropism. *Proc Natl Acad Sci U S A*

22. Kabanova, A., J. Marcandalli, T. Zhou, S. Bianchi, U. Baxa, Y. Tsybovsky, D. Lilleri, C. Silacci-Fregni, M. Foglierini, B. M. Fernandez-Rodriguez, A. Druz, B. Zhang, R. Geiger, M. Pagani, F. Sallusto, P. D. Kwong, D. Corti, A. Lanzavecchia, and L. Perez. 2016. Platelet-derived growth factor-α receptor is the cellular receptor for human cytomegalovirus gHgLgO trimer. *Nat Microbiol* 2016:

23. Martinez-Martin, N., J. Marcandalli, C. S. Huang, C. P. Arthur, M. Perotti, M. Foglierini, H. Ho, A. M. Dosey, S. Shriver, J. Payandeh, A. Leitner, A. Lanzavecchia, L. Perez, and C. Ciferri. 2018. An Unbiased Screen for Human Cytomegalovirus Identifies Neuropilin-2 as a Central Viral Receptor. *Cell*

24. Laib Sampaio, K., C. Stegmann, I. Brizic, B. Adler, R. J. Stanton, and C. Sinzger. 2016. The contribution of pUL74 to growth of human cytomegalovirus is masked in the presence of RL13 and UL128 expression. *J Gen Virol*

25. Stegmann, C., D. Hochdorfer, D. Lieber, N. Subramanian, D. Stöhr, K. Laib Sampaio, and C. Sinzger. 2017. A derivative of platelet-derived growth factor receptor alpha binds to the trimer of human cytomegalovirus and inhibits entry into fibroblasts and endothelial cells. *PLoS Pathog* 13: e1006273.

26. Wille, P. T., A. J. Knoche, J. A. Nelson, M. A. Jarvis, and D. C. Johnson. 2010. A human cytomegalovirus gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter fibroblasts and epithelial and endothelial cells. *J Virol* 84: 2585-2596.

27. Zhou, M., J. M. Lanchy, and B. J. Ryckman. 2015. Human cytomegalovirus gH/gL/gO promotes the fusion step of entry into all cell types whereas gH/gL/UL128-131 broadens virus tropism through a distinct mechanism. *J Virol*

28. Liu, J., T. S. Jardetzky, A. L. Chin, D. C. Johnson, and A. L. Vanarsdall. 2018. The human cytomegalovirus trimer and pentamer promote sequential steps in entry into epithelial and endothelial cells at cell surfaces and endosomes. *J Virol*

29. Luo, M. H., P. H. Schwartz, and E. A. Fortunato. 2008. Neonatal neural progenitor cells and their neuronal and glial cell derivatives are fully permissive for human cytomegalovirus infection. *J Virol* 82: 9994-10007.

30. Nogalski, M. T., G. C. Chan, E. V. Stevenson, D. K. Collins-McMillen, and A. D. Yurochko. 2013. The HCMV gH/gL/UL128-131 Complex Triggers the Specific Cellular Activation Required for Efficient Viral Internalization into Target Monocytes. *PLoS Pathog* 9: e1003463.

31. Ryckman, B. J., M. A. Jarvis, D. D. Drummond, J. A. Nelson, and D. C. Johnson. 2006. Human cytomegalovirus entry into epithelial and endothelial cells depends on genes UL128 to UL150 and occurs by endocytosis and low-pH fusion. *J Virol* 80: 710-722.

32. Auerbach, M., D. Yan, A. Fouts, M. Xu, A. Estevez, C. D. Austin, F. Bazan, and B. Feierbach. 2013. Characterization of the guinea pig CMV gH/gL/GP129/GP131/GP133 complex in infection and spread. *Virology* 441: 75-84.

33. Coleman, S., J. Hornig, S. Maddux, K. Y. Choi, and A. McGregor. 2015. Viral Glycoprotein Complex Formation, Essential Function and Immunogenicity in the Guinea Pig Model for Cytomegalovirus. *PLoS One* 10: e0135567.

34. Lemmermann, N. A., A. Krmpotic, J. Podlech, I. Brizic, A. Prager, H. Adler, A. Karbach, Y. Wu, S. Jonjic, M. J. Reddehase, and B. Adler. 2015. Non-redundant and Redundant Roles of Cytomegalovirus

683   gH/gL Complexes in Host Organ Entry and Intra-tissue Spread. *PLoS Pathog* 11: e1004640.

684 35. Scrivano, L., C. Sinzger, H. Nitschko, U. H. Koszinowski, and B. Adler. 2011. HCMV spread and cell
685   tropism are determined by distinct virus populations. *PLoS Pathog* 7: e1001256.

686 36. Wagner, F. M., I. Brizic, A. Prager, T. Trsan, M. Arapovic, N. A. Lemmermann, J. Podlech, M. J.
687   Reddehase, F. Lemnitzer, J. B. Bosse, M. Gimpfl, L. Marcinowski, M. Macdonald, H. Adler, U. H.
688   Koszinowski, and B. Adler. 2013. The Viral Chemokine MCK-2 of Murine Cytomegalovirus Promotes
689   Infection as Part of a gH/gL/MCK-2 Complex. *PLoS Pathog* 9: e1003493.

690 37. Yunis, J., H. E. Farrell, K. Bruce, C. Lawler, O. Wyer, N. Davis-Poynter, I. Brizić, S. Jonjić, B. Adler,
691   and P. G. Stevenson. 2018. Murine Cytomegalovirus glycoprotein O promotes epithelial cell infection. *J*
692   *Virol*

693 38. Murrell, I., P. Tomasec, G. S. Wilkie, D. J. Dargan, A. J. Davison, and R. J. Stanton. 2013. Impact of
694   sequence variation in the UL128 locus on production of human cytomegalovirus in fibroblast and
695   epithelial cells. *J Virol* 87: 10489-10500.

696 39. Zhang, L., M. Zhou, R. Stanton, J. Kamil, and B. J. Ryckman. 2018. Expression levels of glycoprotein
697   O (gO) vary between strains of human cytomegalovirus, influencing the assembly of gH/gL complexes
698   and virion infectivity. *J Virol*

699 40. Stanton, R. J., K. Baluchova, D. J. Dargan, C. Cunningham, O. Sheehy, S. Seirafian, B. P. McSharry,
700   M. L. Neale, J. A. Davies, P. Tomasec, A. J. Davison, and G. W. Wilkinson. 2010. Reconstruction of
701   the complete human cytomegalovirus genome in a BAC reveals RL13 to be a potent inhibitor of
702   replication. *J Clin Invest* 120: 3191-3208.

703 41. Sinzger, C., G. Hahn, M. Digel, R. Katona, K. L. Sampaio, M. Messerle, H. Hengel, U. Koszinowski,
704   W. Brune, and B. Adler. 2008. Cloning and sequencing of a highly productive, endotheliotropic virus
705   strain derived from human cytomegalovirus TB40/E. *J Gen Virol* 89: 359-368.

706 42. Murrell, I., C. Bedford, K. Ladell, K. L. Miners, D. A. Price, P. Tomasec, G. W. G. Wilkinson, and R. J.
707   Stanton. 2017. The pentameric complex drives immunologically covert cell-cell transmission of wild-
708   type human cytomegalovirus. *Proc Natl Acad Sci U S A*

709 43. Wu, K., A. Oberstein, W. Wang, and T. Shenk. 2018. Role of PDGF receptor-α during human
710   cytomegalovirus entry into fibroblasts. *Proc Natl Acad Sci U S A*

711 44. Wu, Y., A. Prager, S. Boos, M. Resch, I. Brizic, M. Mach, S. Wildner, L. Scrivano, and B. Adler. 2017.
712   Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-α as a key for entry. *PLoS*
713   *Pathog* 13: e1006281.

714 45. Digel, M., K. L. Sampaio, G. Jahn, and C. Sinzger. 2006. Evidence for direct transfer of cytoplasmic
715   material from infected to uninfected cells during cell-associated spread of human cytomegalovirus. *J*
716   *Clin Virol* 37: 10-20.

717 46. Gerna, G., E. Percivalle, F. Baldanti, S. Sozzani, P. Lanzarini, E. Genini, D. Lilleri, and M. G. Revello.
718   2000. Human cytomegalovirus replicates abortively in polymorphonuclear leukocytes after transfer
719   from infected endothelial cells via transient microfusion events. *J Virol* 74: 5629-5638.

720 47. Silva, M. C., J. Schroer, and T. Shenk. 2005. Human cytomegalovirus cell-to-cell spread in the absence
721   of an essential assembly protein. *Proc Natl Acad Sci U S A* 102: 2081-2086.

722 48. Cui, X., D. C. Freed, D. Wang, P. Qiu, F. Li, T. M. Fu, L. M. Kauvar, and M. A. McVoy. 2017. Impact
723   of antibodies and strain polymorphisms on cytomegalovirus entry and spread in fibroblasts and
724   epithelial cells. *J Virol*

725 49. Gerna, G., E. Percivalle, L. Perez, A. Lanzavecchia, and D. Lilleri. 2016. Monoclonal antibodies to
726   different components of the human cytomegalovirus (HCMV) pentamer gH/gL/pUL128L and trimer
727   gH/gL/gO as well as antibodies elicited during primary HCMV infection prevent epithelial cell
728   syncytium formation. *J Virol*

729 50. Murrell, I., G. S. Wilkie, A. J. Davison, E. Statkute, C. A. Fielding, P. Tomasec, G. W. Wilkinson, and
730   R. J. Stanton. 2016. Genetic Stability of Bacterial Artificial Chromosome-Derived Human
731   Cytomegalovirus during Culture In Vitro. *J Virol* 90: 3929-3943.

732 51. Zhou, M., Q. Yu, A. Wechsler, and B. J. Ryckman. 2013. Comparative analysis of gO isoforms reveals
733   that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and gH/gL/UL128-131 in the
734   virion envelope. *J Virol* 87: 9680-9690.

52. Bogner, E., M. Reschke, B. Reis, E. Reis, W. Britt, and K. Radsak. 1992. Recognition of
compartmentalized intracellular analogs of glycoprotein H of human cytomegalovirus. *Arch Virol* 126:
67-80.
53. Schultz, E. P., J. M. Lanchy, E. E. Ellerbeck, and B. J. Ryckman. 2015. Scanning mutagenesis of human
cytomegalovirus gH/gL. *J Virol*
54. Baldanti, F., M. R. Underwood, C. L. Talarico, L. Simoncini, A. Sarasini, K. K. Biron, and G. Gerna.
1998. The Cys607-->Tyr change in the UL97 phosphotransferase confers ganciclovir resistance to two
human cytomegalovirus strains recovered from two immunocompromised patients. *Antimicrob Agents*
Chemother 42: 444-446.
55. Dargan, D. J., E. Douglas, C. Cunningham, F. Jamieson, R. J. Stanton, K. Baluchova, B. P. McSharry,
P. Tomasec, V. C. Emery, E. Percivalle, A. Sarasini, G. Gerna, G. W. Wilkinson, and A. J. Davison.
2010. Sequential mutations associated with adaptation of human cytomegalovirus to growth in cell
culture. *J Gen Virol* 91: 1535-1546.
56. Britt, W. J., and L. G. Vugler. 1989. Processing of the gp55-116 envelope glycoprotein complex (gB) of
human cytomegalovirus. *J Virol* 63: 403-410.
57. Schoppel, K., E. Hassfurther, W. Britt, M. Ohlin, C. A. Borrebaeck, and M. Mach. 1996. Antibodies
specific for the antigenic domain 1 of glycoprotein B (gpUL55) of human cytomegalovirus bind to
different substructures. *Virology* 216: 133-145.
58. Spaete, R. R., A. Saxena, P. I. Scott, G. J. Song, W. S. Probert, W. J. Britt, W. Gibson, L. Rasmussen,
and C. Pachl. 1990. Sequence requirements for proteolytic processing of glycoprotein B of human
cytomegalovirus strain Towne. *J Virol* 64: 2922-2931.
59. Sinzger, C., K. Schmidt, J. Knapp, M. Kahl, R. Beck, J. Waldman, H. Hebart, H. Einsele, and G. Jahn.
1999. Modification of human cytomegalovirus tropism through propagation in vitro is associated with
changes in the viral genome. *J Gen Virol* 80: 2867-2877.
60. Galitska, G., M. Biolatti, M. De Andrea, A. Leone, A. Coscia, L. Bertolotti, U. Ala, E. Bertino, V.
Dell'Oste, and S. Landolfo. 2018. Biological relevance of Cytomegalovirus genetic variability in
congenitally and postnatally infected children. *J Clin Virol* 108: 132-140.
61. Subramanian, N., Z. Wu, and T. M. Mertens. 2016. Phenotypic characterization of human
cytomegalovirus strains in cell cultures based on their transmission kinetics. *J Gen Virol* 97: 2376-2386.
62. Rasmussen, L., A. Geissler, C. Cowan, A. Chase, and M. Winters. 2002. The genes encoding the gCIII
complex of human cytomegalovirus exist in highly diverse combinations in clinical isolates. *J Virol* 76:
10841-10848.
63. Mattick, C., D. Dewin, S. Polley, E. Sevilla-Reyes, S. Pignatelli, W. Rawlinson, G. Wilkinson, P. Dal
Monte, and U. A. Gompels. 2004. Linkage of human cytomegalovirus glycoprotein gO variant groups
identified from worldwide clinical isolates with gN genotypes, implications for disease associations and
evidence for N-terminal sites of positive selection. *Virology* 318: 582-597.
64. Stanton, R., D. Westmoreland, J. D. Fox, A. J. Davison, and G. W. Wilkinson. 2005. Stability of human
cytomegalovirus genotypes in persistently infected renal transplant recipients. *J Med Virol* 75: 42-46.
65. Jiang, X. J., B. Adler, K. L. Sampaio, M. Digel, G. Jahn, N. Ettischer, Y. D. Stierhof, L. Scrivano, U.
Koszinowski, M. Mach, and C. Sinzger. 2008. UL74 of human cytomegalovirus contributes to virus
release by promoting secondary envelopment of virions. *J Virol* 82: 2802-2812.
66. Day, L. Z., C. Stegmann, E. P. Schultz, J.-M. Lanchy, Q. Yu, and B. J. Ryckman. 2019. Naturally
Occurring Polymorphisms in Human Cytomegalovirus gO Exert Epistatic Influences on Cell-Free and
Cell-To-Cell Spread, and Antibody Neutralization on gH Epitopes. *bioRxiv* 867234.
67. Strive, T., E. Borst, M. Messerle, and K. Radsak. 2002. Proteolytic processing of human
cytomegalovirus glycoprotein B is dispensable for viral growth in culture. *J Virol* 76: 1252-1264.
68. Ourahmane, A., X. Cui, L. He, M. Catron, D. P. Dittmer, A. Al Qaffasaa, M. R. Schleiss, L. Hertel, and
M. A. McVoy. 2019. Inclusion of Antibodies to Cell Culture Media Preserves the Integrity of Genes
Encoding RL13 and the Pentameric Complex Components During Fibroblast Passage of Human
Cytomegalovirus. *Viruses* 11:
69. Hutt-Fletcher, L. M. 2007. Epstein-Barr virus entry. *J Virol* 81: 7825-7832.
70. Cannon, M. J., T. B. Hyde, and D. S. Schmid. 2011. Review of cytomegalovirus shedding in bodily

787   fluids and relevance to congenital cytomegalovirus infection. *Rev Med Virol* 21: 240-255.

788 71. Chou, S., and T. C. Merigan. 1983. Rapid detection and quantitation of human cytomegalovirus in urine
789   through DNA hybridization. *N Engl J Med* 308: 921-925.

790 72. Gerna, G., E. Percivalle, A. Sarasini, and M. G. Revello. 2002. Human cytomegalovirus and human
791   umbilical vein endothelial cells: restriction of primary isolation to blood samples and susceptibilities of
792   clinical isolates from other sources to adaptation. *J Clin Microbiol* 40: 233-238.

793 73. Kitamura, T., Y. Koshino, F. Shibata, T. Oki, H. Nakajima, T. Nosaka, and H. Kumagai. 2003.
794   Retrovirus-mediated gene transfer and expression cloning: powerful tools in functional genomics. *Exp
795   Hematol* 31: 1007-1014.

796 74. Stewart, S. A., D. M. Dykxhoorn, D. Palliser, H. Mizuno, E. Y. Yu, D. S. An, D. M. Sabatini, I. S.
797   Chen, W. C. Hahn, P. A. Sharp, R. A. Weinberg, and C. D. Novina. 2003. Lentivirus-delivered stable
798   gene silencing by RNAi in primary cells. *RNA* 9: 493-501.

799 75. Sancak, Y., T. R. Peterson, Y. D. Shaul, R. A. Lindquist, C. C. Thoreen, L. Bar-Peled, and D. M.
800   Sabatini. 2008. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science*
801   320: 1496-1501.

802 76. Dull, T., R. Zufferey, M. Kelly, R. J. Mandel, M. Nguyen, D. Trono, and L. Naldini. 1998. A third-
803   generation lentivirus vector with a conditional packaging system. *J Virol* 72: 8463-8471.

804 77. Murphy, E., D. Yu, J. Grimwood, J. Schmutz, M. Dickson, M. A. Jarvis, G. Hahn, J. A. Nelson, R. M.
805   Myers, and T. E. Shenk. 2003. Coding potential of laboratory and clinical strains of human
806   cytomegalovirus. *Proc Natl Acad Sci U S A* 100: 14976-14981.

807 78. Tischer, B. K., G. A. Smith, and N. Osterrieder. 2010. En passant mutagenesis: a two step markerless
808   red recombination system. *Methods Mol Biol* 634: 421-430.

809 79. Chee, M., S. A. Rudolph, B. Plachter, B. Barrell, and G. Jahn. 1989. Identification of the major capsid
810   protein gene of human cytomegalovirus. *J Virol* 63: 1345-1353.

811 80. Rooney, J. P., I. T. Ryde, L. H. Sanders, E. H. Howlett, M. D. Colton, K. E. Germ, G. D. Mayer, J. T.
812   Greenamyre, and J. N. Meyer. 2015. PCR based determination of mitochondrial DNA copy number in
813   multiple species. *Methods Mol Biol* 1241: 23-38.

    FIGURE LEGENDS

Figure 1. Comparison of focal spread patterns among distinct HCMV strains. Confluent monolayers of

fibroblasts (A) or epithelial cells (B) were infected at MOI 0.001 with GFP-expressing HCMV TB, TR, or ME.

Foci were documented by fluorescence microscopy at 6, 12, and 18 days post infection. Four representative

fields (10X) are shown for each.

Figure 2. Quantitation of HCMV spread by in fibroblasts and epithelial cells. Confluent 9.5cm^3

monolayers of fibroblasts (A) or epithelial cells (D) were infected at MOI 0.001 with GFP-expressing HCMV

strains TB, TR, or ME. The number of infected cells at 3, 6, 9, and 12 days post infection was determined by

flow cytometry. (B and E) LN GFP+ cells is plotted at each time point shown in panels A and D, respectively.

(C and F) The average spread rate (LN GFP+ cells per day) as the slope (m) of the regression lines in

panels B and E, respectively. All experiments were performed at least three times and error bars represent

standard deviation between all experiments (omitted if smaller than the data marker), and p-values were

generated using unpaired, two-tailed t-tests (*<0.05, **<0.01, ***<0.001).

Figure 3. Antibody neutralization of cell-free HCMV. Equal numbers (genomes/mL) of fibroblast-derived

(A) or epithelial-derived (B) HCMV TB, TR, or ME virions were incubated with multiple concentrations of anti-

gH mAb 14-4b (A) or anti-UL130/131 rabbit sera (B) for 1hr at RT. Remaining infectivity was determined by

titration on the matched producer cell type, and plotted as percent of no antibody control. All experiments

were performed in triplicate and error bars represent standard deviation.

Figure 4. Effects of neutralizing antibodies on spread of HCMV strains in fibroblasts and epithelial

cells. Confluent 9.5cm^3 monolayers of fibroblasts or epithelial cells were infected at MOI 0.001 with GFP-

expressing HCMV strains TB, TR, or ME and allowed to spread in the presence of either anti-gH mAb 14-4b

(A-C), anti-UL130/131 rabbit sera (D-F), or ganciclovir (all panels). The average spread rates over 12 days

(LN GFP+ cells per day; as in Fig 2) for TB, TR, and ME on fibroblasts (A-C) and epithelial cells (D-F) are

shown. Conditions where no spread was detected are designated "n.d.". All experiments were performed

three times and error bars represent standard deviation. P-values were generated by unpaired, two-tailed t-

test vs. mock (*<0.05, **<0.01, ***<0.001).

Fig 5. Production and infectivity of cell-free and cell-associated HCMV progeny virus. Replicate

cultures of fibroblasts or epithelial cells were infected at MOI 1 with TB, TR, or ME. 7 days later infected

cells were quantified by flow cytometry for GFP expression. *(A and C).* Cultures were fractionated into whole

cell (WC), nuclear (N), cytoplasmic (C), and supernatant (S), and viral genomes were determined by qPCR.

The number of HCMV genomes in each fraction is plotted per GFP+ cell. *(B and D).* Infectious units were

determined from whole cell sonicates (WC) and culture supernatants (S) by titration on the matched

producer cell type. Total infectious units are plotted per 1000 GFP+ cells. *All panels.* Error bars represent

the standard deviation of three separate experiments, and p-values were generated using unpaired, two-

tailed t-tests (*<0.05, **<0.01, ***<0.001, ****<0.0001).

Fig 6. Repression of gH/gL/UL128-131 expression from HCMV ME in fibroblasts and epithelial cells.

(A) Tetracycline-repressor protein (tetR) expressing fibroblasts (HFFFtet, nHDFtet) or epithelial cells

(ARPEtet) were transfected with TetO-firefly luciferase reporter and the relative luciferase activity was

measured. HFFFtet were treated with doxycycline (+) or left untreated (-). (B) Non-reducing (top) or

reducing (bottom) immunoblot analyses of MT or ME virions derived from TetR-expressing or control cells,

respectively. Blots were probed with anti-gL antibodies (top) or anti-MCP antibodies (bottom). (C) Specific

infectivity (genomes/IU) of MT and ME virions derived from culture supernatants of TetR-expressing or

control cells, respectively. Titrations of fibroblasts-derived virus were performed of nHDFs, epithelial-derived

virus on ARPEs. All experiments were performed in triplicate and error bars represent standard deviation. P-

values were generated by unpaired, two-tailed t-test (*<0.05, **<0.01, ***<0.001).

Figure 7. Effect of gH/gL/UL128-131 repression on HCMV ME spread in fibroblasts and epithelial

cells. Confluent 9.5cm^3 monolayers of (A) nHDF or nHDFtet, for ME and MT, respectively, (B) HFFFtet (+)

or (-) doxycycline, for ME and MT, respectively, or (C) ARPE or ARPEtet, for ME and MT, respectively were

infected with at MOI 0.001 with GFP-expressing HCMV ME/MT. Spread rates in the presence of either anti-

gH mAb 14-4b (A and B), or anti-UL130/131 rabbit sera (C) were determined as described in Figure 2.

Plotted are averages and standard deviaitons for three independent experiments. P-values were generated

by unpaired, two-tailed t-test (*<0.05, **<0.01).

Figure 8. RL13-expressing fibroblasts and epithelial cell lines. nHDF fibroblasts and ARPE epithelial

cells transduced with lentivectors encoding a C-terminal 6XHis tagged RL13 were analyzed by immunoblot

(A) or flow cytometry (B) using anti-6XHis antibodies. Median fluorescence intensities for RL13-expressing

nHDFs and ARPEs, and control cells are shown for comparison.

Figure 9. Effects of RL13 expression on spread of HCMV in fibroblasts and epithelial cells. Spread

rates of TB, TR, or ME were determined as in Fig. 2 in RL13-expressing fibroblasts (A) or epithelial cells (B)

and respective control cells in both the presence and absence of neutralizing antibodies anti-gH mAb 14-4b

(A), or anti-UL130/131 rabbit sera (B). Plotted are average spread rates and standard deviations for three

independent experiments. P-values were generated by unpaired two-tailed t-tests comparing RL13-

expressing and control cells (*<0.05, **<0.01).

Figure 10. Quantity and infectivity of cell-free HCMV progeny virus from RL13-expressing fibroblasts

and epithelial cells. RL13-expressing fibroblasts (A-B) and epithelial cells (C-D) or control cells were

infected with HMCV strains TB, TR, or ME at MOI 1. At 7 days post infection the number of infected cells

was determined by flow cytometry and culture supernatants were analyzed by qPCR for viral genomes (A

and C) or by titration on matched producer cell type to quantify infectious units. Plotted are averages and

standard deviations for three independent experiments. P-values were generated by unpaired two tailed t-

tests comparing RL13-expressing and control cells (*<0.05, **<0.01, ***<0.001).

Figure 11. Effect of RL13 expression on levels of gH/gL complexes and gB in ME virions. Equal

numbers (genomes/mL) of ME virions derived from either RL13-expressing or control fibroblasts or epithelial

cells were separated by SDS-PAGE under non-reducing (A) or reducing (B and C) conditions and analyzed

by immunoblot with antibodies directed against gL (A), gB (B), or major capsid protein (C).

A   Fibroblasts

B   Epithelial Cells

890
891
892
893   **Figure 1.   Comparison of focal spread patterns among distinct HCMV strains.**   Confluent monolayers of
894   fibroblasts (A) or epithelial cells (B) were infected at MOI 0.001 with GFP-expressing HCMV TB, TR, or ME.
895   Foci were documented by fluorescence microscopy at 6, 12, and 18 days post infection. Four representative
896   fields (10X) are shown for each.
897

898
899
900
Figure 2. Quantitation of HCMV spread by in fibroblasts and epithelial cells. Confluent 9.5cm^2
monolayers of fibroblasts (A) or epithelial cells (D) were infected at MOI 0.001 with GFP-expressing HCMV
strains TB, TR, or ME. The number of infected cells at 3, 6, 9, and 12 days post infection was determined by
flow cytometry. (B and E) LN GFP+ cells is plotted at each time point shown in panels A and D, respectively.
(C and F) The average spread rate (LN GFP+ cells per day) as the slope (m) of the regression lines in
panels B and E, respectively. All experiments were performed at least three times and error bars represent
standard deviation between all experiments (omitted if smaller than the data marker), and p-values were
generated using unpaired, two-tailed t-tests (*<0.05, **<0.01, ***<0.001).
909

910
911
912 **Figure 3. Antibody neutralization of cell-free HCMV.** Equal numbers (genomes/mL) of fibroblast-derived
913 (A) or epithelial-derived (B) HCMV TB, TR, or ME virions were incubated with multiple concentrations of anti-
914 gH mAb 14-4b (A) or anti-UL130/131 rabbit sera (B) for 1hr at RT. Remaining infectivity was determined by
915 titration on the matched producer cell type, and plotted as percent of no antibody control. All experiments
916 were performed in triplicate and error bars represent standard deviation.
917

918
919
920
Figure 4. Effects of neutralizing antibodies on spread of HCMV strains in fibroblasts and epithelial
cells. Confluent 9.5cm^2 monolayers of fibroblasts or epithelial cells were infected at MOI 0.001 with GFP-
expressing HCMV strains TB, TR, or ME and allowed to spread in the presence of either anti-gH mAb 14-4b
(A-C), anti-UL130/131 rabbit sera (D-F), or ganciclovir (all panels). The average spread rates over 12 days
(LN GFP+ cells per day; as in Fig 2) for TB, TR, and ME on fibroblasts (A-C) and epithelial cells (D-F) are
shown. Conditions where no spread was detected are designated "n.d.". All experiments were performed
three times and error bars represent standard deviation. P-values were generated by unpaired, two-tailed t-
test vs. mock (*<0.05, **<0.01, ***<0.001).
929

Fig 5. Production and infectivity of cell-free and cell-associated HCMV progeny virus. Replicate
cultures of fibroblasts or epithelial cells were infected at MOI 1 with TB, TR, or ME. 7 days later infected
cells were quantified by flow cytometry for GFP expression. *(A and C).* Cultures were fractionated into whole
cell (WC), nuclear (N), cytoplasmic (C), and supernatant (S), and viral genomes were determined by qPCR.
The number of HCMV genomes in each fraction is plotted per GFP+ cell. *(B and D).* Infectious units were
determined from whole cell sonicates (WC) and culture supernatants (S) by titration on the matched
producer cell type. Total infectious units are plotted per 1000 GFP+ cells. *All panels.* Error bars represent
the standard deviation of three separate experiments, and p-values were generated using unpaired, two-
tailed t-tests (*<0.05, **<0.01, ***<0.001, ****<0.0001).

941
942

A

B

C

943
944
945
Fig 6. Repression of gH/gL/UL128-131 expression from HCMV ME in fibroblasts and epithelial cells.
(A) Tetracycline-repressor protein (tetR) expressing fibroblasts (HFFFtet, nHDFtet) or epithelial cells
(ARPEtet) were transfected with TetO-firefly luciferase reporter and the relative luciferase activity was
measured. HFFFtet were treated with doxycycline (+) or left untreated (-). (B) Non-reducing (top) or
reducing (bottom) immunoblot analyses of MT or ME virions derived from TetR-expressing or control cells,
respectively. Blots were probed with anti-gL antibodies (top) or anti-MCP antibodies (bottom). (C) Specific
infectivity (genomes/IU) of MT and ME virions derived from culture supernatants of TetR-expressing or
control cells, respectively. Titrations of fibroblasts-derived virus were performed of nHDFs, epithelial-derived
virus on ARPEs. All experiments were performed in triplicate and error bars represent standard deviation. P-
values were generated by unpaired, two-tailed t-test (*<0.05, **<0.01, ***<0.001).
956

957
958
Figure 7. Effect of gH/gL/UL128-131 repression on HCMV ME spread in fibroblasts and epithelial
cells.   Confluent 9.5cm^2 monolayers of (A) nHDF or nHDFtet, for ME and MT, respectively, (B) HFFFtet (+)
or (-) doxycycline, for ME and MT, respectively, or (C) ARPE or ARPEtet, for ME and MT, respectively were
infected with at MOI 0.001 with GFP-expressing HCMV ME/MT. Spread rates in the presence of either anti-
gH mAb 14-4b (A and B), or anti-UL130/131 rabbit sera (C) were determined as described in Figure 2.
Plotted are averages and standard deviations for three independent experiments. P-values were generated
by unpaired, two-tailed t-test (*<0.05, **<0.01).
966
967
968

969
970
971
972 **Figure 8. RL13-expressing fibroblasts and epithelial cell lines.** nHDF fibroblasts and ARPE epithelial
973 cells transduced with lentivectors encoding a C-terminal 6XHis tagged RL13 were analyzed by immunoblot
974 (A) or flow cytometry (B) using anti-6XHis antibodies. Median fluorescence intensities for RL13-expressing
975 nHDFs and ARPEs, and control cells are shown for comparison.
976
977

A Fibroblasts

B Epithelial cells

Figure 9. Effects of RL13 expression on spread of HCMV in fibroblasts and epithelial cells. Spread rates of TB, TR, or ME were determined as in Fig. 2 in RL13-expressing fibroblasts (A) or epithelial cells (B) and respective control cells in both the presence and absence of neutralizing antibodies anti-gH mAb 14-4b (A), or anti-UL130/131 rabbit sera (B). Plotted are average spread rates and standard deviations for three independent experiments. P-values were generated by unpaired two-tailed t-tests comparing RL13-expressing and control cells (*<0.05, **<0.01).

Figure 10. Quantity and infectivity of cell-free HCMV progeny virus from RL13-expressing fibroblasts and epithelial cells. RL13-expressing fibroblasts (A-B) and epithelial cells (C-D) or control cells were infected with HMCV strains TB, TR, or ME at MOI 1. At 7 days post infection the number of infected cells was determined by flow cytometry and culture supernatants were analyzed by qPCR for viral genomes (A and C) or by titration on matched producer cell type to quantify infectious units. Plotted are averages and standard deviation for three independent experiments. P-values were generated by unpaired two tailed t-tests comparing RL13-expressing and control cells (*<0.05, **<0.01, ***<0.001).

Figure 11. Effect of RL13 expression on levels of gH/gL complexes and gB in ME virions. Equal numbers (genomes/mL) of ME virions derived from either RL13-expressing or control fibroblasts or epithelial cells were separated by SDS-PAGE under non-reducing (A) or reducing (B and C) conditions and analyzed by immunoblot with antibodies directed against gL (A), gB (B), or major capsid protein (C).