1   A limited role for autophagy during arbovirus infection of mosquito cells

3   Doug E. Brackney[1*] and Maria A. Correa[1]

7   1) Center for Vector-Borne and Zoonotic Diseases, Department of Environmental Sciences, The

8   Connecticut Agricultural Experiment Station, New Haven, Connecticut 06504, USA

14   * Corresponding Authors: Doug E. Brackney (doug.brackney@ct.gov)

Abstract:

Macroautophagy is an evolutionarily conserved cellular process critical for maintaining cellular

homeostasis. It can additionally function as an innate immune response to viral infection as has

been demonstrated for a number of arthropod-borne (arbo-) viruses. Arboviruses are maintained

in a transmission cycle between vertebrate hosts and invertebrate vectors yet the majority of

studies assessing autophagy-arbovirus interactions have been limited to the mammalian host.

Therefore we evaluated the role of autophagy during arbovirus infection of the invertebrate

vector using the tractable Aag2 *Aedes aegypti* mosquito cell culture system. Our data

demonstrates that autophagy is significantly induced in mosquito cells upon infection with two

flaviviruses, dengue virus (DENV) and Zika virus (ZIKV), as well as an unrelated mosquito-

borne virus, chikungunya virus (CHIKV; *Togaviridae*). While assessing the role of autophagy

during arbovirus infection, we observed a somewhat paradoxical outcome. Both induction and

suppression of autophagy via torin 1 and spautin-1, respectively, resulted in increased viral titers

for all three viruses, yet suppression of autophagy-related genes had no effect. Interestingly,

chemical modulators of autophagy had either no effect or opposite effects in another widely used

mosquito cell line, C6/36 *Aedes albopictus* cells. Together, our data reveals a limited role for

autophagy during arbovirus infection of mosquito cells. Further, our findings suggest that

commonly used chemical modulators of autophagy alter mosquito cells in such a way as to

promote viral replication; however, it is unclear if this occurs directly through autophagic

manipulation or other means.

Author Summary:

Arthropod-borne (arbo) viruses, specifically those transmitted by Aedes aegypti mosquitoes,

cause significant morbidity and mortality and pose a continued public health threat worldwide.

Many of these viruses lack vaccines or therapeutics and current mosquito control strategies are

underperforming. For these reasons, identifying vulnerabilities within the transmission cycle that

can be targeted will be critical to the development of novel control interventions. Autophagy is a

highly conserved cellular pathway and previous studies manipulating this pathway have shown

promise in minimizing viral infections in mammalian hosts. In this study we examined

arbovirus-autophagy interactions within vector mosquitoes. The goal was to elucidate the role of

autophagy during infection of mosquitoes in hopes of identifying critical interactions that can be

targeted by novel approaches to block infection of and transmission by vector mosquitoes.

Introduction:

Arthropod-borne (arbo) viruses, specifically those of the families *Flaviviridae* and

Togaviridae, cause significant morbidity and mortality and pose a continued public health threat

worldwide. This is highlighted by the sustained transmission of dengue virus (DENV;

Flaviviridae, *Flavivirus*) throughout the tropics, the emergent epidemics of Zika virus (ZIKV;

Flaviviridae, *Flavivirus*), chikungunya virus (CHIKV; *Togaviridae*, *Alphavirus*) and West Nile

virus (WNV; *Flaviviridae*, *Flavivirus*) in the Americas, and the sporadic outbreaks of yellow

fever virus (YFV; *Flaviviridae*, *Flavivirus*) in Africa and South America. Arboviruses are

maintained in a transmission cycle between vertebrate hosts and invertebrate vectors. Because

therapeutics and vaccines are lacking for many of these viruses, public education and vector

control strategies are the only defenses available to minimize disease burden and epidemic risk.

However, as insecticide resistance is becoming more commonplace it is critical that novel

approaches for preventing and treating arboviral diseases, targeting either vector or host, be

developed.

Macroautophagy (herein referred to as autophagy) is an essential cellular process

required for maintaining homeostasis and plays a crucial role in development, cell differentiation

and immunity (1, 2). Consequently, abnormal autophagic activity has been linked to a number of

pathologies, including cancer and neurodegenerative diseases (3). This evolutionarily conserved

pathway mediates the degradation and recycling of cellular components. Upon induction, protein

aggregates, damaged or dysfunctional organelles and foreign bodies are sequestered into cup-

shaped double-membrane vesicles termed phagophores or isolation membranes (4).

Appropriation of cargo can occur in a non-selective manner known as bulk autophagy where

cytoplasmic material is indiscriminately engulfed or through selective autophagy, which

specifically targets poly-ubiquitin tagged proteins (5). Through a series of kinase signaling

cascades and recruitment events the phagophore elongates, eventually encompassing the cargo

and forming an autophagosome (4). Subsequently, the mature autophagosomes fuse with

lysosomes resulting in the acidification of the newly formed autophagolysosome and degradation

of the cargo (4). Targeting aspects of the autophagy pathway has become a promising strategy

for combatting a number of diseases (6, 7).

Numerous stimuli have the ability to activate autophagy including innate immunity and

cellular stress. It is therefore not surprising that viral infections often induce an autophagic

response. In fact, studies have demonstrated that both flaviviruses and alphaviruses induce

autophagy during infection and that this can occur through a number of mechanisms, all of which

culminate in the inhibition of the Akt-TOR signaling pathway (8-14). During alphavirus

infection, autophagy functions as an antiviral defense. Previous studies have demonstrated that

activation of the autophagy pathway via recombinant Beclin 1 reduces Sindbis virus and CHIKV

titers and improves clinical outcomes in mice. This process is mediated by the autophagy cargo

receptor p62, which recognizes poly-ubiquitin tagged viral capsid proteins and transports them to

maturing autophagosomes for degradation (10, 13, 15-18). In comparison, flavivirus-autophagy

interactions are much more nuanced (19). For instance, during DENV infection, p62 actively

targets DENV proteins for degradation and replication complexes associated with the ER

(endoplasmic reticulum) are targeted by reticulophagy; however, DENV benefits from an

energetically favorable environment as a result of virus-induced lipophagy and efficient

processing of mature virions (9, 19-21). While it is clear that arboviruses closely interact with

autophagy during infection, these studies have been limited to either non-vector organisms or

mammals. As a result we know very little about the role of autophagy during arbovirus infection

of the other half of the transmission cycle, the invertebrate vector (22).

To address this knowledge gap we investigated the role of autophagy during DENV,

ZIKV and CHIKV infection of Aag2 mosquito cells derived from the primary vector *Aedes*

aegypti. Using an anti-Atg8 (mammalian homolog LC3) specific antibody and recombinant

Atg8-EGFP provided *in trans*, we assessed autophagy induction and flux by immunoblot and

microscopy. We observed an increase in the accumulation of Atg8 positive puncta and increased

conversion of Atg8 to Atg8-PE. Further, we found that chemical inhibition and induction of

autophagy both increased viral titers, while suppression of autophagy related genes had no effect.

Interestingly, chemical modulators had minimal effects on DENV titers in another commonly

used mosquito cell line, *Aedes albopictus* C6/36 cells, and spautin-1 inhibition of autophagy

significantly decreased in DENV titers in mammalian cells as previously reported (21). Together,

these data reveal a limited role for autophagy during DENV, ZIKV and CHIKV infection of

mosquito cells and highlight differences in autophagy-virus interactions between cell culture

systems. Further, our data suggest that outcomes associated with commonly used chemical

modulators of autophagy are cell-dependent and may result from cell-specific interactions with

the chemicals.

Materials and Methods

Cell Lines & Virus Strains

Autophagy was modeled in three different cell lines, the mosquito-derived *Aedes*

albopictus C6/36 (ATCC; American Type Culture Collection) and *Aedes aegypti* Aag2 cells

(Generously provided by Dr. Gregory Ebel, Colorado State University) and mammalian cell line

BHK-21 clone 15 (Syrian golden hamster kidney cells)(Generously provided by Dr. Rushika

Perrera, Colorado State University) . The two mosquito cell lines were maintained at 28°C in the

presence of CO_2, and the BHK cells were maintained at 37°C with CO_2. All cells were grown in

media containing 10% fetal bovine serum, sodium bicarbonate, 100 U/ml penicillin, 100 μg/ml

streptomycin, 0.25 μg/ml amphotericin B, L-glutamine, and non-essential amino acids. This

media was used in all transfections, infections and plaque assays. Cell infections were carried out

using three different viruses from two major arbovirus families *Flaviviridae* and *Togaviridae*.

DENV2 strain Jam 1409 (Jamaica) was passaged on C6/36 cells, ZIKV strain PRVABC 59

(Asian lineage-derived American strain, Puerto Rico, 2015) was passaged on Vero cells (African

green monkey kidney cells) and C6/36 cells, and CHIKV strain R99659 (Caribbean lineage, British Virgin Islands, 2013) was passaged on Vero cells.

Pharmacological Modulators

To assess the effect of pharmacological modulators on autophagy, the following drugs were used in cell culture infections of C6/36, Aag2, and BHK21 cells: torin 1, an autophagy inducer that works by inhibiting mTOR kinase, and spautin 1 (Tocris Bioscience), which inhibits autophagy by interfering with USP10 and USP13, two ubiquitin-specific peptidases, resulting in the degradation of class III phosphatidylinositol-3 kinase complexes (23). Bafilomycin A1 (InvivoGen), an inhibitor of V-ATPase, was also used in inhibition of late stage autophagy in infectious and non-infectious immunoblotting of Aag2 cells. All drugs were prepared in dimethyl sulfoxide (DMSO) and stored at -20°C.

Immunoblots

Two days after Aag2 cells were seeded at 1.5×10^6 cells/well in a 12-well plate, cells were treated with 1% DMSO, 1 µM bafilomycin A1, 1 µM torin-1 or 10 µM spautin-1. Cells were harvested 24 hours later using RIPA buffer. Total protein in cell lysates was quantified using a Pierce BCA Protein Assay kit (ThermoFisher) on a microplate reader. An equal amount of protein from each sample was separated on a 16% Tris-glycine gel and transferred to a polyvinylidene difluoride (PVDF) membrane. Autophagy was detected using a rabbit anti-Atg8 antibody at a 1:8,000 dilution in PBS + 5% BSA + 1% Tween. Polyclonal anti-Atg8 antibody was generated by ProSci Inc. as previously described (24). The blot was then probed with horseradish peroxidase (HRP) goat anti-rabbit antibody at a 1:20,000 dilution. Actin was detected using a rabbit anti-β-Actin antibody (Abcam) in PBS + 5% BSA + 1% Tween at a

1:8,000 dilution. The secondary probe was performed using the same secondary antibody as above. Membranes were developed in SuperSignal West Pico chemiluminescent substrate (Thermo Fisher) for 5 minutes at room temperature. Band intensities were quantified using ImageJ.

The effect of virus infection in combination with pharmacological modulators on autophagy was also assayed via western blot. Aag2 cells were prepared as above and infected with DENV, ZIKV, or CHIKV at a multiplicity of infection (MOI) of 0.5. After one hour of infection at 28°C, DMSO, bafilomycin A1, 1 µM torin-1, or 10 µM spautin-1 were added to the wells. For DENV and ZIKV, cells were harvested 48 hours post infection (hpi) using RIPA buffer and tested for Atg8 expression as above. CHIKV samples were similarly harvested at 24 hpi. Band intensities of Atg8-PE and β-actin were determined by ImageJ. Autophagy induction was quantified by calculating the ratio of Atg8-PE to β-actin for each sample. The fold change in Atg-PE induction for each treatment was determined by comparing it to the DMSO control. All experiments were completed in triplicate.

Autophagy Gene Suppression

Aedes aegypti Atg5, Atg14, Atg8 and luciferase genes were amplified using primer sets containing the T7 promotor (Table S1). Amplicons were PCR purified and a T7 Megascript kit was used to synthesize dsRNA molecules for transfection per the manufacturer's recommendations (Ambion). Two days prior to transfection, Aag2 cells were seed at 1.5×10^6 cells/well in 12-well plates. 1 µg/well of dsRNA was combined with OPTI-MEM and Lipofectamine 2000 transfection reagent (Invitrogen) and allowed to incubate at room temperature for 20 minutes. The media on the Aag2 cells was discarded and replaced with 1 mL of Opti-MEM. The lipid-nucleic complexes were then added to the wells containing OPTI-MEM

and were incubated at 28°C for four hours, after which the media was discarded and again

replaced with the standard 10% FBS media. Cells were harvested 48 hours after transfection by

manually scraping the cell monolayer using a 1 ml pipette tip. The cells were pelleted and re-

suspended in QVL Lysis buffer from an Omega Bio-tek Viral RNA Extraction kit or RIPA

buffer. Total RNA was then extracted following the manufacturer's instructions. Extractions

were DNase treated and purified via phenol/chloroform extraction.

Silencing efficiency of Atg5 and Atg14 was determined through RT-qPCR with

Universal SYBR Green master mix (Bio-Rad), using luciferase samples as the non-targeting

control group and GAPDH as a reference gene (Table S1). Silencing efficiency of Atg8 was

determined by Western blotting. Two days after dsAtg8 was transfected into Aag2 cells, cells

were infected with DENV as above. Cells were harvested in RIPA buffer 24 hpi.

Immunoblotting was performed as previously described.

Virus titration

Two days after Aag2 cells were seeded at 1.5×10^6 cells/well in a 12-well plate, cells were

infected with DENV, ZIKV, or CHIKV at an MOI of 0.1. After infection for one hour at 28°C,

cells were treated in duplicate with 1% DMSO, 1 µM bafilomycin A1, 1 µM torin-1 or 10 µM

spautin-1. Virus supernatants were collected 24 and 48 hours after drug application. Finally,

Aag2 cells were seeded and transfected with dsLuciferase, dsAtg5, and dsAtg14 as previously

described. Two days after transfection, the cells were infected with DENV and virus

supernatants were harvested at 24 and 48 hpi.

The effect of cell line on DENV virus titers was determined by following the above

protocol in a mammalian cell line (BHK21 c15) and a mosquito cell line with a dysfunctional

RNA interference pathway (C6/36). C6/36 cells were seeded at 1.5×10^6 cells/well and BHK cells were seeded at 5×10^5 cells/ well in 12 well plates. The following day, both cell lines were infected with DENV and treated with 1% DMSO, 1 µM torin-1 or 10 µM spautin-1. Titers were harvested at 48 hpi. Each experiment was completed in triplicate with two biological replicates per experimental replicate.

For titration of DENV and ZIKV samples, BHK21 c15 cells were grown to a confluent monolayer in 12-well plates and infected with 10-fold serial dilutions of virus for 1 hour at room temperature and overlaid with a mixture of the standard media and 5% methyl cellulose. After incubation at 37°C for four days, the cells were fixed with 7.4% formaldehyde and stained with gentian violet. Viral titers were determined by counting plaques. For titration of CHIKV, the above protocol was followed in Vero cells.

Confocal Microscopy

In order to track autophagy during virus infection we quantified Atg8+ puncta by confocal microscopy. While we possess a polyclonal anti-Atg8 antibody, it did not produce reliable and consistent results. Therefore, we generated an expression vector to provide EGFP tagged Atg8 *in trans*. This was achieved by cloning the full length *Ae. aegypti* Atg8 gene (AAEL007162) into the pIEx-EGFP vector 3' to EGFP by Gibson Assembly. The generation of pIEx-EGFP has been described previously (25). Aag2 cells were seeded on coverslips at a density of 1.5×10^6 cells/well in a 12 well plate and 24 hrs. later transfected with 1.5 µg/ well of pIEx-EGFP-Atg8 using Lipofectamine 2000 transfection reagent according to the manufacturer's instructions. Two days post-transfection, cells were infected (MOI 0.1) and/ or chemically treated. Cells were fixed in 4% paraformaldehyde 24 hrs. post-treatment, washed 3X in warm PBS and mounted on microscope slides using ProLong Gold + DAPI. Microscopy was

performed on a Leica SP5 confocal microscope at 100x magnification with a 3x zoom. Because

the efficiency of transfection is relatively low in Aag2 cells, images were captured by scanning

the slide and identifying field of views that contained multiple cells with EGFP expression. Five

images per sample per experiment were acquired. Approximately 50 cells were imaged per

treatment. Experiments were completed in triplicate and all identifying information was blinded

to the microscopist. Quantification of puncta per cell was completed using the ImageJ software

with the Puncta Analyzer plug-in.

Statistical Analysis

Immunoblot fold-change differences and puncta abundance were analyzed by one-way ANOVA

with a Sidak's multiple comparisons test. All viral titer data was analyzed by one-way ANOVA

with a Dunnett's multiple comparisons test. Suppression of Atg5 and Atg14 was calculated with

a two-tailed t-test.

Results

Autophagy in Mosquito Cells

Because autophagy has been understudied in mosquitoes it is unclear if traditional methods for

manipulating and analyzing autophagy, such as the use of chemical compounds, will be effective

in mosquito cells. Consequently, we verified the ability of commonly used chemical compounds

to modulate autophagy in Aag2 cells. Treatment of Aag2 cells with bafilomycin A1 (inhibits

autophagosome/ lysosome fusion), torin-1 (induces autophagy) and spautin-1 (blocks autophagy)

alone had no discernible effects on the ratio of Atg8-PE to β-actin compared to the DMSO

control cells as determined by immunoblot (Fig. S1). However, the combination of torin-1 and

bafilomycin A1 resulted in a significant increase in Atg8-PE to β-actin ratios. By inhibiting

autophagolysosome acidification we were able to observe signs of autophagic activity suggesting

that Aag2 cells have a high rate of autophagic flux and providing evidence that torin-1 and

bafilomycin A1 are functioning as expected with regards to autophagy. As a result of the high

autophagic flux, the addition of spautin-1 had no effect on baseline autophagy levels. However,

when added to cells treated with torin-1 and bafilomycin A1, spautin-1 reduced the ratio of Atg8-

PE to β-actin thereby validating its ability to inhibit autophagy in Aag2 cells (Fig. S1).

Arbovirus Induced Autophagy

Numerous studies have demonstrated that arboviruses induce autophagy upon infection of

mammalian cells (12, 16, 17, 19-21). To examine if autophagy is similarly induced upon

arbovirus infection in mosquito cells, Aag2 cells were infected with DENV-2, ZIKV and CHIKV

and autophagic activity quantified. Infection with either DENV-2, ZIKV or CHIKV alone had no

effect on the ratio of Atg8-PE to β-actin as determined by immunoblot (Fig. 1A-B, 2A-B, 3A-B).

As previously determined, our inability to observe autophagy induction may be due to the high

rate of autophagic flux associated with these cells. Consequently, addition of bafilomycin A1 to

the DENV-2 infections significantly increased Atg8-PE to β-actin ratios (Fig. 1A-B). Similar

increases in the ratio could be observed for CHIKV infections, although these results were not

significant (Fig. 3A-B). Interestingly this did not increase ratios during ZIKV infection compared

to the bafilomycin A1 control group (Fig. 2A-B). To further validate these findings, Aag2 cells

were transfected with a plasmid constitutively expressing Atg8-EGFP and infected in the

presence or absence of bafilomycin A1. As with the immunoblot, DENV infections in the

absence of bafilomycin A1 resulted in a marginal increase in autophagic activity, while the

addition of bafilomycin A1 significantly increased Atg8+ puncta/ cell compared to the controls

(Fig. 1C-D). Similarly, increases in Atg8+ puncta/ cell were observed during ZIKV and CHIKV

infection in the presence of bafilomysin A1. It should be noted that infection status of each cell

could not be verified at the time of microscopy due to the rate of virus replication in these cells.

Samples were harvested 24 hpi, yet viral antigen is in low abundance at this early time point

making it difficult to detect signal above background auto-fluorescence. This limitation made it

difficult to correlate virus replication with Atg8+ puncta accumulation. Nevertheless, the totality

of the microscopy and immunoblot data provide confidence that, like mammalian systems,

arboviruses induce autophagy in mosquito cells.

Autophagy Arbovirus Interactions

Autophagy plays a central role in both innate and adaptive immunity (1). Numerous studies have

demonstrated that autophagy can be antiviral; however, some viruses have evolved mechanisms

to either circumvent autophagy or usurp components of the pathway for their own benefit (26).

To ascertain the role of autophagy during arbovirus infection of mosquitoes, the impacts of

pharmacological modulators of autophagy on arbovirus replication in Aag2 mosquito cells was

assessed. If autophagy were functioning in an antiviral manner, inhibition of autophagy would

result in increased viral titers and induction would result in decreased titers. Indeed the

application of spautin-1 resulted in significantly increased DENV-2 titers at 24 and 48 hpi, ZIKV

titers at 48 hpi, and CHIKV titers at 24 and 48 hpi (Fig. 4). Interestingly, induction of autophagy

with torin-1 also resulted in significantly increased titers for all three viruses at all time points

(Fig. 4). Further, blockage of V-ATPase activity with bafilomycin A1 resulting in an increase in

autophagolysosomal pH levels significantly increased DENV-2 and ZIKV titers at all time

points, but not those of CHIKV (Fig. 4). Due to the unexpected and paradoxical findings, we

examined the effects of autophagy through suppression of three genes within the autophagy

pathway: Atg5 (required for phagophore elongation), Atg8 (required for phagophore elongation

and membrane curvature) and Atg14 (essential autophagy regulator). Treatment of Aag2 cells

with gene specific dsRNA resulted in significant suppression of Atg5 and Atg14 as determined

by RT-qPCR as well as depletion of Atg8 as determined by immunoblot (Fig. S2). Suppression

of the autophagy related genes had no discernible effect on DENV-2 titers 24 or 48 hpi compared

to the control (Fig. 5A). Likewise suppression of Atg5 and Atg14 had no effect on CHIKV titers

at 24 or 48 hpi (Fig. 5B). Together these results suggest that autophagy has little effect on

arbovirus replication in Aag2 cells. Further, it suggests that, in addition to regulating autophagy,

commonly used chemical modulators of autophagy can have beneficial side effects on viral

replication through potentially non-autophagy mediated mechanisms.

Autophagy Response to Arboviruses in Other Systems

Due to the unexpected results associated with chemical treatment in Aag2 cells, we examined if

the observed outcomes were consistent in other host systems. Others have previously shown that

spautin-1 mediated inhibition of autophagy in mammalian cells, specifically BHK-21, drastically

reduces DENV-2 titers (21). In an effort to recapitulate these findings we treated DENV-2

infected BHK-21 cells with spautin-1 and observed a significant reduction in DENV-2 titers

(Fig. 6). Induction of autophagy with torin-1 had no effect (Fig. 6). Next we wanted to determine

if our results in Aag2 cells were similar in another commonly used mosquito cell line, C6/36

Aedes albopictus cells. As before, treatment of Aag2 cells with both torin-1 and spautin-1

resulted in significantly increased DENV-2 titers; however, we observed a slight, but significant

decrease in titers upon torin-1 treatment and no effect upon spautin-1 treatment in C6/36 cells (Fig. 7). This data suggests that these commonly used chemical compounds can have profoundly different effects on host cell-virus interactions.

Discussion

Manipulation of the autophagy pathway has become a promising approach for treating a number of diseases including viral infections (6, 7, 18). Because arboviruses are maintained in a two-host system, the invertebrate vector and vertebrate host, there exists the possibility that autophagy can be manipulated in one or both systems in order to reduce disease burden and/ or transmission. Despite this two-host dependency, our understanding of the autophagy-arbovirus interface is primarily informed by one host, mammals (22). It is known that virus infections can induce an autophagic response in Diptera, specifically *Drosophila*; however, there are conflicting reports as to the role of autophagy during infection (27-30). Aside from having an intact and functional autophagy pathway, little is known about autophagy in mosquitoes (24). In an attempt to close this knowledge gap, this study characterized the autophagic response to arbovirus infection in the tractable Aag2 *Aedes aegypti* cell culture system. Our findings suggest that while autophagy is induced, it seems to have a limited role during arbovirus infection of mosquito cell culture. Further, we demonstrate that while commonly used chemicals can modulate the autophagy pathway in mosquito cells, their effects on viruses are largely cell line dependent.

In agreement with a previous report, we found spautin-1 treatment significantly reduced the release of infectious DENV particles during infection of mammalian cells (21). In addition, Mateo et al. also observed that levels of extracellular viral RNA remained unchanged compared

to the controls and that DENV particles recovered from spautin-1 treated cells retained their 'pr'

peptide after exocytosis (21). During assembly, the viral nucleocapsid composed of genomic

RNA and capsid protein, is packaged into a host-cell-derived envelope containing 180 copies of

each of the E and prM structural proteins. Subsequently, immature virions undergo maturation in

the trans-golgi network whereby the 'pr' peptide is proteolytically removed from M by host furin

(31). 'pr' remains in association with the virion through exocytosis after which it disassociates.

Improper cleavage or disassociation can result in the production of immature particles which

have been shown to be less infectious than the fully mature virions (32). Based on this evidence

it was hypothesized that either retention of the 'pr' peptide or altered virion confirmation

diminished DENV infectiousness after spautin-1 treatment (21). However, this outcome appears

to be limited to mammalian cells as we observed increased infectiousness after spautin-1

treatment of infected Aag2 mosquito cells. The reasons for this are unknown. A previous study

demonstrated that furin substrate specificity differs between vertebrate and invertebrate cells,

which may explain why there is a higher proportion of immature virions produced in mosquito

cells compared to mammalian cells (33). Interestingly, there doesn't appear to be a reduction in

infectiousness. One potential explanation could be that DENV-2 adopts different conformational

arrangements in mosquito and mammalian cells. In fact, at 28°C, the temperature at which

mosquito cells are cultured, mature DENV virions contain 90 E-glycoprotein homodimers

arranged in a tightly packed antiparallel formation with a smooth appearance (34). Conversely, at

37°C, E-glycoprotein homodimers adopt a very different conformation in which virions appear

bumpy with large gaps allowing visualization of the underlying lipid membrane bilayer (34). It is

possible that 'pr' cleavage and disassociation are not required to form infectious virions at 28°C

due to the observed conformational rearrangements and therefore, spautin-1 has no adverse

effects and potentially beneficial effects on the production of infectious virus particles. This

could explain the differences observed between mosquito cells and mammalian cells upon

spautin-1 treatment.

Paradoxically, chemical induction and suppression of autophagy and inhibition of

autophagolysosomal acidification all resulted in increased DENV-2, ZIKV and CHIKV titers in

Aag2 cells. In an attempt to reconcile these conflicting results, we performed RNAi suppression

of autophagy genes. For both DENV-2 and CHIKV, no effects were observed upon suppression

of Atg5, Atg14 or Atg8 despite high levels of suppression. Such results suggest that autophagy

plays a limited role in mediating arbovirus infection of mosquito cells and that the effects

conferred by the chemicals may be indirectly beneficial. Alternatively, it could suggest that

>80% suppression was not sufficient to diminish autophagic activity and therefore, the RNAi

suppression infection assays were not sensitive enough to identify a role for autophagy during

arbovirus infection. A third possible explanation could be that mosquitoes encode functional

redundancy in key autophagic genes. It is known that Atg8 is part of a larger family of homologs

that include genes such as GABARAP and GABARAPL. Recent work has shown that autophagy

remains functional upon deletion of any of these homologs individually (35). Further, humans

encode four isoforms of Atg4 and that elimination of all four is necessary to eliminate cleavage

of pro-LC3/GABARAP prior to lipidation (36). While additional homologs and/ or isoforms of

Atg5 and Atg14 have not been identified in other systems, functional redundancy may exist in

mosquitoes.

In this study we observed a significant increase in viral titers upon spautin-1 and torin-1

treatment in Aag2 cells; however, we observed no effect in another mosquito cell line, C6/36

cells. While both are mosquito cell lines, there are significant differences that could affect the

chemical-arbovirus-cell interface. Aag2 cell are derived from *Ae. aegypti* mosquitoes whereas

C6/36 cells originate from *Ae. albopictus* mosquitoes; however, due to the evolutionary

conservation of autophagy it is unlikely that there are significant species-level differences in

autophagic activity. A potentially more important factor could be the tissue source of the cells

(37). Mosquito cell culture is initiated by grinding up larvae and recovering cells amenable to

growth in culture. Consequently, the specific cell type of the cells in culture is unknown. It could

be that the differences arise because they are derived from different tissues and therefore respond

differently to DENV in the presence of chemicals. In addition, previous reports have

demonstrated that Aag2 cells are persistently infected with multiple insect-only viruses while

C6/36 cells are not (38). It is possible that additional viruses in the system could alter how the

cells respond to both chemicals and arboviruses. Finally, the primary antiviral immune response,

RNAi, is functionally impaired in C6/36 cells (39). How this would impact the host cells'

response to chemicals targeting the autophagy pathway during viral infection is unclear;

however, previous studies have demonstrated that autophagy can selectively degrade

components of the RNAi pathway (40, 41). If RNAi is actively targeting viruses during

infection, than induction with torin-1 could lead to increased viral replication in Aag2 cells but

not C6/36 cells. Conversely, suppression of autophagy with spautin-1 would in theory increase

available RNAi machinery thereby enabling a more potent RNAi response and reduction of viral

titers; however, this is not what was observed. This may reflect the fact that RNAi genes are not

significantly upregulated during arbovirus infection so suppression of autophagy has little effect

on the normal levels of RNAi while increased autophagic activity could reduce activity allowing

the viruses to replicate to higher titers (42). Further research will be needed to fully elucidate

autophagy-RNAi interactions and how they impact arbovirus infection of mosquitoes.

Comparing our results to what has been reported in mammalian systems reveals that at

least for DENV and ZIKV the general patterns are consistent. The increase in titers upon TOR-

inhibition with torin-1 suggests that both DENV-2 and ZIKV benefit from downstream pathways

regulated by TOR, one of which is autophagy. We observed increased flavivirus titers in Aag2

cells upon torin-1 treatment suggesting that flaviviruses benefit from an autophagic state and is

consistent with what has been found in mammalian cells (9, 21). However, increased titers were

also observed upon suppression of autophagy by spautin-1 suggesting an antiviral role. While

these outcomes seem contradictory, they are consistent to the current model for DENV-

autophagy interactions. In mammalian cells, DENV and DENV replication complexes are

targeted for degradation by autophagy while simultaneously benefitting from more efficient

mature virion processing and increased energy availability as a result of virus-induced lipophagy

(9, 19-21). While the flavivirus data was consistent with what has been observed in mammalian

systems, the CHIKV data was not. Numerous studies have found that autophagy functions in an

antiviral manner during infection with members of the family *Togaviridae*, specifically

alphaviruses (10, 13, 15, 17). While inhibition of autophagy did result in increased CHIKV titers

as would be expected, induction also increased titers. This discrepancy may reflect a difference

in the role of autophagy during CHIKV infection of mosquito and mammalian cells.

This study provides an important first-step in understanding autophagy-arbovirus

interactions within mosquitoes and highlights differences between the mammalian and mosquito

autophagic response to infection. However, future work assessing these interactions within

mosquitoes will be needed due to the limitations of our cell culture system. As previously

mentioned, the origin of mosquito cell culture is unknown and therefore, it is difficult to

determine if, in the context of natural infection, virus-autophagy interactions may function

differently. Further, during infection, arboviruses must infect and replicate in a diversity of

tissues and cell types before being transmitted. Previous studies have shown that autophagy can

function differently between cell types and tissue types, it is therefore possible that during the

course of mosquito infection autophagy might interact differently with viruses depending on the

tissue and cell type (37). In addition, a recent study highlighted the genomic differences observed

between C6/36 *Aedes albopictus* cells and *Aedes albopictus* mosquitoes (43). The cells used in

this study have been in culture for decades and may not accurately recapitulate the cellular

environment of *Aedes aegypti*. Regardless, this study better defines autophagy-arbovirus

interactions in a medically relevant vector system and provides insights that will be important for

future *in vivo* studies.

Acknowledgements

We would like to thank Dr. Gregory Ebel and Benjamin Dodd for their thoughtful insights and early support of this work. This work was supported in part by grants from the Centers for Disease Control and Prevention (U01/CK000509-01) and the National Institute of Health, National Institute of Allergy and Infectious Diseases (AI099042).

References

1. **Jang YJ, Kim JH, Byun S.** 2019. Modulation of Autophagy for Controlling Immunity. Cells **8**.
2. **Mizushima N, Levine B.** 2010. Autophagy in mammalian development and differentiation. Nat Cell Biol **12**:823-830.
3. **Jiang P, Mizushima N.** 2014. Autophagy and human diseases. Cell Res **24**:69-79.
4. **Levine B, Klionsky DJ.** 2004. Development by self-digestion: molecular mechanisms and biological functions of autophagy. Dev Cell **6**:463-477.

1   5.   **Gatica D, Lahiri V, Klionsky DJ.** 2018. Cargo recognition and degradation by selective autophagy. Nat Cell Biol **20:**233-242.

6.   **Rubinsztein DC, Codogno P, Levine B.** 2012. Autophagy modulation as a potential therapeutic target for diverse diseases. Nat Rev Drug Discov **11:**709-730.

7.   **Zhan L, Li J, Wei B.** 2018. Autophagy therapeutics: preclinical basis and initial clinical studies. Cancer Chemother Pharmacol **82:**923-934.

8.   **Blazquez AB, Martin-Acebes MA, Saiz JC.** 2014. Amino acid substitutions in the non-structural proteins 4A or 4B modulate the induction of autophagy in West Nile virus infected cells independently of the activation of the unfolded protein response. Front Microbiol **5:**797.

9.   **Heaton NS, Randall G.** 2010. Dengue Virus-Induced Autophagy Regulates Lipid Metabolism. Cell Host & Microbe **8:**422-432.

10.   **Joubert P-E, Werneke SW, de la Calle C, Guivel-Benhassine F, Giodini A, Peduto L, Levine B, Schwartz O, Lenschow DJ, Albert ML.** 2012. Chikungunya virus–induced autophagy delays caspase-dependent cell death. The Journal of Experimental Medicine **209:**1029-1047.

11.   **Khongwichit S, Wikan N, Abere B, Thepparit C, Kuadkitkan A, Ubol S, Smith DR.** 2016. Cell-type specific variation in the induction of ER stress and downstream events in chikungunya virus infection. Microb Pathog **101:**104-118.

12.   **Liang Q, Luo Z, Zeng J, Chen W, Foo SS, Lee SA, Ge J, Wang S, Goldman SA, Zlokovic BV, Zhao Z, Jung JU.** 2016. Zika Virus NS4A and NS4B Proteins Deregulate Akt-mTOR Signaling in Human Fetal Neural Stem Cells to Inhibit Neurogenesis and Induce Autophagy. Cell Stem Cell **19:**663-671.

13.   **Liang XH, Kleeman LK, Jiang HH, Gordon G, Goldman JE, Berry G, Herman B, Levine B.** 1998. Protection against fatal Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. J Virol **72:**8586-8596.

14.   **Sharma M, Bhattacharyya S, Nain M, Kaur M, Sood V, Gupta V, Khasa R, Abdin MZ, Vrati S, Kalia M.** 2014. Japanese encephalitis virus replication is negatively regulated by autophagy and occurs on LC3-I- and EDEM1-containing membranes. Autophagy **10:**1637-1651.

15.   **Eng KE, Panas MD, Murphy D, Karlsson Hedestam GB, McInerney GM.** 2012. Accumulation of Autophagosomes in Semliki Forest Virus-Infected Cells Is Dependent on Expression of the Viral Glycoproteins. Journal of Virology **86:**5674-5685.

16.   **Judith D, Mostowy S, Bourai M, Gangneux N, Lelek M, Lucas-Hourani M, Cayet N, Jacob Y, Prevost MC, Pierre P, Tangy F, Zimmer C, Vidalain PO, Couderc T, Lecuit M.** 2013. Species-specific impact of the autophagy machinery on Chikungunya virus infection. EMBO Rep **14:**534-544.

17.   **Orvedahl A, MacPherson S, Sumpter Jr R, Tallóczy Z, Zou Z, Levine B.** 2010. Autophagy Protects against Sindbis Virus Infection of the Central Nervous System. Cell Host & Microbe **7:**115-127.

18.   **Shoji-Kawata S, Sumpter R, Leveno M, Campbell GR, Zou Z, Kinch L, Wilkins AD, Sun Q, Pallauf K, MacDuff D, Huerta C, Virgin HW, Helms JB, Eerland R, Tooze SA, Xavier R, Lenschow DJ, Yamamoto A, King D, Lichtarge O, Grishin NV, Spector SA, Kaloyanova DV, Levine B.** 2013. Identification of a candidate therapeutic autophagy-inducing peptide. Nature **advance online publication**.

19.   **Metz P, Chiramel A, Chatel-Chaix L, Alvisi G, Bankhead P, Mora-Rodriguez R, Long G, Hamacher-Brady A, Brady NR, Bartenschlager R.** 2015. Dengue Virus Inhibition of Autophagic Flux and Dependency of Viral Replication on Proteasomal Degradation of the Autophagy Receptor p62. J Virol **89:**8026-8041.

20.   **Lennemann NJ, Coyne CB.** 2017. Dengue and Zika viruses subvert reticulophagy by NS2B3-mediated cleavage of FAM134B. Autophagy **13:**322-332.

21. **Mateo R, Nagamine CM, Spagnolo J, Mendez E, Rahe M, Gale M, Jr., Yuan J, Kirkegaard K.** 2013. Inhibition of cellular autophagy deranges dengue virion maturation. J Virol **87:**1312-1321.

22. **Brackney DE.** 2017. Implications of autophagy on arbovirus infection of mosquitoes. Curr Opin Insect Sci **22:**1-6.

23. **Liu J, Xia H, Kim M, Xu L, Li Y, Zhang L, Cai Y, Norberg HV, Zhang T, Furuya T, Jin M, Zhu Z, Wang H, Yu J, Li Y, Hao Y, Choi A, Ke H, Ma D, Yuan J.** 2011. Beclin1 Controls the Levels of p53 by Regulating the Deubiquitination Activity of USP10 and USP13. Cell **147:**223-234.

24. **Bryant B, Raikhel AS.** 2011. Programmed Autophagy in the Fat Body of *Aedes aegypti* Is Required to Maintain Egg Maturation Cycles. Plos One **6:**e25502.

25. **Scott JC, Brackney DE, Campbell CL, Bondu-Hawkins V, Hjelle B, Ebel GD, Olson KE, Blair CD.** 2010. Comparison of Dengue Virus Type 2-Specific Small RNAs from RNA Interference-Competent and –Incompetent Mosquito Cells. PLoS Negl Trop Dis **4:**e848.

26. **Kudchodkar SB, Levine B.** 2009. Viruses and autophagy. Reviews in Medical Virology **19:**359-378.

27. **Lamiable O, Arnold J, de Faria IJ, Olmo RP, Bergami F, Meignin C, Hoffmann JA, Marques JT, Imler JL.** 2016. Analysis of the Contribution of Hemocytes and Autophagy to Drosophila Antiviral Immunity. J Virol **90:**5415-5426.

28. **Moy Ryan H, Gold B, Molleston Jerome M, Schad V, Yanger K, Salzano M-V, Yagi Y, Fitzgerald Katherine A, Stanger Ben Z, Soldan Samantha S, Cherry S.** 2014. Antiviral Autophagy Restricts Rift Valley Fever Virus Infection and Is Conserved from Flies to Mammals. Immunity **40:**51-65.

29. **Nakamoto M, Moy Ryan H, Xu J, Bambina S, Yasunaga A, Shelly Spencer S, Gold B, Cherry S.** 2012. Virus Recognition by Toll-7 Activates Antiviral Autophagy in Drosophila. Immunity **36:**658-667.

30. **Shelly S, Lukinova N, Bambina S, Berman A, Cherry S.** 2009. Autophagy Is an Essential Component of Drosophila Immunity against Vesicular Stomatitis Virus. Immunity **30:**588-598.

31. **Dowd KA, Pierson TC.** 2018. The Many Faces of a Dynamic Virion: Implications of Viral Breathing on Flavivirus Biology and Immunogenicity. Annu Rev Virol **5:**185-207.

32. **Zybert IA, van der Ende-Metselaar H, Wilschut J, Smit JM.** 2008. Functional importance of dengue virus maturation: infectious properties of immature virions. J Gen Virol **89:**3047-3051.

33. **Cano-Monreal GL, Williams JC, Heidner HW.** 2010. An arthropod enzyme, Dfurin1, and a vertebrate furin homolog display distinct cleavage site sequence preferences for a shared viral proprotein substrate. J Insect Sci **10:**29.

34. **Zhang X, Sheng J, Plevka P, Kuhn RJ, Diamond MS, Rossmann MG.** 2013. Dengue structure differs at the temperatures of its human and mosquito hosts. Proc Natl Acad Sci U S A **110:**6795-6799.

35. **Nguyen TN, Padman BS, Usher J, Oorschot V, Ramm G, Lazarou M.** 2016. Atg8 family LC3/GABARAP proteins are crucial for autophagosome–lysosome fusion but not autophagosome formation during PINK1/Parkin mitophagy and starvation. The Journal of Cell Biology **215:**857-874.

36. **Agrotis A, Pengo N, Burden JJ, Ketteler R.** 2019. Redundancy of human ATG4 protease isoforms in autophagy and LC3/GABARAP processing revealed in cells. Autophagy **15:**976-997.

37. **Chapin HC, Okada M, Merz AJ, Miller DL.** 2015. Tissue-specific autophagy responses to aging and stress in C. elegans. Aging (Albany NY) **7:**419-434.

38. **Weger-Lucarelli J, Rückert C, Grubaugh ND, Misencik MJ, Armstrong PM, Stenglein MD, Ebel GD, Brackney DE.** 2018. Adventitious viruses persistently infect three commonly used mosquito cell lines. Virology **521:**175-180.

1   39.   **Brackney DE, Scott JC, Sagawa F, Woodward JE, Miller NA, Schilkey FD, Mudge J, Wilusz J,**
2   **Olson KE, Blair CD, Ebel GD.** 2010. C6/36 Aedes albopictus cells have a dysfunctional antiviral
3   RNA interference response. PLoS Negl Trop Dis **4:**e856.
4   40.   **Derrien B, Baumberger N, Schepetilnikov M, Viotti C, De Cillia J, Ziegler-Graff V, Isono E,**
5   **Schumacher K, Genschik P.** 2012. Degradation of the antiviral component ARGONAUTE1 by the
6   autophagy pathway. Proceedings of the National Academy of Sciences **109:**15942-15946.
7   41.   **Gibbings D, Mostowy S, Jay F, Schwab Y, Cossart P, Voinnet O.** 2012. Selective autophagy
8   degrades DICER and AGO2 and regulates miRNA activity. Nature cell biology **14:**1314-1321.
42.   **Campbell C, Keene K, Brackney D, Olson K, Blair C, Wilusz J, Foy B.** 2008. *Aedes aegypti* uses
RNA interference in defense against Sindbis virus infection. BMC Microbiology **8:**47.
43.   **Miller JR, Koren S, Dilley KA, Puri V, Brown DM, Harkins DM, Thibaud-Nissen F, Rosen B, Chen**
X-G, Tu Z, Sharakhov IV, Sharakhova MV, Sebra R, Stockwell TB, Bergman NH, Sutton GG,
Phillippy AM, Piermarini PM, Shabman RS. 2018. Analysis of the Aedes albopictus C6/36
genome provides insight into cell line utility for viral propagation. GigaScience **7:**1-13.

21   **Tables and Figures:**

22   Table 1: Primer List

Gene	Primer	Sequence (5'-3')	Accession #
AaeAtg5	T7-Forward	TAA TAC GAC TCA CTA TAG GGT CCG ATG AAA CCG ATG TC	XM_001661191
	T7-Reverse	TAA TAC GAC TCA CTA TAG GGC ATC GAA CTG GAA TGT G	
	RT-qPCR-Forward	AGT TCG ATG TTA TGC CGA GG	
	RT-qPCR-Reverse	GAT AGC TGA GGT GTT CCG AG	
AaeAtg14	T7-Forward	TAA TAC GAC TCA CTA TAG GGT TTC TCG AGC AGT GAC GGA	XM_001652367
	T7-Reverse	TAA TAC GAC TCA CTA TAG GCG CAA TTC GTT CAA CTG GCA	
	RT-qPCR-Forward	ACT CGA AAC GGC ACT TCC AT	
	RT-qPCR-Reverse	TGT GGC GTA GTT GCT TCT CG	
AaeAtg8	T7-Forward	TAA TAC GAC TCA CTA TAG GAT GAA ATT TCA ATA CAA G	XM_001652521
	T7-Reverse	TAA TAC GAC TCA CTA TAG GAC TTG TTT CCA TAC ACA TTC	

AaeGAPDH	RT-qPCR-Forward	ACG TGA ACA GAC GCT AGT TAT	XM_011494724
	RT-qPCR-Reverse	GTG GGT CGA ATC GTA CTT GAA	

Figure Legends:

Figure 1: Dengue virus 2 induces autophagy in *Aedes aegypti* Aag2 cells. A) Representative

immunoblot of Aag2 lysates upon DENV-2 infection (M.O.I. 0.5) and/ or chemical treatment

(1% DMSO, 1 µM bafilomycin A1, 1 µM torin-1 or 10 µM spautin-1) 48 hpi. B) Fold-change in

the ratio of Atg8-PE to β-actin band intensities as determined by ImageJ. Includes data from four

experimental replicates. C) Representative confocal microscopy images of Atg8-EGFP

expressing Aag2 cells ± DENV-2 infection (M.O.I. 0.1) and ±1 µM bafilomycin A1 24 hpi. D)

The number of Atg8+ puncta were quantified using the ImageJ Puncta Analyzer plug-in from

~50 Atg8-EGFP expressing Aag2 cells ± DENV-2 infection (M.O.I. 0.1) and ±1 µM bafilomycin

A1 24 hpi. Combined data from three blinded experimental replicates. Data were analyzed by

one-way ANOVA with a Sidak's multiple comparisons test. (*) $p<0.05$, (**) $p<0.01$, (***)

$p<0.001$.

Figure 2: Zika virus induces autophagy in *Aedes aegypti* Aag2 cells. A) Representative

immunoblot of Aag2 lysates upon ZIKV infection (M.O.I. 0.5) and/ or chemical treatment (1%

DMSO, 1 µM bafilomycin A1, 1 µM torin-1 or 10 µM spautin-1) 48 hpi. B) Fold-change in the

ratio of Atg8-PE to β-actin band intensities as determined by ImageJ. Includes data from four

experimental replicates. C) Representative confocal microscopy images of Atg8-EGFP

expressing Aag2 cells ± ZIKV infection (M.O.I. 0.1) and ±1 µM bafilomycin A1 24 hpi. D) The

number of Atg8+ puncta were quantified using the ImageJ Puncta Analyzer plug-in from ~50

Atg8-EGFP expressing Aag2 cells ± ZIKV infection (M.O.I. 0.1) and ±1 µM bafilomycin-A1 24

hpi. Combined data from three blinded experimental replicates. Data were analyzed by One-way

ANOVA with a Sidak's multiple comparisons test. (*) $p<0.05$, (**) $p<0.01$, (***) $p<0.001$.

Figure 3: Chikungunya virus induces autophagy in *Aedes aegypti* Aag2 cells. A)

Representative immunoblot of Aag2 lysates upon CHIKV infection (M.O.I. 0.5) and/ or

chemical treatment (1% DMSO, 1 µM bafilomycin A1, 1 µM torin-1 or 10 µM spautin-1) 24

hpi. B) Fold-change in the ratio of Atg8-PE to β-actin band intensities as determined by ImageJ.

Includes data from four experimental replicates. C) Representative confocal microscopy images

of Atg8-EGFP expressing Aag2 cells ± CHIKV infection (M.O.I. 0.1) and ±1 µM bafilomycin

A1 24 hpi. D) The number of Atg8+ puncta were quantified using the ImageJ Puncta Analyzer

plug-in from ~50 Atg8-EGFP expressing Aag2 cells ± CHIKV infection (M.O.I. 0.1) and ±1 µM

bafilomycin A1 24 hpi. Combined data from three blinded experimental replicates. Data were

analyzed by one-way ANOVA with a Sidak's multiple comparisons test. (*) $p<0.05$, (**)

$p<0.01$, (***) $p<0.001$.

Figure 4: Pharmacological modulators of autophagy uniformly increase DENV, ZIKV and

CHIKV titers in Aag2 cells. *Aedes aegypti* Aag2 cells were infected with A) DENV-2, B)

ZIKV, and C) CHIKV followed by chemical treatment (1% DMSO, 1 µM bafilomycin A1, 1 µM

torin-1 or 10 µM spautin-1). Samples were collected for titration 24 and 48 hpi for DENV-2 and

CHIKV and 48 hpi for ZIKV. Data was analyzed by one-way ANOVA with a Dunnett's multiple

comparisons test. (*) $p<0.05$, (**) $p<0.01$, (***) $p<0.001$.

Figure 5: Suppression of autophagy related genes has no effect on DENV-2 or CHIKV

titers in Aag2 cells. *Aedes aegypti* Aag2 cells were treated with dsRNA against Atg5, Atg14,

Atg8, and non-specific control luciferase genes two days prior to infection with either A) DENV-

2 or B) CHIKV. Samples were collected for titration 24 and 48 hpi. Data was analyzed by one-

way ANOVA with a Dunnett's multiple comparisons test. (*) $p<0.05$, (**) $p<0.01$, (***)

$p<0.001$.

Figure 6. Pharmacological inhibition of autophagy decreases DENV titers in BHK-21 cells.

BHK-21 c15 cells were infected with DENV-2 and then treated with either torin-1 (1 µM),

spautin-1 (10 µM), or 1% DMSO as a control. Samples were collected 48 hpi for titration. Data

was analyzed by one-way ANOVA with a Dunnett's multiple comparisons test. (*) $p<0.05$, (**)

$p<0.01$, (***) $p<0.001$.

Figure 7. Pharmacological modulators of autophagy increase DENV titers in Aag2 cells,

but not C6/36 cells. Both Aag2 and *Aedes albopictus* C6/36 cells were infected with DENV-2

and then treated with either torin-1 (1 µM), spautin-1 (10 µM), or 1% DMSO as a control.

Samples were collected 48 hpi for titration. Data was analyzed by one-way ANOVA with a

Dunnett's multiple comparisons test. (*) $p<0.05$, (**) $p<0.01$, (***) $p<0.001$.

Figure S1: Common pharmacological manipulators of autophagy function as expected in mosquito cells. A) Representative immunoblot of Aag2 lysates upon chemical treatment (1% DMSO, 1 μM bafilomycin A1, 1 μM torin-1 or 10 μM spautin-1) 24 hpi. B) Fold-change in the ratio of Atg8-PE to β-actin band intensities as determined by ImageJ. Includes data from three experimental replicates. Data were analyzed by one-way ANOVA with a Sidak's multiple comparisons test. (*) $p < 0.05$, (**) $p < 0.01$, (***) $p < 0.001$.

Figure S2: Efficient silencing of autophagy genes in mosquito cells: Aag2 cells were treated with dsRNA targeting Atg5, Atg14, and Atg8 and assayed for suppression. Silencing efficiency of A) Atg5 and B) Atg14 was determined by $\Delta\Delta C_T$ analysis with luciferase samples as the non-targeting control group and GAPDH as a reference gene. Data was analyzed with a two-tailed t-test. C) Silencing efficiency of Atg8 was determined by immunoblot. Two days after dsAtg8 was transfected into Aag2 cells, cells were infected with DENV as above. Cells were harvested in RIPA buffer 24 hpi. Immunoblotting was performed as previously described.

Figure 1

PE

n

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7