1   Evolutionary and biochemical analyses reveal conservation of the Brassicaceae telomerase

2   ribonucleoprotein complex

4   Kelly Dew-Budd[2], Julie Cheung[2], Anna Nelson-Dittrich, Kyle Palos, Evan S. Forsythe, Andrew

5   D. L. Nelson and Mark A. Beilstein[1]

7   School of Plant Sciences, University of Arizona, Tucson, Arizona 85721

9   [1]Corresponding Author: Mark A. Beilstein

10   School of Plant Sciences

11   University of Arizona

12   Tucson, Arizona 85721

13   mbeilstein@email.arizona.edu

16   [2]These authors contributed equally

Abstract:

The telomerase ribonucleoprotein complex (RNP) is essential for genome stability and performs

this role through the addition of repetitive DNA to the ends of chromosomes. The telomerase

enzyme is composed of a reverse transcriptase (TERT), which utilizes a template domain in an

RNA subunit (TER) to reiteratively add telomeric DNA at the ends of chromosomes. Multiple

TERs have been identified in the model plant *Arabidopsis thaliana.* Here we combine a

phylogenetic and biochemical approach to understand how the telomerase RNP has evolved in

Brassicaceae, the family that includes *A. thaliana*. Because of the complex phylogenetic pattern

of template domain loss and alteration at the previously characterized *A. thaliana* TER loci,

TER1 and *TER2*, across the plant family Brassicaceae, we bred double mutants from plants

with a template deletion at *AtTER1* and T-DNA insertion at *AtTER2*. These double mutants

exhibited no telomere length deficiency, a definitive indication that neither of these loci encode a

functional telomerase RNA. Moreover, we determined that the telomerase components *TERT*,

Dyskerin, and the *KU* heterodimer are under strong purifying selection, consistent with the idea

that the TER with which they interact is also conserved. To test this hypothesis further, we

analyzed the substrate specificity of telomerase from species across Brassicaceae and

determined that telomerase from close relatives bind and extend substrates in a similar manner,

supporting the idea that TERs in different species are highly similar to one another and are likely

encoded from an orthologous locus. Lastly, TERT proteins from across Brassicaceae were able

to complement loss of function *tert* mutants *in vivo*, indicating TERTs from other species have

the ability to recognize the native TER of *A. thaliana*. Finally, we immunoprecipitated the

telomerase complex and identified associated RNAs via RNA-seq. Using our evolutionary data

we constrained our analyses to conserved RNAs within Brassicaceae that contained a template

domain. These analyses revealed a highly expressed locus whose disruption by a T-DNA

resulted in a telomeric phenotype similar to the loss of other telomerase core proteins, indicating

that the RNA has an important function in telomere maintenance.

Introduction

Chromosomes in most eukaryotes are capped by tandem TG-rich DNA repeats called

telomeres. The telomere repeat is remarkably conserved across deep evolutionary divergences.

For example, the repeat sequence TTTAGGG present in the majority of examined plants differs

by a single nucleotide from the vertebrate repeat TTAGGG [1]. Telomeric DNA repeats are

recognized and bound by a suite of proteins that are essential for chromosome end protection

and telomere replication [2], while telomere length is maintained by a ribonucleoprotein (RNP)

complex, which minimally consists of a telomerase reverse transcriptase (TERT) and a

telomerase RNA (TER). A template region within TER is complementary to approximately 1.5x

the telomere repeat and is used by TERT to synthesize telomeric DNA at the chromosome

terminus [3]. Although TERT and TER are sufficient to catalyze telomerase activity *in vitro*, *in*

vivo TER serves as a scaffold for the assembly of essential accessory proteins that have a

variety of functions including RNP biogenesis and recruitment of the RNP to the chromosome

end [4]. Many of these accessory proteins do not interact directly, but only associate with one

another via their interactions with TER [5,6]. Thus, TER is essential both as a template for

telomere synthesis, and as a core scaffolding molecule in telomerase assembly.

TERT proteins from distantly related species across the eukaryotic tree are readily

identifiable by homology searches based on amino acid sequence [3]. Given the essential role

of TER in telomere repeat addition and binding accessory proteins, a logical expectation would

be that TER would also display high levels of sequence conservation. However, TERs from

different eukaryotic lineages are highly variable at the nucleotide level and appear to be entirely

non-homologous, and thus independently evolved in each of the major eukaryotic lineages (i.e.,

animals, ciliates, fungi, and plants) [7,8]. Interestingly, most described TERs have converged on

similar structural features that likely result from the shared requirement to bind both TERT and

telomerase accessory proteins [6]. Two such core TER features include a pseudoknot, which is

necessary for activity, and a stem loop domain (termed CR4/CR5, stem loop IV, and three way

junction (TWJ) in vertebrates, ciliates, ascomycetes, respectively), which is necessary for TERT

binding [9–12].

Bioinformatics approaches to recover TER have limits across major lineages of the

eukaryotic crown group, but within eukaryotic clades TERs have been successfully recovered

using a variety of sequence similarity based approaches. For example, in vertebrates Chen et

al. (2000) recovered TER from species across the ~450 million year radiation using sequence

similarity searches and positional conservation (synteny) to identify eight conserved domains

[10]. Similarly, budding yeast and its closest relatives in *Saccharomyces sensu stricto*, a

radiation spanning ~20 million years of evolution, encode TER at a syntenic locus, a fact that

permitted bioinformatic detection of structurally conserved domains through signatures of

covariation [13]. In the Pezizomycotina (Ascomycota), with the exception of *Saccharomyces*

s.s., TERs display sufficient conservation to facilitate identification by BLAST using the

Neurospora crassa TER as query. Similarly, using a combination of template sequence

identification and structural motif modeling from Pezizomycotina TERs, Qi et al. (2013)

recovered TERs from the more distantly related Taphrinomycotina [12]. Thus, while biochemical

approaches have been required to identify TER in each major crown group lineage of

eukaryotes, bioinformatics approaches have aided discovery of TER within lineages.

Until recently, the only known functional TER in plants was from *Arabidopsis thaliana*.

Cifuentes-Rojas et al. (2011) found that *A. thaliana* was unusual among studied eukaryotes

because it encoded two TERs: *TER1* and *TER2* [14]. *TER1* was hypothesized to serve the

canonical function in providing a template for telomere addition by telomerase *in vivo*. *TER2* has

the same template domain as *TER1*, but it is encoded at a different locus and has been

hypothesized to have an alternative function in regulating telomerase activity during genome

stress caused in part by double strand DNA breaks [15,16]. Both *AtTER1* and *AtTER2* were

shown to assemble *in vivo* into a telomerase RNP that includes *TERT*, although the accessory

proteins with which they assemble differ [14]. Importantly, the *TER2* RNP has not been

observed to contribute to telomere length maintenance.

Attempts to use bioinformatics approaches to identify the TER-encoding locus or loci

similar to *AtTER1* or *AtTER2* from other Brassicaceae have yielded surprising results. For

example, in 15 sampled species of the family, spanning 60 million years of evolution [17], there

is a single locus with sequence similarity to both *AtTER1* and *AtTER2* [18]. Sequence alignment

of the recovered *AtTER1/2-like* loci from the 15 sampled species revealed changes at the

template domain in four and the complete absence of a template-like domain in three. In order

for the four loci with template mutations to encode for TER, a corresponding change in the

telomere repeat should be observed [18]. However, genomic data from several of these

species, including *Arabidopsis lyrata* and *Capsella rubella*, indicate the plant telomere repeat is

conserved [19], suggesting the presence of an alternative TER locus. Although TER is known to

evolve rapidly at the sequence level in other systems, the rapid rate of novel TER incorporation

suggested by the identification of *AtTER1* is unprecedented. Recently, Fajkus et al. (2019)

showed that *AtTER1* mutants do not exhibit telomere shortening, and thus does not act as a

bona fide telomerase RNA [20].

Here we generate double mutants at *AtTER1* and *AtTER2* and verify that *AtTER2* does

not play a compensatory role in telomere length maintenance in the absence of *AtTER1*. Due to

the previously documented rapid evolution at the *AtTER1/2-like* locus [18], we sought to

determine the evolutionary conservation of the telomerase complex using a series of

comparative evolutionary and biochemical analyses on the protein constituents of telomerase.

Interestingly, rather than concluding that TER and components of telomerase are evolving

rapidly as the evolutionary history of *AtTER1/2* suggested, we find that the protein components

of telomerase are highly conserved, as is the biochemical activity of telomerase. We performed

an RNA immunoprecipitation using core protein components of the telomerase holoenzyme,

followed by RNA-sequencing, and then used our evolutionary data to constrain possible

template-bearing TER candidates to those that were conserved across the Brassicaceae family.

This alternative approach led to the identification of the conserved plant TER discovered by

Fajkus et al. (2019). Our data indicate that telomerase core components are highly conserved

and our data complements the findings in Fajkus et al. (2019) in verifying that the biochemical

activity of the holoenzyme is also conserved. Finally, we explore the possibility that the

previously documented *AtTER-like* locus in Brassicaceae shares the *AtTER2* function of

modulating telomerase activity during genotoxic stress.

Results and Discussion

Double *AtTER1/AtTER2* mutants lack a telomere phenotype. Beilstein et al. (2012) showed

that the *TER1/2-like* locus was not conserved throughout the Brassicaceae family, or even in

other species of the genus *Arabidopsis*. This led us to re-examine the conservation of *AtTER1*

among the ecotypes of *A. thaliana*. Using genomic data from 1001genomes.org, we generated

an alignment of the *AtTER1* locus using MEGA5 and sorted through the template domain in

Geneious (Biomatters, Inc.) [21,22]. We found that *AtTER1* had a mutated template domain in

41/853 *A. thaliana* ecotypes (TCCCAAA**T** -> TCCCAAA**A**), which would preclude the use of an

AtTER1 transcript for telomere elongation in these ecotypes. Until the recently published work of

Fajkus et al. (2019), previous characterization of the *AtTER1* phenotype was performed using

an RNAi knock-down approach [14]; therefore, we generated a template deletion mutant at

AtTER1.

A telomerase RNA cannot perform its templating function without an intact template

domain, and thus we independently used CRISPR-Cas9 nuclease and CRISPR-Cas9-D10

nickase to create full template deletions at the *AtTER1* locus. We recovered two alleles that

lacked the template domain in its entirety (Fig 1A), *Atter1 Δ18* and *Atter1 Δ22*. First generation

homozygous mutants showed no reduction in telomere length as measured by telomere

restriction fragment length analysis (TRF) (Fig 1B). We then propagated both *ter1* homozygous

template nulls for several generations and measured telomere length in order to determine

whether there was progressive loss over time. We observed no reduction in telomere length for

either template-null mutant allele, regardless of generation (Fig 1B). Thus, our data definitively

indicate that *AtTER1* alone does not function as the template for telomere elongation,

confirming the findings in Fajkus et al. 2019.

Fig 1. Double knockout of *TER1* and *TER2* does not decrease telomere length. (A) Two

Atter1 mutants were created using either CRISPR-Cas9 nuclease or nickase. The complete

deletion of the template domain precludes the use of *AtTER1* as the template for telomere

extension. (B) TRF analyses were performed on individual plants with two biological replicates

in a population segregating the indicated *ter1* allele (Δ18 or Δ22), and in successive generations

(G4 and G5). (C) TRF analyses were performed on two sets of double mutant (*ter1 Δ22/ter2-2*

and *ter1 Δ18/ter2-3*) populations segregating the indicated mutations. A '+' indicates a

homozygous wild type individual for the indicated gene and a '-' represents a homozygous

mutant for the indicated gene.

To rule out the possibility that *AtTER2* acts in a compensatory role to elongate telomeres

in the absence of *AtTER1*, we crossed both *Atter1* template null alleles (*Atter1 Δ18* and *Atter1*

Δ22), with two *Atter2* T-DNA mutants (*Atter2-2* and *Atter2-3*), to create homozygous double

mutants and then measured telomere length by TRF. Neither single mutants for either gene, nor

double mutants showed a decrease in telomere length (Fig 1C). Taken together, the lack of

decrease in telomere length in both the single and double mutants indicates that *A. thaliana*

requires neither the *AtTER1* nor the *AtTER2* locus to elongate telomeres.

Telomerase RNP evolution is under purifying selection in Brassicaceae. Relatively high

levels of conservation of the telomerase RNA within major eukaryotic lineages has allowed for

the identification of distantly related TERs within those lineages. It was the unprecedented lack

of conservation of the template domain at *AtTER1* within *A. thaliana* and *AtTER1/2-like* locus

within other species of Brassicaceae that motivated further exploration of telomerase RNP

evolution and function. Numerous genome expansions and contractions within the Brassicaceae

family [23] have created opportunities for gene neofunctionalization or subfunctionalization,

potentially leading to incorporation of novel subunits into the telomerase RNP in Brassicaceae,

including the potential for alternative TERs [18]. If novel subunits have been incorporated into

the telomerase RNP, we might expect the core subunits to undergo structural or chemical

changes associated with the accommodation of novel binding partners. These changes would

be expected to arise under positive selection. Alternatively, if the telomerase RNP has not

undergone this type of change, we would expect the core subunits to evolve under purifying

selection. With this in mind, we sought to determine if major protein components of the

telomerase RNP exhibited signatures of positive selection, indicating underlying changes to the

complex, or purifying selection, indicating a stable RNP. We tested for positive selection in

TERT along several branches within the Brassicaceae. In total, we tested five branches

indicated by roman numerals in Figure 2 using the branch-sites test in PAML 4.4b [24]. In each

case we recovered no evidence that *TERT* evolved under elevated rates of non-synonymous to

synonymous substitutions (Fig 2), suggesting that *TERT* remains under strong purifying

selection.

Fig 2. Tests for signatures of positive selection in TERT evolution in Brassicaceae.

Branches throughout the Brassicaceae phylogeny, indicated by italicized roman numerals and

bolded branches, were tested for positive selection, Likelihood ratio tests were performed on the

bolded branches using PAML (codeml) to compare the likelihood scores of models of evolution

that either include site classes with ω values ≥ 1 (Alt = Alternative model) or explicitly exclude

these site classes (Null = Null model). Alternative model likelihood scores that are ≥ 1.92 better

than null model likelihood scores indicate a significant signal of positive selection ($p<0.05$) along

the specified branch (denoted as *i-vi* on the tree).

In addition to *TERT*, we also searched for evidence of positive selection in three other

genes that encode telomerase accessory factors in *A. thaliana*. Dyskerin is a pseudouridine

synthase necessary for maturation of rRNA that has also been shown to be an essential

component of telomerase in *A. thaliana* and many other eukaryotes [25]. KU is a heterodimer

composed of KU70 and KU80 that associates with telomerase in humans, budding yeast, and

A. thaliana via Ku80-telomerase interactions [15,26,27]. KU plays dual roles as a key factor in

double-strand DNA break repair via the non-homologous end-joining pathway, as well as

telomere maintenance [28,29]. With the exception of the branches leading to *C. rubella* (iii) and

S. irio/B. rapa (vi) for *DYSKERIN* and *A. lyrata KU70*, we found no evidence of positive selection

(S1 Figure). Thus, contrary to our initial expectations, the protein components of telomerase are

not responding at the molecular level to genome duplications or contractions, such as those that

occurred in species like *B. rapa*. These findings support the idea that a highly conserved TER

locus may be present and functional in Brassicaceae.

An evolutionary analysis of Brassicaceae telomerase enzymology recapitulates the

accepted organismal phylogeny. We next sought to take an indirect, but more fine-scale

evolutionary look at how conserved TERs are in Brassicaceae, specifically in the region within

and adjacent to the template domain. The ability of telomerase to bind and extend a substrate

comes from TERT-substrate and TER-substrate interactions [8]. In addition, telomere synthesis

requires Watson-Crick base-pairing between the 3' end of the DNA substrate and the beginning

of the template domain within TER [30]. Thus, changes within and surrounding the template

domain can lead to differential alignment on a telomere substrate or premature product

dissociation, particularly when these changes occur within the 3' end of the template region [31–

33]. With this rationale in mind, we set out to determine if the origin of the structurally similar

Brassicaceae TERs came from one evolutionary event (i.e., an alternative TER locus with

shared ancestry for all Brassicaceae) or multiple independent evolutionary events (i.e.,

alternative TER loci in each species that converged rapidly on a similar TER structure).

To distinguish between these two possibilities, telomerase substrate utilization was

performed in ten species across Brassicaceae (Fig 3; S2 Figure). To generate a profile of

substrate utilization, we used the telomere repeat amplification protocol (TRAP) on partially

purified plant extracts incubated with a suite of oligos of different lengths and 3' end

composition, ranging from one to three nucleotides capable of Watson-Crick base-pairing with a

TER template (Fig 3; S2 Figure) [34]. Radio-labeled products were separated on a

polyacrylamide sequencing gel and compared against an oligo expected to produce the shortest

telomere permutation based on previous observations (N_{15}-GGG; Fig 3) [35]. Observed

differences in size between products relative to this oligo were measured and recorded as a

value of 0-6 using *A. thaliana* as the baseline (Fig 3, S2A Figure). A substrate utilization profile

was then generated for nine other species within Brassicaceae, including several close relatives

to *A. thaliana* (three biological replicates per species; summarized in Fig 3; S2B Figure). This

profile was then compared against the phylogenetic tree reflecting the known relationships

among these species (Fig 3).

Fig 3. Substrate utilization by Brassicaceae telomerase RNPs closely resembles the

organismal phylogeny. Substrate utilization profile for telomerase RNPs across ten

Brassicaceae. Left, the accepted organismal phylogeny of species examined. Right, substrate

utilization profile for all tested species. Telomerase extracts were incubated with a suite of oligos

of varying length and nucleotide composition. The site of binding and number of nucleotides

added prior to the first full repeat dictate the size of products produced. The lengths of each

product were determined by comparing against the shortest permutation (N15-GGG). Observed

product differences relative to N15-GGG were calculated for all species and oligo combinations.

The intensity with which the numbered boxes are shaded corresponds to the degree to which

the results differ from A. thaliana.

The substrate utilization profiles we generated for Brassicaceae telomerase closely

recapitulated the evolutionary history of the family (Fig 3). The Arabidopsis clade, along with *C.*

rubella, all utilized the suite of oligos in the same manner (White boxes; Fig 3). Importantly, *A.*

arabicum, which represents the earliest diverging lineage in Brassicaceae [36], shares the

profile of *A. thaliana*, despite 55 million years of evolution. Thus, these data support a model for

shared ancestry of a TER locus in all Brassicaceae.

Distantly related Brassicaceae TERTs assemble into a functional telomerase RNP in *A.

thaliana.** To confirm our evolutionary analyses indicating that both the protein and RNA

components of the telomerase RNP are likely highly conserved across Brassicaceae, we

attempted to complement *A. thaliana tert* null lines with the genomic version of *TERT* from four

species in the group: *Capsella rubella, Cardamine hirsuta, Eutrema salsugineum*, and

Schrenkiella parvula [37]. The ability to complement *A. thaliana tert* null mutants with TERTs

from divergent species would indicate high conservation of telomerase across Brassicaceae. In

A. thaliana, plants can survive for approximately nine generations in the absence of TERT [38].

During this time, telomeres progressively shorten until they reach a length sufficient to elicit a

DNA damage response that prohibits further cell division. We determined the degree to which

the *Attert* null was complemented by each of the transgenes using a chromosome arm-specific

telomere length assay known as PETRA [39,40]. In the wild type *A. thaliana* (Col-0)

background, telomeres for all three arms tested were between the typical 2-4 kilobase size

range (Figs 4A and 4C) [39]. As expected, telomeres in the unselected sixth generation *Attert* -/-

mutants were between 0.5-2 kilobases in length (Figs 4A and 4C). The *AtTERT* transgene was

able to complement the mutant background and restore telomeres back to wild type range (Figs

4A and 4C; three independent biological replicates for each construct). Telomeres in these lines

were elongated further in the second generation, suggesting successful and complete

complementation with this construct (Figs 4A and 4C). We observed similar or slightly better

complementation with TERT from *S. parvula*, one of the most distantly related species in our

study (Figs 4B and 4C) [17]. Interestingly, despite observing transcription of the *E. salsugineum,*

C. hirsuta, and *C. rubella TERT* transgenes (S4 Figure), and telomerase activity in these

complementation lines, we did not observe telomere elongation with these constructs (S3

Figure). However, with the exception of the 1L chromosome arm in *C. rubella*, there was no

significant decrease in telomere length between the two generations we tested (S3B Figure),

suggesting some degree of partial complementation.

Fig 4. *S. parvula* TERT complements loss of function *Attert* -/- mutants. (*A*) PETRA results

of the unselected, untransformed (US) control, wild type *A. thaliana* (Col.), and *Attert* -/- mutants

transformed with the *AtTERT* construct. Chromosome arms are depicted by 1L, 3L, and 5L and

indicate the left arm of chromosomes 1, 3, and 5, respectively. T1 and T2 indicate the first and

second generation after transformation. At least three independent T1 transformants were

analyzed and then propagated to the second generation. (B) PETRA results of complementation

of *Attert* -/- mutants by *S. parvula* (Sp) TERT. (C) Quantification of PETRA results from (A) and

(B).

These data may represent an inability on the part of some of these *TERT*s to fully

reconstitute telomerase in *A. thaliana*. However, *C. rubella, C. hirsuta,* and *E. salsugineum* all

have short endogenous telomeres, ranging from 1-3 Kb [19]. *S. parvula*, in contrast, has

telomeres closer in length to *A. thaliana*. Thus, what appears to be partial complementation in *A.*

thaliana may reflect some feature in the *C. rubella, C. hirsuta*, and *E. salsugineum* TERT protein

that governs the production of shorter telomeres. The observed complementation in *A. thaliana*

by TERTs from other Brassicaceae supports the hypothesis of a common TER secondary

structure in all Brassicaceae.

Identification of an alternative telomerase RNA by immunoprecipitation of the telomerase

RNP followed by RNA-seq. CRISPR induced template deletion mutants of both previously

described TER loci in *A. thaliana* indicated that neither is a bona fide TER. We performed a

series of immunoprecipitation (IP) experiments designed to purify the telomerase RNP from

floral tissue, seedling tissue, and plant cell culture, all of which express *TERT* at a high level

[37]. We tracked telomerase activity from our IP and input fractions using TRAP, and extracted

RNA from IP fractions with verified telomerase activity. Following cDNA synthesis we pooled

and sequenced the libraries from IP experiments that used either an anti-TERT or anti-POT1A

antibody. Sequencing yielded 164,505,451 total paired-end reads across the 20 IP experiments.

The resulting reads were mapped to the *A. thaliana* genome (TAIR 10) and long non-coding

RNAs (lncRNAs) were identified using Evolinc (Fig 5A) [41,42]. Interestingly, no reads mapped

to the *AtTER1* locus. We winnowed candidate TER loci by sorting through Evolinc predicted

lncRNAs to find those capable of generating the telomeric repeat and those that were

conserved across Brassicaceae, in accordance with our evolutionary analyses (Figs 2-4)

suggesting a highly conserved TER. From the candidate set, we identified five highly-expressed

loci exhibiting conservation across the family (Fig 5B). The locus with the highest average TPM

among those conserved in at least three of our representative Brassicaceae is the same locus

that was identified as a telomerase RNA, named *TR*, which was shown to be conserved across

land plants [20]. This locus was also previously identified as a lncRNA associated with hypoxic

stress, termed *AtR8*. Hereafter we will refer to this locus as *AtTR/R8* [43].

Fig 5. Computational identification of a putative telomerase RNA. (A) Flowchart showing analysis pipeline for RNA-immunoprecipitation. (B) List of the five telomerase RNA candidates identified through the analysis pipeline and ranked by highest average transcripts per million (TPM). (C) TRF of *AtTR/R8* T-DNA insertion lines from wildtype through third generation (G3) homozygous insertion lines with each lane representing separate individuals.

To confirm the involvement of this lncRNA in telomere maintenance, we obtained T-DNA mutants of *AtTR/R8* and confirmed homozygosity. We grew these lines for three generations, measuring telomere length using a TRF with each generation (Fig 5C). We found that homozygous *Attr/r8* mutants showed discrete banding patterns similar to *Attert* mutants. This phenotype is a hallmark of telomerase deficiency [38] and indicates an important role for *AtTR/R8* in telomere maintenance.

The *AtTER-like* locus is transcribed in other Brassicaceae. The relatively high expression of *AtTER2* and its reported role in the negative regulation of telomerase in *A. thaliana* [15] raises the possibility that the transcript produced from the *AtTER1/2-like* locus may be acting as a telomerase interacting RNA (TIR) similar to *AtTER2*. To address whether the *AtTER1/2-like* loci recovered from other species of Brassicaceae could potentially encode a TIR, we tested whether or not we could detect transcripts consistent with a TIR, but independent of the transcription of the *RAD52-1A* mRNA. For our analyses we chose four species with varying phylogenetic distances from *A. thaliana*: *C. rubella* (diverged ~20 MYA), *C. hirsuta* (diverged ~35 MYA), *E. salsugineum* (diverged ~ 43 MYA), and *S. parvula* (diverged ~ 43 MYA) [17]. The *AtTER1/2-like* locus partially overlaps *RAD52-1A* in each of these species, therefore we performed RT-PCR on *RAD52-1A* in each species in order to map the intron-exon boundaries (Fig 6A). Following cloning and sequencing of *RAD52-1A* mRNA, we designed reverse primers that bind within the first intron of *RAD52-1A* and used it in combination with a forward primer in the predicted 5' UTR (Fig. 6A and 6B). We amplified and sequenced transcripts from RT-PCR

products generated using this forward primer with a reverse primer in the first intron of *RAD52-*

1A. Our results indicate that either a lncRNA is produced at the locus in the tested species or

splice variants for *RAD52-1A* exist. Thus, whether the amplified RNAs are entirely distinct from

the transcript produced from *RAD52-1A* and if they are TIRs performing a similar role as

AtTER2 remains an open question.

Fig 6. The *AtTER1/2-like* locus is transcribed in four other Brassicaceae species. (A)

Schematic of the *AtTER1/2-like* locus in the Brassicaceae species examined in this study. The

overlapping *RAD52-1A* locus is also shown. Known transcription start sites (or ones shown in

this study) are indicated by a solid arrow. Conserved but unverified TER transcription stop sites

are indicated by dashed vertical black lines. Exons are shown by dark grey boxes, while introns

are light grey. Primer binding sites for determining putative TER expression are shown as a

black arrow beneath each locus. Expected product length is shown. Exon/intron lengths and

positions are shown to scale. (B) Transcriptional analysis of the putative TER loci. RT-PCR was

performed to determine TER expression using primers designed in the 5' UTR and within the

second intron of *RAD52 1A* as shown in (A). RT-PCR was performed on *RAD52-1A* to

determine intron/exon boundaries.

Conclusions

Our results indicate that the original identification of the telomerase RNA in *A. thaliana* was

incorrect, in agreement with the findings of Fajkus et al. (2019). Moreover, our data indicate that

the true TER locus is highly conserved in the family, and that the telomerase holoenzyme

across the family shares both key protein components as well as biochemical activity.

Independently through RNA-seq of AtTERT and AtPOT1A pull downs, we identified *AtTR/R8*, a

highly conserved lncRNA within the Brassicaceae that has the potential to act as the canonical

TER. These results are further confirmed by recent work in Asparagaceae and a number of

other taxa [20]. Fajkus et al. (2019) indicated that the same locus is conserved in a variety of

species spanning land plant history, indicating a single evolutionary origin of plant TER. Our

work expands on the findings in Fajkus et al. (2019) by showing a direct interaction of this RNA

with the telomerase holoenzyme in multiple tissues and stages in development in *A. thaliana*.

Further, we showed that despite sequence changes to any potential telomerase RNAs, TERT

proteins from across the Brassicaceae family can at least partially rescue an *Attert* mutant. This

finding points to the evolutionary stability of the entire telomerase RNP complex. Finally,

whether *AtTER2* has a role as a telomerase regulatory RNA remains an open question. We

detected transcripts from the *AtTER1/2-like* locus in other Brassicaceae, but were unable to

verify if these represent splice variants of *RAD52A-1* or a distinct lncRNA product. One

important consideration is that the phenotypes observed for *AtTER2* under genomic stress may

be due to a partial protein product encoded from this locus in *A. thaliana*. Regardless, future

work on telomerase regulation is required to distinguish between these hypotheses.

Materials and Methods

Plant Material and Propagation. *A. thaliana* (Col-0) seed from Dr. Dorothy Shippen, Texas

A&M University; *C. rubella* from Dr. Steven Wright, University of Toronto; *C. hirsuta* seed from

Dr. Angela Hay, Max Planck Institute for Plant Breeding Research; *E. salsugineum* seed from

Dr. Karen Schumaker, University of Arizona; *S. parvula* seed from Drs. Maheshi Dassanayake

and Dong-Ha Oh, Louisiana State University were used in this research. Standard Arabidopsis

growth conditions were used for all species.

CRISPR-Cas9 Mutation Analyses. The suite of CRISPR plasmids from Schiml, et al. (2014)

were obtained from the Ohio State University Arabidopsis Biological Resources Center (ABRC)

and used for cloning of CRISPR-Cas9 and CRISPR-Cas9-D10A vectors, following the

procedure described in Schiml, et al. (2014) [44]. A protospacer (5'-

GGGTTTAGTTGTCGTCTGAT-3') overlapping the template domain of *AtTER1* was used in

conjunction with both Cas9 and Cas9-D10A; in addition, a second protospacer (5'-

TTGTCCGGCGACAGAAATGG-3') targeting 33 base pairs downstream of the template domain

was used with Cas9-D10A. T2's were screened using PCR of the *TER1* locus followed by

HindIII restriction digest for deletions of the entire template domain.

Atter2 mutants *ter2-2* (SALK_121147) and *ter2-3* (SALK_140126) were obtained from

the ABRC then crossed to the two *ter1* alleles, *ter1 Δ22* and *ter1 Δ18*, respectively. The progeny

resulting from the crosses were hemizygous for a *ter2* allele and heterozygous for a *ter1* allele.

By selfing these individuals we obtained a population segregating the mutations for both genes

allowing direct comparisons between full siblings in TRF analyses.

Nuclear Protein Isolation, TRAP, TRF and RT-PCR. Nuclear extracts were obtained from

~10g of seedling tissue as described elsewhere [14]. TRAP was performed as described

previously [25]. Briefly, 50ng of total protein, sourced from flower tissue, was added to a 25ul

final reaction volume containing ~3uC ^{32}P-dGTP, Gotaq hot start mastermix (Promega), and

400nM Forward TRAP primer. This mixture was placed at 37°C for 45 minutes, followed by the

addition of 400nM TRAP Reverse primer. For a list of primers see Supplemental Table 1. PCR

followed with an initial 95°C step for 3', followed by 95°C for 15", 60°C for 30", 72°C for 1' 30"

and 35 cycles. TRAP products were resolved on a 6% acrylamide gel (19:1) with 7M urea. For

RNA extraction, SDS extraction buffer (final concentration of: 2mM Tris pH 7.5, 200uM EDTA

pH 8.0, 0.05% SDS) was added to 1ml of fraction supplemented with enough 3M sodium

acetate (300mM final concentration). Following vortexing, this mixture was phenol/chloroform

extracted and ethanol precipitated according to standard protocols. Terminal restriction

fragment (TRF) length analyses were performed as described in Nelson et al. (2014). In brief,

genomic DNA was extracted and digested with the restriction enzyme Tru1I, followed by a

Southern blot using the telomeric repeat as a probe. RT-PCR was performed as described

above using primers listed in Supplemental Table 1.

Positive Selection Analyses. We obtained CDS sequences for *TERT*, *Dyskerin*, *Ku70*, and

Ku80 from 15 taxa (Brassicaceae or close relatives) from publicly available genomes using

CoGe, TAIR, Phytozome, and SALK [41,45,46]. *Branch-sites* likelihood ratio tests were

performed on the indicated branches of each gene tree using PAML (codeml) [24] to compare

the likelihood scores of models of evolution that either include site classes with ω values ≥ 1

(denoted Alt = Alternative model) or explicitly exclude these site classes (denoted Null = null

model). Alternative model Likelihood scores that are ≥ 1.92 better than null model Likelihood

scores indicate a significant signal of positive selection ($p < 0.05$) along the specified branch

(see critical values from chi-squared distribution table at one degree of freedom). For *Ku70* and

Dyskerin, *L. alabamica* was removed from our analysis because the inferred phylogeny was not

congruent with the accepted organismal phylogeny. The *C. hirsuta* branch was not tested for

Ku70/80 and *Dyskerin*, as genomic sequence was not available for these genes.

Substrate Specificity Analysis. Substrate specificity was tested by replacing the TRAP

Forward G primer with alternative forward primers, keeping the reverse primer the same. These

products were resolved on a 6% acrylamide sequencing gel. The space between two bands

generated from the N15-GGG oligo was divided into seven quadrants (using ImageJ, NIH).

Where bands in other lanes migrate relative to these quadrants determines the value given to

them for "observed product difference."

***tert -/-* Complementation.** Full length genomic *TERT* constructs were PCR amplified from each

species tested. These constructs were designed to include 2 Kb upstream and 0.5 Kb

downstream sequence to include appropriate regulatory regions. Constructs were cloned into

the promoter-less Gateway (Invitrogen) vector pH7WG using In-Fusion (Clontech) and then

sequence verified. *Agrobacterium tumefaciens* (strain GV3101) was used to transform this

binary vector into a fifth generation *tert* -/- *A. thaliana* background using the floral dip method

[47]. At least three independent transformed lines were identified for each construct. RT-PCR

was performed on each line to ensure transcription of the *TERT* transgene using primers

denoted in Supplemental Table 1. PETRA was performed as described in Heacock et al. (2004)

using subtelomeric primers specific for chromosome arms 1L, 3L, and 5L (see S1 Table for

primer sequence). Radioactive images were quantified using ImageJ (NIH).

RNA Immunoprecipitation (RIP) of the Telomerase Complex and RNA-seq Library

Preparation: 5.5-day old *A. thaliana* accession Columbia (Col-0) seedlings, 5-day old *A.*

thaliana T87 cell culture, and *A. thaliana* Col-0 flowers were used for IP experiments. Cell

culture was concentrated using Miracloth and the resulting powder was dried, packed, and

stored at -80 °C for IP. Samples were pulverized in liquid nitrogen and approximately 100 mg of

tissue powder was resuspended in 5 mL of cold RIP buffer (200 mM Tris-HCl pH 9, 110 mM

potassium acetate, 0.5 % Triton X-100, 0.1 % Tween 20, 2.5 mM dithiothreitol (DTT), 1.5 %

Protease Inhibitor Cocktail (Sigma-Aldrich P9599), 40 units/µL RNaseOUT (ThermoFisher

p10777019)) by pipetting. The 5 mL of homogenate was then centrifuged at 1,500 x g for 2

minutes at 4°C. From the resulting supernatant, 250 µL was kept as RIP input, and 5 µL was

kept for the TRAP assay.

Immunoprecipitation was performed by washing 150 µL of Protein A Dynabeads

(ThermoFisher, 10002D) twice with PBS (137 mM NaCl, 2.7 mM KCl, 10 mM sodium phosphate

dibasic, 2 mM sodium phosphate monobasic, pH 7.4) supplemented with 0.02% Tween 20.

Then, 5-10 µL of GFP (Abcam, ab290), TERT, or POT1A antibody was diluted to 600 µL with

PBS supplemented with 0.02 % *Tween* 20, 0.2 mg/mL Salmon Sperm DNA (ThermoFisher,

15632011), and 0.25 mg/mL BSA. Washed beads were resuspended in the antibody solution

and allowed to rotate at room temperature for 90 minutes.

After 90 minutes, the beads were separated from the antibody-PBS solution and

resuspended in 2 mL of the RIP extract. The bead-RIP mixture was rotated at 4°C for 3 hours.

After 3 hours, beads were washed 6 times with 1 mL cold RIP buffer and washed twice with 1

mL of cold TMG buffer (10 mM Tris-Acetate pH8, 1 mM $MgCl_2$, 1 mM DTT, 10 % glycerol).

Beads were resuspended in 150 μL of cold TMG buffer, 5 μL was kept for TRAP assay.

To purify the RNA associated with the telomerase RNA complex, we added 1 mL of

TRIzol Reagent (ThermoFisher, 15596026) to the 145 μL of remaining beads at the end of the

RIP experiment to extract RNA following manufacturer's instructions. RNA was treated with

TURBO DNase (ThermoFisher, AM2238) to degrade contaminating genomic DNA following the

manufacturer's instructions. RNA was purified from the DNase reaction by ethanol precipitation.

The RNA libraries were prepared using the YourSeq FT v1.5 kit (Amaryllis Nucleics) following

manufacturer's instructions. RNA samples checked for integrity using a Bioanalyzer throughout

preparation. RNA was sequenced using Illumina NextSeq 500 PE x 75bp.

Computational Analysis of RNA-seq Data. RNA purifications from POT1A and TERT

antibodies generated 164,505,451 total paired-end reads across 20 pull-down experiments.

Reads were mapped to the *A. thaliana* TAIR 10 reference genome using the Read Mapping and

Transcript Assembly workflow (RMTA v2.5.1.2) (github.com/Evolinc/RMTA) through the

CyVerse Discovery Environment (cyverse.org). RMTA utilizes Hisat2 [48] for read mapping, and

Stringtie [49] for transcript assembly as a seamless workflow. Default RMTA parameters were

used, besides specifying paired-end reads and changing the maximum intron length from

500,000 bp to 50,000. Read mapping for all experiments achieved at least 89% overall

alignment rates. Candidate loci were identified using Evolinc v1.7.5 [42]. Evolinc identifies and

annotates novel long intergenic noncoding RNAs (lincRNAs) using assembled transcripts from

RMTA as GTF files as well as information regarding surrounding annotated genes using a

reference genome annotation. Candidate lincRNAs from the 20 pull-down experiments were

merged with the reference genome annotation to a single annotation file using Evolinc merge for

subsequent analyses.

To quantify the expression of candidate loci, featureCounts v1.6.0 [50] was used

(specifying paired-end reads, feature types: exon, and gene attribute: gene_id) taking BAM files

from RMTA as input along with the merged annotation file. Gene expression was normalized

across the 20 pull-down experiments by converting counts to transcripts per million (TPM) which

accounts for feature length and sequencing depth.

To begin filtering for candidate telomerase RNAs, genes were only selected if they had

an average TPM (Transcripts per Kilobase Million) over 1 across the 20 experiments (78,487

features to 16,473 features). Candidates overlapping known genes, from TAIR 10, were then

discarded (16,473 features to 1,780 features). Candidate loci were then analyzed for putative

template domains by mapping all permutations of the telomere repeat "TTTAGGG" and its

reverse complement from 9 to 19 nt in length (154 possible permutations) to the 1,780

candidates using the software Geneious Prime 2019.2.1 (BioMatters, Inc.), resulting in 225

remaining candidate loci containing at least a 9 nt template domain.

Conservation analyses on the 225 candidate loci was performed using Evolinc-II which

identifies orthologous loci across a defined set of genomes, in this case from representative

species across the Brassicaceae family *(Arabidopsis thaliana, Arabidopsis lyrata, Capsella*

rubella, Brassica oleracea, Brassica rapa, Eutrema salsugineum, Schrenkiella parvula, and

Aethionema arabicum) as well as one species from the sister family, *Tarenaya hassleriana* in

Cleomaceae. Evolinc-II was used with an E-value cutoff of E-5 to allow for sequence divergent

orthologs to be identified. Finally, TER candidates were removed if a template containing

ortholog was not identified in at least 3 relatives, yielding 5 candidates.

Telomere Length of AtTR/R8 T-DNA Lines. The T-DNA line (FLAG-410H04) was obtained

from the Versailles INRA collection. Plants homozygous for the T-DNA were bred for three

generations to show the long term impact of a *AtTR/R8* loss on telomeres. The wildtype

comparison used for the TRF was a homozygous wildtype sibling obtained from a population

segregating for the FLAG-410H04 T-DNA insertion.

Acknowledgements

We thank Dr. Dorothy Shippen for the kind gift of anti-TERT and anti-POT1A antibodies, and

Jairui Song from the Shippen Lab for assistance with our immunoprecipitation experiments. In

addition, we thank the PaBeBaWoMo research group at the University of Arizona for insightful

and helpful comments during the completion of this project and members of the Beilstein lab for

help with plant cultivation, tissue collection, and moral support.

References

1.   Watson JM, Riha K. Comparative biology of telomeres: Where plants stand. FEBS Lett.

2010;584: 3752–3759. doi:10.1016/j.febslet.2010.06.017

2.   de Lange T. How Telomeres Solve the End-Protection Problem. Science (80-).

2009;326: 948–952. doi:10.1126/science.1170633

3.   Autexier C, Lue NF. The Structure and Function of Telomerase Reverse Transcriptase.

Annu Rev Biochem. 2006;75: 493–517. doi:10.1146/

4.   Egan ED, Collins K. Biogenesis of telomerase ribonucleoproteins. RNA. 2012;18: 1747–

1759. doi:10.1261/rna.034629.112

5.   Collins K. The biogenesis and regulation of telomerase holoenzymes. Nat Rev Mol Cell

Biol. 2006;7: 484–494. doi:10.1038/nrm1961

6.   Theimer CA, Feigon J. Structure and function of telomerase RNA. Curr Opin Struct Biol.

2006;16: 307–318. doi:10.1016/J.SBI.2006.05.005

7.   Gunisova S, Elboher E, Nosek J, Gorkovoy V, Brown Y, Lucier J-F, et al. Identification

and comparative analysis of telomerase RNAs from Candida species reveal conservation

of functional elements. RNA. 2009;15: 546–59. doi:10.1261/rna.1194009

8.   Blackburn EH, Collins K. Telomerase: an RNP enzyme synthesizes DNA. Cold Spring

Harb Perspect Biol. 2011;3. doi:10.1101/cshperspect.a003558

9.   Lingner J, Hendrick LL, Cech TR. Telomerase RNAs of different ciliates have a common

secondary structure and a permuted template. Genes Dev. 1994;8: 1984–1998.

doi:10.1101/gad.8.16.1984

10.   Chen J-L, Blasco MA, Greider CW. Secondary Structure of Vertebrate Telomerase RNA.

Cell. 2000;100: 503–514. doi:10.1016/S0092-8674(00)80687-X

11.   Tzfati Y. A novel pseudoknot element is essential for the action of a yeast telomerase.

Genes Dev. 2003;17: 1779–1788. doi:10.1101/gad.1099403

12.   Qi X, Li Y, Honda S, Hoffmann S, Marz M, Mosig A, et al. The common ancestral core of

vertebrate and fungal telomerase RNAs. Nucleic Acids Res. 2013;41: 450–462.

doi:10.1093/nar/gks980

13.   Dandjinou AT, Lévesque N, Larose S, Lucier J-F, Elela SA, Wellinger RJ. A

Phylogenetically Based Secondary Structure for the Yeast Telomerase RNA. Curr Biol.

2004;14: 1148–1158. doi:10.1016/j.cub.2004.05.054

14.   Cifuentes-Rojas C, Kannan K, Tseng L, Shippen DE. Two RNA subunits and POT1a are

components of Arabidopsis telomerase. Proc Natl Acad Sci U S A. 2011;108: 73–78.

doi:10.1073/pnas.1013021107

15.   Cifuentes-Rojas C, Nelson ADL, Boltz KA, Kannan K, She X, Shippen DE. An alternative

telomerase RNA in Arabidopsis modulates enzyme activity in response to DNA damage.

Genes Dev. 2012;26: 2512–23. doi:10.1101/gad.202960.112

16.   Xu H, Nelson ADL, Shippen DE. A Transposable Element within the Non-canonical

Telomerase RNA of Arabidopsis thaliana Modulates Telomerase in Response to DNA

Damage. Zhou J-Q, editor. PLOS Genet. 2015;11: e1005281.

doi:10.1371/journal.pgen.1005281

17. Beilstein MA, Nagalingum NS, Clements MD, Manchester SR, Mathews S. Dated

molecular phylogenies indicate a Miocene origin for Arabidopsis thaliana. Proc Natl Acad

Sci U S A. 2010;107: 18724–8. doi:10.1073/pnas.0909766107

18. Beilstein MA, Brinegar AE, Shippen DE. Evolution of the Arabidopsis telomerase RNA.

Front Genet. 2012;3: 1–8. doi:10.3389/fgene.2012.00188

19. Nelson ADL, Forsythe ES, Gan X, Tsiantis M, Beilstein MA. Extending the model of

Arabidopsis telomere length and composition across Brassicaceae. Chromosom Res.

2014;22: 153–166. doi:10.1007/s10577-014-9423-y

20. Fajkus P, Peška V, Závodník M, Fojtová M, Fulnečková J, Dobias Š, et al. Telomerase

RNAs in land plants. Nucleic Acids Res. 2019; doi:10.1093/nar/gkz695

21. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et al.

1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell.

2016;166: 481–491. doi:10.1016/j.cell.2016.05.063

22. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular

Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and

Maximum Parsimony Methods. Mol Biol Evol. 2011;28: 2731–2739.

doi:10.1093/molbev/msr121

23. Bowers JE, Chapman BA, Rong J, Paterson AH. Unravelling angiosperm genome

evolution by phylogenetic analysis of chromosomal duplication events. Nature. 2003;422:

433–438. doi:10.1038/nature01521

24. Bielawski J, Yang Z. A Maximum Likelihood Method for Detecting Functional Divergence

at Individual Codon Sites, with Application to Gene Family Evolution. J Mol Evol. 2004;59:

121–132. doi:10.1007/s00239-004-2597-8

25. Kannan K, Nelson ADL, Shippen DE. Dyskerin Is a Component of the Arabidopsis

Telomerase RNP Required for Telomere Maintenance. Mol Cell Biol. 2008;28: 2332–

2341. doi:10.1128/MCB.01490-07

26. Stellwagen AE. Ku interacts with telomerase RNA to promote telomere addition at native

and broken chromosome ends. Genes Dev. 2003;17: 2384–2395.

doi:10.1101/gad.1125903

27. Ting NSY. Human Ku70/80 interacts directly with hTR, the RNA component of human

telomerase. Nucleic Acids Res. 2005;33: 2090–2098. doi:10.1093/nar/gki342

28. Jackson SP. Sensing and repairing DNA double-strand breaks. Carcinogenesis. 2002;23:

687–696. doi:10.1093/carcin/23.5.687

29. Bertuch AA, Lundblad V. The Ku Heterodimer Performs Separable Activities at Double-

Strand Breaks and Chromosome Termini. Mol Cell Biol. 2003;23: 8202–8215.

doi:10.1128/MCB.23.22.8202-8215.2003

30. Qi X, Xie M, Brown AF, Bley CJ, Podlevsky JD, Chen JJ-L. RNA/DNA hybrid binding

affinity determines telomerase template-translocation efficiency. EMBO J. 2012;31: 150–

161. doi:10.1038/emboj.2011.363

31. Gilley D, Blackburn EH. Specific RNA residue interactions required for enzymatic

functions of Tetrahymena telomerase. Mol Cell Biol. 1996;16: 66–75.

doi:10.1128/MCB.16.1.66

32. Wang H. A novel specificity for the primer-template pairing requirement in Tetrahymena

telomerase. EMBO J. 1998;17: 1152–1160. doi:10.1093/emboj/17.4.1152

33. Förstemann K, Lingner J. Telomerase limits the extent of base pairing between template

RNA and telomeric DNA. EMBO Rep. 2005;6: 361–366. doi:10.1038/sj.embor.7400374

34. Fitzgerald MS, McKnight TD, Shippen DE. Characterization and developmental patterns

of telomerase expression in plants. Proc Natl Acad Sci U S A. 1996;93: 14422–7.

doi:10.1073/pnas.93.25.14422

35. Fitzgerald MS, Shakirov E V., Hood EE, McKnight TD, Shippen DE. Different modes of

de novo telomere formation by plant telomerases. Plant J. 2001;26: 77–87.

doi:10.1046/j.1365-313x.2001.01010.x

36. Beilstein MA, Al-Shehbaz IA, Kellogg EA. Brassicaceae phylogeny and trichome

evolution. Am J Bot. 2006;93: 607–619. doi:10.3732/ajb.93.4.607

37. Fitzgerald MS, Riha K, Gao F, Ren S, McKnight TD, Shippen DE. Disruption of the

telomerase catalytic subunit gene from Arabidopsis inactivates telomerase and leads to a

slow loss of telomeric DNA. Proc Natl Acad Sci U S A. 1999;96: 14813–8.

doi:10.1073/pnas.96.26.14813

38. Riha K, McKnight TD, Griffing LR, Shippen DE. Living with genome instability: Plant

responses to telomere dysfunction. Science (80-). 2001;291: 1797–1800.

doi:10.1126/science.1057110

39. Heacock M, Spangler E, Riha K, Puizina J, Shippen DE. Molecular analysis of telomere

fusions in Arabidopsis: multiple pathways for chromosome end-joining. EMBO J. 2004;23:

2304–2313. doi:10.1038/sj.emboj.7600236

40. Heacock ML, Idol RA, Friesner JD, Britt AB, Shippen DE. Telomere dynamics and fusion

of critically shortened telomeres in plants lacking DNA ligase IV. Nucleic Acids Res.

2007;35: 6490–6500. doi:10.1093/nar/gkm472

41. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, et al. The arabidopsis

information resource: Making and mining the "gold standard" annotated reference plant

genome. genesis. 2015;53: 474–485. doi:10.1002/dvg.22877

42. Nelson ADL, Devisetty UK, Palos K, Haug-Baltzell AK, Lyons E, Beilstein MA. Evolinc: A

Tool for the Identification and Evolutionary Comparison of Long Intergenic Non-coding

RNAs. Front Genet. 2017;8: 52. doi:10.3389/fgene.2017.00052

43. Wu J, Okada T, Fukushima T, Tsudzuki T, Sugiura M, Yukawa Y. A novel hypoxic stress-

responsive long non-coding RNA transcribed by RNA polymerase III in Arabidopsis. RNA

Biol. 2012;9: 302–313. doi:10.4161/rna.19101

44.   Schiml S, Fauser F, Puchta H. The CRISPR/Cas system can be used as nuclease for in

planta gene targeting and as paired nickases for directed mutagenesis in Arabidopsis

resulting in heritable progeny. Plant J. 2014;80: 1139–1150. doi:10.1111/tpj.12704

45.   Lyons E, Pedersen B, Kane J, Alam M, Ming R, Tang H, et al. Finding and Comparing

Syntenic Regions among Arabidopsis and the Outgroups Papaya, Poplar, and Grape:

CoGe with Rosids. Plant Physiol. 2008;148: 1772–1781. doi:10.1104/pp.108.124867

46.   Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a

comparative platform for green plant genomics. Nucleic Acids Res. 2012;40: D1178–

D1186. doi:10.1093/nar/gkr944

47.   Zhang X, Henriques R, Lin S-S, Niu Q-W, Chua N-H. Agrobacterium-mediated

transformation of Arabidopsis thaliana using the floral dip method. Nat Protoc. 2006;1:

641–646. doi:10.1038/nprot.2006.97

48.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. Nat Methods. 2015;12: 357–360. doi:10.1038/nmeth.3317

49.   Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. StringTie

enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol.

2015;33: 290–295. doi:10.1038/nbt.3122

50.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–930.

doi:10.1093/bioinformatics/btt656

Figure Legends

Supplemental Fig 1. Positive selection analysis of Dyskerin, Ku70, and Ku80 in

Brassicaceae. Roman numerals correspond to branches tested in Fig 2. Alternative model

likelihood scores that are significantly greater than null model likelihood scores indicate positive

selection (p<0.05) along the specified branch and are denoted in bold.

Supplemental Fig 2. Substrate utilization of telomerase from different Brassicaceae

telomerase RNPs. (*A*). Expected binding between AtTER1 template and each oligo used in the

experiment. The first set of nucleotides added by this template before the first full repeat is

shown in bold. Expected product differences are calculated relative to oligo #4, which is the

shortest oligo provided in the assay and therefore serves as the baseline. (*B*) A subset of the

gel image from the substrate utilization assay for each species tested. A dashed line is drawn

through the center of a band for oligo N15-GGG, which serves as the baseline for calculating

observed product differenced for the other oligos.

Supplemental Fig 3. Test of complementation of *Attert* -/- background with genomic TERT

constructs from other Brassicaceae. (*A*) PETRA results for Attert -/- lines transformed with *C.*

rubella TERT (Cr), *C. hirsuta* TERT (Ch), and *E. salsugineum* TERT (Es). (*B*) Quantification of

results shown in (*A*).

Supplemental Fig 4. Confirmation of expression of TERT transgenes for

complementation experiments. RNA was extracted from floral tissue from each background.

RT-PCR was performed on both the 3' end of the TERT transgene (shown) and the full-length

construct (not shown). GAPDH was used to determine quality of RNA and as an approximate

loading control. "+" indicates a positive RT reaction, whereas "-" indicates no reverse

transcriptase was added.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

4 Flower samples
4 TERT IPs

12 Cell culture Samples
[5 days old]
8 POT1A IPs & 4 TERT IPs

4 Seedling samples
[5.5 days old]
4 TERT IPs

24,431,607
paired-end
reads

76,210,945
paired-end
reads

55,676,420
paired-end
reads

8,186,479
paired-end
reads

Map, assemble, and
annotate novel transcripts
[RMTA, Evolinc-1]

Count and normalize reads to TPM
[featureCounts]

78,487 features total

Keep anything with TPM > 1

16,473 features total

Remove known genes

1,780 features total

Find transcripts with template
repeat 9-19 nt in length

225 transcripts total

Remove transcripts with < 3 template
containing orthologs in selected
Brassicaceae species
[Evolinc-2]

5 candidates

B

Rank	No. of Reads	Average TPM	Locus Length	Template (length)	Chromosome: Position	Longest ORF
#1	4,572	347.05	296 bp	CTAAACCCT 9 nt	2: 12619075-12619370	48 bp
#2	2,276	75.77	846 bp	CCTAAACCCT 10 nt	2: 3496512-3497357	162 bp
#3	389	34.98	242 bp	AAACCCTAAACCC 13 nt	3: 380473-380714	78 bp
#4	349	17.47	541 bp	AAACCCTAA 9 nt	4: 16020834-16021374	141 bp
#5	158	14.80	270 bp	AAACCCTAA 9 nt	2: 8742377-8742646	45 bp

C

AtTR/R8 T-DNA

Wild Type   G1 G2   G3

5000 bp
4000 bp
3000 bp
2000 bp
1500 bp

Figure 6