Rest functional brain maturation during the first year of life

Hervé Lemaître[a,b†*] and Pierre Augé[a†], Ana Saitovitch[a], Jean-Marc Tacchella[a], Ludovic Fillon[a],

Raphael Calmon[a], Raphaël Lévy[a], David Grévent[a], Francis Brunelle[a], Nathalie Boddaert[a],

Monica Zilbovicius[a]

Author affiliation

"a" INSERM UA10, Department of Pediatric Radiology, Hôpital Necker Enfants Malades, AP-

HP, Imagine Institute (UMR 1163), Paris Descartes University, Sorbonne Paris Cité University.

"b" Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives (CNRS

UMR 5293), Université de Bordeaux.

† Both authors contributed equally for this work

*Correspondence and material request should be addressed to Hervé Lemaître

(herve.lemaitre@u-bordeaux.fr)

Abstract

The first year of life is a key period of brain development, characterized by dramatic structural and functional modifications. Here, we measured rest cerebral blood flow (CBF) modifications throughout babies' first year of life using arterial spin labeling magnetic resonance imaging sequence in 52 infants, from 3 to 12 months of age. Overall, global rest CBF significantly increased during this age span. In addition, we found marked regional differences in local functional brain maturation. While primary sensorimotor cortices and insula showed early maturation, temporal and prefrontal region presented great rest CBF increase across the first year of life. Moreover, we highlighted a late and remarkably synchronous maturation of the prefrontal and posterior superior temporal cortices. These different patterns of regional cortical rest CBF modifications reflect a timetable of local functional brain maturation and are consistent with baby's cognitive development within the first year of life.

Keywords:

Infants brain maturation, Rest cerebral blood flow, Neurodevelopment, ASL-MRI

Introduction

The human brain is still immature at birth and undergoes dynamic structural and functional processes throughout life. During the first year, the maturation of neural networks is a complex process that is particularly important to the baby's acquisition of cognitive and motor skills (Kagan and Herschkowitz 2005). At the cortical level, development comprises both gross morphometric changes and microstructural progression (Dubois et al. 2014). The first year of life is therefore a critical phase of postnatal brain development.

Historically, much of what we know about the intricate processes of early brain development comes from post-mortem studies in human fetuses, neonates, and non-human primates (Goldman-Rakic 1987; Kostovic et al. 2002; Innocenti and Price 2005). With the increasing availability of high-quality neuroimaging techniques, studying early human brain development in vivo in unprecedented detail is now feasible (Partridge et al. 2004; Fransson et al. 2007; Ball et al. 2014; van den Heuvel et al. 2015). These advances have led to exciting new insights into both healthy and atypical macroscale brain network development and have paved the way to bridge the gap between the brain's neurobiological architecture and its behavioral repertoire.

At the structural level, in neonates and infants, studies of cortical morphological development have focused on the modification of gray matter volume (Knickmeyer et al. 2008; Gilmore et al. 2012), gyrification (Li et al. 2014), deep sulcal landmark maturation (Meng et al. 2014), thickness and surface area maturation (Lyall et al. 2015), as well as folding and fiber density (Nie et al. 2014). Structural brain imaging studies showed an increase in the gray matter volume during the first years of life (Knickmeyer et al. 2008), consistent with post-mortem studies, indicating rapid development of synapses and spines during this period (Huttenlocher and Dabholkar 1997; Petanjek et al. 2011; Webster et al. 2011). Indeed, throughout late gestation, rapid synaptogenesis results in an over-abundance of synapses (up to 150% of adult values) that are subsequently pruned throughout childhood and adolescence (Huttenlocher 1979). During the first year of life, synaptogenesis is one of the most important maturational processes, and its timetable differs across cortical regions. Gilmore et al. described a posterior to anterior gradient of gray matter growth throughout the first year of life (Gilmore et al. 2007), consistent with regional differences that have been

described in post-mortem studies, showing an increase in synaptic density, and therefore

synaptogenesis, earlier in the sensory cortex and later in the prefrontal cortex (Huttenlocher

and Dabholkar 1997). In general, studies have suggested a complex pattern of development

that varies based on anatomical location and cortical metrics. In addition, across early

development, cortical maturation exhibits regionally specific asymmetry between the left

and right hemispheres (Li et al. 2014; Nie et al. 2014). These changes continue throughout

childhood and adolescence, with cortical thickness following different trajectories of

thinning depending on the region, cortex type and gender (Sowell et al. 2004; Shaw et al.

2008).

At the functional level, early brain development has been investigated using mainly three

different approaches. Pioneer studies measuring metabolism and rest cerebral blood flow

(CBF) were followed by activation studies using functional MRI and more recently resting

state MRI studies investigating functional connectivity.

Rest cerebral metabolism and blood flow are an index of synaptic density, which allows the

in vivo study of functional brain maturation using positron emission tomography (PET) and

single-photon emission computed tomography (SPECT) (Leenders et al. 1990). These studies

revealed that infants' brains showed higher rest metabolism in subcortical structures and in

the sensorimotor cortex than in other regions (Chugani and Phelps 1986). In the newborn,

the highest degree of glucose metabolism is in the primary sensory and motor cortex,

cingulate cortex, thalamus, brain stem, cerebellar vermis and hippocampal region. During

the first months of life, rest metabolism and CBF increase firstly within the primary sensory

cortices, followed by the associative sensory cortices and finally within the prefrontal cortex

at the end of the first year (Chugani and Phelps 1986; Chugani et al. 1987; Chiron et al.

1992). At 2 to 3 months of age, glucose utilization increases in the parietal, the temporal and

the primary visual cortices, basal ganglia, and cerebellar hemispheres. Between 6 and 12

months of age, glucose utilization increases in the frontal cortex. These metabolic changes

correspond to the emergence of motor and cognitive abilities during the first year of life.

However, these studies were limited by very low spatial resolution of the brain imaging

devices. In addition, these techniques required administration of ionizing radiation and,

therefore, have limited application in the pediatric population.

Following these pioneer studies, functional MRI studies have used blood oxygen level-
dependent (BOLD) contrast to measure brain activity. Task-based fMRI contributed to
present-day knowledge about brain maturation shortly after birth (Dehaene-Lambertz et al.
2006; Arichi et al. 2010; Allievi et al. 2016). These studies have provided important
background on the brain's responses to sensory input during the early developmental phases
of brain-behavior interactions. Adult-like activation patterns were observed in response to a
variety of sensory stimuli, including tactile and proprioceptive stimulation (passive hand
movement) (Erberich et al. 2006; Arichi et al. 2010) as well as auditory (Anderson et al. 2001)
and olfactory (the odor of infant formula) (Arichi et al. 2013). Functional MRI studies in 2- to
3-month-old infants demonstrated left-lateralized activation of perisylvian regions, including
the superior temporal gyrus, angular gyrus and Broca's area, in response to native language
speech. The response followed a hierarchical pattern, with auditory regions being activated
first, followed by superior temporal regions, the temporal poles and Broca's area in the
inferior frontal cortex; a pattern that is highly consistent with language organization in the
mature brain (Dehaene-Lambertz et al. 2006).

More recently, BOLD signal has been used to conduct resting state fMRI (rs-fMRI) as a
measure of temporal coherence between brain regions. This technique has provided, by
studying infants during the first years of life, insight into the maturation of multiple resting
state networks (RSNs). Results show that the rate at which correlations within and between
RSNs develop differs by network and closely reflect known rates of cortical development
based on histological evidence (Smyser and Neil 2015). The sensorimotor (SM) and attention
(AN) networks seem to be the earliest developing networks with their within-network
synchronization largely established before birth. This replicates several reports showing the
bilateral symmetric, adult-like topology of both networks at birth (Gao et al. 2013; Lin et al.
2013) or even prenatally (Smyser et al. 2010), indicating significant prenatal development of
these 2 networks. In the brains of term babies, rs-fMRI studies employing seed-based
connectivity or independent component analyses have identified specific functional
networks, including primary visual, auditory, sensorimotor networks and default mode and
executive-control networks involved in heteromodal functions (Fransson et al. 2007; Doria et
al. 2010; Smyser et al. 2010). Network analyses based on graph theory further revealed that
the functional connectomes of infant brains already exhibited the small-world structure.

Distinct from the adults, however, the hubs were largely confined to primary sensorimotor

regions (Fransson et al. 2011; Gao et al. 2011). Taken together, these findings provide

important insights into the early brain functional maturation process.

The emergence of arterial spin labeling (ASL), a technique that provides both non-invasive

and regional cerebral blood flow quantification, offers new opportunities to investigate local

rest brain function in neonates and children. ASL perfusion MRI uses magnetically labeled

arterial blood water as a nominally diffusible flow tracer. By labeling the blood water

proximal to the target imaging region, the perfusion signal is subsequently calculated by

comparison with a separate image acquired using a control pulse without labeling the blood

flow to remove the static background signal and control for magnetization transfer effects

(Williams et al. 1992). Therefore, ASL MRI non-invasively assesses brain perfusion and allows

for a quantitative measurement of rest CBF without the administration of contrast material

or exposure to ionizing radiation (Detre and Alsop 1999). Compared with BOLD, ASL has

several benefits by providing (1) an absolute quantitative measure of brain function through

CBF signaling, (2) not a derived measure from the BOLD signal affected by numerous

physiological and noise contributions, and (3) an increased spatial specificity to neuronal

activity due to the capillary/tissue origin of the ASL signal (Detre et al. 2009; Chen et al.

2015). Then, ASL MRI has been used to measure rest CBF to better characterize brain

maturation in children after the first year of life (Biagi et al. 2007; Paniukov et al. 2020).

Indeed, in a very recent study, Paniukov et al. (Paniukov et al. 2020) followed longitudinally

children between 2 and 7 years and showed a constant CBF increase across different regions

of the prefrontal, temporal, parietal and occipital cortex. However, knowledge about the

first year of life remains rather scarce. Wang et al. (Wang et al. 2008) by comparing 7- and

13-month old infants showed a regional CBF increase in the hippocampi, anterior cingulate,

amygdala, occipital lobe and auditory cortex. Duncan et al. (Duncan et al. 2014) studied

infants from 3 to 5 months and described a significantly greater rest CBF in the orbitofrontal,

subgenual and inferior occipital regions.

In this study, in order to characterize CBF developmental trajectories, we have measured

age-related changes of local rest CBF at the voxel level and regionally throughout the first

year of life using ASL perfusion MRI. We hypothesized that this crucial age range is

characterized by different regional patterns of brain development, mainly between primary

and associative regions, that in turn reflect cognitive development of the baby during the
first year.

Materials and Methods

Subject. Eighty-five babies from the Necker-Enfants-Malades hospital were initially included
in this study. The inclusion criteria were normal clinical multimodal MRI, absence of
prematurity, neurological or cranial pathology, parent's consanguinity or abnormal
psychomotor development. Were included infants presenting syndromes that are not
originally neurological, mainly dermatological or ophthalmological, but request an MRI to
discard infrequent associated brain abnormalities, that may be present in a small percentage
of cases (see SI Appendix, Table S1). Normal psychomotor development was assured in
follow-up consultations. Our final sample included 52 babies (29 girls) from 3 to 12 months
of age in our study, including 10 babies at 3 and 4 months (90 to 120 days), 14 at 5 and 6
months (120 to 180 days), 14 at 7 and 8 months (180 to 240 days), 7 at 9 and 10 months
(240 to 300 days), 7 at 11 and 12 months (300 to 375 days). The Ethical Committee of French
Public Hospitals approved this study and the written informed consent was obtained for all
participants.

MRI acquisition. All MRI exams included whole brain T1-weighted and ASL sequences and
were acquired on a General Electric Signa 1.5T MRI scanner in the Necker-Enfants-Malades
hospital (See SI Appendix, Table S2 and SI Methods for details). Due to the age of the babies,
all of them received premedication before their MRI (pentobarbital, 7.5 mg/kg) to prevent
motion artifacts. It has been shown that barbiturates do not have any influence on the
regional distribution of CBF or on default mode resting state network (Werner 1995;
Zilbovicius et al. 2000; Mishra 2002; Fransson et al. 2007; Doria et al. 2010).

Data processing and treatment. MRI images were pre-processed using Statistical Parametric
Mapping (SPM8 software, Welcome Department of Cognitive Neurology London
www.fil.ion.ucl.ac.uk/spm/software/spm8) implemented in Matlab (Mathworks Inc.,
Sherborn, MA, USA) and analyzed using a voxel-based approach (See SI Appendix, SI
Methods for details). Native 3D-T1-weighted images were segmented into gray matter,
white matter and cerebrospinal fluid using the Infant Brain Probability Templates
(https://irc.cchmc.org/software/infant.php). The unified segmentation enables spatial

normalization, tissue segmentation and bias correction within the same generative model

(Ashburner and Friston 2005). A postprocessing visual quality control was performed by two

independent investigators (PA and JMT) on the grey matter maps to ensure the quality of

the segmentation. The ASL images were co-registered to the corresponding native gray

matter images using a normalized mutual information cost function (separation: 4 and 2

mm, tolerance: from 0.02 to 0.001, histogram smoothing: 7 x 7 mm). After visual inspection

of the co-registration, the ASL images were then spatially normalized using the deformation

matrices from the segmentation process. The resulting ASL images were smoothed using an

isotropic Gaussian filter of 10 mm. ASL acquisition provides a high-quality image of

quantitative CBF. Motion in ASL acquisition is mainly characterized by signal outside of the

brain, often recognizable as signal from layers of skin or fat, that can be detected by on-the-

fly expert visual analysis. Therefore, we performed a two steps quality control. The first one

by an expert radiologist right after acquisition (NB) and the second one by an imaging

processing expert engineer (HL) before pre-processing to discard images with artifacts such

as motion, aliasing, ghosting, spikes, low signal to noise ratio.

Image analysis. We normalized rest CBF within the ASL images by the mean CBF measured

within the basal ganglia to avoid major variations in rest CBF due to cardiac blood flow (Licht

et al. 2004; Varela et al. 2012) and blood pressure labilities (Hardy et al. 1997). The basal

ganglia was specifically chosen in our study as it is one of the earliest structures to matures

(Chugani and Phelps 1986) and regression analyses did not show any age-related variations

in the rest CBF of this region within our age range (beta= 5.2 x 10^{-4} unit/day, $t_{(50)}$ = 0.045, p =

0.96). The regional rest CBF was expressed as percentage of basal ganglia rest CBF and

presented in arbitrary unit.

We then performed whole-brain voxel-wise analyses of the 52 images within the general

linear model framework using SPM8. Age was entered as covariate in a multiple regression

model. The analyses were constrained to gray matter tissue only by thresholding the analysis

mask to 40% of the mean gray matter image of our sample.

We also extracted mean rest CBF from 92 regions of interest (hemispheres and regions)

using the AAL parcellation toolbox (Tzourio-Mazoyer et al. 2002). We matched the AAL

parcellation to our sample by spatially normalizing the MNI single subject MRI brain to the

Infant Brain Probability Templates. In addition, a further analysis was performed by selecting

and merging regions of interested based on their relevance in term of development. We

selected the hippocampus, the amygdala, the thalamus, the primary visual and auditory

cortices, the insula, the superior temporal cortex. We formed the sensorimotor cortex by

merging the precentral and postcentral regions, and the prefrontal cortex by merging the

inferior, middle and superior frontal regions and the gyrus rectus. All analyses were

performed using R version 3.6.1 (http://cran.r-project.org) and ggplot2 (3.2.1) and lme4 (1.1-

21) packages. Age-related regressions were assessed using linear mixed models to account

for the intra-subject left and right hemisphere measurements. Age, hemisphere and age-by-

hemisphere interaction were entered as fixed effects and subject as nested random effect.

We corrected for multiple comparisons using Bonferroni correction by multiplying the p

values by the number of regions. Finally, we performed inter-regional correlation analyses of

the rest CBF values to derive a Pearson's correlation coefficient for each pair of regions and

to build an inter-regional correlation matrix. The ordering of the correlation matrix was

based on correlations with the first principal component of the same matrix.

Results

The relative values of global rest CBF increased with age from 3 to 12 months in the right (b

= 0.0010 unit/day, $t_{(55.71)}$ = 6.64, p = 1.36E-08) and in the left hemisphere (b = 0.00078

unit/day, $t_{(55.71)}$ = 5.34, p = 1.74E-06) with a greater age-related increase in the right as

compared to the left (p = 0,0074, see Figure 1 and Table 1).

<Table 1><Figure 1>

Qualitative analysis of the whole-brain voxel-wise maps showed a regionally heterogeneous

age-related increase of the relative rest CBF values (see Figure 2 and SI Appendix, Movies S1

to S4). The highest rest CBF at 3 months were observed within the sensorimotor and the

primary visual cortices. The age-related increase in rest CBF progressed spatially from these

regions. From the calcarine fissure, the rest CBF increased toward the visual associative

regions up to the supramarginal and the precuneus regions. From the primary motor and

sensory cortices, the rest CBF increased toward both the anterior and the posterior part of

the brain. Anteriorly, through the anterior cingulate and the prefrontal cortices; posteriorly,

through the insula and the superior temporal cortices. In contrast, the rest CBF was stable

within the thalamus, the amygdala and the hippocampus. Between 9 and 12 months, the

rest CBF increase was predominantly seen in the temporal and the prefrontal cortices. The

regional right over left rest CBF asymmetry remained present throughout the whole studied

period.

<Figure 2>

Quantitative analysis within the predefined regions of interest showed different trajectories

of local rest functional maturation (see Table 1 and Figure 1). First, in a subset of subcortical

regions including the hippocampus (right: b = 0.00026 unit/day, $t_{(69.66)}$ = 1.76, p = 0.74; left: b

= -0.00015 unit/day, $t_{(69.66)}$ = -1.01, p = 1), the amygdala (right: b = -0.00008 unit/day, $t_{(83.74)}$ =

-0.60, p = 1; left: b = -0.00019 unit/day, $t_{(83.74)}$ = -1.36, p = 1) and the thalamus (right: b = -

0.00030 unit/day, $t_{(65.29)}$ = -2.52, p = 0,13; left: b = -0.00044 unit/day, $t_{(65.29)}$ = -3.62, p =

0.0051), the age-related rest CBF maturation through the first year of life remained stable

indicating already matured regions at 3 months old. Second, the subset of cortical regions

including the primary visual (right: b = 0.00094 unit/day, $t_{(60.07)}$ = 4.33, p = 5.2E-04; left: b =

0.00028 unit/day, $t_{(60.07)}$ = 3.91, p = 0.0021) and primary auditory cortices (right: b = 0.00030

unit/day, $t_{(73.77)}$ = 1.60, p = 0.94; left: b = 0.00028 unit/day, $t_{(73.77)}$ = 1.48, p = 1), the insula

(right: b = 0.00043 unit/day, $t_{(84.2)}$ = 3.37, p = 0.010, left: b = 0.00024 unit/day, $t_{(84.2)}$ = 1.86, p

= 0.59) and the sensorimotor cortex (right: b = 0.00052 unit/day, $t_{(59.12)}$ = 2.50, p = 0.14; left:

b = 0.00025 unit/day, $t_{(59.12)}$ = 1.18, p = 1) presented a small age-related rest CBF increase

indicating early maturational process. Third, a subset of cortical regions including the

prefrontal (right: b = 0.00132 unit/day, $t_{(60.12)}$ = 6.80, p = 4.86E-08; left: b = 0.00118 unit/day,

$t_{(60.12)}$ = 6.10, p = 7.36E-07) and the superior temporal cortices (right: b = 0.00088 unit/day,

$t_{(63.74)}$ = 5.10, p = 2.94E-05; left: b = 0.00072 unit/day, $t_{(63.74)}$ = 4.18, p = 8.03E-04) presented a

high age-related rest CBF increase indicating late maturational process. Faster right over left

age-related rest CBF increase was more pronounced within the hippocampus (p = 0.014).

Finally, the rest CBF inter-regional correlation matrix between the predefined regions

showed a cluster of highly correlated regions with the following rank order: Superior

temporal, prefrontal, insula, sensorimotor and primary auditory cortices (see Figure 3).

<Figure 3>

The age-related rest CBF changes computed for the exhaustive list of 45 regions of interest

are available in SI Appendix, Table S3.

Discussion

Our study shows for the first time the dynamics of local rest functional brain maturation

throughout the first year of life using a non-invasive imaging method. Global rest CBF

increased significantly from 3 to 12 months of age and this increase was more pronounced in

the right than in the left hemisphere. Qualitative and quantitative analyses revealed marked

regional differences in local functional brain maturation. Subcortical structures such as basal

ganglia, thalamus, amygdala and hippocampus cortices are stable at 3 months. At the

cortical level, we observed two different maturational trajectories: first, a set of regions with

a low age-related rest CBF increase between 3 to 12 months, including the primary

auditory/visual cortices, the sensorimotor cortex and the insula. A second set of regions a

high age-related increase rest CBF increase between 3 to 12 months, including the superior

temporal and prefrontal cortices.

The increase in global rest CBF from 3 to 12 months of age that we describe here is

consistent with pioneers PET and SPECT studies showing increase in rest metabolism and

CBF during the same period (Chugani et al. 1987; Chiron et al. 1992). Furthermore, we

highlighted a hemispheric functional maturational asymmetry, with greater right than left

global rest CBF increase during the first year. This agrees with previous studies that showed

greater right than left rest CBF for these regions at birth (Lin et al. 2013) and from 1 year to 3

years old (Chiron et al. 1997), supporting the hypothesis that the right hemisphere

functionally matures earlier than the left.

Globally, our findings are in accordance with results from prior research based on histology,

structural and rest functional brain imaging that has revealed distinct maturation trajectories

of cortical regions and brain networks over the first year of life (Chugani and Phelps 1986;

Gilmore et al. 2012; Smyser and Neil 2015; Zhang et al. 2019). Firstly, at the histological

level, post-mortem data showed that the time course of synaptogenesis differs across

cortical regions. Indeed, a burst of synapse formation occurs between 3 and 4 months within

primary visual, auditory cortices somatosensory cortices, which appeared already mature at

3 months of life (Marin-Padilla 1970; Huttenlocher 1979, 1990; Michel and Garey 1984). In

non-human primates, Rakic and colleagues have shown a synchronic synaptogenesis in the visual, somatosensory, motor, and prefrontal areas (Rakic et al. 1986). In humans, synaptogenesis in the prefrontal cortex begins about the same time as in visual cortex, but it does not reach its peak period until age 8 months, continuing thereafter through the second year of life (Kostovic et al. 1995; Huttenlocher and Dabholkar 1997). Using quantitative electron microscopy in non-human primates, this remarkable overproduction followed by elimination of synapses in the prefrontal cortex have also been described (Bourgeois et al. 1994). These congruent findings strengthen our results as synaptic density is coupled to rest CBF. Secondly, concerning myelination, microstructural MRI maturational studies described a global maturation pattern characterized by early maturation of the sensorimotor cortex, followed by the other sensory cortices and then the associative cortices, including the prefrontal cortex (Dubois et al. 2008; Deoni et al. 2011). Finally, recent data obtained with resting-state functional MRI studies allowed to describe maturational changes of functional networks during the first year of life (Smyser et al. 2010; Gao et al. 2015; Smyser and Neil 2015; Wen et al. 2019). Especially, Gao et al. have described a maturation sequence starting with primary sensorimotor/auditory and visual then attention/default-mode, and finally executive control, prefrontal, networks (Gao et al. 2015). These different sequences of functional network maturation fit with and complement our results. Therefore, data coming from our study and previous rs-fMRI studies contribute to map a timetable of functional brain maturation during the first year of life.

Importantly, the spatial resolution of the ASL images allowed an accurate mapping of the age-related rest CBF changes. Consequently, we were able to describe insular local functional maturational evolution, which reaches its one-year pattern rather early, during the first months of life. A well-established literature and recently anatomical and resting-state functional MRI studies describe early human cortical development in areas close to the insula and radiating outward (Alcauter et al. 2015). This early insular maturation fits with its role in the integration of interoceptive stimuli, such as coolness, warmth and distension of the bladder, stomach or rectum (Craig 2009), but also in the integration of external stimuli, notably pain (Mazzola et al. 2009). In addition, it is highly pertinent, since the insula is a key structure for the baby's development and essential to baby's survival.

The spatial resolution improvement also allowed us to describe for the first time a
remarkably synchronous increase in rest CBF between the prefrontal and superior temporal
cortices (see Figure 3), both main components of the called "social brain" (Brothers et al.
1990). The late maturation of the prefrontal cortex had been previously described by
structural and functional brain imaging studies (Chugani and Phelps 1986; Gilmore et al.
2012). Noticeably, we describe here a late maturation within the posterior temporal regions
during the first year of life, particularly within the posterior superior temporal sulcus, a
region known to be highly implicated in social cognition (Zilbovicius et al. 2006).
Interestingly, the late and synchronous maturation of these two cortical structures
corresponds to the remarkable development of the baby's social skills through the first year
of life.

To the best of our knowledge, only 2 studies using ASL imaging have focused on brain
development during the first year of life, but a comparison with our results is limited due to
important differences in their methodological approaches. Duncan et al. studied a sample of
61 infants within a very narrow age-range from 3 to 5 months (Duncan et al. 2014). Their
main results describing a significantly greater rest CBF in the sensorimotor and occipital
regions compared with the dorsolateral prefrontal in this age-range are in accordance with
our results. In the second study, combining region of interest (ROI) and whole-brain analyses
on rest CBF, Wang et al. investigated a group of 8 7-month-old infants to a group of 8 13-
month-old infants (Wang et al. 2008). Although they showed rest CBF increase in the 13-
month-old group compared to the 7-month-old group mainly located in the frontal lobe,
they did not examine directly the age-related rest CBF slopes.

This study has some limitations. First, we used a linear model for data analysis. Although
cubic and quadratic fitting models did not improve our statistical models, it is improbable
that a linear model exactly fits functional cortical maturation. This issue can be addressed in
future studies by adding more and older subjects to further investigate the postnatal brain
rest functional maturation trajectory. Second, due to their age, all infants received light
premedication before the MRI to prevent motion artifacts, and all the scans were acquired
during sleep. No significant influence neither on the regional distribution of CBF (Fransson et
al. 2007; Doria et al. 2010; Carsin-Vu et al. 2018) nor in the default-mode network
connectivity (Greicius et al. 2008) has been reported to this premedication. Third, it is

important to stress that CBF is an indirect surrogate marker of focal neural activity by
providing two important metabolic substrate, oxygen and glucose, that are critically
important to neurons, synapses and astrocytes. Although, this neuro-hemodynamic coupling
plays an important role in brain development through dendritic sprouting, axonal growth,
synapse formation and vascular patterning, it does not necessarily imply a maturity of the
resulting neural activity. Finally, our study was performed in a clinical pediatric population.
To ensure that it could be comparable with a non-clinical population, we discarded all clinical
indications for MRI that could affect brain anatomy, function and further
neurodevelopmental disorders. In addition, all scans were strictly normal, and follow-up
confirmed a normal psychomotor development.

Defining typical trajectories of brain maturation provides references for a better
understanding of neurodevelopmental disorders and preterm effects on further brain
maturation. Because CBF reflects regional changes in synaptic density, ASL offers a
noninvasive approach to studying local brain function. Furthermore, the recent possibility to
implement ASL perfusion-based functional connectivity in conjunction to regular resting
state BOLD connectivity should be investigated for characterizing spatiotemporal and
quantitative properties of cerebral networks during brain maturation (Chen et al. 2015). In
conclusion, to our knowledge, our study is the first to describe and characterize dynamics
local functional brain maturation during the first year of life and provide insight into an
important and vulnerable neurodevelopmental period.

<u>Notes</u>

The authors declare no competing interests. The data that support the findings of this study
are available on request from the corresponding author. The data are not publicly available
due to them containing information that could compromise research participant consent.

Corresponding author:

Hervé Lemaître, PhD (herve.lemaitre@u-bordeaux.fr), Groupe d'Imagerie
Neurofonctionnelle, Institut des Maladies Neurodégénératives (CNRS UMR 5293), Université
de bordeaux, Centre Broca Nouvelle-Aquitaine, 146 rue Léo Saignat - CS 61292 - Case 28,
33076 Bordeaux cedex, France

References

Alcauter S, Lin W, Keith Smith J, Gilmore JH, Gao W. 2015. Consistent anterior-posterior segregation
of the insula during the first 2 years of life. Cereb Cortex. 25:1176–1187.
Allievi AG, Arichi T, Tusor N, Kimpton J, Arulkumaran S, Counsell SJ, Edwards AD, Burdet E. 2016.
Maturation of Sensori-Motor Functional Responses in the Preterm Brain. Cereb Cortex.
26:402–413.
Anderson AW, Marois R, Colson ER, Peterson BS, Duncan CC, Ehrenkranz RA, Schneider KC, Gore JC,
Ment LR. 2001. Neonatal auditory activation detected by functional magnetic resonance
imaging. Magn Reson Imaging. 19:1–5.
Arichi T, Gordon-Williams R, Allievi A, Groves AM, Burdet E, Edwards AD. 2013. Computer-controlled
stimulation for functional magnetic resonance imaging studies of the neonatal olfactory
system. Acta Paediatr. 102:868–875.
Arichi T, Moraux A, Melendez A, Doria V, Groppo M, Merchant N, Combs S, Burdet E, Larkman DJ,
Counsell SJ, Beckmann CF, Edwards AD. 2010. Somatosensory cortical activation identified by
functional MRI in preterm and term infants. Neuroimage. 49:2063–2071.
Ashburner J, Friston KJ. 2005. Unified segmentation. Neuroimage. 26:839–851.
Ball G, Aljabar P, Zebari S, Tusor N, Arichi T, Merchant N, Robinson EC, Ogundipe E, Rueckert D,
Edwards AD, Counsell SJ. 2014. Rich-club organization of the newborn human brain. Proc
Natl Acad Sci U S A. 111:7456–7461.
Biagi L, Abbruzzese A, Bianchi MC, Alsop DC, Del Guerra A, Tosetti M. 2007. Age dependence of
cerebral perfusion assessed by magnetic resonance continuous arterial spin labeling. J Magn
Reson Imaging. 25:696–702.
Bourgeois JP, Goldman-Rakic PS, Rakic P. 1994. Synaptogenesis in the prefrontal cortex of rhesus
monkeys. Cerebral Cortex (New York, NY: 1991). 4:78–96.
Brothers L, Ring B, Kling A. 1990. Response of neurons in the macaque amygdala to complex social
stimuli. Behav Brain Res. 41:199–213.
Carsin-Vu A, Corouge I, Commowick O, Bouzille G, Barillot C, Ferre JC, Proisy M. 2018. Measurement
of pediatric regional cerebral blood flow from 6 months to 15 years of age in a clinical
population. Eur J Radiol. 101:38–44.
Chen JJ, Jann K, Wang DJ. 2015. Characterizing Resting-State Brain Function Using Arterial Spin
Labeling. Brain Connect. 5:527–542.
Chiron C, Jambaque I, Nabbout R, Lounes R, Syrota A, Dulac O. 1997. The right brain hemisphere is
dominant in human infants. Brain. 120 (Pt 6):1057–1065.
Chiron C, Raynaud C, Mazière B, Zilbovicius M, Laflamme L, Masure MC, Dulac O, Bourguignon M,
Syrota A. 1992. Changes in regional cerebral blood flow during brain maturation in children
and adolescents. J Nucl Med. 33:696–703.
Chugani HT, Phelps ME. 1986. Maturational changes in cerebral function in infants determined by
18FDG positron emission tomography. Science. 231:840–843.
Chugani HT, Phelps ME, Mazziotta JC. 1987. Positron emission tomography study of human brain
functional development. Ann Neurol. 22:487–497.
Craig ADB. 2009. How do you feel--now? The anterior insula and human awareness. Nat Rev
Neurosci. 10:59–70.
Dehaene-Lambertz G, Hertz-Pannier L, Dubois J, Meriaux S, Roche A, Sigman M, Dehaene S. 2006.
Functional organization of perisylvian activation during presentation of sentences in
preverbal infants. Proc Natl Acad Sci U S A. 103:14240–14245.
Deoni SCL, Mercure E, Blasi A, Gasston D, Thomson A, Johnson M, Williams SCR, Murphy DGM. 2011.
Mapping infant brain myelination with magnetic resonance imaging. J Neurosci. 31:784–791.
Detre JA, Alsop DC. 1999. Perfusion magnetic resonance imaging with continuous arterial spin
labeling: methods and clinical applications in the central nervous system. Eur J Radiol.
30:115–124.

Detre JA, Wang J, Wang Z, Rao H. 2009. Arterial spin-labeled perfusion MRI in basic and clinical
neuroscience. Curr Opin Neurol. 22:348–355.
Doria V, Beckmann CF, Arichi T, Merchant N, Groppo M, Turkheimer FE, Counsell SJ, Murgasova M,
Aljabar P, Nunes RG, Larkman DJ, Rees G, Edwards AD. 2010. Emergence of resting state
networks in the preterm human brain. Proc Natl Acad Sci U S A. 107:20015–20020.
Dubois J, Dehaene-Lambertz G, Kulikova S, Poupon C, Huppi PS, Hertz-Pannier L. 2014. The early
development of brain white matter: a review of imaging studies in fetuses, newborns and
infants. Neuroscience. 276:48–71.
Dubois J, Dehaene-Lambertz G, Soares C, Cointepas Y, Le Bihan D, Hertz-Pannier L. 2008.
Microstructural correlates of infant functional development: example of the visual pathways.
J Neurosci. 28:1943–1948.
Duncan AF, Caprihan A, Montague EQ, Lowe J, Schrader R, Phillips JP. 2014. Regional cerebral blood
flow in children from 3 to 5 months of age. AJNR Am J Neuroradiol. 35:593–598.
Erberich SG, Panigrahy A, Friedlich P, Seri I, Nelson MD, Gilles F. 2006. Somatosensory lateralization
in the newborn brain. Neuroimage. 29:155–161.
Fransson P, Aden U, Blennow M, Lagercrantz H. 2011. The functional architecture of the infant brain
as revealed by resting-state fMRI. Cereb Cortex. 21:145–154.
Fransson P, Skiold B, Horsch S, Nordell A, Blennow M, Lagercrantz H, Aden U. 2007. Resting-state
networks in the infant brain. Proc Natl Acad Sci U S A. 104:15531–15536.
Gao W, Alcauter S, Elton A, Hernandez-Castillo CR, Smith JK, Ramirez J, Lin W. 2015. Functional
Network Development During the First Year: Relative Sequence and Socioeconomic
Correlations. Cereb Cortex. 25:2919–2928.
Gao W, Gilmore JH, Giovanello KS, Smith JK, Shen D, Zhu H, Lin W. 2011. Temporal and spatial
evolution of brain network topology during the first two years of life. PLoS One. 6:e25278.
Gao W, Gilmore JH, Shen D, Smith JK, Zhu H, Lin W. 2013. The synchronization within and interaction
between the default and dorsal attention networks in early infancy. Cereb Cortex. 23:594–
603.
Gilmore JH, Lin W, Prastawa MW, Looney CB, Vetsa YS, Knickmeyer RC, Evans DD, Smith JK, Hamer
RM, Lieberman JA, Gerig G. 2007. Regional gray matter growth, sexual dimorphism, and
cerebral asymmetry in the neonatal brain. J Neurosci. 27:1255–1260.
Gilmore JH, Shi F, Woolson SL, Knickmeyer RC, Short SJ, Lin W, Zhu H, Hamer RM, Styner M, Shen D.
2012. Longitudinal development of cortical and subcortical gray matter from birth to 2 years.
Cereb Cortex. 22:2478–2485.
Goldman-Rakic PS. 1987. Development of cortical circuitry and cognitive function. Child Dev. 58:601–
622.
Greicius MD, Kiviniemi V, Tervonen O, Vainionpaa V, Alahuhta S, Reiss AL, Menon V. 2008. Persistent
default-mode network connectivity during light sedation. Hum Brain Mapp. 29:839–847.
Hardy P, Varma DR, Chemtob S. 1997. Control of cerebral and ocular blood flow autoregulation in
neonates. Pediatr Clin North Am. 44:137–152.
Huttenlocher PR. 1979. Synaptic density in human frontal cortex - developmental changes and
effects of aging. Brain Res. 163:195–205.
Huttenlocher PR. 1990. Morphometric study of human cerebral cortex development.
Neuropsychologia. 28:517–527.
Huttenlocher PR, Dabholkar AS. 1997. Regional differences in synaptogenesis in human cerebral
cortex. J Comp Neurol. 387:167–178.
Innocenti GM, Price DJ. 2005. Exuberance in the development of cortical networks. Nat Rev Neurosci.
6:955–965.
Kagan J, Herschkowitz N. 2005. A young mind in a growing brain, A young mind in a growing brain.
Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers.
Knickmeyer RC, Gouttard S, Kang C, Evans D, Wilber K, Smith JK, Hamer RM, Lin W, Gerig G, Gilmore
JH. 2008. A structural MRI study of human brain development from birth to 2 years. J
Neurosci. 28:12176–12182.

Kostovic I, Judas M, Petanjek Z, Simic G. 1995. Ontogenesis of goal-directed behavior: anatomo-
functional considerations. Int J Psychophysiol. 19:85–102.
Kostovic I, Judas M, Rados M, Hrabac P. 2002. Laminar organization of the human fetal cerebrum
revealed by histochemical markers and magnetic resonance imaging. Cereb Cortex. 12:536–
544.
Leenders KL, Perani D, Lammertsma AA, Heather JD, Buckingham P, Healy MJ, Gibbs JM, Wise RJ,
Hatazawa J, Herold S, et al. 1990. Cerebral blood flow, blood volume and oxygen utilization.
Normal values and effect of age. Brain. 113 (Pt 1):27–47.
Li G, Wang L, Shi F, Lyall AE, Lin W, Gilmore JH, Shen D. 2014. Mapping longitudinal development of
local cortical gyrification in infants from birth to 2 years of age. J Neurosci. 34:4228–4238.
Licht DJ, Wang J, Silvestre DW, Nicolson SC, Montenegro LM, Wernovsky G, Tabbutt S, Durning SM,
Shera DM, Gaynor JW, Spray TL, Clancy RR, Zimmerman RA, Detre JA. 2004. Preoperative
cerebral blood flow is diminished in neonates with severe congenital heart defects. J Thorac
Cardiovasc Surg. 128:841–849.
Lin PY, Roche-Labarbe N, Dehaes M, Fenoglio A, Grant PE, Franceschini MA. 2013. Regional and
hemispheric asymmetries of cerebral hemodynamic and oxygen metabolism in newborns.
Cereb Cortex. 23:339–348.
Lyall AE, Shi F, Geng X, Woolson S, Li G, Wang L, Hamer RM, Shen D, Gilmore JH. 2015. Dynamic
Development of Regional Cortical Thickness and Surface Area in Early Childhood. Cereb
Cortex. 25:2204–2212.
Marin-Padilla M. 1970. Prenatal and early postnatal ontogenesis of the human motor cortex: a golgi
study. I. The sequential development of the cortical layers. Brain Res. 23:167–183.
Mazzola L, Isnard J, Peyron R, Guénot M, Mauguière F. 2009. Somatotopic organization of pain
responses to direct electrical stimulation of the human insular cortex. Pain. 146:99–104.
Meng Y, Li G, Lin W, Gilmore JH, Shen D. 2014. Spatial distribution and longitudinal development of
deep cortical sulcal landmarks in infants. Neuroimage. 100:206–218.
Michel AE, Garey LJ. 1984. The development of dendritic spines in the human visual cortex. Hum
Neurobiol. 3:223–227.
Mishra LD. 2002. Cerebral blood flow and anaesthesia: a review. Indian Journal of Anaesthesia.
46:87–95.
Nie J, Li G, Wang L, Shi F, Lin W, Gilmore JH, Shen D. 2014. Longitudinal development of cortical
thickness, folding, and fiber density networks in the first 2 years of life. Hum Brain Mapp.
35:3726–3737.
Paniukov D, Lebel RM, Giesbrecht G, Lebel C. 2020. Cerebral blood flow increases across early
childhood. Neuroimage. 204:116224.
Partridge SC, Mukherjee P, Henry RG, Miller SP, Berman JI, Jin H, Lu Y, Glenn OA, Ferriero DM,
Barkovich AJ, Vigneron DB. 2004. Diffusion tensor imaging: serial quantitation of white
matter tract maturity in premature newborns. Neuroimage. 22:1302–1314.
Petanjek Z, Judas M, Simic G, Rasin MR, Uylings HB, Rakic P, Kostovic I. 2011. Extraordinary neoteny
of synaptic spines in the human prefrontal cortex. Proc Natl Acad Sci U S A. 108:13281–
13286.
Rakic P, Bourgeois JP, Eckenhoff MF, Zecevic N, Goldman-Rakic PS. 1986. Concurrent overproduction
of synapses in diverse regions of the primate cerebral cortex. Science (New York, NY).
232:232–235.
Shaw P, Kabani NJ, Lerch JP, Eckstrand K, Lenroot R, Gogtay N, Greenstein D, Clasen L, Evans A,
Rapoport JL, Giedd JN, Wise SP. 2008. Neurodevelopmental trajectories of the human
cerebral cortex. J Neurosci. 28:3586–3594.
Smyser CD, Inder TE, Shimony JS, Hill JE, Degnan AJ, Snyder AZ, Neil JJ. 2010. Longitudinal analysis of
neural network development in preterm infants. Cereb Cortex. 20:2852–2862.
Smyser CD, Neil JJ. 2015. Use of resting-state functional MRI to study brain development and injury in
neonates. Semin Perinatol. 39:130–140.

Sowell ER, Thompson PM, Leonard CM, Welcome SE, Kan E, Toga AW. 2004. Longitudinal mapping of
cortical thickness and brain growth in normal children. J Neurosci. 24:8223–8231.
Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B, Joliot
M. 2002. Automated anatomical labeling of activations in SPM using a macroscopic
anatomical parcellation of the MNI MRI single-subject brain. Neuroimage. 15:273–289.
van den Heuvel MP, Kersbergen KJ, de Reus MA, Keunen K, Kahn RS, Groenendaal F, de Vries LS,
Benders MJ. 2015. The Neonatal Connectome During Preterm Brain Development. Cereb
Cortex. 25:3000–3013.
Varela M, Groves AM, Arichi T, Hajnal JV. 2012. Mean cerebral blood flow measurements using phase
contrast MRI in the first year of life. NMR Biomed. 25:1063–1072.
Wang Z, Fernandez-Seara M, Alsop DC, Liu WC, Flax JF, Benasich AA, Detre JA. 2008. Assessment of
functional development in normal infant brain using arterial spin labeled perfusion MRI.
Neuroimage. 39:973–978.
Webster MJ, Elashoff M, Weickert CS. 2011. Molecular evidence that cortical synaptic growth
predominates during the first decade of life in humans. Int J Dev Neurosci. 29:225–236.
Wen X, Zhang H, Li G, Liu M, Yin W, Lin W, Zhang J, Shen D. 2019. First-year development of modules
and hubs in infant brain functional networks. Neuroimage. 185:222–235.
Werner C. 1995. [Effects of analgesia and sedation on cerebrovascular circulation, cerebral blood
volume, cerebral metabolism and intracranial pressure]. Anaesthesist. 44 Suppl 3:S566-572.
Williams DS, Detre JA, Leigh JS, Koretsky AP. 1992. Magnetic resonance imaging of perfusion using
spin inversion of arterial water. Proc Natl Acad Sci U S A. 89:212–216.
Zhang H, Shen D, Lin W. 2019. Resting-state functional MRI studies on infant brains: A decade of gap-
filling efforts. Neuroimage. 185:664–684.
Zilbovicius M, Boddaert N, Belin P, Poline JB, Remy P, Mangin JF, Thivard L, Barthelemy C, Samson Y.
2000. Temporal lobe dysfunction in childhood autism: a PET study. Positron emission
tomography. Am J Psychiatry. 157:1988–1993.
Zilbovicius M, Meresse I, Chabane N, Brunelle F, Samson Y, Boddaert N. 2006. Autism, the superior
temporal sulcus and social perception. Trends Neurosci. 29:359–366.
573

Table 1: Age-related changes of the rest CBF values between 3 and 12 months of age.

	Hemisphere	estimate (unit/day)	95 % Confidence Interval	t(degree of freedom) = t-value	p-value (age)	p-value (age x hemisphere)
Whole brain	Right	0.0010	[6.87e-04, 1.26e-02]	t(55.71) = 6.64	1.36E-08	0.0074
	Left	0.00078	[4.97e-04, 1.07e-02]	t(55.71) = 5.34	1.74E-06	
Hippocampus†	Right	0.00026	[-2.75, 5.46E-04]	t(69.66) = 1.76	0.74	0.014
	Left	-0.00015	[-4.35E-04, 1.38E-04]	t(69.66) = -1.01	1	
Amygdala†	Right	-0.00008	[-3.55E-04, 1.86E-04]	t(83.74) = -0.60	1	1
	Left	-0.00019	[-4.60E-04, 8.11]	t(83.74) = -1.36	1	
Thalamus†	Right	-0.00030	[-5.36E-04, -6.79]	t(65.29) = -2.52	0.13	1
	Left	-0.00044	[-6.69E-04, -2.01E-04]	t(65.29) = -3.62	0.0051	
Primary visual cortex†	Right	0.00094	[5.17E-04, 1.37E-03]	t(60.07) = 4.33	5.2E-04	1
	Left	0.00085	[4.26E-04, 1.28E-03]	t(60.07) = 3.91	0.0021	
Primary auditory cortex†	Right	0.00030	[-6.59, 6.72E-04]	t(73.77) = 1.60	1	1
	Left	0.00028	[-8.89, 6.49E-04]	t(73.77) = 1.48	1	
Insula†	Right	0.00043	[1.84E-04, 6.85E-04]	t(84.2) = 3.37	0.010	1
	Left	0.00024	[-1.08, 4.90E-04]	t(84.2) = 1.86	0.59	
Sensorimotor cortex†	Right	0.00052	[1.13E-04, 9.22E-04]	t(59.12) = 2.50	0.14	0.25
	Left	0.00025	[-1.59E-04, 6.49E-04]	t(59.12) = 1.18	1	
Prefrontal cortex†	Right	0.00132	[9.38E-04, 1.69E-03]	t(60.12) = 6.80	4.86E-08	1
	Left	0.00118	[8,03E-04, 1.56E-03]	t(60.12) = 6.10	7.36E-07	
Superior Temporal cortex†	Right	0.00088	[5.43E-04, 1.21E-03]	t(63.74) = 5.10	2.94E-05	1
	Left	0.00072	[3.85E-04, 1.06E-03]	t(63.74) = 4.18	8.03E-04	

†: p- values Bonferroni corrected for the number of sub-parts of the brain. rest CBF values are normalized by the rest CBF measured within the basal ganglia and presented in arbitrary unit.

Figure 1: Age-related changes of the rest CBF values in predefined regions of interest

between 3 and 12 months of age. The whole brain in red, subset of stable subcortical regions

(hippocampus, amygdala and thalamus) in purple, subset of early maturing cortical regions

(primary visual and auditory cortices, insula and sensorimotor cortex) in green, subset of late

maturing cortical regions (prefrontal and superior temporal cortices) in blue. Each dot

represents a subject, and each line represents the estimated regression based on a linear

model for the left (empty dots and dashed line) and right (filled dots and solid line)

hemispheres. The rest CBF values are normalized by the rest CBF measured within the basal

ganglia and presented in arbitrary unit.

Figure 2: rest CBF values at 3, 6, 9 and 12 months of age displayed on the medial and lateral

view of the left and right hemispheres. The rest CBF values are normalized by the rest CBF

measured within the basal ganglia and presented in arbitrary unit. Surface rendering was

done using mri_vol2surf from freesurfer (https://surfer.nmr.mgh.harvard.edu/).

Figure 3: Correlogram of the correlation matrix for the rest CBF values in the predefined

regions of interest. Size and color of the circle represent the Pearson's correlation

coefficients. Correlation ordering is based on correlations with the first principal components

of the same matrix (i.e. similarity measure).

Left

Right

2.0

1.5

1.0

0.5

Rest CBF
relative value

Age (months)