A Computational Framework to Study the Primary Lifecycle

Metabolism of *Arabidopsis thaliana*

Running title: A Lifecycle Metabolic Modeling Framework

Authors: Wheaton L. Schroeder[1] and Rajib Saha[□*]

[1]Department of Chemical and Biomolecular Engineering, University of Nebraska – Lincoln,

USA

Correspondence:

Dr. Rajib Saha

rsaha2@unl.edu

Keywords: stoichiometric model, *Arabidopsis thaliana*, metabolism, core metabolism, carbon

metabolism, lifecycle, multi-tissue

Language Style: American English

Article Type: Research Article

Article Length: 9539 words (excluding abstract, summary, references, captions)

Number of Figures: 5

Number of Tables: 1

Abstract

Stoichiometric Models of metabolism have proven valuable tools for increased understanding of

metabolism and accuracy of synthetic biology interventions to achieve desirable phenotypes.

Such models have been used in conjunction with optimization-based and have provided

"snapshot" views of organism metabolism at specific stages of growth, generally at exponential

growth. This approach has limitations in that metabolic history of the modeled system cannot be

studied. The inability to study the complete metabolic history has limited stoichiometric

metabolic modeling only to the static investigations of an inherently dynamic process. In this

work, we have sought to address this limitation by introducing an optimization-based

computational framework and applying to a stoichiometric model of the model plant *Arabidopsis*

thaliana of four linked sub-models of leaf, root, seed, and stem tissues which models the core

carbon metabolism through the lifecycle of arabidopsis (named as p-ath780). Uniquely, this

framework and model considers diurnal metabolism, changes in tissue mass, carbohydrate

storage, and loss of plant mass to senescence and seed dispersal. p-ath780 provide "snapshots" of

core-carbon metabolism at one hour intervals of growth, in order to show the evolution of

metabolism and whole-plant growth across the lifecycle of a single representative plant. Further,

it can simulate important growth stages including seed germination, leaf development, flower

production, and silique ripening. The computational framework has shown broad agreement with

published experimental data in tissue mass yield, maintenance cost, senescence cost, and whole-

plant growth checkpoints. Having focused on core-carbon metabolism, it serves as a scaffold for

lifecycle models of other plant systems, to further increase the sophistication of *in silico*

metabolic modeling, and to increase the range of hypotheses which can be investigated *in silico*.

As an example, we have investigated the effect of alternate growth objectives on this plant over

the lifecycle.

Author Summary

In an attempt to study the evolution of metabolism across the lifecycle of plants, in this work we

have created an optimization-based framework for the *in silico* modeling of plant metabolism

across the lifecycle of a model plant. We then applied this framework to four core-carbon tissue-

level (namely, leaf, root, seed, and stem) stoichiometric models of the model plant species

Arabidopsis thaliana, and further informed this framework with a wide array of published *in vivo*

data to increase model and framework accuracy. Unique to the p-ath780 model, comparted to

other models of plant metabolism, is the simultaneous considerations of diurnal metabolism,

carbohydrate storage, changes in tissue mass (including losses), and changes in metabolism with

respect to plant growth stage. This provides a more complete picture of plant metabolism and

allows for a wider array of future studies of plant metabolism, particularly since we have only

modeled the core carbon metabolism of *A. thaliana,* allowing this work to serve as a framework

for studies of other plant systems.

Introduction

The use of synthetic biology for the engineering of uni- and multi-cellular organisms to enhance

desirable phenotypes in microbe, plant, and animal systems, has been well established and has

been capable of affecting the lives of millions of individuals, such as in the case of artemisinin

production in yeast or enhancing nutritional value of agricultural products [1-2]. Synthetic

biology techniques have been applied to many plant systems such as tomatoes [3], rice [4], and

maize [5] to produce enhanced phenotypes often with application to human nutrition [2], pest

resistance [5], and resilience to abiotic stresses [6]. Many of these efforts have focused on a

genetic understanding and manipulation of the plant system (or plant tissue) in question, having

relied on intuitive interventions such as changes in regulation, insertion of new gene(s), and

deletion of gene(s) from competing pathway(s) [2,5,6]. Alternatively, computation-based

systems biology approaches, such as the use of stoichiometric genome-scale models (GSMs) of

metabolism, have predicted non-intuitive genetic interventions [7] by accounting for Gene-

Protein-Reaction (GPR) links and understanding how a gene knockout, or a change in gene

regulation, affects the entire system through tools such as Flux Balance Analysis (FBA) [8],

OptKnock [9], and OptForce [10]. Other tools are built upon previously existing tools, such as

dynamic FBA (dFBA), which performs FBA over windows of time by solving a non-dynamic

linear or a static linear problem, both of which integrate system variables over discrete time

windows to solve to metabolite concentration, in addition to reaction flux [11]. Such tools have

led to enhanced mechanistic understanding for exploring the system-wide effects of synthetic

biology interventions especially in a microbial or a fungal system, such as *E. coli* [10],

cyanobacteria [12], and yeast [13].

Stoichiometric global plant models, which treat the metabolism of the plant as a single unit, have

been developed for *Arabidopsis thaliana* (hereafter arabidopsis) [14-17], *Zea mayz* (maize) [18],

Sorghum bicolor (sorghum) [19], *Saccharum officinarum* (sugarcane) [19], *Brassica napus*

(rapeseed) [19], and *Oryza sativa* (rice) [20]. These models have sought to analyze metabolic

maintenance, response to abiotic stimuli, enzyme regulation changes, and metabolism as a whole

at steady state (or pseudo-steady state). In addition, tissue-specific single-unit models have been

reconstructed for various arabidopsis tissues [21], a maize leaf [22], and a barley seed [23] to

better understand how present metabolites, metabolic pathways, and nutrient availability differ

between tissues. Multi-tissue models have been created to characterize whole-plant metabolism

for arabidopsis [16] and barley [17] and subsequently to study whole-plant metabolic response to

the diurnal cycle and the source-to-sink relationship of leaves and seeds [16,17]. These studies

either have considered metabolism at a single point [14,15,18-20], having taken a metabolic

"snapshot" of a single point in growth time (often in the exponential growth phase) or have

considered a single diurnal cycle [16]. This approach has been inherently limited in that

metabolism is a dynamic and cumulative process. To clarify, metabolic state is dependent on

both on external factors, such as availability of light, carbon sources, and availability of

micronutrients, which these "snapshots" have captured, but also are dependent on metabolic

history. These limitations have been inconsequential for single-cell systems in that laboratory

apparatuses have held single-cell cultures at an exponential growth state; therefore, the

"snapshot" approach has given good approximation of metabolism in these steady-state systems.

In contrast, multi-cellular organisms, such as plants, will have passed through multiple and

distinct stages of growth throughout its lifecycle [24], and the organism cannot be held at a

steady state growth point. For this study, we have chosen *Arabidopsis thaliana* as the multi-

cellular organism for several reasons. Firstly, since the advent of modern genetics, arabidopsis

has served as a model plant species in that it has a small genome; therefore, arabidopsis has been

well studied. Secondly, arabidopsis has a limited number of basic tissues which will have

required the construction of a tissue-level model. Thirdly, arabidopsis has at least two distinct

metabolic modes dependent on the availability of light. When studying the effects of a synthetic

biology intervention on a plant system, such as arabidopsis, understanding the evolution of

metabolism throughout the plant lifecycle can increase understanding of the cumulative effect of

a synthetic biology intervention. The multi-tissue Edinburgh forest model, which has made use

of Ordinary Differential Equations (ODEs) rather than stoichiometric matrices, has modeled the

lifecycle of a tree for the purposes of studying lumber yield [16,17]; however, the intent of the

aforementioned model has not been to consider individual reactions or genetic interventions, and

therefore the GPR links which are central to the sought understanding and testing hypotheses

when using SMs have not been included.

In this work, a core carbon stoichiometric metabolic model of arabidopsis has been reconstructed

which consists of major primary carbon metabolism pathways, including, but not limited to,

photosynthesis; the citrate cycle; starch and sucrose synthesis; fatty acid synthesis and

degradation; and amino acid synthesis. The multi-tissue arabidopsis stoichiometric model,

referred to as p- ath780 has 1033 total (and 633 unique) reactions (R), 1157 total (and 325

unique) metabolites (M), and accounts for 780 genes (G) including 42 chloroplastic and 11

mitochondrial genes. The model p-ath780 (**p**lant-scale primary **a**rabidopsis **th**aliana model

including **780** genes) consists of four tissue-level models of metabolism: leaf (R: 537, M: 479,

and G: 703), root (R: 130, M: 126, and G: 250), seed (R: 428, M: 411, and G: 529), and stem (R:

160, M: 140, and G: 250). The models are linked to one another and their respective environment

by a comprehensive Flux Balance Analysis (FBA)-based [8] optimization framework [25] which

considers both inter-tissue and environmental interactions. These four tissues have been chosen

for model reconstruction to represent core plant functions. The root has been chosen and

reconstructed for nutrient uptake and growth; the leaf for photosynthesis, carbon fixation, and as

a source tissue for plant nutrition; the seed for metabolite storage and a sink tissue for metabolic

investment; and the stem for metabolic transport and acting as a conduit for all metabolic

interactions between other tissues. The dFBA method determines metabolite concentrations at

the start and end points of the time frame [11], whereas the method developed does not focus on

concentrations and considers multiple points within the time interval to make more accurate

time-derivative estimates of steps in plant and tissue masses, as well as plant maintenance and

senescence costs. The optimization framework of the p-ath780 model has taken a series of

metabolic "snapshots" of arabidopsis metabolism throughout the lifecycle of a single

representative plant subject to diurnal status, carbohydrate storage/uptake, changes in tissue mass

(including losses), changes in relative tissues masses (due to growth stages), and changes in

metabolism with respect to plant growth stage. p-ath780 has taken "snapshots" at hour intervals,

and information from these snapshots have advanced plant and tissue masses forward one hour,

when the next "snapshot" is taken. The series of "snapshots" produced by p-ath780 has given a

framework for the investigation of the central metabolism of arabidopsis across its lifecycle.

Several different objectives for this optimization-based framework have been investigated, with

the default framework being the maximization of plant growth. Other alternative objectives

investigated have included linear photonic efficiency and seed fatty acid production. This

framework with the default objective has shown general agreement with experimental data and is

potentially useful as an initial framework for other plant systems.

Results

Reconstruction of arabidopsis primary carbon metabolism in tissue-specific models. Figure 1

shows an overview of the workflow designed for determining the optimal reaction rates and mass

step for each "snapshot" (top), how these "snapshots" have been advanced from one time point

to the next; and how tissues have interacted at various stages of growth along with listing some

important characteristics of a given growth stage that differs from other stages (bottom). In order

to track the important metabolic interactions and transactions within and between major tissues

of arabidopsis plant, namely seed, leaf, root, and stem, corresponding tissue-level metabolic

models have been reconstructed. Model files for each tissue can be found in Supplemental Files

1 (seed), 2 (leaf), 3 (root), and 4 (stem). Figure 2 shows a summary of the distribution of model

reactions across KEGG-defined pathways of each tissue model and an overview of reasons for

reaction inclusion through confidence scoring (see Method section) [26]. Figure 2(A)

summarizes the pathways common to all tissues, Figure 2(B) summarizes the pathways common

to seed and leaf tissues, and Figure 2(C-G) graphically summarize the sources of reactions in

each tissue model and p-ath780 as a whole through confidence scores (see methods section) [26].

First, the seed model has been reconstructed based on gene annotations and available MFA data

[27] and then tissue model reactions have been distributed across five compartments based on

literature evidence (see list of works cited in Supplemental File 5): extracellular space, cytosol,

non-green plastid, inner mitochondria, and outer mitochondria. Next, transport and exchange

reactions have been added to the model based on literature evidence (see list of works cited in

Supplemental File 5) or modeling necessity to increase model connectivity [26]. The leaf model

has been reconstructed using common reactions and pathways from the seed model and having

added new pathways and functions essential to the major functions of the leaf tissue, such as

photosynthesis [18]. In addition to the five compartments in the seed model, the leaf model

contains chloroplast and thylakoid compartments. Similarly, by having extracted common

reactions/pathways from the seed model, the root and stem models have been reconstructed. The

root and stem models have been focused primarily on nutrient uptake (root) and transport (root

and stem). Both these models contain necessary transport/exchange reactions to ensure model

connectivity and to facilitate their roles in transport processes. The stem and root models have all

the subcellular compartments present in the seed model. Once initial reconstructions have been

accomplished, thermodynamically infeasible cycles in addition to atom and charge imbalances

have been resolved [26] and tissue-specific biomass equations based on literature information

have been defined [18,27,28].

Figure 1. The design-build-test used cycle in constructing the p-ath780 model, where each box

represents a step used in this cycle. The numbers in the lower left corner of each box indicates

the approximate order in which these steps are undertaken for this cycle. This cycle is repeated

until the *in silico* representation of *Arabidopsis thaliana* that is p-ath780 converges satisfactorily

with *in vivo* experimental data as described in the results section.

Figure 2. A heuristic look at the four tissue models in terms of number of reactions in various

KEGG-defined pathways which provides some clarity as to the metabolic functions of each

model (A and B) and in terms of the sources of included model reactions, indicated by

confidence scores (C through F). A) A bar graph showing tissue model reaction counts in

KEGG-defined pathways (with the exceptions being the user defined pathways of exchange and

transport) common to most or all tissue (threshold: at least one model has at least three reactions

in that pathway). B) Additional KEGG-defined pathways common to the seed and/or leaf model

as these models contain more complete metabolism. C) - F) The source of each reaction included

in the models through confidence scores for each tissue. G) The source of all reactions included

in the p-ath780 model. See the methods section discussion related to confidence scores in this

model.

Development and tuning of the p-ath780 model. Once these core tissue models have been

reconstructed and curated, these have been linked within a comprehensive FBA-based

optimization framework (provided in Supplemental File 6) for *in silico* representation of

metabolic behavior across the arabidopsis lifecycle. This framework has next been applied to the

p-ath780 model that includes all four tissue-specific models and has 1033 total reactions, 1157

total metabolites, and 780 unique genes. Further details of the model development steps can be

found in the methods section. The seed and leaf tissue have been selected to model an important

source-to-sink relationship, whereas the stem and root tissues have been included to model

nutrient transport and nutrient uptake in arabidopsis, respectively. The FBA-based framework

has defined constraints related to tissue interactions and whole-plant growth heuristics based on

experimental data, and also helped align *in silico* growth with experimentally determined *in vivo*

growth through the modified design-build-test cycle shown in Figure 3, which will be discussed

in greater detail later in this subsection.

Figure 3. A simplified workflow of the calculations made to estimate the plant mass step size

taken from one "snapshot" to the next (large top box) and visual representation of how these

"snapshots" are strung together and grouped into stages and transitions (large bottom box). Each

"snapshot" has been represented visually as small boxes containing initial time point, a figure

highlighting major metabolic interactions, initial plant mass, step estimate, and cumulative

relative growth rate. The contained figures show some major metabolic interactions across the

plant system boundaries (full-headed arrows crossing the dashed system boundary) and indicates

which tissues interact (single-headed arrows with circles to indicate a shared metabolic pool

between the two tissues) for the given stage of growth which is indicated by the beveled box

below the group of "snapshots". The beveled boxes below a group of snapshots indicate the stage

name (or transition name), the time points in growth which this stage encompasses, and some

distinguishing characteristics of that stage.

The output of this framework has given metabolic "snapshots", consisting of plant mass, growth

rate, and flux rate of each reaction, at one-hour intervals across 61 days of growth, as the plant

disperses all new seeds (through silique shattering) by 61 days after germination (DAG) [24].

After 61 DAG the plant begins to desiccate, eventually resulting in plant death [24]. The p-

ath780 model is not used to model plant metabolism after 61 DAG because no *in vivo* data has

been found in literature concerning the metabolomics of plant death and desiccation. This

optimization-based framework has allowed for the sampling of changes in central carbon

metabolism at different stages in the arabidopsis lifecycle (see Figure 3). For all the following

analyses, the objective of this framework at each point has been the maximization of the sum of

all tissue biomass production rates, unless otherwise indicated.

In having determined the mass steps taken for each hour intervals, three FBA-like calculations at

0, $^1/_3$, and $^2/_3$ hours past the hour have been made to increase the accuracy of the derivative

estimate by an explicit numerical integration method calculating the mass step at each hour

interval using Heunn's rule for the third order Runge-Kutta method (see methods for greater

detail). Figure 4 shows a more detailed workflow for each individual step in the form of a

workflow diagram. Optimal flux points at every whole hour have been saved as the optimal flux

rates at that growth point. To evaluate these balanced flux estimates, Flux Variability Analysis

(FVA) [29] has been performed, at nine points, selected to represent each non-transition growth

stage and diurnal status in those stages, subject to all growth constraints and a growth rate

equivalent to the optimal growth rate (see methods for enumeration of these points).

Figure 4. The workflow diagram of the p-ath780 model, including inputs (orange), outputs

(green), and internal workflow (blue). The inputs for the p-ath780 model include each individual

tissue model, a file of growth specifications, and a list of point at which to take metabolic

"snapshots". The internal workflow has read these inputs and then used them to construct model

objects (bold, Times New Roman text) which are used to perform FBA, to solve for what the

plant mass step is from the current to the next "snapshot", and to perform FVA. For each

iteration, the time of the snapshot is stepped forward 1/3 step (hour), the FBA model object is

solved, the mass step is calculated, and process is repeated. Every third iteration (e.g. where step

= 3), Heun's method for a third order Runge-Kutta is used to estimate the plant mass step from

the previous whole hour to the next whole hour and FVA is performed on the model at the

previous whole hour using saved values. Once all iterations are complete (e.g. model is at final

time point), then the output files are written.

The simulations of the p-ath780 model has been advanced through several growth stages using

time point for changes in growth stage taken from experimental data [24]. In the seed

germination stage, uptake of fatty and amino acids from seed storage has been modeled as a

constant rate of fatty acid usage which results in all stored fatty and amino acids being depleted

by the end of the seed germination stage [30]. The 12 hours of light and 12 hours of dark diurnal

rhythm has been chosen to match experimental conditions for studies on starch and sucrose

storage/uptake dependence on the diurnal cycle [31]. These patterns have been fit to a sine wave

model constraint with $\pm 1\%$ tolerance. In growth stages when plant tissue ratios have been

constant, the tissue mass ratio values had been taken from values typical for herbaceous plants

[32].

Returning to the design-build-test cycle used to improve the p-ath780 model, experimental data

related to plant growth and plant growth stages have been collected from a variety of literature

sources to serve as checks for the accuracy of the modeled system [24,33]. The first set of

experimental data has included mass data, including whole plant and individual tissue. At

approximately 17, 24, and 31 DAG the total dry plant mass should be between 0.5 and 2.0 mg; 2

and 5 mg; and 10 and 30 mg, respectively [33]. Once the design-build-test cycle has been

completed, the p-ath780 model has shown a total dry plant mass of 0.554 mg at 17 days (408

hours), 3.74 mg at 24 days (576 hours), and 25.2 mg at 31 days (744 hours) after germination,

demonstrating growth consistent with *in vivo* data. Furthermore, the relative growth rate for the

first 31 days of plant growth has been reported as between 0.21 and 0.25 day^{-1} [33], and the final

p-ath780 has shown a relative growth rate of 0.246 over this time period. To adjust model

behavior in latter stages of growth, tissue-specific mass data has been obtained from literature.

Specifically, the dry weight of the stem, the leaves, and the seeds has been reported as

approximately 188 mg (standard deviation 39.3 mg), 163.7 mg (standard deviation 52.0 mg), and

127.9 mg (standard deviation 52.7 mg), respectively [24]. As p-ath780 models both plant growth

and loss of seed (and other) mass in the silique ripening stage, the peak mass of each of these

tissues has been comparted to this data. In the final p-ath780 model, the peak mass of the stem,

leaves, and seeds has been determined as 189 mg, 177 mg, and 130, respectively, all of which are

within one standard deviation of the experimental value (see the methods section for how tissue

masses are determined). In summary, through the results of the design-build-test cycle

implemented, *in silico* tissue and plant mass values are similar to *in vivo* data, thus showing

strong agreement with respect to growth trends.

In early rounds of model reconstruction, it has been noticed that the plant model's photosynthesis

is too efficient at fixing carbon. This is due to the fact that plants do not make full use of

available light source(s), but the reconstructed metabolic model had been. Published *in vivo* data

which has been used in the modeling and verification of p-ath780 made use of fluorescent lights,

which have tight transmission spectra peaks at 544 and 609 nm [34]. In contrast, peak

absorbance for plant leaves is at approximately 440 and 680 nm [35]. The problem of the

availability of light has been addressed by scaling the transmission of the fluorescent lights by

the absorbance of plant leaves. This has left approximately 21.06% of light transmitted by the

fluorescent bulbs usable by the plant (see Supplemental File 5 and methods). An additional

restriction, namely biomass yield, has also been placed on metabolic efficiency in plant systems.

Biomass yield has been defined as the carbon fraction of biomass produced appearing in new

growth for each unit of carbon used for growth [36]. This yield value accounts for repair of

existing biomass and replacement of lost biomass. Experimentally, this value has been identified

as generally between 0.7 and 0.85 [37]. Here, for p-ath780 mass values to align with

experimental data, two separate mass yield vales have been set at 0.32 and 0.23 for when the

plant system lacks and has seed tissue respectively. This represents an incongruity with

experimental evidence, although this value is still in the same order of magnitude as

experimental evidence. All files necessary for p-ath780 have been included with this work in

Supplemental Files 7 through 16. The *in silico* results of the final p-ath780 model can be found

in Supplemental File 17.

In silico *Plant Growth under* Alternative Objective Functions. A total of six different objective

functions for p-ath780 have been investigated and a summary of that investigation has been

shown in Table 1. In all cases, root and stem tissue objectives have been defined as biomass

production, where the leaf and seed objective functions are varied. When any tissue has a non-

biomass objective, that objective is weighted by some scaling factor (either α for light-based

objectives or β for fatty acid-based objectives) to ensure the new terms do not dominate or be

insignificant compared to biomass (e.g. be an order or magnitude or more different) and to

investigate the different effects of weight factors. The first row (green) of Table 1 contains the *in*

vivo arabidopsis data which has been used as targets and verification of the p-ath780 model. The

second row (blue) contains the *in silico* data from p-ath780 where the objective for all tissues is

biomass production (default objective), and has summarized the findings of the preceding

subsection. For the mathematical definition of this and other objective functions discussed here

see the methods section. The next two objective functions presented (grey) we have considered

set the seed tissue-level objective as the maximization of fatty acid stored in the seed tissue at

two different weight factor values (β). At low β values, this causes a scavenging of carbon

wasted in the plant metabolism which is then diverted to the seed fatty acid production without a

change in plant growth. At high β values, this alternate seed objective results in stunted plant

growth as carbon used elsewhere is diverted to the seed tissue. This alternate objective function

has no effect when the plant does not have seed tissue present. Photonic efficiency for the leaf

tissue has been attempted as an alternative objective function and results are reported in the next

two grey rows; however, depending on the weight of the photonic efficiency parameter, the

model is generally photophobic (no light has been uptaken) or grows as normal. In all

photophobic growth cases, the plant mass of the model eventually becomes negative, leading to

nonsense in later time points, thus the results of these α values are not reported. No α value has

been identified which produced a result between these extremes (photophobic and normal

growth), and these attempts are not included in this work. The full output of each reported result

can be found in Supplemental File 17. The final objective function investigated (and reported in

Table 1) is one which combined the leaf linear photonic efficiency objective and the seed fatty

acid storage objective at a moderate weight value (values enumerated in Table 1). As with other

linear photonic efficiency objectives, the amount of light uptaken by the plant is unaffected, and

as with other investigations of the seed fatty acid objective, the plant growth is stunted when the

seed tissue is present. In summary, the p-ath780 model is robust to small and moderate

perturbations in the objective related to photonic efficiency, fails with large perturbations to

photonic efficiency objectives, and results in continuously changeable growth levels to

metabolite production objectives.

Discussion

In the current work, a multi-tissue core metabolism stoichiometric model, including leaf, root,

seed, and stem tissues, of *Arabidopsis thaliana* has been reconstructed (Figure 3), and linked in

an FBA-based optimization framework (Figure 1). This framework has been embedded in a

workflow (Figure 1) which has simulated how plant metabolism evolves over time with respect

to the presence or absence of light, the transition to different growth stages, and the gain or loss

of tissues (such as seed). This model has incorporated a wide variety of data which has not been

incorporated in other stoichiometric modeling efforts such as the effect of plant mass, the effect

of tissue mass difference on tissue interactions, whole-plant growth heuristics such as yield, the

availability of usable light, and biomass-based plant maintenance (as opposed to ATP-based).

The tissue models taken together with these literature-based constraints has been named the p-

ath780 model. The whole-plant growth characteristics of p-ath780 have shown general

agreement with experimental data, particularly with respect to whole plant mass at certain

growth milestones and lifecycle tissue yields.

The design-build-test cycle used to develop and tune p-ath780, shown in Figure 3, has been

implemented. As a result, in the final p-ath780 model, *in silico* predictions compared well to *in*

vivo data, particularly plant, leaf, seed, and stem masses, with the exception of biomass yield.

The incongruity between *in* vivo and *in silico* biomass yield has likely resulted from the p-ath780

model only having included primary carbon metabolism, which in turn means that plant biomass

has been built entirely from generally less metabolically expensive primary metabolites. This had

resulted in too efficient biomass production, hence the lower yield for the model. This

discrepancy in biomass yield has served to highlight the large effect of secondary metabolism on

plant growth and has served as a correction factor on the model due to the lack of modeled

secondary metabolism. In addition, likely the plant mass yield is lower when seed tissue is

present because flower tissue is metabolically expensive yet is not modeled in this work. In

additional, biomass drains for plant senescence and maintenance have been included [36,37].

Once the model has been developed, six different objective functions have been applied to it,

including the default objective of maximizing plant growth, linear photonic efficiency, and seed

fatty acid production. In summary, the p-ath780 model is robust to small and moderate

perturbations in the objective related to photonic efficiency, breaks when large perturbations are

made to the photonic efficiency objective, and is capable of some fine tuning with respect to

metabolite production objectives. The behavior of p-ath780 with respect to the linear photonic

efficiency objective function is due to multiple factors. First, as the partially photophobic case

exists, this suggests that seed tissue is the most metabolically expensive tissue to create. This is

as expected because the seed tissue requires storage of high-energy molecules such as fatty acids,

proteins, and sugars to feed its embryo when dispersed. Further, the rate of biomass production

for all tissues are linked in the optimization-based framework. Secondly, seed tissue has a target

fraction of overall plant mass which it must grow to for each hour interval of the flower

development stage. If seed tissue is too metabolically expensive to produce, relative to the cost to

uptake more light, it appears that the solution strategy then becomes to decrease the mass of

other tissues while leaving the growth of the seed tissue to be minimal. This can result in sharp

changes in mass which falls outside the realm of stability for Heunn's third order Runge-Kutta

rule, resulting in predictions with no biological relevance. Thirdly, biomass composition and

metabolic cost is not dependent on the amount of light uptaken, so the biomass cost is constant

with respect to light uptaken so there is no steady equilibrium between the two terms except at

the extremes. Fourthly, minimizing light uptake and maximizing biomass growth as objectives

are competitive, increase light uptake results in increased growth in stoichiometric models. In

contrast, this is not an issue for maximizing fatty acid production as biomass partially consists of

fatty acids, therefore these two objectives can be complimentary to some degree and a variety of

β values can be used without the model failing to find a solution. Theoretically, there exists some

highly-specific value of α at which the cost to the light needed to drive growth is balanced with

the rate of production of new biomass, but this is an unsteady equilibrium which when the value

of α is slightly perturbed finds the new equilibrium at either extreme. Therefore, the value of α

might be imagined as a fulcrum between the two terms as illustrated in Figure 5. Figure 5(A)

restates the linear photonic efficiency objective equation, and Figure 5(A-D) represents the

action of α as a fulcrum. Figure 5(D) in particular illustrates why the p-ath780 model is robust to

changes in the value of α below this theoretical balance point.

Figure 5. This figure highlights how the weight factor, α, fails to reach some kind of equilibrium

between light uptake and biomass production in the linear photonic efficiency objective. A)

Restates the linear photonic efficiency objective function. B) Shows the theoretical balance

which might exist between light uptake and biomass production at some highly specific value of

α. C) Shows how as slight increase in α from that point causes light to outweigh biomass in

terms of influence on the objective function value. This results in photophobic growth as light

"outweighs" growth. D) Shows how as slight decrease in α from that point causes light to

outweigh biomass in terms of influence on the objective function value. This results in normal

growth as growth "outweighs" light.

This work does not account for diurnal rhythms in the transcriptome of *Arabidopsis thaliana* for

several reasons. Firstly, the majority of transcriptomic studies have focused on the regulatory

network of proteins which regulate metabolism based on the availability of light and rhythm

[43,44], rather than considering metabolic proteins which are represented in the p-ath780 model.

Secondly, tissue-specific diurnal transcriptomic information is only available for the leaf tissue

[43,44]. Further, these experiments generally consider a single point in the growth cycle of

arabidopsis under specific growth conditions. The framework of p-ath780 is already highly

constrained, and that the inclusion of too much data will invariably cause model failure. This is

because *in vivo* experiments, in general and those used in this work, often occur under different

conditions, at different points in the plant lifecycle, have different methods to some degree, or

even seem quantitatively difference due to the noise inherent to biological systems, making

alignment of quantitative *in vivo* data from too many sources impossible. In this work, we have

decided to use data which described a wide range of time point in arabidopsis growth, such as

biomass yield, relative growth rate, growth up to a certain time point, and overall tissue yield,

rather than data which may be specific to a single point in growth, such as transcriptomics.

This work provides the basis for much future development and sophistication. For instance, the

current p-ath780 model could be further sophisticated by adding the secondary metabolism of the

plant system, which is a considerable resource drain in many plant systems. Further, at present

several simplifications are made regarding tissues, particularly related to seed tissue. For

instance, the model currently assumes that when the plant is flowering, that flower biomass and

metabolism is roughly equivalent to that of the seed. While this resulted in a simpler model, this

model cannot be then used to investigate certain metabolic hypothesis such as the cost to the

plant resulting from flower pigmentation, pollen, and nectar production. Future work will include

producing models for other plant tissues, such as flowers. In addition, as this is a core carbon

metabolism model, it is likely quite similar to the core metabolism of other plant systems;

therefore, the p-ath780 model can serve as a basis for the development of lifecycle models for

other plant systems, particularly annual eudicots which are of agricultural interest, such as rice

(*Oryza sativa*), potatoes (*Solanum tuberosum*), tomatoes (*Solanum lycopersicum*), and soybeans

(*Glycine max*).

Methods

Overview of the reconstruction of core metabolic models of leaf, root, seed, and stem tissues.

The seed tissue model. The general workflow which has been used for the development of the

four core tissue models has been illustrated in Figure 3. We have developed the seed model first,

with the central metabolic pathways based on a Metabolic Flux Analysis (MFA) of four seed

genotypes published previously [27]. We have then manually filled gaps in this model with

reactions based on literature and genomic evidence [26] or with reactions being necessary for

ensuring model connectivity. The stoichiometric coefficients of biomass precursors have been

determined using sink reactions, dry biomass weight composition, and amino acid mass ratios

provided in a previous work [27] (see Supplemental File 22). The resultant seed tissue model has

focused on storage, respiration, and growth, and consists of 428 reactions, 529 genes, and 411

metabolites (included as Supplemental File 1).

The leaf tissue model. Next, we have reconstructed the leaf model by taking common

reactions/pathways from the seed model and adding synthesis pathways for amino acids that are

not synthesized in the seed, in addition to photosynthesis, carbon fixation, gluconeogenesis, and

transport reactions. We have then developed the biomass equation for the leaf tissue using that of

a previously published Arabidopsis model [18] (see Supplemental File 22). The resultant leaf

tissue model has focused on photosynthesis, respiration, gas exchange, fatty acid synthesis, and

growth, and contains of 537 reactions, 703 genes, and 479 metabolites. We have included the

leaf model with this work as Supplemental File 2.

The root and stem tissue models. We have constructed the root and stem models, similarly, by

extracting common reactions/pathways from the seed model and adding necessary transport and

exchange reactions. Then exchange reactions have been added to allow the root to be linked to

micronutrient uptake processes from the soil, and the stem to be involved in inter-tissue transport

processes. In the absence of Arabidopsis-specific estimates, the dry weight composition of

switchgrass (*Panicum virgatum*) root and stem [28] have been assumed to be equivalent to the

biomass composition of these tissues in Arabidopsis. Due to the low detail level of the dry

weight composition analysis, the biomass of root and stem tissues have been composed entirely

of carbohydrates. The resultant root tissue model has focused on nutrient uptake, transport, and

growth, consisting of 130 reactions, 250 genes, and 126 metabolites, while the stem tissue model

focuses on transport and growth, consisting of 160 reactions, 250 genes, and 140 metabolites.

We have included the root and stem models with this work as Supplemental Files 3 and 4

respectively.

Confidence scoring. Reaction confidence scores have been defined in a manner consistent with a

previously published protocol [26]. Additional information on confidence scoring of the p-

ath780 model can be found in Supplemental File 22.

Curation of all these tissue models. For all four models, we have balanced (both in terms of

elements and charge) all model reactions and have resolved thermodynamically infeasible cycles

by removing reactions, breaking composite reactions, and adding metabolic costs to transport

reactions. For all these tissue models, GPR links have been established through a largely

automated workflow utilizing the KEGG API for the majority of reactions using the code

included in Supplemental File 18. This is followed by having manually curated the GPR links

and/or inclusion rational of reactions with non-KEGG identifiers. The count of tissue model

reactions present in KEGG-defined pathways is shown in Figure 2(A), showing pathways

common to most all tissue models, and Figure 2(B), showing pathways common to seed and leaf

tissues, these figures have been created using code included in Supplemental Files 19, 20, and

21. The results of this automated workflow can be found in Supplemental File 5. Sources for

reactions included in leaf, root, seed, and stem models are shown in Figure 2(C-F), respectively

through confidence scoring (see confidence score section). Similarly, the confidence scores for

all reactions in the p-ath780 model have been reported in Figure 2(G).

Overview of the developed of the optimization-based framework of p-ath780. The models have

next been linked using well-known computational framework known for modeling microbial

communities [25]. An objective function for each of these models has then been defined,

specifying the maximization of the tissue-level biomass production rate followed by adding

constraints for simulating growth in light and dark conditions. Next, literature information

including embryo mass [30], initial tissue masses [38], growth stages [24], time points at which

growth stages occur [24], constraints to link tissue growth rates to appropriate tissue ratios,

transpiration [33,39], leaf surface area [28], usability of provided light [31,34,35], and defining

changes in tissue mass ratios [24,40] has been integrated into these models, which are typically

overlooked in most other SMs. In this work, we have decided to simulate arabidopsis biomass

across 61 days (1464 hours) of growth, as all plant seeds are dispersed by day 61, and after

which *in vivo* data on plant growth and mass is sparse. More specific details can be found in the

following sub-sections. The full optimization-based framework used in this work has been

provided in Supplemental File 6, and further requires Supplemental Files 7 through 16.

Generalized statement of the FBA-like optimization-based framework used. The optimization-

based framework used can be stated in general terms as follows. The FBA-based framework

which determines the optimal rates of flux through each reactions can be stated as follows (using

the default objective).

$$maximize \; Z = v_{leaf,biomass} + v_{root,biomass} + v_{stem,biomass} \qquad (1)$$

Subject to

$$10 \geq \sum_{j \in J} S_{ij} v_j \geq 0 \qquad\qquad \forall i \in I \qquad (2)$$

$$M_{leaf} v_{CO_2,in,leaf}$$

$$\geq M_{root} v_{CO_2,out,root} + M_{seed} v_{CO_2,out,seed} \qquad (3)$$

$$+ M_{stem} v_{CO_2,out,stem} \; (when \; light \; available)$$

$$M_{leaf} v_{O_2,out,leaf}$$

$$\geq M_{root} v_{O_2,in,root} + M_{seed} v_{O_2,in,seed} \tag{4}$$

$$+ M_{stem} v_{O_2,in,stem} \; (when \; light \; available)$$

$$v_{m,in,root} \leq M \qquad\qquad\qquad \forall m \in Micro \quad (5)$$

$$M_{root} v_{m,out,root} \leq M_{root} v_{m,in,root} \qquad\qquad \forall m \in Micro \quad (6)$$

$$M_{root} v_{m,out,root} = M_{stem} v_{m,in,stem} \qquad\qquad \forall m \in Micro \quad (7)$$

$$M_{stem} v_{m,out,stem} \leq M_{stem} v_{m,in,stem} \qquad\qquad \forall m \in Micro \quad (8)$$

$$M_{stem} v_{m,out,stem} = M_{leaf} v_{m,in,leaf} + M_{seed} v_{m,in,seed} \qquad \forall m \in Micro \quad (9)$$

$$M_{stem} v_{x,in,stem} = M_{leaf} v_{x,out,leaf} \qquad\qquad \forall x \in X \quad (10)$$

$$M_{stem} v_{x,out,stem} = M_{stem} v_{x,in,stem} \qquad\qquad \forall x \in X \quad (11)$$

$$M_{stem} v_{x,out,stem} = M_{seed} v_{x,in,seed} \qquad\qquad \forall x \in X \quad (12)$$

$$M_{stem} v_{sucrose,in,stem} = M_{leaf} v_{sucrose,out,leaf} \tag{13}$$

$$M_{stem} v_{sucrose,out,stem} \leq M_{stem} v_{sucrose,in,stem} \tag{14}$$

$$M_{stem} v_{sucrose,out,stem} = M_{root} v_{sucrose,in,root} + M_{seed} v_{sucrose,in,seed} \tag{15}$$

$$v_{starch,store,leaf} = s_{la} \sin\left(s_{lf}(n + s_{lx})\right) \tag{16}$$

$$v_{starch,store,stem} = s_{sa1} \sin\left(s_{sf1}(n + s_{sx1})\right) \tag{17}$$

$$v_{sucrose,store,stem} = s_{sa2} \sin\left(s_{sf2}(n + s_{sx2})\right) \tag{18}$$

$$v_{water,in,root} \leq M \tag{19}$$

$$M_{root} v_{water,out,root} = M_{stem} v_{water,in,stem} \tag{20}$$

$$M_{stem} v_{water,out,stem} = M_{leaf} v_{water,in,leaf} + M_{seed} v_{water,in,seed} \tag{21}$$

$$v_{water,in,leaf} = t_{leaf} \; (when \; light \; available) \tag{22}$$

$$v_{root,biomass} = \ln(\mu) + v_{leaf,biomass} \tag{23}$$

$$\mu = \frac{x_{root}M_{leaf}}{x_{leaf}M_{root}} \tag{24}$$

$$v_{seed,biomass} = \begin{cases} 0 & M_{seed} = 0, s = 0 \\ \ln(\theta) + v_{leaf,biomass} & M_{seed} \neq 0, s \neq 0 \\ -v_{leaf,biomass} & M_{seed} \neq 0, s = 0 \\ v_{leaf,biomass} & M_{seed} = 0, s \neq 0 \end{cases} \tag{25}$$

$$\mu\theta = \frac{x_{seed}M_{leaf}}{x_{leaf}M_{seed}} \tag{26}$$

$$v_{stem,biomass} = \ln(\lambda) + v_{leaf,biomass} \tag{27}$$

$$\lambda = \frac{x_{stem}M_{leaf}}{x_{leaf}M_{stem}} \tag{28}$$

$$v_{seed,biomass\ loss} = \begin{cases} -v_{seed,biomass} & n \geq n_{flowering}\ n \geq n_{seed\ loss} \\ 0 & n \leq n_{seed\ loss} \end{cases} \tag{29}$$

$$v_{scen} = \left(\frac{k_m + k_s}{24}\right) M_{tissue} \tag{30}$$

Where *I* is the set of metabolites; *J* is the set of reactions; *Micro* is the set of micronutrients

(phosphate, ammonium, and sulfate) and is a subset of *I*; *X* is the set of amino acids which are

synthesized in the leaf tissue and exported to other tissues; and *Y* is the set of sugars stored by

various plant tissues. In addition, *M* is defined as a very large number, t_{leaf} is the mass-

normalized transpiration rate from the leaf; s_{la}, s_{sa1}, and s_{sa2} are the amplitudes of the sine wave

modeling of starch storage in the leaf and stem and sucrose storage in the stem, respectively; s_{lf},

s_{sf1}, s_{sf2} are the frequencies of the sine wave modeling of starch storage in the leaf and stem and

sucrose storage in the stem, respectively; s_{lx}, s_{sx1}, and s_{sx2} are the x-intercept shifts of the sine

wave modeling of starch storage in the leaf and stem and sucrose storage in the stem,

respectively; *s* is the level of seeding of the model; x_{tissue} is the fraction of total plant mass

accounted for by that tissue; and M_{tissue} is the mass of the given tissue. The following subsections

will explain the constraints used in the FBA framework.

Defining model objective functions. For most analyses and results, the objective function of p-

ath780 has been to maximize the sum of the biomass production rates for all four tissues

according to the following equation (referred to as the default objective).

$$maximize\ z = v_{growth\ leaf} + v_{growth\ root} + v_{growth\ seed} + v_{growth\ stem} \tag{31}$$

Where z has been defined the objective function and $v_{growth\ tissue}$ is defined as the rate of

biomass production, in units of h^{-1}, of the tissue referenced. This objective function is

approximately equivalent to having maximized the growth rate (change in mass per unit time) of

the plant as a whole. This objective function has led to one major issue, namely how to avoid the

model producing only the metabolically "cheapest" tissue which could result in the maximum

objective value but is biologically unrealistic. This is addressed by equations (23) through (28)

and will be further discussed later.

It has been noted that the maximization of plant biomass has not been the only feasible objective

function for plant SM system, for instance one alternate objective function is the maximization

of plant photonic efficiency [15,16]. This objective has generally been framed as minimizing the

amount of light used by the plant system, given a required growth rate [15,16]. As it has been

assumed that the only (significant) photosynthetic tissue in the p-ath780 model is the leaf tissue,

only the objective with relation to the leaf tissue has been altered. As a result, the leaf tissue term

in equation (1) has been replaced with a photonic efficiency term in the following equation.

$$maximize\ z = -\alpha\left(v_{photon\ uptake}\right) + v_{growth\ root} + v_{growth\ seed} + v_{growth\ stem} \qquad (32)$$

Where α has been defined a correction factor to scale $v_{photon\ uptake}$ to be on the same order of

magnitude as the growth rates for each other tissue.

An alternative objective function has also been defined for the seed tissue. Specifically, as fatty

acids have been shown to one of the most prominent forms of carbon storage in the seed tissue

[38], the alternate objective function is the maximization of seed fatty acid stores. This has

resulted in seed objective function as follows:

$$maximize\ z = v_{growth\ root} + v_{growth\ root} + \beta \sum v_{FA\ sinks} + v_{growth\ stem} \qquad (33)$$

579   Where the new seed flux term is defined as the sum of fatty acid storing (sink) reactions and β

580   serves to reduce this term to be equal in order of magnitude to the other objectives. Similar to α,

581   β has been determined through trial and error. One additional objective function has been studied

582   which combines linear photonic efficiency and fatty acid storage:

$$maximize\ z = -\alpha\left(v_{photon\ uptake}\right) + v_{growth\ root} + \beta \sum v_{FA\ sinks} + v_{growth\ stem} \qquad (34)$$

Equation (33) has combined three separate objective types: linear photonic efficiency (leaf),

biomass production (root and stem), and fatty acid storage (seed). It should be noted that the

objective for the root and stem tissues are always to maximize biomass production. For more

details see Supplemental File 22.

Mass balance. In this model, the mass balance, equation (2), is allowed some flexibility for the

storage of metabolites in the plant tissue up to 10 mmol per gDW hour. This has been found to

be necessary in the design-build-test cycle so that all points would be feasible.

Enforcing net CO_2 consumption O_2 production. In equations (3) and (4), it is required that the net

effect plant metabolism is carbon fixation and oxygen production, since this is a well-known role

of plant systems (see Supplemental File 22).

Enforcing logical flow of micronutrients. In equations (5) through (9) the logical flow of

micronutrients is dictated. Equation (5) ensures that the uptake rate is bounded, equation (6)

ensures that the rate of each micronutrient exported by the root to the other tissues is less than or

equal to that uptaken by the root from the soil, allowing for the root to use a portion of the uptake

nutrients. Equation (7) ensures that all micronutrient exported by the root is uptaken by the stem,

and equation (8) is essentially the same as equation (6), but for the stem tissue. Finally, equation

(9) ensures that the micronutrients exported by the stem are all given to other tissues, specifically

leaf or seed. Both sides of the equation in equations (6) through (9) are multiplied by each tissues

mass to convert the units of the constraint from mmol per gDW hour to mmol per hour as each

tissue has a different mass value in gDW. For more details see Supplemental File 22.

Enforcing logical flow of amino acids. Similar to micronutrients, the logical flow of amino acids

has been defined explicitly via equations (10) to (12), as having been synthesized in the leaf

tissue and exported to seed tissue. This is because seed tissue has not been shown to produce all

needed amino acids in the MFA study consulted [27], and the root and stem models do not

require amino acids for biomass production in the defined biomass composition. Essentially,

these constraints ensure that all amino acids exported by the leaf are uptaken by the stem

[equation (6)]; that these amino acids are not stored in the stem [equation (7)]; and that all amino

acids are exported by the stem to the seed tissue. For more details see Supplemental File 22.

Enforcing logical flow of sucrose. As with amino acids, sucrose is modeled as being produced in

the leaf tissue and exported to other tissues. In contrast to amino acids, sucrose is necessary for

all tissue models. Therefore, equation (13) is analogous to equation (10), equation (14) allows

the stem to use the sucrose it receives from the leaf, unlike equation (11), and equation (15)

exports sucrose both to the seed and root, unlike equation (12). For more details see

Supplemental File 22.

Enforcing diurnal patterns of carbohydrate storage. Plants store carbohydrates in leaf and stem

tissues in the form of starch (leaf and stem) and sucrose (stem) in a pattern where the rates of

storage may be modeled by a sine wave with a period of 24 hours [31,41]. The calculations for

defining the necessary parameters, parameters s_{la}, s_{sa1}, s_{sa2}, s_{lf}, s_{sf1}, s_{sf2}, s_{lx}, s_{sx1}, and s_{sx2} in

equations (16) through (18), of the fit sine waves (see Supplemental Files 5 and 22).

Enforcing logical flow of water. The flow of water in the p-ath780 model is constrained as similar

to that of micronutrients and are defined in equations (19) through (22), but without the

equivalents of equations (6) and (8), and with the addition of a transpiration constraint. This

difference is because oxidative phosphorylation in tissues creates water. Hence, in tissues

without significant photosynthetic activity, water might be produced in this model, and the

largest usage to which plants put water, to maintain turgor pressure, is not modeled as SMs use

gDW as a basis of calculation, rather than fresh weight (see Supplemental File 22).

Defining the relationship between tissue growth rates. To avoid the aforementioned problem of

having p-ath780 produce only the "cheapest" biomass, the growth rates of all four tissues have

been linked by a series of constraints which ensure that they grow at rates which maintain the

desired tissue mass ratios. The rate of biomass production determined by a SM is the exponential

growth rate of the biological system being modeled [8]; therefore, plant mass can be defined as:

$$M_{tiss,n+1} = M_{tis,n} e^{v_{tis\ growth,n} t} \quad \forall tis \in [leaf, root, seed, stem] \tag{35}$$

646   Where M has been defined as the plant mass at time $n + 1$, $v_{plant\ growth,n}$ is defined as the rate

647   of plant growth at time n, and M_{n-1} is defined as the plant mass at time n. Further, the ratio of

648   the masses to two tissues can be defined with reference to a single tissue, such as leaf, in the

649   following manner:

$$M_{tiss,n} = \frac{x_{tis}}{x_{leaf}} M_{leaf,n} \quad \forall tis \in [leaf, root, seed, stem] \tag{36}$$

652   By having substituted the former equation into the latter and simplifying the result (see

653   Supplemental File 22), linear equations have been written to constrain biomass production rates

654   of root, seed, and stem tissues with respect to leaf tissue as follows:

$$v_{tiss,n} = \ln\left(\frac{x_{tis,n+1}M_{leaf,n}}{x_{leaf,n+1}M_{tis,n}}\right) + v_{leaf,n} \quad \forall tis \in [root, seed, stem] \tag{37}$$

This constraint we have added to the SM model, as equations (23) through (28) in order to ensure

that all tissues do have biomass production (or loss) and that it is in an amount which will result

in tissue masses in the correct proportions.

Ensuring non-productive loss of seed mass in silique shattering. A constraint has been found

necessary to enforce that metabolites associated with the loss of seed biomass, modeled by the

biomass production constraint having reversed flow, when seeds are being lost during silique

shattering (in the Silique Ripening Stage) and are not recycled into other parts of plant

metabolism. This constraint just does that by forcing recovered metabolites into the biomass loss

reaction of the seed tissue.

Defining model maintenance and senescence costs. An important consideration in any SM is

the definition of a maintenance cost, which is typically defined as ATP hydrolysis [26]. Biomass-

based maintenance and senescence costs have been defined as they have been suggested as more

accurate or applicable for plant systems [36,42], but have not yet been used in an SM. We have

defined maintenance and senescence costs as a biomass drain on each tissue scaled by tissue

mass in equation (30). A maintenance cost value of k_m=0.03 day^{-1} has been defined which is in

an order of magnitude typical for plant systems [42], and the same value has been defined for

plant senescence, k_s, as this parameter appears to generally be of the same order of magnitude

[36,42]. These rates are then converted into their per hour equivalent and scaled by tissue mass to

enforce these constraints. Only a single constraint has been defined for both phenomena as both

are biomass drains whose effect is additive. Literature evidence, including pictorial evidence of

plant phenotype at various growth stages, appears to suggest that the rate of plant senescence

increases drastically as the flowering production stage finishes and the Silique Ripening phases

begin (in literature, growth stage 0.65 to 9.70) [24]. Further, it appears that the plant no longer

maintains current mass, but allows tissues to die and desiccate [24]. This has been included in the

p-ath780 model in that plant senescence is increased by two orders of magnitude and plant

maintenance is set to zero following the end of the Flower Production stage. This results in a

growth curve in-line with *in vivo* evidence (see Table 1).

Other constraints enforced on the FBA-like optimization framework. There are other constraints

enforced on the optimization framework discussed above that are more difficult or cumbersome

to state mathematically and are therefore discussed here.

Defining the usage of seed stores by the seedling. A seedling's source of carbon is primarily its

reserves of stored carbohydrates, proteins, and lipids. Namely, it has been shown that seeds have

stores of approximately 0.425 μg of sucrose, 6 μg of fatty acids, and 6 μg of proteins (modeled

here as component amino acids) available [30]. As no information concerning the pattern of

usage of the seed storage has been found, it has been assumed that the stores are utilized at a

constant rate during the duration of the seed germination period and that all the storage is fully

consumed by 88.5 hours after germination, which has been defined the point at which the

cotyledons are fully open and leaf development intensifies [24]. The rate at which the seedling

should have uptaken the seed storage has been determined by identifying the moles (mmol) of

each major component of the seed storage and dividing by the time over which the seedling

consumes those. This has resulted in a mmol/h quantity. See Supplemental File 5 for this

calculation. This quantity has then been scaled by plant mass to result in a mmol/gDW□h

quantity, which is used to bound the uptake rates of seed store metabolites. As the leaf has

proven the most metabolically active tissue, it is assumed that the leaf tissue of an arabidopsis

seedling uptakes the stored fatty acids, amino acids, and carbohydrates which is provided for

seedling growth during the Seed Germination stage when the leaves have no access to light (see

Figure 1, Seed Germination).

Defining initial plant and tissue ratios. As the model advances plant and tissue masses with

respect to time, the establishment of initial mass for plant and tissues has become important in

this framework. Experimental evidence has shown that arabidopsis seeds have a fresh weight

(FW) of 25.3 μg and have only about 7% water content [30]. The embryo itself is assumed equal

to the seed mass less the mass of seed stores of sucrose (0.425 μg), Fatty Acids (6 μg), and

proteins (6 μg) [30]. Having assumed that the dry matter content ratio holds for the embryo as

well, this has left approximately 11.0 μg dry weight (DW) for the embryo. As information on the

ratio of tissue masses in arabidopsis has not been documented in literature, the general ratio for

herbaceous plants has been used as a starting point, namely 0.46:0.24:0.3 leaf:root:stem FW [32].

This ratio has been converted to DW ratio for stoichiometric modeling. Experimental data has

shown that the dry matter content of leaf tissue is 0.212 DW/FW, of root tissue is 0.170 DW/FW,

and of the stem tissue is 0.176 DW/FW [44]. Having converted the FW ratios to DW ratios has

given the ratio of 0.511:0.267:0.211 leaf:root:stem DW. While the dry matter content of an

embryonic arabidopsis is much higher than that of a mature plant (the source of the utilized dry

matter content ratios), this DW tissue ratio has non-the-less been assumed to be accurate for the

embryo due to lack of evidence to the contrary.

Defining stage times. Time points which define the transition between different stages of growth

have been taken from a single source of experimental evidence [24]. Stage transitions selected

include the transition to stage 0.70 (Seed Germination to Leaf Development transition in Figure

3), stage 6.00 (Leaf Development to Flower Production transition in Figure 3), and stage 8.00

(Flower Production to Silique Ripening transition in Figure 3). Not all lifecycle stage transitions

for which there is experimental evidence have been incorporated into this model. In some cases,

this has been due to a lack of metabolic relevance, such as the transition from stage 1.04 to stage

1.05 where the plant transitions from 4 rosette leaves to 5 rosette leaves that are greater than

1mm in length. This has not been important to the p-ath780 model as a ratio of plant mass to leaf

surface area ratio is used instead [33] (see Supplemental File 5). Others cannot be modeled by

the current framework tissues such as stage 5.10 which is when the first flower bud is visible

[24], as the current p-ath780 model has no flower bud tissue. The length of the seed ripening

stage is also determined by experimental evidence [24].

Defining the change in tissue mass ratios with growth stage. Using available literature evidence,

two endpoints for the plant tissue mass ratios have been defined when no seeds are present and

all seeds are produced [24,38]. The transition between these states are assumed to be linear with

respect to a parameter called seeding, defined above as *s*. These relationships are then modeled

as:

$$x_{leaf} = c_{leaf} * seeding + x_{leaf,0} \tag{38}$$

$$x_{root} = c_{root} * seeding + x_{root,0} \tag{39}$$

$$x_{seed} = c_{seed} * seeding + x_{seed,0} \tag{40}$$

$$x_{stem} = c_{stem} * seeding + x_{stem,0} \tag{41}$$

$$c_{leaf} = -0.2514; c_{root} = -0.02862; c_{seed} = 0.2030; c_{stem} = 0.07698$$

$$x_{leaf,0} = 0.511; x_{root,0} = 0.267; x_{seed,0} = 0; x_{stem,0} = 0.211$$

Where x_{tissue} has been defined as the tissue mass fraction with respect to the total mass of the

plant, c_{tissue} is defined as the change in tissue mass fraction with respect to seeding, and x_{tissue}

is defined as the initial mass fraction of each tissue. The gain in the seeding parameter has been

assumed to be linear with time and is fit to experimental time point describing the fraction of

flowers produced [24] (see Supplemental Files 5 and 22).

Defining the availability of light. The amount of light available to the model to use for

photosynthesis has been defined initially by literature sources used for other constraints [31], and

scaled by the transmittance of that light source (fluorescent lights) [34] and the absorbance of

arabidopsis leaves [34] and surface area to plant mass of arabidopsis leaves [33] to define the

amount of light usable by the plant system, which has been approximately estimated to be 4.00

mmol/gDW plant·h. This value has been shown to be 21.50% of the total photons output by the

fluorescent light (see Supplemental Files 5 and 22).

Defining the FVA for the p-ath780 model. A Flux Variability Analysis (FVA) model has been

defined for growth both in light and dark growth. All flux bounds and constraints are the same

and the FBA models, but the objective function is defined as:

$$maximize/minimize\ z = \gamma_j v_j \qquad\qquad (42)$$

Where the FVA model solution has been iterated for each reaction j, and γ_j has been valued at 1

for the current reaction whose maximum and minimum are to be investigated and 0 for all others

and is stepped through first maximizing and then minimizing each reaction. Due to restrictions of

the time allowed for model solutions, nine points has been selected at which to perform FVA.

These points are 70 hours after germination (HAG, seed germination stage, dark), 408 HAG (leaf

development stage, light), 576 HAG (leaf development stage, light), 590 HAG (leaf development

stage, dark), 800 HAG (flower production stage, light), 810 HAG (flower production stage,

dark), 1156 HAG (flower production to silique ripening transition), 1200 HAG (silique ripening

stage, light), and 1220 HAG (silique ripening stage, dark).

Defining the mass step between time points. Using the biomass production rates calculated by

the FBA-like optimization framework, a Constrained Non-linear System (CNS) of equations can

be defined to advance the plant mass by treating the growth rates as constants. This system of

equations has been derived from the basic principles of FBA (e.g. that growth rates are

exponential rates of growth) through a sequence of simplifications and assumptions which can be

found in Supplemental File 22, and therefore will not be elaborated on here. The end result is

shown below for a given time point *t*.

$$\frac{dM_{plant}}{dt}\bigg|_{n\to n+1/3} = \frac{e^{v_{leaf,bio}}M_{leaf,n}}{x_{leaf,n}}\left[x_{leaf,n}v_{leaf,bio,n} + \frac{d}{dt}\left(v_{leaf,bio,n}\right) + \xi(t)\right] \tag{43}$$

$$\xi(t) = x_{root}\psi(t) + x_{seed}\pi(t) + x_{stem}\kappa(t) + x_{leaf}\left(\psi(t)\zeta(t) + \pi(t)\rho(t) + \kappa(t)\iota(t)\right)\frac{d}{dt}(s)$$

$$+ x_{root}\frac{d}{dt}\left(\ln(\mu(t))\right) + x_{seed}\frac{d}{dt}\left(\ln(\theta(t))\right) + x_{stem}\frac{d}{dt}\left(\ln(\lambda(t))\right) \tag{44}$$

$$\psi(t) = \ln(\mu(t)) + v_{leaf,bio,n} \tag{45}$$

$$\pi(t) = \ln(\theta(t)) + v_{leaf,bio,n} \tag{46}$$

$$\kappa(t) = \ln(\lambda(t)) + v_{leaf,bio,n} \tag{47}$$

$$\zeta = \frac{c_{root}\left(c_{leaf}s + x_{leaf,n}\right) - c_{leaf}\left(c_{root}s + x_{root,n}\right)}{c_{leaf}^2 s^2 + 2c_{leaf}sx_{leaf,n} + x_{leaf,n}^2} \tag{48}$$

$$\rho = \frac{c_{seed}\left(c_{leaf}s + x_{leaf,n}\right) - c_{leaf}(c_{seed}s)}{c_{leaf}^2 s^2 + 2c_{leaf}sx_{leaf,n} + x_{leaf,n}^2} \tag{49}$$

$$\iota = \frac{c_{stem}\left(c_{leaf}s + x_{leaf,n}\right) - c_{leaf}\left(c_{stem}s + x_{stem,n}\right)}{c_{leaf}^2 s^2 + 2c_{leaf}sx_{leaf,n} + x_{leaf,n}^2} \tag{50}$$

$$\frac{d}{dt}\left(v_{leaf\ growth,n}\right) = \frac{3v_{leaf\ growth,n} - 4v_{leaf\ growth,n-1/3} + v_{leaf\ growth,n-2/3}}{2\left[\frac{1}{3}h\right]} \tag{51}$$

Where, μ, θ, λ are parameters defined in equations (24), (26), and (28), The above system of nine

equations has nine corresponding variables: the mass step [LHS of equation (43)], ξ, ψ, π, ζ, ρ, ι,

and the time derivative of the leaf growth rate [LHS of equation (53)].

Equation (51), as shown above, comes from a backwards finite difference formula of error order

h^2. For the purposes of increased model accuracy and stability, the FBA-like framework is solved

every $^1/_3$ of an hour to more accurately calculate the mass step at each hour (error is

approximately one ninth of that when using full hour values as estimates). As equation (51) is an

estimate (rather than an exact equation), should CNS not find a feasible solution, it is further

relaxed using a tolerance parameter (*tol*) to a pair of inequalities:

$$\left| \frac{d}{dt}\left(v_{leaf\ growth,n}\right) \right|$$

$$\geq (1 - tol) \left| \frac{3v_{leaf\ growth,n} - 4v_{leaf\ growth,n-1/3} + v_{leaf\ growth,n-2/3}}{2\left[\frac{1}{3}h\right]} \right| \quad (52)$$

$$+ O\left(\left[\frac{1}{3}h\right]^2\right)$$

$$\left| \frac{d}{dt}\left(v_{leaf\ growth,n}\right) \right|$$

$$\leq (1 + tol) \left| \frac{3v_{leaf\ growth,n} - 4v_{leaf\ growth,n-\frac{1}{3}} + v_{leaf\ growth,n-\frac{2}{3}}}{2\left[\frac{1}{3}h\right]} \right| \quad (53)$$

$$+ O\left(\left[\frac{1}{3}h\right]^2\right)$$

Where, *tol* begins at 0.00 and increases by 0.10 for each iteration if a solution is not found. This

results in an mixed-integer non-linear programming (MINLP) problem (8 equality constraints

and 9 variables), and the BARON solver has been used to attempt to solve the model. In all cases

where a solution has not been found via a CNS solver, a solution has been found using the

MINLP solver at *tol* = 0.10. The above set of equations [either equations (43) through (51) as a

CNS problem or (43) through (50), (52), and (53)] is solved three times to make estimates of the

803   LHS of equation (42) usable in Heunn's rule for explicit third-order Runge-Kutta method. For

804   why this method has been used (see Supplemental File 22).

$$\left.\frac{dM_{plant}}{dt}\right|_{n\rightarrow n+1/3} = k_1 \tag{54}$$

$$\left.\frac{dM_{plant}}{dt}\right|_{n1/3\rightarrow n+2/3} = k_2 \tag{55}$$

$$\left.\frac{dM_{plant}}{dt}\right|_{n+2/3\rightarrow n+1} = k_3 \tag{56}$$

807   After each partial step, the plant and tissues masses are updated for the next solution. These mass

808   step estimates are then combined using Heunn's rule for explicit third-order Runge-Kutta

809   method, where the new mass is calculated as follows.

$$M_{plant,n+1} = M_{plant,n} + h\left(\frac{1}{4}k_1 + \frac{3}{4}k_3\right) \tag{57}$$

812   And the mass of each individual tissue is then updated as follows:

$$M_{leaf,n+1} = x_{leaf,n+1}M_{plant,n+1} \tag{58}$$

$$M_{root,n+1} = x_{root,n+1}M_{plant,n+1} \tag{59}$$

$$M_{seed,n+1} = x_{seed,n+1}M_{plant,n+1} \tag{60}$$

$$M_{stem,n+1} = x_{stem,n+1}M_{plant,n+1} \tag{61}$$

Why p-ath780 updates overall plant mass rather than solving the above problem for each

individual tissue is discussed in Supplemental File 22.

Software platforms used. See Supplemental file 23 for which programming language various

supplemental files utilize. For Python code, version 3.3 is used; for Perl, version 5.26 for

Supplemental Files 18 through 21 and Strawberry Perl version 5.24.0.1 is used for Supplemental

File 8; GAMS code utilizes version 24.7.4. All GAMS and Python code, in addition code

included in Supplemental File 8 is run using the Holland Computing Center at the University of

Nebraska, Lincoln. Supplemental Files 18 through 21 utilize the additional module the LWP (the

world-wide web library for Perl) module 6.39, and have been run on a windows desktop

computer.

Code availability. The authors declare that the code supporting the findings of this study is

available within the article's Supplementary Information files.

Abbreviations used. For the convenience of our readers, a list of abbreviations used is given

below:

GPR: Gene-Protein-Reaction

SM: Stoichiometric Model

FBA: Flux Balance Analysis

FVA: Flux Variability Analysis

LP: Linear Problem

CNS: Constrained Non-linear System

838 MINLP: Mixed Integer Non-Linear Problem

839 arabidopsis: *Arabidopsis thaliana*

840 LHS: Left-Hand Side

841 wrt: with respect to

842 gDW: grams Dry Weight

843 DW: Dry Weight

844 gFW: grams Fresh Weight

845 FW: Fresh Weight

846 MFA: Metabolic Flux Analysis

847 KEGG: Kyoto Encyclopedia of Genes and Genomes

848 DAG: Days After Germination

849 HAG: Hours After Germination

851 **Acknowledgement.** This work has been completed utilizing the Holland Computing Center of

852 the University of Nebraska, which receives support from the Nebraska Research Initiative. The

853 authors gratefully acknowledge funding from UNL Faculty Startup Grant 21-1106-4038.

855 **Author contributions.** Experiments have been conceived by R.S. and W.L.S. W.L.S. performed

856 the experiments and analyzed the data. R.S. and W.L.S. contributed analysis tools. R.S. and

857 W.L.S. wrote the manuscript.

Supplementary information is provided. To help readers navigate the extensive set of included

data and replicate this study, Supplemental File 23 provides an overview of the included

supplemental files and lays out the file structure to use in conjunction with the p-ath780 model.

Conflict of interest: The authors declare no conflicts of interest.

References

1. Beyer P, Al-Babili S, Ye X, Lucca P, Schaub P, Welsch R, et al. Golden rice: introducing the

β carotene biosynthesis pathway into rice endosperm by genetic engineering to defeat

vitamin A deficiency. J Nutr. 2002;132:506S-510S.

2. Hall RD, Brouwer ID, Fitzgerald MA. Plant metabolomics and its potential application for

human nutrition. Physiol Plant. 2008;132(2):162-175.

3. Gonzali S, Mazzucato A, Perata P. Purple as a tomato: towards high anthocyanin tomatoes.

Trends Plant Sci. 2009;14(5):237-241.

4. Paddon CJ, Keasling Semi-synthetic artemisinin: a model for the use of synthetic biology in

pharmaceutical development. Nat Rev Microbiol. 2014;12(5):355–367.

5. Hilder VA, Boulter D. Genetic engineering of crop plants for insect resistance - a critical

review. Crop Prot. 1999;18(3):177–191.

6. Chen THH, Murata N. Enhancement of tolerance of abiotic stress by metabolic engineering

of betaines and other compatible solutes. Curr Opin Plant Biol. 2002;5(3):250–257.

7. Srinivasan S, Cluett WR, Mahadevan R. Constructing kinetic models of metabolism at

genome-scales: A review. Biotechnol J. 2015;1359:1345–1359.

8. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nat Biotechnol.

2010;28(3):245–248.

9. Burgard AP, Pharkya P, Maranas CD. OptKnock: a bilevel programming framework for

identifying gene knockout strategies for microbial strain optimization. Biotechnol Bioeng.

2003;84(6):647–657.

10. Ranganathan S, Suthers PF, Maranas CD. OptForce: An optimization procedure for

identifying all genetic manipulations leading to targeted overproductions. *PLoS Comput Biol*

2010;6(4):1-11.

11. Mahadevan R, Edwards JS, Francis DJ. Dynamic flux balance analysis of diaxic growth in

Escherichia coli. Biophys J. 2002;83(3):1331–1340.

12. Saha R, Liu D, Hoynes-O'Connor A, Liberton M, Yu J, Bhattacharyya-Pakrasi M, et al.

Diurnal regulation of cellular processes in the Cyanobacterium synechocystis sp. strain PCC

6803: insights from transcriptomic, fluxomic and physiological analyses. MBio. 2016;7(3):1–

14.

13. Ng CY, Jung M, Lee J, Oh M-K. Production of 2,3-butanediol in Saccharomyces cerevisiae

by in silico aided metabolic engineering. Microb Cell Fact. 2012;11(68):1-14.

14. Poolman MG, Miguet L, Sweetlove LJ, Fell DA. A genome-scale metabolic model of

Arabidopsis and some of its properties. Plant Physiol. 2009;151(3):1570–81.

15. Gomes de Oliveira Dal'Molin C, Quek LE, Palfreyman RW, Brumbley SM, Nielsen LK.

AraGEM, a genome-scale reconstruction of the primary metabolic network in Arabidopsis.

Plant Physiol. 2010;152:579–589.

16. Gomes de Oliveira Dal'Molin C, Quek LE, Saa PA, Nielsen LK. A multi-tissue genome-

scale metabolic modeling framework for the analysis of whole plant systems. Front Plant Sci.

2015;6(4):1–12.

17. Grafahrend-Belau E, Junker A, Eschenröder A, Müller J, Schreiber F, Junker B. Multiscale

metabolic modeling: dynamic flux balance analysis on a whole-plant scale. Plant Physiol.

2013;163(2):637–647.

18. Saha R, Suthers PF, Maranas CD. Zea mays iRS1563: a comprehensive genome-scale

metabolic reconstruction of maize metabolism. PLoS One. 2011;6(7):1-12.

19. Gomes de Oliveira Dal'Molin C, Quek LE, Palfreyman RW, Brumbley SM, Nielsen LK.

C4GEM, a genome-scale metabolic model to study C4 plant metabolism. Plant Physiol.

2010;154(4):1871–1885.

20. Poolman MG, Kundu S, Shaw R, Fell DA. Responses to light intensity in a genome-scale

model of rice metabolism. Plant Physiol. 2013;162(2):1060–1072.

21. Mintz-Oron S, Meir S, Malitsky S, Ruppin E, Ahoroni A, Shlomi T. Reconstruction of

Arabidopsis metabolic network models accounting for subcellular compartmentalization and

tissue-specificity. Proc Natl Acad Sci. 2012;109(1):339–344.

22. Simons M, Saha R, Guillard L, Clément G, Armengaud P, Cañas R, et al. Nitrogen-use

efficiency in maize (Zea mays L.): from "omics" studies to metabolic modelling. J Exp Bot.

2014;65(19):5657–5671.

23. Grafahrend-Belau E, Schreiber F, Koschutzki D, Junker BH. Flux balance analysis of barley

seeds: a computational approach to study systemic properties of central metabolism. Plant

Physiol. 2009;149(1):585–598.

24. Boyes DC, Zayed AM, Ascenzi R, McCaskill AJ, Hoffman NE, Davis KR, et al. Growth

stage-based phenotypic analysis of Arabidopsis: a model for high throughput functional

genomics in plants. Plant Cell. 2001;13:1499–1510.

25. Zomorrodi AR, Maranas CD. OptCom: A multi-level optimization framework for the

metabolic modeling and analysis of microbial communities. *PLoS Comput Biol* 2012;8(2):1–

13.

26. Thiele I, Palsson BØ. A protocol for generating a high-quality genome-scale metabolic

reconstruction. Nat Protoc. 2010;5(1):93–121.

27. Lonien J, Schwender J. Analysis of metabolic flux phenotypes for two Arabidopsis mutants with severe impairment in seed storage lipid synthesis. Plant Physiol. 2009;151(3):1617–1634.

28. Johnson JMF, Barbour NW, Weyers SL. Chemical composition of crop biomass impacts its decomposition. Soil Sci Soc Am J. 2007;71(1):155-162.

29. Thiele I, Gudmundsson S. Computationally efficient flux variability analysis. BMC Bioninformatics. 2010;11(489):1-3.

30. Tomaz T, Bagard M, Pracharoenwattana I, Lindén P, Lee CP, Carroll AJ, et al. Mitochondrial malate dehydrogenase lowers leaf respiration and alters photorespiration and plant growth in Arabidopsis. Plant Physiol. 2010;154(3):1143–1157.

31. Hendrik Poorte A, Nagel O. The role of biomass allocation in the growth response of plants to different levels of light, CO2, nutrients and water: a quantitative review. Aust J Plant Physiol. 2000;27(189):595–607.

32. Poorter H, Nagel O. The role of biomass allocations in the growth respose of plants to different levels of light, CO2, nutrients and water: a quantitative review. Aust J Plant Physiol. 2000;27:595-607.

33. Clauss MJ, Aarssen LW. Phenotypic placity of size-fecundity relationships in Arabidopsis thaliana. J Ecol. 1994;82(3):447–455.

34. Bläsing OE, Gibon Y, Günther M, Höhne M, Morcuende R, Osuna D, et al. Sugars and Circadian Regulation Make Major Contributions to the Global Regulation of Diurnal Gene Expression in Arabidopsis. Plant Cell. 2007;17:3257–3281.

35. Baleja R, Šumpich J, Bos P, Helštýnová B, Sokanský K, Novák T. Comparison of LED properties, compact fluorescent bulbs and bulbs in residential areas. Proceedings of the 16th

International Scientific Conference on Electrical Power Engengineering; 2015 May 20-22;

Kouty nad Desnou, Czech Republic: IEEE, 2015.

36. Cannell MGR, Thornley JHM. Modelling the components of plant respiration: representation

and realism. Ann Bot. 1999;85:55–67

37. Cannell MGR, Thornley JHM. Modelling the components of plant respiration: some guiding

principles. Ann Bot. 1999:85:45–54.

38. Baud S, Boutin J, Miquel M, Lepiniec L, Rochat C. An integrated overview of seed

development in Arabidopsis thaliana ecotype WS. *Plant Physiol Biochem* 2002;40:151–160.

39. Li B, Suzuki JI, Hara T. Latitudinal variation in plant size and relative growth rate in

Arabidopsis thaliana. Oecologia. 1998;115:293–301.

40. Solovchenko AE, Merzlyak MN. Screening of visible and UV radiation as a photoprotective

mechanism in plants. Russ J Plant Physiol. 2008;55(6):719–737.

41. Yazdanbakhsh N, Sulpice R, Graf A, Stitt M, Fisahn J. Circadian control of root elongation

and C partitioning in Arabidopsis thaliana. Plant, Cell Environ. 2011;34(6):877–894.

42. Amthor JS. The role of maintenance respiration in plant growth. Plant, Cell Environ.

1984;7(8):561–569.

43. McClung CR. Plant circadian rhythms. The Plant Cell. 2006;18:792-803.

44. Oakenfull RJ, Davis SJ. Shining a light on the Arabidopsis circandian clock. Plant Cell

Envirn. 2017;40:2571-2585.

Supporting Information Captions

Supplemental_File_1.txt: This file is a text file (extension ".txt") which contains the seed tissue

model of p-ath780. This file is referenced as "p-ath780Seed.txt" by other supplemental files,

particularly code, and this correct name should replace the default name for attached code to run

properly.

Supplemental_File_2.txt: This file is a text file which contains the leaf tissue model of p-

ath780. This file is referenced as "p-ath780Leaf.txt" by other supplemental files, particularly

code, and this correct name should replace the default name for attached code to run properly.

Supplemental_File_3.txt: This file is a text file which contains the root tissue model of p-

ath780. This file is referenced as "p-ath780Root.txt" by other supplemental files, particularly

code, and this correct name should replace the default name for attached code to run properly.

Supplemental_File_4.txt: This file is a text file which contains the stem tissue model of p-

ath780. This file is referenced as "p-ath780Stem.txt" by other supplemental files, particularly

code, and this correct name should replace the default name for attached code to run properly.

Supplemental_File_5.xlsx: This file is a Microsoft Excel file which store a wide variety of

information concerning the p-ath780 model. This include the manually-curated GPR results for

each tissue model, the calculations pertaining to the determination of the biomass equation for

each tissue model, calculations for various parameters used in the p-ath780 model to incorporate

literature data, and calculations pertaining to the diurnal storage and uptake of carbohydrates.

Supplemental_File_6.txt: This file is the GAMS code for the p-ath780 model itself. It is

generally named "p-ath780.gms".

1001

1002   **Supplemental_File_7.txt:** This is an executable Python code which takes the input of a model

1003   (such as Supplemental Files 1 through 4) and outputs a number of files which can be read by

1004   GAMs code. Generally, this code is named "convert.py". This code requires slight modifications

1005   depending on which model file is to be converted (see in-code comments).

1006

1007   **Supplemental_File_8.txt:** This is an executable Perl code which takes the results of converting

1008   each tissue model file using Supplemental File 11 and creates some of the necessary inputs for

1009   the p-ath780 GAMS code. This is generally referenced as "makeGrowthInputs.pl".

1010

1011   **Supplemental_File_9.txt:** This is a text file which contains a list of the names of parameters

1012   used to defined p-ath780 model growth. This is referred to by other files as

1013   "growthSpecsNames.txt".

1014

1015   **Supplemental_File_10.txt:** This is a text file which contains the actual specifications used for

1016   growing by the p-ath780 model. This is referred to by other files as "growthSpecs.txt",

1017   importantly it is referred to as this by Supplemental File 6.

1018

1019   **Supplemental_File_11.txt:** This is a text file containing the list of time points to iterate over for

1020   each day, e.g. this contains each hour of the day, beginning at 0 and ending at 23. This file is

1021   referenced by others as "timepointsH.txt", importantly it is referred to as this by Supplemental

1022   File 6.

1023

1024   **Supplemental_File_12.txt:** This is a text file containing a list of days to solve the model over, in

1025   this case from day 0 to day 61. This file is referred to by others as "timepoints.txt", importantly it

1026   is referred to as this by Supplemental File 6.

1027

1028   **Supplemental_File_13.txt:** This is a text file containing a list of data labels for much of the data

1029   saved at each time point (combination of day and hour) and reported on in the troubleshooting

1030   file. This file is referred to by others as "timeData.txt", importantly it is referred to as this by

1031   Supplemental File 6.

1032

1033   **Supplemental_File_14.txt:** This is a text file which lists the time at which the sun rises (or light

1034   is made available) each day. At present, light is made available at a default time of 0. This file is

1035   referred to by others as "sunrise.txt", importantly it is referred to as this by Supplemental File 6.

1036

1037   **Supplemental_File_15.txt:** This is a text file which lists the time at which the sun sets (or light

1038   is no longer made available) each day. At present, light is made available at a default time of 0.

1039   This file is referred to by others as "sunset.txt", importantly it is referred to as this by

1040   Supplemental File 6.

1041

1042   **Supplemental_File_16.txt:** This file basically converts the set time of day to a parameter of

1043   equal value. Necessary because mathematical operations cannot be performed on sets. This file is

1044   referred to by other files as "timeofday.txt", importantly it is referred to as this by Supplemental

1045   File 6.

1046

Supplemental_File_17.xlsx: This is a Microsoft Excel file which contains the results of the p-

ath780 model for various alternative objective functions. The sheet tabs indicate which

alternative objective function the data corresponds to. The key is as follows:

lpe_g_g_g_X: linear photonic efficiency objective for the leaf, growth objective for other tissues.

The number which replaces "X" indicates the run number.

g_g_g_g: Growth objective for all tissues.

nlpe_X: Non-linear photonic efficiency objective, the number which replaces the "X" denotes

the run number.

g_g_fa_g_X: Fatty acid storage objective for the seed tissue, growth objective for all other, the

number which replaces the "X" denotes the run number.

1057

1058   **Supplemental_File_18.txt:** This is an executable Perl code file which is used to automatically

1059   curate the Gene-Protein-Reaction (GPR) links for all tissue models using the KEGG API

1060   (advanced programming interface, rest.kegg.jp). Inside the documentation of the code is the

1061   instructions for adapting it to investigate the GPR links for each tissue. Generally, this file is

1062   named "RxnstoGenes.pl". This file requires the LWP Perl package.

1063

1064   **Supplemental_File_19.tex:** This file is a comma separated values file. This file contains a list of

1065   73 KEGG pathways for which to get the list of associated reactions. This file is referred to by

1066   code as ".csv", and must have the proper name for the code to function properly. Generally

1067   referred to as "PathRxns.csv".

1068

1069   **Supplemental_File_20.txt:** This file is an executable Perl code file (extension of ".pl") which

1070   automatically generates the lists of reactions associated with various KEGG pathways. The

1071   KEGG pathways used are listed in Supplemental File 19. This file is generally named

1072   "PathGetRxnsComps.pl".

1073

1074   **Supplemental_File_21.txt:** This file is an executable Perl code file which is used to

1075   automatically read the files created from Supplemental Files 7 to give counts of how many

1076   reactions a model has which belong to each of the pathways indicated by Supplemental File 7.

1077   Generally, this file is named "ModelPathComp.pl".

1078

1079   **Supplemental_File_22.docx:** This file is a Microsoft Word file which contains all the

1080   calculations simplifications, and rational used for the determination of the function for whole-

1081   plant mass step with respect to time.

1082

1083   **Supplemental_File_23.docx:** A Microsoft Word file designed to help navigate other files

1084   provided as well as to outline the general file structure and barebones workflow used with the p-

1085   ath780 model to make model implementation easier.

1086

Table Legends

Table 1: This table compares the critical mass-based metrics which have been used for *in silico* to

in vivo data comparison across different objective functions for the p-ath780 model. The final

two columns are for metrics which have been targets of change when applying different

objective functions. The green row is the *in vivo* data ranges which have served as targets for *in*

silico model behavior. The blue row is the behavior of the p-ath780 model with the usual

objective function used in all other analyses (maximizing biomass production of all tissues). The

grey rows are alternate objective functions we have explored. The photonic efficiency objective

functions for the leaf tissue have caused lower fractions of available light used, while the fatty

acid production objective for the seed tissue has caused greater diversion of carbon resources

toward fatty acid production and storage. All alternative objective functions have resulted in

lower   mass   production   *in*   *silico*.

Tissue-Level Objectives			Total Plant Mass (mg)			Relative Growth Rate (RGR)	Peak Tissue Mass (mg)			Resource utilization or production (median %)	
Leaf	Seed	Root & Stem	17 DAG	24 DAG	31 DAG	Whole-plant RGR (day^{-1})	Leaf	Stem	Seed	Of usable light used in light	Of carbon uptaken by seed to fatty acid storage
In vivo data	*In vivo* data	*In vivo* data	0.5-2	2-8	10-30	$0.21 - 0.25$	163.7 ± 52.0	188 ± 39.3	127.9 ± 52.7	?	?
Biomass production	Biomass production	Biomass production	0.554	3.74	25.2	0.246	177	189	130.	100%	0%
Biomass production	Fatty acid Storage $(\beta = 1E - 6)$	Biomass production	0.554	3.74	25.2	0.246	177	189	130.	100%	61.6%
Biomass production	Fatty acid Storage $(\beta = 2E - 2)$	Biomass production	0.554	3.74	25.2	0.246	94.1	100.	68.9	100%	61.7%
Linear	Biomass	Biomass	0.554	3.74	25.2	0.246	177	189	130.	100%	0%

photonic efficiency $(\alpha = 25)$	production	production									
Linear photonic efficiency $(\alpha = 50)$	Biomass production	Biomass production	0.554	3.74	25.2	0.246	177	189	130.	100%	0%
Linear photonic efficiency $(\alpha = 50)$	Fatty acid Storage $(\beta = 1E - 2)$	Biomass production	0.554	3.74	25.2	0.246	133	142	97.7	100%	61.7%

1099

Core Network Model Construction Workflow

① Use 'omics data to determine reactions present in the organism.

Transcriptome
Genome
Proteome
Reactome A + B → C

Seed Core Model
- Metabolite storage
- Biomass production
- Respiration
- Fatty acid metabolism

First reconstructed

Leaf Core Model
- Photosynthesis
- Respiration
- Gas exchange
- Biomass Production
- Amino acid synthesis
- Fatty acid metabolism

Second reconstructed

Stem Core Model
- Metabolite transport
- Biomass production
- Essential metabolism

Third reconstructed

Root Core Model
- Nutrient uptake
- Biomass production
- Essential metabolism

Fourth reconstructed

② Develop core metabolic network models for each selected tissue.

Seed Germination
Leaf Development
Flower Production
Silique Ripening

Sample Growth Parameters:
- Biomass yeild
- Senescense rate
- Maintenance cost
- Tissue mass ratios
- Seeding gain rate
- Seeding loss rate
- Seeding loss rate
- Others (see File S10)

Program workflow:
- Inputs
- Decision branches
- Models outputs

Fig. 4

③ Use literature to identify, major growth stages, tissue interactions, and growth parameters. Design program workflow.

Fig. 1

④ Use the model and optimization framework to create the series of snapshots for lifecycle metabolism.

p-ath780
Experiment

Comparisons:
- Tissue mass
- Reaction flux ranges (FVA)
- Plant mass
- Growth trends

⑤ Compare metabolic lifecylce results to experimental literature data.