1   **Measuring genome sizes using read-depth, k-mers, and flow**

2 **cytometry: methodological comparisons in beetles (Coleoptera)**

3 James M. Pflug,*,[1] Valerie Renee Holmes,[†] Crystal Burrus,[†] J. Spencer Johnston,[†] and David R.
4 Maddison*
5
6 *Department of Integrative Biology, Oregon State University, Corvallis, OR 97331, USA
7 †Department of Entomology, Texas A&M University, College Station, TX 77843, USA
8

[1] Corresponding author:
3029 Cordley Hall, Department of Integrative Biology, Oregon State University, Corvallis, OR 97331 USA.
  E-mail: pflugja@gmail.com

Running Title: Comparing Genome Size Methods
Corresponding author: James Pflug
3029 Cordley Hall, Department of Integrative Biology, Oregon State University, Corvallis, OR
97331 USA.
1 (417) 529-7851
E-mail: pflugja@gmail.com
15

Keywords:
Genome size
Coleoptera
k-mer
Flow cytometry
Read mapping

ABSTRACT

Measuring genome size across different species can yield important insights into evolution of the genome and allow for more informed decisions when designing next-generation genomic sequencing projects. New techniques for estimating genome size using shallow genomic sequence data have emerged which have the potential to augment our knowledge of genome sizes, yet these methods have only been used in a limited number of empirical studies. In this project, we compare estimation methods using next-generation sequencing (k-mer methods and average read depth of single-copy genes) to measurements from flow cytometry, the gold standard for genome size measures, using ground beetles (Carabidae) and other members of the beetle suborder Adephaga as our test system. We also present a new protocol for using read-depth of single-copy genes to estimate genome size. Additionally, we report flow cytometry measurements for five previously unmeasured carabid species, as well as 21 new draft genomes and six new draft transcriptomes across eight species of adephagan beetles. No single sequence-based method performed well on all species, and all tended to underestimate the genome sizes, although only slightly in most samples. For one species, *Bembidion haplogonum*, most sequence-based methods yielded estimates half the size suggested by flow cytometry. This discrepancy for k-mer methods can be explained by a large number of repetitive sequences, but we have no explanation for why read-depth methods yielded results that were also strikingly low.

INTRODUCTION

The advent of modern genomics and the resulting deluge of data from next generation

sequencing (NGS) has been a tremendous boon to the biological sciences. In spite of this, many

foundational questions about genomes have remained largely unanswered. One such question is

why genomes vary so much in size: there is an over 3,000-fold difference between the smallest

and largest genomes in animals (Gregory, 2001). Revealing the myriad evolutionary causes

behind this variation has proven to be a difficult and enduring challenge (Cavalier-Smith 1978,

Elliott and Gregory 2015). One limitation to understanding genome size evolution is the relative

lack of knowledge of genome sizes in some of the larger clades of life, such as the arthropods

(Hanrahan and Johnston 2011).

Knowledge of genome size is desirable for multiple reasons. It can be important to

understand phenomena such as whole genome duplication and polyploidy (Allen 1983, Némorin

et al. 2013), genome reduction driven by changes in selective pressure (Johnston et al. 2004), or

proliferation of non-coding DNA sequence (Gregory 2005). Genome size has also been observed

to correlate with a variety of developmental factors, such as egg size and cell division rate

(Gregory 2001). Knowledge about genome size can also be valuable in species delimitation.

Differences sufficient to reproductively isolate a population into separate species may be difficult

to distinguish using traditional morphological or DNA sequence data; however, such differences

may be more apparent once genome sizes are taken into consideration alongside other evidence

(Gregory 2005, Leong-S□kornic□ková 2007).

Traditional methods for determining genome size, such as Feulgen densitometry and flow

cytometry, involve staining cells with a DNA-specific dye and comparing the results to stained

cells from a standard reference of a known genome size. These methods are well-tested and

generally reliable (Chen et al. 2015, Hanrahan and Johnston 2011). Flow cytometry in particular

is considered the "gold standard" for estimating genome size (Mounsey et al. 2012). However,

these techniques rely on live, fixed, or frozen tissues with largely intact cells, effectively limiting

study to organisms that can be raised in the lab or easily found in nature and transported to the

lab (Hanrahan & Johnston, 2011). This can be an insurmountable problem if the specimens are

difficult to collect, endangered or extinct, or only known from museum collections. NGS can

potentially provide a relatively simple alternative for bioinformatically estimating genome size;

however, the accuracy of these methods has not been extensively studied.

The first of these methods uses k-mer distributions. K-mers are unique subsequences of a

particular length, k, from a larger DNA sequence. For example, the DNA sequence AACCTG

can be decomposed into four unique k-mers that are three bases long (referred to as 3-mers):

AAC, ACC, CCT, and CTG. Any set of DNA sequences, including unassembled short reads

produced by NGS, can be broken down into its constituent k-mers. Each unique k-mer can be

assigned a value for coverage based on the number of times it occurs in a sequence (e.g., if the 3-

mer CTG is found a total of 20 times, it would have a coverage of 20). The distribution of

coverages for all k-mers from a sequence can be plotted to produce a k-mer frequency

distribution. For k-mers generated from genomic sequencing reads with negligible levels of

sequence artifacts (sequencing errors, repeats, or coverage bias), the distribution of k-mer

frequencies will approximate a Poisson distribution, with the peak centered on the average

sequencing depth for the genome (Li & Waterman, 2003). The value of k varies among analyses,

though values ranging between 17 and 35 are typical (Liu et al., 2013; Chen et al., 2015).

Techniques for estimating genome size using k-mer distributions generally work best

when the average coverage is greater than 10X (Williams et al. 2013), but newer methods with

more comprehensive models for addressing sequencing errors and repetitive sequence are

showing promise at coverages as low as 0.5X (Hozza, Vinar□, & Brejová, 2015). Examples of

accurate k-mer based genome size estimates exist for a variety of organisms, including giant

pandas (Li et al. 2010), cultivated potatoes (Potato Genome Sequencing Consortium, 2011), the

agricultural pest *Bemisia tabaci* (Chen et al. 2015), and oyster (Zhang et al. 2012). However,

these methods can also produce ambiguous or incorrect estimates. K-mer analysis of genomic

reads from a male milkweed bug produced estimates that were 60Mb to 1110Mb higher than the

approximately 930Mb flow cytometry genome estimate (Panfilio et al, 2018), with the

magnitude of this overestimation increased at larger values of k. A separate study on the *Bemisia*

tabaci genome (Guo et al. 2015) found that k-mer estimates of one particular biotype were about

60Mb larger than those given by flow cytometry. An alternative approach to inferring genome

size from sequence data is to map NGS reads onto a set of putative single-copy genes using a

reference-based assembler to determine the average coverage for the set of genes as a whole, and

use that average as an estimate of coverage for the entire genome (Desvillechabrol 2016, Kanda

et al. 2015).

Despite the potential value of sequence-based genome size estimation methods, little

empirical research to verify them has been conducted. In most instances, these methods are

incidental to the overall project and are only applied to a single individual or species. In this

study, we perform three of these sequence-based genome size estimation techniques, focusing on

beetles in the suborder Adephaga, and compare the results to genome size estimates derived from

flow cytometry.

MATERIALS AND METHODS

Taxon sampling and specimen processing: Specimens were collected in Oregon and California

(Tables S1, S2). Specimens assessed with flow cytometry were collected live and chilled, and

their heads removed and stored at -80°C. The remaining portions of the beetles were stored in

95%-100% ethanol and retained as vouchers. Eight of these specimens were also sequenced

(Table S2). No flow cytometry was performed on *Amphizoa insolens*, *Omoglymmius hamatus*,

and *Trachypachus gibbsii* as sufficient numbers of specimens could not be collected at the time

of the study. These three species were only assessed using sequence-based methods.

To insure sufficient low-copy-number reference sequences would be available for read

mapping, six transcriptomes were also sequenced (Table S3). These transcriptomes were derived

from whole-body RNA extractions from individual beetles conspecific to those used for genomic

sequencing with the exception of *Lionepha casta* DNA4602, which is a close relative to

Lionepha "Waterfalls." Specimens used for transcriptome sequencing were either stored in

RNAlater (Thermo Fisher Scientific, Waltham, MA) or kept alive until RNA extraction (Table

S4). The reference transcriptome for *Amphizoa insolens* was obtained from the NCBI SRA

(accession number: SRR5930489).

DNA extraction and shallow genome sequencing: DNA was extracted from all specimens

using a Qiagen DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany). DNA was extracted

from muscle tissue in most cases; however, reproductive tissue (one *Lionepha* "Waterfalls" and

three *Bembidion haplogonum* specimens) and, in one case (*Bembidion lividulum* DNA4146), the

entire body was extracted instead. The extractions for *Lionepha* "Waterfalls" DNA5435 and

Lionepha "Waterfalls" DNA5436 were derived from different tissues of the same specimen

(reproductive and muscle, respectively) to determine if any obvious qualitative differences

resulted from use of different tissue types.

The DNA was quantified using a Qubit Fluorometer (Life Technologies, Carlsbad, CA)

with a Quant-iT dsDNA HS Assay Kit, and DNA fragment length distributions with a 2100

Bioanalyzer (Agilent Technologies) using the High Sensitivity DNA Analysis Kit. The samples

were sheared for 10 minutes (30 seconds on, 30 seconds off) to a length of approximately 300bp

using a Bioruptor Pico Sonication System (Diagenode, Denville, NJ).

Libraries were prepared using either a NEBNext DNA Ultra II kit (New England

BioLabs) or an Illumina TruSeq DNA Sample Prep Kit (Table S5). Illumina sequencing was

performed at the Oregon State University Center for Genomic Research and Biocomputing (OSU

CGRB) on either a Hiseq 2500 or HiSeq 3000 (Table S5).

RNA extraction and transcriptome sequencing: All transcriptomic specimens were extracted

using Trizol Reagent (Thermo Fisher Scientific) and a Qiagen RNeasy Mini Kit. Whole beetles

were homogenized in liquid nitrogen, with the genitalia as well as a single antenna, leg, and

elytron being retained as a voucher for each individual. RNA extractions were quantified using a

Qubit Fluorometer (Life Technologies) with a Qubit RNA BR Assay Kit (Thermo Fisher

Scientific).

The RNA library for *Bembidion haplogonum* DNA3229 was prepared using an Illumina

TruSeq RNA Sample Prep Kit at HudsonAlpha Institute for Biotechnology (Huntsville, AL). For

all other specimens, mRNA was isolated using NEBNext Poly(A) mRNA Magnetic Isolation

Module (New England Biolabs, Ipswich, MA), and libraries were constructed with NEBNext

Ultra RNA Library Prep Kit for Illumina (New England Biolabs). The fragment size distribution

of each library was characterized with a 2100 Bioanalyzer (Agilent Technologies, Santa Clara,

CA) using the High Sensitivity DNA Analysis Kit and 1μl of sample.

Bembidion haplogonum DNA3229 was sequenced on an Illumina HiSeq 2000 at the

HudsonAlpha Institute for Biotechnology. The remaining transcriptome libraries were run on

either an Illumina HiSeq 2500 maintained by the OSU CGRB, or on an Illumina HiSeq 2500 at

the Oregon Health and Science University's Massively Parallel Sequencing Shared Resource

(Table S4).

Read processing and de novo assembly: Slightly different protocols were used to assemble

genomes and transcriptomes. The reads of one representative of each of the eight species were

imported into CLC Genomics Workbench (GW) v9.5.3 (CLC Bio-Qiagen, Aarhus, Denmark),

and low quality and Illumina adapter contaminated reads were removed using the "Trim

Sequences" tool with a quality limit parameter of 0.05 and an ambiguity limit of 2. *De novo*

genome assembly was performed in GW using an automatic word and bubble size.

Transcriptome reads were quality and adapter trimmed using the Agalma workflow (Dunn,

Howison, and Zapata 2013), and then assembled using Trinity (Grabherr et al. 2011).

For genome size estimation using k-mer methods or read mapping, reads were

preprocessed. To insure consistent read length for read mapping and k-mer genome size

estimation, the raw reads were reprocessed using BBduk v37.62 from the BBTools package

(Bushnell 2017). Reads containing Illumina adapters or other NGS artifact sequences, or with an

average quality score below 10, were discarded.

The relative quality of assemblies was assessed by identifying single-copy orthologs with

BUSCO v3 (Waterhouse et al. 2017) using the Endopterygota odb9 reference data set. In order to

identify and quantify repetitive elements, a random sample of 500,000 read pairs was generated

for each of the eight assemblies and analyzed with RepeatExplorer V2 (Novàk et al. 2013) with

default parameters and using the Metazoa V3 database.

Given that mitochondrial DNA can make up a substantial portion of the DNA present in a

cell (Moraes 2001), all libraries were screened for mitochondrial sequence. Mitochondrial DNA

sequences were assembled for each genome using NOVOPlasty v2.7.1 (Dierckxsens 2016).

NOVOPlasty produced complete circularized mitochondrial genomes for five of the eight

libraries (*Amphizoa insolens* DNA3784, *Chlaenius sericeus* DNA4821, *Omoglymmius hamatus*

DNA3783, *Pterostichus melanarius* DNA3787), while the remaining two libraries (*Bembidion*

haplogonum DNA2544 and *Lionepha* "Waterfalls" DNA3782) produced a single large (>16,000

bp) and several small (<2,500 bp) contigs. The six complete mitochondrial genomes and the two

largest contigs from DNA2544 and DNA3782 were combined in a single FASTA file. BBmap

was used to map reads from each library to these mitochondrial reference sequences (minid=0.7),

and the unmapped reads were used for subsequent "No Mito" read mapping analyses, which are

the primary focus of this paper unless otherwise stated. Although this will also remove from

consideration nuclear copies of mitochondrial DNA ("numts") that are similar enough to the

current mitochondrial genome, the fraction of reads that match mtDNA is low enough (0.14–

2.81%; see below) that removing numts can have at most a minimal effect on the estimate of

genome size, especially as most of the reads that match mtDNA are presumably from the

mitochondria.

Flow cytometry: Genome size was determined following methods in Johnston, Bernardini, and

Hjelmen (2019). One half of the head of each frozen adult sample was placed in ice-cold

Galbraith buffer, along with the head of a female *Drosophila virilis* strain maintained in the

laboratory of JSJ (1C = 328 Mb). All specimens of *Bembidion haplogonum* also contained the

head of a *Periplaneta americana* (1C = 3324 Mb) to act as a second internal standard. Combined

heads of the sample and standard were ground using 15 strokes of the "A" pestle in a 2 ml

Kontes Dounce tissue grinder and filtered through a 20μm nylon mesh. The DNA in the nuclei

released by grinding was stained for 2 hours under dark refrigeration with 25 μg/ml propidium

iodide. The mean red PI fluorescence of stained nuclei was quantified using a Beckman-Coulter

(Brea, CA) CytoFlex flow cytometer with a solid-state laser emitting at 488 nm. The total

quantity of DNA in the sample was calculated as the ratio of red fluorescence of sample/standard

(mean red fluorescence of the 2C peak of the sample divided by the mean fluorescence of the 2C

peak of the standard) times the 1C amount of DNA in the standard. To increase precision, the

genome size for each sample was estimated as the average from 2 technical replicates from the

two halves of the head of each individual. Up to 5 individuals of each sex were scored to produce

biological replicates, allowing calculation of the standard error of the mean genome size

estimate. The genome size is reported as 1C, the mean amount of DNA in Mb in a haploid

nucleus.

K-mer distribution: The optimal k-mer length for genome size estimation has not been

extensively tested, and the choice of k may impact the accuracy of estimates. At least one study

observed that k lengths above 16 resulted in an overestimation of genome size (Panfilio et al.

2017), while another observed little variation at different lengths of k (Sun et al. 2018). The

authors of GenomeScope recommend k=21 as a good tradeoff between computation speed and

accuracy (Vurture *et al.* 2017), though values between 17 and 27 have been used in other studies

(Zhang et al. 2015, Chen et al. 2015, Sun et al. 2018). To ensure that the length of k was not

affecting the estimates, all k-mer analyses were performed with values of k ranging from 13 to

31, at steps of two, for the four species represented by multiple specimens (*Bembidion*

haplogonum, Chlaenius sericeus, Lionepha "Waterfalls", and Pterostichus melanarius). One-

way ANOVA F-tests were performed for each combination of species and k-mer method

(Figures S1–S4, Sup File 1).

The input for all k-mer based analyses in this study were plain text histograms depicting

the number of k-mers at a given frequency. These were generated using the filtered Illumina

reads with Jellyfish v2.0 (Marçais & Kingsford 2011), and genome sizes were estimated using

two programs: GenomeScope (Vurture *et al.* 2017) and CovEST (Hozza et al. 2015). Histograms

were analyzed using GenomeScope with no maximum k-mer coverage (maximum k-mer

coverage = -1). GenomeScope uses an analytical model to estimate properties of a genome,

including genome size, average coverage, and heterozygosity, from the distribution of k-mer

frequencies. The k-mer histograms were also analyzed with CovEST using the "basic" and

"repeats" models. CovEST employs a model that accounts for error rate and repeat structure, and

has been demonstrated to perform well on low-coverage datasets (Hozza et al. 2015).

Read mapping: Two separate sets of loci were selected for use as reference sequences, both

containing putatively low-copy-number nuclear protein-coding genes. All loci consisted solely of

protein-coding, exonic sequence to exclude introns or other non-coding sequence unexpectedly

affecting read mapping. The first set contained the 74 low-copy-number nuclear genes

(Supplemental File 2). This set of loci (here referred to as "Regier") was initially selected for

studying arthropod evolution (Regier *et al.* 2008), and has been used for validating NGS results

in several previous studies of carabid beetles (Kanda *et al.* 2015; Sproul & Maddison 2017). The

reference sequences of these genes for each species were obtained by performing a pairwise

search, using Exonerate v2.4.0 (Slater & Birney 2005), of the corresponding transcriptome and

the *Bembidion haplogonum* references from Kanda et al. (2015). Because no transcriptome was

available for *Omoglymmius* or a near relative, reference sequences were generated directly from

the assembled *Omoglymmius hamatus* DNA3783 genome.

Read mapping was repeated with a second non-overlapping set of putatively single-copy

genes from OrthoDB v9.1 (Zdobnov et al. 2016); this set is here referred to as "ODB"

(Supplemental File 3). Loci from six coleopteran genomes were obtained (from *Agrilus*

planipennis, GCF_000699045.1; *Anoplophora glabripennis*, GCF_000390285.2; *Dendroctonus*

ponderosae, GCF_000355655.1; *Leptinotarsa decemlineata*, GCF_000500325.1; *Onthophagus*

taurus, GCF_000648695.1; *Tribolium castaneum*, GCF_000002335.3) by filtering for genes

present in >80% of species at the Endopterygota level, resulting in a total of 135 loci

(Supplemental File 4). Next, the loci were used as reference sequences for an orthologous gene

search of the eight adephagan transcriptomes created here using the Orthograph software

package (Petersen et al. 2017). Orthograph uses profile hidden Markov models derived from the

amino acid sequences of predetermined ortholog groups from related species to identify

orthologous genes in novel NGS data. These putative orthologs are then validated by a reciprocal

BLAST using the same set of orthologous gene groups.

Read mapping for both data sets was performed using BBmap v37.62 with the minid

parameter set to 0.7. Preprocessed reads from each of the genomic libraries were mapped to the

each of the gene sets. Sequencing coverage for each library was estimated by calculating the

mean coverage across all loci. The raw coverage data was generated using BBmap's per-scaffold

coverage output (using the *basecov* option). We generated plots showing coverage at each

nucleotide position of the reference loci (Figures 1, S5) and observed that the coverage profile

varied across each locus, with the first and last 60 to 70 bases having substantially lower

coverage than the central region of the locus. Some variation in coverage is inevitable due to the

stochastic nature of shotgun sequencing (Linder et al. 2013, Desvillechabrol et al. 2016);

however, the decreased coverage towards the flanks is likely an artifact of the read mapping

procedure, since BBMap does not map a read if only a small portion of the read overlaps with

the reference sequence. This causes the average coverage of the locus to be underestimated to

some degree, resulting in the average coverage estimate for all loci to be underestimated.

To compensate, a custom python script (https://github.com/JMPflug/gsec) was made to

trim low coverage, flanking regions of the read mapping and produce corrected coverage

estimates. The script begins with one "per-base" coverage file, containing the coverage at each

base position of all loci in the reference set, for each sequenced library. This type of file can be

generated by several read mapping utilities, including BBMap, Samtools, and Bedtools. The

script then excludes a user-selected number of base positions at the beginning and end of each

locus. The coverage profiles of the Regier set mappings for several specimens (Figures 1, S5)

indicated that the drop-off in relative coverage begins at roughly 75 bases from the ends of each

locus; we chose this value as the number of bases excluded by the script from each end. To

prevent outlier loci from skewing the calculated means, we included loci with coverage and

length values within three interquartile ranges of the median; loci with more extreme coverage or

length values were removed. The script then recalculates the average per-base coverage of each

locus in the reference set and uses these values to calculate the average genome coverage for the

library across all loci.

These average coverage values are then used as an estimate of the average coverage of

the genome, and genome size is calculated using the Lander-Waterman equation (Lander and

Waterman 1988),

$$G = \frac{LN}{C}$$

where C is the coverage estimated by read mapping, L is read length, N is the total

number of reads, and G is the haploid genome length.

Cytogenetic methods: A total of 14 males of *Bembidion haplogonum* and three males of

Lionepha "Waterfalls" were examined for chromosome number and size; specimens came from

the same localities studied for genome size. In brief, Feulgen staining followed by squashing was

used; more details are given in Maddison (1985; 2008). First and second meiotic metaphase and

anaphase were studied to determine chromosome number and sex chromosome system.

Data Availability: Raw genomic reads were deposited in the Short Read Archive of NCBI's

Genbank database under accession numbers SRR8518612 to SRR8518632 (Table 1), and

transcriptome reads were deposited to the NCBI SRA under accession numbers SRR8801541 to

SRR8801545 (Table S3). The custom python script used is available at:

https://github.com/JMPflug/gsec. All supplemental figures and tables are available on Figshare:

https://TBD. The supplemental figures are:

Figures S1-S4. Genome size estimates (in Mb) from GenomeScope and CovEST at different

values of K for S1. *Bembidion haplogonum*, S2. *Chlaenius sericeus*, S3. *Lionepha*

"Waterfalls", and S4. *Pterostichus melanarius*.

Figure S5. Average per-base read coverage for the first 200 bases in loci of the Regier set for

four specimens.

Figure S6. Relative red fluorescence and the number of nuclei counted at each level fluorescence

level of representative *Chlaenius sericeus, Lionepha "Waterfalls", and *Pterostichus*

melanarius compared to a *Drosophila virilis* standard.

Figure S7. Percent of reads inferred to contain repetitive elements as inferred by RepeatExplorer

from a sample of 500,000 read pairs for each of eight beetle species, with reads classified to

major group of repetitive elements.

The supplemental tables are:

Table S1. Information on genomic specimens sequenced.

Table S2. Flow Cytometry values for specimens examined.

Table S3. Information on transcriptomic specimens sequenced.

Table S4. Additional details for methods used on transcriptomic sequencing specimens.

Table S5. Additional details for methods used on genomic sequencing specimens.

Table S6. Results of BUSCO analysis on eight genome assemblies using the 2442 gene

Endopterygota odb9 reference set.

Table S7. Results of BUSCO analysis on six transcriptome assemblies using the 2442 gene

Endopterygota odb9 reference set.

Table S8. GenomeScope results.

Table S9. Summary of read mapping genome size estimates for the Regier and OrthoDB gene

sets using three different read filtering methods.

Table S10. CovEST genome size (in Mb) and coverage estimates for two models, Basic and

Repeat, performed using a k value of 21.

Table S11. Regier read mapping mean coverages before and after removing outliers using the

3*IQR rule.

Table S12. OrthoDB read mapping mean coverages before ("Untrimmed") and after ("IQR

Trim") removing outliers more than three interquartiles from the median.

Table S13. Comparison of total repetitive DNA sequence (in Mb) estimated by RepeatExplorer

and GenomeScope.

Table S14. GenomeScope genome size estimates using *Drosophila melanogaster* embryo,

salivary gland, and ovarian cell data set from Yarosh and Spradling (2014).

RESULTS

Flow cytometry: The genome sizes of the five species of carabid beetles studied (Tables 1, S2) vary over a five-fold range. *Bembidion haplogonum* possessed the largest genome, with estimates of 1C = 2,193.4♂ / 2,118.1♀Mb (Figure 2), while the genome of *Chlaenius sericeus* was the smallest with 1C = 408.4♂ / 391.5♀Mb (Figure S6). The genome of the former is over twice the size of the largest previously measured carabid beetle, *Calosoma scrutator,* which was estimated to be 1C = 1,017.1Mb (Hanrahan and Johnston 2011), and ranks as the 8th largest beetle genome out of the nearly 300 analyzed Coleoptera in the Animal Genome Size Database (Gregory 2001).

Sequencing and Assembly: Approximately 3.5 billion reads were generated across all 21

genomes, resulting in a total 473.6 billion bases. These were used to create draft assemblies of

the eight species of adephagan beetles studied. The total assembled scaffold lengths varied

between 153.1Mb for *Omoglymmius hamatus* DNA3783, and 665.1Mb for *Bembidion*

haplogonum DNA2544 (Table 2). The assembly lengths are substantially smaller than the

genome sizes inferred by both sequence-based and flow cytometric methods, indicating that

large portions of the genomes could not be assembled. The number of genes in each assembly

identified by BUSCO as "complete" (i.e., a putative orthologous gene found to similar to one of

the 2442 BUSCO gene groups, and whose length is within two standard deviations of the mean

length of the genes in that BUSCO group) varied greatly across the genome assemblies (Table

S6). 1,923 (78.7%) of the 2442 genes were completely found in *Chlaenius sericeus* DNA4821,

while *Omoglymmius hamatus* DNA3783 contained 178 (7.3%) complete genes.

A total of over 570 million reads were generated from the six transcriptomes (Table 3).

The number of transcripts assembled ranged from 22,330 with *Bembidion lividulum* DNA4279

to 57,119 with *Bembidion haplogonum* DNA3229. As anticipated, BUSCO was able to locate

many more genes in the transcriptomic assemblies (Table S7).

RepeatExplorer classified between 23.9% and 66.6% of the sampled reads as originating

from repetitive elements (Figure S7). Three samples, all from the carabid tribe Bembidiini,

consisted of over 50% repetitive DNA: *Bembidion haplogonum* DNA2544, *Lionepha*

"Waterfalls" DNA3782, and *Bembidion lividulum* DNA4161. Although a variety of repeat

families were identified, including Ty3/Gypsy, Ty1/Copia, Penelope, and LINEs,

RepeatExplorer was unable to classify the overwhelming majority of these repetitive sequences

(70.1% to 99.3%). 10.3% of the *Bembidion lividulum* DNA4161 reads were classified as

ribosomal DNA, a finding consistent with other studies on this species (Sproul and Maddison

2017). The genomic libraries consisted of between 0.14% to 2.81% mitochondrial DNA.

K-mer distribution: The estimated genome size changed with k-mer length for both CovEST

models, but not for GenomeScope. The length of k had a significant effect on estimated genome

size in all four species (one-way ANOVA, p-value < 0.001) for both "repeat" and "basic"

models. A non-significant difference in GenomeScope genome size estimates was observed at

varying sizes of k for all species (one-way ANOVA, p-value > 0.40).

Post-hoc Tukey-Kramer tests of the CovEST results revealed that estimates using a k of

less than 17 differed significantly (p-value < 0.05) from those using k values above 21 for all

species (Figures S1–S4, Sup. File 1). Values of k less than 17 always yielded much lower

genome size estimates than suggested by both flow cytometry and read mapping, while larger k

values (19 to 31) produced estimates that were more consistent with the results of other

estimation methods. Given these observations, and as the computational expense of generating k-

mer graphs increases as k becomes larger (Marçais & Kingsford 2011), a k value of 21 was

selected as the standard k value for this study. All subsequent discussion about k-mer based

estimates in this study, including tables and figures, refer to analyses performed using this value

unless otherwise stated.

GenomeScope converged and produced genome size estimates for 12 of the 21 specimens

(Tables S6, S8). GenomeScope failed to converge for *Amphizoa insolens* DNA3784, *Bembidion*

lividulum DNA4161, and four of the five *Pterostichus melanarius* samples (JMP059, JMP060,

JMP061, and JMP062). *Omoglymmius hamatus* DNA3783 and *Pterostichus melanarius*

DNA3787 did converge but yielded implausibly low genome size estimates (74Mb and 74.6MB,

respectively). The specimens that failed to converge lacked obvious coverage peaks in their k-

mer histograms, suggesting they did not have sufficient coverage for the GenomeScope model.

Coverage estimates of these specimens using read mapping provide support for this idea, as all

specimens that did not converge had an estimated coverage less than 26X, while only one

specimen whose coverage was less than 26X, *Lionepha* "Waterfalls" DNA3782, converged

(Tables S9, S10). A similar situation was observed for both CovEST models. All non-converging

specimens were estimated to have a coverage less than 13X for the "basic" model and less than

22X for the "repeat" model, with *Lionepha* "Waterfalls" DNA3782 again being the only

specimen to converge with a coverage below this level. This result is consistent the Vurture et al.

(2017)'s recommended minimum coverage of at least 25X coverage for an accurate estimate.

The successful analyses produced genome size estimates ranging from 1,113.7Mb for *Bembidion*

haplogonum DNA2544 to 264.0Mb for *Trachypachus gibbsii* DNA3786.

CovEST produced estimates for all 21 samples (Tables S6, S10), though the two models

behaved differently. The "repeats" model yielded estimates approximately double the size of the

"basic" model estimates, and the "repeat" model always produced the largest estimate among all

the sequence-based methods.

Read Mapping: There was no statistically significant difference between the estimates using the

Regier and ODB gene sets (two-sample t-test, p=0.1611). Removing mitochondrial reads

resulted, on average, in a modest increase in genome size estimates for both Regier (1.31%, s =

0.75%) and ODB (1.2%, s = 0.66%) gene sets (Table S9).

In most samples, we observed that a small number of loci with notably high coverage was

removed by the gsec script. In all but one instance, excluding these loci had a minimal impact on

the average coverage (Tables S11, S12). The single exception, *Omoglymmius hamatus*

DNA3783, contained three Regier set loci with substantially higher coverage (>160X) than the

rest of the set (Figure 3), which ranged between 2.10X to 9.27X. The inclusion of these three loci

nearly doubled the estimated genome size. The high-coverage of these loci suggests that all or

part of these genes are multi-copy in this species, or their coverage is being inflated by

extraneous reads from an undetected source such as a pseudogene or contaminant DNA.

Comparison of Flow Cytometry and Sequence-based Genome Size Estimates: Accuracy of

the sequence-based genome size estimation methods depended greatly on the species being

analyzed (Figures 4, 5, Table 4, S6). In most cases, sequence-based methods underestimated size

of the genome as measured by flow cytometry. In addition, estimates produced by the various

sequence-based methods sometimes differed from each other. The largest discrepancy among

methods was observed with the large genome of *Bembidion haplogonum*. The average flow

cytometry measurement of male *B. haplogonum* was 2,118.1Mb; however, read mapping,

GenomeScope, and the "basic" CovEST model estimated the genome to be approximately half

that size (932.0Mb to 1134.1Mb). A similar, though less pronounced, pattern was observed with

the next largest genome, of *Pterostichus melanarius*. In contrast, the CovEST "repeats" estimate

was within 10% of the flow cytometric value in three of the four *B. haplogonum* samples (Table

S6). For *Chlaenius sericeus*: read mapping, GenomeScope, and CovEST "basic" underestimated

genome size by a modest 4.50% on average, while the CovEST "repeats" model produced the

only overestimate, inflating the genome by an average of 78.0%. Sequence-based estimates of

Lionepha "Waterfalls" varied depending on the method but were generally lower than the flow

cytometric value.

Cytogenetic results: The chromosomes of *Bembidion haplogonum* are visibly larger than those

of *Lionepha* "Waterfalls" (Figure 6), which agrees with both flow cytometry and sequence-based

estimates. Males of *Bembidion haplogonum* have 11 pairs of autosomes and an XY pair of sex

chromosomes; males of *Lionepha* "Waterfalls" have 12 pairs of autosomes and have a single X

chromosome with no Y chromosome.

 DISCUSSION

K-mer Analysis: GenomeScope was unable to converge on nine of the 21 samples in this study,

which is likely attributable to the relatively low coverage of those samples. All five specimens of

Pterostichus melanarius had a coverage of <15X, and all five failed to converge. Of the samples

that did converge, the estimates for four of the five *Chlaenius sericeus* were within 6% of the

flow cytometric value. The estimates of the three *Lionepha* "Waterfalls" samples were not as

close, ranging between 6.8% and 17.5% lower than the true value. Curiously, the estimates for

all four *Bembidion haplogonum* were gross underestimates, around half the size of the flow

cytometry measurements. The coverage estimated by GenomeScope was double the value

expected for a species with a 2,118Mb haploid genome, leading to the underestimated genome

size. The k-mer histograms of *Bembidion haplogonum* DNA2544, the specimen with by far the

most reads, showed a distinct bimodal profile, with a large peak at 40X coverage and a shorter

peak at 80X. This is potentially indicative of a highly heterozygous genome (Vurture *et al.*

2017). Interestingly, GenomeScope also estimated that all four *Bembidion haplogonum* genomes

consisted of upwards of 75% repetitive sequence, which is similar to the RepeatExplorer

estimate (Table S13). A previous study has found that repeats, as well as high heterozygosity and

sequencing errors, can decrease the accuracy of genome size estimation using k-mer frequency

(Liu et al. 2013). This suggests the large number of repetitive sequences in the *Bembidion*

haplogonum genome may be at least partially responsible for the observed underestimates.

The striking difference between the estimates of the two CovEST models, with "repeat"

consistently giving estimates twice the size of "basic," suggests that neither model is well suited

to all genomes. Genome size estimates of the five *Chlaenius sericeus* specimens were closer to

the true value using the "basic" model, while the opposite was true of the five *Pterostichus*

melanarius specimens. The CovEST "repeat" model was the sequence-based method which

came closest to correctly estimating the 2,100Mb genome of *Bembidion haplogonum* with

estimates ranging between 1,480Mb to 2,140Mb. This is an interesting result given that

Bembidion haplogonum was shown by RepeatExplorer and GenomeScope to have the most

repeat-rich genome of the eight species analyzed, suggesting that *a priori* knowledge of the

amount of repetitive content of a genome may be necessary in order to select an appropriate

sequence-based genome size estimation approach.

Read Mapping: The method we used to infer coverage from read mapping data (averaging the

coverage across many single-copy loci) is relatively simple. Despite its simplicity, this approach

managed to perform well in some species, particularly those with smaller genomes such as

Chlaenius sericeus and *Lionepha* "Waterfalls." Read mapping proved to be inconsistent when

estimating genome size in *Pterostichus melanarius*. As with GenomeScope, it consistently

underestimated *Bembidion haplogonum* by approximately half.

The reason for the underestimation of *B. haplogonum* genome size based on single-copy

read coverage is unclear. Unlike k-mer based methods, which can struggle to assess highly

repetitive genomes, the read mapping approach used in this study only infers coverage from

single-copy exonic regions. In principle, as long as the selected loci are truly single-copy and the

resulting sequence data exhibit no biases in regions sequenced, the estimated coverage should

approach the true coverage. It follows that the presence of repetitive sequences elsewhere in the

genome should have no effect on this estimate. In actual genomes, evolutionary processes

responsible for increasing the size of the genome, such as segmental duplication followed by

neofunctionalization or pseudogenization, can lead to the proliferation of genes with similar

sequences (Rastogi and Liberles 2005, Levasseur and Pontarotti 2011). Such processes can

complicate the selection of a single-copy reference gene set, especially if the group of organisms

being studied lacks genomic resources.

We considered the possibility that using a larger set of reference loci from *Bembidion*

haplogonum may yield a better coverage estimate. To test this, we repeated read mapping on

Bembidion haplogonum DNA2544 using the 1421 single-copy orthologs annotated by BUSCO

and "No Mito" reads. However, this gave coverage and genome size estimates (94.89X and

1,115.9Mb, respectively) that were very similar to the Regier (96.18X, 1,101Mb) and ODB

(93.36X, 1,134Mb) gene sets, which casts doubt on insufficient references as the cause.

Comparison of Flow Cytometry and Sequence-based Genome Size Estimates: No single

sequence-based estimation method proved to be accurate in all cases. Species with large

genomes, such as *Bembidion haplogonum* and *Pterostichus melanarius*, appear to present the

greatest difficulty for inference of genome size by sequence-based means alone. It is tempting to

speculate that the same factors responsible for inaccuracy of k-mer methods are at work in read

mapping, especially given that the two methods often yielded underestimates of similar

magnitude; however, it is unclear what those factors may be, and they are not necessarily of

equal consequence to the two methods. Although both methods use the same underlying reads to

estimate coverage, they rely on somewhat different assumptions and portions of the genome. For

example, read mapping focuses on coding regions, but k-mer analysis analyzes the entire

genomes (Vurture et al. 2017).

The underestimation of the *Bembidion haplogonum* genome by half by most sequence-

based methods, including read mapping, is particularly puzzling. A potential explanation that

could have accounted for this is recent whole genome duplication or copy number variation;

however, cytological examination showed no evidence of genome duplication, nor are such

events known in *Bembidion* or near relatives (Maddison 1985, Serrano 1981, 1998). *Bembidion*

chromosomes do, however, possess several unusual characteristics. The number and shape of

chromosomes is remarkably consistent among species, with almost all species having males with

$2n = 22+XY$, a value shared by nearly all *Bembidion*, including *B. haplogonum* (Maddison

1985). Each autosome consists primarily of a large heterochromatic central region flanked by

small euchromatic tails, and males exhibit achiasmatic meiosis (Maddison 1985, Serrano 1981,

1998). Several species of *Bembidion*, including *Bembidion lividulum*, are also known to possess

highly replicated rDNA regions (Sproul and Maddison 2017). It is possible that some of these

chromosomal properties are involved in the underestimation of genome size by sequence-based

methods, but this is purely speculative, as a mechanism that could account for this unknown.

Genomic data can exhibit a variety of sequencing biases (Benjamini and Speed 2012,

Desvillechabrol et al. 2016), which could cause certain portions of the genome to be

underrepresented relative to the rest, leading to misestimation of genome size. For example,

consider an organism with a genome largely composed of AT-rich repetitive sequences. Because

a highly AT- or GC-rich sequence is less likely to be amplified during library preparation

(Benjamini and Speed 2012, Chen et al. 2013, Ross et al. 2013), the resulting library would

contain fewer reads derived from these repeats and more from the more AT-balanced protein-

coding regions, violating a core assumption of the sequence-based genome size estimation

methods (i.e. all bases have an equal probability of being sequenced). The resulting coverage

estimates would be artificially increased, causing a decrease in the subsequent genome size

estimates.

This particular issue, however, does not seem to be the case with *Bembidion haplogonum*,

as the GC content of the reads identified as repetitive elements by RepeatExplorer (33.0%) is

nearly identical to the overall GC content of the library (32.2%). It was also unremarkable

compared to the GC content of the other seven species analyzed, which was 32.3% on average

and ranged from 29.0% to 36.5%.

A more interesting possibility is that some biological process is physically causing parts

of the genome to be disproportionately represented within nuclei. Flow cytometric estimates

compare the unreplicated DNA amount in the G1 peak of the sample with the G1 peak of the

standard. The G1 peak is typically the major peak. If the major peak is G2 rather than G1 the

estimate will be a 2C value or twice the value assigned as a 1C. An increase in DNA per nucleus

is also created by underreplication, a process by which portions of the genome are replicated

more slowly during cell growth and division, resulting in fewer copies of certain parts of the

chromosomes (Belyaeva et al. 2008, Spradling and Orr-Weaver 1987). The most well-known

example of this occurs in the polytene salivary glands of *Drosophila* (Belyakin 2005, Yarosh and

Spradling 2014). Using the NGS genome data of Yarosh and Spradling (2014), we repeated

several of the sequence-based analyses employed in this study (Table S14), and observed that

genomic sequence from polytene salivary gland cells produced genome size estimates

significantly (two-sample t-test, p=0.000028) lower than those derived from unreplicated

embryonic cells, by an average of 36%. While cytological study of *Bembidion haplogonum*

effectively rules out any polytene-like chromosomal structures in the tissues sequenced for this

study, it is possible some other process, perhaps related to the unusual characteristics of

Bembidion chromosomes, is causing significant underreplication in the genome.

CONCLUSIONS

Increasing our knowledge of genome sizes across the tree of life is important for a deeper

understanding of genomic evolution, but the pace of this increase is currently very slow. In this

paper we have presented the first published flow cytometry estimates for five species of carabid

beetles; although this nearly doubles the number of carabids with genome size estimates, we now

have flow-cytometric estimates for less than 0.03% of carabid species.

With the explosion of short-read sequence data, and the development of methods to infer

genome sizes from these, the pace may increase, if the sequence-based methods are accurate

enough. We found that some of the sequence-based estimation techniques we investigated were

consistent with flow cytometry in some species, but no single technique was uniformly

congruent with flow cytometry. Flow cytometry should be the preferred option for estimating

genome size when live material and adequate resources are available. In cases where this is not

possible, especially when working with rare or extinct organisms, sequence-based methods can

provide an initial estimate of the size of a genome, and they may be well suited for cases when

the genome is likely to be small and non-repetitive. However, our work shows that these

techniques can also be misleading, particularly in large or highly repetitive genomes. Further

study on more genomes with different properties is necessary to better understand the strengths

and weaknesses of these methods.

ACKNOWLEDGEMENTS

We wish to thank Kipling W. Will for providing specimens of *Chlaenius sericeus*, and

John S. Sproul for providing specimens and Illumina genomic data for *Bembidion lividulum*. We

are also grateful to Mark Dasenko at the OSU CGRB for performing all the Illumina sequencing

for this project, as well as constructing several Illumina libraries. This project was partially

supported by Harold E. and Leona M. Rice Endowment Fund at Oregon State University and

NSF Grant DEB-1702062 to JMP and DRM.

LITERATURE CITED

Allen, S. K., 1983 Flow cytometry: Assaying experimental polyploid fish and shellfish.

Aquaculture 33: 317–328.

Belyaeva, E. S., E. N. Andreyeva, S. N. Belyakin, E. I. Volkova, and I. F. Zhimulev, 2008

Intercalary heterochromatin in polytene chromosomes of *Drosophila melanogaster*.

Chromosoma 117: 411–418.

Belyakin, S. N., G. K. Christophides, A. A. Alekseyenko, E. V. Kriventseva, E. S. Belyaeva *et*

al., 2005 Genomic analysis of *Drosophila* chromosome underreplication reveals a link

between replication control and transcriptional territories. Proceedings of the National

Academy of Sciences 102: 8269–8274.

Benjamini, Y., and T. P. Speed, 2012 Summarizing and correcting the GC content bias in high-

throughput sequencing. Nucleic Acids Research 40: e72–e72.

Bi, K., T. Linderoth, D. Vanderpool, J. M. Good, R. Nielsen *et al.*, 2013 Unlocking the vault:

next-generation museum population genomics. Molecular Ecology 22: 6018–6032.

Bock, D. G., N. C. Kane, D. P. Ebert, and L. H. Rieseberg, 2014 Genome skimming reveals the

origin of the Jerusalem Artichoke tuber crop species: neither from Jerusalem nor an

artichoke. New Phytologist 201: 1021–1030.

Buchfink, B., C. Xie, and D. H. Huson, 2015 Fast and sensitive protein alignment using

DIAMOND. Nature Methods 12: 59–60.

Bushnell, B., 2014 BBTools software package. URL http://sourceforge.net/projects/bbmap.

Cavalier-Smith, T., 1978 Nuclear volume control by nucleoskeletal DNA, selection for cell

volume and cell growth rate, and the solution of the DNA C-value paradox. Journal of Cell

Science 34: 247–278.

Chen, W., D. K. Hasegawa, K. Arumuganathan, A. M. Simmons, W. M. Wintermantel *et al.*,

2015 Estimation of the whitefly *Bemisia tabaci* genome size based on k-mer and flow

cytometric analyses. Insects 6: 704–715.

Chen, Y.-C., T. Liu, C.-H. Yu, T.-Y. Chiang, and C.-C. Hwang, 2013 Effects of GC bias in next-

generation-sequencing data on de novo genome assembly. PLOS ONE 8: e62856.

Desvillechabrol, D., C. Bouchier, S. Kennedy, and T. Cokelaer, 2016 Detection and

characterization of low and high genome coverage regions using an efficient running

median and a double threshold approach. bioRxiv 092478.

Dierckxsens, N., P. Mardulyn, and G. Smits, 2017 NOVOPlasty: de novo assembly of organelle

genomes from whole genome data. Nucleic Acids Research 45: e18–e18.

Dunn, C. W., M. Howison, and F. Zapata, 2013 Agalma: an automated phylogenomics

workflow. BMC Bioinformatics 14: 330.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson *et al.*, 2011 Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nature

Biotechnology 29: 644–652.

Gregory, T.R. 2019 Animal Genome Size Database. http://www.genomesize.com

Gregory, T. R., 2001 Coincidence, coevolution, or causation? DNA content, cell size, and the C-

value enigma. Biological Reviews 76: 65–101.

Gregory, T. R., 2005 Genome Size Evolution in Animals, pp. 3–87 in *The Evolution of the*

Genome, edited by T. R. Gregory. Academic Press, Burlington.

Hanrahan, S. J., and J. S. Johnston, 2011 New genome size estimates of 134 species of

arthropods. Chromosome Research 19: 809.

Hare, E. E., and J. S. Johnston, 2011 Genome size determination using flow cytometry of

propidium iodide-stained nuclei, pp. 3–12 in *Molecular Methods for Evolutionary Genetics*,

edited by V. Orgogozo and M. V. Rockman. Methods in Molecular Biology, Humana Press,

Totowa, NJ.

Hozza, M., T. Vinař, and B. Brejová, 2015 How big is that genome? Estimating genome size and

coverage from k-mer abundance spectra, pp. 199–209 in *String Processing and Information*

Retrieval, edited by C. Iliopoulos, S. Puglisi, and E. Yilmaz. Lecture Notes in Computer

Science, Springer International Publishing.

Johnston, J. S., A. Bernardini, and C. E. Hjelmen, 2019 Genome size estimation and quantitative

cytogenetics in insects, pp. 15–26 in *Insect Genomics: Methods and Protocols*, edited by S.

J. Brown and M. E. Pfrender. Methods in Molecular Biology, Springer New York, New

York, NY.

Johnston, J. S., L. D. Ross, L. Beani, D. P. Hughes, and J. Kathirithamby, 2004 Tiny genomes

and endoreduplication in Strepsiptera. Insect Molecular Biology 13: 581–585.

Kanda, K., J. M. Pflug, J. S. Sproul, M. A. Dasenko, and D. R. Maddison, 2015 Successful

recovery of nuclear protein-coding genes from small insects in museums using illumina

sequencing. PLOS ONE 10: e0143929.

Lander, E. S., and M. S. Waterman, 1988 Genomic mapping by fingerprinting random clones: A

mathematical analysis. Genomics 2: 231–239.

Leong-Škorničková, J., O. Šída, V. Jarolímová, M. Sabu, T. Fér *et al.*, 2007 Chromosome

numbers and genome size variation in Indian species of *Curcuma* (Zingiberaceae). Annals

of Botany 100: 505–526.

Levasseur, A., and P. Pontarotti, 2011 The role of duplications in the evolution of genomes

highlights the need for evolutionary-based approaches in comparative genomics. Biology

Direct 6: 11.

Li, R., W. Fan, G. Tian, H. Zhu, L. He *et al.*, 2010 The sequence and *de novo* assembly of the

giant panda genome. Nature 463: 311–317.

Li, X., and M. S. Waterman, 2003 estimating the repeat structure and length of DNA Sequences

using ℓ-tuples. Genome Research 13: 1916–1922.

Lindner, M. S., M. Kollock, F. Zickmann, and B. Y. Renard, 2013 Analyzing genome coverage

profiles with applications to quality control in metagenomics. Bioinformatics 29: 1260–

1267.

Liu, B., Y. Shi, J. Yuan, X. Hu, H. Zhang *et al.*, 2013 Estimation of genomic characteristics by

analyzing k-mer frequency in *de novo* genome projects. arXiv:1308.2012 [q-bio].

Maddison, D. R., 1985 Chromosomal diversity and evolution in the ground beetle genus

Bembidion and related taxa (Coleoptera: Carabidae: Trechitae). Genetica 66: 93–114.

Malé, P.-J. G., L. Bardon, G. Besnard, E. Coissac, F. Delsuc *et al.*, 2014 Genome skimming by

shotgun sequencing helps resolve the phylogeny of a pantropical tree family. Molecular

Ecology Resources 14: 966–975.

Marçais, G., and C. Kingsford, 2011 A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. Bioinformatics 27: 764–770.

Menzel, P., K. L. Ng, and A. Krogh, 2016 Fast and sensitive taxonomic classification for

metagenomics with Kaiju. Nature Communications 7: 11257.

Moraes, C. T., 2001 What regulates mitochondrial DNA copy number in animal cells? TRENDS

in Genetics 17: 199–205.

Mounsey, K. E., C. Willis, S. T. Burgess, D. C. Holt, J. McCarthy *et al.*, 2012 Quantitative PCR-

based genome size estimation of the astigmatid mites *Sarcoptes scabiei*, *Psoroptes ovis* and

Dermatophagoides pteronyssinus. Parasites & Vectors 5: 3.

Némorin, A., J. David, E. Maledon, E. Nudol, J. Dalon *et al.*, 2013 Microsatellite and flow

cytometry analysis to help understand the origin of *Dioscorea alata* polyploids. Annals of

Botany 112: 811–819.

Novák, P., P. Neumann, J. Pech, J. Steinhaisl, and J. Macas, 2013 RepeatExplorer: a Galaxy-

based web server for genome-wide characterization of eukaryotic repetitive elements from

next-generation sequence reads. Bioinformatics 29: 792–793.

Ozerov, M. Y., F. Ahmad, R. Gross, L. Pukk, S. Kahar *et al.*, 2018 highly continuous genome

assembly of Eurasian perch (*Perca fluviatilis*) using linked-read sequencing. G3: Genes,

Genomes, Genetics 8: 3737–3743.

Panfilio, K. A., I. M. V. Jentzsch, J. B. Benoit, D. Erezyilmaz, Y. Suzuki *et al.*, 2019 Molecular

evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the

milkweed bug genome. Genome Biology 20: 64.

703 Petersen, M., K. Meusemann, A. Donath, D. Dowling, S. Liu *et al.*, 2017 Orthograph: a versatile

704   tool for mapping coding nucleotide sequences to clusters of orthologous genes. BMC

705   Bioinformatics 18: 111.

706 Rastogi, S., and D. A. Liberles, 2005 Subfunctionalization of duplicated genes as a transition

707   state to neofunctionalization. BMC Evolutionary Biology 5: 28.

708 Regier, J. C., J. W. Shultz, A. R. Ganley, A. Hussey, D. Shi *et al.*, 2008 Resolving arthropod

709   phylogeny: exploring phylogenetic signal within 41 kb of protein-coding nuclear gene

710   sequence. Systematic Biology 57: 920–938.

711 Ross, M. G., C. Russ, M. Costello, A. Hollinger, N. J. Lennon *et al.*, 2013 Characterizing and

712   measuring bias in sequence data. Genome Biology 14: R51.

713 Serrano, J., 1981 A chromosome study of Spanish Bembidiidae and other Caraboidea

714   (Coleoptera, Adephaga). Genetica 57: 119–129.

715 Serrano, J., Galian, J., 1998. A review of karyotypic evolution and phylogeny of carabid beetles

716   (Coleoptera), pp. 191-228. in *Phylogeny and classification of Caraboidea (Coleoptera:*

717   *Adephaga),* by Ball, G.E., Casale, A., Vigna Taglianti, A. (Eds.). Proceedings of a

718   symposium (28 August, 1996, Florence, Italy). 20 International Congress of Entomology.

719   Museo Regionale di Scienze Naturali, Torino.

720 Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov, 2015

721   BUSCO: assessing genome assembly and annotation completeness with single-copy

722   orthologs. Bioinformatics 31: 3210–3212.

723 Slater, G. S. C., and E. Birney, 2005 Automated generation of heuristics for biological sequence

724   comparison. BMC Bioinformatics 6: 31.

Sota, T., J. Konuma, M. Fujiwara, and E. Shoguchi, 2013 Genome sizes of three species in the

subtribe Carabina (Coleoptera: Carabidae). Entomological Science 16: 122–124.

Spradling, A., and T. Orr-Weaver, 1987 Regulation of DNA replication during *Drosophila*

development. Annual Review of Genetics 21: 373–403.

Sproul, J. S., and D. R. Maddison, 2017 Cryptic species in the mountaintops: species

delimitation and taxonomy of the *Bembidion breve* species group (Coleoptera: Carabidae)

aided by genomic architecture of a century-old type specimen. Zoological Journal of the

Linnean Society 183: 556–583.

Sun, H., J. Ding, M. Piednoël, and K. Schneeberger, 2017 findGSE: estimating genome size

variation within human and *Arabidopsis* using k-mer frequencies. Bioinformatics 34: 550–

557.

The Potato Genome Sequencing Consortium, 2011 Genome sequence and analysis of the tuber

crop potato. Nature 475: 189–195.

Trost, B., S. Walker, Z. Wang, B. Thiruvahindrapuram, J. R. MacDonald *et al.*, 2018 A

comprehensive workflow for read depth-based identification of copy-number variation from

whole-genome sequence data. The American Journal of Human Genetics 102: 142–155.

Vurture, G. W., F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang *et al.*, 2017

GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics 33:

2202–2204.

Waterhouse, R. M., M. Seppey, F. A. Simão, M. Manni, P. Ioannidis *et al.*, 2017 BUSCO

applications from quality assessments to gene prediction and phylogenomics. Molecular

Biology and Evolution 35: 543–548.

Weitemier, K., S. C. Straub, R. C. Cronn, M. Fishbein, R. Schmickl *et al.*, 2014 Hyb-Seq:

Combining target enrichment and genome skimming for plant phylogenomics. Applications

in Plant Sciences 2: 1400042.

Williams, D., W. L. Trimble, M. Shilts, F. Meyer, and H. Ochman, 2013 Rapid quantification of

sequence repeats to resolve the size, structure and contents of bacterial genomes. BMC

Genomics 14: 537.

Yarosh, W., and A. C. Spradling, 2014 Incomplete replication generates somatic DNA

alterations within *Drosophila* polytene salivary gland cells. Genes & Development 28:

1840–1855.

Zdobnov, E. M., F. Tegenfeldt, D. Kuznetsov, R. M. Waterhouse, F. A. Simao *et al.*, 2016

OrthoDB v9. 1: cataloging evolutionary and functional annotations for animal, fungal,

plant, archaeal, bacterial and viral orthologs. Nucleic Acids Research 45: D744–D749.

Zhang, G., X. Fang, X. Guo, L. Li, R. Luo *et al.*, 2012 The oyster genome reveals stress

adaptation and complexity of shell formation. Nature 490: 49.

FIGURE CAPTIONS

Figure 1. Read mapping coverage at the beginning and end of each of the Regier set

loci using reads from *Bembidion haplogonum* DNA2544. The black line indicates the

average relative coverage along the length of the locus, and the blue line shows 75

base positions from either end of the locus.

Figure 2. Relative red fluorescence and the number of nuclei counted at each

fluorescence level of representative male (A) and female (B) *Bembidion haplogonum.*

Bars around each peak represent statistical gates that provide the total nuclei in that

peak, average channel number of nuclei in the peak, and the coefficient of variation

(CV). *D. virilis* standard 1C=328 Mb, *P. americana* standard 1C=3,338 Mb.

Figure 3. Boxplot of average genomic read mapping coverages for each of the Regier

(A) and OrthoDB (B) genes for eight representative specimens. Red dots indicate outlier

genes with coverage outside three interquartiles from the median.

Figure 4. Summary of genome size estimates using flow cytometry and sequence-

based methods for the eight adephagan species. Flow cytometry measurements are

averages of multiple individuals (see Tables TX02 and S3). CovEST Basic, CovEST

Repeat, and GenomeScope analyses were conducted using a k value of 21. Sequence-

based estimates were obtained from different individual specimens (Table TX01) than

those analyzed with flow cytometry.

Figure 5. Summary of genome size estimates using flow cytometry and sequence-

based methods for the 13 samples. CovEST Basic, CovEST Repeat, and

GenomeScope analyses were conducted using a k value of 21.

a. Samples made with DNA extracted from different tissues of the same individual.

Figure 6. Meiotic first metaphase cells. (A) Three cells of male *Bembidion haplogonum*

(B) Two cells of male *Lionepha* "Waterfalls". Photographs are at the same scale. Scale

bar 10μm.

	Female				Male			
	Genome Size	N	SD	SE	Genome Size	N	SD	SE
Bembidion haplogonum	2193.41	9	26.26	8.75	2118.05	9	27.77	9.26
Bembidion lividulum	837.39	1	-	-	831.70	5	6.57	2.94
Chlaenius sericeus	408.37	6	12.43	5.08	391.50	4	4.01	2.00
Lionepha "Waterfall"	597.60	1	-	-	585.70	5	5.41	2.42
Pterostichus melanarius	1040.65	4	18.42	9.21	1000.93	4	19.16	9.58

Table 1. Average flow cytometry genome size measurements. Values given in Mb. SD indicates standard deviation; SE indicates standard error.

	Read Length	Total Reads	Reads After Trim	N50	Average Contig (bp)	Maximum Contig (bp)	Contig Count	Total Assembled Bases
Amphizoa insolens DNA3784	101	59,590,984	59,182,421	355	299	79,486	905,365	270,465,134
Bembidion haplogonum DNA2544	151	717,036,094	713,035,405	525	460	140,578	1,438,922	661,385,315
Bembidion lividulum DNA4161	101	55,816,204	51,442,737	673	543	16,894	280,998	152,578,104
Chlaenius sericeus DNA4821	151	77,809,004	75,284,196	1,411	893	68,744	340,400	304,113,552
Lionepha "Waterfalls" DNA3782	101	80,214,318	79,962,073	4,114	1,481	47,683	124,458	184,268,766
Omoglymmius hamatus DNA3783	101	80,209,476	79,995,111	366	306	13,857	1,298,649	397,294,330
Pterostichus melanarius DNA3787	101	69,486,602	69,476,201	341	289	17,107	1,160,880	335,949,421
Trachypachus gibbsii DNA3786	101	88,000,000	86,431,416	1,258	588	65,825	472,429	277,787,110

Table 2. Summary of results for genomes assembled with CLC Genomics Workbench.

	Read Length	Reads Examined	Transcripts	N50	Average contig length	Total assembled bases
Bembidion haplogonum DNA3229	50	425,577,514	57,119	1274	898.66	35,767,428
Bembidion lividulum DNA4279	101	22,400,532	22,330	1727	1177.05	20,375,906
Chlaenius sericeus JMPR007	101	22,103,790	24,327	1764	1146.54	22,566,122
Lionepha casta DNA4602	100	30,791,800	26,994	1963	1338.76	23,724,243
Pterostichus melanarius DNA4765	100	38,061,712	34,153	2184	1338.91	30,509,665
Trachypachus gibbsii DNA4436	101	31,571,972	29,159	1823	1209.21	26,260,335

Table 3. Summary of results for transcriptomes assembled with Trinity.

	Flow Cytometry	Regier Mapping	ODB Mapping	GenomeScope	CovEST Basic	CovEST Repeat
Amphizoa insolens DNA3784	-	710.1	610.7	-	376.78	728.47
Bembidion haplogonum DNA2544	2118.1[b]	1,291.1	1,241.0	1113.67	932.04	2140.04
Bembidion haplogonum DNA5427	2118.1[b]	1,114.9	1,113.4	1010.67	827.88	1980.24
Bembidion haplogonum DNA5428	2118.1[b]	897.9	866.4	945.51	813.44	1924.27
Bembidion haplogonum DNA5433	2118.1[b]	983.3	946.4	908.41	758.09	1480.15
Bembidion lividulum DNA4161	831.7[b]	603.1	645.8	-	359.94	790.81
Chlaenius sericeus DNA4821	391.5	411.5	438.5	414.46	386.92	608.44
Chlaenius sericeus JMP068	385.8	390.1	395.7	374.46	410.91	751.89
Chlaenius sericeus JMP069	390.1	389.5	393.4	374.67	395.86	704.51
Chlaenius sericeus JMP070	396.5	354.5	393.1	-	409.81	796.39
Chlaenius sericeus JMP071	393.6	383.1	406.7	377.09	412.78	624.53
Lionepha "Waterfalls" DNA3782	585.7[b]	769.1	663.1	545.72	442.32	659.97
Lionepha "Waterfalls" DNA5435[a]	585.7[b]	518.1	535.9	483.39	340.38	423.66
Lionepha "Waterfalls" DNA5436[a]	585.7[b]	522.1	513.9	493.63	346.52	578.96
Omoglymmius hamatus DNA3783	-	1,525.7	1,552.1	-	627.13	1188.94
Pterostichus melanarius DNA3787	1000.9	1,144.9	984.6	-	475.11	1221.83
Pterostichus melanarius JMP059	1045.1	760.3	860.5	-	586.67	1127.86
Pterostichus melanarius JMP060	1018.3	855.5	787.4	-	597.29	1145.04
Pterostichus melanarius JMP061	1068.0	859.9	746.4	-	650.22	1454.54
Pterostichus melanarius JMP062	1031.2	1,103.7	927.6	-	678.61	1181.38
Trachypachus gibbsii DNA3786	-	342.0	315.5	264.08	314.81	525.13

Table 4. Summary of genome size estimates using flow cytometry and sequence-based methods. Values given in Mb. Flow cytometry Flow cytometry was not performed on *Trachypachus gibbsii* DNA3786, *Amphizoa insolens* DNA3784, and *Omoglymmius hamatus* DNA3782. CovEST Basic, CovEST Repeat, and GenomeScope analyses were conducted using a k value of 21. Cells in the GenomeScope column containing dashes indicate that sample failed to converge.

a. Samples made with DNA extracted from different tissues of the same individual.
b. Flow cytometry measurements for sample are species averages of multiple individuals (see Tables 2 and S3).