1

A dual function of the IDA peptide in regulating cell separation and modulating plant immunity at the molecular level

Vilde Olsson Lalun[1], Maike Breiden[§2], Sergio Galindo-Trigo[1§], Elwira Smakowska-Luzan[3], Rüdiger Simon[3] and Melinka A. Butenko[1*]

[1]Section for Genetics and Evolutionary Biology, Department of Biosciences, University of Oslo, 0316 Oslo, Norway

[2]Institute for Developmental Genetics and Cluster of Excellence on Plant Sciences, Heinrich Heine University, Universitätsstraße 1, 40225 Düsseldorf, Germany

[3]Gregor Mendel Institute (GMI), Austrian Academy of Sciences, Vienna Biocenter (VBC), Dr Bohr-Gasse 3, 1030 Vienna, Austria

[§]Equal contribution

*Corresponding author; email: m.a.butenko@ibv.uio.no

Abstract:

The abscission of floral organs and emergence of lateral roots in Arabidopsis is regulated by the peptide ligand INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) and the receptor protein kinases HAESA (HAE) and HAESA-LIKE 2 (HSL2). During these cell separation processes, the plant induces defense-associated genes to protect against pathogen invasion. However, the molecular coordination between abscission and immunity has not been thoroughly explored. Here we show that IDA induces a receptor-dependent release of cytosolic calcium ions (Ca^{2+}) and apoplastic production of reactive oxygen species, which are signatures of early defense responses. In addition, we find that IDA promotes late defense responses by the transcriptional upregulation of genes known to be involved in immunity. When comparing the IDA induced early immune responses to known immune responses, such as those elicited by flagellin22 treatment, we observe both similarities and differences. We propose a molecular mechanism by which IDA promotes signatures of an immune response in cells destined for separation to guard them from pathogen attack.

Introduction

All multicellular organisms require tightly regulated cell-to-cell communication during development and

adaptive responses. In addition to hormones, plants use small-secreted peptide ligands to regulate

these highly coordinated events of growth, development and responses to abiotic and biotic stress

(Matsubayashi, 2011; Olsson et al., 2018). In plants, organ separation or abscission, involves cell

separation between specialized abscission zone (AZ) cells and enables the removal of unwanted or

diseased organs in response to endogenous developmental signals or environmental stimuli.

Arabidopsis thaliana (Arabidopsis) abscise floral organs (stamen, petals and sepals) after pollination, and

this system has been used as a model to study the regulation of cell separation and abscission (Bleecker

& Patterson, 1997). Precise cell separation also occurs during the emergence of new lateral root (LR)

primordia, root cap sloughing, formation of stomata and radicle emergence during germination

(Roberts et al., 2000). In Arabidopsis, the INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) peptide

mediated signaling system ensures the correct spatial and temporal abscission of sepals, petals, and

stamens. The IDA protein shares a conserved C-terminal domain with the other 7 IDA-LIKE (IDL)

members (Butenko et al., 2003, Vie et al., 2015) which is processed to a functional 12 amino acid

hydroxylated peptide (Butenko et al., 2014). The abscission process is regulated by the production and

release of IDA, which binds the genetically redundant plasma membrane (PM) localized receptor kinases

(RKs), HAESA (HAE) and HAESA-LIKE 2 (HSL2). IDA binding to HAE and HSL2 promotes receptor

association with members of the co-receptor SOMATIC EMBRYOGENESIS RECEPTOR KINASE (SERK)

family and further downstream signaling leading to cell separation events. (Cho et al., 2008; Meng et

al., 2016; Santiago et al., 2016; Stenvik et al., 2008). A deficiency in the IDA signaling pathway prevents

the expression of genes encoding secreted cell wall remodeling and hydrolase enzymes thus hindering

floral organs to abscise (Butenko et al., 2003; Cho et al., 2008; Kumpf et al., 2013; Xiangzong Meng et

al., 2016; Niederhuth et al., 2013; Stenvik et al., 2008; Isaiah Taylor & John C. Walker, 2018).

Interestingly, components of the IDA signaling pathway control different cell separation events during

Arabidopsis development. IDA signaling through HAE and HSL2 regulates cell wall remodeling genes in

the endodermal, cortical, and epidermal tissues overlaying the LR primordia during LR emergence

(Kumpf et al., 2013). In addition, IDL1 signals through HSL2 to regulate cell separation during root cap

sloughing (Shi et al., 2018).

Plant cell walls act as physical barriers against pathogenic invaders. The cell-wall processing and

remodeling that occurs during abscission leads to exposure of previously protected tissue. This provides

an entry point for phytopathogens thereby increasing the need for an effective defense system in these

cells (Agustí et al., 2009). Indeed, it has been shown that AZ cells undergoing cell separation express

defense genes and it has been proposed that they function in protecting the AZ from infection after

abscission has occurred (Cai & Lashbrook, 2008; Niederhuth et al., 2013). *mIDA* elicits the production of

Reactive Oxygen Species (ROS) in *Nicotiana benthamiana* (*N.benthamiana*) leaves transiently expressing

the HAE or HSL2 receptor (Butenko et al., 2014). As a consequence of defense responses, ROS occurs in

the apoplast and is involved in regulating the cell wall structure by producing cross-linking of cell wall

components in the form of hydrogen peroxide (H_2O_2) or act as cell wall loosening agents in the form of

hydroxyl radical ($\cdot$OH)radicals (Kärkönen & Kuchitsu, 2015). ROS may also serve as a direct defense

response by their highly reactive and toxic properties (Kärkönen & Kuchitsu, 2015). In addition, ROS may

act as a secondary signaling molecule enhancing intracellular defense responses as well as contributing

to the induction of defense related genes (Reviewed in (Castro et al., 2021)). In Arabidopsis, apoplastic

ROS is mainly produced by a family of NADPH oxidases located in the plasma membrane known as

RESPIRATORY BURST OXIDASE PROTEINs (RBOHs) (Torres & Dangl, 2005). The production of ROS is often

found closely linked to an increase in the concentration of cytosolic Ca^{2+} ($[Ca^{2+}]_{cyt}$), where the interplay

between these signaling molecules is observed in immunity across kingdoms (Kadota et al., 2015;

Steinhorst & Kudla, 2013). Interestingly, ROS and $[Ca^{2+}]_{cyt}$ are known signaling molecules acting

downstream of several peptide ligand-receptor pairs. This includes the endogenous PAMPs ELICITOR

PEPTIDE(PEP)1, PEP2 and PEP3 which function as elicitors of systemic responses against pathogen attack

and herbivory (Huffaker et al., 2006; Ma et al., 2012; Qi et al., 2010), as well as the pathogen derived

peptide ligands flagellin22 (flg22) and Ef-Tu which bind and activate defense related RKs in the PM of

the plant cell (Felix et al., 1999; Gómez-Gómez et al., 1999; Ranf et al., 2011).

The interplay between ROS and $[Ca^{2+}]_{cyt}$ has been well studied in the flg22 induced signaling system. The

flg22 peptide interacts with the extracellular domain of the FLAGELLIN SENSING2 (FLS2) receptor kinase,

which rapidly leads to an increase in $[Ca^{2+}]_{cyt}$ (Ranf et al., 2011). In a resting state, the co-receptor

BRASSINOSTEROID INSENSITIVE1-ASSOCIATED RECEPTOR KINASE1/SERK3) (BAK1) and the cytoplasmic

kinase BOTRYTIS-INDUCED KINASE1 (BIK1) form a complex with FLS2. Upon binding of flg22 to the FLS2

receptor, BIK1 is released from the receptor complex and activates intracellular signaling molecules

through phosphorylation, among other, the PM localized RBOHD (Kadota et al., 2014; Li et al., 2014).

flg22 induced apoplastic ROS production binds to PM localized Ca^{2+} channels on neighboring cells,

activating Ca^{2+} influx into the cytosol. The cytosolic Ca^{2+} may bind to CALCIUM DEPENDENT PROTEIN

KINASE5 (CDPKs) regulating RBOHD activity through phosphorylation. The Ca^{2+} dependent

phosphorylation of RBOH enhances ROS production, which in turn leads to a propagation of the

ROS/Ca^{2+} signal through the plant tissue (Dubiella et al., 2013)

A rapid increase in $[Ca^{2+}]_{cyt}$ is not only observed in peptide-ligand induced signaling. Similar $[Ca^{2+}]_{cyt}$

changes are observed as responses to developmental signals, such as extracellular auxin and

environmental signals such as high salt conditions and drought(reviewed in (Kudla et al., 2018)). Also,

$[Ca^{2+}]_{cyt}$ signals are observed in single cells during pollen tube and root hair growth (Monshausen et al.,

2008; Sanders et al., 1999). How cells decipher the $[Ca^{2+}]_{cyt}$ changes into a specific cellular response is

still largely unknown, however the cell contains a large toolkit of Ca^{2+} sensor proteins that may affect

cellular function through changes in protein phosphorylation and gene expression. Each Ca^{2+} binding

protein has specific affinities for Ca^{2+} allowing them to respond to different Ca^{2+} concentrations

providing a specific cellular response (Geiger et al., 2010; Scherzer et al., 2012).

Due to our previous studies showing IDA induced apoplastic ROS production (Butenko et al., 2014), and

the known link between ROS and $[Ca^{2+}]_{cyt}$ response observed for other ligands such as flg22 (Dubiella et

al., 2013), we sought to investigate if IDA induces a release of $[Ca^{2+}]_{cyt}$. Here we report that IDA is able

to induce an intracellular release of Ca^{2+} in Arabidopsis root tips as well as in the AZ. Therefore, we

aimed to understand how the observed production of ROS and Ca^{2+} is involved in the IDA-HAE/HSL2

signaling pathway, either by being a part of the developmental process of abscission or as a part of a

suggested involvement of IDA in regulating plant defense (Patharkar et al., 2017). We show that a range

of biotic and abiotic signals can induce *IDA* expression and we demonstrate that the IDA signal

modulates responses of plant immune signaling. We propose that, in addition to regulating cell

separation, IDA plays a role in enhancing a defense response in tissues undergoing cell separation thus

providing the essential need for enhanced defense responses in tissue during the cell separation event.

<u>Results:</u>

IDA induces apoplastic production of ROS and elevation in cytosolic calcium-ion concentration

We have previously shown that a 12 amino acids functional IDA peptide, hereby referred to as mIDA

(Sup Table 1), elicits the production of a ROS burst in *N.benthamiana* leaves transiently expressing *HAE*

or *HSL2* (Butenko et al., 2014). To investigate whether mIDA could elicit a ROS burst in Arabidopsis, a

luminol-dependent ROS assay on *hae hsl2* mutant rosette leaves expressing the full length HAE receptor

under a constitutive promoter (*35S:HAE-YFP*) was performed. *hae hsl2 35S:HAE-YFP* leaf discs treated

with mIDA emitted extracellular ROS, whereas no response was observed in wild-type (WT) leaves most

likely due to the absence of *HAE* expression (Sup Fig. 1). Since rapid production of ROS is often linked to

an elevation in $[Ca^{2+}]_{cyt}$ (Steinhorst & Kudla, 2013), we aimed to investigate if mIDA could induce a

[Ca^{2+}]$_{cyt}$ response. We imaged Ca^{2+} in 10 days-old roots expressing the cytosolic localized fluorescent

Ca^{2+} sensor R-GECO1 (Keinath et al., 2015). As a positive control for [Ca^{2+}]$_{cyt}$ release we added 1mM

extracellular ATP (eATP), which leads to a [Ca^{2+}]$_{cyt}$ elevation in the root tip within a minute after

application (Fig. 1a) (Breiden et al., 2021). After application of 1µM mIDA, we detected an increase in

[Ca^{2+}]$_{cyt}$ (Fig. 1a and Movie 1). R-GECO1 fluorescence intensities normalized to background intensities

(ΔF/F) were measured from a region of interest (ROI) covering the meristematic and elongation zone of

the root and revealed that the response starts 4-5 minutes after application of the mIDA peptide and

lasts for 7-8 minutes (Fig. 1a). The signal initiated in the root meristematic zone from where it spread

toward the elongation zone and root tip, a second wave was observed in the meristematic zone

continuing with Ca^{2+} spikes. The signal amplitude was at a maximum within the elongation zone and

decreased as the signal spread (Fig. 1a).

The [Ca^{2+}]$_{cyt}$ response in roots to the bacterial elicitor flg22 has previously been studied using the R-

GECO1 sensor (Keinath et al., 2015). To better understand the specificity of the mIDA induced Ca^{2+}

response, we aimed to compare the Ca^{2+} dynamics in mIDA treated roots to those treated with flg22.

We observed striking differences in the onset and distribution of the Ca^{2+} signals. Analysis of roots

treated with 1 µM flg22 showed that the Ca^{2+} signal initiated in the root elongation zone from where it

spread toward the meristematic zone as a single wave and that the signal amplitude was at a maximum

within the elongation zone and decreased as the signal spread (Fig. 1b, Movie 2). These observations

indicate differences in tissue specificity of Ca^{2+} responses between mIDA and flg22, which we

hypothesize depend on the cellular distribution of their cognate receptors. Indeed, when investigating

the promoter activity of the *HAE*, *HSL2* and *FLS2* receptors by cloning the promoters fused to the nuclear

localized YFP-derived fluorophore Venus protein (*pHAE:Venus-H2B*, *pHSL2:Venus-H2B* and

pFLS2:Venus-H2B) we observed a different pattern of fluorescent nuclei in the roots (Fig. 1c,d,e).

pHAE:Venus-H2B lines had fluorescent expression in the epidermis and stele of the elongation zone (Fig.

1c) and *pHSL2:Venus-H2B* lines in the lateral root cap, root tip and root meristem (Fig. 1d); while

fluorescent nuclei were observed in the stele of the elongation zone in *pFLS2:Venus-H2B lines* (Fig. 1e).

The different patterns of fluorescent nuclei observed for the three different receptor constructs could

indeed explain the differences in the Ca^{2+} signatures triggered by mIDA and flg22. However, the

receptors promoter activity show a broader expression pattern compared to the respective ligand

induced [Ca^{2+}]$_{cyt}$ responses, indicating additional regulating factors for the observed [Ca^{2+}]$_{cyt}$ response.

The flg22 induced [Ca^{2+}]$_{cyt}$ response is completely abolished in the *fls2* mutant (Ranf et al., 2011). To

investigate the dependency of the mIDA-induced response on HAE and HSL2, a cytosolic localized

Aequorin-based luminescence Ca^{2+} sensor (Aeq) was introduced into the *hae hsl2* mutant background

(Knight et al., 1991). The Aeq sensor was chosen due to the well-established use of this sensor in mutant

studies (Ranf et al., 2011) and due to problems with transgene silencing of the R-GECO1 reporter in *hae*

hsl2 (not shown). The mIDA triggered increase in $[Ca^{2+}]_{cyt}$ was completely abolished in *hae hsl2 Aeq*

seedlings compared to WT *Aeq* expressing seedlings, indicating that the response observed is receptors

dependent (Sup Fig. 2a). Removing the last Asparagine of the mIDA peptide (IDAΔN69; Sup Table 1)

renders it inactive (Butenko et al., 2014). Seedlings treated with 1 µM IDAΔN69 did not show any

increase in $[Ca^{2+}]_{cyt}$ (Sup Fig. 2a). To investigate if the $[Ca^{2+}]_{cyt}$ increase triggered by mIDA was dependent

on extracellular Ca^{2+}, we pre-treated seedling with $LaCl_3$ and Ethylene glycol-bis (2-aminoethylether)- N,

N, N′,N′-tetraacetic acid (EGTA). $LaCl_3$ and EGTA are inhibitors that block plasma membrane localized

cation channels and chelate Ca^{2+} in the extracellular space, respectively (Knight et al., 1997). The mIDA

induced response was abolished in *Aeq*-seedlings pre-incubated in 2 mM EGTA or 1 mM $LaCL_3$ (Sup Fig.

2b), indicating that the mIDA dependent $[Ca^{2+}]_{cyt}$ response depends on Ca^{2+} from the extracellular space.

Next we set out to investigate whether mIDA would trigger an increase in $[Ca^{2+}]_{cyt}$ in floral AZ cells. We

utilized the Aeq expressing line and monitored $[Ca^{2+}]_{cyt}$ changes in flowers at different developmental

stages (see Sup Fig. 3a for developmental stages [floral positions]) to 1 µM mIDA. Interestingly, only

flowers at a the stage where there is an initial weakening of the AZ cell walls showed an increase in

$[Ca^{2+}]_{cyt}$ (Sup Fig. 3). Flowers treated with mIDA prior to cell wall loosening showed no increase in

luminescence (Sup Fig. 3), indicating that the mIDA triggered Ca^{2+} release in flowers correlates with the

onset of the abscission process and the increase in *HAE* and *HSL2* expression at the AZ (Cai & Lashbrook,

2008; Patharkar & Walker, 2015). Using plants expressing the R-GECO1 sensor we performed a detailed

investigation of the mIDA induced $[Ca^{2+}]_{cyt}$ response in flowers at position 6 which is the position where

initial cell wall loosening occurs. The AZ region was analyzed for signal intensity values and revealed that

the Ca^{2+} signal was composed of one wave (Fig. 2, Movie 3). The signal initiated close to the nectaries

and spread throughout the AZ and further into the floral receptacle. Flowers treated with 1 µM IDAΔN69

did not show any increase in $[Ca^{2+}]_{cyt}$ in the AZ or receptacle, while a clear Ca^{2+} wave could be detected

in the whole AZ, receptacle and proximal pedicel after treatment with eATP (Sup Fig.4, Sup Movie 2).

Detailed investigation of *HAE* and *HSL2* promoter activity in flowers at position 6 shows restricted

promoter activity to the AZ cells (Fig. 2b). The observed promoter activity correlates with the observed

mIDA induced $[Ca^{2+}]_{cyt}$ response.

The ROS producing enzymes, RBOHD and RBOHF, are not involved in the developmental process of

abscission.

ROS and Ca^{2+} are secondary messengers that can play a role in developmental processes involving cell

wall remodeling, but are also important players in plant immunity (Kärkönen & Kuchitsu, 2015). We

therefore set out to investigate if the IDA induced production of ROS and increase in $[Ca^{2+}]_{cyt}$ function in

the developmental process of abscission or if these signaling molecules form part of IDA modulated

plant immunity.

The IDA induced Ca^{2+} response depends on Ca^{2+} from extracellular space (Sup Fig. 2b). Ca^{2+} transport

over the plasma membrane is enabled by a variety of Ca^{2+} permeable channels, including the CYCLIC

NUCLEOTIDE GATED CHANNELs (CNGCs). Various CNGCs are known to be involved in peptide ligand

signaling in plants, including the involvement of CNGC6 and CNGC9 in CLAVATA3/EMBRYO

SURROUNDING REGION40 signaling in roots (Breiden et al., 2021), and CNGC17 In phytosulfokine

signaling (Ladwig et al., 2015). Using publicly available expression data we identified multiple CNGCs

expressed in the AZ of Arabidopsis (Cai & Lashbrook, 2008) (Sup Fig. 5a) and investigated if plants

carrying mutations in the *CNGSs* genes expressed in AZs showed a defect in floral organ abscission. We

observed no deficiency in floral abscission in the *cngc* mutants (Sup Fig. 5b), in contrast to what is

observed in the *hae hsl2* mutant where all floral organs are retained throughout the inflorescence. Ca^{2+}

partially activates members of the NOX family of NADPH oxidases (RBOHs), key producers of ROS

(Kadota et al., 2015). Two members of this family, *RBOHD* and *RBOHF* have high transcriptional levels in

the floral AZs (Cai & Lashbrook, 2008) and have been reported to be important in the cell separation

event during floral abscission in Arabidopsis (Lee et al., 2018). We therefore set out to investigate if ROS

production in AZ cells was dependent on RBOHD and RBOHF and to quantify to which degree these

NADPH oxidases were necessary for organ separation. We treated AZs with the ROS indicator, H2DCF-

DA, which upon contact with ROS has fluorescent properties, and observed ROS in both WT and to a

lower extent in *rbohd rbohdf* flowers. In addition, we observed normal floral abscission in the *rbohd*

rbohf double mutant flowers, indicated that neither ROS production in the AZ nor cell separation was

dependent on RBOHD and RBOHF (Sup Fig. 5c,d). These results are in stark contrast to what has

previously been reported, where cell separation during floral abscission was shown to be dependent on

RBOHD and RBOHF (Lee et al., 2018). We used a stress transducer to quantify the force needed to

remove petals, the petal breakstrength (pBS) (Stenvik et al., 2008), from the receptacle of gradually

older flowers along the inflorescence of WT and *rbohd rbohf* flowers. Measurements showed a

significant lower pBS value for the *rbohd rbohf* mutant compared to WT at the developmental stage

where cell loosening normally occurs, indicating premature cell wall loosening (Sup Fig. 5e).

Furthermore, *rbohd rbohf* petals abscised one position earlier than WT (Sup Fig. 5e). In conclusion, we

found no RBOHD and RBOHF dependency for cell separation during floral abscission (Crick et al., 2022).

Due to a known function of IDA in inducing ROS production (Sup fig 1), these results points towards a

role for IDA in modulating additional responses to the process of cell separation.

IDA is upregulated by biotic and abiotic factors

In Arabidopsis, infection with the pathogen *P.syringae* induces cauline leaf abscission. Interestingly, stress induced cauline leaf abscission was reduced in plants with mutations in components of the IDA-HAE/HSL2 signaling pathway (Patharkar et al., 2017; Patharkar & Walker, 2016). In addition, upregulation of defense genes in floral AZ cells during the abscission process is altered in the *hae hsl2* mutant (Cai & Lashbrook, 2008; Niederhuth et al., 2013), indicating that the IDA-HAE/HSL2 signaling system may be involved in regulating defense responses in cells undergoing cell separation. To further investigate the involvement of IDA in stress responses we investigated if *IDA* is upregulated by biotic and abiotic elicitors. Transgenic plants containing the *β-glucuronidase (GUS)* gene under the control of the *IDA* promoter (*pIDA:GUS*) have previously been reported to exhibit *GUS* expression in cells overlaying newly formed LR primordia (Butenko et al., 2003; Kumpf et al., 2013). To investigate the expression of the *IDA* gene in response to biotic stress *pIDA:GUS* seedlings were exposed to the bacterial elicitor flg22 and fungal chitin. Compared to the control seedlings, the seedlings exposed to flg22 and chitin showed a significant increase in *GUS* expression in LR primordia (Fig. 3a). Also, when comparing untreated root tips to those exposed to flg22 and chitin we observed a GUS signal in the primary root of flg22 and chitin treated roots (Fig. 3b). We used the fluorogenic substrate 4-methylumbelliferyl β-D-glucuronide (4-MUG) which is hydrolysed to the fluorochrome 4-methyl umbelliferone (4-MU) to quantify GUS-activity (Blazquez, 2007). A significant increase in fluorescence was observed for flg22 and chitin treated samples compared to untreated controls, (Fig. 3b). Exposure of *pIDA:GUS* seedlings to IDAΔN69 did not increase GUS activity in the root, supporting the specific increment of *pIDA:GUS* expression to flg22 peptide (Fig. 3). Spatial *pIDA:GUS* expression was also monitored upon abiotic stress. When *pIDA:GUS* seedling were treated with the osmotic agent mannitol or exposed to salt stress by NaCl, both treatments simulating dry soil, a similar increment in GUS signal was observed surrounding the LR primordia (Fig. 3). However, no *pIDA:GUS* expression was detected in the primary root (Fig. 3b). Taken together these results indicate an upregulation of *IDA* in tissue involved in cell separation processes such as the LR primordia and, interestingly, the root cap upon biotic stress. The process of cell separation is a fast and time-restricted response where cells previously protected by outer cell layers are rapidly exposed to the environment. These cells need strong and rapid protection against the environment, and we therefore aimed to investigate if IDA is involved in modulating immune responses in addition to inducing ROS and $[Ca^{2+}]_{cyt}$ responses.

mIDA does not induce production of callose as a long term defense response

We aimed to investigate a possible role of IDA in regulating long-term defense responses in the plant,
such as the production of callose. Callose deposition is a resulting hallmark of the late plant immunity
responses that together with other events such as stomatal closure and production of ethylene leads to
inhibition of pathogen multiplication and containment of disease (Reviewed in (Wang et al., 2021)).
Callose deposition is highly increased in response to flg22 (Gómez-Gómez et al., 1999). We treated WT
seedlings with 1 µM mIDA and measured the area of callose deposition in the cotyledons, but detected
no difference to water treated controls (Fig. 4a,b). However, flg22 treated WT seedlings showed
increased callose deposition, which was not observed in flg22 treated *fls2* seedlings (Fig. 4a,b). These
results indicate that mIDA, in contrast to flg22, is not involved in promoting deposition of callose as a
long-term defense response.

mIDA triggers expression of defense-associated marker genes associated with innate immunity.

A -well-known part of a plants immune response is the enhanced transcription of genes involved in
immunity. Transcriptional reprogramming mostly mediated by WRKY transcription factors takes place
during microbe-associated molecular pattern (MAMP)- triggered immunity and is essential to mount an
appropriate host defense response (Birkenbihl et al., 2017). We selected well-established defense-
associated marker genes: *FLG22-INDUCED RECEPTOR-LIKE KINASE1* (*FRK1*), a specific and early immune-
responsive gene activated by multiple MAMPs (Asai et al., 2002; He et al., 2006), *MYB51*, a WRKY target
gene, previously shown to increase in response to the MAMP flg22 (Frerigmann et al., 2016) resulting
in biosynthesis of the secondary metabolite indolic glucosinolates; and an endogenous danger peptide,
ELICITOR PEPTIDE 3 (PEP3) (Huffaker et al., 2006). Seven days-old WT seedlings treated with 1µM mIDA
were monitored for changes in transcription of the aforementioned genes by RT-qPCR compared to
untreated controls at two time points, 1h and 12h after treatment. All genes showed a significant
transcriptional elevation 1 h after mIDA treatment (Fig. 5a). This in accordance with previous reports,
where the expression of *FRK1* and *PEP3* was monitored in response to bacterial elicitors (He et al., 2006;
Huffaker et al., 2006). After 12 h, *FRK1* and *PEP3* transcription in mIDA treated tissue was not
significantly different from untreated tissue; while transcription of MYB51 in mIDA treated tissue
remained elevated after 12 h but was significantly lower compared to 1 h after mIDA treatment (Fig.
5a). Interestingly, the observed increase in transcription of the defense genes *MYB51* and *PEP3*, was
only partially reduced in the *hae hsl2* mutant (Sup Fig. 6). In contrast, *FRK1* showed higher
transcriptional levels in the *hae hsl2* mutant compared to WT (Sup Fig. 6). These results indicate that
also other receptors than HAE and HSL2 are involved in the mIDA-induced enhancement of the defense
genes *FRK1, MYB51* and *PEP3*.

We then analyzed a plant line expressing nuclear localized YFP from the MYB51 promoter
(*pMYB51:YFP*[N]) (Poncini et al., 2017) treated with mIDA. Enhanced expression of p*MYB51:YFP* was
predominantly detected in the meristematic zone of the root after 7 h of mIDA treatment, compared to
a non-treated control (Fig. 5b). Moreover, comparison of mIDA induction with the elicitor-triggered
response of *pMYB51:YFP*[N] to flg22 showed similar temporal expression in the root (Sup Fig. 7) .
Together, these data show that IDA can trigger a rapid increase in the expression of key genes involved
in immunity.

We next investigated if the mIDA induced transcription of defense related genes was similar to that
induced by flg22. Seven days-old Col-0 WT seedlings were treated for 1h with mIDA, flg22 or a
combination of mIDA and flg22 to explore potential additive effects. The relative increase in
transcription of the genes tested is similar when seedlings are treated with mIDA and flg22 (Fig. 6).
However, when treating seedlings with both peptides the relative transcription of the genes of interest
increased substantially. To our surprise, the combined enhanced transcription when seedling where co-
treated with both flg22 and mIDA exceeds a simple additive effect of the two peptides, suggesting that
IDA has a role in enhancing the defense response triggered by flg22. Based on these results we propose
a role for IDA in enhancing immune responses in tissue undergoing cell separation.

Discussion:

Several plant species can abscise infected organs to limit colonization of pathogenic microorganisms
thereby adding an additional layer of defense to the innate immune system (Kissoudis et al., 2016;
Patharkar et al., 2017). Also, in a developmental context, there is an induction of defense associated
genes in AZ cells during cell separation and prior to the formation of the suberized layer that functions
as a barrier rendering protection to pathogen attacks (Cai & Lashbrook, 2008; Niederhuth et al., 2013;
Roberts et al., 2000; I. Taylor & J. C. Walker, 2018). Interestingly, the induction of defense genes during
floral organ abscission is altered in *hae hsl2* plants (Niederhuth et al., 2013). It is largely unknown how
molecular components regulating cell separation and how the IDA-HAE/HSL2 signaling pathway,
contributes to modulation of plant immune responses. Here we show that the mature IDA peptide,
mIDA, is involved in activating known defense responses in Arabidopsis, including the production of ROS,
the intracellular release of Ca^{2+} and the transcription of genes known to be involved in defense (Sup Fig.
1, Fig. 1, Fig. 2, Fig. 5). We suggest a role for IDA in ensuring optimal cellular responses in tissue
undergoing cell separation, regulating both development and defense.

The intracellular event occurring downstream of IDA is still partially unknown. Here we report the

release of cytosolic Ca^{2+} as a new signaling component. Roots and AZs expressing the Ca^{2+} sensor R-

GECO1, showed a $[Ca^{2+}]_{cyt}$ in response to mIDA. Moreover, the increased $[Ca^{2+}]_{cyt}$ in response to mIDA in

flowers occurred only at stages where cell separation was taking place, linking the observed $[Ca^{2+}]_{cyt}$

response to the temporal point of abscission and the need for an induced local defense response. During

floral development the expression of the *HAE* and *HSL2* receptors increase prior to the onset of

abscission (Cai & Lashbrook, 2008) and may account for the temporal mIDA induced $[Ca^{2+}]_{cyt}$ response

observed.

Based on these results, it is likely that the expression of the receptors is the limiting factor for a cell's

ability to respond to the mIDA peptide with a $[Ca^{2+}]_{cyt}$ response. However, promoter activity of the

receptors are observed in a broader area of the root than the observed $[Ca^{2+}]_{cyt}$ response. As an example,

pHSL2 is found to have a localized activation in the root tip, and when HSL2 is activated by IDL1 be

involved in regulation cell separation during root cap sloughing (Shi et al., 2018). IDL1 and IDA have a

high degree of similarity in the protein sequence, and *IDL1* expressed under the *IDA* promoter can fully

rescue the abscission phenotype observed in the *ida* mutant (Stenvik et al., 2008). We can assume that

IDL1 activates similar signaling components in the root cap as those used by IDA. Despite the expression

of HSL2 and the involvement of the receptor in root cap sloughing, no $[Ca^{2+}]_{cyt}$ response is observed in

this region when plants are treated with the IDA peptide. The absence of an observed $[Ca^{2+}]_{cyt}$ response

in the root tip indicates that $[Ca^{2+}]_{cyt}$ is most likely not involved in the cell separation process during root

cap sloughing, or that cells involved in this process only are able to induce a $[Ca^{2+}]_{cyt}$ response at a

specific developmental time point. Thus, in addition to receptor availability, other factors must be

involved in regulating the $[Ca^{2+}]_{cyt}$ response. It is intriguing that the mIDA cellular output is similar to that

of flg22 (Kadota et al., 2015; Kadota et al., 2014) yet shows some distinct differences. The spatial

distribution of the Ca^{2+} wave originated by mIDA differs from that of flg22. The *rbohd rbohf* mutant

shows ROS production in the AZ, indicating that other NADPH oxidases or cell wall peroxidases

expressed in the AZ may be involved in ROS production, in contrast to the importance of RBOHD RBOHF

in flg22 signaling (Kadota et al., 2015). This suggests that while IDA and flg22 share many components

for their signal transduction, such as the BAK1 co-receptor (Chinchilla et al., 2007; X. Meng et al., 2016)

and MAPK cascade (Asai et al., 2002; Cho et al., 2008), there are other specific components that are

likely important to provide differences in signaling. The difference in callose deposition observed in

mIDA and flg22 cotyledons supports this.

Interestingly, FLS2 arrangement into nanoscale domains at the PM is dependent on FERONIA (FER). FLS2

becomes more disperse and mobile in *fer* mutants, implying a role for FER in regulating FLS2 activity

(Gronnier et al., 2022). Also altering of the cell wall affects FLS2 nanoscale organization (McKenna et al.,

2019). Formation of nanoscale domains containing the HAE and HSL2 receptors may be an important

signaling step in the IDA-HAE/HSL2 signaling pathway. An intriguing thought is a possible co-localization

of HAE and HSL2 with FLS2 in nanoscale domains, making a signaling unit possible to involve multiple

different receptors responsible for the signaling outcome observed. In this paper, we show that co-

treatment with mIDA and flg22 induces an enhanced transcription of defense related genes. How FLS2,

HAE and HSL2 communicate at the plasma membrane of the responsive cells may give the answers to

the molecular mechanisms behind the augmented transcriptional regulation observed when co-

treatment of IDA and flg22 is performed (Fig. 6). We can investigate possible interaction partners of

HAE and HSL2 looking at available data from a sensitized high-throughput interaction assay between

extracellular domains (ECDs) of 200 Leucine Rich Repeat (LRR) (Smakowska-Luzan et al., 2018).

Interestingly, LRR-RLKs known to play a function in biotic or abiotic stress responses, such as RECEPTOR-

LIKE KINASE7 (RLK7), STRUBBELIG-RECEPTOR FAMILY3 , RECEPTOR-LIKE PROTEIN KINASE 1 and FRK1,

are found to interact mainly with HSL2, whereas HAE shows interaction with LRRs mainly involved in

development (Sup Fig. 8). RLK7 has recently been identified as an additional receptor for CEP4, involved

in regulating cell surface immunity and response to nitrogen starvation, indication a role of RLK7 as an

additional receptor regulating defense responses in known peptide-receptor complexes (Rzemieniewski

et al., 2022). It would be interesting to investigate a possible role of RLK7 in IDA-HAE/HSL2 signaling as

a possible signaling partner in the observed IDA-induced early defense responses.

In this paper, we propose a model where the IDA-HAE/HSL2 signaling pathway modulates defense

responses in tissues undergoing cell separation. It is beneficial for the plant to ensure an upregulation

of defense responses in tissues undergoing cell separation as a precaution against pathogen attack. This

tissue is a major entry route for pathogens, and by combining the need of both the IDA peptide and the

pathogen molecular pattern, such as flg22, for the induction of a strong defense response, the plant

ensures maximum immunity in the most prone cells without spending energy inducing this in all cells

(Fig. 7). Such a molecular mechanism, where the presence of mIDA in tissue exposed to stress leads to

maximum activation of the immune responses, will ensure maximal protection of infected cells during

cell separation, cells which are major potential entry routes for invading pathogens.

Movies

Movie 1: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 µM mIDA. 10 days-old roots

were treated with 1 µM mIDA and changes in $[Ca^{2+}]_{cyt}$ was recorded over time. Response shown as

normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie

corresponds to measurements shown in Fig. 1a. Representative response from 10 roots. Time shown as,

hh:mm:ss.

Movie 2: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 µM flg22 10 days-old roots were

treated with 1 µM flg22 and changes in $[Ca^{2+}]_{cyt}$ was recorded over time. Response shown as normalized

fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds

to measurements shown in Fig. 1b. Representative response from 10 roots. Time shown as, hh:mm:ss.

Movie 3: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing AZ in response to 1 µM mIDA. Flowers at position 6 were

treated with 1 µM mIDA and changes in $[Ca^{2+}]_{cyt}$ was recorded over time.   Response shown as normalized

fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds

to measurements shown in Fig. 2a. Representative response from 10 flowers. Time shown as, hh:mm:ss.

Sup Movie 1: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 mM eATP. 1 mM eATP will

induce a strong increase in $[Ca^{2+}]_{cyt}$ and was added as a last treatment to all roots acting as a positive control.

Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of

5 seconds. Movie corresponds to measurements shown in Fig. 1a. Representative response from 10 roots.

Time shown as, hh:mm:ss.

Sup Movie 2: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing AZ in response to to 1 µM of the inactive IDA peptide,

$IDA^{\Delta N69}$. Flowers at position 6 were treated with 1 µM $IDA^{\Delta N69}$ and changes in $[Ca^{2+}]_{cyt}$ was recorded over

time.   Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame

rate of 5 seconds. Movie corresponds to measurements shown in Sup Fig. 3. Representative response from

10 flowers. Time shown as, hh:mm:ss.

**Methods**

Accession numbers of genes studied in this work:

HSL2 At5g65710, IDA At1g68765, FLS2 At5g46330, HAE At4g28490, MYB51 At1g18570, RBOHD

At5g47910, RBOHF At1g64060, PEP3 At5g64905, WRKY33 At2g38470, FRK1 At2g19190. Plant lines used

in this work: Ecotype Colombia-0 (Col-0) was used as wild type (WT). Mutant line: _hae_ (SALK_021905),

hsl2, (SALK_030520), _fls2_ (SALK_062054), _rbohd_ (SALK_070610), _rbohf_ (SALK_059888), _cngc1_

(SAIL_443), *cngc2* (SALK_066908), *cngc4* (SALK_081369), *cngc5* (SALK_149893), *cngc6* (SALK_064702),

cngc9 (SAIL_736), *cngc12* (SALK_093622). SALK lines were provided from Nottingham Arabidopsis Stock

Centre (NASC).

Plant lines

The pIDA:GUS, pHSL2:Venus-H2B and pMYB51:YFPN lines have been described previously (Kumpf et al.,

2013; Poncini et al., 2017; Shi et al., 2018).The promoters of HAE (1601 bp (Kumpf et al., 2013) and HSL2

(2300 bp (Sto et al., 2015))were available in the pDONRZeo vector (Thermo Fischer Scientific).

Sequences corresponding to the FLS2 promoter (988 bp (Robatzek et al., 2006)) were amplified from

WT DNA (primers are listed in table 2) and cloned into the pDONRZeo vector (Thermo Fischer Scientific).

All promoter constructs were further recombined into the promotor:Venus (YFP)-H2B and

Cerulean(CFP)-H2B destination vectors (Somssich et al., 2016) using the Invitrogen Gateway cloning

system (Thermo Fischer Scientific). Constructs were transformed into *Agrobacterium tumefaciens*

(A.tumefaciens) C58 and the floral dip method (Clough & Bent, 1998) was used to generate transgenic

lines. Single-copy homozygous plant lines were selected and used in this study. The CDS of HAE was

cloned into the pEarleyGate101 destination vector (Earley et al., 2006) using the Invitrogen Gateway

cloning system and transformed into A.tumefaciens C58 and further used to generate the 35S:HAE:YFP

lines.

Growth conditions

Plants were grown in long day conditions (8h dark and 16h light) at 22°C. Seeds were surface sterilized

and plated out on MS-2 plates, stratified for 24 h at 4 °C and grown on plates for 7 days before

transferred to soil.

Peptide sequences

Peptides used in this study were ordered from BIOMATIK. Peptide sequences are listed in

Supplementary Table 1.

Primers

Primers for genotyping and generation of constructs were generated using VectorNTI. Gene specific

primers for RT-qPCR were generated using Roche Probe Library Primer Design. All primers are listed in

Supplementary Table 2.

Histochemical GUS assay

Seven days-old seedlings were pre-incubated for 12 h in liquid MS-2 medium containing stimuli of
interest; 1 µM peptide (table 1), 60 mM Mannitol (M4125 – Sigma), 20 µg/mL Chitin (C9752 – Sigma),
50 mM NaCl and then stained for GUS activity following the protocol previously described (Stenvik et
al., 2008). Roots were pictured using a Zeiss Axioplan2 microscope with an AxioCam HRc, 20x air
objective. The assay was performed on 10 individual roots and the experiment was repeated 3 times.

<u>Fluorescent GUS assay</u>

Seven days-old seedlings were pre-incubated for 12 h in liquid MS-2 medium with or without stimuli of
interest; 1 µM peptide (table 1), 60 mM Mannitol (M4125 – Sigma), 20 µg/mL Chitin (C9752 – Sigma),
50 mM NaCl. After treatment, 10 seedlings were incubated in wells containing 1 mL reaction mix
described in (Blazquez, 2007) containingcontaining: 10 mM EDTA (pH 8.0), 0,1 % SDS, 50 mM Sodium
Phosphate (pH 7.0), 0,1 % Triton X-100, 1 mM 4-MUG (, M9130-Sigma) and incubated at 37 °C for 6 h.
Six 100 µl aliquots from each well were transferred to individual wells in a microtiter plate and the
reaction was stopped by adding 50 µl of stop reagent (1 M Sodium Carbonate) to each well.
Fluorescence was detected by the use of a Wallac 1420 VICTOR2 microplate luminometer (PerkinElmer)
using an excitation wavelength of 365 nm and a filter wavelength of 430 nm. Each experiment was
repeated 3 times.

<u>Confocal laser microscopy of roots and flowers expressing promoter:Venus/YFP-H2B and promoter:</u>
<u>Cerulean-H2B construct</u>

Imaging of 7 days-old roots was performed on a LSM 880 Airyscan confocal microscope equipped with
two flanking PMTs and a central 32 array GaAsP detector. Images were acquired with Zeiss Plan-
Apochromat 20x/0.8 WD=0.55 M27 objective and excited with laser light of 405 nm, 488 nm and 561
nm. Roots were stained by 1 µM propidium iodide for 10 min and washed in dH$_2$O before imaging.
Imaging of flowers was performed on an Andor Dragonfly spinning disk confocal using an EMCCD iXon
Ultra detector. Images were acquired with a 10x plan Apo NA 0,45 dry objective and excited with laser
light of 405 nm, 488 nm and 561 nm. Maximum intensity projections of z-stacks were acquired with
step size of 1,47 µm. Image processing was performed in FIJI 51. These steps are: background
subtraction, gaussian blur/smooth, brightness/contrast. Imaging was performed at the NorMIC Imaging
platform.

<u>Calcium imaging using the R-GECO1 sensor</u>

$[Ca^{2+}]_{cyt}$ in roots were detected using WT plants expressing the cytosolic localized single-fluorophore

based Ca^{2+} sensor, R-GECO1 (Keinath et al., 2015). Measurements were performed using a confocal

laser scanning microscopy Leica TCS SP8 STED 3X using a 20x multi NA 0.75 objective. Images were

recorded with a frame rate of 5 seconds at 400 Hz. Seedling mounting was performed as described in

(Krebs & Schumacher, 2013). The plant tissues were incubated overnight in half strength MS, 21°C and

continuous light conditions before the day of imaging. $[Ca^{2+}]_{cyt}$ in the abscission zone were detected

using WT plants expressing R-GECO1 (Keinath et al., 2015). Abscission zones of position 7 were used.

Mounting of the abscission zones were performed using the same device as for seedlings (Krebs &

Schumacher, 2013). The abscission zones were incubated in half strength MS medium for 1 h before

imaging. Measurements were performed with a Zeiss LSM880 Airyscan using a Plan-Apochromat 10×

air NA 0.30 objective. Images were recorded with a frame rate of 10 seconds. R-GECO1 was excited

with a white light laser at 561 nm and its emission was detected at 590 nm to 670 nm using a HyD

detector. Laser power and gain settings were chosen for each experiment to maintain comparable

intensity values. For mIDA and flg22 two-fold concentrations were prepared in half strength MS. mIDA

or flg22 were added in a 1:1 volume ratio to the imaging chamber (final concentration 1 µM). ATP was

prepared in a 100-fold concentration in half strength MS and added as a last treatment in a 1:100

volume ratio (final concentration 1 mM) to the imaging chamber as a positive control for activity of the

R-GECO1 sensor (Movie 3). Image processing was performed in FIJI. These steps are: background

subtraction, gaussian blur, MultiStackReg v1.45 (http://bradbusse.net/sciencedownloads.html), 32-bit

conversion, threshold. Royal was used as a look up table. Fluorescence intensities of indicated ROIs were

obtained from the 32-bit images (Krebs & Schumacher, 2013). Normalization was done using the

following formula $\Delta F/F = (F-F0)/F0$ where F0 represents the mean of at least 1 min measurement

without any treatment. R-GECO1 measurements were performed at the Center for Advanced imaging

(CAi) at HHU and at NorMIC Imaging platform at the University of Oslo.

<u>Calcium measurements using the Aequorin (pMAQ2) sensor</u>

$[Ca^{2+}]_{cyt}$ in seedlings and flowers were detected using WT plants expressing p35S-apoaequorin (pMAQ2)

located to cytosol (Aeq) (Knight et al., 1991; Ranf et al., 2011). Aequorin luminescence was measured

as previously described (Ranf et al., 2012). Emitted light was detected by the use of a Wallac 1420

VICTOR2 microplate luminometer (PerkinElmer). Differences in Aeq expression levels due to seedling

size and expression of sensor were corrected by using luminescence at specific time point (L)/Max

Luminescence (Lmax). Lmax was measured after peptide treatment by individually adding 100 µL 2 M

$CaCl_2$ to each well and measuring luminescence constantly for 180 seconds (Ranf et al., 2012). 2 M $CaCl_2$

disrupts the cells and releases the Aeq sensor into the solution where it will react with Ca^{2+} and release

the total possible response in the sample (Lmax) in form of a luminescent peak. A final concentration of

1 µM mIDA was added to each wells at the start of measurements. For inhibitor treatements, Aeq-

seedlings were incubated in in 2mM EGTA (Sigma-Aldrich) or 1mM LaCL$_3$ (Sigma-Aldrich) O/N before

measurements. For seedlings 3 independent experiments were performed with 12 replications in each

experiment. For flowers 3 independent experiments were performed with 4-6 replications in each

experiment.

<u>Measurements of reactive oxygen species (ROS)</u>

ROS production was monitored by the use of a luminol-dependent assay as previously described

(Butenko et al., 2014) using a Wallac 1420 VICTOR2 microplate luminometer (PerkinElmer). Arabidopsis

leaves expressing *35S:HAE:YFP* were cut into leaf discs and incubated in water overnight before

measurements. A final concentration of 1 µM mIDA was added to each well at the start of

measurements. All measurements were performed on 6 leaf discs and each experiment was repeated

3 times.

<u>ROS stain (H2DCF-DA)</u>

Flowers at position 6 were gently incubated in staining solution (25 µM (2',7'-dichlorodihydrofluorescein

diacetate) (H2DCF-DA) (Sigma-Aldrich, D6883), 50 mM KCL, 10 mM MES) for 10 min and further washed

3 times in wash solution (50 mM KCL, 10 mM MES). For the *hae hsl2* mutant the floral organs were

forcibly removed immediately before imaging. Imaging was done using a Dragonfly Airy scan spinning

disk confocal microscope, excited by a 488 nm laser. A total of 9 flowers per genotype were imaged.

The experiment was repeated 2 independent times.

<u>Callose deposition staining and quantification</u>

Callose deposition experiments were conducted as per (Luna et al., 2011) with modifications. Seeds

from WT and *fls2* were vapor-phase sterilized for 4 h. Seeds were transferred to 12-well plates

containing 1 mL of filter-sterilized MS medium and 0. 5 % MES with a final pH of 5.7, and stratified in

the dark at 4 °C for 2 days. Plates were consecutively transferred to a growth chamber with 16 h of light

(150µE m-2 s-1)/8h darkness cycle at 22°C. On the seventh day, growth medium was refreshed under

sterile conditions. On the eight day the treatments were applied: addition of 10 µL of water (mock); 10

µL of 100 µM flg22; or 10 µL of 100 µM mIDA (final peptide concentrations of 1 µM). After 24 h the

growth medium was replaced with 9 5% ethanol and incubated overnight. Ethanol was replaced with

an 8 M NaOH softening solution and incubated for 2 h. Seedlings were washed 3 times in water and

immediately transferred to the aniline blue staining solution (100 mM KPO4-KOH, pH 11; 0.1 % aniline

blue). After 2 h of staining, the seedlings were mounted on 50 % glycerol and imaged under a ZEISS

epifluorescence microscope with UV filter (BP 365/12 nm; FT 395 nm; LP 397 nm). Image analysis was

performed on FIJI (Schindelin et al., 2012). In brief, cotyledons were cropped and measured the leaf

area. Images were thresholded to remove autofluorescence and area of depositions measured to

calculate the ratio of callose deposition area per cotyledon area unit. Between 9-12 seedlings were

analyzed per genotype x treatment combination.

<u>Real time quantitative PCR (RT-qPCR)</u>

Seven days-old Arabidopsis seedlings (WT, *hae hsl2*) grown vertically on ½ sucrose MS-2 plates were

transferred to liquid ½ MS-2 medium (non-treated) and liquid ½ MS-2 medium containing 1 µM peptide

and incubated in growth chambers for 1 h or 12 h. Seedlings were flash-frozen in liquid nitrogen before

total RNA was extracted using SpectrumTM Plant Total RNA Kit (SIGMA Aldrich). cDNA synthesis was

performed as previously described (Grini et al., 2009). RT-qPCR was performed according to protocols

provided by the manufacturer using FastStart Essential DNA Green Master (Roche) and LightCycler96

(Roche) instrument. ACTIN2 was used to normalize mRNA levels as described in (Grini et al., 2009). Three

biological replicates and 4 technical replicates including standard curves were performed for each

sample.

<u>Petal Break strength (pBS)</u>

The force required to remove a petal at a given position on the inflorescence was measured in gram

equivalents using a load transducer as previously described (Stenvik et al., 2008). Plants were grown

until they had at least 15 positions on the inflorescence. A minimum of 15 petals per position were

measured. pBS measurements were performed on WT, *rbohd rbohf* (SALK_070610 SALK_059888) and

hae hsl2 (SALK_021905 SALK_030520) plants.

<u>Statistical methods</u>

Two tailed students t-test ($p < 0.05$) was used to identify significant differences in the fluorescent GUS

assay by comparing treated samples to untreated samples of the same plant line. Statistical analysis of

the RT-qPCR results was performed on all replicas using ONE-WAY or TWO-WAY ANOVA (as stated in

the figure text) and post-hoc Tukey's test ($p < 0.05$). Two tailed students t-test with ($p < 0.05$) was used

in the petal break strength (pBS) measurements to identify significant differences from WT at a given

position on the inflorescence.

<u>**Acknowledgements**</u>

We thank M.K. Anker, I.M. Stø, V. Iversen and R. Falleth for technical assistance in the laboratory and

phytotrone. We thank the NorMic Imaging platform for the use and technical support. This work was

supported by the Research Council of Norway (grant 230849) to V.O. and M.A.B. Work by R. Simon and

and M.B. was supported through CEPLAS.

<u>Author contributions</u>

V.O. generated Arabidopsis lines and constructs, tested IDA expression to biotic and abiotic stress,

performed gene expression studies, phenotypic analysis of mutants and ROS measurements. V.O. and

M.B. performed Ca^{2+} measurements S.G-T. performed the callose deposition assays. V.O., S. G-T., M.B.,

R.Simon and M.A.B designed experiments, analyzed data, and drafted the manuscript. V.O. and M.A.B

wrote the paper with input from all authors.

<u>Competing interests</u>

The authors have no competing interests.

<u>Materials and correspondence</u>

Correspondence to Melinka A. Butenko, m.a.butenko@ibv.uio.no

Supplementary Tables

Supplementary Table 1: Peptide sequences

Peptide	Amino acid sequence
mIDA	PIPPSA**o**SKRHN
flg22	QRLSTGSRINSAKDDAAGLQIA
IDA^{ΔN69}	PIPPSA**o**SKRH

602   (**o** = hydroxyproline)

604 ## Supplementary Table 2: Primers sequences and function

Function	Primer name	Primer sequence 5'-3'
Cloning promoter FLS2	PromFLS2 -988bp Attb1	5'GGGGACAAGTTTGTACAAAAAAGCAGGCTTA GAAGTTGTGAATTGTGAT'3
Cloning promoter FLS2	PromFLS2 -988bp Attb2	5'GGGGACCACTTTGTACAAGAAAGCTGGGTA GGTTTAGACTTTAGAAGA'3
Genotype hsl2	Hsl2 LP	5'CGTCTTGAGCTAGCCAACAAC'3
Genotype hsl2	Hsl2 RP	5'GTCCAATCAAGTGGAGAAACG'3
Genotype hae	Haesa LP	5'CACCTTCCTTCTCTCCATTCC'3
Genotype hae	Haesa RP	5'GTTCGAGAAGTGACAAGCGAG'3
Genotype SALK-lines	Lbb1-	5'GCGTGGACCGCTTGCTGCAACT'3
Genotype rbohd SALK_070610C	LP_SALK_070610C	5'TTTCAACGCCTTTTGGTACAC'3
Genotype rbohd SALK_070610C	RP_SALK_070610C	5'GTTACCTATTCTTTTGCCGGG'3
Genotype rbohf SALK_059888	LP_SALK_059888	5'CAAAGAGCTCTTCGTGGTTTG'3
Genotype rbohf SALK_059888	RP_SALK_059888	5'TCTCTATTGTATCTTGTGTCACCG'3
Genotype fls2 SALK_062054	FLS2_SALK_062054 Fw	5'GGTTCGATTCCTTCTGGAATC'3
Genotype fls2 SALK_062054	FLS2_SALK_062054 Rv	5'CCTGAGTTTTTGAAGCTTCCC'3
qPCR	33.FRK1 Fw	5'AACTTAGGAGACTATTTGGCAGGTAA'3
qPCR	33.FRK1 Rv	5'TGCATCTAATGATATCTTCAACCTCT'3
qPCR	63.PEP3 Fw	5'GCGAGGAAGATGAGAGTATCG'3
qPCR	63.PEP3 Rv	5'TCAATGGTCATGCCATCTTCT'3
qPCR	91.MYB51 Fw	5'GGCCAATTATCTTAGACCTGACA'3
qPCR	91.MYB51 Rv	5'CCACGAGCTATAGCAGACCATT'3
qPCR reference gene	act2int2 sense	5'CCCTGAGGAGCACCCAGTTCTACTC'3

	act2int2 antisense	5'CCGCAAGATCAAGACGAAGGATAGC'3
qPCR reference gene		

References:

Agustí, J., Merelo, P., Cercós, M., Tadeo, F. R., & Talón, M. (2009). Comparative transcriptional survey between laser-microdissected cells from laminar abscission zone and petiolar cortical tissue during ethylene-promoted abscission in citrus leaves. *BMC Plant Biology*, *9*(1), 127. https://doi.org/10.1186/1471-2229-9-127

Asai, T., Tena, G., Plotnikova, J., Willmann, M. R., Chiu, W. L., Gomez-Gomez, L., Boller, T., Ausubel, F. M., & Sheen, J. (2002). MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature*, *415*(6875), 977-983. https://doi.org/10.1038/415977a

Birkenbihl, R. P., Kracher, B., & Somssich, I. E. (2017). Induced Genome-Wide Binding of Three Arabidopsis WRKY Transcription Factors during Early MAMP-Triggered Immunity. *Plant Cell*, *29*(1), 20-38. https://doi.org/10.1105/tpc.16.00681

Blazquez, M. (2007). Quantitative GUS Activity Assay in Intact Plant Tissue. *CSH Protoc*, *2007*, pdb.prot4688. https://doi.org/10.1101/pdb.prot4688

Bleecker, A. B., & Patterson, S. E. (1997). Last exit: senescence, abscission, and meristem arrest in Arabidopsis. *Plant Cell*, *9*(7), 1169-1179. https://doi.org/10.1105/tpc.9.7.1169

Breiden, M., Olsson, V., Blümke, P., Schlegel, J., Gustavo-Pinto, K., Dietrich, P., Butenko, M. A., & Simon, R. (2021). The Cell Fate Controlling CLE40 Peptide Requires CNGCs to Trigger Highly Localized Ca2+ Transients in Arabidopsis thaliana Root Meristems. *Plant Cell Physiol*, *62*(8), 1290-1301. https://doi.org/10.1093/pcp/pcab079

Butenko, M. A., Patterson, S. E., Grini, P. E., Stenvik, G.-E., Amundsen, S. S., Mandal, A., & Aalen, R. B. (2003). *INFLORESCENCE DEFICIENT IN ABSCISSION* Controls Floral Organ Abscission in Arabidopsis and Identifies a Novel Family of Putative Ligands in Plants. *Plant Cell*, *15*, 2296-2307.

Butenko, M. A., Wildhagen, M., Albert, M., Jehle, A., Kalbacher, H., Aalen, R. B., & Felix, G. (2014). Tools and Strategies to Match Peptide-Ligand Receptor Pairs. *Plant Cell*, *26*(5), 1838-1847. https://doi.org/10.1105/tpc.113.120071

Cai, S., & Lashbrook, C. C. (2008). Stamen abscission zone transcriptome profiling reveals new candidates for abscission control: enhanced retention of floral organs in transgenic plants overexpressing Arabidopsis ZINC FINGER PROTEIN2. *Plant Physiol*, *146*(3), 1305-1321. https://doi.org/10.1104/pp.107.110908

Castro, B., Citterico, M., Kimura, S., Stevens, D. M., Wrzaczek, M., & Coaker, G. (2021). Stress-induced reactive oxygen species compartmentalization, perception and signalling. *Nat Plants*, *7*(4), 403-412. https://doi.org/10.1038/s41477-021-00887-0

Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nurnberger, T., Jones, J. D., Felix, G., & Boller, T. (2007). A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature*, *448*(7152), 497-500. https://doi.org/10.1038/nature05999

Cho, S. K., Larue, C. T., Chevalier, D., Wang, H., Jinn, T. L., & Zhang, S. (2008). Regulation of floral organ abscission in Arabidopsis thaliana. *Proc Natl Acad Sci USA.*, *105*. https://doi.org/10.1073/pnas.0805539105

Clough, S. J., & Bent, A. F. (1998). Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *The Plant Journal*, *16*(6), 735-743.

Crick, J., Corrigan, L., Belcram, K., Khan, M., Dawson, J. W., Adroher, B., Li, S., Hepworth, S. R., & Pautot, V. (2022). Floral organ abscission in Arabidopsis requires the combined activities of three TALE homeodomain transcription factors. *J Exp Bot*, *73*(18), 6150-6169. https://doi.org/10.1093/jxb/erac255

Dubiella, U., Seybold, H., Durian, G., Komander, E., Lassig, R., Witte, C. P., Schulze, W. X., & Romeis, T. (2013). Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. *Proc Natl Acad Sci U S A*, *110*(21), 8744-8749. https://doi.org/10.1073/pnas.1221294110

Earley, K. W., Haag, J. R., Pontes, O., Opper, K., Juehne, T., Song, K., & Pikaard, C. S. (2006). Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J*, *45*(4), 616-629. https://doi.org/10.1111/j.1365-313X.2005.02617.x

Felix, G., Duran, J. D., Volko, S., & Boller, T. (1999). Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. *The Plant Journal*, *18*(3), 265-276. https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-313X.1999.00265.x?sid=nlm%3Apubmed

Frerigmann, H., Pislewska-Bednarek, M., Sanchez-Vallet, A., Molina, A., Glawischnig, E., Gigolashvili, T., & Bednarek, P. (2016). Regulation of Pathogen-Triggered Tryptophan Metabolism in Arabidopsis thaliana by MYB Transcription Factors and Indole Glucosinolate Conversion Products. *Mol Plant*, *9*(5), 682-695. https://doi.org/10.1016/j.molp.2016.01.006

Geiger, D., Scherzer, S., Mumm, P., Marten, I., Ache, P., Matschi, S., Liese, A., Wellmann, C., Al-Rasheid, K., & Grill, E. (2010). Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca2+ affinities. *Proceedings of the National Academy of Sciences*, *107*(17), 8023-8028.

Gómez-Gómez, L., Felix, G., & Boller, T. (1999). A single locus determines sensitivity to bacterial flagellin in Arabidopsis thaliana. *Plant J*, *18*(3), 277-284. https://doi.org/10.1046/j.1365-313x.1999.00451.x

Grini, P. E., Thorstensen, T., Alm, V., Vizcay-Barrena, G., Windju, S. S., Jorstad, T. S., Wilson, Z. A., & Aalen, R. B. (2009). The ASH1 HOMOLOG 2 (ASHH2) histone H3 methyltransferase is required for ovule and anther development in Arabidopsis. *PLoS One*, *4*(11), e7817. https://doi.org/10.1371/journal.pone.0007817

Gronnier, J., Franck, C. M., Stegmann, M., DeFalco, T. A., Abarca, A., von Arx, M., Dünser, K., Lin, W., Yang, Z., Kleine-Vehn, J., Ringli, C., & Zipfel, C. (2022). Regulation of immune receptor kinase plasma membrane nanoscale organization by a plant peptide hormone and its receptors. *Elife*, *11*. https://doi.org/10.7554/eLife.74162

He, P., Shan, L., Lin, N. C., Martin, G. B., Kemmerling, B., Nurnberger, T., & Sheen, J. (2006). Specific bacterial suppressors of MAMP signaling upstream of MAPKKK in Arabidopsis innate immunity. *Cell*, *125*(3), 563-575. https://doi.org/10.1016/j.cell.2006.02.047

Huffaker, A., Pearce, G., & Ryan, C. A. (2006). An endogenous peptide signal in Arabidopsis activates components of the innate immune response. *Proc Natl Acad Sci U S A*, *103*(26), 10098-10103. https://doi.org/10.1073/pnas.0603727103

Kadota, Y., Shirasu, K., & Zipfel, C. (2015). Regulation of the NADPH Oxidase RBOHD During Plant Immunity. *Plant and Cell Physiology*, *56*(8), 1472-1480. https://doi.org/10.1093/pcp/pcv063

Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, J. D., Shirasu, K., Menke, F., Jones, A., & Zipfel, C. (2014). Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. *Mol Cell*, *54*(1), 43-55. https://doi.org/10.1016/j.molcel.2014.02.021

Keinath, N. F., Waadt, R., Brugman, R., Schroeder, J. I., Grossmann, G., Schumacher, K., & Krebs, M. (2015). Live Cell Imaging with R-GECO1 Sheds Light on flg22- and Chitin-Induced Transient [Ca(2+)]cyt Patterns in Arabidopsis. *Mol Plant*, *8*(8), 1188-1200. https://doi.org/10.1016/j.molp.2015.05.006

Kissoudis, C., Sunarti, S., van de Wiel, C., Visser, R. G., van der Linden, C. G., & Bai, Y. (2016). Responses to combined abiotic and biotic stress in tomato are governed by stress intensity and resistance mechanism. *J Exp Bot*, *67*(17), 5119-5132. https://doi.org/10.1093/jxb/erw285

Knight, H., Trewavas, A. J., & Knight, M. R. (1997). Calcium signalling in Arabidopsis thaliana responding
to drought and salinity. *Plant J*, *12*(5), 1067-1078. https://doi.org/10.1046/j.1365-
313x.1997.12051067.x
Knight, M. R., Campbell, A. K., Smith, S. M., & Trewavas, A. J. (1991). Transgenic plant aequorin reports
the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature*, *352*(6335),
524-526. https://doi.org/10.1038/352524a0
Krebs, M., & Schumacher, K. (2013). Live cell imaging of cytoplasmic and nuclear Ca2+ dynamics in
Arabidopsis roots. *Cold Spring Harbor Protocols*, *2013*(8), pdb. prot073031.
Kudla, J., Becker, D., Grill, E., Hedrich, R., Hippler, M., Kummer, U., Parniske, M., Romeis, T., &
Schumacher, K. (2018). Advances and current challenges in calcium signaling. *New Phytol*,
218(2), 414-431. https://doi.org/10.1111/nph.14966
Kumpf, R. P., Shi, C. L., Larrieu, A., Sto, I. M., Butenko, M. A., Peret, B., Riiser, E. S., Bennett, M. J., &
Aalen, R. B. (2013). Floral organ abscission peptide IDA and its HAE/HSL2 receptors control cell
separation during lateral root emergence. *Proc Natl Acad Sci U S A*, *110*(13), 5235-5240.
https://doi.org/10.1073/pnas.1210835110
Kärkönen, A., & Kuchitsu, K. (2015). Reactive oxygen species in cell wall metabolism and development
in plants. *Phytochemistry*, *112*, 22-32.
https://doi.org/https://doi.org/10.1016/j.phytochem.2014.09.016
Ladwig, F., Dahlke, R. I., Stuhrwohldt, N., Hartmann, J., Harter, K., & Sauter, M. (2015). Phytosulfokine
Regulates Growth in Arabidopsis through a Response Module at the Plasma Membrane That
Includes CYCLIC NUCLEOTIDE-GATED CHANNEL17, H+-ATPase, and BAK1. *Plant Cell*, *27*(6),
1718-1729. https://doi.org/10.1105/tpc.15.00306
Lee, Y., Yoon, T. H., Lee, J., Jeon, S. Y., Lee, J. H., Lee, M. K., Chen, H., Yun, J., Oh, S. Y., Wen, X., Cho, H.
K., Mang, H., & Kwak, J. M. (2018). A Lignin Molecular Brace Controls Precision Processing of
Cell Walls Critical for Surface Integrity in Arabidopsis. *Cell*, *173*(6), 1468-1480.e1469.
https://doi.org/10.1016/j.cell.2018.03.060
Li, L., Li, M., Yu, L., Zhou, Z., Liang, X., Liu, Z., Cai, G., Gao, L., Zhang, X., Wang, Y., Chen, S., & Zhou, J.
M. (2014). The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD
to control plant immunity. *Cell Host Microbe*, *15*(3), 329-338.
https://doi.org/10.1016/j.chom.2014.02.009
Luna, E., Pastor, V., Robert, J., Flors, V., Mauch-Mani, B., & Ton, J. (2011). Callose deposition: a
multifaceted plant defense response. *Mol Plant Microbe Interact*, *24*(2), 183-193.
https://doi.org/10.1094/mpmi-07-10-0149
Ma, Y., Walker, R. K., Zhao, Y., & Berkowitz, G. A. (2012). Linking ligand perception by PEPR pattern
recognition receptors to cytosolic Ca2+ elevation and downstream immune signaling in plants.
Proc Natl Acad Sci U S A, *109*(48), 19852-19857. https://doi.org/10.1073/pnas.1205448109
Matsubayashi, Y. (2011). Post-translational modifications in secreted peptide hormones in plants.
Plant Cell Physiol, *52*(1), 5-13. https://doi.org/10.1093/pcp/pcq169
McKenna, J. F., Rolfe, D. J., Webb, S. E. D., Tolmie, A. F., Botchway, S. W., Martin-Fernandez, M. L.,
Hawes, C., & Runions, J. (2019). The cell wall regulates dynamics and size of plasma-membrane
nanodomains in Arabidopsis. *Proc Natl Acad Sci U S A*, *116*(26), 12857-12862.
https://doi.org/10.1073/pnas.1819077116
Meng, X., Zhou, J., Tang, J., Li, B., de Oliveira, M. V., Chai, J., He, P., & Shan, L. (2016). Ligand-Induced
Receptor-like Kinase Complex Regulates Floral Organ Abscission in Arabidopsis. *Cell Rep*, *14*(6),
1330-1338. https://doi.org/10.1016/j.celrep.2016.01.023
Meng, X., Zhou, J., Tang, J., Li, B., de Oliveira, M. V. V., Chai, J., He, P., & Shan, L. (2016). Ligand-Induced
Receptor-like Kinase Complex Regulates Floral Organ Abscission in Arabidopsis. *Cell reports*,
14(6), 1330-1338. https://doi.org/10.1016/j.celrep.2016.01.023
Monshausen, G. B., Messerli, M. A., & Gilroy, S. (2008). Imaging of the Yellow Cameleon 3.6 indicator
reveals that elevations in cytosolic Ca2+ follow oscillating increases in growth in root hairs of
Arabidopsis. *Plant Physiol*, *147*(4), 1690-1698. https://doi.org/10.1104/pp.108.123638

Niederhuth, C., Patharkar, O. R., & Walker, J. (2013). Transcriptional profiling of the Arabidopsis abscission mutant hae hsl2 by RNA-Seq. *BMC Genomics*, *14*(1), 37. http://www.biomedcentral.com/1471-2164/14/37

Olsson, V., Joos, L., Zhu, S., Gevaert, K., Butenko, M. A., & De Smet, I. (2018). Look Closely, the Beautiful May Be Small: Precursor-Derived Peptides in Plants. *Annu Rev Plant Biol*. https://doi.org/10.1146/annurev-arplant-042817-040413

Patharkar, O. R., Gassmann, W., & Walker, J. C. (2017). Leaf shedding as an anti-bacterial defense in Arabidopsis cauline leaves. *PLoS Genet*, *13*(12), e1007132. https://doi.org/10.1371/journal.pgen.1007132

Patharkar, O. R., & Walker, J. C. (2015). Floral organ abscission is regulated by a positive feedback loop. *Proc Natl Acad Sci U S A*, *112*(9), 2906-2911. https://doi.org/10.1073/pnas.1423595112

Patharkar, O. R., & Walker, J. C. (2016). Core Mechanisms Regulating Developmentally Timed and Environmentally Triggered Abscission. *Plant Physiol*, *172*(1), 510-520. https://doi.org/10.1104/pp.16.01004

Poncini, L., Wyrsch, I., Dénervaud Tendon, V., Vorley, T., Boller, T., Geldner, N., Métraux, J. P., & Lehmann, S. (2017). In roots of Arabidopsis thaliana, the damage-associated molecular pattern AtPep1 is a stronger elicitor of immune signalling than flg22 or the chitin heptamer. *PLoS One*, *12*(10), e0185808. https://doi.org/10.1371/journal.pone.0185808

Qi, Z., Verma, R., Gehring, C., Yamaguchi, Y., Zhao, Y., Ryan, C. A., & Berkowitz, G. A. (2010). Ca2+ signaling by plant Arabidopsis thaliana Pep peptides depends on AtPepR1, a receptor with guanylyl cyclase activity, and cGMP-activated Ca2+ channels. *Proc Natl Acad Sci U S A*, *107*(49), 21193-21198. https://doi.org/10.1073/pnas.1000191107

Ranf, S., Eschen-Lippold, L., Pecher, P., Lee, J., & Scheel, D. (2011). Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns. *Plant J*, *68*(1), 100-113. https://doi.org/10.1111/j.1365-313X.2011.04671.x

Ranf, S., Grimmer, J., Poschl, Y., Pecher, P., Chinchilla, D., Scheel, D., & Lee, J. (2012). Defense-related calcium signaling mutants uncovered via a quantitative high-throughput screen in Arabidopsis thaliana. *Mol Plant*, *5*(1), 115-130. https://doi.org/10.1093/mp/ssr064

Robatzek, S., Chinchilla, D., & Boller, T. (2006). Ligand-induced endocytosis of the pattern recognition receptor FLS2 in Arabidopsis. *Genes Dev*, *20*(5), 537-542. https://doi.org/10.1101/gad.366506

Roberts, J. A., Whitelaw, C. A., Gonzalez-Carranza, Z. H., & McManus, M. T. (2000). Cell Separation Processes in Plants- Models, Mechanisms and Manipulation. *Annals of Botany*, *86*, 223-235.

Rzemieniewski, J., Leicher, H., Lee, H. K., Broyart, C., Nayem, S., Wiese, C., Maroschek, J., Camgöz, Z., Lalun, V. O., Djordjevic, M. A., Vlot, A. C., Hückelhoven, R., Santiago, J., & Stegmann, M. (2022). Phytocytokine signaling integrates cell surface immunity and nitrogen limitation. *bioRxiv*, 2022.2012.2020.521212. https://doi.org/10.1101/2022.12.20.521212

Sanders, D., Brownlee, C., & Harper, J. F. (1999). Communicating with calcium. *Plant Cell*, *11*(4), 691-706. https://doi.org/10.1105/tpc.11.4.691

Santiago, J., Brandt, B., Wildhagen, M., Hohmann, U., Hothorn, L. A., & Butenko, M. A. (2016). Mechanistic insight into a peptide hormone signaling complex mediating floral organ abscission. *Elife.*, *5*. https://doi.org/10.7554/eLife.15075

Scherzer, S., Maierhofer, T., Al-Rasheid, K. A. S., Geiger, D., & Hedrich, R. (2012). Multiple calcium-dependent kinases modulate ABA-activated guard cell anion channels. *Mol Plant*, *5*(6), 1409-1412. https://doi.org/10.1093/mp/sss084

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods*, *9*(7), 676-682. https://doi.org/10.1038/nmeth.2019

Shi, C.-L., von Wangenheim, D., Herrmann, U., Wildhagen, M., Kulik, I., Kopf, A., Ishida, T., Olsson, V., Anker, M. K., Albert, M., Butenko, M. A., Felix, G., Sawa, S., Claassen, M., Friml, J., & Aalen, R.

B. (2018). The dynamics of root cap sloughing in Arabidopsis is regulated by peptide signalling. *Nat Plants*, *4*(8), 596-604. https://doi.org/10.1038/s41477-018-0212-z

Smakowska-Luzan, E., Mott, G. A., Parys, K., Stegmann, M., Howton, T. C., Layeghifard, M., Neuhold, J., Lehner, A., Kong, J., Grunwald, K., Weinberger, N., Satbhai, S. B., Mayer, D., Busch, W., Madalinski, M., Stolt-Bergner, P., Provart, N. J., Mukhtar, M. S., Zipfel, C., . . . Belkhadir, Y. (2018). An extracellular network of Arabidopsis leucine-rich repeat receptor kinases. *Nature*, *553*(7688), 342-346. https://doi.org/10.1038/nature25184

Somssich, M., Bleckmann, A., & Simon, R. (2016). Shared and distinct functions of the pseudokinase CORYNE (CRN) in shoot and root stem cell maintenance of Arabidopsis. *J Exp Bot*, *67*(16), 4901-4915. https://doi.org/10.1093/jxb/erw207

Steinhorst, L., & Kudla, J. (2013). Calcium and reactive oxygen species rule the waves of signaling. *Plant Physiol*, *163*(2), 471-485. https://doi.org/10.1104/pp.113.222950

Stenvik, G. E., Tandstad, N. M., Guo, Y., Shi, C. L., Kristiansen, W., & Holmgren, A. (2008). The EPIP peptide of INFLORESCENCE DEFICIENT IN ABSCISSION is sufficient to induce abscission in Arabidopsis through the receptor-like kinases HAESA and HAESA-LIKE2. *Plant Cell.*, *20*. https://doi.org/10.1105/tpc.108.059139

Sto, I. M., Orr, R. J., Fooyontphanich, K., Jin, X., Knutsen, J. M., Fischer, U., Tranbarger, T. J., Nordal, I., & Aalen, R. B. (2015). Conservation of the abscission signaling peptide IDA during Angiosperm evolution: withstanding genome duplications and gain and loss of the receptors HAE/HSL2. *Front Plant Sci*, *6*, 931. https://doi.org/10.3389/fpls.2015.00931

Taylor, I., & Walker, J. C. (2018). Transcriptomic evidence for distinct mechanisms underlying abscission deficiency in the Arabidopsis mutants haesa/haesa-like 2 and nevershed. *BMC Res Notes*, *11*(1), 754. https://doi.org/10.1186/s13104-018-3864-x

Torres, M. A., & Dangl, J. L. (2005). Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Curr Opin Plant Biol*, *8*(4), 397-403. https://doi.org/10.1016/j.pbi.2005.05.014

Vie, A. K., Najafi, J., Liu, B., Winge, P., Butenko, M. A., Hornslien, K. S., Kumpf, R., Aalen, R. B., Bones, A. M., & Brembu, T. (2015). The IDA/IDA-LIKE and PIP/PIP-LIKE gene families in Arabidopsis: phylogenetic relationship, expression patterns, and transcriptional effect of the PIPL3 peptide. *Journal of Experimental Botany*, *66*(17), 5351-5365. https://doi.org/10.1093/jxb/erv285

Wang, Y., Li, X., Fan, B., Zhu, C., & Chen, Z. (2021). Regulation and Function of Defense-Related Callose Deposition in Plants. *Int J Mol Sci*, *22*(5). https://doi.org/10.3390/ijms22052393

Main figures - A dual function of the IDA peptide in regulating cell

 separation and modulating plant immunity at the molecular level

Fig. 1: mIDA-induced [Ca²⁺]cyt release in Arabidopsis roots correlates with *pHAE* and *pHSL2* activity

a, Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI) (upper panel, outlined in red) in the meristematic and elongation zone of the root. Fluorescence intensities (ΔF/F) over time of the whole root represented in a heat map (lower panel). Shown are cytosolic calcium concentration ([Ca²⁺]cyt) dynamics in the ROI in response to 1 mM mIDA over time. (see also Movie 1). Red lines at 5 minutes (min) indicates application of mIDA peptide or application of eATP at 22 min. Representative response from 10 roots. The increase in [Ca²⁺]cyt response propagates through the roots as two waves. **b,** For comparison; Normalized R-GECO1 fluorescence intensities (ΔF/F) measured from regions of interest (ROI) (outlined in red, upper panel) in response to 1µM flg22 over time. Fluorescence intensities (ΔF/F) over time of the whole root represented in a heat map (lower panel). Red line at 0 min (min) indicates application of flg22 peptide. Representative response from 10 roots. The increase in [Ca²⁺]cyt response propagates through the roots as a single wave seen as normalized R-GECO1 fluorescence intensities (ΔF/F) shown as a heat map (see also movie 2). **c,d,e,** Expression of the receptors **c,** *pHAE:Venus-H2B* **d,** *pHSL2:Venus-H2B* and **e,** *pFLS2:Venus-H2B* in 7 days-

Fig. 2 : mIDA-induced [Ca²⁺]$_{cyt}$ release in Arabidopsis abscission zones.

a, Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI)

(outlined in red) in floral abscission zone (AZ)s. Shown are cytosolic calcium concentration ([Ca²⁺]$_{cyt}$)

dynamics in the ROI in response to 1 μM mIDA over time (see also Movie 3). Representative response

from 10 flowers. Red lines at 5 minutes (min) indicates application of mIDA peptide or application of

eATP at 35 min (for AZs), respectively. The increase in [Ca²⁺]$_{cyt}$ response propagates through the AZ as a

single wave. **b**,Expression of *pHAE:Venus-H2B* and *pHSL2:Venus-H2B* in flowers at position 6 (See Sup

Fig. 4a for positions) (arrowhead indicates AZ). Representative pictures of n = 8, scale bar = 100 μm,

maximum intensity projections of z-stacks. **b** Expression of *pHAE:Venus-H2B* and *pHSL2:Venus-H2B* in

flower position 6 (See Sup. Fig.4a for positions) (arrowhead indicates AZ). Representative picture of n =

8. Scale bar = 100 μm, maximum intensity projections of z-stacks.

Fig. 3: *IDA* is induced by biotic and abiotic stress

Representative pictures of *pIDA:GUS* expression after 12 h treatment with 1 µM flg22, chitin, NaCl, Mannitol and 1 µM IDAΔN69 in **a,** cells surrounding emerging lateral roots, and **b,** in the main root . **c,** Normalized emitted fluorescence of fluorochrome 4-methyl umbelliferone (4-MU) in 7 days-old seedlings after 12 h treatment with 1 µM flg22, chitin, NaCl, Mannitol and 1 µM of the inactive IDA peptide, IDAΔN69. Normalized to 1 on No Treatment sample. Controls were not subjected to any stimuli (No treatment). n = 6, experiment repeated 3 times. * = significantly different from the non-treated (No treatment) sample (p < 0.05, student t-test, two tailed). Controls were not subjected to any stimuli (No treatment). **a,b,** Representative picture of n = 10, experiment repeated 3 times, scale bar = 50 µm.

Fig. 4: Callose deposition is not induced by mIDA

a, No callose deposition could be detected in Col-0 WT and *fls2* mutants treated with water (mock treatment) or 1 μM mIDA. Callose deposition could be observed in cotyledons of eight day old Col-0 WT seedlings treated with 1 μM flg22. No callose deposition could be detected in *fls2* mutant treated with flg22. Representative images of 9-12 seedlings per genotype. **b**, Total callose deposition area for the different genotypes treated with water (mock treatment), 1 μM mIDA or 1 μM flg22. Statistical significant difference at (p < 1*10^-6). N = 9-12

Fig. 5: mIDA induced expression of defense-associated marker genes

a, Transcripts of *FRK1*, *MYB51*, and *PEP3* in WT Col-0 seedlings exposed to 1 µM mIDA for 1 h (light gray) and to 1 µM mIDA for 12 h (dark gray) compared to untreated tissue (black). RNA levels were measured by RT-qPCR analysis. *ACTIN* was used to normalize mRNA levels. n = 4, three biological replicates Statistical analyses was performed comparing induction times of individual genes using one-way ANOVA and post-hoc Tukey's test ($p < 0.05$). b,*pMYB51:YFPN* expression is enhanced in roots after 7 h exposure to 1 µM mIDA peptide compared to untreated roots (control), scale bar = 50 µm, maximum intensity projections of z-stacks, magenta = propidium iodide stain.

Fig. 6: mIDA and flg22 co-treatement extremely enhances the expression of defense-associated marker genes

a, RT-qPCR data showing transcription of *FRK1*, *MYB51*, and *PEP3* in Col-0 WT seedlings exposed to 1 μM mIDA, 1 μM flg22 or a combination of 1 μM mIDA and 1 μM flg22 for 1 h compared to untreated seedlings (No treatment control). *ACTIN* was used to normalize mRNA levels. Figure represent 3 biological replicates with 4 technical replicates. Statistical analyses comparing No Treatment to peptide treated samples was performed using one-way ANOVA and post-hoc Tukey's test ($p < 0.05$).

Fig. 7: IDA regulates cell separation processes, but is also involved in major transcription of defense genes upon pathogen attack.

a, IDA and the IDL peptides control cell separation processes during plant development and in response to abiotic and biotic stress (Butenko et al. 2003; Patharkar and Walker 2016; Patharkar et al. 2017; Shi et al. 2018). Tissue undergoing cell separation includes floral organ abscission, cauline leaf abscission, emerging of lateral roots and root cap sloughing. b, During normal conditions, IDA control floral organ abscission and emergence of lateral roots by relaying a signal through receptor complexes including HAE, HSL2, SERK1 and BAK1 to modulate the expression of cell wall remodeling (CWR) genes as well as moderately expression of defense genes. IDA activates a receptor dependent production of ROS and an increase in $[Ca^{2+}]_{cyt}$. b, Upon stress, such as pathogen attack, the activation of HAE and HSL2 acts in addition to activation of defense related receptors, such as FLS2, to enhance the expression of defense related genes significantly. This ensure optimal protection of cells undergoing cell separation, which may be major entry routes during a pathogen attack.

931 # Supplementary figures - A dual function of the IDA peptide in
932 # regulating cell separation and modulating plant immunity at the
933 # molecular level

Sup Fig. 1: IDA induces a production of ROS in Arabidopsis

a, ROS production detected by the luminol-based assay was monitored over time as RLU. ROS

production from *hae hsl2* leaf disks expressing *35S:HAE-YFP* in response to 1 μM mIDA (black) and in

WT control (blue). **b**, Fluorescent image of *hae hsl2* rosette leaves expressing *35S:HAE:YFP*. Signal is

detected in the plasma membrane of cells in the epidermal layer. Representative picture. Scale bar =

50 μm. **c**, *35S:HAE-YFP* complements the *hae hsl2* mutant abscission phenotype of observed. Scale = 1

cm.

Sup Fig. 2 : The IDA triggered increase in $[Ca2+]_{cyt}$ is receptor dependent and is abolished in the

presence of Ca^{2+} inhibitors

a, Increase in $[Ca^{2+}]_{cyt}$ in seedlings expressing the cytosolic localized *Aequorin*-based luminescence Ca^{2+}

sensor (*Aeq*) measured in relative light units (RLU) treated with 1 µM mIDA (black). No response is

observed in *Aeq* seedlings treated with 1 µM IDAΔN69 (gray) or in *hae hsl2 Aeq* seedlings treated with

1 µM mIDA (blue). **b,** Increase in $[Ca^{2+}]_{cyt}$ in seedlings expressing the cytosolic localized *Aequorin*-based

luminescence Ca^{2+} sensor (*Aeq*) measured in relative light units (RLU) treated with 1 µM mIDA . No

response is observed in *Aeq* seedlings treated with 1 mM EGTA (blue) or 1mM $LaCl_3$ (grey). Curves

represent average of 3 independent experiments with 4-6 replicates in each experiment.

Sup Fig. 3: mIDA induces a Ca²⁺ response in flowers

a, Arabidopsis flowers and siliques at different developmental stages. Flowers along the main

inflorescence are counted from the first flower with visible white petals at the top of the inflorescence

and is defined as p 1. Subsequently older flowers along the inflorescence (p0-p8). Cell wall remodeling

in AZ cells, accompanied with a reduction in petal breakstrength (the force required to remove a petal

from the receptacle) is initiated at p6 (red arrow). By p7 AZ cells have undergone cell separation and

the floral organs have abscised. p=position **b**, Increase in $[Ca^{2+}]_{cyt}$ in flowers expressing the cytosolic

localized *Aequorin*-based luminescence Ca²⁺ sensor (*Aeq*) measured in relative light units (RLU) treated

with 1 μM mIDA. A $[Ca^{2+}]_{cyt}$ increase is observed in flowers where there is an initial weakening of the

cell walls (p5-6) (black). No increase in luminescence was observed in *Aeq* expressing flowers of an

earlier developmental stage (p3) (gray). Curves show representative measurements. Three

independent experiments were performed with 4-6 replicates in each experiment.

Sup Fig. 4: mIDAΔN69 does not induce a [Ca²⁺]$_{cyt}$ release in Arabidopsis abscission zones (AZ)

Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI)

(outlined in red) in the floral AZs. Shown are cytosolic calcium concentration ([Ca²⁺]$_{cyt}$) dynamics in the

ROI in response to application of 1 μM mIDAΔN69 or eATP over time. Red lines at 5 minutes (min)

indicates application of mIDAΔN69 peptide or application of eATP at 30 min (see also Movie 4).

Representative response from 10 flowers. The increase in [Ca²⁺]$_{cyt}$ response to eATP propagates

through the AZ as a single wave seen as normalized R-GECO1 fluorescence intensities (ΔF/F) shown as

a heat map.

Sup Fig. 5: Investigation of CNGCs and RBOHs in the abscission process

a, Relative expression of *CNGC1*, *CNGC2*, *CNGC4*, *CNGC5*, *CNGC6*, *CNGC9*, *CNGC12, RBOHD* and *RBOHF*

in the floral organs during the onset of abscission (Cai and lashbrook 2008). X-axis represents the

developmental stages of abscission where position 12 represent flowers where the stamen and petals

are of the same length, position 13 represent flowers at anthesis, position 15 represent flowers with

stigma extended above the anthers. Floral organ abscission occurs at position 15. Relative expression

of *HAE* and *HSL2* added for comparison. Nomenclature for the abscission process and relative

expression values taken from (Cai and Lashbrook 2008). **b,** No floral organ abscission phenotype is

observed in the single *cngc* mutant plants compared to the *hae hsl2* mutant which retains the floral

organ indefinitely. Scale bar = 2cm, 5 weeks old plants. **c,** ROS production in the AZ (white arrow head)

detected by using the fluorescent dye H_2DCF-DA in WT and *rbohd rbohf* flowers at position 6 (See

Supplementary Fig. 5a for positions). Scale bar = 100 µm, maximum intensity projections of z-stacks.

Representative pictures from 9 flowers. **d**, Representative pictures of WT and *rbohd rbohf*

inflorescences, scale bar for inflorescences = 1 cm. **e**, Petal break strength (pBS) measurements of WT

and the *rbohd rbohf* mutant at position 1-6 along the inflorescence. Force required to remove petals

from the receptacle represented in gram equivalent. * = significantly different from WT at the given

position ($p < 0.05$, student t-test, two tailed).

1003

Sup Fig 6: mIDA induced expression of defense-associated marker genes is only partially dependent on

the HAE and HSL2 receptors

RT-qPCR data showing transcription of *FRK1*, *MYB51*, and *PEP3* in Col-0 WT and *hae hsl2* seedlings

exposed to 1 μM mIDA for 1 h compared to untreated seedlings (No treatment). *ACTIN* was used to

normalize mRNA levels. Figure represent 3 biological replicates with 4 technical replicates. Statistical

analyses was performed comparing samples within same gene, using two-way ANOVA and post-hoc

Tukey's test ($p < 0.05$).

1011

1012

1013

Sup Fig. 7: mIDA and flg22 induced expression of *pMYB51:YFP*

Representative microscopic pictures of 7-days-old *pMYB51:YFPN* roots from seedlings transferred to

liquid medium with 1 µM flg22 or 1 µM mIDA for 1, 3, 5 and 7 h using a fluorescent microscope.

Control seedlings were transferred to medium with no peptide and imaged in the same time frame.

Fluorescent nuclei could be observed in a similar temporal pattern in roots subjected to mIDA and

flg22. Scale bar = 100 µm, single plane images using a Zeiss Axioplan2 microscope with an AxioCam

HRc, 20x air objective and a YFP filter (Excitation: 500/20, Beamsplitter: FT515, Emission: 535/50), no

imaging processing was performed, t = time in h. n = 3, experiment repeated 3 times.

1022

1023

1024

Sup Fig. 8: HAE (yellow) and HSL2 (red) display very distinct repertoire of immune and growth-related

interacting LRR-RKs.

HEA and HSL2 specific interacting partners determined using a sensitized high-throughput

extracellular domain interaction assay (Smakowska-Luzan et al,.2018). HSL2 and HAE show distinct

interacting LRR-RKs where only HSL2 was found to interact with RLK7 (red halo). * indicates receptors

known to play a function in biotic and/or abiotic responses in planta based on previous published

work, dark blue nodes refer to bidirectional interactions with GeoMean ≥2.5, light blue nodes- refer to

bidirectional interactions with GeoMean ≤2.5, and grey nodes refer to unidirectional interactions.

1033

1034