1

A dual function of the IDA peptide in regulating cell separation and modulating plant immunity at the molecular level

Vilde Olsson Lalun[1], Maike Breiden[§2], Sergio Galindo-Trigo[1§], Elwira Smakowska-Luzan[3], Rüdiger Simon[3] and Melinka A. Butenko[1*]

[1]Section for Genetics and Evolutionary Biology, Department of Biosciences, University of Oslo, 0316 Oslo, Norway

[2]Institute for Developmental Genetics and Cluster of Excellence on Plant Sciences, Heinrich Heine University, Universitätsstraße 1, 40225 Düsseldorf, Germany

[3]Gregor Mendel Institute (GMI), Austrian Academy of Sciences, Vienna Biocenter (VBC), Dr Bohr-Gasse 3, 1030 Vienna, Austria

[§]Equal contribution

*Corresponding author; email: m.a.butenko@ibv.uio.no

Abstract:

The abscission of floral organs and emergence of lateral roots in Arabidopsis is regulated by the peptide ligand INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) and the receptor protein kinases HAESA (HAE) and HAESA-LIKE 2 (HSL2). During these cell separation processes, the plant induces defense-associated genes to protect against pathogen invasion. However, the molecular coordination between abscission and immunity has not been thoroughly explored. Here we show that IDA induces a release of cytosolic calcium ions (Ca^{2+}) and apoplastic production of reactive oxygen species, which are signatures of early defense responses. In addition, we find that IDA promotes late defense responses by the transcriptional upregulation of genes known to be involved in immunity. When comparing the IDA induced early immune responses to known immune responses, such as those elicited by flagellin22 treatment, we observe both similarities and differences. We propose a molecular mechanism by which IDA promotes signatures of an immune response in cells destined for separation to guard them from pathogen attack.

Introduction

All multicellular organisms require tightly regulated cell-to-cell communication during development and adaptive responses. In addition to hormones, plants use small-secreted peptide ligands to regulate these highly coordinated events of growth, development and responses to abiotic and biotic stress (Matsubayashi, 2011; Olsson et al., 2018). In plants, organ separation or abscission, involves cell separation between specialized abscission zone (AZ) cells and enables the removal of unwanted or diseased organs in response to endogenous developmental signals or environmental stimuli. *Arabidopsis thaliana* (Arabidopsis) abscise floral organs (stamen, petals and sepals) after pollination, and this system has been used as a model to study the regulation of cell separation and abscission (Bleecker & Patterson, 1997). Precise cell separation also occurs during the emergence of new lateral root (LR) primordia, root cap sloughing, formation of stomata and radicle emergence during germination (Roberts et al., 2000). In Arabidopsis, the INFLORESCENCE DEFICIENT IN ABSCISSION (IDA) peptide mediated signaling system ensures the correct spatial and temporal abscission of sepals, petals, and stamens. The IDA protein shares a conserved C-terminal domain with the other 7 IDA-LIKE (IDL) members (Butenko et al., 2003, Vie et al., 2015) which is processed to a functional 12 amino acid hydroxylated peptide (Butenko et al., 2014). The abscission process is regulated by the production and release of IDA, which binds the genetically redundant plasma membrane (PM) localized receptor kinases (RKs), HAESA (HAE) and HAESA-LIKE 2 (HSL2). IDA binding to HAE and HSL2 promotes receptor association with members of the co-receptor SOMATIC EMBRYOGENESIS RECEPTOR KINASE (SERK) family and further downstream signaling leading to cell separation events. (Cho et al., 2008; Meng et al., 2016; Santiago et al., 2016; Stenvik et al., 2008). Recently, it has been shown that IDA and IDL family members also bind and activate HSL1 in the regulation of leaf epidermal pattering, indicating subfunctionalisation within this clade of receptors (Roman et al., 2022) and opens for the possibility that HAE or HSL2 could have additional functions than regulating cell separation events.

A deficiency in the IDA signaling pathway prevents the expression of genes encoding secreted cell wall remodeling and hydrolase enzymes thus hindering floral organs to abscise (Butenko et al., 2003; Cho et al., 2008; Kumpf et al., 2013; Xiangzong Meng et al., 2016; Niederhuth et al., 2013; Stenvik et al., 2008; Isaiah Taylor & John C. Walker, 2018). Interestingly, components of the IDA signaling pathway control different cell separation events during Arabidopsis development. IDA signaling through HAE and HSL2 regulates cell wall remodeling genes in the endodermal, cortical, and epidermal tissues overlaying the LR primordia during LR emergence (Kumpf et al., 2013). In addition, IDL1 signals through HSL2 to regulate cell separation during root cap sloughing (Shi et al., 2018).

Plant cell walls act as physical barriers against pathogenic invaders. The cell-wall processing and remodeling that occurs during abscission leads to exposure of previously protected tissue. This provides an entry point for phytopathogens thereby increasing the need for an effective defense system in these

cells (Agustí et al., 2009). Indeed, it has been shown that AZ cells undergoing cell separation express

defense genes and it has been proposed that they function in protecting the AZ from infection after

abscission has occurred (Cai & Lashbrook, 2008; Niederhuth et al., 2013). IDA elicits the production of

Reactive Oxygen Species (ROS) in *Nicotiana benthamiana* (*N.benthamiana*) leaves transiently expressing

the HAE or HSL2 receptor (Butenko et al., 2014). As a consequence of defense responses, ROS occurs in

the apoplast and is involved in regulating the cell wall structure by producing cross-linking of cell wall

components in the form of hydrogen peroxide (H_2O_2) or act as cell wall loosening agents in the form of

hydroxyl radical ($\cdot$OH)radicals (Kärkönen & Kuchitsu, 2015). ROS may also serve as a direct defense

response by their highly reactive and toxic properties (Kärkönen & Kuchitsu, 2015). In addition, ROS may

act as a secondary signaling molecule enhancing intracellular defense responses as well as contributing

to the induction of defense related genes (Reviewed in (Castro et al., 2021)). In Arabidopsis, apoplastic

ROS is mainly produced by a family of NADPH oxidases located in the plasma membrane known as

RESPIRATORY BURST OXIDASE PROTEINs (RBOHs) (Torres & Dangl, 2005). The production of ROS is often

found closely linked to an increase in the concentration of cytosolic Ca^{2+} ($[Ca^{2+}]_{cyt}$), where the interplay

between these signaling molecules is observed in immunity across kingdoms (Kadota et al., 2015;

Steinhorst & Kudla, 2013). Interestingly, ROS and $[Ca^{2+}]_{cyt}$ are known signaling molecules acting

downstream of several peptide ligand-receptor pairs. This includes the endogenous PAMPs ELICITOR

PEPTIDE(PEP)1, PEP2 and PEP3 which function as elicitors of systemic responses against pathogen attack

and herbivory (Huffaker et al., 2006; Ma et al., 2012; Qi et al., 2010), as well as the pathogen derived

peptide ligands flagellin22 (flg22) and elongation factor thermo unstable(Ef-Tu) which bind and activate

defense related RKs in the PM of the plant cell (Felix et al., 1999; Gómez-Gómez et al., 1999; Ranf et al.,

2011).

The interplay between ROS and $[Ca^{2+}]_{cyt}$ has been well studied in the flg22 induced signaling system. The

flg22 peptide interacts with the extracellular domain of the FLAGELLIN SENSING2 (FLS2) receptor kinase,

which rapidly leads to an increase in $[Ca^{2+}]_{cyt}$ (Ranf et al., 2011). In a resting state, the co-receptor

BRASSINOSTEROID INSENSITIVE1-ASSOCIATED RECEPTOR KINASE1/SERK3) (BAK1) and the cytoplasmic

kinase BOTRYTIS-INDUCED KINASE1 (BIK1) form a complex with FLS2. Upon binding of flg22 to the FLS2

receptor, BIK1 is released from the receptor complex and activates intracellular signaling molecules

through phosphorylation, among other, the PM localized RBOHD (Kadota et al., 2014; Li et al., 2014).

The flg22 induced apoplastic ROS produced binds to PM localized Ca^{2+} channels on neighboring cells,

activating Ca^{2+} influx into the cytosol. The cytosolic Ca^{2+} may bind to CALCIUM DEPENDENT PROTEIN

KINASE5 (CDPKs) regulating RBOHD activity through phosphorylation. The Ca^{2+} dependent

phosphorylation of RBOH enhances ROS production, which in turn leads to a propagation of the

ROS/Ca^{2+} signal through the plant tissue (Dubiella et al., 2013)

A rapid increase in $[Ca^{2+}]_{cyt}$ is not only observed in peptide-ligand induced signaling. Similar $[Ca^{2+}]_{cyt}$

changes are observed as responses to developmental signals, such as extracellular auxin and

environmental signals such as high salt conditions and drought(reviewed in (Kudla et al., 2018)). Also,

$[Ca^{2+}]_{cyt}$ signals are observed in single cells during pollen tube and root hair growth (Monshausen et al.,

2008; Sanders et al., 1999). How cells decipher the $[Ca^{2+}]_{cyt}$ changes into a specific cellular response is

still largely unknown, however the cell contains a large toolkit of Ca^{2+} sensor proteins that may affect

cellular function through changes in protein phosphorylation and gene expression. Each Ca^{2+} binding

protein has specific affinities for Ca^{2+} allowing them to respond to different Ca^{2+} concentrations

providing a specific cellular response (Geiger et al., 2010; Scherzer et al., 2012).

Given our previous studies showing IDA induced apoplastic ROS production (Butenko et al., 2014), and

the known link between ROS and $[Ca^{2+}]_{cyt}$ response observed for other ligands such as flg22 (Dubiella et

al., 2013), we sought to investigate if IDA induces a release of $[Ca^{2+}]_{cyt}$. Here we report that IDA is able

to induce an intracellular release of Ca^{2+} in Arabidopsis root tips as well as in the AZ. Pursuing the tight

connection between ROS and Ca^{2+} in other defense signaling systems we explored if the the observed

production of ROS and Ca^{2+} in the IDA-HAE/HSL2 signaling pathway cold be linked to the involvement of

IDA in regulating plant defense (Patharkar et al., 2017). We show that a range of biotic and abiotic

signals can induce *IDA* expression and we demonstrate that the IDA signal modulates responses of plant

immune signaling. We propose that, in addition to regulating cell separation, IDA plays a role in

enhancing a defense response in tissues undergoing cell separation thus providing the essential need

for enhanced defense responses in tissue during the cell separation event.

Results:

IDA induces apoplastic production of ROS and elevation in cytosolic calcium-ion concentration

We have previously shown that a 12 amino acids functional IDA peptide, hereby referred to as mIDA

(Sup Table 1), elicits the production of a ROS burst in *N.benthamiana* leaves transiently expressing *HAE*

or *HSL2* (Butenko et al., 2014). To investigate whether mIDA could elicit a ROS burst in Arabidopsis, a

luminol-dependent ROS assay on *hae hsl2* mutant rosette leaves expressing the full length HAE receptor

under a constitutive promoter (*35S:HAE-YFP*) was performed. *hae hsl2 35S:HAE-YFP* leaf discs treated

with mIDA emitted extracellular ROS, whereas no response was observed in wild-type (WT) leaves (Sup

Fig.1a-c). We investigated the activity of the *HAE* and *HSL2* promoter in 22 days old Arabidopsis rosette

leaves by cloning the promoters fused to the nuclear localized YFP-derived fluorophore Venus protein

(*pHAE:Venus-H2B*, *pHSL2:Venus-H2B*) and could observe a decrease in promoter activity in the oldest

rosette leaves giving a possible explanation to the lack of mIDA induced ROS in true leaves (Sup Fig.

1d,e). Since rapid production of ROS is often linked to an elevation in $[Ca^{2+}]_{cyt}$ (Steinhorst & Kudla, 2013),

we aimed to investigate if mIDA could induce a $[Ca^{2+}]_{cyt}$ response. We imaged Ca^{2+} in 10 days-old roots

expressing the cytosolic localized fluorescent Ca^{2+} sensor R-GECO1 (Keinath et al., 2015). As a positive

control for $[Ca^{2+}]_{cyt}$ release we added 1 mM extracellular ATP (eATP), which leads to a $[Ca^{2+}]_{cyt}$ elevation

in the root tip within a minute after application (Fig. 1a) (Breiden et al., 2021). Following application of

1 µM mIDA, we detected an increase in $[Ca^{2+}]_{cyt}$ (Fig. 1a and Movie 1). R-GECO1 fluorescence intensities

normalized to background intensities (ΔF/F) were measured from a region of interest (ROI) covering the

meristematic and elongation zone of the root and revealed that the response starts 4-5 minutes after

application of the mIDA peptide and lasts for 7-8 minutes (Fig. 1a). The signal initiated in the root

meristematic zone from where it spread toward the elongation zone and root tip, a second wave was

observed in the meristematic zone continuing with Ca^{2+} spikes. The signal amplitude was at a maximum

within the elongation zone and decreased as the signal spread (Fig. 1a).

The $[Ca^{2+}]_{cyt}$ response in roots to the bacterial elicitor flg22 has previously been studied using the R-

GECO1 sensor (Keinath et al., 2015). To better understand the specificity of the mIDA induced Ca^{2+}

response, we aimed to compare the Ca^{2+} dynamics in mIDA treated roots to those treated with flg22.

We observed striking differences in the onset and distribution of the Ca^{2+} signals. Analysis of roots

treated with 1 µM flg22 showed that the Ca^{2+} signal initiated in the root elongation zone from where it

spread toward the meristematic zone as a single wave and that the signal amplitude was at a maximum

within the elongation zone and decreased as the signal spread (Fig. 1b, Movie 2). These observations

indicate differences in tissue specificity of Ca^{2+} responses between mIDA and flg22, which we

hypothesize depend on the cellular distribution of their cognate receptors. Indeed, when investigating

the promoter activity of the *HAE*, *HSL2* and *FLS2* receptors by the use of nuclear localized transcriptional

reporter lines we observed a different pattern of fluorescent nuclei in the roots (Fig. 1c,d,e).

pHAE:Venus-H2B lines had fluorescent expression in the epidermis and stele of the elongation zone (Fig.

1c) and *pHSL2:Venus-H2B* lines in the lateral root cap, root tip and root meristem (Fig. 1d); while

fluorescent nuclei were observed in the stele of the elongation zone in *pFLS2:Venus-H2B lines* (Fig. 1e).

The different patterns of fluorescent nuclei observed for the three different receptor constructs could

indeed explain the differences in the Ca^{2+} signatures triggered by mIDA and flg22. However, the

receptors promoter activity show a broader expression pattern compared to the respective ligand

induced $[Ca^{2+}]_{cyt}$ responses, indicating additional regulating factors for the observed $[Ca^{2+}]_{cyt}$ response.

To further investigate the mIDA induced Ca^{2+} response, we used a cytosolic localized Aequorin-based

luminescence Ca^{2+} sensor (Aeq) (Knight et al., 1991). The Aeq sensor was chosen due to the well-

established use of this sensor in studies investigating peptide induced Ca^{2+} responses (Ranf et al., 2011)

Removing the last Asparagine of the mIDA peptide (IDAΔN69; Sup Table 1) renders it inactive (Butenko

et al., 2014). 7 days old *Aeq*-Seedlings treated with 1 µM IDAΔN69 did not show any increase in $[Ca^{2+}]_{cyt}$

(Sup Fig. 4a). To investigate if the $[Ca^{2+}]_{cyt}$ increase triggered by mIDA was dependent on extracellular

Ca^{2+}, we pre-treated seedling with $LaCl_3$ and Ethylene glycol-bis (2-aminoethylether)- N, N, N',N'-

tetraacetic acid (EGTA). $LaCl_3$ and EGTA are inhibitors that block plasma membrane localized cation

channels and chelate Ca^{2+} in the extracellular space, respectively (Knight et al., 1997). The mIDA induced

response was abolished in *Aeq*-seedlings pre-incubated in 2 mM EGTA or 1 mM $LaCL_3$ (Sup Fig. 4b),

indicating that the mIDA dependent $[Ca^{2+}]_{cyt}$ response depends on Ca^{2+} from the extracellular space.

Next we set out to investigate whether mIDA would trigger an increase in $[Ca^{2+}]_{cyt}$ in floral AZ cells. We

utilized the Aeq expressing line and monitored $[Ca^{2+}]_{cyt}$ changes in flowers at different developmental

stages (see Sup Fig. 5a for developmental stages [floral positions]) to 1 µM mIDA. Interestingly, only

flowers at a the stage where there is an initial weakening of the AZ cell walls showed an increase in

$[Ca^{2+}]_{cyt}$ (Sup Fig. 5). Flowers treated with mIDA prior to cell wall loosening showed no increase in

luminescence (Sup Fig. 5), indicating that the mIDA triggered Ca^{2+} release in flowers correlates with the

onset of the abscission process and the increase in *HAE* and *HSL2* expression at the AZ (Cai & Lashbrook,

2008; Patharkar & Walker, 2015). Using plants expressing the R-GECO1 sensor we performed a detailed

investigation of the mIDA induced $[Ca^{2+}]_{cyt}$ response in flowers at position 6 which is the position where

initial cell wall loosening occurs. The AZ region was analyzed for signal intensity values and revealed that

the Ca^{2+} signal was composed of one wave (Fig. 2, Movie 3). The signal initiated close to the nectaries

and spread throughout the AZ and further into the floral receptacle. Flowers treated with 1 µM IDAΔN69

did not show any increase in $[Ca^{2+}]_{cyt}$ in the AZ or receptacle, while a clear Ca^{2+} wave could be detected

in the whole AZ, receptacle and proximal pedicel after treatment with eATP (Sup Fig.7, Sup Movie 2).

Detailed investigation of *HAE* and *HSL2* promoter activity in flowers at position 6 shows restricted

promoter activity to the AZ cells (Fig. 2b). The observed promoter activity correlates with the observed

mIDA induced $[Ca^{2+}]_{cyt}$ response

193 The ROS producing enzymes, RBOHD and RBOHF, are not involved in the developmental process of

194 abscission.

ROS and Ca^{2+} are secondary messengers that can play a role in developmental processes involving cell

wall remodeling, but are also important players in plant immunity (Kärkönen & Kuchitsu, 2015). We

therefore set out to investigate if the IDA induced production of ROS and increase in $[Ca^{2+}]_{cyt}$ function in

the developmental process of abscission or if these signaling molecules form part of IDA modulated

plant immunity.

The IDA induced Ca^{2+} response depends on Ca^{2+} from extracellular space (Sup Fig. 4b). Ca^{2+} transport

over the plasma membrane is enabled by a variety of Ca^{2+} permeable channels, including the CYCLIC

NUCLEOTIDE GATED CHANNELs (CNGCs). Various CNGCs are known to be involved in peptide ligand

signaling in plants, including the involvement of CNGC6 and CNGC9 in CLAVATA3/EMBRYO

SURROUNDING REGION40 signaling in roots (Breiden et al., 2021), and CNGC17 In phytosulfokine

signaling (Ladwig et al., 2015). Using publicly available expression data we identified multiple CNGCs

expressed in the AZ of Arabidopsis (Cai & Lashbrook, 2008) (Sup Fig. 8a) and investigated if plants

carrying mutations in the *CNGSs* genes expressed in AZs showed a defect in floral organ abscission. We

observed no deficiency in floral abscission in the *cngc* mutants (Sup Fig. 8b), in contrast to what is

observed in the *hae hsl2* mutant where all floral organs are retained throughout the inflorescence. Ca^{2+}

partially activates members of the NOX family of nicotinamide adenine dinucleotide phosphate (NADPH)

oxidases (RBOHs), key producers of ROS (Kadota et al., 2015). Two members of this family, *RBOHD* and

RBOHF have high transcriptional levels in the floral AZs (Cai & Lashbrook, 2008) and have been reported

to be important in the cell separation event during floral abscission in Arabidopsis (Lee et al., 2018). We

therefore set out to investigate if ROS production in AZ cells was dependent on RBOHD and RBOHF and

to quantify to which degree these NADPH oxidases were necessary for organ separation. We treated

AZs with the ROS indicator, H2DCF-DA, which upon contact with ROS has fluorescent properties, and

observed ROS in both WT and to a lower extent in *rbohd rbohdf* flowers. In addition, we observed normal

floral abscission in the *rbohd rbohf* double mutant flowers, indicating that the developmental

progression of cell separation is not dependent on RBOHD and RBOHF (Sup Fig. 8c,d). These results are

in stark contrast to what has previously been reported, where cell separation during floral abscission

was shown to be dependent on RBOHD and RBOHF (Lee et al., 2018). We used a stress transducer to

quantify the force needed to remove petals, the petal breakstrength (pBS) (Stenvik et al., 2008), from

the receptacle of gradually older flowers along the inflorescence of WT and *rbohd rbohf* flowers.

Measurements showed a significant lower pBS value for the *rbohd rbohf* mutant compared to WT at the

developmental stage where cell loosening normally occurs, indicating premature cell wall loosening

(Sup Fig. 8e). Furthermore, *rbohd rbohf* petals abscised one position earlier than WT (Sup Fig. 8e).

Similar to Crick et al 2022,, we found no RBOHD and RBOHF dependent delay or absence of cell

separation during floral abscission (Crick et al., 2022). Due to a known function of IDA in inducing ROS

production (Sup Fig. 1), these results points towards a role for IDA in modulating additional responses

to the process of cell separation.

IDA is upregulated by biotic and abiotic factors

In Arabidopsis, infection with the pathogen *P.syringae* induces cauline leaf abscission. Interestingly,

stress induced cauline leaf abscission was reduced in plants with mutations in components of the IDA-

HAE/HSL2 signaling pathway (Patharkar et al., 2017; Patharkar & Walker, 2016). In addition,

upregulation of defense genes in floral AZ cells during the abscission process is altered in the *hae hsl2*

mutant (Cai & Lashbrook, 2008; Niederhuth et al., 2013), indicating that the IDA-HAE/HSL2 signaling

system may be involved in regulating defense responses in cells undergoing cell separation. We further

investigated if *IDA* is upregulated by biotic and abiotic elicitors. Transgenic plants containing the *β-*

glucuronidase (GUS) gene under the control of the *IDA* promoter (*pIDA:GUS*) have previously been

reported to exhibit *GUS* expression in cells overlaying newly formed LR primordia (Butenko et al., 2003;

Kumpf et al., 2013). To investigate the expression of the *IDA* gene in response to biotic stress, *pIDA:GUS*

seedlings were exposed to the bacterial elicitor flg22 and fungal chitin. Compared to the control

seedlings, the seedlings exposed to flg22 and chitin showed a significant increase in *GUS* expression in

LR primordia (Fig. 3a). Also, when comparing untreated root tips to those exposed to flg22 and chitin

we observed a GUS signal in the primary root of flg22 and chitin treated roots (Fig. 3b). We used the

fluorogenic substrate 4-methylumbelliferyl β-D-glucuronide (4-MUG) which is hydrolysed to the

fluorochrome 4-methyl umbelliferone (4-MU) to quantify GUS-activity (Blazquez, 2007). A significant

increase in fluorescence was observed for flg22 and chitin treated samples compared to untreated

controls, (Fig. 3b), which was not observed when exposing seedlings to IDAΔN69 indicating that

increased *IDA* expression is not triggered simply by the application of a peptide (Fig. 3). Spatial *pIDA:GUS*

expression was also monitored upon abiotic stress. When *pIDA:GUS* seedling were treated with the

osmotic agent mannitol or exposed to salt stress by NaCl, both treatments simulating dry soil, a similar

increment in GUS signal was observed surrounding the LR primordia (Fig. 3). However, no *pIDA:GUS*

expression was detected in the primary root (Fig. 3b). Taken together these results indicate an

upregulation of *IDA* in tissue involved in cell separation processes such as the LR primordia and,

interestingly, the root cap upon biotic stress. The process of cell separation is a fast and time-restricted

response where cells previously protected by outer cell layers are rapidly exposed to the environment.

These cells need strong and rapid protection against the environment, and we therefore aimed to

investigate if IDA is involved in modulating immune responses in addition to inducing ROS and $[Ca^{2+}]_{cyt}$

responses.

mIDA can induce production of callose as a long term defense response

We aimed to investigate a possible role of IDA in regulating long-term defense responses in the plant, such as the production of callose. Callose deposition is a resulting hallmark of the late plant immunity responses that together with other events such as stomatal closure and production of ethylene leads to inhibition of pathogen multiplication and containment of disease (reviewed in (Wang et al., 2021)). Callose deposition is highly increased in response to flg22 (Gómez-Gómez et al., 1999). Promoter activity studies of *pHAE* and *pHSL2* indicates a weak activity of the promoters in the cotelydons, (Sup Fig. 1d) and we thus treated WT seedlings with 1 μM mIDA and measured the callose depositions per area in the cotyledons. We detected no difference to water treated controls indicating that either callose deposition is not a cellular outcome for mIDA treatment (Fig. 4a,b), or that receptor availability in the cotyledons is too low to observe a response We further tested if mIDA could induce callose depositions in cotyledons expressing the full length HAE receptor under a constitutive promoter (*35S:HAE-YFP*). An increase in the callose depositions per area upon mIDA treatment compared to water treated controls was observed in these lines (Fig. 4a,b). As a control, flg22 treated WT seedlings showed increased callose deposition, which was not observed in flg22 treated *fls2* seedlings (Fig. 4a,b). These results indicate that mIDA, is capable of promoting a HAE dependent deposition of callose as a long-term defense response.

mIDA triggers expression of defense-associated marker genes associated with innate immunity.

A -well-known part of a plants immune response is the enhanced transcription of genes involved in immunity. Transcriptional reprogramming mostly mediated by WRKY transcription factors takes place during microbe-associated molecular pattern (MAMP)- triggered immunity and is essential to mount an appropriate host defense response (Birkenbihl et al., 2017). We selected well-established defense-associated marker genes: *FLG22-INDUCED RECEPTOR-LIKE KINASE1* (*FRK1*), a specific and early immune-responsive gene activated by multiple MAMPs (Asai et al., 2002; He et al., 2006), *MYB51*, a WRKY target gene, previously shown to increase in response to the MAMP flg22 (Frerigmann et al., 2016) resulting in biosynthesis of the secondary metabolite indolic glucosinolates; and an endogenous danger peptide, ELICITOR PEPTIDE 3 (PEP3) (Huffaker et al., 2006). Seven days-old WT seedlings treated with 1 μM mIDA were monitored for changes in transcription of the aforementioned genes by RT-qPCR compared to untreated controls at two time points, 1 h and 12 h after treatment. All genes showed a significant transcriptional elevation 1 h after mIDA treatment (Fig. 5a). This in accordance with previous reports, where the expression of *FRK1* and *PEP3* was monitored in response to bacterial elicitors (He et al., 2006; Huffaker et al., 2006). After 12 h, *FRK1* and *PEP3* transcription in mIDA treated tissue was not

significantly different from untreated tissue; while transcription of *MYB51* in mIDA treated tissue
remained elevated after 12 h but was significantly lower compared to 1 h after mIDA treatment (Fig.
5a). Interestingly, the observed increase in transcription of the defense genes *MYB51* and *PEP3*, was
only partially reduced in the *hae hsl2* mutant (Sup Fig. 9). In contrast, *FRK1* showed higher
transcriptional levels in the *hae hsl2* mutant compared to WT (Sup Fig. 9). These results indicate that
also other receptors than HAE and HSL2 are involved in the mIDA-induced enhancement of the defense
genes *FRK1*, *MYB51* and *PEP3*.

We then analyzed a plant line expressing nuclear localized YFP from the MYB51 promoter
(*pMYB51:YFP^N*) (Poncini et al., 2017) treated with mIDA. Enhanced expression of p*MYB51:YFP* was
predominantly detected in the meristematic zone of the root after 7 h of mIDA treatment, compared to
a non-treated control (Fig. 5b). Moreover, comparison of mIDA induction with the elicitor-triggered
response of *pMYB51:YFP^N* to flg22 showed similar temporal expression in the root (Sup Fig. 10) .
Together, these data show that IDA can trigger a rapid increase in the expression of key genes involved
in immunity. We next investigated if the mIDA induced transcription of defense related genes was
similar to that induced by flg22. Seven days-old Col-0 WT seedlings were treated for 1h with mIDA, flg22
or a combination of mIDA and flg22 to explore potential additive effects. The relative increase in
transcription of the genes tested is similar when seedlings are treated with mIDA and flg22 (Fig. 6a).
However, when treating seedlings with both peptides the relative transcription of the genes of interest
increased substantially. To our surprise, the combined enhanced transcription when seedling where co-
treated with both flg22 and mIDA exceeds a simple additive effect of the two peptides. To investigate if
the observed increase is specific to a combination of mIDA and flg22, we investigated if co-treatment
with mIDA and PIP1, an endogenous DAMP peptide known to amplify the immune response triggered
by flg22, (Hou et al., 2014) gave a similar effect. Seedlings were treated for 1h with mIDA, PIP1 or a
combination of the peptides to explore additive effects. In contrast to what we observed with flg22, no
enhanced transcription of *FRK1*, *MYB51* and *PEP3* were observed after co-treatment with PIP1 and mIDA
(Sup Fig. 11). These results suggest a role of IDA in enhancing the defense response triggered by flg22.
To explore whether cells capable of undergoing cell separation in response to IDA and IDL peptides
could be expressing *HAE* or *HSL2* in combination with *FLS2* we made transcriptional reporter lines
expressing each of the receptors in fusion with either the nuclear localized YFP-derived fluorophore
Venus protein or the RFP-derived Tomato protein (*pRECEPTOR:Venus/Tomato-H2B*) and crossed the
lines to each other. Plants expressing both constructs were inspected for fluorescent nuclei in 7 day-old
roots as well as in floral organs of floral position 7. Fluorescent nuclei with overlapping Venus and
Tomato expression were observed both in the vascular tissue of the root, in cells surrounding LRs as
well as in cells of the AZs of plants expressing *pFLS2-Venus-H2B* and *pHAE-Tomat-H2B* (Fig. 6b). Similar

observations were done in the AZ of lines expressing *pFLS2-Venus-H2B* and *pHSL2-Tomat-H2B lines*,

however in these lines overlapping fluorescent nuclei in roots were mainly observed in the epidermal

cells surrounding the LR and in the root tip, as well as in the root cap (Fig. 6c). Based on these results

we propose a role for IDA in enhancing immune responses in tissue undergoing cell separation, possibly

by enhancing cellular responses activated by immune receptors, such as FLS2.

Discussion:

Several plant species can abscise infected organs to limit colonization of pathogenic microorganisms

thereby adding an additional layer of defense to the innate immune system (Kissoudis et al., 2016;

Patharkar et al., 2017). Also, in a developmental context, there is an induction of defense associated genes

in AZ cells during cell separation and prior to the formation of the suberized layer that functions as a barrier

rendering protection to pathogen attacks (Cai & Lashbrook, 2008; Niederhuth et al., 2013; Roberts et al.,

2000; I. Taylor & J. C. Walker, 2018). Interestingly, the induction of defense genes during floral organ

abscission is altered in *hae hsl2* plants (Niederhuth et al., 2013). It is largely unknown how molecular

components regulating cell separation and how the IDA-HAE/HSL2 signaling pathway, contributes to

modulation of plant immune responses. Here we show that the mature IDA peptide, mIDA, is involved in

activating known defense responses in Arabidopsis, including the production of ROS, the intracellular

release of Ca^{2+}, the production of callose and the transcription of genes known to be involved in defense

(Sup Fig. 1, Fig. 1, Fig. 2, Fig. 5). We suggest a role for IDA in ensuring optimal cellular responses in tissue

undergoing cell separation, regulating both development and defense.

The intracellular event occurring downstream of IDA is still partially unknown. Here we report the

release of cytosolic Ca^{2+} as a new signaling component. Roots and AZs expressing the Ca^{2+} sensor R-

GECO1, showed a $[Ca^{2+}]_{cyt}$ in response to mIDA. Moreover, the increased $[Ca^{2+}]_{cyt}$ in response to mIDA in

flowers occurred only at stages where cell separation was taking place, linking the observed $[Ca^{2+}]_{cyt}$

response to the temporal point of abscission and the need for an induced local defense response. During

floral development the expression of the *HAE* and *HSL2* receptors increase prior to the onset of

abscission (Cai & Lashbrook, 2008) and may account for the temporal mIDA induced $[Ca^{2+}]_{cyt}$ response

observed.

Based on these results, it is likely that the expression of the receptors is the limiting factor for a cells ability

to respond to the mIDA peptide with a $[Ca^{2+}]_{cyt}$ response. However, promoter activity of the receptors are

observed in a broader area of the root than the observed $[Ca^{2+}]_{cyt}$ response. As an example, *pHSL2* is found

to have a localized activation in the root tip, and when HSL2 is activated by IDL1 be involved in the

regulation of cell separation during root cap sloughing (Shi et al., 2018). IDL1 and IDA have a high degree

of similarity in the protein sequence, and *IDL1* expressed under the *IDA* promoter can fully rescue the

abscission phenotype observed in the *ida* mutant (Stenvik et al., 2008). We can assume that IDL1 activates

similar signaling components in the root cap as those used by IDA. Despite the expression of HSL2 and the

involvement of the receptor in root cap sloughing, no $[Ca^{2+}]_{cyt}$ response is observed in this region when

plants are treated with the IDA peptide. The absence of an observed $[Ca^{2+}]_{cyt}$ response in the root tip

indicates that $[Ca^{2+}]_{cyt}$ is most likely not involved in the cell separation process during root cap sloughing, or

that cells involved in this process only are able to induce a $[Ca^{2+}]_{cyt}$ response at a specific developmental

time point. Thus, in addition to receptor availability, other factors must be involved in regulating the $[Ca^{2+}]_{cyt}$

response. It is intriguing that the mIDA cellular output is similar to that of flg22 (Kadota et al., 2015; Kadota

et al., 2014) yet shows some distinct differences. The spatial distribution of the Ca^{2+} wave originated by

mIDA differs from that of flg22. The *rbohd rbohf* mutant shows ROS production in the AZ, indicating that

other NADPH oxidases or cell wall peroxidases expressed in the AZ may be involved in ROS production, in

contrast to the importance of RBOHD RBOHF in flg22 signaling (Kadota et al., 2015). This suggests that while

IDA and flg22 share many components for their signal transduction, such as the BAK1 co-receptor

(Chinchilla et al., 2007; X. Meng et al., 2016) and MAPK cascade (Asai et al., 2002; Cho et al., 2008), there

are other specific components that are likely important to provide differences in signaling. Despite the

observation of the numerous defense related responses to mIDA, we cannot exclude that the mIDA induced

$[Ca^{2+}]_{cyt}$ and rises in ROS levels may also have an intrinsic developmental role such as modulating cell wall

properties and cell expansion similar to what is observed during FERONIA (FER) signaling (Dünser et al.,

2019; Wei Feng et al., 2018). To further decipher the importance of the mIDA induced Ca^{2+} and ROS in

defense it will be important to identify the specific signaling components responsible for this cellular output

in addition to investigating the susceptibility of mutants to pathogen exposure.

Interestingly, FLS2 arrangement into nanoscale domains at the PM is dependent on FER. FLS2 becomes

more disperse and mobile in *fer* mutants, implying a role for FER in regulating FLS2 activity (Gronnier et

al., 2022). Also altering of the cell wall affects FLS2 nanoscale organization (McKenna et al., 2019).

Formation of nanoscale domains containing the HAE and HSL2 receptors may be an important signaling

step in the IDA-HAE/HSL2 signaling pathway. From our promoter analysis we clearly see overlapping

activity of the *HAE* and *HSL2* promoters with the *FLS2* promoter *in planta*, providing the possibility that

the receptors are localized at the PM of the same cells and can have coordinated signaling events. An

intriguing thought is a possible co-localization of HAE and HSL2 with FLS2 in nanoscale domains, making

a signaling unit possible to involve multiple different receptors responsible for the signaling outcome

observed. In this paper, we show that co-treatment with mIDA and flg22 induces an enhanced

transcription of defense related genes. How FLS2, HAE and HSL2 communicate at the plasma membrane

of the responsive cells may give the answers to the molecular mechanisms behind the augmented

transcriptional regulation observed when co-treatment of mIDA and flg22 is performed (Fig. 6). We can

investigate possible interaction partners of HAE and HSL2 looking at available data from a sensitized

high-throughput interaction assay between extracellular domains (ECDs) of 200 Leucine Rich Repeat

(LRR) (Smakowska-Luzan et al., 2018). Interestingly, LRR-RLKs known to play a function in biotic or

abiotic stress responses, such as RECEPTOR-LIKE KINASE7 (RLK7), STRUBBELIG-RECEPTOR FAMILY3 ,

RECEPTOR-LIKE PROTEIN KINASE 1 and FRK1, are found to interact mainly with HSL2, whereas HAE

shows interaction with LRRs mainly involved in development (Sup Fig. 12). RLK7 has recently been

identified as an additional receptor for CEP4, involved in regulating cell surface immunity and response

to nitrogen starvation, indication a role of RLK7 as an additional receptor regulating defense responses

in known peptide-receptor complexes (Rzemieniewski et al., 2022). It would be interesting to investigate

a possible role of RLK7 in IDA-HAE/HSL2 signaling as a signaling partner in the observed IDA-induced

defense responses.

In this paper, we propose a model where the IDA-HAE/HSL2 signaling pathway modulates defense

responses in tissues undergoing cell separation. It is beneficial for the plant to ensure an upregulation

of defense responses in tissues undergoing cell separation as a precaution against pathogen attack. This

tissue is a major entry route for pathogens, and by combining the need of both the IDA peptide and the

pathogen molecular pattern, such as flg22, for the induction of a strong defense response, the plant

ensures maximum immunity in the most prone cells without spending energy inducing this in all cells

(Fig. 7). Such a molecular mechanism, where the presence of mIDA in tissue exposed to stress leads to

maximum activation of the immune responses, will ensure maximal protection of infected cells during

cell separation, cells which are major potential entry routes for invading pathogens.

Movies

Movie 1: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 μM mIDA. 10 days-old roots were treated with 1 μM mIDA and changes in $[Ca^{2+}]_{cyt}$ was recorded over time. Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds to measurements shown in Fig. 1a. Representative response from 10 roots. Time shown as, hh:mm:ss.

Movie 2: $[Ca^{2+}]_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 μM flg22 10 days-old roots were treated with 1 μM flg22 and changes in $[Ca^{2+}]_{cyt}$ was recorded over time. Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds to measurements shown in Fig. 1b. Representative response from 7 roots. Time shown as, hh:mm:ss.

Movie 3: [Ca^{2+}]$_{cyt}$ dynamics in R-GECO1 expressing AZ in response to 1 µM mIDA. Flowers at position 6 were treated with 1 µM mIDA and changes in [Ca^{2+}]$_{cyt}$ was recorded over time. Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds to measurements shown in Fig. 2a. Representative response from 8 flowers. Time shown as, hh:mm:ss.

Sup Movie 1: [Ca^{2+}]$_{cyt}$ dynamics in R-GECO1 expressing root tip in response to 1 mM eATP. 1 mM eATP will induce a strong increase in [Ca^{2+}]$_{cyt}$ and was added as a last treatment to all roots acting as a positive control. Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds to measurements shown in Fig. 1a. Representative response from 10 roots. Time shown as, hh:mm:ss.

Sup Movie 2: [Ca^{2+}]$_{cyt}$ dynamics in R-GECO1 expressing AZ in response to to 1 µM of the inactive IDA peptide, IDA$^{\Delta N69}$. Flowers at position 6 were treated with 1 µM IDA$^{\Delta N69}$ and changes in [Ca^{2+}]$_{cyt}$ was recorded over time. Response shown as normalized fluorescence intensities (ΔF/F). Images were recorded with a frame rate of 5 seconds. Movie corresponds to measurements shown in Sup Fig. 6. Representative response from 7 flowers. Time shown as, hh:mm:ss.

Methods

Accession numbers of genes studied in this work:

HSL2 At5g65710, IDA At1g68765, FLS2 At5g46330, HAE At4g28490, MYB51 At1g18570, RBOHD At5g47910, RBOHF At1g64060, PEP3 At5g64905, WRKY33 At2g38470, FRK1 At2g19190. Plant lines used in this work: Ecotype Colombia-0 (Col-0) was used as wild type (WT). Mutant line: *hae* (SALK_021905), *hsl2*, (SALK_030520), *fls2* (SALK_062054), *rbohd* (SALK_070610), *rbohf* (SALK_059888), *cngc1* (SAIL_443), *cngc2* (SALK_066908), *cngc4* (SALK_081369), *cngc5* (SALK_149893), *cngc6* (SALK_064702), *cngc9* (SAIL_736), *cngc12* (SALK_093622). SALK lines were provided from Nottingham Arabidopsis Stock Centre (NASC).

Plant lines

The pIDA:GUS, pHSL2:Venus-H2B and pMYB51:YFPN lines have been described previously (Kumpf et al., 2013; Poncini et al., 2017; Shi et al., 2018).The promoters of HAE (1601 bp (Kumpf et al., 2013) and HSL2

(2300 bp (Sto et al., 2015))were available in the pDONRZeo vector (Thermo Fischer Scientific).

Sequences corresponding to the FLS2 promoter (988 bp (Robatzek et al., 2006)) were amplified from

WT DNA (primers are listed in table 2) and cloned into the pDONRZeo vector (Thermo Fischer Scientific).

All promoter constructs were further recombined into the promotor:Venus (YFP)-H2B and Tomato-H2B

destination vectors (Somssich et al., 2016) using the Invitrogen Gateway cloning system (Thermo Fischer

Scientific). Constructs were transformed into *Agrobacterium tumefaciens* (A.tumefaciens) C58 and the

floral dip method (Clough & Bent, 1998) was used to generate transgenic lines. Single-copy homozygous

plant lines were selected and used in this study. The CDS of HAE was cloned into the pEarleyGate101

destination vector (Earley et al., 2006) using the Invitrogen Gateway cloning system and transformed

into A.tumefaciens C58 and further used to generate the *35S:HAE:YFP* lines.

<u>Growth conditions</u>

Plants were grown in long day conditions (8 h dark and 16 h light) at 22 °C. Seeds were surface sterilized

and plated out on MS-2 plates, stratified for 24 h at 4 °C and grown on plates for 7 days before

transferred to soil.

<u>Peptide sequences</u>

Peptides used in this study were ordered from BIOMATIK. Peptide sequences are listed in

Supplementary Table 1.

<u>Primers</u>

Primers for genotyping and generation of constructs were generated using VectorNTI. Gene specific

primers for RT-qPCR were generated using Roche Probe Library Primer Design. All primers are listed in

Supplementary Table 2.

<u>Histochemical GUS assay</u>

Seven days-old seedlings were pre-incubated for 12 h in liquid MS-2 medium containing stimuli of

interest; 1 µM peptide (table 1), 60 mM Mannitol (M4125 – Sigma), 20 µg/mL Chitin (C9752 – Sigma),

50 mM NaCl and then stained for GUS activity following the protocol previously described (Stenvik et

al., 2008). Roots were pictured using a Zeiss Axioplan2 microscope with an AxioCam HRc, 20x air

objective. The assay was performed on 10 individual roots and the experiment was repeated 3 times.

<u>Fluorescent GUS assay</u>

Seven days-old seedlings were pre-incubated for 12 h in liquid MS-2 medium with or without stimuli of

interest; 1 µM peptide (table 1), 60 mM Mannitol (M4125 – Sigma), 20 µg/mL Chitin (C9752 – Sigma),

50 mM NaCl. After treatment, 10 seedlings were incubated in wells containing 1 mL reaction mix
described in (Blazquez, 2007) containing: 10 mM EDTA (pH 8.0), 0,1 % SDS, 50 mM Sodium Phosphate
(pH 7.0), 0,1 % Triton X-100, 1 mM 4-MUG (, M9130-Sigma) and incubated at 37 °C for 6 h. Six 100 µl
aliquots from each well were transferred to individual wells in a microtiter plate and the reaction was
stopped by adding 50 µl of stop reagent (1 M Sodium Carbonate) to each well. Fluorescence was
detected by the use of a Wallac 1420 VICTOR2 microplate luminometer (PerkinElmer) using an
excitation wavelength of 365 nm and a filter wavelength of 430 nm. Each experiment was repeated 3
times.

<u>Confocal laser microscopy of roots and flowers expressing promoter:Venus/YFP-H2B and promoter:</u>
<u>Tomato-H2B construct</u>

Imaging of 7 days-old roots was performed on a LSM 880 Airyscan confocal microscope equipped with
two flanking PMTs and a central 32 array GaAsP detector. Images were acquired with Zeiss Plan-
Apochromat 20x/0.8 WD=0.55 M27 objective and excited with laser light of 405 nm, 488 nm and 561
nm. Roots were stained by 1 µM propidium iodide for 10 min and washed in dH_2O before imaging.
Imaging of flowers was performed on an Andor Dragonfly spinning disk confocal using an EMCCD iXon
Ultra detector. Images were acquired with a 10x plan Apo NA 0,45 dry objective and excited with laser
light of 405 nm, 488 nm and 561 nm. Maximum intensity projections of z-stacks were acquired with
step size of 1,47 µm. Image processing was performed in FIJI 51. These steps are: background
subtraction, gaussian blur/smooth, brightness/contrast. Imaging was performed at the NorMIC Imaging
platform.

<u>Calcium imaging using the R-GECO1 sensor</u>

$[Ca^{2+}]_{cyt}$ in roots were detected using WT plants expressing the cytosolic localized single-fluorophore
based Ca^{2+} sensor, R-GECO1 in Col-0 background (Keinath et al., 2015). Measurements were performed
using a confocal laser scanning microscopy Leica TCS SP8 STED 3X using a 20x multi NA 0.75 objective.
Images were recorded with a frame rate of 5 seconds at 400 Hz. Seedling mounting was performed as
described in (Krebs & Schumacher, 2013). The plant tissues were incubated overnight in half strength
MS, 21°C and continuous light conditions before the day of imaging. $[Ca^{2+}]_{cyt}$ in the abscission zone were
detected using WT plants expressing R-GECO1 (Keinath et al., 2015). Abscission zones of position 7 were
used. Mounting of the abscission zones were performed using the same device as for seedlings (Krebs
& Schumacher, 2013). The abscission zones were incubated in half strength MS medium for 1 h before
imaging. Measurements were performed with a Zeiss LSM880 Airyscan using a Plan-Apochromat 10×

air NA 0.30 objective. Images were recorded with a frame rate of 10 seconds. R-GECO1 was excited

with a white light laser at 561 nm and its emission was detected at 590 nm to 670 nm using a HyD

detector. Laser power and gain settings were chosen for each experiment to maintain comparable

intensity values. For mIDA and flg22 two-fold concentrations were prepared in half strength MS. mIDA

or flg22 were added in a 1:1 volume ratio to the imaging chamber (final concentration 1 µM). ATP was

prepared in a 100-fold concentration in half strength MS and added as a last treatment in a 1:100

volume ratio (final concentration 1 mM) to the imaging chamber as a positive control for activity of the

R-GECO1 sensor (Movie 3). Image processing was performed in FIJI. These steps are: background

subtraction, gaussian blur, MultiStackReg v1.45 (http://bradbusse.net/sciencedownloads.html), 32-bit

conversion, threshold. Royal was used as a look up table. Fluorescence intensities of indicated ROIs were

obtained from the 32-bit images (Krebs & Schumacher, 2013). Normalization was done using the

following formula $\Delta F/F = (F-F0)/F0$ where F0 represents the mean of at least 1 min measurement

without any treatment. R-GECO1 measurements were performed at the Center for Advanced imaging

(CAi) at HHU and at NorMIC Imaging platform at the University of Oslo.

Calcium measurements using the Aequorin (pMAQ2) sensor

$[Ca^{2+}]_{cyt}$ in seedlings and flowers were detected using WT plants expressing p35S-apoaequorin (pMAQ2)

located to cytosol (Aeq) (Knight et al., 1991; Ranf et al., 2011). Aequorin luminescence was measured

as previously described (Ranf et al., 2012). Emitted light was detected by the use of a Wallac 1420

VICTOR2 microplate luminometer (PerkinElmer). Differences in Aeq expression levels due to seedling

size and expression of sensor were corrected by using luminescence at specific time point (L)/Max

Luminescence (Lmax). Lmax was measured after peptide treatment by individually adding 100 µL 2 M

$CaCl_2$ to each well and measuring luminescence constantly for 180 seconds (Ranf et al., 2012). 2 M $CaCl_2$

disrupts the cells and releases the Aeq sensor into the solution where it will react with Ca^{2+} and release

the total possible response in the sample (Lmax) in form of a luminescent peak. A final concentration of

1 µM mIDA was added to each wells at the start of measurements. For inhibitor treatments, Aeq-

seedlings were incubated in in 2 mM EGTA (Sigma-Aldrich) or 1 mM $LaCL_3$ (Sigma-Aldrich) O/N before

measurements. For seedlings 3 independent experiments were performed with 12 replications in each

experiment. For flowers 3 independent experiments were performed with 4-6 replications in each

experiment.

Measurements of reactive oxygen species (ROS)

ROS production was monitored by the use of a luminol-dependent assay as previously described

(Butenko et al., 2014) using a Wallac 1420 VICTOR2 microplate luminometer (PerkinElmer). Arabidopsis

leaves expressing *35S:HAE:YFP* were cut into leaf discs and incubated in water overnight before

measurements.   A final concentration of 1 μM mIDA was added to each well at the start of

measurements. All measurements were performed on 6 leaf discs and each experiment was repeated

3 times.

<u>ROS stain (H2DCF-DA)</u>

Flowers at position 6 were gently incubated in staining solution (25 μM (2',7'-dichlorodihydrofluorescein

diacetate) (H2DCF-DA) (Sigma-Aldrich, D6883), 50 mM KCL, 10 mM MES) for 10 min and further washed

3 times in wash solution (50 mM KCL, 10 mM MES). For the *hae hsl2* mutant the floral organs were

forcibly removed immediately before imaging. Imaging was done using a Dragonfly Airy scan spinning

disk confocal microscope, excited by a 488 nm laser. A total of 9 flowers per genotype were imaged.

The experiment was repeated 2 independent times.

<u>Callose deposition staining and quantification</u>

Callose deposition experiments were conducted as per (Luna et al., 2011) with modifications. Seeds

from WT and *fls2* were vapor-phase sterilized for 4 h. Seeds were transferred to 12-well plates

containing 1 mL of filter-sterilized MS medium and 0. 5 % MES with a final pH of 5.7, and stratified in

the dark at 4 °C for 2 days. Plates were consecutively transferred to a growth chamber with 16 h of light

(150μE m-2 s-1)/8h darkness cycle at 22°C. On the seventh day, growth medium was refreshed under

sterile conditions. On the eight day the treatments were applied: addition of 10 μL of water (mock); 10

μL of 100 μM flg22; or 10 μL of 100 μM mIDA (final peptide concentrations of 1 μM). After 24 h the

growth medium was replaced with 9 5% ethanol and incubated overnight. Ethanol was replaced with

an 8 M NaOH softening solution and incubated for 2 h. Seedlings were washed 3 times in water and

immediately transferred to the aniline blue staining solution (100 mM KPO4-KOH, pH 11; 0.01 % aniline

blue). After 2 h of staining, the seedlings were mounted on 50 % glycerol and imaged under a ZEISS

epifluorescence microscope with UV filter (BP 365/12 nm; FT 395 nm; LP 397 nm). Image analysis was

performed on FIJI (Schindelin et al., 2012). In brief, cotyledons were cropped and measured the leaf

area. Images were thresholded to remove autofluorescence and area of depositions measured to

calculate the ratio of callose deposition area per cotyledon area unit. Between 9-12 seedlings were

analyzed per genotype x treatment combination.

<u>Real time quantitative PCR (RT-qPCR)</u>

Seven days-old Arabidopsis seedlings (WT, *hae hsl2*) grown vertically on ½ sucrose MS-2 plates were

transferred to liquid ½ MS-2 medium (non-treated) and liquid ½ MS-2 medium containing 1 μM peptide

(Supplementary Table 1) and incubated in growth chambers for 1 h or 12 h. Seedlings were flash-frozen

in liquid nitrogen before total RNA was extracted using SpectrumTM Plant Total RNA Kit (SIGMA Aldrich).

cDNA synthesis was performed as previously described (Grini et al., 2009). RT-qPCR was performed

according to protocols provided by the manufacturer using FastStart Essential DNA Green Master

(Roche) and LightCycler96 (Roche) instrument. ACTIN2 was used to normalize mRNA levels as described

in (Grini et al., 2009). Two-three biological replicates and 4 technical replicates including standard curves

were performed for each sample.

<u>Petal Break strength (pBS)</u>

The force required to remove a petal at a given position on the inflorescence was measured in gram

equivalents using a load transducer as previously described (Stenvik et al., 2008). Plants were grown

until they had at least 15 positions on the inflorescence. A minimum of 15 petals per position were

measured. pBS measurements were performed on WT, *rbohd rbohf* (SALK_070610 SALK_059888) and

hae hsl2 (SALK_021905 SALK_030520) plants.

<u>Statistical methods</u>

Two tailed students t-test ($p < 0.05$) was used to identify significant differences in the fluorescent GUS

assay by comparing treated samples to untreated samples of the same plant line. Statistical analysis of

the RT-qPCR results was performed on all replicas using ONE-WAY or TWO-WAY ANOVA (as stated in

the figure text) and post-hoc Tukey's test ($p < 0.05$). Two tailed students t-test with ($p < 0.05$) was used

in the petal break strength (pBS) measurements to identify significant differences from WT at a given

position on the inflorescence.

Acknowledgements

We thank M.K. Anker, I.M. Stø, V. Iversen and R. Falleth for technical assistance in the laboratory and

phytotrone. We thank the NorMic Imaging platform for the use and technical support. This work was

supported by the Research Council of Norway (grant 230849) to V.O.Lalun and M.A.Butenko Work by R.

Simon and and M.Breiden was supported through CEPLAS.

Author contributions

V.O.L generated Arabidopsis lines and constructs, tested IDA expression to biotic and abiotic stress,

performed gene expression studies, phenotypic analysis of mutants and ROS measurements. V.O.L and

M.B. performed Ca^{2+} measurements. V.O.L, S. G-T., M.B., R.S and M.A.B designed experiments, analyzed

data, and drafted the manuscript. V.O.L and M.A.B wrote the paper with input from all authors.

Competing interests

The authors have no competing interests.

Materials and correspondence

Correspondence to Melinka A. Butenko, m.a.butenko@ibv.uio.no

Supplementary Tables

Supplementary Table 1: Peptide sequences

Peptide	Amino acid sequence
mIDA	PIPPSA**o**SKRHN
flg22	QRLSTGSRINSAKDDAAGLQIA
IDA$^{\Delta N69}$	PIPPSA**o**SKRH
PIP1	RFVKHSG**o**SPSGPGH

632 (**o** = hydroxyproline)

634 ## Supplementary Table 2: Primers sequences and function

Function	Primer name	Primer sequence 5'-3'
Cloning promoter FLS2	PromFLS2 -988bp Attb1	5'GGGGACAAGTTTGTACAAAAAAGCAGGCTTA GAAGTTGTGAATTGTGAT'3
Cloning promoter FLS2	PromFLS2 -988bp Attb2	5'GGGGACCACTTTGTACAAGAAAGCTGGGTA GGTTTAGACTTTAGAAGA'3
Genotype hsl2	Hsl2 LP	5'CGTCTTGAGCTAGCCAACAAC'3
Genotype hsl2	Hsl2 RP	5'GTCCAATCAAGTGGAGAAACG'3

Genotype hae	Haesa LP	5'CACCTTCCTTCTCTCCATTCC'3
Genotype hae	Haesa RP	5'GTTCGAGAAGTGACAAGCGAG'3
Genotype SALK-lines	Lbb1-	5'GCGTGGACCGCTTGCTGCAACT'3
Genotype rbohd SALK_070610C	LP_SALK_070610C	5'TTTCAACGCCTTTTGGTACAC'3
Genotype rbohd SALK_070610C	RP_SALK_070610C	5'GTTACCTATTCTTTTGCCGGG'3
Genotype rbohf SALK_059888	LP_SALK_059888	5'CAAAGAGCTCTTCGTGGTTTG'3
Genotype rbohf SALK_059888	RP_SALK_059888	5'TCTCTATTGTATCTTGTGTCACCG'3
Genotype fls2 SALK_062054	FLS2_SALK_062054 Fw	5'GGTTCGATTCCTTCTGGAATC'3
Genotype fls2 SALK_062054	FLS2_SALK_062054 Rv	5'CCTGAGTTTTTGAAGCTTCCC'3
qPCR	33.FRK1 Fw	5'AACTTAGGAGACTATTTGGCAGGTAA'3
qPCR	33.FRK1 Rv	5'TGCATCTAATGATATCTTCAACCTCT'3
qPCR	63.PEP3 Fw	5'GCGAGGAAGATGAGAGTATCG'3
qPCR	63.PEP3 Rv	5'TCAATGGTCATGCCATCTTCT'3
qPCR	91.MYB51 Fw	5'GGCCAATTATCTTAGACCTGACA'3
qPCR	91.MYB51 Rv	5'CCACGAGCTATAGCAGACCATT'3
qPCR reference gene	act2int2 sense	5'CCCTGAGGAGCACCCAGTTCTACTC'3
qPCR reference gene	act2int2 antisense	5'CCGCAAGATCAAGACGAAGGATAGC'3

References:

Agustí, J., Merelo, P., Cercós, M., Tadeo, F. R., & Talón, M. (2009). Comparative transcriptional survey
between laser-microdissected cells from laminar abscission zone and petiolar cortical tissue
during ethylene-promoted abscission in citrus leaves. *BMC Plant Biology*, *9*(1), 127.
https://doi.org/10.1186/1471-2229-9-127
Asai, T., Tena, G., Plotnikova, J., Willmann, M. R., Chiu, W. L., Gomez-Gomez, L., Boller, T., Ausubel, F.
M., & Sheen, J. (2002). MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature*,
415(6875), 977-983. https://doi.org/10.1038/415977a
Birkenbihl, R. P., Kracher, B., & Somssich, I. E. (2017). Induced Genome-Wide Binding of Three
Arabidopsis WRKY Transcription Factors during Early MAMP-Triggered Immunity. *Plant Cell*,
29(1), 20-38. https://doi.org/10.1105/tpc.16.00681
Blazquez, M. (2007). Quantitative GUS Activity Assay in Intact Plant Tissue. *CSH Protoc*, *2007*,
pdb.prot4688. https://doi.org/10.1101/pdb.prot4688
Bleecker, A. B., & Patterson, S. E. (1997). Last exit: senescence, abscission, and meristem arrest in
Arabidopsis. *Plant Cell*, *9*(7), 1169-1179. https://doi.org/10.1105/tpc.9.7.1169
Breiden, M., Olsson, V., Blümke, P., Schlegel, J., Gustavo-Pinto, K., Dietrich, P., Butenko, M. A., & Simon,
R. (2021). The Cell Fate Controlling CLE40 Peptide Requires CNGCs to Trigger Highly Localized
Ca2+ Transients in Arabidopsis thaliana Root Meristems. *Plant Cell Physiol*, *62*(8), 1290-1301.
https://doi.org/10.1093/pcp/pcab079
Butenko, M. A., Patterson, S. E., Grini, P. E., Stenvik, G.-E., Amundsen, S. S., Mandal, A., & Aalen, R. B.
(2003). *INFLORESCENCE DEFICIENT IN ABSCISSION* Controls Floral Organ Abscission in
Arabidopsis and Identifies a Novel Family of Putative Ligands in Plants. *Plant Cell*, *15*, 2296-
2307.
Butenko, M. A., Wildhagen, M., Albert, M., Jehle, A., Kalbacher, H., Aalen, R. B., & Felix, G. (2014). Tools
and Strategies to Match Peptide-Ligand Receptor Pairs. *Plant Cell*, *26*(5), 1838-1847.
https://doi.org/10.1105/tpc.113.120071
Cai, S., & Lashbrook, C. C. (2008). Stamen abscission zone transcriptome profiling reveals new
candidates for abscission control: enhanced retention of floral organs in transgenic plants
overexpressing Arabidopsis ZINC FINGER PROTEIN2. *Plant Physiol*, *146*(3), 1305-1321.
https://doi.org/10.1104/pp.107.110908
Castro, B., Citterico, M., Kimura, S., Stevens, D. M., Wrzaczek, M., & Coaker, G. (2021). Stress-induced
reactive oxygen species compartmentalization, perception and signalling. *Nat Plants*, *7*(4),
403-412. https://doi.org/10.1038/s41477-021-00887-0
Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nurnberger, T., Jones, J. D., Felix, G., & Boller, T.
(2007). A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence.
Nature, *448*(7152), 497-500. https://doi.org/10.1038/nature05999
Cho, S. K., Larue, C. T., Chevalier, D., Wang, H., Jinn, T. L., & Zhang, S. (2008). Regulation of floral organ
abscission in Arabidopsis thaliana. *Proc Natl Acad Sci USA.*, *105*.
https://doi.org/10.1073/pnas.0805539105
Clough, S. J., & Bent, A. F. (1998). Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *The Plant Journal*, *16*(6), 735-743.
Crick, J., Corrigan, L., Belcram, K., Khan, M., Dawson, J. W., Adroher, B., Li, S., Hepworth, S. R., & Pautot,
V. (2022). Floral organ abscission in Arabidopsis requires the combined activities of three TALE
homeodomain transcription factors. *J Exp Bot*, *73*(18), 6150-6169.
https://doi.org/10.1093/jxb/erac255
Dubiella, U., Seybold, H., Durian, G., Komander, E., Lassig, R., Witte, C. P., Schulze, W. X., & Romeis, T.
(2013). Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for
rapid defense signal propagation. *Proc Natl Acad Sci U S A*, *110*(21), 8744-8749.
https://doi.org/10.1073/pnas.1221294110

Dünser, K., Gupta, S., Herger, A., Feraru, M. I., Ringli, C., & Kleine-Vehn, J. (2019). Extracellular matrix
sensing by FERONIA and Leucine-Rich Repeat Extensins controls vacuolar expansion during
cellular elongation in Arabidopsis thaliana. *The EMBO journal*, *38*(7), e100353.
https://doi.org/10.15252/embj.2018100353
Earley, K. W., Haag, J. R., Pontes, O., Opper, K., Juehne, T., Song, K., & Pikaard, C. S. (2006). Gateway-
compatible vectors for plant functional genomics and proteomics. *Plant J*, *45*(4), 616-629.
https://doi.org/10.1111/j.1365-313X.2005.02617.x
Felix, G., Duran, J. D., Volko, S., & Boller, T. (1999). Plants have a sensitive perception system for the
most conserved domain of bacterial flagellin. *The Plant Journal*, *18*(3), 265-276.
https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-
313X.1999.00265.x?sid=nlm%3Apubmed
Feng, W., Kita, D., Peaucelle, A., Cartwright, H. N., Doan, V., Duan, Q., Liu, M. C., Maman, J., Steinhorst,
L., Schmitz-Thom, I., Yvon, R., Kudla, J., Wu, H. M., Cheung, A. Y., & Dinneny, J. R. (2018). The
FERONIA Receptor Kinase Maintains Cell-Wall Integrity during Salt Stress through Ca(2+)
Signaling. *Curr Biol*, *28*(5), 666-675.e665. https://doi.org/10.1016/j.cub.2018.01.023

Frerigmann, H., Pislewska-Bednarek, M., Sanchez-Vallet, A., Molina, A., Glawischnig, E., Gigolashvili, T.,
& Bednarek, P. (2016). Regulation of Pathogen-Triggered Tryptophan Metabolism in
Arabidopsis thaliana by MYB Transcription Factors and Indole Glucosinolate Conversion
Products. *Mol Plant*, *9*(5), 682-695. https://doi.org/10.1016/j.molp.2016.01.006
Geiger, D., Scherzer, S., Mumm, P., Marten, I., Ache, P., Matschi, S., Liese, A., Wellmann, C., Al-Rasheid,
K., & Grill, E. (2010). Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with
distinct Ca2+ affinities. *Proceedings of the National Academy of Sciences*, *107*(17), 8023-8028.
Gómez-Gómez, L., Felix, G., & Boller, T. (1999). A single locus determines sensitivity to bacterial flagellin
in Arabidopsis thaliana. *Plant J*, *18*(3), 277-284. https://doi.org/10.1046/j.1365-
313x.1999.00451.x
Grini, P. E., Thorstensen, T., Alm, V., Vizcay-Barrena, G., Windju, S. S., Jorstad, T. S., Wilson, Z. A., &
Aalen, R. B. (2009). The ASH1 HOMOLOG 2 (ASHH2) histone H3 methyltransferase is required
for ovule and anther development in Arabidopsis. *PLoS One*, *4*(11), e7817.
https://doi.org/10.1371/journal.pone.0007817
Gronnier, J., Franck, C. M., Stegmann, M., DeFalco, T. A., Abarca, A., von Arx, M., Dünser, K., Lin, W.,
Yang, Z., Kleine-Vehn, J., Ringli, C., & Zipfel, C. (2022). Regulation of immune receptor kinase
plasma membrane nanoscale organization by a plant peptide hormone and its receptors. *Elife*,
11. https://doi.org/10.7554/eLife.74162
He, P., Shan, L., Lin, N. C., Martin, G. B., Kemmerling, B., Nurnberger, T., & Sheen, J. (2006). Specific
bacterial suppressors of MAMP signaling upstream of MAPKKK in Arabidopsis innate immunity.
Cell, *125*(3), 563-575. https://doi.org/10.1016/j.cell.2006.02.047
Hou, S., Wang, X., Chen, D., Yang, X., Wang, M., Turrà, D., Di Pietro, A., and Zhang, W. (2014). The
secreted peptide PIP1 amplifies immunity through Receptor-Like Kinase 7. PLoS Pathog 10:
e1004331.
Huffaker, A., Pearce, G., & Ryan, C. A. (2006). An endogenous peptide signal in Arabidopsis activates
components of the innate immune response. *Proc Natl Acad Sci U S A*, *103*(26), 10098-10103.
https://doi.org/10.1073/pnas.0603727103
Kadota, Y., Shirasu, K., & Zipfel, C. (2015). Regulation of the NADPH Oxidase RBOHD During Plant
Immunity. *Plant and Cell Physiology*, *56*(8), 1472-1480. https://doi.org/10.1093/pcp/pcv063
Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, J. D., Shirasu, K.,
Menke, F., Jones, A., & Zipfel, C. (2014). Direct regulation of the NADPH oxidase RBOHD by the
PRR-associated kinase BIK1 during plant immunity. *Mol Cell*, *54*(1), 43-55.
https://doi.org/10.1016/j.molcel.2014.02.021
Keinath, N. F., Waadt, R., Brugman, R., Schroeder, J. I., Grossmann, G., Schumacher, K., & Krebs, M.
(2015). Live Cell Imaging with R-GECO1 Sheds Light on flg22- and Chitin-Induced Transient

[Ca(2+)]cyt Patterns in Arabidopsis. *Mol Plant*, *8*(8), 1188-1200. https://doi.org/10.1016/j.molp.2015.05.006

Kissoudis, C., Sunarti, S., van de Wiel, C., Visser, R. G., van der Linden, C. G., & Bai, Y. (2016). Responses to combined abiotic and biotic stress in tomato are governed by stress intensity and resistance mechanism. *J Exp Bot*, *67*(17), 5119-5132. https://doi.org/10.1093/jxb/erw285

Knight, H., Trewavas, A. J., & Knight, M. R. (1997). Calcium signalling in Arabidopsis thaliana responding to drought and salinity. *Plant J*, *12*(5), 1067-1078. https://doi.org/10.1046/j.1365-313x.1997.12051067.x

Knight, M. R., Campbell, A. K., Smith, S. M., & Trewavas, A. J. (1991). Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature*, *352*(6335), 524-526. https://doi.org/10.1038/352524a0

Krebs, M., & Schumacher, K. (2013). Live cell imaging of cytoplasmic and nuclear Ca2+ dynamics in Arabidopsis roots. *Cold Spring Harbor Protocols*, *2013*(8), pdb. prot073031.

Kudla, J., Becker, D., Grill, E., Hedrich, R., Hippler, M., Kummer, U., Parniske, M., Romeis, T., & Schumacher, K. (2018). Advances and current challenges in calcium signaling. *New Phytol*, *218*(2), 414-431. https://doi.org/10.1111/nph.14966

Kumpf, R. P., Shi, C. L., Larrieu, A., Sto, I. M., Butenko, M. A., Peret, B., Riiser, E. S., Bennett, M. J., & Aalen, R. B. (2013). Floral organ abscission peptide IDA and its HAE/HSL2 receptors control cell separation during lateral root emergence. *Proc Natl Acad Sci U S A*, *110*(13), 5235-5240. https://doi.org/10.1073/pnas.1210835110

Kärkönen, A., & Kuchitsu, K. (2015). Reactive oxygen species in cell wall metabolism and development in plants. *Phytochemistry*, *112*, 22-32. https://doi.org/https://doi.org/10.1016/j.phytochem.2014.09.016

Ladwig, F., Dahlke, R. I., Stuhrwohldt, N., Hartmann, J., Harter, K., & Sauter, M. (2015). Phytosulfokine Regulates Growth in Arabidopsis through a Response Module at the Plasma Membrane That Includes CYCLIC NUCLEOTIDE-GATED CHANNEL17, H+-ATPase, and BAK1. *Plant Cell*, *27*(6), 1718-1729. https://doi.org/10.1105/tpc.15.00306

Lee, Y., Yoon, T. H., Lee, J., Jeon, S. Y., Lee, J. H., Lee, M. K., Chen, H., Yun, J., Oh, S. Y., Wen, X., Cho, H. K., Mang, H., & Kwak, J. M. (2018). A Lignin Molecular Brace Controls Precision Processing of Cell Walls Critical for Surface Integrity in Arabidopsis. *Cell*, *173*(6), 1468-1480.e1469. https://doi.org/10.1016/j.cell.2018.03.060

Li, L., Li, M., Yu, L., Zhou, Z., Liang, X., Liu, Z., Cai, G., Gao, L., Zhang, X., Wang, Y., Chen, S., & Zhou, J. M. (2014). The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to control plant immunity. *Cell Host Microbe*, *15*(3), 329-338. https://doi.org/10.1016/j.chom.2014.02.009

Luna, E., Pastor, V., Robert, J., Flors, V., Mauch-Mani, B., & Ton, J. (2011). Callose deposition: a multifaceted plant defense response. *Mol Plant Microbe Interact*, *24*(2), 183-193. https://doi.org/10.1094/mpmi-07-10-0149

Ma, Y., Walker, R. K., Zhao, Y., & Berkowitz, G. A. (2012). Linking ligand perception by PEPR pattern recognition receptors to cytosolic Ca2+ elevation and downstream immune signaling in plants. *Proc Natl Acad Sci U S A*, *109*(48), 19852-19857. https://doi.org/10.1073/pnas.1205448109

Matsubayashi, Y. (2011). Post-translational modifications in secreted peptide hormones in plants. *Plant Cell Physiol*, *52*(1), 5-13. https://doi.org/10.1093/pcp/pcq169

McKenna, J. F., Rolfe, D. J., Webb, S. E. D., Tolmie, A. F., Botchway, S. W., Martin-Fernandez, M. L., Hawes, C., & Runions, J. (2019). The cell wall regulates dynamics and size of plasma-membrane nanodomains in Arabidopsis. *Proc Natl Acad Sci U S A*, *116*(26), 12857-12862. https://doi.org/10.1073/pnas.1819077116

Meng, X., Zhou, J., Tang, J., Li, B., de Oliveira, M. V., Chai, J., He, P., & Shan, L. (2016). Ligand-Induced Receptor-like Kinase Complex Regulates Floral Organ Abscission in Arabidopsis. *Cell Rep*, *14*(6), 1330-1338. https://doi.org/10.1016/j.celrep.2016.01.023

Meng, X., Zhou, J., Tang, J., Li, B., de Oliveira, M. V. V., Chai, J., He, P., & Shan, L. (2016). Ligand-Induced Receptor-like Kinase Complex Regulates Floral Organ Abscission in Arabidopsis. *Cell reports*, *14*(6), 1330-1338. https://doi.org/10.1016/j.celrep.2016.01.023

Monshausen, G. B., Messerli, M. A., & Gilroy, S. (2008). Imaging of the Yellow Cameleon 3.6 indicator reveals that elevations in cytosolic Ca2+ follow oscillating increases in growth in root hairs of Arabidopsis. *Plant Physiol*, *147*(4), 1690-1698. https://doi.org/10.1104/pp.108.123638

Niederhuth, C., Patharkar, O. R., & Walker, J. (2013). Transcriptional profiling of the Arabidopsis abscission mutant hae hsl2 by RNA-Seq. *BMC Genomics*, *14*(1), 37. http://www.biomedcentral.com/1471-2164/14/37

Olsson, V., Joos, L., Zhu, S., Gevaert, K., Butenko, M. A., & De Smet, I. (2018). Look Closely, the Beautiful May Be Small: Precursor-Derived Peptides in Plants. *Annu Rev Plant Biol*. https://doi.org/10.1146/annurev-arplant-042817-040413

Patharkar, O. R., Gassmann, W., & Walker, J. C. (2017). Leaf shedding as an anti-bacterial defense in Arabidopsis cauline leaves. *PLoS Genet*, *13*(12), e1007132. https://doi.org/10.1371/journal.pgen.1007132

Patharkar, O. R., & Walker, J. C. (2015). Floral organ abscission is regulated by a positive feedback loop. *Proc Natl Acad Sci U S A*, *112*(9), 2906-2911. https://doi.org/10.1073/pnas.1423595112

Patharkar, O. R., & Walker, J. C. (2016). Core Mechanisms Regulating Developmentally Timed and Environmentally Triggered Abscission. *Plant Physiol*, *172*(1), 510-520. https://doi.org/10.1104/pp.16.01004

Poncini, L., Wyrsch, I., Dénervaud Tendon, V., Vorley, T., Boller, T., Geldner, N., Métraux, J. P., & Lehmann, S. (2017). In roots of Arabidopsis thaliana, the damage-associated molecular pattern AtPep1 is a stronger elicitor of immune signalling than flg22 or the chitin heptamer. *PLoS One*, *12*(10), e0185808. https://doi.org/10.1371/journal.pone.0185808

Qi, Z., Verma, R., Gehring, C., Yamaguchi, Y., Zhao, Y., Ryan, C. A., & Berkowitz, G. A. (2010). Ca2+ signaling by plant Arabidopsis thaliana Pep peptides depends on AtPepR1, a receptor with guanylyl cyclase activity, and cGMP-activated Ca2+ channels. *Proc Natl Acad Sci U S A*, *107*(49), 21193-21198. https://doi.org/10.1073/pnas.1000191107

Ranf, S., Eschen-Lippold, L., Pecher, P., Lee, J., & Scheel, D. (2011). Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns. *Plant J*, *68*(1), 100-113. https://doi.org/10.1111/j.1365-313X.2011.04671.x

Ranf, S., Grimmer, J., Poschl, Y., Pecher, P., Chinchilla, D., Scheel, D., & Lee, J. (2012). Defense-related calcium signaling mutants uncovered via a quantitative high-throughput screen in Arabidopsis thaliana. *Mol Plant*, *5*(1), 115-130. https://doi.org/10.1093/mp/ssr064

Robatzek, S., Chinchilla, D., & Boller, T. (2006). Ligand-induced endocytosis of the pattern recognition receptor FLS2 in Arabidopsis. *Genes Dev*, *20*(5), 537-542. https://doi.org/10.1101/gad.366506

Roberts, J. A., Whitelaw, C. A., Gonzalez-Carranza, Z. H., & McManus, M. T. (2000). Cell Separation Processes in Plants- Models, Mechanisms and Manipulation. *Annals of Botany*, *86*, 223-235.

Roman, A. O., Jimenez-Sandoval, P., Augustin, S., Broyart, C., Hothorn, L. A., & Santiago, J. (2022). HSL1 and BAM1/2 impact epidermal cell development by sensing distinct signaling peptides. *Nat Commun*, *13*(1), 876. https://doi.org/10.1038/s41467-022-28558-4

Rzemieniewski, J., Leicher, H., Lee, H. K., Broyart, C., Nayem, S., Wiese, C., Maroschek, J., Camgöz, Z., Lalun, V. O., Djordjevic, M. A., Vlot, A. C., Hückelhoven, R., Santiago, J., & Stegmann, M. (2022). Phytocytokine signaling integrates cell surface immunity and nitrogen limitation. *bioRxiv*, 2022.2012.2020.521212. https://doi.org/10.1101/2022.12.20.521212

Sanders, D., Brownlee, C., & Harper, J. F. (1999). Communicating with calcium. *Plant Cell*, *11*(4), 691-706. https://doi.org/10.1105/tpc.11.4.691

Santiago, J., Brandt, B., Wildhagen, M., Hohmann, U., Hothorn, L. A., & Butenko, M. A. (2016).
Mechanistic insight into a peptide hormone signaling complex mediating floral organ
abscission. *Elife.*, *5*. https://doi.org/10.7554/eLife.15075

Scherzer, S., Maierhofer, T., Al-Rasheid, K. A. S., Geiger, D., & Hedrich, R. (2012). Multiple calcium-
dependent kinases modulate ABA-activated guard cell anion channels. *Mol Plant*, *5*(6), 1409-
1412. https://doi.org/10.1093/mp/sss084

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,
C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P.,
& Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods*,
9(7), 676-682. https://doi.org/10.1038/nmeth.2019

Shi, C.-L., von Wangenheim, D., Herrmann, U., Wildhagen, M., Kulik, I., Kopf, A., Ishida, T., Olsson, V.,
Anker, M. K., Albert, M., Butenko, M. A., Felix, G., Sawa, S., Claassen, M., Friml, J., & Aalen, R.
B. (2018). The dynamics of root cap sloughing in Arabidopsis is regulated by peptide signalling.
Nat Plants, *4*(8), 596-604. https://doi.org/10.1038/s41477-018-0212-z

Smakowska-Luzan, E., Mott, G. A., Parys, K., Stegmann, M., Howton, T. C., Layeghifard, M., Neuhold,
J., Lehner, A., Kong, J., Grunwald, K., Weinberger, N., Satbhai, S. B., Mayer, D., Busch, W.,
Madalinski, M., Stolt-Bergner, P., Provart, N. J., Mukhtar, M. S., Zipfel, C., . . . Belkhadir, Y.
(2018). An extracellular network of Arabidopsis leucine-rich repeat receptor kinases. *Nature*,
553(7688), 342-346. https://doi.org/10.1038/nature25184

Somssich, M., Bleckmann, A., & Simon, R. (2016). Shared and distinct functions of the pseudokinase
CORYNE (CRN) in shoot and root stem cell maintenance of Arabidopsis. *J Exp Bot*, *67*(16), 4901-
4915. https://doi.org/10.1093/jxb/erw207

Steinhorst, L., & Kudla, J. (2013). Calcium and reactive oxygen species rule the waves of signaling. *Plant*
Physiol, *163*(2), 471-485. https://doi.org/10.1104/pp.113.222950

Stenvik, G. E., Tandstad, N. M., Guo, Y., Shi, C. L., Kristiansen, W., & Holmgren, A. (2008). The EPIP
peptide of INFLORESCENCE DEFICIENT IN ABSCISSION is sufficient to induce abscission in
Arabidopsis through the receptor-like kinases HAESA and HAESA-LIKE2. *Plant Cell.*, *20*.
https://doi.org/10.1105/tpc.108.059139

Sto, I. M., Orr, R. J., Fooyontphanich, K., Jin, X., Knutsen, J. M., Fischer, U., Tranbarger, T. J., Nordal, I.,
& Aalen, R. B. (2015). Conservation of the abscission signaling peptide IDA during Angiosperm
evolution: withstanding genome duplications and gain and loss of the receptors HAE/HSL2.
Front Plant Sci, *6*, 931. https://doi.org/10.3389/fpls.2015.00931

Taylor, I., & Walker, J. C. (2018). Transcriptomic evidence for distinct mechanisms underlying abscission
deficiency in the Arabidopsis mutants haesa/haesa-like 2 and nevershed. *BMC Res Notes*,
11(1), 754. https://doi.org/10.1186/s13104-018-3864-x

Torres, M. A., & Dangl, J. L. (2005). Functions of the respiratory burst oxidase in biotic interactions,
abiotic stress and development. *Curr Opin Plant Biol*, *8*(4), 397-403.
https://doi.org/10.1016/j.pbi.2005.05.014

Vie, A. K., Najafi, J., Liu, B., Winge, P., Butenko, M. A., Hornslien, K. S., Kumpf, R., Aalen, R. B., Bones,
A. M., & Brembu, T. (2015). The IDA/IDA-LIKE and PIP/PIP-LIKE gene families in Arabidopsis:
phylogenetic relationship, expression patterns, and transcriptional effect of the PIPL3 peptide.
Journal of Experimental Botany, *66*(17), 5351-5365. https://doi.org/10.1093/jxb/erv285

Wang, Y., Li, X., Fan, B., Zhu, C., & Chen, Z. (2021). Regulation and Function of Defense-Related Callose
Deposition in Plants. *Int J Mol Sci*, *22*(5). https://doi.org/10.3390/ijms22052393

 # Main figures - A dual function of the IDA peptide in regulating cell

 separation and modulating plant immunity at the molecular level

Fig. 1: mIDA-induced [Ca^{2+}]$_{cyt}$ release in Arabidopsis roots correlates with *pHAE* and *pHSL2* activity

a, Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI) (upper panel, outlined in red) in the meristematic and elongation zone of the root. Fluorescence intensities (ΔF/F) over time of the whole root represented in a heat map (lower panel). Shown are cytosolic calcium concentration ([Ca^{2+}]$_{cyt}$) dynamics in the ROI in response to 1 µM mIDA over time. (see also Movie 1). Red lines at 5 minutes (min) indicates application of mIDA peptide or application of eATP at 22 min. Representative response from 10 roots (Sup Fig. 2). The increase in [Ca^{2+}]$_{cyt}$ response propagates through the roots as two waves. **b,** For comparison; Normalized R-GECO1 fluorescence intensities (ΔF/F) measured from regions of interest (ROI) (outlined in red, upper panel) in response to 1µM flg22 over time. Fluorescence intensities (ΔF/F) over time of the whole root represented in a heat map (lower panel). Red line at 0 min (min) indicates application of flg22 peptide. Representative response from 7 roots (Sup Fig. 3). The increase in [Ca^{2+}]$_{cyt}$ response propagates through the roots as a single wave seen as normalized R-GECO1 fluorescence intensities (ΔF/F) shown as a heat map (see also movie 2). **c,d,e,** Expression of the receptors **c,** *pHAE:Venus-H2B* **d,** *pHSL2:Venus-H2B* and **e,** *pFLS2:Venus-H2B* in 7 days-old roots. Representative pictures of n = 8. Scale bar = 50 µm, single plane image, magenta = propidium iodide stain.

Fig. 2 : mIDA-induced $[Ca^{2+}]_{cyt}$ release in Arabidopsis abscission zones.

a, Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI)

(outlined in red) in floral abscission zone (AZ)s. Shown are cytosolic calcium concentration ($[Ca^{2+}]_{cyt}$)

dynamics in the ROI in response to 1 µM mIDA over time (see also Movie 3). Representative response

from 8 flowers (Sup. Fig 6). Red lines at 5 minutes (min) indicates application of mIDA peptide or

application of eATP at 35 min (for AZs), respectively. The increase in $[Ca^{2+}]_{cyt}$ response propagates

through the AZ as a single wave. **b**, Expression of *pHAE:Venus-H2B* and *pHSL2:Venus-H2B* in flowers at

position 6 (See Sup Fig. 5a for positions) (arrowhead indicates AZ). Representative pictures of n = 8,

scale bar = 100 µm, maximum intensity projections of z-stacks.

Fig. 3: *IDA* is induced by biotic and abiotic stress

Representative pictures of *pIDA:GUS* expression after 12 h treatment with 1 μM flg22, chitin, NaCl,

Mannitol and 1 μM IDAΔN69 in **a,** cells surrounding emerging lateral roots, and **b,** in the main root . **c,**

Normalized emitted fluorescence of fluorochrome 4-methyl umbelliferone (4-MU) in 7 days-old

seedlings after 12 h treatment with 1 μM flg22, chitin, NaCl, Mannitol and 1 μM of the inactive IDA

peptide, IDAΔN69. Normalized to 1 on No Treatment sample. Controls were not subjected to any stimuli

(No treatment). n = 6, experiment repeated 3 times. * = significantly different from the non-treated (No

treatment) sample ($p < 0.05$, student t-test, two tailed). Controls were not subjected to any stimuli (No

treatment). **a,b,** Representative picture of n = 10, experiment repeated 3 times, scale bar = 50 μm.

Fig. 4: mIDA induce callose deposition in Arabidopsis cotelydons expressing 35S::HAE-YFP

Callose deposition in Col-0 WT, 35S:HAE in Col-0 and *fls2* treated with water (mock treatment), 1 μM

mIDA or 1uM flg22. **a**, Callose deposition could be observed in cotyledons of eight day old Col-0 WT and

35S:HAE in seedlings treated with 1 μM flg22, and to a smaller extend in 35S:HAE treated with 1uM

mIDA. No callose deposition could be detected in *fls2* mutant treated with…… Representative images

of 9-12 seedlings per genotype. Scale bar 500 μm. **b**, Total callose depositions for the different

genotypes treated with water (mock treatment), 1 μM mIDA or 1 μM flg22. Statistically significant

difference at ($p < 0.05$). N = 9-12

Fig. 5: mIDA induced transcription of defense-associated marker genes

a, Transcripts of *FRK1*, *MYB51*, and *PEP3* in WT Col-0 seedlings exposed to 1 μM mIDA for 1 h (red) and

to 1 μM mIDA for 12 h (blue) compared to untreated tissue (black). RNA levels were measured by RT-

qPCR analysis. *ACTIN* was used to normalize mRNA levels Figure represent three biological replicates

with four technical replicates. Statistical analyses was performed comparing induction times of

individual genes using one-way ANOVA and post-hoc Tukey's test (p < 0.05). **b**,*pMYB51:YFPN* expression

is enhanced in roots after 7 h exposure to 1 μM mIDA peptide compared to untreated roots (control),

scale bar = 50 μm, maximum intensity projections of z-stacks, magenta = propidium iodide stain.

Fig. 6: mIDA and flg22 co-treatment extremely enhances the transcription of defense-associated marker

genes

a, RT-qPCR data showing transcription of *FRK1*, *MYB51*, and *PEP3* in Col-0 WT seedlings exposed to 1

µM mIDA, 1 µM flg22 or a combination of 1 µM mIDA and 1 µM flg22 for 1 h compared to untreated

seedlings (No treatment control). *ACTIN* was used to normalize mRNA levels. Figure represent three

biological replicates with four technical replicates. Statistical analyses comparing No Treatment to

peptide treated samples was performed using one-way ANOVA and post-hoc Tukey's test ($p < 0.05$). **b,**

Microscopic analysis of 7-days-old *pFLS2:Venus-H2B pHAE:Tomato-H2B* expressing plants of root tip

(upper panel) and lateral root (middle panel) of 7 days old plants, and flowers at position 6 (lower

panel). Fluorescent nuclei representing co-expression of the Venus and Tomato marker could be

observed in cells surrounding emerging LRs, in the stele of the root, as well as in the abscission zone.

c, Microscopic analysis of 7-days-old *pFLS2:Venus-H2B pHSL2:Tomato-H2B* expressing plants of root

tip (upper panel) and lateral root (middle panel) of 7 days old plants, and flowers at position 6 (lower

panel). Fluorescent nuclei representing co-expression of the Venus and Tomato marker could be

observed in the root tip, in cells surrounding emerging LRs, as well as in the abscission zone. Root

pictures scale bar = 50 µm, single plane image. Abscission zone images= maximum intensity

projections of z-stacks. Scale = 50µm.

Fig. 7: IDA regulates cell separation processes, but is also involved in major transcription of defense genes upon pathogen attack.

a, IDA and the IDL peptides control cell separation processes during plant development and in response to abiotic and biotic stress (Butenko et al. 2003; Patharkar and Walker 2016; Patharkar et al. 2017; Shi et al. 2018). Tissue undergoing cell separation includes floral organ abscission, cauline leaf abscission, emerging of lateral roots and root cap sloughing. **b,** During normal conditions, IDA control floral organ abscission and emergence of lateral roots by relaying a signal through receptor complexes including HAE, HSL2, SERK1 and BAK1 to modulate the expression of cell wall remodeling (CWR) genes as well as moderately expression of defense genes. IDA activates a receptor dependent production of ROS and an increase in $[Ca^{2+}]_{cyt}$. **b,** Upon stress, such as pathogen attack, the activation of HAE and HSL2 acts in addition to activation of defense related receptors, such as FLS2, to enhance the expression of defense related genes significantly. This ensure optimal protection of cells undergoing cell separation, which may be major entry routes during a pathogen attack.

Supplementary figures - A dual function of the IDA peptide in regulating cell separation and modulating plant immunity at the molecular level

Sup Fig. 1: IDA induces ROS production in Arabidopsis

a, ROS production detected by the luminol-based assay was monitored over time as RLU. ROS production from *hae hsl2* leaf disks expressing *35S:HAE-YFP* in response to 1 μM mIDA (black) and in WT control (blue). **b**, Fluorescent image of *hae hsl2* rosette leaves expressing *35S:HAE:YFP*. Signal is detected in the plasma membrane of cells in the epidermal layer. Representative picture. Scale bar = 50 μm. **c**, *35S:HAE-YFP* complements the *hae hsl2* mutant abscission phenotype of observed. Scale = 1 cm. **d,e**, Microscopic analysis of *pHAE:Venus-H2B* (upper panel) and *pHSL2-Venus-H2B* (lower panel) in **d**, cotelydons of 7 days old seedlings and in **e**, rosette leaves of a 22 days old Arabidopsis plant containing 7 true leaves. Expression in each leaf is shown from youngest (left) to oldest (right).Scale = 100 μm

Sup Fig. 2: mIDA-induced [Ca^{2+}]$_{cyt}$ release in Arabidopsis roots

998 Shown are cytosolic calcium concentration ([Ca^{2+}]$_{cyt}$) dynamics in the ROI in response to 1 μM mIDA

999 over time. Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from a region of

1000 interest (ROI) containing the meristematic and elongation zone of the root.

1001

1002

1003   **Sup Fig. 3: flg22-induced $[Ca^{2+}]_{cyt}$ release in Arabidopsis roots**

1004   Shown are cytosolic calcium concentration ($[Ca^{2+}]_{cyt}$) dynamics in the ROI in response to 1 µM flg22

1005   over time. Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from a region of

1006   interest (ROI) containing the meristematic and elongation zone of the root.

1007

1008

1009

1010

1011

1012

1013

Sup Fig. 4 : The IDA triggered increase in $[Ca2+]_{cyt}$ is abolished in the presence of Ca^{2+} inhibitors

a, Increase in $[Ca^{2+}]_{cyt}$ in 7 days old seedlings expressing the cytosolic localized *Aequorin*-based

luminescence Ca^{2+} sensor (*Aeq*) measured in relative light units (RLU) treated with 1 µM mIDA (black).

No response is observed in *Aeq* seedlings treated with 1 µM IDAΔN69 (gray) or in *hae hsl2 Aeq*

seedlings treated with 1 µM mIDA (blue). **b,** Increase in $[Ca^{2+}]_{cyt}$ in 7 days old seedlings expressing the

cytosolic localized *Aequorin*-based luminescence Ca^{2+} sensor (*Aeq*) measured in relative light units

(RLU) treated with 1 µM mIDA . No response is observed in *Aeq* seedlings treated with 1 mM EGTA

(blue) or 1mM $LaCl_3$ (grey). Curves represent average of 3 independent experiments with 4-6

replicates in each experiment.

1023

1024

Sup Fig. 5: mIDA induces a Ca²⁺ response in flowers

a, Arabidopsis flowers and siliques at different developmental stages. Flowers along the main

inflorescence are counted from the first flower with visible white petals at the top of the inflorescence

and is defined as p 1. Subsequently older flowers along the inflorescence (p0-p8). Cell wall remodeling

in AZ cells, accompanied with a reduction in petal breakstrength (the force required to remove a petal

from the receptacle) is initiated at p6 (red arrow). By p7 AZ cells have undergone cell separation and

the floral organs have abscised. P=position **b**, Increase in [Ca²⁺]$_{cyt}$ in flowers expressing the cytosolic

localized *Aequorin*-based luminescence Ca²⁺ sensor (*Aeq*) measured in relative light units (RLU) treated

with 1 µM mIDA. A [Ca²⁺]$_{cyt}$ increase is observed in flowers where there is an initial weakening of the

cell walls (p5-6) (black). No increase in luminescence was observed in *Aeq* expressing flowers of an

earlier developmental stage (p3) (gray). Curves show representative measurements. Three

independent experiments were performed with 4-6 replicates in each experiment.

1037

1038

1039   **Sup Fig. 6: mIDA-induced $[Ca^{2+}]_{cyt}$ release in Arabidopsis abscission zones.** Normalized R-GECO1

1040   fluorescence intensities (ΔF/F) were measured from regions of interest (ROIs) in floral abscission zone

1041   (AZ)s. Shown are cytosolic calcium concentration ($[Ca^{2+}]_{cyt}$) dynamics in the ROI1-5 in response to 1 μM

1042   mIDA over time.

1043

1044

1045

Sup Fig. 7 : mIDAΔN69 does not induce a $[Ca^{2+}]_{cyt}$ release in Arabidopsis abscission zones (AZ)

Normalized R-GECO1 fluorescence intensities (ΔF/F) were measured from regions of interest (ROI)

(outlined in red) in the floral AZs. Shown are cytosolic calcium concentration ($[Ca^{2+}]_{cyt}$) dynamics in the

ROI in response to application of 1 μM mIDAΔN69 or eATP over time. Red lines at 5 minutes (min)

indicates application of mIDAΔN69 peptide or application of eATP at 30 min (see also Movie 4).

Representative response from 10 flowers. The increase in $[Ca^{2+}]_{cyt}$ response to eATP propagates

through the AZ as a single wave seen as normalized R-GECO1 fluorescence intensities (ΔF/F) shown as

a heat map.

1054

1055

1056

1057

Sup Fig. 8: Investigation of CNGCs and RBOHs in the abscission process

a, Relative expression of *CNGC1*, *CNGC2*, *CNGC4*, *CNGC5*, *CNGC6*, *CNGC9*, *CNGC12, RBOHD* and *RBOHF*

in the floral organs during the onset of abscission (Cai and lashbrook 2008). X-axis represents the

developmental stages of abscission where position 12 represent flowers where the stamen and petals

are of the same length, position 13 represent flowers at anthesis, position 15 represent flowers with

stigma extended above the anthers. Floral organ abscission occurs at position 15. Relative expression

of *HAE* and *HSL2* added for comparison. Nomenclature for the abscission process and relative

expression values taken from (Cai and Lashbrook 2008). b, No floral organ abscission phenotype is

observed in the single *cngc* mutant plants compared to the *hae hsl2* mutant which retains the floral

organ indefinitely. Scale bar = 2cm, 5 weeks old plants. c, ROS production in the AZ (white arrow head)

detected by using the fluorescent dye H_2DCF-DA in WT and *rbohd rbohf* flowers at position 6 (See

Supplementary Fig. 5a for positions). Scale bar = 100 µm, maximum intensity projections of z-stacks.

Representative pictures from 9 flowers. **d**, Representative pictures of WT and *rbohd rbohf*

inflorescences, scale bar for inflorescences = 1 cm. **e**, Petal break strength (pBS) measurements of WT

and the *rbohd rbohf* mutant at position 1-6 along the inflorescence. Force required to remove petals

from the receptacle represented in gram equivalent. * = significantly different from WT at the given

position ($p < 0.05$, student t-test, two tailed).

1075

Sup Fig. 9: mIDA induced enhanced transcription of defense-associated marker genes is only partially

dependent on the HAE and HSL2 receptors

RT-qPCR data showing transcription of *FRK1*, *MYB51*, and *PEP3* in Col-0 WT and *hae hsl2* seedlings

exposed to 1 µM mIDA for 1 h compared to untreated seedlings (No treatment). *ACTIN* was used to

normalize mRNA levels. Figure represent 3 biological replicates with 4 technical replicates. Statistical

analyses was performed comparing samples within same gene, using two-way ANOVA and post-hoc

Tukey's test ($p < 0.05$).

1083

1084

1085

Sup Fig. 10: mIDA and flg22 induced expression of *pMYB51:YFP*

Representative microscopic pictures of 7-days-old *pMYB51:YFPN* roots from seedlings transferred to

liquid medium with 1 µM flg22 or 1 µM mIDA for 1, 3, 5 and 7 h using a fluorescent microscope.

Control seedlings were transferred to medium with no peptide and imaged in the same time frame.

Fluorescent nuclei could be observed in a similar temporal pattern in roots subjected to mIDA and

flg22. Scale bar = 100 µm, single plane images using a Zeiss Axioplan2 microscope with an AxioCam

HRc, 20x air objective and a YFP filter (Excitation: 500/20, Beamsplitter: FT515, Emission: 535/50), no

imaging processing was performed, t = time in h. n = 3, experiment repeated 3 times.

1094

1095

1096

1097

Sup Fig. 11 : mIDA and PIP1 co-treatment do not enhances transcription of defense-associated

marker genes

RT-qPCR data showing transcription of *FRK1*, *MYB51*, and *PEP3* in Col-0 WT seedlings exposed to 1 µM

mIDA, 1 µM PIP1 or a combination of 1 µM mIDA and 1 µM PIP1 for 1 h compared to untreated

seedlings (No treatment control). *ACTIN* was used to normalize mRNA levels. Figure showing extended

results from experiment shown in Fig 6. For comparison, No treatment and mIDA treated samples

were included in the figure (same results presented in Fig 6). Scale on y axis was kept as in Fig 6.

Figure represents 2-3 biological replicates with 4 technical replicates. Statistical analyses comparing

the samples was performed using one-way ANOVA and post-hoc Tukey's test ($p < 0.05$).

1107

1108

1109

Sup Fig. 12: HAE (yellow) and HSL2 (red) display very distinct repertoire of immune and growth-related

interacting LRR-RKs.

HEA and HSL2 specific interacting partners determined using a sensitized high-throughput

extracellular domain interaction assay (Smakowska-Luzan et al,.2018). HSL2 and HAE show distinct

interacting LRR-RKs where only HSL2 was found to interact with RLK7 (red halo). * indicates receptors

known to play a function in biotic and/or abiotic responses in planta based on previous published

work, dark blue nodes refer to bidirectional interactions with GeoMean ≥2.5, light blue nodes- refer to

bidirectional interactions with GeoMean ≤2.5, and grey nodes refer to unidirectional interactions.

1118

1119

1120

1121

1122