Non-parallel transcriptional divergence during parallel adaptation

Running title: Transcriptional mechanisms of parallelism

Eva K Fischer[1*,2], Youngseok Song[3], Kimberly A Hughes[4], Wen Zhou[3], Kim L Hoke[2]

[1] Department of Evolution, Ecology, and Behavior, University of Illinois, Urbana, IL 61801, USA

[2] Department of Biology, Colorado State University, Fort Collins, CO 80523, USA

[3] Department of Statistics, Colorado State University, Fort Collins, CO 80523, USA

[4] Department of Biological Science, Florida State University, Tallahassee, FL 32306 USA

*Corresponding author:

Eva K Fischer

efischer@illinois.edu

Department of Evolution, Ecology, and Behavior

University of Illinois at Urbana-Champaign

Morrill Hall, 505 S. Goodwin Avenue

Urbana, IL 61801

Abstract

How underlying mechanisms bias evolution toward predictable outcomes remains an area of active debate. In this study, we leveraged phenotypic plasticity and parallel adaptation across independent lineages of Trinidadian guppies (*Poecilia reticulata*) to assess the predictability of gene expression evolution during parallel adaptation. Trinidadian guppies have repeatedly and independently adapted to high- and low-predation environments in the wild. We combined this natural experiment with a laboratory breeding design to attribute transcriptional variation to the genetic influences of population of origin and developmental plasticity in response to rearing with or without predators. We observed substantial gene expression plasticity as well as the evolution of expression plasticity itself across populations. Genes exhibiting expression plasticity *within* populations were more likely to also differ in expression *between* populations, with the direction of population differences more likely to be opposite those of plasticity. While we found more overlap than expected by chance in genes differentially expressed between high- and low-predation populations from distinct evolutionary lineages, the majority of differentially expressed genes were not shared between lineages. Our data suggest alternative transcriptional configurations associated with shared phenotypes, highlighting a role for transcriptional flexibility in the parallel phenotypic evolution of a species known for rapid adaptation.

Key words: *Poecilia reticulata*, transcriptomics, phenotypic plasticity, adaptation, parallel evolution

Introduction

Phenotypic evolution is biased by the mechanisms that link genetic variation to phenotypic variation, i.e. the genotype-phenotype map (Alberch, 1991; Pigliucci, 2010). The complex nature of the genotype-phenotype map, the existence of 'many-to-one' mappings across hierarchical levels of biological organization, and the multigenic nature of complex traits make it challenging to assess how mechanistic properties shape evolutionary outcomes (Boyle, Li, & Pritchard, 2017; Potticary, Morrison, & Badyaev, 2020). A deeper understanding of these phenomena will further our insight into the forces limiting and potentiating the evolution of complex traits and deepen our knowledge of how organisms respond to novel and changing environments.

Studies focusing on the independent evolution of shared traits can reveal biases in mechanisms of adaptation by asking whether similar mechanism are associated with repeated, independent evolutionary transitions. Shared phenotypes arising in response to similar selection pressures are referred to as "parallel" or "convergent" depending on the evolutionary relatedness of populations and author preference (see Glossary). Different definitions have been discussed and debated elsewhere (e.g. Arendt & Reznick, 2008; Losos, 2011; Wake, Wake, & Specht, 2011), and we use "parallel" here following the terminology conventionally applied to our study system (see below).

Understanding whether shared phenotypes also share underlying genetic and molecular mechanisms is a long-stranding focus of evolutionary biology (reviewed in Bolnick, Barrett, Oke, Rennison, & Stuart, 2018; Stern, 2013). Compelling examples of parallelism and convergence demonstrate that similar phenotypes may share underlying neural, physiological, molecular, and/or genetic mechanisms, even across highly divergent taxa (e.g. Insel and Young, 2000; Manceau et al., 2010; Pankey et al., 2014; Rosenblum et al., 2010). Similar mechanisms underlying parallel evolutionary transitions suggest the genotype-phenotype map is constrained by limits on the possible ways to construct adaptive phenotypes (Losos, 2011). In this case, phenotypic evolution may rely on a limited number of mechanistic paths, repeatedly targeting

those mechanisms that yield the greatest phenotypic responses with the smallest pleiotropic

costs. Indeed, the recent view of complex traits as "omnigenic" highlights the potential for near

ubiquitous pleiotropy to constrain evolutionary trajectories to a very limited number of paths (Boyle

et al., 2017).

In contrast, non-shared mechanisms giving rise to shared, independently derived

phenotypes suggest that mechanistic flexibility may facilitate evolution by providing multiple,

alternative 'solutions' to a given adaptive problem, thereby providing alternative paths by which

organisms can reach adaptive peaks (Badyaev & Morrison, 2018; Drion, O'Leary, & Marder, 2015;

Grashow, Brookings, & Marder, 2009). Empirical studies demonstrate that different mechanistic

'solutions' can give rise to shared phenotypes in closely related species, among populations of

the same species, or even among individuals of the same population (e.g. Abouheif and Wray,

2002; Crawford and Oleksiak, 2007; Drion et al., 2015; Grashow et al., 2009; Mandic et al., 2018).

Mechanistic flexibility may facilitate evolution by allowing organisms to reach the same adaptive

peak from alternative starting points (e.g. distinct genetic backgrounds), and/or buffering

organisms from pleiotropic effects via compensatory mechanisms that exploit the alternative

paths created by many-to-one mappings. Whether the potentially "omnigenic" nature of complex

traits contributes to constraints due to widespread pleiotropy or facilitates alternative paths is an

area ripe for exploration (Boyle et al., 2017).

In sum, empirical evidence for both shared and distinct mechanisms underlying parallel

phenotypic evolution leaves open the question of when and why either pattern dominates. Further

addressing this question is a focus of theoretical and empirical work, with implications for

evolutionary biology, conservation, and medicine (reviewed in Bolnick et al., 2018). In view of the

potential importance of many-to-one mappings, gene expression studies provide promising

starting points for surveying broadscale patterns of (non)parallelism in the mechanisms underlying

complex traits. Despite this promise and substantial recent interest in (non)parallel evolution

(Bolnick et al., 2018; Kaeuffer, Peichel, Bolnick, & Hendry, 2011; Oke et al., 2016; Paccard et al.,

2020; Stuart et al., 2017), gene expression mechanisms underlying parallel evolution have only

recently begun to be explored (Ghalambor et al., 2015; Manousaki et al., 2013; Pankey et al.,

2014).

Does phenotypic plasticity bias phenotypic evolution?

One feature of biological systems that may channel evolution into particular paths is

environmentally induced phenotypic plasticity. First, both plastic and evolutionary processes may

rely on a limited number of mechanistic paths, repeatedly targeting those mechanisms that yield

the greatest phenotypic responses with the smallest pleiotropic costs. In this case, we expect

plastic and evolutionary responses to share underlying mechanisms. Theory predicts that plastic

and evolutionary changes will be in the same direction when plasticity in a novel environment is

adaptive, increasing immediate survival and allowing time for evolutionary divergence via co-

option of mechanisms involved in environmentally induced responses (Baldwin, 1896;

Ghalambor, McKay, Carroll, & Reznick, 2007; Lande, 2009; West-Eberhard, 2003). In contrast,

plastic and evolutionary changes will be in opposite directions when plasticity in a novel

environment is non-adaptive, thereby increasing the strength of selection, or when plastic

responses 'overshoot' adaptive optima and are compensated by selection (Conover, Duffy, &

Hice, 2009; Ghalambor et al., 2007; Grether, 2005; Velotta & Cheviron, 2018). Importantly,

plasticity may facilitate adaptation under either scenario and empirical studies document both

adaptive (e.g. Fraser et al., 2014; Gleason & Burton, 2015; Li et al., 2018; Mäkinen, Papakostas,

Vøllestad, Leder, & Primmer, 2016; Scoville & Pfrender, 2010; Shaw et al., 2014; Wang & Althoff,

2019) and non-adaptive (e.g. Dayan, Crawford, & Oleksiak, 2015; Ghalambor et al., 2015; W. Ho

& Zhang, 2018; Pespeni et al., 2013; Schaum, Rost, Millar, & Collins, 2013) plasticity associated

with adaptive evolution. Finally, plastic and evolutionary processes may rely on shared

mechanistic paths because the mechanisms mediating phenotypic plasticity promote the

accumulation of cryptic genetic variation that is released under new environmental conditions,

thereby fostering associations between plasticity and divergence (Draghi & Whitlock, 2012;

Espinosa-Soto, Martin, & Wagner, 2011).

In the present study, we combine measures of phenotypic plasticity and parallel adaptation

to ask whether flexibility in underlying mechanisms shapes evolutionary trajectories. While the

potentially important role of phenotypic plasticity in this process has been pointed out, few studies

have explicitly considered the role of environmentally induced plasticity in parallelism (for rare

examples see Oke et al., 2016; Torres-Dowdal, Handelsman, Reznick, & Ghalambor, 2012), nor

asked how plasticity itself evolves (West-Eberhard, 2003). By comparing transcriptional

mechanisms associated with adaptation within and across parallel lineages, we are able to assess

whether plasticity biases evolutionary outcomes, to ask whether parallel phenotypic evolution is

associated with shared gene expression patterns, and to speculate on the role of mechanistic

flexibility in adaptive evolution.

Guppies as a model system

Trinidadian guppies (*Poecilia reticulata*) have become a model system in ecology and

evolutionary biology due to repeated, independent adaptation of natural guppy populations to

distinct predation environments (Endler, 1995; Haskins, Haskins, McLaughlin, & Hewitt, 1961; D.

Reznick, Butler IV, & Rodd, 2001). In Trinidad, high- and low-predation population pairs from

different river drainages represent independent evolutionary lineages (Barson, Cable, & Van

Oosterhout, 2009; Fraser, Künstner, Reznick, Dreyer, & Weigel, 2015; Gilliam, Fraser, & Alkins-

Koo, 1993; Willing et al., 2010) in which colonization of low-predation environments has led to

parallel, adaptive changes in life history traits, morphology, and behavior (Endler, 1995;

Magurran, 2005; D. A. Reznick, Bryga, & Endler, 1990; D. Reznick et al., 2001; D. N. Reznick,

1997). For example, low-predation guppies shoal less tightly (Huizinga, Ghalambor, & Reznick,

2009; Magurran & Seghers, 1990b, 1991), escape more slowly (Ghalambor, Reznick, & Walker,

2004), are slower to re-commence movement following a predator encounter (Elvidge,

Ramnarine, & Brown, 2014; Harris, Ramnarine, Smith, & Pettersson, 2010), and perform fewer

predator inspections (Magurran & Seghers, 1990a, 1994) than their high-predation counterparts.

Building on decades of work comparing high- and low-predation populations in the wild,

more recent laboratory experiments have deployed breeding designs to disentangle genetic and

environmental influences of predation on phenotypic differences. These recent studies

demonstrate that a combination of genetic and environmental influences shape guppy life history

(Torres Dowdall et al., 2012), morphology (Fischer, Soares, Archer, Ghalambor, & Hoke, 2013;

Handelsman, Ruell, Torres-Dowdall, & Ghalambor, 2014; Ruell et al., 2013; Torres-Dowdal et al.,

2012), physiology (Fischer, Harris, Hofmann, & Hoke, 2014; Handelsman et al., 2013), and

behavior (Fischer, Ghalambor, & Hoke, 2016b; Huizinga et al., 2009; Torres-Dowdal et al., 2012).

Yet despite many years of work on whole-organism phenotypes, the mechanisms underlying

adaptive phenotypic differences in guppies remain largely unexplored. A single study

characterized brain gene expression in multiple populations from the same lineage during the

earliest stages of adaptation (~3 generations after colonization of low-predation environments)

and found a negative relationship between phenotypic plasticity and adaptive divergence

(Ghalambor et al., 2015).

Transcriptional mechanisms of plasticity, evolution, and parallel adaptation

In the present study, we take advantage of parallel phenotypic evolution in independent

lineages of Trinidadian guppies (*Poecilia reticulata*) to explore patterns of flexibility and constraint

in transcriptional mechanisms mediating repeated adaptation. We compare the effects of genetic

background (high- versus low-predation populations) and developmental environment (rearing

with and without predator cues) on brain gene expression patterns in two parallel, independent

evolutionary lineages. These lineages diverged at least 600,000 years ago, with subsequent,

more recent colonization of low-predation environments by high-predation ancestors within each

river drainage (Fajen & Breden, 1992; Fraser et al., 2015; Willing et al., 2010). We compare whole-

brain gene expression patterns *within* and *between* lineages based on evolutionary history with

and rearing with predators. We focus on gene expression in the brain because the previous gene

expression study in guppies used brain tissue (Ghalambor et al., 2015) and because of previous

evidence for behavioral plasticity in our focal populations (Fischer et al., 2016b).

We test four hypotheses concerning the influences of underlying mechanisms on adaptive

evolutionary outcomes. **Hypothesis 1** (H1), that genes with significant expression plasticity are

more likely to show genetic divergence in expression; **Hypothesis 2** (H2), that the direction of

plastic responses predicts the direction of genetic divergence; **Hypothesis 3** (H3), that gene

expression plasticity itself evolves; and **Hypothesis 4** (H4), that parallel phenotypic adaptation

across independent lineages relies on parallel gene expression changes. To test our hypotheses

we (H1) quantify the extent of overlap in genes exhibiting significant expression plasticity as well

as genetic expression divergence within each drainage, (H2) examine associations in the

direction of developmental and genetic expression differences within each lineage, (H3) ask

whether expression plasticity itself evolves, and (H4) assess whether parallel phenotypic

adaptation between lineages relies on similar changes in gene expression profiles.

A number of non-mutually exclusive, interrelated predictions follow from our hypotheses.

If developmental plasticity in gene expression does indeed predict genetic divergence in gene

expression, we expect the magnitude (H1) and direction (H2) of gene expression plasticity to be

associated with the magnitude and direction of expression divergence between high- and low-

predation populations within a river drainage (i.e. within an evolutionary lineage). If gene

expression plasticity itself evolves (H3), we expect the extent and/or direction of plastic responses

to differ between high- and low-predation populations within a lineage. Finally, if only a few gene

expression configurations (i.e. mechanistic paths) can give rise to shared adaptive phenotypes,

we expect parallel adaptation to low-predation habitats to be characterized by parallel expression

changes in set of genes shared between lineages (H4). In contrast, if transcriptional mechanisms

are flexible, we expect gene expression divergence between high- and low-predation populations

within each river to occur in largely non-overlapping gene sets. Taken together, our results allow

us to assess the flexibility of transcriptional mechanisms of adaptation across timescales – from

developmental plasticity, to genetic divergence within rivers, to parallel adaptation across

lineages.

Material and Methods

Fish collection and rearing

We used a laboratory breeding design to disentangle the influences of evolutionary history

with predators and developmental experience with predators on brain gene expression patterns.

We established lab populations of guppies from high- and low-predation populations pairs

collected from the Aripo and Quare river drainages in 2012 and 2014, respectively. To maintain

the genetic variation of the original wild-caught fish, we established 20 - 30 unique family lines for

each population (i.e., for each generation a single female from each family was crossed to a single

male from another family) (D. N. Reznick & Bryga, 1987). We used second-generation lab born

fish from these unique family lines in this study to minimize environmental and maternal effects.

At birth, we split second-generation siblings into rearing environments with (pred+) or without

(pred-) predator chemical cues, and they remained in these environments until the completion of

the experiment (as in Fischer et al., 2016). We used only mature males in this study. To maximize

the range of genetic variation captured among focal fish, all males in a given experimental group

(i.e. population and rearing environment) were from distinct families; however, our split-brood

design allowed us to simultaneously leverage genetic similarity among siblings and control for the

effects of genetic background on developmental responses. Figure 1 provides an overview of our

experimental design and interpretation of comparisons.

All guppies were individually housed in 1.5 liter tanks on a 12:12 hour light cycle at

Colorado State University. Fish were fed a measured food diet once daily, receiving Tetramin™

tropical fish flake paste and hatched *Artemia* cysts on an alternating basis. Prior to tissue

collection for this study, behavioral and hormone data were collected in an identical manner for

all fish and these results are reported elsewhere (Fischer et al., 2016b). All experimental methods

were approved by the Colorado State University Animal Care and Use Committee (Approval #12-

3818A).

Tissue collection and processing

We collected brain tissue from the Aripo and Quare lineage males described above in

2013 and 2015, respectively. To standardize any effects of recent experience, behavior, and

circadian rhythm on gene expression, we collected whole brains within 10 minutes of lights-on in

the morning. We collected tissue from a maximum of four fish each day, one from each

experimental group per day, collected in a random order. We interpret our transcriptional data as

baseline, in the sense that fish were minimally stimulated prior to tissue collection, and expression

levels should therefore reflect genetic background and developmental experience more strongly

than responses to immediate environmental context. Fish were anesthetized by immersion in ice

water followed by rapid decapitation. Whole brains were removed, flash frozen in liquid nitrogen,

and stored at -80°C until further processing. Tissue collection took <2 minutes per fish, rapid

enough to minimize changes in gene expression responses to handling and dissection.

We extracted total RNA from brain tissue using the Qiagen RNeasy Lipid Tissue Mini Kit

(Qiagen, Germany) following manufacturer guidelines. We prepared separate sequencing

libraries for each individual using the NEBNext Ultra RNA Library Prep Kit for Illumina (New

England Biolabs, Massachusetts, USA) following manufacturer instructions. Libraries were

sequenced on an Illumina HiSeq 2000 at the Florida State University College of Medicine

Translational Science Laboratory (Tallahassee, Florida) in May 2014 (Aripo dataset) and January

2016 (Quare dataset). For the Aripo dataset, 40 samples (N=10 per group) were combined with

unique barcodes into eight samples per pool and each pool was sequenced on a single lane. For

the Quare dataset, 60 samples (N=12-16 per group) were combined into three pools with 20

samples per pool and each pool was sequenced in two separate lanes. Blocks of individuals run

in the same week were processed and sequenced together, and experimental groups were

balanced across sequencing lanes.

Transcript abundance estimation

To quantify gene expression, we mapped RNA sequencing reads to the guppy genome.

We received 1.45 billion 100-bp paired-end reads that passed the HiSeq quality filter, averaging

14.5 million reads per sample. We used Rcorrector to amend Illumina sequencing errors (Song &

Florea, 2015), and removed adapter sequences and trimmed reads for high-quality sequence

using Trim Galore! (Babraham Bioinformatics, Babraham Institute). Following developer

recommendations, we used a quality score of 33, a stringency of 5, and a minimum read length

of 36bp. We aligned corrected, trimmed reads from both datasets to the guppy genome

(http://uswest.ensembl.org/Poecilia_reticulata; downloaded November 2019) and estimated

transcript abundance using STAR (Dobin et al., 2013) with default parameters. On average, 91%

(range: 88.8% – 94.2%) of sequences per individual mapped to the guppy genome assembly.

Data filtering and screening

Prior to analysis, we performed data filtering and quality screening of gene expression

data. Preliminary cluster analyses revealed retinal contamination in a subset of our Aripo dataset

brain samples. While opsins are expressed at low levels in the brain, the very high expression

levels (>10,000 copies) in three samples pointed to retinal contamination. To deal with this issue,

we devised a sample filtering and screening procedure to remove genes in which expression

differences between samples were likely dominated by retinal contamination. Briefly, we first

filtered genes with low expression, then we used contigs annotated as known retinal genes

(Rhodopsin, red/green-sensitive opsins, blue-sensitive opsins) as seed contigs to identify other

contamination-related transcripts based on high positive correlations of expression levels with

seed genes. We calculated the gene-wise sum of correlations between candidate genes and seed

genes and performed multiple hypothesis testing using a false discovery rate (FDR) controlling

procedure. The nominal level of FDR was set to $\alpha=0.2$ to remove presumptive contaminant

contigs. Using this approach, we identified 1,559 contigs as presumptive retina-enriched genes

(~ 3% of all contigs in our final assembly) which we removed from both datasets in all subsequent

analyses (Table S1). More detailed descriptions of statistical procedures used for filtering are in

the Supplemental Methods.

Differential expression analysis

To test our specific hypotheses, we first needed to identify transcripts differentially

expressed between experimental groups, i.e. fish with differences in evolutionary history with

predators and/or developmental experience with predators (Figure 1). Due to differences in the

timing of fish rearing, sample processing, and sequencing, we did not combine Aripo and Quare

datasets for statistical analyses. We instead performed analysis in an identical fashion for both

drainages and subsequently conducted separate analyses to test hypotheses concerning

mechanistic parallelism across drainages (see below). Standard differential expression analysis

packages could not accommodate the random effects in our experimental design (i.e. family and

week; see below) but including these effects improved model fit and reflected our split-brood

experimental design, in which siblings were split into different rearing environments at birth, and

tissue was collected from fish in balanced groups of four in different weeks (see above). Fitting a

generalized linear mixed model with negative binomial link to each data set, revealed that a subset

of genes had statistically significant random effects due to either family, week, or both. Model

comparisons between the generalized linear model without random effects and the generalized

linear mixed model including week and family effects supported incorporating random effects, as

likelihood ratio tests indicated better fit of the mixed models for 391 out of 19,004 genes in the

Aripo dataset and 1,194 out of 19,902 genes in the Quare dataset. For consistency, we therefore

performed differential expression analysis using generalized linear mixed models with random

effects for all genes. Additional details of linear mixed model procedures are in the Supplemental

Methods.

We normalized read counts using DESeq2 in R (Love, Huber, & Anders, 2014) and

performed differential expression analysis using the lme4 package in R (github.com/lme4).

Normalized count data were modeled using a generalized linear mixed model with negative

binomial distribution as described above. We included population of origin, rearing environment,

and their interaction as fixed effects. In addition, we included family (siblings split across rearing

environments) and week (tissue was collected from animals in balanced blocks across multiple

weeks) as random effects. A Wald's test provided p-values for main effects and interaction effects

for each gene (Lehmann & Romano, 2005). We adjusted p-values for multiple hypothesis testing

using a direct approach for FDR correction (Storey, 2002) as implemented in the fdrtool package

in R (Strimmer, 2008). We considered transcripts differentially expressed (DE) if the adjusted p-

value for an effect was below 0.05.

To examine whether differential expression calls were influenced by transcript abundance,

we compared mean and median counts of DE and non-DE genes using two sample t-tests and

Wilcoxon rank sum tests. We also compared overall gene-wise variance between DE and non-

DE groups with respect to main effects of population of origin or rearing environment using Siegel-

Tukey tests and Kolmogorov-Smirnov tests. As sample sizes differed between our two datasets,

we also performed a down-sampling simulation to test for the effects of sample size on DE gene

number (Supplemental Methods).

We performed GO term enrichment analysis for all sets of DE transcripts using 'Biological

Processes' in the topGO package in R (Alexa & Rahnenfuhrer, 2016) and GO annotation

information from the guppy genome (http://uswest.ensembl.org/Poecilia_reticulata; downloaded

November 2019).

Statistical tests of specific hypotheses

To test Hypothesis 1, we evaluated overlap in transcripts differentially expressed between

populations (i.e. population DE genes) and between rearing conditions (i.e. rearing DE genes).

To test the hypothesis that initial plastic responses following colonization of low-predation

environments might drive subsequent divergence, we specifically compared genes with

significant ancestral plasticity (i.e. differences between high-predation siblings reared with and

without predators; see Glossary and Figure 1) to genes with significant genetic expression

divergence (high- versus low-predation fish reared in the derived, predator-free environment)

within each drainage. We used FDR-corrected post-hoc tests of simple effects to identify DE

genes in these comparisons and estimate log-fold gene expression changes.

To test Hypothesis 2, we compared the association between directions of plasticity and

divergence (concordant vs. non-concordant) in (1) those genes with *both* significant ancestral

plasticity and significant population differences, and (2) *all* transcripts with significant population

differences in expression, including those without significant expression plasticity. The second

comparison addressed the possibility that even subthreshold expression plasticity in response to

rearing environment could influence patterns of expression divergence, especially given the high

false negative rate in our conservative DE analyses. We used log-fold expression changes to

determine whether gene expression differences were in the same/concordant direction (i.e.

upregulated in high-predation populations and in response to rearing with predator cues, or down-

regulated in high-predation populations and in response to rearing with predator cues) or

opposite/non-concordant direction (upregulated in high-predation populations but down-regulated

in response to rearing with predators, or vice versa).

Statistical approaches to test Hypotheses 1 and 2 that compare gene-wise contrasts with

respect to a common experimental group (high predation fish reared without exposure to

predators) are biased and nontrivial to correct, (for illustration of this point see a classical example

in the Supplementary Methods and the discussion by Mallard, Jakšić, & Schlötterer, 2018). We

therefore instead evaluated whether significant effects of rearing condition in the ancestral

population tended to be associated with significant population effects across all genes rather than

evaluating individual genes. To do so, we focused on (a) the probability of significant population

simple effects conditional on significant rearing effects in the high-predation population; and (b)

the probability of significant population simple effects conditional on non-significant rearing effects

(Cao, Zhou, Breidt, & Peers, 2020; Hwang & Liu, 2010; Storey, 2007). We tested whether the

conditional probabilities of gene expression divergence between high- and low-predation

populations differed for genes that exhibited ancestral plasticity and those that did not (Hypothesis

1). We further tested whether conditional probabilities for upregulated or downregulated genes in

the low-predation population compared to the high-predation population differed depending on

the direction of plasticity (Hypothesis 2). To draw inference on these conditional probabilities

across all genes, we adapted a parametric bootstrap method (Efron & Tibshirani, 1993) similar to

the approach implemented by (W. C. Ho & Zhang, 2019). As the number of genes is large, the

estimated conditional probabilities are accurate and have statistical properties that make the use

of parametric bootstrap valid. We calculated parametric bootstrap confidence intervals for these

differences in conditional probabilities as detailed in the Supplemental Methods, with confidence

intervals that do not include zero indicating a significant association between gene expression

divergence and plasticity.

To test Hypothesis 3, we examined the subset of transcripts with significant interaction

effects in our differential expression analysis (see above). We grouped genes with significant

interaction effects into categories for the evolution of plasticity outlined by Renn and Schumer

(2013) based on *post hoc* differences between rearing environments within a population (p<0.05):

(1) Assimilated: plasticity in the ancestral high-predation population but a loss of plasticity in the

derived low-predation population; (2) Accommodated: a change in the degree, but not the

direction, of plasticity in the derived as compared to the ancestral population; (3) Reversed:

opposing directions of plasticity in high- versus low-predation populations; (4) Evolved plastic: no

plasticity in the ancestral high-predation population but an emergence of plasticity in the derived

low-predation population; and (5) Unclassified: genes that had a significant interaction effect but

no significant *post hoc* rearing differences and therefore could not be unambiguously classified

into one of the above categories. All statistical tests and data visualization were performed in R

(version 3.5.1; The R Foundation for Statistical Computing).

To test Hypothesis 4, we evaluated overlap in DE transcript identity and expression

direction between the Aripo and Quare lineages. We used chi-square tests of independence to

test for greater than chance overlap *between* drainages in those gene sets differentially expressed

based on rearing environment or population of origin *within* drainages. We also used chi-square

tests to test for biases in concordant versus non-concordant expression between drainages. We

performed two additional comparisons to address potential bias in our conclusions due to high

false negative rates in gene expression data (Rice, Schork, & Rao, 2008). First, we performed

the same analysis as before at two less stringent p-value cut-offs (p=0.1 and 0.2). Second, for

comparisons of expression direction, we made an additional, more conservative comparison by

including not only those genes differentially expressed in *both* drainages (i.e. the intersection of

DE gene lists), but also those genes differentially expressed in *either* drainage (i.e. the union of

DE gene lists) in the analysis. We again used log-fold expression differences to call gene

expression as being in the same/concordant or opposite/non-concordant direction between

drainages.

Results

Differential gene expression and GO term enrichment

In order to test our specific hypotheses, we first summarized differential expression

patterns in both datasets. In the Aripo drainage, 766 genes were differentially expressed (DE)

between high- and low-predation populations, 1,118 genes were differentially expressed between

pred- and pred+ fish, and 825 genes had interaction effects (Fig. 2; Table S2). DE genes were

enriched for GO categories including DNA integration, immune function, and morphogenesis. In

the Quare drainage, 3,831 genes were differentially expressed between high- and low-predation

populations, 791 genes were differentially expressed between pred- and pred+ fish, and 586

genes had interaction effects (Fig. 2; Table S3). DE genes were enriched for GO categories

including immune function, developmental processes, and neurogenesis. Complete results of GO

enrichment analyses for population of origin, rearing, and interaction effects are in the

Supplemental Information (Tables S4 & S5).

Given the marked difference in the total number of DE genes between datasets, we looked

for evidence that this pattern might arise from technical variation, expression bias, or differences

in sample size between the two datasets. While we cannot rule out some influence of technical

variation, we found no evidence for differences in sequence quality, read alignment statistics, or

variance in transcript abundance to suggest that variation in the number of DE transcripts between

drainages was of technical origin. Nor did we find evidence that identification of DE transcripts

was biased by transcript expression level: mean and median gene expression were significantly

lower in DE (population, rearing, and/or interaction effects) as compared to non-DE transcripts in

both the Aripo (Fig. S1; mean: DE=263, non-DE=344, $p<0.0001$; median: DE=65, non-DE=107,

$p<0.0001$), and Quare (Fig. S1; mean: DE=264, non-DE=334, $p<0.0001$; median: DE=86, non-

DE=101, $p<0.0001$) datasets. A down-sampling simulation provided no evidence for a reliance of

the greater number of population DE genes in the Quare dataset on the larger sample size of this

dataset (Figure S2). In sum, these tests suggest that identification of DE transcripts was not

influenced by relatively greater expression magnitude or variance of these transcripts, nor by

sample size differences.

Expression plasticity biases the propensity for genetic expression divergence

Evolutionary theory and empirical studies suggest that environmentally induced plasticity

may facilitate adaptation to novel environments. To test Hypothesis 1, we asked whether genes

with significant ancestral expression plasticity were more likely to show genetic divergence in

expression. We examined the extent of overlap in gene sets (1) differentially expressed between

high-predation fish reared with and without predators, to those (2) differentially expressed

between populations adapted to high- versus low-predation environments. In both drainages, the

DE genes with significant ancestral plasticity were more likely to show significant population

divergence than expected by chance (Aripo: 264 transcripts, 95% CI for differences in conditional

probability of significant divergence for plastic genes compared to conditional probability of

divergence for non-plastic genes: [0.125, 0.150]; Quare: 409 transcripts, 95% CI: [0.211,0.239]),

supporting an association between plastic responses and genetic divergence.

Expression plasticity biases the direction of genetic expression divergence

To test Hypothesis 2, we asked whether the directions of expression plasticity and genetic

expression divergence were associated. We did so for those genes with both significant ancestral

plasticity and population differences, as well as the larger set of all transcripts with significant

population differences (i.e. including those with and without significant rearing differences). We

performed the second, more conservative comparison because we reasoned that even

subthreshold expression plasticity could influence the direction of adaptive divergence. Within

both drainages, we found a signature of non-adaptive plasticity associated with genetic

expression divergence. Approximately 84% of DE genes exhibited population and rearing

expression changes in non-concordant directions in the Aripo drainage (95% CI for difference in

probability of same versus opposite expression direction: [-0.564,-0.499]), and 68% of DE genes

exhibiting non-concordant expression in the Quare drainage (95% CI: [-0.520,-0.456]) (Fig. 3).

Patterns were consistent in both drainages when we considered the larger set of all genes with

significant population expression differences (Aripo: 80% of genes with non-concordant

expression, 95% CI: [-0.811,-0.752]; Quare: 66% of genes with non-concordant expression, 95%

CI: [-0.803,-0.733]).

Evolution of expression plasticity

We next explored the evolution of expression plasticity itself to test Hypothesis 3. We

identified genes with evolved expression plasticity as those in which the direction and/or degree

of rearing effects depended on population of origin, i.e. those genes with significant population-

by-rearing interaction effects. The number of genes with significant expression plasticity that

evolved (i.e. significant rearing and interaction effects) was approximately a quarter to a third of

the number of genes with significant expression plasticity that did not evolve (i.e. significant

rearing effect but no interaction; 263 (24%) in Aripo and 286 (36%) in Quare) (Fig. 2).

To further characterize how expression plasticity evolved, we subdivided genes with

significant population-by-rearing interaction effects into one of five categories: assimilated,

accommodated, reversed, evolved plastic, or unclassified (Fig. 4A; based on Renn and Schumer,

2013). We found many transcripts that exhibited evolution of plasticity, with all five categories

represented in both datasets (Fig. 4B). Of the genes with significant interaction effects, in the

Aripo drainage 263 (32%) transcripts showed patterns of expression assimilation, 81 (10%)

transcripts showed patterns of expression accommodation, 113 (14%) transcripts exhibited

reversed plasticity, and 182 (22%) transcripts evolved plasticity in the derived population. In the

Quare drainage, 164 (28%) transcripts showed patterns of expression assimilation, 110 (19%)

transcripts showed patterns of expression accommodation, 128 (22%) transcripts exhibited

reversed plasticity, and 95 (16%) transcripts evolved plasticity in the derived population (Fig. 4B).

Parallelism in gene expression patterns across drainages

To test Hypothesis 4 that shared mechanisms mediate parallel phenotypic adaptation

across drainages, we asked whether differentially expressed genes *within* drainages were

overlapping in identity and expression direction *between* drainages. Of the genes that diverged

between high- and low-predation populations within a drainage (i.e., population main effect), 174

were overlapping between drainages (Table S6), more than expected by chance (χ^2=8.02,

p=0.0046). However, the direction of expression divergence among these overlapping genes was

not associated across the two drainages (χ^2=0.16, p=0.686): 91 genes (52%) had concordant

expression changes and 83 genes (48%) had non-concordant expression changes between

drainages (Fig. 5). To control for the influence of false negatives on our conclusions, we repeated

this comparison at less stringent significance adjusted p-value cut-offs. At a p<0.1, we found

marginally significant overlap in gene identify (347 overlapping genes; χ^2=3.38, p=0.06), but no

association in expression direction (χ^2=0.65, p=0.42), and at p<0.2 we found no significant overlap

or association in expression direction. These patterns remained unchanged when we restricted

the analysis to those genes with strongest evidence for adaptive divergence (see Methods and

Table S8).

Of the genes that showed significant expression plasticity (i.e. main effect of rearing), 64

transcripts were overlapping between drainages (Table S7), more than expected by chance

(χ^2=14.06, p=0.0001). Of these 43 (67%) had plastic changes in the same direction (i.e.

concordant) and 21 (33%) had expression changes in opposite directions (i.e. non-concordant)

between drainages, a stronger signature of concordant expression between river drainages than

expected by chance (χ^2=7.56, p=0.006) (Fig. 5). At less stringent p-value cut-offs for differential

expression we found no more overlap in gene identity than expected by chance (p<0.1: 108

overlapping genes; χ^2=0.05, p=0.81; p<0.2: 276 overlapping genes; χ^2=1.36, p=0.24).

Discussion

We examined transcriptomic differences associated with adaptive divergence and rearing

environment across repeated, independent evolutionary lineages in guppies. Within lineages, we

observed phenotypic plasticity in gene expression patterns as well as the evolution of gene

expression plasticity itself. Plastic genes were more likely to exhibit population differences in

expression, and these population differences were more likely to be in the opposite, rather than

the same, direction as ancestral expression plasticity. Comparing across lineages, DE gene sets

for population and rearing within each lineage were largely non-overlapping (>75% non-shared

identity) between lineages. Nonetheless, the small number of DE genes that did overlap between

drainages was more than expected by chance. Taken together, our findings suggest that a small

number of core genes may be repeatedly targeted during colonization of low-predation

environments, but that largely non-shared, alternative transcriptional solutions are associated with

parallel phenotypic adaptation across lineages. Combining studies of phenotypic plasticity and

parallel adaptation, we highlight a role for transcriptional flexibility in the parallel phenotypic

evolution of a species known for rapid adaptation.

Impacts of gene expression plasticity on expression evolution

A growing number of studies are using transcriptomic and proteomic surveys to address

the long-standing debate on the role of plasticity in evolution, with contrasting results favoring the

alternative hypotheses that adaptive (Fraser et al., 2014; Gleason & Burton, 2015; Li et al., 2018;

Mäkinen et al., 2016; Scoville & Pfrender, 2010; Shaw et al., 2014; Wang & Althoff, 2019) versus

non-adaptive (Dayan et al., 2015; Ghalambor et al., 2015; W. Ho & Zhang, 2018; Pespeni et al.,

2013; Schaum et al., 2013) plasticity facilitates adaptation. We recently proposed that non-

adaptive plasticity dominates during the earliest stages of rapid evolution, and that adaptive

plasticity may contribute to subsequent fine-tuning of phenotypes (Fischer et al., 2016a). In line

with this prediction and our previous findings, we report here that the strong signature of non-

adaptive plasticity in brain gene expression (89% of transcripts) observed in guppy populations in

the earliest stages of adaptation to low-predation environments (Ghalambor et al., 2015) is

present but weaker in long-established low-predation populations. Furthermore, in the current

study, this signature of non-adaptive plasticity was more apparent in the more recently diverged

Aripo lineage (80% of genes) as compared to the older Quare lineage (65% of genes). Our

findings highlight the need to explicitly consider how plasticity relates to divergence throughout

successive stages of adaptation in order to build a more holistic understanding of the role of

plasticity in adaptation.

Evolution of plasticity in gene expression

Diverse transcriptional patterns could accompany evolved differences in behavioral

plasticity (Renn & Schumer, 2013), but relevant data characterizing the evolution of gene

expression plasticity has been lacking. We found that plasticity itself evolved in approximately one

third of genes with significant ancestral expression plasticity in both lineages. As was observed

for transcriptomic evolution of gill tissue in stickleback fish (Gibbons, Metzger, Healy, & Schulte,

2017), plasticity evolution showed no consistent pattern, with genes gaining, losing, and switching

the direction of expression plasticity between ancestral and derived populations in both lineages.

The lack of consistency in these patterns and the dearth of studies that have characterized the

evolution of gene expression plasticity make these patterns difficult to interpret at present. Some

of the diversity is likely associated with adaptive phenotypic divergence between high- and low-

predation populations, in which initially plastic behavioral shifts may become fixed, eliminated, or

altered over time. For example, evolution in gene expression plasticity could reflect the gains,

losses, and switches behavioral plasticity in these populations (Fischer et al., 2016b).

Alternatively, compensatory and homeostatic mechanisms could promote diversity among plastic

responses in gene expression without altering higher-level phenotypic traits such as morphology

and behavior (Badyaev, 2018; Fischer, Ghalambor, & Hoke, 2016a; Renn & Schumer, 2013).

Finally, because fish in low-predation habitats experience relaxed selection on predator-induced

plasticity in conjunction with low effective populations sizes, some of the evolution of

transcriptional plasticity we report likely arose as a product of genetic drift, rather than selection

for altered plastic responses (Lynch, 2007). Documenting the evolution of expression plasticity is

an important first step, and additional studies are needed to understand the ubiquity and

evolutionary sources of these patterns.

Transcriptomic signatures of population divergence in two independent evolutionary lineages

We identified many genes differentially expressed between high- and low-predation

populations in each river drainage. The absolute number of population DE transcripts was smaller

in the Aripo drainage as compared to the Quare drainage, but the number of developmental and

interaction differences was greater in the Aripo drainage. Our analyses provided no evidence that

technical differences in expression magnitude, expression variance, or sample size influenced

these differences between datasets in the number of genes reaching statistical significance. We

therefore propose that differences in DE gene number between drainages are biological in origin,

as high- and low-predation populations in the Quare drainage show greater genetic divergence

than those in the Aripo drainage (Willing et al., 2010). Indeed, we note that the larger number of

genes in the Quare dataset come from a large difference in the number of genes with population,

but not rearing or interaction, effects. In sum, we suggest that the difference in the number of

differentially expressed transcripts between datasets is primarily driven by the greater degree of

genetic divergence between high- and low-predation populations within the Quare as compared

to the Aripo river drainage (Willing et al., 2010).

When we compared DE gene sets between drainages, we found a small, but significant,

number of population and rearing DE genes shared across lineages. However, the majority of DE

genes were non-shared across drainages for both population (Aripo: 78%; Quare: 96%), and

rearing (Aripo: 94%; Quare: 92%) effects, and we note that this pattern persisted at less stringent

p-values cutoffs. Rearing DE genes showed a significant association in expression direction

across lineages, with 67% of genes concordantly differentially expressed between lineages. In

contrast, genes associated with genetic divergence between high- and low-predation populations

in both drainages showed no signature of expression concordance across lineages. We suggest

that these patterns across drainages point to a small number of core genes that exhibit

predictable, plastic expression responses upon colonization of low-predation environments, but

that lineage-specific selection pressures, differences in genetic background, non-adaptive

processes (e.g. drift, inbreeding, founder effects), and alternative gene expression responses give

rise to largely non-overlapping, non-concordant expression differences associated with parallel

phenotypic adaptation.

A previous study in guppies performed a similar comparison of gene expression changes

associated with adaptation to low-predation environments (Ghalambor et al., 2015) and found a

strong signal of concordant differential expression in genes differentially expressed based on

population of origin. Whereas the present study compared long-term natural population

divergence across drainages, Ghalambor et al. (2015) characterized the early stages of

adaptation of experimentally introduced low-predations populations derived from founders from a

single high-predation source population within the same drainage. These contrasting findings in

comparisons of population pairs within the same drainage versus across genetically distinct

drainages highlight the impacts of standing genetic variation within the source population on

mechanisms of divergence (Feiner, Rago, While, & Uller, 2017; Thompson, Osmond, & Schluter,

2019), particularly at early stages of evolution (Barrett & Schluter, 2008): while alternative

transcriptional 'solutions' are possible, shared genetic background appears to bias evolutionary

outcomes toward shared patterns. These findings are exciting as they extend to a mechanistic

level the growing recognition of and interest in the non-parallel aspects of parallel adaptation,

even in systems considered classical examples of parallel/convergent evolution (Bolnick et al.,

2018; Manousaki et al., 2013; Paccard et al., 2020; Stuart et al., 2017).

Both adaptive and non-adaptive processes may contribute to the combination of shared

and distinct transcriptional mechanisms we find associated with parallel, adaptive life-history,

morphological, and behavioral phenotypes across lineages in guppies. First, as described above,

differences in standing genetic variation likely influence which mechanisms are available to

selection in response to common environmental conditions in different drainages (Barrett &

Schluter, 2008; Thompson et al., 2019). Second, low-predation populations are typically

established by a small number of individuals (Barson et al., 2009; Fraser et al., 2015; Willing et

al., 2010), making them susceptible to the unpredictable, non-adaptive influences of founder

effects, genetic drift, and/or inbreeding on gene expression divergence; although we note that

work in three-spine stickleback suggests deviations from parallel evolution tend to result from

selection rather than non-adaptive processes (Kaeuffer et al., 2011). Third, the large number of

significantly evolved genes that did not overlap between drainages may also represent adaptive

responses to drainage- or site-specific environmental factors other than predation (Fitzpatrick,

Torres-Dowdall, Reznick, Ghalambor, & Chris Funk, 2014; Zandonà et al., 2011). Finally,

compensatory or homeostatic gene expression responses may arise in response to any of the

above factors, leading to alternative transcriptional configurations associated with similar higher-

level phenotypes.

In sum, we suggest that genetic similarity among ancestral populations may channel low-

predation populations within the same drainage toward shared transcriptional solutions (as in

Ghalambor et al. 2015), while differences in standing genetic variation, drainage-specific

environmental conditions, founder effects, and alternative compensatory changes can result in

distinct mechanistic paths to arrive at shared organism-level phenotypes. We cannot definitely

distinguish causal from non-adaptive and compensatory gene expression differences under these

scenarios – indeed, it is likely a combination of these factors that contribute to distinct

transcriptional patterns associated with parallel adaptation. Nonetheless, alternative

transcriptional patterns suggest that mechanistic flexibility and 'many-to-one' mapping of gene

expression to organism level phenotypes may facilitate adaptation.

Conclusions

We assessed the extent to which adaptation to common environments targets predictable

changes in transcript abundance across independent evolutionary events. Within lineages, genes

with greater expression plasticity were more likely to diverge in expression between populations,

but in the direction opposite of developmental changes. Despite a small number of differentially

expressed genes shared across drainages, parallel adaptation to low-predation environments in

independent lineages was associated with divergence in largely non-overlapping transcripts.

While identification of shared genes is generally used as the starting point for work exploring

mechanisms of parallel adaptation – and our work here indeed suggests that a small number of

core genes may indeed be critical during the earliest stages of adaptation – we propose that

parallel evolutionary transitions are not limited to a small set of possible transcriptional

mechanisms in guppies. Instead, our results highlight the potential for alternative transcriptional

solutions associated with parallel, adaptive trait evolution even within a single species.

Transcriptional network versatility, in which diverse alterative network configurations can produce

common network outputs and behavioral phenotypes, may allow underlying networks to

simultaneously accommodate the influences of selection, drift, and genetic background and

thereby facilitate evolution in a species known for rapid adaptation to novel environments.

Acknowledgements: We thank the members of the Colorado State University Guppy Group for

fish care, especially Sarah E Westrick and Kimberly E Dolphin for help with tissue collection. We

thank Laura R Stein for assistance with tissue processing and Cameron K Ghalambor for fruitful

discussions and comments on earlier versions of the manuscript. All high-powered computing

was performed on the Odyssey computing cluster supported by the FAS Science Division

Research Computing Group at Harvard University. This work was supported by the National

Science Foundation DDIG-1311680 (to EKF), RCN IOS-1256839 (to EKF), IOS-1354755 (to

KLH), IOS-1354775 (to KAH), IOS-0934451 (to KAH), DEB-0846175 (to CKG), IIS-1545994 (to

WZ), and US Department of Energy DE-SC0018344 (to WZ).

Author contributions: EKF and KLH conceived of the study; EKF collected samples and

performed molecular work; EKF and KAH performed gene expression mapping, transcript

abundance estimation, and preliminary differential expression analyses; YS and WZ devised

and performed statistical analyses with input from EKF and KLH; EKF and KLH wrote the

manuscript with contributions from all authors. All authors gave final approval for publication.

Data Accessibility: Raw sequencing reads are available through the NCBI SRA repository

(PRJNA601479). R code for statistical analyses is available on GitHub.

References
Abouheif, E., & Wray, G. (2002). Evolution of the gene network underlying wing polymorphism
in ants. *Science*, *297*(2002), 249–252. doi: 10.1126/science.1071468
Alberch, P. (1991). From genes to phenotype: dynamical systems and evolvability. *Genetica*,
84, 5–11. doi: 10.1007/BF00123979
Alexa, A., & Rahnenfuhrer, J. (2016). *topGO: Enrichment Analysis for Gene Ontoloty*. R
package version 2.30.1.
Arendt, J., & Reznick, D. (2008). Convergence and parallelism reconsidered: what have we
learned about the genetics of adaptation? *Trends in Ecology and Evolution*, *23*(1), 26–32.
doi: 10.1016/j.tree.2007.09.011
Badyaev, A. V. (2018). Evoulutionary transitions in controls reconcile adaptation with continuity
of evolution. *Seminars in Cell and Developmental Biology*, *88*, 36–45. doi:
10.1016/j.semcdb.2018.05.014
Badyaev, A. V, & Morrison, E. S. (2018). Emergent buffering balances evolvability and
robustness in the evolution of phenotypic flexibility. *Evolution*, *72*(3), 647–662. doi:
10.1111/evo.13441
Baldwin, J. M. (1896). A New Factor in Evolution. *The American Naturalist*, *30*, 441–451; 536–
553.
Barrett, R. D. H., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in
Ecology and Evolution*, *23*(1), 38–44. doi: 10.1016/j.tree.2007.09.008
Barson, N. J., Cable, J., & Van Oosterhout, C. (2009). Population genetic analysis of
microsatellite variation of guppies (Poecilia reticulata) in Trinidad and Tobago: Evidence for
a dynamic source-sink metapopulation structure, founder events and population
bottlenecks. *Journal of Evolutionary Biology*, *22*(3), 485–497. doi: 10.1111/j.1420-
9101.2008.01675.x
Bolnick, D. I., Barrett, R. D. H., Oke, K. B., Rennison, D. J., & Stuart, Y. E. (2018). (Non)Parallel
evolution. *Annual Review of Ecology, Evolution, and Systematics*. doi: 10.1146/annurev-
ecolsys-110617-062240
Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An Expanded View of Complex Traits: From
Polygenic to Omnigenic. *Cell*. doi: 10.1016/j.cell.2017.05.038
Cao, M., Zhou, W., Breidt, F. J., & Peers, G. (2020). Large scale maximum average power
multiple inference on time-course count data with application to RNA-seq analysis.
Biometrics. 76(1). 9-22. doi: 10.1111/biom.13144
Conover, D. O., Duffy, T. A., & Hice, L. A. (2009). The covariance between genetic and
environmental influences across ecological gradients: Reassessing the evolutionary
significance of countergradient and cogradient variation. *Annals of the New York Academy*
of Sciences, *1168*, 100–129. doi: 10.1111/j.1749-6632.2009.04575.x
Crawford, D. L., & Oleksiak, M. F. (2007). The biological importance of measuring individual
variation. *Journal of Experimental Biology*, *210*(9), 1613–1621. doi: 10.1242/jeb.005454
Dayan, D. I., Crawford, D. L., & Oleksiak, M. F. (2015). Phenotypic plasticity in gene expression
contributes to divergence of locally adapted populations of Fundulus heteroclitus.
Molecular Ecology, *24*(13), 3345–3359. doi: 10.1111/mec.13188
Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R.
(2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21. doi:
10.1093/bioinformatics/bts635
Draghi, J. A., & Whitlock, M. C. (2012). Phenotypic plasticity facilitates mutational variance,
genetic variance, and evolvability along the major axis of environmental variation.
Evolution, *66*(9), 2891–2902. doi: 10.1111/j.1558-5646.2012.01649.x
Drion, G., O'Leary, T., & Marder, E. (2015). Ion channel degeneracy enables robust and tunable
neuronal Firing rates. *Proceedings of the National Academy of Sciences of the United*
States of America. doi: 10.1073/pnas.1516400112

Efron, B., & Tibshirani, R. J. (1993). An introduction to the bootstrap. Monographs on Statistics
and Applied Probability, No. 57. Chapman and Hall, London, 436 p. In *Monographs on*
Statistics and Applied Probability.
Elvidge, C. K., Ramnarine, I., & Brown, G. E. (2014). Compensatory foraging in Trinidadian
guppies: Effects of acute and chronic predation threats. *Current Zoology*, *60*(3), 323–332.
doi: 10.1093/czoolo/60.3.323
Endler, J. A. (1995). Multiple-trait coevolution and environmental gradients in guppies. *Trends in*
Ecology & Evolution, *10*(1), 22–29. doi: 10.1016/S0169-5347(00)88956-9
Espinosa-Soto, C., Martin, O. C., & Wagner, A. (2011). Phenotypic plasticity can facilitate
adaptive evolution in gene regulatory circuits. *BMC Evolutionary Biology*, *11*(5), 1–14. doi:
10.1186/1471-2148-11-5
Fajen, A. V. A., & Breden, F. (1992). Mitochondrial DNA sequence variation among natural
populations of the Trinidad guppy, Poecilia reticulata. *Evolution*, *46*(5), 1457–1465.
Feiner, N., Rago, A., While, G. M., & Uller, T. (2017). Signatures of selection in embryonic
transcriptomes of lizards adapting in parallel to cool climate. *Evolution*, *72*(1), 67–81. doi:
10.1111/evo.13397
Fischer, E. K., Ghalambor, C. K., & Hoke, K. L. (2016a). Can a Network Approach Resolve How
Adaptive vs Nonadaptive Plasticity Impacts Evolutionary Trajectories? *Integrative and*
Comparative Biology, *56*(5), 877–888. doi: 10.1093/icb/icw087
Fischer, E. K., Ghalambor, C. K., & Hoke, K. L. (2016b). Plasticity and evolution in correlated
suites of traits. *Journal of Evolutionary Biology*, *29*(5), 991–1002. doi: 10.1111/jeb.12839
Fischer, E. K., Harris, R. M., Hofmann, H. A., & Hoke, K. L. (2014). Predator exposure alters
stress physiology in guppies across timescales. *Hormones and Behavior*, *65*(2), 165–172.
doi: 10.1016/j.yhbeh.2013.12.010
Fischer, E. K., Soares, D., Archer, K. R., Ghalambor, C. K., & Hoke, K. L. (2013). Genetically
and environmentally mediated divergence in lateral line morphology in the Trinidadian
guppy (Poecilia reticulata). *Journal of Experimental Biology*, *216*(16), 3132–3142. doi:
10.1242/jeb.081349
Fitzpatrick, S. W., Torres-Dowdall, J., Reznick, D. N., Ghalambor, C. K., & Chris Funk, W.
(2014). Parallelism Isn't Perfect: Could Disease and Flooding Drive a Life-History Anomaly
in Trinidadian Guppies? *The American Naturalist*, *183*(2), 290–300. doi: 10.1086/674611
Fraser, B. A., Janowitz, I., Thairu, M., Travis, J., Hughes, K. A., Fraser, B. A., … Hughes, K. A.
(2014). Phenotypic and genomic plasticity of alternative male reproductive tactics in sailfin
mollies. *Proceedings of the Royal Society B: Biological Sciences*, *281*, 20132310.
Fraser, B. A., Künstner, A., Reznick, D. N., Dreyer, C., & Weigel, D. (2015). Population
genomics of natural and experimental populations of guppies (Poecilia reticulata).
Molecular Ecology, *24*(2), 389–408. doi: 10.1111/mec.13022
Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K. A.
(2015). Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
nature. *Nature*, *525*(7569), 372–375. doi: 10.1038/nature15256
Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-
adaptive phenotypic plasticity and the potential for contemporary adaptation in new
environments. *Functional Ecology*, *21*(3), 394–407. doi: 10.1111/j.1365-2435.2007.01283.x
Ghalambor, C. K., Reznick, D. N., & Walker, J. A. (2004). Constraints on adaptive evolution:
The functional trade-off between reproduction and fast-start swimming performance in the
Trinidadian guppy (Poecilia reticulata). *American Naturalist*, *164*(1), 38–50. doi:
10.1086/421412
Gibbons, T. C., Metzger, D. C., Healy, T. M., & Schulte, P. M. (2017). Gene expression plasticity
in response to salinity acclimation in threespine stickleback ecotypes from different salinity
habitats. *Molecular Ecology*, *26*, 2711–2725. doi: 10.1111/mec.14065
Gilliam, J. F., Fraser, D. F., & Alkins-Koo, M. (1993). Structure of a Tropical Stream Fish

Community : A Role for Biotic Interactions. *Ecology*, *74*(6), 1856–1870.

Gleason, L. U., & Burton, R. S. (2015). RNA-seq reveals regional differences in transcriptome
response to heat stress in the marine snail Chlorostoma funebralis. *Molecular Ecology*,
24(3), 610–627. doi: 10.1111/mec.13047

Grashow, R., Brookings, T., & Marder, E. (2009). Reliable neuromodulation from circuits with
variable underlying structure. *Proceedings of the National Academy of Sciences*, *106*(28),
11742–11746. doi: 10.1073/pnas.0905614106

Grether, G. F. (2005). Environmental Change, Phenotypic Plasticity, and Genetic
Compensation. *The American Naturalist*, *166*(4), E115–E123. doi: 10.1086/432023

Handelsman, C. A., Ruell, E. W., Torres-Dowdall, J., & Ghalambor, C. K. (2014). Phenotypic
Plasticity Changes Correlations of Traits Following Experimental Introductions of
Trinidadian Guppies (Poecilia reticulata). *Integrative and Comparative Biology*, *54*(5), 794–
804. doi: 10.1093/icb/icu112

Handelsman, C. A., Broder, E. D., Dalton, C. M., Ruell, E. W., Myrick, C. A., Reznick, D. N., &
Ghalambor, C. K. (2013). Predator-induced phenotypic plasticity in metabolism and rate of
growth: Rapid adaptation to a novel environment. *Integrative and Comparative Biology*,
53(6), 975–988. doi: 10.1093/icb/ict057

Harris, S., Ramnarine, I. W., Smith, H. G., & Pettersson, L. B. (2010). Picking personalities
apart: Estimating the influence of predation, sex and body size on boldness in the guppy
Poecilia reticulata. *Oikos*. doi: 10.1111/j.1600-0706.2010.18028.x

Haskins, C., Haskins, E., McLaughlin, J., & Hewitt, R. (1961). Polymorphism and population
structure in Lebistes reticulatus, an ecological study. In F. Blair (Ed.), *Vertebrate speciation*
(pp. 320–395). Austin: University of Austin Press.

Ho, W. C., & Zhang, J. (2019). Genetic Gene Expression Changes during Environmental
Adaptations Tend to Reverse Plastic Changes Even after the Correction for Statistical
Nonindependence. *Molecular Biology and Evolution*, *36*(3), 604–612. doi:
10.1093/molbev/msz002

Ho, W., & Zhang, J. (2018). Evolutionary adaptations to new environments generally reverse
plastic phenotypic changes. *Nature Communications*, *9*(350), 1–11. doi: 10.1038/s41467-
017-02724-5

Huizinga, M., Ghalambor, C. K., & Reznick, D. N. (2009). The genetic and environmental basis
of adaptive differences in shoaling behaviour among populations of Trinidadian guppies,
Poecilia reticulata. *Journal of Evolutionary Biology*, *22*(9), 1860–1866. doi: 10.1111/j.1420-
9101.2009.01799.x

Hwang, J. T. G., & Liu, P. (2010). Optimal tests shrinking both means and variances applicable
to microarray data analysis. *Statistical Applications in Genetics and Molecular Biology*. doi:
10.2202/1544-6115.1587

Insel, T. R., & Young, L. J. (2000). Neuropeptides and the evolution of social behavior. *Current
Opinion in Neurobiology*, *10*(6), 784–789. doi: 10.1016/S0959-4388(00)00146-X

Kaeuffer, R., Peichel, C. L., Bolnick, D. I., & Hendry, A. P. (2011). Parallel and nonparallel
aspects of ecological, phenotypic, and genetic divergence across replicate population pairs
of lake and stream stickleback. *Evolution*, *66*(2), 402–418. doi: 10.5061/dryad.k987h

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446. doi:
10.1111/j.1420-9101.2009.01754.x

Lehmann, E. L., & Romano, J. P. (2005). *Testing Statistical Hypotheses* (Third Edit). New York:
Springer.

Leinonen, T., Cano, J. M., Mäkinen, H., & Merilä, J. (2006). Contrasting patterns of body shape
and neutral genetic divergence in marine and lake populations of threespine sticklebacks.
Journal of Evolutionary Biology, *19*, 1803–1812. doi: 10.1111/j.1420-9101.2006.01182.x

Li, L., Li, A., Song, K., Meng, J., Guo, X., Li, S., … Xu, F. (2018). Divergence and plasticity

shape adaptive potential of the Pacific oyster. *Nature Ecology Evolution*, *2*, 1751–1760.
doi: 10.1038/s41559-018-0668-2
Losos, J. B. (2011). Convergence, adaptation, and constraint. *Evolution*, *65*(7), 1827–1840. doi:
10.1111/j.1558-5646.2011.01289.x
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21. doi:
10.1186/s13059-014-0550-8
Lynch, M. (2007). The frailty of adaptive hypotheses for the origins of organismal complexity.
Proceedings of the National Academy of Sciences, *104*, 8597–8604. doi: 10.17226/11790
Magurran, A. E. (2005). Evolutionary ecology: the Trinidadian guppy. In *Oxford Series in*
Ecology and Evolution (Vol. 19). doi: 10.1093/acprof
Magurran, A. E., & Seghers, B. H. (1990a). Population differences in predator recognition and
attack cone avoidance in the guppy Poecilia reticulata. *Animal Behaviour*, *40*(3), 443–452.
doi: 10.1016/S0003-3472(05)80524-X
Magurran, A. E., & Seghers, B. H. (1990b). Population Differences in the Schooling Behaviour
of Newborn Guppies, Poecilia reticulata. *Ethology*, *84*, 334–342. doi: 10.1111/j.1439-
0310.1990.tb00807.x
Magurran, A. E., & Seghers, B. H. (1991). Variation in Schooling and Aggression Amongst
Guppy (Poecilia Reticulata) Populations in Trinidad. *Behaviour*, *188*(3–4), 214–234. doi:
10.1163/156853991X00292
Magurran, A. E., & Seghers, B. H. (1994). Predator Inspection Behaviour Covaries with
Schooling Tendency Amongst Wild Guppy, Poecilia Reticulata, Populations in Trinidad.
Behaviour, *128*(1–2), 121–134. doi: 10.1163/156853994X00073
Mäkinen, H., Papakostas, S., Vøllestad, L. A., Leder, E. H., & Primmer, C. R. (2016). Plastic
and evolutionary gene expression responses are correlated in European grayling
(Thymallus thymallus) subpopulations adapted to different thermal environments. *Journal*
of Heredity, *107*(1), 82–89. doi: 10.1093/jhered/esv069
Mallard, F., Jakšić, A. M., & Schlötterer, C. (2018). Contesting the evidence for non-adaptive
plasticity. *Nature*. doi: 10.1038/nature25496
Manceau, M., Domingues, V. S., Linnen, C. R., Rosenblum, E. B., & Hoekstra, H. E. (2010).
Convergence in pigmentation at multiple levels: mutations, genes and function.
Philosophical Transactions of the Royal Society B: Biological Sciences, *365*(1552), 2439–
2450. doi: 10.1098/rstb.2010.0104
Mandic, M., Ramon, M. L., Gerstein, A. C., Gracey, A. Y., & Richards, J. G. (2018). Variable
gene transcription underlies phenotypic convergence of hypoxia tolerance in sculpins. *BMC*
Evolutionary Biology, *18*, 163. doi: 10.1186/s12862-018-1275-1
Manousaki, T., Hull, P. M., Kusche, H., MacHado-Schiaffino, G., Franchini, P., Harrod, C., …
Meyer, A. (2013). Parsing parallel evolution: Ecological divergence and differential gene
expression in the adaptive radiations of thick-lipped Midas cichlid fishes from Nicaragua.
Molecular Ecology. doi: 10.1111/mec.12034
Marder, E., & Goaillard, J. M. (2006). Variability, compensation and homeostasis in neuron and
network function. *Nature Reviews Neuroscience*, *7*, 563–574. doi: 10.1038/nrn1949
Oke, K. B., Bukhari, M., Kaeuffer, R., Rolshausen, G., Räsänen, K., Bolnick, D. I., … Hendry, A.
P. (2016). Does plasticity enhance or dampen phenotypic parallelism? A test with three
lake-stream stickleback pairs. *Journal of Evolutionary Biology*. doi: 10.1111/jeb.12767
Paccard, A., Hanson, D., Stuart, Y. E., Von Hippel, F. A., Kalbe, M., Klepaker, T., … Barrett, R.
D. H. (2020). Repeatability of Adaptive Radiation Depends on Spatial Scale: Regional
Versus Global Replicates of Stickleback in Lake Versus Stream Habitats. *Journal of*
Heredity. doi: 10.1093/jhered/esz056
Pankey, M. S., Minin, V. N., Imholte, G. C., Suchard, M. A., & Oakley, T. H. (2014). Predictable
transcriptome evolution in the convergent and complex bioluminescent organs of squid.

875 *Proceedings of the National Academy of Sciences of the United States of America*,
876  *111*(44), E4736–E4742. doi: 10.1073/pnas.1416574111
877 Pespeni, M. H., Sanford, E., Gaylord, B., Hill, T. M., Hosfelt, J. D., Jaris, H. K., … Palumbi, S. R.
878  (2013). Evolutionary change during experimental ocean acidification. *Proceedings of the*
879  *National Academy of Sciences*, *110*(17), 6937–6942. doi: 10.1073/pnas.1220673110
880 Pigliucci, M. (2010). Genotype-phenotype mapping and the end of the "genes as blueprint"
881  metaphor. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*,
882  557–566. doi: 10.1098/rstb.2009.0241
883 Potticary, A. L., Morrison, E. S., & Badyaev, A. V. (2020). Turning induced plasticity into refined
884  adaptations during range expansion. *Nature Communications*. doi: 10.1038/s41467-020-
885  16938-7
886 Renn, S. C. P., & Schumer, M. E. (2013). Genetic accommodation and behavioural evolution :
887  insights from genomic studies. *Animal Behaviour*, *85*(5), 1012–1022. doi:
888  10.1016/j.anbehav.2013.02.012
889 Reznick, D. A., Bryga, H., & Endler, J. A. (1990). Experimentally induced life-history evolution in
890  a natural population. *Nature*, *346*(6282), 357–359. doi: 10.1038/346357a0
891 Reznick, D., Butler IV, M. J., & Rodd, H. (2001). Life-History Evolution in Guppies. VII. The
892  Comparative Ecology of High- and Low-Predation Environments. *The American Naturalist*,
893  *157*(2), 126–140. doi: 10.1086/318627
894 Reznick, D. N. (1997). Life history evolution in guppies (Poecilia reticulata): Guppies as a model
895  for studying the evolutionary biology of aging. *Experimental Gerontology*, *32*(3), 245–258.
896  doi: 10.1016/S0531-5565(96)00129-5
897 Reznick, D. N., & Bryga, H. (1987). Life-History Evolution in Guppies (Poecilia reticulata): 1.
898  Phenotypic and Genetic Changes in an Introduction Experiment. *Evolution*, *41*(6), 1370.
899  doi: 10.2307/2409101
900 Rice, T. K., Schork, N. J., & Rao, D. C. (2008). Methods for Handling Multiple Testing.
901  *Advances in Genetics*, *60*, 293–308. doi: 10.1016/S0065-2660(07)00412-9
902 Rosenblum, E. B., Rompler, H., Schoneberg, T., & Hoekstra, H. E. (2010). Molecular and
903  functional basis of phenotypic convergence in white lizards at White Sands. *Proceedings of*
904  *the National Academy of Sciences*, *107*(5), 2113–2117. doi: 10.1073/pnas.0911042107
905 Ruell, E. W., Handelsman, C. A., Hawkins, C. L., Sofaer, H. R., Ghalambor, C. K., & Angeloni,
906  L. (2013). Fear, food and sexual ornamentation: plasticity of colour development in
907  Trinidadian guppies. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1758),
908  20122019–20122019. doi: 10.1098/rspb.2012.2019
909 Schaum, E., Rost, B., Millar, A. J., & Collins, S. (2013). Variation in plastic responses of a
910  globally distributed picoplankton species to ocean acidification. *Nature Climate Change*,
911  *3*(3), 298–302. doi: 10.1038/nclimate1774
912 Scoville, A. G., & Pfrender, M. E. (2010). Phenotypic plasticity facilitates recurrent rapid
913  adaptation to introduced predators. *Proceedings of the National Academy of Sciences*,
914  *107*(9), 4260–4263. doi: 10.1073/pnas.0912748107
915 Shaw, J. R., Hampton, T. H., King, B. L., Whitehead, A., Galvez, F., Gross, R. H., … Stanton, B.
916  A. (2014). Natural selection canalizes expression variation of environmentally induced
917  plasticity-enabling genes. *Molecular Biology and Evolution*, *31*(11), 3002–3015. doi:
918  10.1093/molbev/msu241
919 Siegal, M. L., & Leu, J.-Y. (2014). On the Nature and Evolutionary Impact of Phenotypic
920  Robustness Mechanisms. *Annual Review of Ecology, Evolution, and Systematics*, *45*(1),
921  495–517. doi: 10.1146/annurev-ecolsys-120213-091705
922 Song, L., & Florea, L. (2015). Rcorrector: Efficient and accurate error correction for Illumina
923  RNA-seq reads. *GigaScience*, *4*(1), 1–8. doi: 10.1186/s13742-015-0089-y
924 Stern, D. L. (2013). The genetic causes of convergent evolution. *Nature Reviews Genetics*,
925  *14*(11), 751–764. doi: 10.1038/nrg3483

Storey, J. D. (2002). A direct approach to false discovery rates. *Journal of the Royal Statistical*
Society. Series B: Statistical Methodology, *64*, 479–498. doi: 10.1111/1467-9868.00346
Storey, J. D. (2007). The optimal discovery procedure: A new approach to simultaneous
significance testing. *Journal of the Royal Statistical Society. Series B: Statistical*
Methodology. doi: 10.1111/j.1467-9868.2007.005592.x
Strimmer, K. (2008). fdrtool: A versatile R package for estimating local and tail area-based false
discovery rates. *Bioinformatics*, *24*(12), 1461–1462. doi: 10.1093/bioinformatics/btn209
Stuart, Y. E., Veen, T., Weber, J. N., Hanson, D., Ravinet, M., Lohman, B. K., … Bolnick, D. I.
(2017). Contrasting effects of environment and genetics generate a continuum of parallel
evolution. *Nature Ecology and Evolution*. doi: 10.1038/s41559-017-0158
Thompson, K. A., Osmond, M. M., & Schluter, D. (2019). Parallel genetic evolution and
speciation from standing variation. *Evolution Letters*, *3*(2), 129–141. doi: 10.1002/evl3.106
Torres-Dowdal, J., Handelsman, C. a, Reznick, D. N., & Ghalambor, C. K. (2012). Local
adaptation and the evolution of of phenotypic plasticity in Trinidadian guppies (Poecilia
reticulata). *Evolution*, *66*(11), 3432–3443. doi: 10.5061/dryad.84gf5
Torres Dowdall, J., Handelsman, C. A., Ruell, E. W., Auer, S. K., Reznick, D. N., & Ghalambor,
C. K. (2012). Fine-scale local adaptation in life histories along a continuous environmental
gradient in Trinidadian guppies. *Functional Ecology*, *26*(3), 616–627. doi: 10.1111/j.1365-
2435.2012.01980.x
van der Vaart, A. W., & Wellner, J. A. (1996). Weak Convergence and Empirical Processes:
With Applications to Statistics. In *Springer Science+Business Media, LLC*.
Velotta, J. P., & Cheviron, Z. A. (2018). Remodeling Ancestral Phenotypic Plasticity in Local
Adaptation : A New Framework to Explore the Role of Genetic Compensation in the
Evolution of Homeostasis. *Integrative and Comparative Biology*, *58*(6), 1098–1110. doi:
10.1093/icb/icy117
Wagner, A. (2011). *The origins of evolutionary innovations: a theory of transformative change in*
living systems. Oxford: Oxford University Press.
Wake, D. B., Wake, M. H., & Specht, C. D. (2011). Homoplasy : From Detecting Pattern and
Mechanism of Evolution. *Science*, *331*(April), 1032–1035. doi: 10.1126/science.1188545
Wang, S. P., & Althoff, D. M. (2019). Phenotypic plasticity facilitates initial colonization of a
novel environment. *Evolution*, *73*(2), 303–316. doi: 10.1111/evo.13676
West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. doi: 10.2002/ajpa.20219
Willing, E. M., Bentzen, P., Van Oosterhout, C., Hoffmann, M., Cable, J., Breden, F., … Dreyer,
C. (2010). Genome-wide single nucleotide polymorphisms reveal population history and
adaptive divergence in wild guppies. *Molecular Ecology*, *19*(5), 968–984. doi:
10.1111/j.1365-294X.2010.04528.x
Zandonà, E., Auer, S. K., Kilham, S. S., Howard, J. L., López-Sepulcre, A., O'Connor, M. P., …
Reznick, D. N. (2011). Diet quality and prey selectivity correlate with life histories and
predation regime in Trinidadian guppies. *Functional Ecology*, *25*(5), 964–973. doi:
10.1111/j.1365-2435.2011.01865.x
966

Glossary

convergent evolution	The evolution of similar phenotypes in multiple independent populations, in response to similar selection pressures, but from different initial conditions.
parallel evolution	The evolution of similar phenotypes in multiple independent populations, in response to similar selection pressures, and from similar initial conditions.
genetic divergence	Differences between populations adapted to distinct environments. Here specifically, genetic differences in gene expression between high- and low-predation fish. These differences are identified through a common garden breeding design.
phenotypic plasticity	The potential for multiple phenotypes to arise from a single genotype, e.g. in response to environmental changes.
ancestral plasticity	The phenotypic plasticity present in individuals/populations adapted to ancestral environmental conditions when exposed to environmental conditions reflecting the derived environment. Here specifically, the plastic response of high-predation fish to rearing in predator free environments. This plasticity may be adaptive or non-adaptive (see below).
adaptive plasticity	Phenotypic plasticity that brings a trait closer to the optimum in the novel environment as compared to the trait value in the ancestral environment. Here specifically, gene expression plasticity in high-predation fish reared without predators that mirrors the direction of expression divergence between high- and low-predation adapted populations.
non-adaptive plasticity	Phenotypic plasticity that moves a trait further away from the optimum in the novel environment as compared to the trait value in ancestral environment. Here specifically, gene expression plasticity in high-predation fish reared without predators that is opposite the direction of expression divergence between high- and low-predation adapted populations.
concordant expression differences	When gene expression differences among parallel comparisons are in the same direction, i.e. either both up- or down-regulated across comparisons. Here specifically, when gene expression differences between low-predation populations and high-predation ancestors are either upregulated or downregulated in distinct drainages.
non-concordant expression differences	When gene expression differences among parallel comparisons are in opposite directions, i.e. up-regulated in one comparison and down-regulated in the other, or vice versa. Here specifically, when gene expression differences between high- and low-predation population in distinct drainages are in opposite directions.

Figures

Figure 1. Overview of experimental design and interpretation. (A) Schematic shows Aripo and Quare river drainages in the Northern Range Mountains of Trinidad. We sampled population pairs from these two drainages, which represent distinct evolutionary lineages. (B) Wild-caught fish from each high- and low-predation population pair were reared to the second generation in the lab to control for maternal and environmental effects. At birth, second generation siblings from each population were split into rearing environments with (pred+) and without (pred-) predation cues. (C) This rearing design allows us to discern lineage effects (contrast between drainages), genetic divergence (contrast between different populations in the same environment), phenotypic plasticity (contrast between genetically similar fish in different rearing condition), and their interactions.

Figure 2. Summary of differential expression analyses. We identified many transcripts in both the Aripo (left) and Quare (right) drainages whose expression levels differed based on evolutionary history of predation (population of origin; orange), developmental experience of predation (rearing; yellow), and their interaction (grey).

Figure 3. Relationship between ancestral plasticity and evolved expression divergence. Genes exhibiting significant ancestral expression plasticity (i.e. different between high-predation fish reared with and without predator cues) were more likely to also differ in expression between high- and low-predation populations reared without predator cue exposure in both the Aripo (top) and Quare (bottom) lineages. Each circle represents expression differences in a single transcript with statistically significant ancestral plasticity in the high-predation population as well as significant expression differences between high- and low-predation populations. Plastic expression changes were more likely to be in the opposite (white circles) than in the same (grey circles) direction as genetic expression divergence. The number of transcripts in each quadrant is indicated on the graphs. Note that axes differ between plots in order to best visualize variation within each dataset.

Figure 4. Evolution of transcript expression plasticity. (A) We grouped transcripts with significant interaction effects into one of four categories based on patterns of evolution in expression plasticity: (1) Assimilated: plasticity in the ancestral high-predation population but a loss of plasticity in the derived low-predation population; (2) Accommodated: a change in the degree, but no the direction, of plasticity in the derived as compared to the ancestral population; (3) Evolved plastic: no plasticity in the ancestral high-predation population but an emergence of plasticity in the derived low-predation population; (4) Reversed: opposing directions of plasticity in high-versus low-predation populations. We categorized remaining transcripts that had a significant main interaction effect, but no significant *post hoc* rearing differences as unclassified. Adapted from Renn & Schumer (2013). (B) All categories were represented in both Aripo and Quare datasets.

Figure 5. Overlap in population and rearing expression differences across evolutionary lineages. Of the transcripts differentially expressed based on population of origin (top), 124 were overlapping between drainages. Of these, 66 were concordantly differentially expressed (i.e. expression divergence in the same direction in both drainages; dark orange circles) and 58 were not (i.e. expression divergence in opposite directions between drainages; light orange circles), no different than expected by chance. Of the transcripts differentially expressed based on rearing environment (bottom), 35 were overlapping in the two drainages. Of these, 19 were concordantly differentially expressed (dark yellow circles) and six were not (light yellow circles), more non-concordance than expected by chance. The number of differentially expressed transcripts in each quadrant is indicated on the graphs and transcripts differentially expressed in one but not both drainages are shown in grey.

Supplemental Information

Table S1. Transcripts filtered from the analysis following identification as retinal contaminants. Provided as Excel file.

Table S2. Transcripts significantly differentially expressed between high- and low-predation populations in the Aripo drainage. Provided as Excel file.

 Table S3. Transcripts significantly differentially expressed between high- and low-predation populations in the Quare drainage. Provided as Excel file.

Table S4. Results of GO enrichment analyses for population, rearing, and interaction effects in the Aripo drainage. Provided as Excel file.

Table S5. Results of GO enrichment analyses for population, rearing, and interaction effects in the Quare drainage. Provided as Excel file.

Table S6. Population DE transcripts overlapping between Aripo and Quare drainages. 185 transcripts were DE in both drainages, of these 89 had expression differences in the same direction in both datasets (concordant differential expression). Provided as Excel file.

Table S7. Rearing DE transcripts overlapping between Aripo and Quare drainages. 16 transcripts were DE in both drainages, of these 7 had expression differences in the same direction in both drainages (i.e. concordant differential expression). Provided as Excel file.

Table S8. Pst estimates for the Aripo and Quare drainages.

Figure S1. Differences in means but not variance of differentially expressed (DE) and non-differentially expressed (non-DE) transcripts. DE transcripts had significantly higher means in the Aripo drainage and lower means in the Quare drainage as compared to non-DE transcripts. Due to the large range of expression values, mean differences are plotted on a log scale (y-axis).

Figure S2. Outcome of down-sampling simulations. The Quare dataset was randomly down-sampled to generate a sample size equivalent to that of the Aripo dataset (40 samples, 10 per group). The histogram shows the distribution of the number of population DE genes obtained from 500 random down-samples. The number of DE genes in the full dataset (red dashed line) is not extreme compared to the distribution of the down-sampled datasets, indicating that sample size is not a critical factor for the difference in DE gene number between the two datasets.