1

Individual Learning Phenotypes Drive Collective Cognition

Chelsea N. Cook[1*], Natalie J. Lemanski[2], Thiago Mosqueiro[2], Cahit Ozturk[1], Jürgen

Gadau[3], Noa Pinter-Wollman[2◊], Brian H. Smith[1◊]

Affiliations

[1] School of Life Sciences, Arizona State University, Tempe, USA

[2] Department of Ecology and Evolutionary Biology, University of California Los Angeles,

Los Angeles, USA

[3] Institute for Evolution and Biodiversity, University of Münster, Münster, Germany

[◊]Co-Senior Authors

[*]Corresponding Author, cncook1@asu.edu, @PhDBee

Summary

Collective cognition allows animal groups to accomplish many tasks that could

not be accomplished as effectively alone, such as schools of fish avoiding predators [1],

flocks of birds moving thousands of miles across the Earth [2], and honey bee colonies

collecting food from millions of flowers [3]. Individuals in groups utilize local information

to quickly adjust to ecological changes, then implicitly or explicitly communicate that

information with group members to form a collective response [4–6]. Collective cognition

emerges from interactions between individuals that vary in their cognitive abilities [3,4].

Individual cognitive variation influences how animals pay attention to or share

information [7–9], although little is known about how variation in individual cognition

drives collective cognition. Here, we show that individual differences in a cognitive task

scales to shape collective foraging behavior in honey bees by utilizing a naturally

variable and heritable learning behavior called latent inhibition (LI) [10]. We artificially

selected two distinct phenotypes: high LI bees that are better at ignoring previously

unrewarding familiar stimuli, and low LI bees that learn previously unrewarding and

novel stimuli equally well. Colonies comprised of high LI individuals preferred to visit the

familiar food location, while low LI colonies visit novel and familiar food locations

equally. However, in colonies of mixed learning phenotypes, the low LI bees showed a

preference to visiting familiar feeders, which contrasts with their behavior when in a

uniform group. We show that the shift in the feeder preference of low LI bees is driven

by foragers of the high LI phenotype dancing more intensely and attracting more

followers. Our results reveal that cognitive abilities of individuals drives emergent

collective behavior.

Key Words: Collective cognition, collective behavior, learning, latent inhibition,

honey bees, foraging,

Results and Discussion

To uncover how individual variation in learning impacts group-level behaviors, we

investigated how an individual's latent inhibition (LI) may scale to influence collective

foraging behavior in honey bee colonies. Although LI has been mostly studied in

vertebrates [11–15] and is associated with attention disorders in humans[16]. Honey

bees (*Apis meillifera*) also exhibit[17,18] and show variation in LI[10]. Despite several

studies showing heritable variation in latent inhibition[10,19], and how it is relevant in

individual ecological contexts such as predator avoidance[11,12,20], it remains unclear

as to whether or how this variation functions in ecologically relevant decisions in

complex social environments. Foraging honey bees vary in their expression of LI;

scouts tend to exhibit high LI and ignore familiar odors, while recruits who tend to exhibit

low LI tend to learn familiar and novel odors equally well[21]. LI is heritable in honey

bees[18]. Therefore, we selected queens and drones (haploid males) for expression of

LI and created selected lines from singly inseminated queens with like performing

drones to produce two distinct lines of workers that exhibit similar LI to their parents[22].

We created 24 colonies composed of single cohorts of only low, only high, 50/50 mixed

high and low LI workers, as well as age-matched non-selected control bees. In semi-

natural foraging conditions, we counted the number of forager visits, first visits, and

revisits to the familiar or novel food locations. To explore the mechanisms underlying

how individual variation in LI affects collective foraging, we quantified the round

recruitment dance activity in 6 mixed colonies while the colonies visited novel and

familiar feeders. These field assays allowed us to simultaneously quantify individual and

colony level foraging behaviors.

To ensure workers in different social environments exhibited the predicted

heritable LI phenotype, we evaluated the LI score of foragers after 21 days in either

their natal colony or a control colony. We marked 1000 individuals from each selected

line (high or low LI) on the day of emergence. We then placed 500 individuals back into

their natal colony and 500 individuals into an established control colony of equal size

with an open mated queen, i.e. workers with a variety of learning phenotypes. We

monitored the colonies until marked bees began to make foraging flights (~21 days). We

then collected marked foragers as they returned to the colony and brought them into the

laboratory to evaluate their LI. We avoided pollen foragers as they tend to exhibit

different learning behavior compared to nectar foragers[22]. We found that foragers

retained the expected LI based on the LI of their parents, regardless of whether they

were housed with same or with variable learning phenotypes. Foragers from the high

and low lines differed in expression of LI as expected (GLM: χ^2 = 4.84, df=1, p=0.027,

Figure 1). We did not detect an effect of the identity of the colony in which the bees

were housed on LI phenotype (χ^2 = 3.28, df=2, p=0.193, Figure 1).

100
101

Figure 1: Social environment does not alter expression of genetically selected
latent inhibition. LI scores of individuals from high LI lines that spent their adult life
either in high LI only colonies (red, n=36) or in a control colony with a variety of LI
phenotypes from an open mated queen (red with gray vertical lines, n=18);
individuals from low LI lines that spent their adult life either in low LI only colonies
(yellow, n=52) or in control colonies (yellow with gray vertical lines, n=10). In this and
subsequent figures, the large black dot is the mean, the white box is the interquartile

range (IQR), whiskers extend to 1.5*IQR, and the small points beyond the whiskers are outliers. Shaded areas show the distribution of the data. Here, and in all following figures, yellow are low LI colonies and individuals, gray are control colonies and individuals, and red are high LI colonies and individuals. Attention is critical for many individual behaviors, including finding the correct mate or prey.

To determine how the learning phenotypes influenced colony-level foraging behavior, we placed small single-cohort (same age bees) colonies into a flight cage and monitored foraging activity. We evaluated 4 colony types each week: one control colony consisting of approximately 1300 age-matched bees from open mated queens; one colony consisting of 650 workers from high LI queens plus 650 age-matched control bees; one colony consisting of 650 workers from low LI queens plus 650 aged-matched control bees; and one 50/50 mixed colony with 325 workers from each LI line plus 650 aged-matched control bees. In the last 3 types, the supplemented 650 age-matched bees from open mated queens were used to ensure a small but functioning colony as we did not have enough workers from the single-drone-inseminated queens and colonies of just 650 individuals would be too weak to forage. Honey bee division of labor is largely influenced by worker age, so we used age-matched bees to remove any influence that age may have on foraging propensity. On day 1, we trained bees to a feeder inside the tent containing 1M sucrose and an odor, which became the 'familiar' feeder. During the subsequent 3 days, in addition to the familiar feeder, we introduced a single novel feeder each day with a different odor and color, but with the same sugar concentration as the familiar feeder (Figure 2A). To evaluate the collective ability of the colony to find a new feeder, we recorded the number of visits to each feeder by bees from each selected line according to the color of paint on the bees' thorax. We further marked bees with a feeder-specific color on their abdomen when they visited the feeder for the first time to determine if bees revisited that feeder. We repeated this for 6 weeks on 6 colonies for each group type.

Colony composition strongly influenced overall number of visits to the food locations (N = 6 colonies in each line, 24 total, 6172 total visits; GLM: χ^2 = 1270, df = 3, p < 0.0001, Figure 2B). High LI colonies had significantly more visits to all food locations compared to low LI colonies (Tukey post hoc: Z=25.5, p <0.0001, Figure 2A), mixed

colonies (Z=5.18, p<0.0001), and controls (Z=26.6, p<0.0001). Mixed LI colonies also had significantly more visits compared to low (Z=20.7, p < 0.0001) and controls (Z=21.8, p<0.0001). Low LI and control colonies had the fewest total visits and were not significantly different from each other (Z=-1.38, p=0.50).

Foraging in the high, low, and control colonies was largely performed by bees revisiting the feeders. (GLM, χ^2 = 22.32, df =3, p<0.0001, Figure 2C). However, the mixed LI colonies had a significantly lower proportion of revisiting foragers compared to the low (Tukey post hoc: Z=-4.2, p=0.0002), high (Z=-3.1, p=0.01), and control colonies (Z=-3.33, p=0.004). We did not detect significant differences among the other colony types (See Table S3).

A colony's LI phenotype composition determined its preference between the novel and familiar feeders (GLM: Feeder*Colony χ^2 =473.64, df=3, p<0.0001; Figure 2D). High and mixed colonies preferred the familiar feeder over the novel one (Tukey Poshoc: High Familiar:Novel: Z=20.2, p<0.0001; Mixed Familiar:Novel: Z=25.6, p<0.0001). Low LI and control colonies did not show a strong preference for either feeder, visiting them equally (Low Familiar:Novel: Z=-1.24, p=0.92; Control Familiar:Novel: Z=2.03, p=0.46). For full pairwise comparisons, see Table S4).

The number of re-visits to the novel and familiar feeders was different across colony compositions (Figure 2E: Colony*Feeder χ^2 =53.67, p<0.0001). All colonies had a higher proportion of re-visits to the familiar feeder compared to the novel feeder. However, the mixed LI colonies had a much lower proportion of re-visitation to the novel feeders than the other colony types (Table S5). Thus, new foragers in the mixed colonies that visited the novel feeder were less likely to return to it compared to foragers who visited the novel feeders in other colonies.

Figure 2: Colonies constructed from different genetic lines selected for high or low latent inhibition exhibited differences in collective foraging behavior. (A) The experimental set up illustrating the location of feeders in relation to the location of the colony (center, white) within the experimental arena (large rectangle). The familiar feeder (red) was provided on day 1 and on all subsequent days. Novel feeder X (blue) was presented on day 2, novel feeder Y (purple) on day 3, and novel feeder Z (orange) on day 4. See Table S2 for associated odors. Visits to all novel feeders were combined for statistical analysis. (B) Cumulative number of visits of bees to all feeders over time by colony type. Different letters to the right of the lines indicate statistically significant differences according to a post hoc Tukey test. For further illustration of visitation by

each colony on each day, see Figure S1. (C) Percent of re-visits out of the total number
of visits to all feeders by colony type. Here and in all following panels, different letters
above boxes indicate statistically significant differences according to a post hoc Tukey
test. (D) Number of all visits to the familiar feeder (solid boxes) and a novel feeder
(hatched boxes) for each type of colony, when both novel and familiar feeders were
presented simultaneously (days 2-4). (E) Percent of re-visits out of the total number of
visits to either the familiar or the novel feeder by type of colony when both novel and
familiar feeders were presented simultaneously (days 2-4). In C, D and E, horizontal
lines are the median, the boxes are the interquartile range (IQR), whiskers extend to
1.5*IQR, and the small points beyond the whiskers are outliers. N=24 colonies, 6
colonies per group type, 6172 total visits.
190
To determine why the mixed colonies showed a preference for the familiar feeder
(Figure 2D), we examined how individual lines visited each feeder (Figure 3). In 2017,
we tested mixed colonies placed in a flight cage. In 2018, we reselected lines and then
placed mixed colonies into two-frame observation hives to evaluate recruitment dances
along with visitation to the feeders in the flight cages. We found that there was a
significant year effect (Table S6), likely due to reselection and different environmental
conditions. We therefore statistically analyzed each year separately to focus on the
within-year variation between the selected lines.
Low LI and control individuals shift their preference to the familiar feeder when
mixed with high LI bees. In 2017, we found a significant interaction between the
selected line and which feeder foragers visited (GLM: χ^2 =7.79, df=2, p=0.02; Figure
3A). Although low LI and control colonies did not show a preference to a novel or
familiar feeder when they had a uniform colony composition (Figure 2E), when mixed
with high LI individuals, low LI and control individuals exhibited a preference to the
familiar feeder (GLM: Low Familiar:Novel: Z=13.28, p<0.0001; Control Familiar:Novel:
Z=18.32, p<0.0001; Figure 3A). High LI individuals showed a preference to familiar
feeders (GLM: Familiar:Novel: Z=22.03, p<0.0001) just as colonies comprised of only
high LI individuals did (Figure 2E). We found a significant interaction between selected
line and feeder in 2018 (GLM: χ^2 =85.27, df=2, p<0.0001; Figure 3B), with low LI and
control individuals showing preference to the familiar feeder over the novel feeder
(GLM: Low Familiar:Novel: Z=25.05, p<0.0001; Control Familiar:Novel: Z=13.90,
p<0.0001; Figure 3B) similar to high LI individuals preferring the familiar feeders

(Familiar:Novel: Z=18.48, p<0.0001). For full contrasts from 2017 see Table S7, for

2018 see Table S8.

216
217
218
219

Figure 3: Visits of individuals from different genetically selected lines when in a mixed colony.

220
221
222
Figure 3: Visits of individuals from different genetically selected lines when in a
mixed colony. Daily visits to the familiar (solid) and novel (hatched) feeders by
individual bees in mixed colonies from low LI parents (yellow), high LI parents (red) or
open mated queens (grey) in (A) 2017, N=6 mixed colonies, 2347 overall visits and (B)
in mixed colonies from lines that were re-selected in 2018, N=6 colonies, 6272 overall
visits. The horizontal line in the box is the median, the box is 25-75% of the data,
whiskers represent 95% of the data, and diamonds show outliers beyond 95%. Different
letters above boxes indicate statistically significant differences according to a post hoc
Tukey test.

To uncover the behavioral mechanisms that underlie the switching of low LI

individuals from having no feeder preference when in a uniform colony composition to

preferring the familiar feeder when in a mixed colony, we examined the round dance,

the modified waggle dance used at short distances[23], of individuals from each

selected line in mixed colonies as they returned from foraging. Using observation hives

with glass walls, we video recorded bees performing the round dance to recruit other

individuals in the colony to forage. To determine which selected line recruited to each

feeder, we noted the selected line of the dancer (high or low LI) according to the paint

marks on the individuals' thorax and whether the dancer had visited a feeder according

to the paint marks on abdomens. We did not record dancers without abdominal marks

as they were likely collecting from and recruiting to unmonitored water sources. To

determine who the information about a feeder was communicated to, we counted the

number of followers of each dancer and the selected line of the followers. To quantify

the dance intensity, we recorded the duration of the dance, and the number of turns the

dancer made during the first 20 seconds of the dance.

Individuals from the lines differed in their likelihood to perform a round dance

(Chi-square test: χ^2=26.61, df=2, p<0.0001; Figure 4B). Low LI individuals were

significantly more likely to perform a dance compared to high Li individuals (pairwise

chi-square test: p=0.0001) and controls (pairwise chi-squared test: p=0.004). High LI

individuals were just as likely to perform a dance as controls (p=0.36). Individuals

differed in their likelihood to follow a dance based on their selected line (Chi-square test:

χ^2=28.26, df=2, p<0.0001; Figure 4B). Low LI individuals were significantly more likely

to follow a dance compared to high LI bees (pairwise chi-square test: p<0.0001) and

controls (pairwise chi-square test: p<0.003). High LI and control individuals were equally

likely to follow a dance (pairwise chi-square test: p=0.240).

Although the high LI individuals danced less often, high LI dances had

significantly more followers compared to low and control bees (Chi-square test: χ^2=

13.93, df=2, p<0.001; Figure 4C). Low LI bees performed more dances that had no

followers compared to high LI and control dances. We did not detect a statistically

significant difference between the proportion of individuals from each line that followed

each line of dancer (Chi-square test: χ^2= 7.05, df = 4, p= 0.13, Figure 4D). Low LI

individuals spent more time dancing; however they attracted fewer followers than high

and control dancers, indicated by the significant interaction between the LI of the dancer

and dance duration when predicting the number of followers (GLMM: χ^2= 6.42, df=2,

p=0.04; Figure 4E).

The relative attraction of dances of high LI bees could be due to the intensity of

the dance. High LI bees performed more turns per second during their dances (ANOVA:

χ^2=12.8, df=2, p=0.002; Figure 4F). High LI dancers performed an average of 0.59 turns

per second, significantly higher than low LI dancers, who performed an average of 0.52

turns per second (Tukey: t=-3.13, p=0.005). Control bees also performed more turns per

second than low LI bees (Tukey: t=-2.5, p=0.03), but not different than high LI bees, at

an average 0.62 turns per second (Tukey: t=-0.7, p=0.7).

Figure 4: Recruitment dances facilitate integration of information from different genetically selected lines. (A) The experimental set up illustrating the location of feeders in relation to the location of the colony entrance (top right, white) within the experimental arena (large rectangle). The familiar feeder (red) was provided on day 1

and on all subsequent days. Novel feeder X (blue) was presented on day 2, novel
feeder Y (purple) on day 3, and novel feeder Z (orange) on day 4. See table S2 for
associated odors. Visitation to novel feeders were combined for statistical analysis. (B)
Proportion of dances (N=667) or follows (N=1201) across 6 colonies performed by bees
from each line, relative to their abundance in the mixed colony (350 high, 350 low, 700
control). We accounted for the difference in abundance of each selected line by dividing
the number of observed control dancers by 2 before calculating these proportions. (C)
Proportion of dances performed per LI line type that were either followed by at least one
individual (colored) or not followed by any other bees (black). (D) Proportion of dances
by LI line type that were followed (from panel B) broken down by LI of the follower. (E)
Relationship between number of followers and duration of a dance by line. Point and
line colors indicate LI of dancer. Best fit line represents the GLM, shaded area
represents the 95% confidence interval. (F) Rate of turns per second in a dance by line.
The large black dot in the box is the mean, the box is 25-75% of the data, whiskers
represent 95% of the data. The violin shapes illustrate distribution of the data. Different
letters above violins indicate statistically significant differences according to a post hoc
Tukey test.
299
300

By combining techniques from experimental psychology and behavioral ecology,
we have developed a system for investigating how variation in individual learning
behavior drives collective cognition. We utilized this system to demonstrate that a
laboratory-selected heritable learning behavior with natural individual variation scales to
shape the collective performance of a honey bee colony on foraging tasks. In the lab,
high LI honey bees learn to ignore familiar odors that they experienced without
reinforcement, while readily learning novel odors. When a stimulus is rewarding, high LI
bees exhibit increased attention to that information. One interpretation of reduced
learning to a familiar, unrewarding, stimulus is that pre-exposure reduces attention to,
and thus associability of, that stimulus. This interpretation is an extension of conditioned
attention theory[24,25], which proposes that latent inhibition is induced by allowing
animals to focus their attention on important information[14,26,27]. Our observations of
field behavior of low and high LI individuals and colonies are consistent with this
interpretation, whereby high LI individuals have stronger attention capacities to food
compared to low LI individuals. Once high LI individuals have found a food location,
they continue to revisit it, 'attending' more strongly to reinforced feeders over new ones.

The increased impact of the resource on these bees could translate into stronger, more

vigorous dances. In contrast, low LI individuals learn and visit both known and new

feeders equally, dividing their attention across resources and acting more like generalist

foragers. In mixed colonies, this broadened attention by low LI individuals may therefore

make them the perfect audience for the high LI dancers, driving them to prefer feeders

that high LI individuals preferentially visit. In fact, individual phenotypes that emerge

from these lines in natural conditions seem to be collectively complementary "finders"

and "refiners", where high LI bees discover food and direct other bees to that location,

while the low LI bees refine that information, finding the most rewarding resource by

sampling a broader spectrum of available resources (Lemanski et al. In Review). Under

natural conditions, where queens mate with many different drones, most colonies would

possess both types of learners, perhaps more closely resembling our mixed

colonies[28]. Attention is critical for many individual behaviors, including finding the

correct mate or prey[29]. We propose that this diversity of 'attention' aspect of individual

cognitive phenotypes may enhance the overall efficacy with which a group Attention is

critical for many individual behaviors, including finding the correct mate or prey. finds

and exploits resources[30]. In summary, our work indicates that individual cognition

scales to shape the collective cognition of animals solving critical ecologically relevant

tasks.

Acknowledgements: We thank S. Ohrt, E Sezen, N Kulkarni, and A Phillips for help
with data collection. We thank C. Kwapich and D. Charbonneau for comments on drafts
of this manuscript. This grant was funded by NIH NIGMS R01GM113967 to BHS, JG, &
NPW, and NIH NIGMS F32GM126728 to CNC.

Author Contributions: CNC, TM, JG, NPW, BHS conceived of and helped design the
study. CNC, NJL, and TM analyzed the data. CNC and CO created the genetically
selected lines and CO maintained them. CNC carried out data collection and wrote the
first draft of the manuscript. CNC, TM, NJL, NPW, BHS discussed results. All authors
commented on the manuscript.

Declaration of Interests: The authors declare no competing financial interests.

Data Availability: Data will be available on FigShare and code will be available on
Github upon publication. Data and code available upon request by reviewers.

FIGURE LEGENDS

**Figure 1: Social environment does not alter expression of genetically selected
latent inhibition.** LI scores of individuals from high LI lines that spent their adult life
either in high LI only colonies (red, n=36) or in a control colony with a variety of LI
phenotypes from an open mated queen (red with gray vertical lines, n=18);
individuals from low LI lines that spent their adult life either in low LI only colonies
(yellow, n=52) or in control colonies (yellow with gray vertical lines, n=10). In this and
subsequent figures, the large black dot is the mean, the white box is the interquartile
range (IQR), whiskers extend to 1.5*IQR, and the small points beyond the whiskers
are outliers. Shaded areas show the distribution of the data. Here, and in all following
figures, yellow are low LI colonies and individuals, gray are control colonies and
individuals, and red are high LI colonies and individuals. Attention is critical for many
individual behaviors, including finding the correct mate or prey.

**Figure 2: Colonies constructed from different genetic lines selected for high or
low latent inhibition exhibited differences in collective foraging behavior.** (A) The
experimental set up illustrating the location of feeders in relation to the location of the
colony (center, white) within the experimental arena (large rectangle). The familiar

feeder (red) was provided on day 1 and on all subsequent days. Novel feeder X (blue)
was presented on day 2, novel feeder Y (purple) on day 3, and novel feeder Z (orange)
on day 4. See Table S2 for associated odors. Visits to all novel feeders were combined
for statistical analysis. (B) Cumulative number of visits of bees to all feeders over time
by colony type. Different letters to the right of the lines indicate statistically significant
differences according to a post hoc Tukey test. For further illustration of visitation by
each colony on each day, see Figure S1. (C) Percent of re-visits out of the total number
of visits to all feeders by colony type. Here and in all following panels, different letters
above boxes indicate statistically significant differences according to a post hoc Tukey
test. (D) Number of all visits to the familiar feeder (solid boxes) and a novel feeder
(hatched boxes) for each type of colony, when both novel and familiar feeders were
presented simultaneously (days 2-4). (E) Percent of re-visits out of the total number of
visits to either the familiar or the novel feeder by type of colony when both novel and
familiar feeders were presented simultaneously (days 2-4). In C, D and E, horizontal
lines are the median, the boxes are the interquartile range (IQR), whiskers extend to
1.5*IQR, and the small points beyond the whiskers are outliers. N=24 colonies, 6
colonies per group type, 6172 total visits.

Figure 3: Visits of individuals from different genetically selected lines when in a
mixed colony. Daily visits to the familiar (solid) and novel (hatched) feeders by
individual bees in mixed colonies from low LI parents (yellow), high LI parents (red) or
open mated queens (grey) in (A) 2017, N=6 mixed colonies, 2347 overall visits and (B)
in mixed colonies from lines that were re-selected in 2018, N=6 colonies, 6272 overall
visits. The horizontal line in the box is the median, the box is 25-75% of the data,
whiskers represent 95% of the data, and diamonds show outliers beyond 95%. Different
letters above boxes indicate statistically significant differences according to a post hoc
Tukey test.

Figure 4: Recruitment dances facilitate integration of information from different
genetically selected lines. (A) The experimental set up illustrating the location of
feeders in relation to the location of the colony entrance (top right, white) within the
experimental arena (large rectangle). The familiar feeder (red) was provided on day 1
and on all subsequent days. Novel feeder X (blue) was presented on day 2, novel
feeder Y (purple) on day 3, and novel feeder Z (orange) on day 4. See table S2 for
associated odors. Visitation to novel feeders were combined for statistical analysis. (B)
Proportion of dances (N=667) or follows (N=1201) across 6 colonies performed by bees
from each line, relative to their abundance in the mixed colony (350 high, 350 low, 700
control). We accounted for the difference in abundance of each selected line by dividing

the number of observed control dancers by 2 before calculating these proportions. (C)

Proportion of dances performed per LI line type that were either followed by at least one

individual (colored) or not followed by any other bees (black). (D) Proportion of dances

by LI line type that were followed (from panel B) broken down by LI of the follower. (E)

Relationship between number of followers and duration of a dance by line. Point and

line colors indicate LI of dancer. Best fit line represents the GLM, shaded area

represents the 95% confidence interval. (F) Rate of turns per second in a dance by line.

The large black dot in the box is the mean, the box is 25-75% of the data, whiskers

represent 95% of the data. The violin shapes illustrate distribution of the data. Different

letters above violins indicate statistically significant differences according to a post hoc

Tukey test.

SUPPLEMENTAL MATERIALS (at end of document)

Tables S1-S8

Figure S1

METHODS

Obtaining queens and drones

To obtain queens for producing selected lines of a specific LI behavior, we

performed the LI assay as outlined in[10,21] on mature virgin queens 10 days after

emergence. Briefly, we familiarized bees to an odor by puffing it at them 40 times every

5 minutes, then used the proboscis extension reflex to test their ability to learn to

associate a food reward to the familiar versus a novel odor. Tested queens were placed

into individually labelled queen cages and returned to a queenless colony until

insemination, which typically occurred within a week of testing. To obtain fertile drones,

we collected them from their returning unsuccessful mating flights at the entrance of

colonies. We placed them into cages overnight in a queenless colony for LI testing the

next day. After testing, drones were marked for individual identification and placed into a

cage and placed into a queen bank for no longer than 3 days until inseminations

occurred.

Queen Inseminations

We used instrumental insemination to inseminate a queen with sperm from a

single drone. We inseminated a high LI queen with a high LI drone, and a low LI queen

with a low LI drone[31,32]. LI varies across individuals. However, for this behavioral

selection, we used the highest and lowest LI scoring individuals to create the high and

low colonies, respectively. We then introduced queens to small queenless colonies,

then allowed the queens to produce workers for 1 month. Colonies were checked

weekly to eliminate the possibility of supersedure.

Cohoused Worker Preparation and Testing

To test the LI of foragers from each LI line, we placed frames of capped pupae

from 3 high and 3 low LI colonies into 34°C incubators for 18 hours. After 18 hours, we

used water based acrylic paint pens (Montana brand) to mark the abdomens of the

eclosed bees with a color indicating their natal colony. Half of the bees were then

returned to their natal colony and half were placed into an established control colony of

an open-mated queen. Fewer bees were recovered from the established colony as

many are recognized as non-nestmates and rejected. After 2 weeks, colonies were

monitored every day until marked bees began to forage, ~21 days after emergence.

Returning nectar foragers were collected and tested for LI.

Field Colony Experimental Setup

To explore the colony-level foraging behavior of the LI lines, we set up 4

treatment colonies for each of the colony types: a high LI colony, a low LI colony, a

50/50 mixed colony, and a control. We ran weekly field experiments for 6 weeks. For

ease of identification, we always marked individuals from high LI colonies red, orange,

and pink, and individuals from low LI colonies green, blue, yellow, and white. We

continued to mark emerging bees from the same frames until we had 650 bees to form

a colony, which took typically 2-3 days. To achieve relatively normal conditions for

typical honey bee behavior, we supplemented workers from an unselected colony

(control bees), who were not marked. For colony set up, see Table S1. Bees were then

placed into 4 different treatment colonies consisting of ~1300 bees: high plus controls,

low plus controls, 50/50 mixed high/low plus controls, and only control colony. Bees

were provided a honeycomb and remained inside for 5 days before being placed for

field experimentation. We then placed nucleus colonies into outdoor flight cages (3.05m

x 12m) and replaced the honeycomb frame with an empty frame to induce foraging the

night before the experiment. Water was provided as needed. We ran high, low, mixed,

and control colonies concurrently in 4 different tents.

We used a familiar and novel feeder foraging assay to characterize colony level

foraging behavior[33] . We placed a feeder with 1M sucrose on Day 1, which remained

in the same location all week and became the 'familiar' feeder (Figure 2). We then

placed one novel feeder in different locations each day (Day 2 (X), Day 3 (Y), and Day 4

(Z)). Feeders had unique colors and unique odors and remained consistent throughout

the experiment (Table S2).

Mixed Colony Round Dance Preparation and Data Collection

To evaluate round dance behavior of each of the selected lines, we created 6

50/50 mixed colonies as detailed above. To induce foraging behavior, we placed the

colonies in a climate controlled indoor room for 10 days to allow bees to age which

increases foraging behavior. After 10 days, we then placed all bees from each colony

into a two-frame observation hive with glass walls. All comb surfaces were visible. We

video recorded round dance behavior using a Panasonic HC-WXF991K, starting the

recording 15 minutes before feeders were placed in the flight cage. For distances from

the colony entrance, see figure 4A. We followed the same feeder placement pattern

across 4 days, from Monday to Thursday, in Figure 4A. Round dance data was then

extracted visually from watching videos. We recorded the LI line of the dancer according

to the color marking on her thorax color, the feeder she visited according to the color

mark on her abdomen (which also distinguished her as having visited a feeder),

duration of the round dance, the LI line of the round dance followers, and the number of

turns in a dance during the first 20 s of the dance, or less if the dance ended before 20 s

elapsed. As the feeders were less than 12 m away from the colony, bees performed

round dances which lack distinct 'runs' and often have incomplete turns. Video watchers

were blind to the thorax and abdomen color associations between LI line and feeder

visitation, respectively.

Data Analysis Methods

To test whether bees exhibited a similar LI score as their parents regardless of

where they were housed after emergence, we used a generalized linear model. We

used LI score as the response variable, which fit a log-linear distribution, so we used a

gaussian family with a log link. Our fixed predictor variables were the line from which the

bees originated (high or low) and the colony type that they were placed in after

emergence (either their natal colony or a control colony).

To evaluate the effect of colony composition on colony-level foraging behavior to

novel and familiar feeders, we performed a general linear model with a gaussian error

distribution on number of visits, with line and feeder as fixed predictor variables, as well

as the interaction between line and feeder. We performed a generalized linear model

with a binomial error distribution with a logit link function on percent revisitation, as it

was a proportion comparing the number of revisits divided by the total number of visits.

Line and feeder were fixed predictor variables, as well as the interaction between the

line and feeder.

To explore whether the selected LI line of a forager bee influenced which feeder

it visited while in the mixed colony, we used a general linear model with a gaussian

error distribution on number of visits, with year, selected line and feeder as a fixed

predictor variables, as well as the interactions between these three. We did find a

significant three-way interaction between year, selected line, and feeder, which we

present in Table S6. Therefore, we treat years independently and performed two

different GLMs with selected line and feeder as our fixed predictor variables, as the

workers from the different years came from a new set of selected queens and drones,

colonies were in nucleus Langstroth hive boxes in 2017 but were placed in observation

colonies in 2018, as well as differences in weather.

To compare the round dance behavior among the selected lines, we examined

the effect of dancer selected line on the duration of the round dance, intensity of dancing,

number of turns by dancers, and number of followers of each dance using generalized

linear models. To analyze whether the duration of the round dance differed across the

learning lines, the duration of the round dance response variable fit a log-normal

distribution, so we used a generalized linear mixed model with a gaussian family and a

log link. The LI of the dancer was our fixed predictor variable. To evaluate which lines

attracted more dancers, we used a chi-square test to compare the proportion of dances

that attracted no followers across the lines. To evaluate whether there were differences

in the number of turns the dancers from each line performed, we used a linear mixed

model because the response variable - the number of turns per second, was normally

distributed. Finally, we used a negative binomial mixed regression model using the

package MASS[34] to understand how duration of a dance and the LI of the dancer

interacted to predict the number of followers. 159 dances out of 908 total dances had no

followers, requiring a zero-inflated model approach. We analyzed only bees that had

paint marks on their abdomens, ensuring that they had visited a feeder.

We used an Analysis of Deviance Wald chi-square test using the function Anova

in the MASS[34] package to further evaluate the overall effect of each fixed predictor

variable and interaction. We used the lme4 package[35] to perform these tests unless

otherwise noted. Post hoc tests were performed to determine the relationships between

the different levels of fixed predictor variables and their interactions using the package

emmeans[36]. We use R[37] for analysis.

REFERENCES

1. Ward, A.J.W., Herbert-Read, J.E., Sumpter, D.J.T., and Krause, J. (2011). Fast and
accurate decisions through collective vigilance in fish shoals. Proc. Natl. Acad. Sci.
108, 2312–2315.

2. Sasaki, T., and Biro, D. (2017). Cumulative culture can emerge from collective
intelligence in animal groups. Nat. Commun. *8*, 15049.

3. Couzin, I.D. (2009). Collective cognition in animal groups. Trends Cogn. Sci. *13*, 36–
43.

4. Sumpter, D.J. (2010). Collective animal behavior (Princeton University Press).

5. Camazine, S., Deneubourg, J.-L., Franks, N.R., Sneyd, J., Bonabeau, E., and
Theraula, G. (2003). Self-organization in biological systems (Princeton university
press).

6. Couzin, I.D., and Krause, J. (2003). Self-Organization and Collective Behavior in
Vertebrates. In Advances in the Study of Behavior (Elsevier), pp. 1–75. Available at:
https://linkinghub.elsevier.com/retrieve/pii/S0065345403010015 [Accessed July 31,
2019].

7. Feinerman, O., and Korman, A. (2017). Individual versus collective cognition in
social insects. J. Exp. Biol. *220*, 73–82.

8. Dornhaus, A., and Franks, N.R. (2008). Individual and collective cognition in ants
and other insects (Hymenoptera: Formicidae). Myrmecol. News *11*, 215–226.

9. Sumpter, D.J.T. (2006). The principles of collective animal behaviour. Philos. Trans.
R. Soc. B Biol. Sci. *361*, 5–22.

10. Chandra, S.B.C., Hosler, J.S., and Smith, B.H. (2000). Heritable variation for latent
inhibition and its correlation with reversal learning in honeybees (Apis mellifera). J.
Comp. Psychol. *114*, 86–97.

11. Mitchell, M.D., McCormick, M.I., Ferrari, M.C.O., and Chivers, D.P. (2011). Friend or
foe? The role of latent inhibition in predator and non-predator labelling by coral reef
fishes. Anim. Cogn. *14*, 707–714.

12. Ferrari, M.C.O., and Chivers, D.P. (2009). Latent inhibition of predator recognition by
embryonic amphibians. Biol. Lett. *5*, 160–162.

13. Lubow, R.E. (1973). Latent inhibition. Psychol. Bull. *79*, 398.

14. Lubow, R.E. (1997). Latent inhibition as a measure of learned inattention: some
problems and solutions. Behav. Brain Res. *88*, 75–83.

15. Lubow, R.E. (1965). Latent inhibition: Effects of frequency of nonreinforced pre-
exposure of the CS. J. Comp. Physiol. Psychol. *60*, 454–457.

16. Schachar, R., Logan, G.D., Robaey, P., Chen, S., Ickowicz, A., and Barr, C. (2007).
Restraint and Cancellation: Multiple Inhibition Deficits in Attention Deficit
Hyperactivity Disorder. J. Abnorm. Child Psychol. *35*, 229–238.

17. Abramson, C.I., and Bitterman, M.E. (1986). Latent inhibition in honeybees. Anim.
Learing Behav. *14*, 184–189.

18. Ferguson, H.J., Cobey, S., and Smith, B.H. (2001). Sensitivity to a change in reward
is heritable in the honeybee, Apis mellifera. Anim. Behav. *61*, 527–534.

19. Brandes, C., and Menzel, R. (1990). Common mechanisms in proboscis extension
conditioning and visual learning revealed by genetic selection in honeybees (Apis
mellifera). J. Comp. Physiol. A *166*, 545–552.

20. Ferrari, M.C.O., and Chivers, D.P. (2011). Learning about non-predators and safe
places: the forgotten elements of risk assessment. Anim. Cogn. *14*, 309–316.

21. Cook, C.N., Mosqueiro, T., Brent, C.S., Ozturk, C., Gadau, J., Pinter-Wollman, N.,
and Smith, B.H. (2018). Individual differences in learning and biogenic amine levels
influence the behavioural division between foraging honeybee scouts and recruits. J.
Anim. Ecol. *88*, 236–246.

22. Latshaw, J.S., and Smith, B.H. (2005). Heritable variation in learning performance
affects foraging preferences in the honey bee (Apis mellifera). Behav. Ecol.
Sociobiol. *58*, 200–207.

23. Waddington, K.D. (1982). Honey bee foraging profitability and round dance
correlates. J. Comp. Physiol. *148*, 297–301.

24. Pavlov, P.I. (1927). Conditioned reflexes: an investigation of the physiological
activity of the cerebral cortex. Ann. Neurosci. *17*, 136.

25. Lubow, R.E., Weiner, I., and Schnur, P. (1981). Conditioned Attention Theory. In
The Psychology of Learning and Motivation: Advances in Research and Theory.
(New York: Academic Press), pp. 1–49. Available at:
http://site.ebrary.com/id/10244511 [Accessed August 25, 2019].

26. Gershman, S.J., Blei, D.M., and Niv, Y. (2010). Context, learning, and extinction.
Psychol. Rev. *117*, 197–209.

27. Lubow, R.E. (1989). Latent inhibition and conditioned attention theory (Cambridge
University Press).

28. Mattila, H.R., Burke, K.M., and Seeley, T.D. (2008). Genetic diversity within
honeybee colonies increases signal production by waggle-dancing foragers. Proc.
R. Soc. B Biol. Sci. *275*, 809–816.

29. Dukas, R. (2002). Behavioural and ecological consequences of limited attention.
Philos. Trans. R. Soc. Lond. B. Biol. Sci. *357*, 1539–1547.

30. Hills, T.T., Todd, P.M., Lazer, D., Redish, A.D., and Couzin, I.D. (2015). Exploration
versus exploitation in space, mind, and society. Trends Cogn. Sci. *19*, 46–54.

31. Cobey, S.W., Tarpy, D.R., and Woyke, J. (2013). Standard methods for instrumental
insemination of *Apis mellifera* queens. J. Apic. Res. *52*, 1–18.

32. Harbo, J.R. (1986). Propagation and instrumental insemination. In Bee Breeding and
Genetics, (Academic Press, Inc), pp. 361–389.

33. Liang, Z.S., Nguyen, T., Mattila, H.R., Rodriguez-Zas, S.L., Seeley, T.D., and
Robinson, G.E. (2012). Molecular Determinants of Scouting Behavior in Honey
Bees. Science *335*, 1225–1228.

34. Ripley, B., Venables, B., Bates, D.M., Hornik, K., Gebhardt, A., Firth, D., and Ripley,
M.B. (2013). Package 'mass.' Cran R.

35. Bates, D., Maechler, M., Bolker, B., and Walker, S. (2014). lme4: Linear mixed-
effects models using Eigen and S4. R Package Version *1*, 1–23.

36. Lenth, R., and Lenth, M.R. (2018). Package 'lsmeans.' Am. Stat. *34*, 216–221.

37. Team, R.C. (2018). R: A language and environment for statistical computing.

SUPPMEMENTARY MATERIALS - Individual Learning Phenotypes Drive Collective

Cognition

Chelsea N. Cook[1*], Natalie J. Lemanski[2], Thiago Mosqueiro[2], Jürgen Gadau[3], Cahit Ozturk[1],

Noa Pinter-Wollman[2†], Brian H. Smith[1†]

Affiliations

[1] School of Life Sciences, Arizona State University, Tempe, USA

[2] Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los

Angeles, USA

[3] Institute for Evolution and Biodiversity, University of Münster, Münster, Germany

[†]Co-Senior Authors

[*]Corresponding Author, cncook1@asu.edu

	Colony Type

	High	Low	Mix	Control
High LI	650	0	325	0
Low LI	0	650	325	0
Control	650	650	650	1300
Totals	1300	1300	1300	1300

700   Table S1: **The number of honey bees in each experimental colony by genetic line.** Each of

701   the 4 created colonies were set up in this way each week. We counted and marked the thorax

702   each bee from the learning lines, and counted but did not mark supplemental control bees.

Day	Feeder Treatment	Odor Added to Feeder	Color of Feeder
Day 1	Familiar	Hexanol	Red
Day 2	Familiar + X	Hexanol + Octanone	Red + Blue
Day 3	Familiar + Y	Hexanol + Geraniol	Red + Pink
Day 4	Familiar + Z	Hexanol + Citranol	Red + Orange

705 **Table S2: The weekly routine of feeder characteristics and placement.** Each feeder had 1M

706 sucrose solution. Color, odor, and location respectively varied by feeder. The treatment

707 sequence was the same each week.

714 **Figure S1: The cumulative visitation to all feeders over time, averaged across days.** The

715 thick colored line is the average, and the gray stepwise lines are visitation on a single day by a

716 single colony. Colored lines are the same data shown in Figure 2B.

contrast	estimate	SE	df	z.ratio	p.value
control - high	0.082	0.084	Inf	0.969	0.767
control - low	-0.065	0.099	Inf	-0.659	0.912
control - mix	0.293	0.088	Inf	3.336	0.005
high - low	-0.147	0.082	Inf	-1.798	0.274

high - mix	0.212	0.068	Inf	3.103	0.010
low - mix	0.358	0.085	Inf	4.198	0.000

721 Table S3: A table of the pairwise post hoc tests of how LI line predicts percent revisitation to all

722 feeders, referenced in figure 2C.

contrast	estimate	SE	df	z.ratio	p.value
constant,control - novel,control	0.163	0.080	Inf	2.031	0.461
constant,control - constant,high	-1.363	0.061	Inf	-22.431	0.000
constant,control - novel,high	-0.249	0.072	Inf	-3.437	0.014
constant,control - constant,low	-0.214	0.073	Inf	-2.931	0.067
constant,control - novel,low	-0.298	0.072	Inf	-4.162	0.001
constant,control - constant,mix	-1.467	0.060	Inf	-24.393	0.000
constant,control - novel,mix	0.219	0.081	Inf	2.701	0.122
novel,control - constant,high	-1.526	0.065	Inf	-23.508	0.000
novel,control - novel,high	-0.411	0.076	Inf	-5.421	0.000
novel,control - constant,low	-0.376	0.076	Inf	-4.925	0.000
novel,control - novel,low	-0.460	0.075	Inf	-6.129	0.000
novel,control - constant,mix	-1.630	0.064	Inf	-25.338	0.000
novel,control - novel,mix	0.057	0.084	Inf	0.675	0.998

constant,high - novel,high	1.115	0.055	Inf	20.194	0.000
constant,high - constant,low	1.150	0.056	Inf	20.555	0.000
constant,high - novel,low	1.065	0.054	Inf	19.661	0.000
constant,high - constant,mix	-0.104	0.038	Inf	-2.756	0.106
constant,high - novel,mix	1.583	0.066	Inf	23.818	0.000
novel,high - constant,low	0.035	0.068	Inf	0.512	1.000
novel,high - novel,low	-0.049	0.067	Inf	-0.736	0.996
novel,high - constant,mix	-1.219	0.055	Inf	-22.358	0.000
novel,high - novel,mix	0.468	0.077	Inf	6.066	0.000
constant,low - novel,low	-0.084	0.068	Inf	-1.248	0.918
constant,low - constant,mix	-1.254	0.055	Inf	-22.690	0.000
constant,low - novel,mix	0.433	0.078	Inf	5.574	0.000
novel,low - constant,mix	-1.170	0.053	Inf	-21.864	0.000
novel,low - novel,mix	0.517	0.076	Inf	6.767	0.000
constant,mix - novel,mix	1.687	0.066	Inf	25.604	0.000

725   Table S4: A table of the pairwise post hoc tests of how the Line*Feeder interaction predicts

726   number of visits, which corresponds to letters in figure 2D.

contrast	estimate	SE	df	z.ratio	p.value
control,constant - high,constant	0.251	0.127	Inf	1.975	0.499

control,constant - low,constant	-0.150	0.151	Inf	-0.996	0.975
control,constant - mix,constant	0.523	0.132	Inf	3.958	0.002
control,constant - control,novel	-0.972	0.147	Inf	-6.635	0.000
control,constant - high,novel	-1.256	0.127	Inf	-9.913	0.000
control,constant - low,novel	-0.881	0.141	Inf	-6.231	0.000
control,constant - mix,novel	-1.394	0.135	Inf	-10.311	0.000
high,constant - low,constant	-0.402	0.125	Inf	-3.208	0.029
high,constant - mix,constant	0.272	0.102	Inf	2.669	0.132
high,constant - control,novel	-1.223	0.120	Inf	-10.209	0.000
high,constant - high,novel	-1.507	0.095	Inf	-15.937	0.000
high,constant - low,novel	-1.133	0.114	Inf	-9.972	0.000
high,constant - mix,novel	-1.645	0.106	Inf	-15.566	0.000
low,constant - mix,constant	0.673	0.130	Inf	5.174	0.000
low,constant - control,novel	-0.822	0.145	Inf	-5.679	0.000
low,constant - high,novel	-1.105	0.125	Inf	-8.873	0.000
low,constant - low,novel	-0.731	0.140	Inf	-5.238	0.000
low,constant - mix,novel	-1.244	0.133	Inf	-9.335	0.000
mix,constant - control,novel	-1.495	0.125	Inf	-11.964	0.000
mix,constant - high,novel	-1.778	0.101	Inf	-17.615	0.000
mix,constant - low,novel	-1.404	0.119	Inf	-11.803	0.000

mix,constant - mix,novel	-1.917	0.111	Inf	-17.197	0.000
control,novel - high,novel	-0.284	0.119	Inf	-2.380	0.251
control,novel - low,novel	0.091	0.135	Inf	0.673	0.998
control,novel - mix,novel	-0.422	0.128	Inf	-3.291	0.022
high,novel - low,novel	0.374	0.113	Inf	3.316	0.021
high,novel - mix,novel	-0.138	0.105	Inf	-1.317	0.893
low,novel - mix,novel	-0.513	0.122	Inf	-4.189	0.001

729   Table S5: A table of the pairwise post hoc tests of how the Line*Feeder interaction predicts

730   percent revisitation, which corresponds to letters in figure 2E.

	LR Chisq	Df	Pr(>Chisq)
BeeType	100.073	2.000	0.000
Feeder	2196.156	1.000	0.000
Year	479.995	1.000	0.000
BeeType:Feeder	47.718	2.000	0.000
BeeType:Year	498.209	2.000	0.000
Feeder:Year	61.341	1.000	0.000
BeeType:Feeder:Year	31.492	2.000	0.000

Table S6: **Individual visitation by bee type differed across two experimental years.** GLM

results showing the three-way interaction between year and the type of bee visiting a feeder

(Figure 3). There is likely a difference in year because of several reasons, including 1) Colonies

were selected from different queens from different breeders in 2017 and 2018 and climactic

conditions were different in 2017 compared to 2018 even though experiments were done in the

same time frame (June-July in 2017, June in 2018).

contrast	estimate	SE	df	z.ratio	p.value
control,familiar - high,familiar	-0.220	0.054	Inf	-4.102	0.001
control,familiar - low,familiar	0.624	0.068	Inf	9.210	0.000
control,familiar - control,novel	1.387	0.076	Inf	18.320	0.000
control,familiar - high,novel	1.425	0.077	Inf	18.564	0.000
control,familiar - low,novel	1.986	0.095	Inf	20.794	0.000
high,familiar - low,familiar	0.844	0.065	Inf	12.916	0.000
high,familiar - control,novel	1.607	0.074	Inf	21.841	0.000
high,familiar - high,novel	1.645	0.075	Inf	22.033	0.000
high,familiar - low,novel	2.206	0.094	Inf	23.512	0.000
low,familiar - control,novel	0.763	0.084	Inf	9.049	0.000
low,familiar - high,novel	0.801	0.085	Inf	9.394	0.000
low,familiar - low,novel	1.362	0.102	Inf	13.290	0.000
control,novel - high,novel	0.038	0.092	Inf	0.413	0.998
control,novel - low,novel	0.599	0.108	Inf	5.546	0.000

high,novel - low,novel	0.561	0.109	Inf	5.159	0.000

743   Table S7: A table of the pairwise GLM contrasts of how the Line*Feeder interaction predicts

744   number of visits by each line in the mixed colonies in 2017, which corresponds to letters in

745   figure 3A.

contrast	estimate	SE	df	z.ratio	p.value
control,familiar - high,familiar	0.389	0.043	Inf	8.949	0.000
control,familiar - low,familiar	-0.227	0.037	Inf	-6.117	0.000
control,familiar - control,novel	0.567	0.041	Inf	13.904	0.000
control,familiar - high,novel	1.451	0.054	Inf	26.775	0.000
control,familiar - low,novel	0.827	0.044	Inf	18.827	0.000
high,familiar - low,familiar	-0.616	0.042	Inf	-14.779	0.000
high,familiar - control,novel	0.178	0.045	Inf	3.947	0.001
high,familiar - high,novel	1.062	0.057	Inf	18.485	0.000
high,familiar - low,novel	0.438	0.048	Inf	9.140	0.000
low,familiar - control,novel	0.793	0.039	Inf	20.441	0.000
low,familiar - high,novel	1.678	0.053	Inf	31.807	0.000

low,familiar - low,novel	1.053	0.042	Inf	25.015	0.000
control,novel - high,novel	0.884	0.055	Inf	15.959	0.000
control,novel - low,novel	0.260	0.045	Inf	5.725	0.000
high,novel - low,novel	-0.625	0.058	Inf	-10.810	0.000

Table S8: A table of the pairwise GLM contrasts of how the Line*Feeder interaction predicts number of visits by each line in the mixed colonies in 2018, which corresponds to letters in figure 3B.