1

Heritable Learning Phenotypes Drive Collective Cognition

Chelsea N. Cook[1*], Natalie J. Lemanski[2], Thiago Mosqueiro[2], Jürgen Gadau[3], Cahit
Ozturk[1], Noa Pinter-Wollman[2◊], Brian H. Smith[1◊]

Affiliations

[1] School of Life Sciences, Arizona State University, Tempe, USA

[2] Department of Ecology and Evolutionary Biology, University of California Los Angeles,
Los Angeles, USA

[3] Institute for Evolution and Biodiversity, University of Münster, Münster, Germany

[◊]Co-Senior Authors

[*]Corresponding Author

ABSTRACT

      Collective cognition allows animal groups to accomplish many tasks that could not be accomplished as effectively alone, such as schools of fish avoiding predators[1], flocks of birds moving thousands of miles across the Earth[2], and honey bee colonies collecting food from millions of flowers[3]. Individuals in groups utilize local information to quickly adjust to ecological changes by implicitly or explicitly communicating information with group members to form collective behavior[4–6]. However, individuals vary in their cognitive abilities, which influences the information each individual pays attention to or shares, thus influencing collective responses[7–9]. Here, we show that individual differences in learning scales to shape collective foraging behavior in honey bees by utilizing a naturally variable and heritable learning behavior called latent inhibition (LI)[10]. We artificially selected two distinct phenotypes: high LI bees that are better at ignoring previously unrewarding familiar stimuli, and low LI bees that learn previously unrewarding and novel stimuli equally well. Colonies comprised of high LI individuals preferred to visit the familiar food location, while low LI colonies visit novel and familiar food locations equally. However, in colonies of mixed learning phenotypes, the low LI bees showed a preference to visiting familiar feeders, which contrasts with their behavior when in a uniform group. We show that the shift in the feeder preference of low LI bees in uniform low LI versus mixed colonies is driven by foragers of the high LI phenotype dancing more intensely and attracting more followers. These results reveal that variation in individual learning phenotypes contributes to collective behavior in social animals.

INTRODUCTION

      To uncover how individual variation in learning impacts group-level behaviors, we investigated how an individual's latent inhibition (LI) may scale to influence collective foraging behavior in honey bee colonies. Although LI has been mostly studied in vertebrates [11–15], honey bees (*Apis meillifera*) also exhibit[16,17] and show variation in LI[10]. Despite several studies showing heritable variation in latent inhibition[10,18], and how it is relevant in individual ecological contexts such as predator avoidance[11,12,19], it remains

unclear as to whether or how this variation functions in ecologically relevant decisions in

complex social environments. Foraging honey bees vary in their expression of LI;

scouts tend to exhibit high LI and ignore familiar odors, while recruits who tend to exhibit

low LI tend to learn familiar and novel odors equally well[20]. LI is heritable in honey

bees[17]. Therefore, we can select queens and drones (haploid males) for expression of

LI and create selected lines from singly inseminated queens with like performing drones

to produce two distinct lines of workers that exhibit similar LI to their parents[21]. We

created 24 colonies composed of single cohorts of only low, only high, 50/50 mixed high

and low LI workers, as well as age-matched non-selected control bees. In semi-natural

foraging conditions, we counted the number of forager visits, first visits, and revisits to

the familiar or novel food locations. To explore the mechanisms underlying how

individual variation in LI affects collective foraging, we quantified the round recruitment

dance activity in 6 mixed colonies while the colonies visited novel and familiar feeders.

These field assays allowed us to simultaneously quantify individual and colony level

foraging behaviors.

RESULTS

To ensure workers in different social environments exhibited the predicted

heritable LI phenotype, we evaluated the LI score of foragers after 21 days in either

their natal colony or a control colony. We marked 1000 individuals from each selected

line (high or low LI) on the day of emergence. We then placed 500 individuals back into

their natal colony and 500 individuals into an established control colony of equal size

with an open mated queen, i.e. workers with a variety of learning phenotypes. We

monitored the colonies until marked bees began to make foraging flights (~21 days). We

then collected marked foragers as they returned to the colony and brought them into the

laboratory to evaluate their LI. We avoided pollen foragers as they tend to exhibit

different learning behavior compared to nectar foragers[21]. We found that foragers

retained the expected LI based on the LI of their parents, regardless of whether they

were housed with same or with variable learning phenotypes. Foragers from the high

and low lines differed in expression of LI as expected (GLM: χ^2 = 4.84, df=1, p=0.027,

Figure 1). We did not detect an effect of the identity of the colony in which the bees

were housed on LI phenotype (χ^2 = 3.28, df=2, p=0.193, Figure 1).

96
97

Figure 1: Social environment does not alter expression of genetically selected latent inhibition. LI scores of individuals from high LI lines that spent their adult life either in high LI only colonies (red, n=36) or in a control colony with a variety of LI phenotypes from an open mated queen (red with gray vertical lines, n=18); individuals from low LI lines that spent their adult life either in low LI only colonies (yellow, n=52) or in control colonies (yellow with gray vertical lines, n=10). In this and subsequent figures, the large black dot is the mean, the white box is the interquartile range (IQR), whiskers extend to 1.5*IQR, and the small points beyond the whiskers are outliers. Shaded areas show the distribution of the data. Here, and in all following figures, yellow are low LI colonies and individuals, gray are control colonies and individuals, and red are high LI colonies and individuals. Here and in subsequent

109 figures, darker hues indicate individual-level behavior and lighter hues indicate
110 colony-level behavior.
111
112
113
To determine how the learning phenotypes influenced colony-level foraging
behavior, we placed small single-cohort (same age bees) colonies into a flight cage and
monitored foraging activity. We evaluated 4 colony types each week: one control colony
consisting of approximately 1300 age-matched bees from open mated queens; one
colony consisting of 650 workers from high LI queens plus 650 age-matched control
bees; one colony consisting of 650 workers from low LI queens plus 650 aged-matched
control bees; and one 50/50 mixed colony with 325 workers from each LI line plus 650
aged-matched control bees. In the last 3 types, the supplemented 650 age-matched
bees from open mated queens were used to ensure a small but functioning colony as
we did not have enough workers from the single-drone-inseminated queens and
colonies of just 650 individuals would be too weak to forage. Honey bee division of labor
is largely influenced by worker age, so we used age-matched bees to remove any
influence that age may have on foraging propensity. On day 1, we trained bees to a
feeder inside the tent containing 1M sucrose and an odor, which became the 'familiar'
feeder. During the subsequent 3 days, in addition to the familiar feeder, we introduced a
single novel feeder each day with a different odor and color, but with the same sugar
concentration as the familiar feeder (Figure 2A). To evaluate the collective ability of the
colony to find a new feeder, we recorded the number of visits to each feeder by bees
from each selected line according to the color of paint on the bees' thorax. We further
marked bees with a feeder-specific color on their abdomen when they visited the feeder
for the first time to determine if bees revisited that feeder. We repeated this for 6 weeks
on 6 colonies for each group type.
Colony composition strongly influenced overall number of visits to the food
locations (N = 6 colonies in each line, 24 total, 6172 total visits; GLM: χ^2 = 1270, df = 3,
p < 0.0001, Figure 2B). High LI colonies had significantly more visits to all food locations
compared to low LI colonies (GLM: Z=25.5, p <0.0001, Figure 2A), mixed colonies
(Tukey post hoc: Z=5.18, p<0.0001), and controls (Z=26.6, p<0.0001). Mixed LI
colonies also had significantly more visits compared to low (Z=20.7, p < 0.0001) and

controls (Z=21.8, p<0.0001). Low LI and control colonies had the fewest total visits and
were not significantly different from each other (Z=-1.38, p=0.50).

Foraging in the high, low, and control colonies was largely performed by bees
revisiting the feeders. (GLM, χ^2 = 22.32, df =3, p<0.0001, Figure 2C). However, the
mixed LI colonies had a significantly lower proportion of revisiting foragers compared to
the low (Tukey post hoc: Z=-4.2, p=0.0002), high (Z=-3.1, p=0.01), and control colonies
(Z=-3.33, p=0.004). We did not detect significant differences among the other colony
types (Supplemental Table 3).

A colony's LI phenotype composition determined its preference between the
novel and familiar feeders (GLM: Feeder*Colony χ^2 =473.64, df=3, p<0.0001; Figure
2D). High and mixed colonies preferred the familiar feeder over the novel one (Tukey
Poshoc: High Familiar:Novel: Z=20.2, p<0.0001; Mixed Familiar:Novel: Z=25.6,
p<0.0001). Low LI and control colonies did not show a strong preference for either
feeder, visiting them equally (Low Familiar:Novel: Z=-1.24, p=0.92; Control
Familiar:Novel: Z=2.03, p=0.46).

The number of re-visits to the novel and familiar feeders was different across
colony compositions (Figure 2E: Colony*Feeder χ^2 =53.67, p<0.0001). All colonies had
a higher proportion of re-visits to the familiar feeder compared to the novel feeder.
However, the mixed LI colonies had a much lower proportion of re-visitation to the novel
feeders than the other colony types (Supplemental Table 4). Thus, new foragers in the
mixed colonies that visited the novel feeder were less likely to return to it compared to
foragers who visited the novel feeders in other colonies.

Figure 2: Colonies constructed from different genetic lines selected for high or low latent inhibition exhibited differences in collective foraging behavior. (A) The experimental set up illustrating the location of feeders in relation to the location of the colony (center, white) within the experimental arena (large rectangle). The familiar feeder (red) was provided on day 1 and on all subsequent days. Novel feeder X (blue) was presented on day 2, novel feeder Y (purple) on day 3, and novel feeder Z (orange) on day 4. See supplemental table 2 for associated odors. Visits to all novel feeders were combined for statistical analysis. (B) Cumulative number of visits of bees to all feeders over time by colony type. Different letters to the right of the lines indicate statistically significant differences according to a post hoc Tukey test (For further

illustration of visitation by each colony on each day, see supplemental figure 1). (C) Percent of re-visits out of the total number of visits to all feeders by colony type. Here and in all following panels, different letters above boxes indicate statistically significant differences according to a post hoc Tukey test. (D) Number of all visits to the familiar feeder (solid boxes) and a novel feeder (hatched boxes) for each type of colony, when both novel and familiar feeders were presented simultaneously (days 2-4). (E) Percent of re-visits out of the total number of visits to either the familiar or the novel feeder by type of colony when both novel and familiar feeders were presented simultaneously (days 2-4). In C, D and E, horizontal lines are the median, the boxes are the interquartile range (IQR), whiskers extend to 1.5*IQR, and the small points beyond the whiskers are outliers. N=24 colonies, 6 colonies per group type, 6172 total visits.

To determine why the mixed colonies showed a preference for the familiar feeder (Figure 2D), we examined how individual lines visited each feeder (Figure 3). In 2017, we tested mixed colonies placed in a flight cage. In 2018, we reselected lines and then placed mixed colonies into two-frame observation hives to evaluate recruitment dances along with visitation to the feeders in the flight cages. We found that there was a significant year effect (Supplemental Table 5), likely due to reselection and different environmental conditions. We therefore statistically analyzed each year separately to focus on the within-year variation between the selected lines.

Low LI and control individuals shift their preference to the familiar feeder when mixed with high LI bees. In 2017, we found a significant interaction between the selected line and which feeder foragers visited (GLM: χ^2=7.79, df=2, p=0.02; Figure 3A). Although low LI and control colonies did not show a preference to a novel or familiar feeder when they had a uniform colony composition (Figure 2E), when mixed with high LI individuals, low LI and control individuals exhibited a preference to the familiar feeder (GLM: Low Familiar:Novel: Z=13.28, p<0.0001; Control Familiar:Novel: Z=18.32, p<0.0001; Figure 3A). High LI individuals showed a preference to familiar feeders (GLM: Familiar:Novel: Z=22.03, p<0.0001) just as colonies comprised of only high LI individuals did (Figure 2E). We found a significant interaction between selected line and feeder in 2018 (GLM: χ^2=85.27, df=2, p<0.0001; Figure 3B), with low LI and control individuals showing preference to the familiar feeder over the novel feeder (GLM: Low Familiar:Novel: Z=25.05, p<0.0001; Control Familiar:Novel: Z=13.90,

209 p<0.0001; Figure 3B) similar to high LI individuals preferring the familiar feeders

210 (Familiar:Novel: Z=18.48, p<0.0001).

212
213
214
215

216
217
218
Figure 3: Visits of individuals from different genetically selected lines when in a
mixed colony. Daily visits to the familiar (solid) and novel (hatched) feeders by
individual bees in mixed colonies from low LI parents (yellow), high LI parents (red) or
open mated queens (grey) in (A) 2017, N=6 mixed colonies, 2347 overall visits and (B)
in mixed colonies from lines that were re-selected in 2018, N=6 colonies, 6272 overall
visits. The horizontal line in the box is the median, the box is 25-75% of the data,
whiskers represent 95% of the data, and diamonds show outliers beyond 95%. Different
letters above boxes indicate statistically significant differences according to a post hoc
Tukey test.

To uncover the behavioral mechanisms that underlie the switching of low LI individuals from having no feeder preference when in a uniform colony composition to preferring the familiar feeder when in a mixed colony, we examined the round dance, the modified waggle dance used at short distances[22], of individuals from each selected line in mixed colonies as they returned from foraging. Using observation hives with glass walls, we video recorded bees performing the round dance to recruit other individuals in the colony to forage. To determine which selected line recruited to each feeder, we noted the selected line of the dancer (high or low LI) according to the paint marks on the individuals' thorax and whether the dancer had visited a feeder according to the paint marks on abdomens. We did not record dancers without abdominal marks as they were likely collecting from and recruiting to unmonitored water sources. To determine who the information about a feeder was communicated to, we counted the number of followers of each dancer and the selected line of the followers. To quantify the dance intensity, we recorded the duration of the dance, and the number of turns the dancer made during the first 20 seconds of the dance.

Individuals from the lines differed in their likelihood to perform a round dance (Chi-square test: χ^2=26.61, df=2, p<0.0001; Figure 4B). Low LI individuals were significantly more likely to perform a dance compared to high Li individuals (pairwise chi-square test: p=0.0001) and controls (pairwise chi-squared test: p=0.004). High LI individuals were just as likely to perform a dance as controls (p=0.36). Individuals differed in their likelihood to follow a dance based on their selected line (Chi-square test: χ^2=28.26, df=2, p<0.0001; Figure 4B). Low LI individuals were significantly more likely to follow a dance compared to high LI bees (pairwise chi-square test: p<0.0001) and controls (pairwise chi-square test: p<0.003). High LI and control individuals were equally likely to follow a dance (pairwise chi-square test: p=0.240).

Although the high LI individuals danced less often, high LI dances had significantly more followers compared to low and control bees (Chi-square test: χ^2= 13.93, df=2, p<0.001; Figure 4C). Low LI bees performed more dances that had no followers compared to high LI and control dances. We did not detect a statistically significant difference between the proportion of individuals from each line that followed

each line of dancer (Chi-square test: χ^2= 7.05, df = 4, p= 0.13, Figure 4D). Low LI

individuals spent more time dancing; however they attracted fewer followers than high

and control dancers, indicated by the significant interaction between the LI of the dancer

and dance duration when predicting the number of followers (GLMM: χ^2= 6.42, df=2,

p=0.04; Figure 4E).

The relative attraction of dances of high LI bees could be due to the intensity of

the dance. High LI bees performed more turns per second during their dances (ANOVA:

χ^2=12.8, df=2, p=0.002; Figure 4F). High LI dancers performed an average of 0.59 turns

per second, significantly higher than low LI dancers, who performed an average of 0.52

turns per second (Tukey: t=-3.13, p=0.005). Control bees also performed more turns per

second than low LI bees (Tukey: t=-2.5, p=0.03), but not different than high LI bees, at

an average 0.62 turns per second (Tukey: t=-0.7, p=0.7).

Figure 4: Recruitment dances facilitate integration of information from different genetically selected lines. (A) The experimental set up illustrating the location of feeders in relation to the location of the colony entrance (top right, white) within the experimental arena (large rectangle). The familiar feeder (red) was provided on day 1

and on all subsequent days. Novel feeder X (blue) was presented on day 2, novel
feeder Y (purple) on day 3, and novel feeder Z (orange) on day 4. See supplemental
table 2 for associated odors. Visitation to novel feeders were combined for statistical
analysis. (B) Proportion of dances (N=667) or follows (N=1201) across 6 colonies
performed by bees from each line, relative to their abundance in the mixed colony (350
high, 350 low, 700 control). We accounted for the difference in abundance of each
selected line by dividing the number of observed control dancers by 2 before calculating
these proportions. (C) Proportion of dances performed per LI line type that were either
followed by at least one individual (colored) or not followed by any other bees (black).
(D) Proportion of dances by LI line type that were followed (from panel B) broken down
by LI of the follower. (E) Relationship between number of followers and duration of a
dance by line. Point and line colors indicate LI of dancer. Best fit line represents the
GLM, shaded area represents the 95% confidence interval. (F) Rate of turns per second
in a dance by line. The large black dot in the box is the mean, the box is 25-75% of the
data, whiskers represent 95% of the data. The violin shapes illustrate distribution of the
data. Different letters above violins indicate statistically significant differences according
to a post hoc Tukey test.
295
296

DISCUSSION

By combining techniques from experimental psychology and behavioral ecology,

we have developed a system for investigating how variation in individual learning

behavior drives collective cognition. We utilized this system to demonstrate that a

laboratory-selected heritable learning behavior with natural individual variation scales to

shape the collective performance of a honey bee colony on foraging tasks. In the lab,

high LI honey bees learn to ignore familiar odors that they experienced without

reinforcement, while readily learning novel odors. When a stimulus is rewarding, high LI

bees exhibit increased attention to that information. One interpretation of reduced

learning to a familiar, unrewarding, stimulus is that pre-exposure reduces attention to,

and thus associability of, that stimulus. This interpretation is an extension of conditioned

attention theory[23,24], which proposes that latent inhibition is induced by allowing animals

to focus their attention on important information[14,25,26]. Our observations of field

behavior of low and high LI individuals and colonies are consistent with this

interpretation, whereby high LI individuals have stronger attention capacities to food

compared to low LI individuals. Once high LI individuals have found a food location,

they continue to revisit it, 'attending' more strongly to reinforced feeders over new ones.

The increased impact of the resource on these bees could translate into stronger, more

vigorous dances. In contrast, low LI individuals learn and visit both known and new

feeders equally, dividing their attention across resources and acting more like generalist

foragers. In mixed colonies, this broadened attention by low LI individuals may therefore

make them the perfect audience for the high LI dancers, driving them to prefer feeders

that high LI individuals preferentially visit. In fact, individual phenotypes that emerge

from these lines in natural conditions seem to be collectively complementary "finders"

and "refiners", where high LI bees discover food and direct other bees to that location,

while the low LI bees refine that information, finding the most rewarding resource by

sampling a broader spectrum of available resources (Lemanski et al. In Review). Under

natural conditions, where queens mate with many different drones, most colonies would

possess both types of learners, perhaps more closely resembling our mixed colonies[27].

We propose that this diversity of 'attention' aspect of individual cognitive phenotypes

may enhance the overall efficacy with which a colony finds and exploits resources[28]. In

summary, our work indicates that individual cognition scales to shape the collective

cognition of animals solving critical ecologically relevant tasks.

METHODS

Obtaining queens and drones

To obtain queens for producing selected lines of a specific LI behavior, we performed the LI assay as outlined in[10,20] on mature virgin queens 10 days after emergence. Briefly, we familiarized bees to an odor by puffing it at them 40 times every 5 minutes, then used the proboscis extension reflex to test their ability to learn to associate a food reward to the familiar versus a novel odor. Tested queens were placed into individually labelled queen cages and returned to a queenless colony until insemination, which typically occurred within a week of testing. To obtain fertile drones, we collected them from their returning unsuccessful mating flights at the entrance of colonies. We placed them into cages overnight in a queenless colony for LI testing the next day. After testing, drones were marked for individual identification and placed into a cage and placed into a queen bank for no longer than 3 days until inseminations occurred.

Queen Inseminations

We used instrumental insemination to inseminate a queen with sperm from a single drone. We inseminated a high LI queen with a high LI drone, and a low LI queen with a low LI drone[29,30]. LI varies across individuals. However, for this behavioral selection, we used the highest and lowest LI scoring individuals to create the high and low colonies, respectively. We then introduced queens to small queenless colonies, then allowed the queens to produce workers for 1 month. Colonies were checked weekly to eliminate the possibility of supersedure.

Cohoused Worker Preparation and Testing

To test the LI of foragers from each LI line, we placed frames of capped pupae from 3 high and 3 low LI colonies into 34°C incubators for 18 hours. After 18 hours, we used water based acrylic paint pens (Montana brand) to mark the abdomens of the eclosed bees with a color indicating their natal colony. Half of the bees were then returned to their natal colony and half were placed into an established control colony of an open-mated queen. Fewer bees were recovered from the established colony as

many are recognized as non-nestmates and rejected. After 2 weeks, colonies were

monitored every day until marked bees began to forage, ~21 days after emergence.

Returning nectar foragers were collected and tested for LI.

Field Colony Experimental Setup

To explore the colony-level foraging behavior of the LI lines, we set up 4

treatment colonies for each of the colony types: a high LI colony, a low LI colony, a

50/50 mixed colony, and a control. We ran weekly field experiments for 6 weeks. For

ease of identification, we always marked individuals from high LI colonies red, orange,

and pink, and individuals from low LI colonies green, blue, yellow, and white. We

continued to mark emerging bees from the same frames until we had 650 bees to form

a colony, which took typically 2-3 days. To achieve relatively normal conditions for

typical honey bee behavior, we supplemented workers from an unselected colony

(control bees), who were not marked. For colony set up, see Supplemental Table 1.

Bees were then placed into 4 different treatment colonies consisting of ~1300 bees:

high plus controls, low plus controls, 50/50 mixed high/low plus controls, and only

control colony. Bees were provided a honeycomb and remained inside for 5 days before

being placed for field experimentation. We then placed nucleus colonies into outdoor

flight cages (3.05m x 12m) and replaced the honeycomb frame with an empty frame to

induce foraging the night before the experiment. Water was provided as needed. We

ran high, low, mixed, and control colonies concurrently in 4 different tents.

We used a familiar and novel feeder foraging assay to characterize colony level

foraging behavior[31] . We placed a feeder with 1M sucrose on Day 1, which remained in

the same location all week and became the 'familiar' feeder (Figure 2). We then placed

one novel feeder in different locations each day (Day 2 (X), Day 3 (Y), and Day 4 (Z)).

Feeders had unique colors and unique odors and remained consistent throughout the

experiment (Supplemental Table 2).

Mixed Colony Round Dance Preparation and Data Collection

To evaluate round dance behavior of each of the selected lines, we created 6

50/50 mixed colonies as detailed above. To induce foraging behavior, we placed the

colonies in a climate controlled indoor room for 10 days to allow bees to age which

increases foraging behavior. After 10 days, we then placed all bees from each colony

into a two-frame observation hive with glass walls. All comb surfaces were visible. We

video recorded round dance behavior using a Panasonic HC-WXF991K, starting the

recording 15 minutes before feeders were placed in the flight cage. For distances from

the colony entrance, see figure 4A. We followed the same feeder placement pattern

across 4 days, from Monday to Thursday, in Figure 4A. Round dance data was then

extracted visually from watching videos. We recorded the LI line of the dancer according

to the color marking on her thorax color, the feeder she visited according to the color

mark on her abdomen (which also distinguished her as having visited a feeder),

duration of the round dance, the LI line of the round dance followers, and the number of

turns in a dance during the first 20 s of the dance, or less if the dance ended before 20 s

elapsed. As the feeders were less than 12 m away from the colony, bees performed

round dances which lack distinct 'runs' and often have incomplete turns. Video watchers

were blind to the thorax and abdomen color associations between LI line and feeder

visitation, respectively.

Data Analysis Methods

To test whether bees exhibited a similar LI score as their parents regardless of

where they were housed after emergence, we used a generalized linear model. We

used LI score as the response variable, which fit a log-linear distribution, so we used a

gaussian family with a log link. Our fixed predictor variables were the line from which the

bees originated (high or low) and the colony type that they were placed in after

emergence (either their natal colony or a control colony).

To evaluate the effect of colony composition on colony-level foraging behavior to

novel and familiar feeders, we performed a general linear model with a gaussian error

distribution on number of visits, with line and feeder as fixed predictor variables, as well

as the interaction between line and feeder. We performed a generalized linear model

with a binomial error distribution with a logit link function on percent revisitation, as it

was a proportion comparing the number of revisits divided by the total number of visits.

Line and feeder were fixed predictor variables, as well as the interaction between the

line and feeder.

To explore whether the selected LI line of a forager bee influenced which feeder

it visited while in the mixed colony, we used a general linear model with a gaussian

error distribution on number of visits, with year, selected line and feeder as a fixed

predictor variables, as well as the interactions between these three. We did find a

significant three-way interaction between year, selected line, and feeder, which we

present in Supplemental Table 5. Therefore, we treat years independently and

performed two different GLMs with selected line and feeder as our fixed predictor

variables, as the workers from the different years came from a new set of selected

queens and drones, colonies were in nucleus Langstroth hive boxes in 2017 but were

placed in observation colonies in 2018, as well as differences in weather.

To compare the round dance behavior among the selected lines, we examined

the effect of dancer selected line on the duration of the round dance, intensity of

dancing, number of turns by dancers, and number of followers of each dance using

generalized linear models. To analyze whether the duration of the round dance differed

across the learning lines, the duration of the round dance response variable fit a log-

normal distribution, so we used a generalized linear mixed model with a gaussian family

and a log link. The LI of the dancer was our fixed predictor variable. To evaluate which

lines attracted more dancers, we used a chi-square test to compare the proportion of

dances that attracted no followers across the lines. To evaluate whether there were

differences in the number of turns the dancers from each line performed, we used a

linear mixed model because the response variable - the number of turns per second,

was normally distributed. Finally, we used a negative binomial mixed regression model

using the package MASS[32] to understand how duration of a dance and the LI of the

dancer interacted to predict the number of followers. 159 dances out of 908 total dances

had no followers, requiring a zero-inflated model approach. We analyzed only bees that

had paint marks on their abdomens, ensuring that they had visited a feeder.

We used an Analysis of Deviance Wald chi-square test using the function Anova

in the MASS[32] package to further evaluate the overall effect of each fixed predictor

variable and interaction. We used the lme4 package[33] to perform these tests unless
otherwise noted. Post hoc tests were performed to determine the relationships between
the different levels of fixed predictor variables and their interactions using the package
emmeans[34]. We use R[35] for analysis.

Data Availability: Data will be available on FigShare and code will be available
on Github upon publication. Data and code available upon request by reviewers.

Ethical Compliance: Honey bees (Apis mellifera) were used in this study.
Queens (reproductive females) and drones (males) were behaviorally selected using lab
assays to create selected lines of colonies. Worker honey bees (non-reproductive
females) were tested in the field. All colonies were kept with typical honey bee
practices. There was no ethics committee involved in approving the animal husbandry
protocol.

REFERENCES

1. Ward, A. J. W., Herbert-Read, J. E., Sumpter, D. J. T. & Krause, J. Fast and
accurate decisions through collective vigilance in fish shoals. *Proc. Natl. Acad. Sci.*
108, 2312–2315 (2011).

2. Sasaki, T. & Biro, D. Cumulative culture can emerge from collective intelligence in
animal groups. *Nat. Commun.* **8**, 15049 (2017).

3. Couzin, I. D. Collective cognition in animal groups. *Trends Cogn. Sci.* **13**, 36–43
(2009).

4. Sumpter, D. J. *Collective animal behavior*. (Princeton University Press, 2010).

5. Camazine, S. *et al. Self-organization in biological systems*. (Princeton university
press, 2003).

6. Couzin, I. D. & Krause, J. Self-Organization and Collective Behavior in Vertebrates.

in *Advances in the Study of Behavior* **32**, 1–75 (Elsevier, 2003).

7. Feinerman, O. & Korman, A. Individual versus collective cognition in social insects.

J. Exp. Biol. **220**, 73–82 (2017).

8. Dornhaus, A. & Franks, N. R. Individual and collective cognition in ants and other

insects (Hymenoptera: Formicidae). *Myrmecol. News* **11**, 215–226 (2008).

9. Sumpter, D. J. T. The principles of collective animal behaviour. *Philos. Trans. R.*

Soc. B Biol. Sci. **361**, 5–22 (2006).

10. Chandra, S. B. C., Hosler, J. S. & Smith, B. H. Heritable variation for latent inhibition

and its correlation with reversal learning in honeybees (Apis mellifera). *J. Comp.*

Psychol. **114**, 86–97 (2000).

11. Mitchell, M. D., McCormick, M. I., Ferrari, M. C. O. & Chivers, D. P. Friend or foe?

The role of latent inhibition in predator and non-predator labelling by coral reef

fishes. *Anim. Cogn.* **14**, 707–714 (2011).

12. Ferrari, M. C. O. & Chivers, D. P. Latent inhibition of predator recognition by

embryonic amphibians. *Biol. Lett.* **5**, 160–162 (2009).

13. Lubow, R. E. Latent inhibition. *Psychol. Bull.* **79**, 398 (1973).

14. Lubow, R. E. Latent inhibition as a measure of learned inattention: some problems

and solutions. *Behav. Brain Res.* **88**, 75–83 (1997).

15. Lubow, R. E. Latent inhibition: Effects of frequency of nonreinforced pre-exposure of

the CS. *J. Comp. Physiol. Psychol.* **60**, 454–457 (1965).

16. Abramson, C. I. & Bitterman, M. E. Latent inhibition in honeybees. *Anim. Learing*

Behav. **14**, 184–189 (1986).

17. Ferguson, H. J., Cobey, S. & Smith, B. H. Sensitivity to a change in reward is

heritable in the honeybee, Apis mellifera. *Anim. Behav.* **61**, 527–534 (2001).

18. Brandes, C. & Menzel, R. Common mechanisms in proboscis extension conditioning

and visual learning revealed by genetic selection in honeybees (Apis mellifera). *J.*

Comp. Physiol. A **166**, 545–552 (1990).

19. Ferrari, M. C. O. & Chivers, D. P. Learning about non-predators and safe places: the

forgotten elements of risk assessment. *Anim. Cogn.* **14**, 309–316 (2011).

20. Cook, C. N. *et al.* Individual differences in learning and biogenic amine levels

influence the behavioural division between foraging honeybee scouts and recruits. *J.*

Anim. Ecol. **88**, 236–246 (2018).

21. Latshaw, J. S. & Smith, B. H. Heritable variation in learning performance affects

foraging preferences in the honey bee (Apis mellifera). *Behav. Ecol. Sociobiol.* **58**,

200–207 (2005).

22. Waddington, K. D. Honey bee foraging profitability and round dance correlates. *J.*

Comp. Physiol. **148**, 297–301 (1982).

23. Pavlov, P. I. Conditioned reflexes: an investigation of the physiological activity of the

cerebral cortex. *Ann. Neurosci.* **17**, 136 (1927).

24. Lubow, R. E., Weiner, I. & Schnur, P. Conditioned Attention Theory. in *The*

Psychology of Learning and Motivation: Advances in Research and Theory. **15**, 1–

49 (Academic Press, 1981).

25. Gershman, S. J., Blei, D. M. & Niv, Y. Context, learning, and extinction. *Psychol.*

Rev. **117**, 197–209 (2010).

26. Lubow, R. E. *Latent inhibition and conditioned attention theory*. **9**, (Cambridge

University Press, 1989).

27. Mattila, H. R., Burke, K. M. & Seeley, T. D. Genetic diversity within honeybee

colonies increases signal production by waggle-dancing foragers. *Proc. R. Soc. B

Biol. Sci.* **275**, 809–816 (2008).

28. Hills, T. T., Todd, P. M., Lazer, D., Redish, A. D. & Couzin, I. D. Exploration versus

exploitation in space, mind, and society. *Trends Cogn. Sci.* **19**, 46–54 (2015).

29. Cobey, S. W., Tarpy, D. R. & Woyke, J. Standard methods for instrumental

insemination of *Apis mellifera* queens. *J. Apic. Res.* **52**, 1–18 (2013).

30. Harbo, J. R. Propagation and instrumental insemination. in *Bee Breeding and

Genetics,* 361–389 (Academic Press, Inc, 1986).

31. Liang, Z. S. *et al.* Molecular Determinants of Scouting Behavior in Honey Bees.

Science **335**, 1225–1228 (2012).

32. Ripley, B. *et al.* Package 'mass'. *Cran R* (2013).

33. Bates, D., Maechler, M., Bolker, B. & Walker, S. lme4: Linear mixed-effects models

using Eigen and S4. *R Package Version* **1**, 1–23 (2014).

34. Lenth, R. & Lenth, M. R. Package 'lsmeans'. *Am. Stat.* **34**, 216–221 (2018).

35. Team, R. C. R: A language and environment for statistical computing. (2018).

Acknowledgements: We thank S. Ohrt, E. Sezen, N. Kulkarni, and A. Phillips for help

with data collection. We thank C. Kwapich and D. Charbonneau for comments on drafts

of this manuscript. This grant was funded by NIH NIGMS R01GM113967 to BHS, JG, &

NPW, and NIH NIGMS F32GM126728 to CNC.

Author Contributions: CNC, TM, JG, NPW, BHS conceived of and helped design the

study. CNC, NJL, and TM analyzed the data. CNC and CO created the genetically

selected lines and CO maintained them. CNC carried out data collection and wrote the

first draft of the manuscript. CNC, TM, NJL, NPW, BHS discussed results. All authors

commented on the manuscript.

Competing Interests: The authors declare no competing financial interests