**Title Page**

**Running title:** Evolutionary history of Brassolini butterflies

**Title:** Mesoamerica is a cradle and the Atlantic Forest is a museum of Neotropical butterfly

diversity: Insights from the evolution and biogeography of Brassolini (Lepidoptera:

Nymphalidae)

**Authors:** Pável Matos-Maraví[1,2,3]*, Niklas Wahlberg[4], André V. L. Freitas[5], Phil DeVries[6,7],

Alexandre Antonelli[1,2,8,9], Carla M. Penz[6]

**Institutions:**

[1] Department of Biological and Environmental Sciences, University of Gothenburg, Carl

Skottbergs gata 22B, 41319 Gothenburg, Sweden

[2] Gothenburg Global Biodiversity Centre, Carl Skottbergs gata 22B, 41319 Gothenburg, Sweden

[3] Biology Centre of the Czech Academy of Sciences, Institute of Entomology, Branišovská

1160/31, 37005 České Budějovice, Czech Republic

[4] Department of Biology, Lund University, Sölvegatan 37, 22362 Lund, Sweden

[5] Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas,

Rua Monteiro Lobato 255, CEP 13.083-862 Campinas, São Paulo, Brazil

[6] Department of Biological Sciences, University of New Orleans, 2000 Lakeshore Drive, New

Orleans, LA 70148, USA

[7] Courtesy Curators of Lepidoptera, Florida Museum of Natural History, 1659 Museum Road,

Gainesville, FL 32611, USA

[8] Royal Botanical Gardens Kew, TW9 3AE Richmond, United Kingdom

[9] Department of Plant Sciences, University of Oxford, South Parks Road, OX1 3RB Oxford,

United Kingdom

**Corresponding author:**

Pável Matos-Maraví; Institute of Entomology, Biology Centre CAS, Branišovská 1160/31, 370

05 České Budějovice, Czech Republic; E-mail: pavel.matos@entu.cas.cz, ORCID:

https://orcid.org/0000-0002-2885-4919

**ORCID ids:**

Pável Matos-Maraví, https://orcid.org/0000-0002-2885-4919

Niklas Wahlberg, https://orcid.org/0000-0002-1259-3363

André V. L. Freitas, https://orcid.org/0000-0002-5763-4990

Phil DeVries, https://orcid.org/0000-0002-5692-6684

Alexandre Antonelli, https://orcid.org/0000-0003-1842-9297

Carla M. Penz, https://orcid.org/0000-0002-2544-3508

**ABSTRACT**

Regional species diversity is ultimately explained by speciation, extinction, and dispersal. Here we estimate dispersal and speciation rates of Neotropical butterflies to propose an explanation for their distribution and diversity of extant species. We focus on the tribe Brassolini (owl butterflies and allies): a Neotropical group that comprises 17 genera and 108 species, most of them endemic to rainforest biomes. We infer a robust species tree using the multispecies coalescent framework and a dataset including molecular and morphological characters. This formed the basis for three changes in Brassolini classification: 1) Naropina, SYN. NOV. is subsumed within Brassolina; 2) *Aponarope*, SYN. NOV. is subsumed within *Narope*; 3) *Selenophanes orgetorix*, COMB. NOV. is reassigned from *Catoblepia* to *Selenophanes*. By applying biogeographical stochastic mapping, we found contrasting species diversification and dispersal dynamics across rainforest biomes, which might be partly explained by the geological and environmental history of each bioregion. Our results reveal a mosaic of biome-specific evolutionary histories within the Neotropics, where butterfly species have diversified rapidly (cradles: Mesoamerica), have accumulated gradually (museums: Atlantic Forest), or have alternately diversified and accumulated (Amazonia). Our study contributes evidence from a major butterfly lineage that the Neotropics are a museum and cradle of species diversity.

**KEYWORDS:** Ancestral state inference – biogeographical stochastic mapping – bioregion delimitation – dispersal – multispecies coalescent – speciation – total-evidence analyses.

## INTRODUCTION

Terrestrial biomes are heterogeneous in terms of species composition and distribution, and some areas have accumulated greater diversity and endemicity than others. A combination of three evolutionary mechanisms explains this biogeographical pattern: species origination, extinction, and dispersal (Goldberg *et al.*, 2005; Jablonski *et al.*, 2006). For example, a species-rich region can attain its present diversity by recent and exceptionally high rates of speciation, becoming a "cradle of diversity", or by the persistence of old lineages via low species extinction rates, becoming a "museum of diversity" (e.g., Stebbins, 1974; Gaston & Blackburn, 1996; Bermingham & Dick, 2001; Richardson *et al.*, 2001). Furthermore, a species-rich region might be a "source of diversity" when dispersal out of the region is high, or a "sink of diversity" when regional species diversity is accrued from external sources via dispersal into the region (Antonelli *et al.*, 2018a). Characterizing the interplay among speciation, extinction, and dispersal is thus highly relevant for understanding the origin and evolution of total regional diversity and endemism (Wiens & Donoghue, 2004; Roy & Goldberg, 2007).

Much of the world's biodiversity and endemism are concentrated in Neotropical rainforests in Central America, the Amazon basin, and the Atlantic Forest in southeastern Brazil. However, there is no consensus on whether these currently separated rainforests were connected once or multiple times, nor for how long (Jaramillo & Cárdenas, 2013). It is also unclear whether Neotropical rainforests represent centers of ecological stability that share a common evolutionary history (Moritz *et al.*, 2000). Mammals and birds in Neotropical biodiversity hotspots, including the Brazilian Atlantic Forest and rainforests in Mesoamerica and the neighboring Chocó in the northwestern side of the Andes (Myers *et al.*, 2000), apparently underwent rapid diversification

(Igea & Tanentzap, 2019). In other groups, a larger number of lineages seem to have dispersed

out of those areas compared to the rest of the Neotropics (e.g., Machado *et al.*, 2018; Toussaint *et*

*al.*, 2019), suggesting that such rainforests are both cradles and sources of diversity (Igea &

Tanentzap, 2019). Other Neotropical regions are, in contrast, museums of diversity given the old

and gradual accumulation of species through time, such as Amazonia for Euptychiina

(Nymphalidae) (Peña *et al.*, 2010) and Troidini butterflies (Papilionidae) (Condamine *et al.*,

2012). It thus seems that the checkered spatial assemblage of rainforest areas where speciation

rates are high plus those where extinction rates are low might explain why the Neotropical region

as a whole is considered both a cradle and museum of diversity (McKenna & Farrell, 2006;

Moreau & Bell, 2013; Antonelli *et al.*, 2018b). However, the impacts of geological and

paleoenvironmental changes in the Neotropics together with the roles of speciation and dispersal

have seldom been jointly elucidated among taxa occurring in individual rainforest areas.

Here we examine biogeographical patterns for butterflies in the tribe Brassolini (Nymphalidae:

Satyrinae), an exclusively Neotropical monophyletic group (Freitas & Brown, 2004; Wahlberg *et*

*al.*, 2009; Espeland *et al.*, 2018; Chazot *et al.*, 2019a) consisting of 17 genera and 108 species

found mainly in the rainforests of Mesoamerica, the northwestern slope (NW) of the Andes

(including e.g., Chocó), the Atlantic Forest, and Amazonia (see Penz, 2007, for an overview).

Studies that focused on individual brassoline genera used adult morphology (Penz, 2008,

2009a,b; Garzón-Orduña & Penz, 2009) and DNA data (Penz, Mohammadi, & Wahlberg, 2011a;

Shirai *et al.*, 2017) to infer species-level phylogenies, but several deep nodes had weak or even

conflicting phylogenetic support in tribal-level analyses (Penz *et al.*, 2013). We infer a time-

calibrated species phylogeny of Brassolini by merging all previous datasets and generating new

DNA and morphological data to infer the diversification and biogeographical histories of these

butterflies. From a biogeographical perspective, if physical connection among Neotropical

biomes existed during the Neogene and Quaternary periods (i.e., the past 23 Myr) (Hoorn *et al.*,

2010), we expect that this will be reflected in the timing and magnitude of dispersal and

speciation trends through time. To assess this expectation, we estimate dispersal rates and

speciation for Brassolini butterflies.

**MATERIAL AND METHODS**

*MOLECULAR DATASET*

Adult specimens were collected in Mexico, Costa Rica, Ecuador, Peru, and Brazil (Supporting

Information, Table S1), and identified to species based on various sources (Casagrande, 2002,

2004; Austin *et al.*, 2007; Penz, 2008, 2009b,a; Garzón-Orduña & Penz, 2009; Penz *et al.*, 2017,

2011a; Penz, Simonsen, & DeVries, 2011b; Chacón *et al.*, 2012). Total DNA was obtained from

two butterfly legs using QIAGEN's DNeasy kits. Based on PCR primers and protocols described

in Wahlberg & Wheat (2008), we Sanger-sequenced fragments of the mitochondrial COI gene

(1,475 bp) and the nuclear genes CAD (850 bp), EF1α (1,240 bp), GAPDH (691 bp), RpS5 (617

bp), and *wingless* (400 bp) (outsourced to Macrogen, Seoul). Our molecular dataset included 64

species in 15 out of 17 Brassolini genera (Penz, 2007); the two missing genera (*Aponarope* and

*Mielkella*) are monotypic. Specifically, we obtained genetic data from 83 individuals classified in

57 Brassolini species, and retrieved COI sequences available from the BOLD database

(http://boldsystems.org) for an additional seven. The analyses also included genetic data of 19

outgroup taxa in the subfamily Satyrinae from public data deposited in GenBank. All DNA

sequences from this study are in GenBank (accession numbers MK551348–MK551551).

*MORPHOLOGICAL AND TOTAL-EVIDENCE DATASET*

Our morphological dataset included 255 discrete characters (201 from adults and 54 from early

stages) for 64 species, 44 of them corresponding to the same species included in the molecular

dataset. The morphological matrix included all 17 Brassolini genera and was concatenated from

previous studies (Penz, 2007, 2008, 2009b,a; Garzón-Orduña & Penz, 2009; Penz *et al.*, 2013)

by eliminating duplicate characters and scoring new data for species missing entries. Specimens

used were the same as listed in the original publications. The combined molecular and

morphological dataset comprised all genera and 84 out of 108 Brassolini species (missing data

coded as "?"). We call this the total-evidence dataset, which we used to infer a time-calibrated

phylogeny.

*DISTRIBUTION DATASET AND BIOREGION DELIMITATION*

We advocate for a data-informed bioregionalization of Brassolini geographical distribution.

Specifically, we expect that parametric bioregionalization will better inform on the best-fit

delimitation of areas for Brassolini compared to any arbitrary assignment of areas within the

hierarchical classification by Morrone (2014a,b) (subregions, dominions or provinces).

We delineated bioregions (*sensu* Vilhena & Antonelli, 2015) using geo-referenced occurrence

points of Brassolini identified with taxonomic certainty at the species level, of which 6,378

points came from GBIF (https://gbif.org; retrieved on 21st January 2019) and 881 points from the

ATLANTIC BUTTERFLIES dataset (Santos *et al.*, 2018). Both databases follow the current

taxonomic understanding of Brassolini, thus, we did not encounter dubious species names. We

used the R v.3.5.3 (R Core Team, 2020) package CoordinateCleaner v.2.0-11 (Zizka *et al.*, 2019)

for automated cleaning of species occurrences. In addition to four *Caligo eurilochus*

observations from tropical butterfly houses in Germany that were removed a priori, the software

flagged 81 potential errors, such as occurrence records in the ocean, geographic coordinates

identical to country centroids, and occurrences in close proximity to biodiversity institutions

such as natural history museums (Supporting Information, Fig. S1). We uploaded the 7,174

vetted geo-referenced occurrences, all within the known natural geographical range of Brassolini

(see below), to the web application Infomap Bioregions (Edler *et al.*, 2017) at

http://bioregions.mapequation.org/. We set the maximum and minimum values for cell sizes to 4

and 2, and for cell capacities to 40 and 4. We used different values of cluster cost, from 0.1 to

3.0, in steps of 0.1. The number of inferred areas remained stable between values of 1.8 and 2.4

and, thus, received the strongest support.

The bioregions identified correspond to formerly recognized biomes, or combinations of them:

1) Mesoamerica plus the NW flank of the Andes (including Chocó), 2) Brazilian Atlantic Forest,

and 3) the remaining tropical South America, including the Amazon drainage basin and the

seasonally dry diagonal biomes Cerrado and Caatinga (Fig. 1). Note that this delimitation based

on different data and approach approximates the Dominions of Morrone (2014b): 1) a joint area

including the Mesoamerican dominion plus the provinces of the Pacific dominion west of the

Andes, in agreement with the strong floristic connection of Central America and Chocó (Pérez-

Escobar *et al.*, 2019), 2) the Parana dominion comprising the Brazilian Atlantic Forest, and 3)

the remaining tropical provinces in South America (Fig. 1).

To inform the biogeographical and within-area speciation rate analyses, we generated a species-

level distributional dataset that includes all 108 valid Brassolini species. We surveyed the

literature to score species occurrences in the defined bioregions because the geo-referenced

dataset included only 72 Brassolini species with occurrence points. To investigate dispersal

patterns across the South American dry diagonal, we kept this area separate (Fig. 1). We used

distributional maps and observations, not always with primary geographic coordinates, from

taxonomic revisions (Bristow, 1981, 1982, 1991; Casagrande, 2002; Furtado & Campos-Neto,

2004; Penz, 2008, 2009a,b; Garzón-Orduña & Penz, 2009; Penz *et al.*, 2017) and published

butterfly inventories in Mesoamerica (DeVries, 1983, 1994; Janzen & Hallwachs, 2009; Basset

*et al.*, 2015), Amazonia (Pereira Martins *et al.*, 2017), Cerrado (Pinheiro & Emery, 2006; Emery

*et al.*, 2006; Silva *et al.*, 2012; Pereira Martins *et al.*, 2017; Dickens *et al.*, 2019), Caatinga

(Zacca & Bravo, 2012), and Atlantic Forest (Santos *et al.*, 2011, 2018; Pérez *et al.*, 2017; Melo

*et al.*, 2019; Soldati *et al.*, 2019). We call this the presence/absence biogeographical dataset.

*PHYLOGENETIC INFERENCE AND DIVERGENCE TIME CALIBRATION*

We ran preliminary phylogenetic analyses using each gene and morphological dataset to check

for tree topology conflicts. The six molecular datasets were partitioned by codon position and the

morphological dataset was partitioned by using homoplasy scores calculated through implied

weighting parsimony (Rosa *et al.*, 2019). We calculated homoplasy measurements $f$ using TNT

v.1.5 (Goloboff & Catalano, 2016) under the default concavity parameter $k = 3$, and we used

these values to subdivide the morphological dataset into 11 partitions (Supporting Information,

Table S3). All seven phylogenetic analyses (6 gene loci and morphology) were run using

MrBayes v.3.2.6 (Ronquist *et al.*, 2012) via the CIPRES Science Gateway v.3.3 (Miller *et al.*,

2010). We performed model averaging over all substitution models within the GTR model

family (Huelsenbeck *et al.*, 2004), taking into account rate variation across sites (+I and +Γ

models) for the molecular partitions, and we applied the Markov (MKv) substitution model

(Lewis, 2001) for the morphological partitions. The Bayesian inferences were run two

independent times for 50 million generations, sampling trees every 5,000[th] generation and

discarding the first 25% of the sampled distribution as burnin. After checking that the log-

probabilities reached a stationary distribution, the average standard deviation of split frequencies

were below 0.005, PSRF values close to 1.000, and the estimated sample sizes (ESS) above 200,

we summarized the post-burnin sampled trees using the 50% majority-rule consensus method.

We inferred time-calibrated species trees using the StarBEAST2 v.0.15.5 package (Ogilvie *et al.*,

2017) available in BEAST v.2.6.3 (Bouckaert *et al.*, 2014). We removed the species *Opsiphanes*

*camena* from the dataset because it had only 325 bp of the COI locus, and its phylogenetic

position was not stable in the preliminary analyses. We used a gene-based partitioning strategy,

each having unlinked tree models in the multispecies coalescent framework. We performed

model averaging over 31 substitution models available in the bModelTest v.1.2.1 package

(Bouckaert & Drummond, 2017). We applied uncorrelated relaxed-clock models for all

partitions because strict molecular clock models were rejected based on the stepping-stone

method (Table S4). We estimated two molecular clock rates, one for the mitochondrial locus and

one for the nuclear loci, by using the gamma distribution for the clock rate priors (shape

parameter ($\alpha$) = 0.01 for mitochondrial clock and 0.001 for nuclear clock; scale parameter ($\beta$) =

1.0). We used the Yule tree model plus two molecular clocks because it received decisive

support, based on path-sampling steps, over other models with different combinations of one,

two, or six clocks, and the Yule and birth-death tree models (Table S5). We added the

morphological dataset onto the species tree and we conditioned the characters to the Markov

(Mk) substitution model (Lewis, 2001). BEAST2 assigns morphological partitions with unlinked

site models based on characters having the same number of states. All morphological characters

were considered to evolve under the same species tree and relaxed clock model with an

uninformative rate prior (1/X distribution).

Inasmuch as there are no described fossils of Brassolini, we relied on secondary calibrations to

time-calibrate the species tree. We conservatively used uniform distributions encompassing the

95% highest posterior density (HPD) from a large fossil-calibrated, genus-level butterfly

phylogeny (Chazot *et al.*, 2019a) to constrain six outgroup nodes (Table 1; Fig. S5).

The analyses were run four independent times for 300 million generations, via CIPRES. We

sampled 20,000 trees from the posterior distribution and discarded the first 25% as burnin. We

checked that the ESS values were above 200 using Tracer v.1.7.1 (Rambaut *et al.*, 2018). We

merged the independent runs using LogCombiner (part of the BEAST v.2.6.3 package) and we

summarized the post-burnin species trees as a maximum clade credibility (MCC) tree in

TreeAnnotator (part of the BEAST v.2.6.3 package).

As supplementary analyses, we inferred total-evidence phylogenetic trees using the

concatenation approach assuming that all gene loci and morphological characters have evolved

under the same tree topology. We carried out gene-tree discordance test to assess the

convenience of the multispecies coalescent framework for our dataset. We performed likelihood-

based tree topology tests to evaluate the agreement of the molecular data with three tree

topologies: 1) the morphology-based systematics of Brassolini (Penz, 2007) suggesting that

*Narope* is sister to remaining genera in the subtribe Brassolina, 2) the multispecies coalescent

species tree, and 3) the concatenation-based tree topology (Supporting Information).

*WITHIN-AREA CLADOGENESIS AND DISPERSAL AMONG BIOREGIONS*

We estimated ancestral geographic ranges and colonization rates among bioregions using the

dispersal-extinction-cladogenesis (DEC) model (Ree *et al.*, 2005) as implemented in the R

package BioGeoBEARS v.1.1.2 (Matzke, 2013). We used the presence/absence biogeographical

dataset and the four defined bioregions: 1) Mesoamerica plus NW Andes, 2) Atlantic Forest, 3)

Amazonia, and the 4) Dry diagonal. The latter was kept separate to estimate dispersal between

Atlantic Forest and Amazonia across a large area of less suitable environments for Brassolini.

We did not constrain any *a priori* dispersal multiplier, nor did we stratify dispersal rates across

the phylogeny. This allowed us to evaluate whether the data alone is in agreement with main

geological and paleogeographical events in the Neotropics. To account for ancestral state

uncertainty, we carried out 100 biogeographical stochastic mappings (BSM; Dupin *et al.*, 2017).

To account for divergence time uncertainty, we used 100 random topologies from the BEAST2

posterior distribution. To account for phylogenetic uncertainty of 25 missing species, we

randomly added such lineages to their currently assigned genera in the 100 posterior

phylogenies. In this case, we assumed that phylogenetic diversity per genus was maximized and

we placed missing taxa in the crown genera given our comprehensive taxonomic sampling. All

genera are monophyletic, but we conservatively joined *Selenophanes* and *Catoblepia* (posterior

probability, PP = 1.0; Fig. 2) when assigning missing species because the monophyly of

*Catoblepia* was only moderately supported (PP = 0.75). We also carried out the biogeographical

analyses using the taxonomically incomplete inferred species trees to rule out any bias in the

approach.

Variation in taxonomic opinion and effort across localities might be an important source of error

in macroevolutionary analyses (Faurby *et al.*, 2016). After analyzing trends in the year of

description of each Brassolini species and the year of the latest revision of their current

taxonomic status (e.g., when former subspecies were raised to species), we did not observe any

taxonomic biases among Neotropical bioregions in terms of taxon discovery, description and

taxonomic opinion (i.e., oversplitting species vs. multiple infraspecific taxa) (Supporting

Information). We therefore estimated the number of dispersal events and within-area-

cladogenesis events by simulating areas on nodes based on the 10,000 pseudoreplicated

biogeographic histories (100 BSM × 100 posterior species trees). We calculated colonization

rates through time as $c_{XtoY}(t_1) = d_{XtoY}(t_1) / Br(t_1)$, where $d_{XtoY}(t_1)$ is the number of inferred

dispersal events from area X to Y in a 2-million years interval, and $Br(t_1)$ is the total length of all

branches in the 2-million-years interval (Antonelli *et al.*, 2018a). In addition, we calculated the

relative number of speciation events in area X through time as $\lambda_X(t_1) = s_X(t_1) / L_X(t_0)$, where

$s_X(t_1)$ is the number of within-area cladogenesis in a 2-million-years interval, and $L_X(t_0)$ is the

number of inferred lineages in area X in the preceding 2-million-years interval ($t_0$), calculated as

the cumulative sum of within-area cladogenesis and dispersal into area X minus local extinction

(Xing & Ree, 2017).

*SPECIATION WITHIN BIOREGIONS*

For a second, independent estimate of within-area speciation, we calculated speciation rates for

all tips and branches in the MCC species tree using BAMM v.2.5.0 (Rabosky, 2014). To account

for the 25 missing species, we generated clade-specific sampling fractions at the genus level

under the assumption of random taxonomic sampling (FitzJohn *et al.*, 2009). We used the R

package BAMMtools v.2.1.6 (Rabosky *et al.*, 2014) to estimate prior values for the evolutionary

rate parameters, and we assigned the expected number of shifts to 1.0, which is a conservative

value that minimizes type I errors and is recommended for phylogenies including ~100 species

(Rabosky, 2014). We ran the analysis for 10 million generations, sampling event data every

1,000 generations. After discarding the first 20% samples, we checked that the ESS value was

higher than 200 and the likelihood estimates reached a stationary distribution.

We used the function "dtRates" in BAMMtools to compute mean speciation rates along branches

in ~2-million years intervals (tau parameter set to 0.05, as the root of phylogeny was 37.9 Mya).

We assigned the most probable inferred area from the 10,000 BSMs to the points where branches

cross time intervals (Igea & Tanentzap, 2019). When the most probable area for a node and its

parental one were the same, we assumed that the branch connecting both nodes occurred in such

an area. When inferred areas for a node and its parental one were different, we assumed that the

dispersal event occurred stochastically along the branch. We computed mean speciation rates

only for cladogenesis events that most probably occurred in a single bioregion.

All the datasets and scripts used in this study can be found in TreeBASE (ID 25040) and Zenodo

(DOI: http://doi.org/10.5281/zenodo.4500393).

**RESULTS**

*PHYLOGENY AND DIVERGENCE TIMES*

Our molecular dataset comprised 5,285 bp, of which 2,193 represented variable sites (Table 2).

Our morphological dataset comprised 219 binary characters, 32 characters with 3 states, and 4

characters with 4 states. The multispecies coalescent model recovered four major clades with

posterior probabilities (PP) of 1.0, namely, clade A, *Dynastor* and *Dasyophthalma*; clade B,

*Opoptera* and *Narope*; clade C, *Caligopsis*, *Eryphanis*, and *Caligo*; clade D, *Penetes*,

*Catoblepia*, *Selenophanes*, *Mielkella*, *Orobrassolis*, *Blepolenis*, and *Opsiphanes*. The maximum

clade credibility (MCC) species tree depicts the following phylogenetic relationships: *Bia*: (clade

A: (clade B: (*Brassolis*: (clade C: clade D)))), with divergence times among these clades between

29 and 34 Mya (entire highest posterior density HPD, 20.24–43.64 Mya). However, the

relationships among clades B, C, D, and the genus *Brassolis* received low support (PP = 0.56–

0.72; Fig. 2). The concatenation-based analyses produced similar divergence times among major

Brassolini clades compared to the multispecies coalescent species tree (between 26 and 30 Mya,

entire HDP, 20.38–39.22 Mya) but recovered higher posterior probabilities for Brassolini deep

nodes when using the molecular (0.97–1.00; Fig. S2) and total-evidence datasets (0.83–0.96; Fig.

S3). Nevertheless, the consensus and MCC tree topologies among all concatenation-based

phylogenetic analyses remained similar, except for the genus *Brassolis* as sister to the clade D,

and clade C sister to these two. The likelihood-based tree topology tests could not resolve this

main discrepancy between the multispecies coalescent and the concatenation-based species tree

topologies (Table S6). The tree topology tests, however, clearly rejected the relationship of

*Narope* as sister to the subtribe Brassolina suggested by previous cladistic analyses of

morphological characters (*p*-value = 0.0003).

Importantly, the total-evidence dataset supported posterior probability of 1.0 for the crown node

of non-monotypic genera, regardless of phylogenetic method, except for two cases: *Narope*

paraphyletic with respect to the monotypic *Aponarope*, and *Catoblepia* paraphyletic with respect

to *Selenophanes* (see *Consequences for Brassolini systematics and classification* in the

Discussion section).

*GEOGRAPHIC RANGE EVOLUTION AND COLONIZATION RATES*

The inferred ranges coming from the 10,000 pseudoreplicated biogeographic histories and

plotted against the MCC species tree in Fig. 2 suggested that Brassolini most probably (*Pr* =

0.50) originated in Amazonia at 38 Mya (entire HPD, 27.68–46.97 Mya). The early rapid

radiation of Brassolini between 29 and 34 Mya also probably took place in Amazonia (*Pr* =

0.44–0.67).

When taking into account phylogenetic and divergence time uncertainties by using the Bayesian

posterior distribution of Brassolini species trees, dispersal out of Amazonia had the highest rates

compared to other bioregions (Fig. 3). Colonization rate from Amazonia to Mesoamerica showed

a sharp increase by ~2–4 Mya and is currently the highest rate among other dispersal types.

Dispersal into Mesoamerica from other bioregions probably occurred throughout the Miocene

but significantly increased at ~6–8 Mya (Fig. S8). Dispersal out of Mesoamerica + NW Andes,

however, was very low through time, and only since the past ~2 Mya did it exponentially

increase toward Amazonia. Dispersal from Amazonia to the dry diagonal increased throughout

the Plio-Pleistocene and has since then equaled dispersal to Atlantic Forest. Dispersal from

Amazonia into the Atlantic Forest was higher in the Oligocene–Miocene transition at ~23–27

Mya than from/to any other Neotropical region. However, for most of the Miocene and Pliocene,

dispersal into or out of the Atlantic Forest from/to any other bioregion was overall low through

time, rendering the Atlantic Forest isolated; only since the Pleistocene ~2 Mya dispersal into and

from the dry diagonal has increased until the present. The dispersal patterns remain similar in

supplementary biogeographical analyses using the taxonomically incomplete inferred species

trees (Supporting Information).

*REGIONAL SPECIATION THROUGH TIME*

The relative rates of within-area cladogenesis estimated in BioGeoBEARS and speciation

estimated in BAMM were congruent (Fig. 4). Speciation rates increased through time in all areas

except in the dry diagonal, which has had near zero endemic cladogenesis. Diversification in the

Atlantic Forest might have begun during the Oligocene-Miocene transition at ~20–24 Mya.

During that time, the cladogenesis rate was slightly above 0.5 (Fig. 4A) and speciation rate began

to increase (Fig. 4B), likely driven by the split of *Penetes* and, to a lesser extent, the stem lineage

of *Dasyophthalma*. However, there seem to be a peak in cladogenesis at ~8–10 Mya, specifically

in the lineages comprising extant *Dasyophthalma*, *Orobrassolis* and *Blepolenis*. Afterwards,

diversification in the Atlantic Forest remained constant throughout most of the Miocene until

~2–4 Mya, when it slightly decreased towards the present. In contrast, speciation in

Mesoamerica + NW Andes began later, at ~8–12 Mya, remained constant during most of the

Miocene and Pliocene, and sharply increased only at ~2 to 4 Ma driven by the origin of extant

endemic lineages within *Caligo*, *Opsiphanes*, and *Eryphanis*. Speciation in Amazonia was

episodically higher than in the Atlantic Forest and Mesoamerica throughout most of the

Miocene, but it also dramatically increased during the Pleistocene (~2 Mya) by the split of

several extant species within *Bia*, *Eryphanis*, and *Catoblepia*. At present, the speciation rate in

Mesoamerica + NW Andes is similar to the Amazonian lineages, both apparently being higher

than in the Atlantic Forest.

**DISCUSSION**

Here we investigated the evolutionary history of the butterfly tribe Brassolini, and studied the

role of regional speciation and dispersal to understand the origin of extant species across

Neotropical rainforest biomes (Fig. 5). By using molecular and morphological datasets and the

multispecies coalescent framework, we propose a revised hypothesis of relationships among

genera, and make three changes to the classification of Brassolini at the tribal and genus levels.

*CONSEQUENCES FOR BRASSOLINI CLASSIFICATION*

Brassolini has been previously subdivided in three subtribes based on shared morphological

similarities: Biina (monotypic), Naropina (*Narope* and *Aponarope*), and Brassolina (remaining

Brassolini genera) (Casagrande, 1995; Penz, 2007). Nonetheless, here *Narope* is sister to

*Opoptera* (clade B in Fig. 2) with strong statistical support, and this clade is nested within the

subtribe Brassolina (*sensu* Casagrande, 1995). These relationships have not been recovered in

any cladistic analysis of Brassolini to date. At the genus level, the main discrepancy among the

multispecies coalescent and the concatenation-based phylogenies was in the phylogenetic

position of the genus *Brassolis*. Morphological character states of adult and early stages in this

genus are highly polymorphic and divergent compared to other Brassolini genera, thus,

obscuring the phylogenetic relationships within *Brassolis* and among this and other lineages

(Garzón-Orduña & Penz, 2009; Penz *et al.*, 2013). Nonetheless, the six molecular markers aided

to resolve the apparent long-branch attraction between *Brassolis* and *Dynastor* seen in

morphological phylogenies (Penz, 2007), and grouped *Brassolis* with clades C and D (*Caligo*

and *Opsiphanes* groups) in both the multispecies coalescent and concatenation-based

phylogenies.

We propose the following changes to the Brassolini classification:

1) The subtribe Naropina Stichel, 1925, **SYN. NOV.** is subsumed within Brassolina Boisduval,

1836, and Brassolini Boisduval, 1836, is redefined to include the subtribes Biina Herrich-

Schäffer, 1864 (monotypic) and Brassolina.

2) The monotypic genus *Aponarope* Casagrande, 1982, **SYN. NOV.** is subsumed within *Narope*

Doubleday, 1849. *Narope sutor* Stichel, 1916, was likely segregated in the monotypic

*Aponarope* by Casagrande (1982) to accommodate the high number of autapomorphic

characters. Using a cladistic analysis of 80 morphological characters, Penz *et al.* (2013)

previously found that *Aponarope* is nested within *Narope* but conservatively retained *Aponarope*

until more evidence increases the support for this relationship. Based on our results, we revise

the genus assignment for *Narope sutor*, **COMB. REST.**, and synonymize *Aponarope*, **SYN.**

**NOV.** to render the genus *Narope* monophyletic.

3) *Selenophanes orgetorix* (Hewitson, 1870), **COMB. NOV.** This species grouped with

members of *Selenophanes* Staudinger, 1887, with high posterior probability (0.99) in our

analysis, which justifies the revised generic assignment. Using COI barcoding sequences, Shirai

*et al.* (2017) also found that the genus *Catoblepia* Stichel, 1902, was paraphyletic with respect to

*Selenophanes* due to the sister relationship between *S. orgetorix* and *Selenophanes* species.

*SPECIES DIVERSIFICATION AND DISPERSAL*

One third of Brassolini species are endemic to rainforests in Mesoamerica, NW slope of the

Andes including the Chocó, and the Brazilian Atlantic Forest. Rapid butterfly speciation has

probably occurred in Mesoamerica and NW Andes since the Pleistocene, suggesting that it acted

as a cradle of diversity. In contrast, low and constant diversification since at least the early

Miocene has shaped Brassolini endemic diversity in the Atlantic Forest. This finding, coupled

with the survival of old lineages found mainly in montane habitats (e.g., *Penetes*,

*Dasyophthalma*, *Orobrassolis*) suggests that the Atlantic Forest acts as a museum of diversity for

Brassolini. Our results also suggest that Brassolini diversified mainly in Amazonia through

episodic increases in cladogenesis rate throughout the Miocene from about 20 Ma (Fig. 4A), and

such diversification events might have been accompanied by increasing dispersal out of this

bioregion.

The diversification and biogeographical history of Brassolini are consistent with the source,

museum, and cradle of diversity scenarios previously proposed for the Amazon rainforest.

Amazonia has been the main source of Neotropical vertebrate and plant species diversity by

means of dispersal during the past 60 million years (Antonelli *et al.*, 2018a). Rapid species

diversification of some butterfly groups occurred there since at least the Oligocene (e.g., Matos-

Maraví *et al.*, 2013; Chazot *et al.*, 2016; Toussaint *et al.*, 2019), and old lineages might have

gradually accumulated in Amazonia since that time (Peña *et al.*, 2010; Condamine *et al.*, 2012).

However, more recent diversification episodes supporting Amazonia as alternating museum and

cradle of diversity could have been triggered by paleoenvironmental changes during the Miocene

(Antonelli *et al.*, 2009; Antonelli & Sanmartín, 2011; Chazot *et al.*, 2019b).

The establishment of the Pebas wetland system in western Amazonia during most of the Miocene

(~8–24 Mya) (Wesselingh & Salo, 2006) has likely hindered dispersal from Amazonia toward

the NW Andes and the Pacific in some butterfly groups (Elias *et al.*, 2009; De-Silva *et al.*, 2016,

2017; Chazot *et al.*, 2019b). In the case of Brassolini, the Pebas wetland system has also likely

hindered dispersal out of Amazonia (Fig. 3) but it might not have drastically limited

diversification (Fig 4). This wetland system was occasionally affected by marine flooding events,

which were probably short-lived, and at least two dated events at ~13.7 and ~17.8 Mya

(Jaramillo *et al.*, 2017a) were concurrent with the early and rapid cladogenesis events of major

Brassolini clades in Amazonia. The Pebas wetland might in fact have increased opportunities for

dispersal across Amazonia and diversification of some typical coastal plants (Bernal *et al.*,

2019), including many of the Brassolini's hostplants in the region: Arecaceae (palms), Poaceae

(bamboos), and Zingiberales such as Marantaceae (Beccaloni *et al.*, 2008; Janzen *et al.*, 2009). It

is plausible that the evolutionary consequences of the Miocene marine incursions in western

Amazonia might have been mixed and rather ephemeral (Musher *et al.*, 2019), restricting

dispersal and diversification in some butterfly lineages such as Ithomiini (Chazot *et al.*, 2019b)

while providing ecological opportunities for diversification of others, such as Brassolini.

The increase in diversification of Amazonian lineages in the Pleistocene might be related to

younger paleoenvironmental changes, such as fluctuating river dynamics mediated by

Quaternary glacial cycles (Ribas *et al.*, 2012). For example, biogeographical structure in the

genus *Bia* seems to be associated with major Amazon-basin rivers (Penz *et al.*, 2017), suggesting

that riverine barriers might have been involved in allopatric differentiation in this genus. Indeed,

in some areas surrounding the Amazon River, the distance between river banks is much larger

than the estimated daily dispersal of species such as *Bia actorion* (Tufto *et al.*, 2012; Penz *et al.*,

2015). Other hypotheses for the diversification of Neotropical butterflies in the Pleistocene have

been put forward, such as rainforests acting as refugia (Garzón-Orduña *et al.*, 2014), but this has

not been corroborated yet in a macroevolutionary framework (Matos-Maraví, 2016). A

phylogeographic evaluation including more samples per species would be necessary to estimate

the influence of river barriers or rainforest refugia in differentiating Brassolini populations, and

potentially to explain the apparent increase in diversification of Amazonian butterflies during the

Pleistocene.

Miocene expansions of wet forests between the Atlantic Forest and Amazonia across the dry

diagonal might have been important determinants of ancient Brassolini diversity. The onset of a

period of global cooling at ~10 to 15 Mya likely drove the expansion of open environments

(Simon *et al.*, 2009; Azevedo *et al.*, 2020), but intermittent forest connections between

Amazonia and the Atlantic Forest have been documented based on palynological data and

biogeographic studies (e.g., Costa, 2003; Werneck, 2011; Prates *et al.*, 2016; Trujillo-Arias *et al.*,

2017). Although evidence of repeated Neogene connection between the Atlantic Forest and

Amazonia is still scarce and restricted to phylogeographic studies (e.g., Ledo & Colli, 2017;

Capurucho *et al.*, 2018), the origin of the modern transcontinental Amazon river by ~10 Mya

might have facilitated dispersal of rainforest taxa (Albert *et al.*, 2018). Indeed, we found that late

Miocene dispersal rates between these wet forest biomes slightly increased through time. In

contrast, the Atlantic Forest has recently become a source of diversity for present dry diagonal as

Brassolini dispersal sharply increased during the Pleistocene, in agreement with previous

phylogeographic evidence (Costa, 2003; Batalha-Filho *et al.*, 2013; Thomé *et al.*, 2016). We

suggest that episodic Miocene wet forest expansions might have represented ecological

opportunities for Brassolini to disperse across Central Brazil, either by increased sizes of wet

forest galleries in the dry diagonal biome, or by fully connecting Amazonia and Atlantic Forest.

By the Plio-Pleistocene, speciation in Mesoamerica and NW Andes seems to have increased and

surpassed the speciation rate in the Atlantic Forest. Biotic dispersal into Mesoamerica from

South America (e.g., Mullen *et al.*, 2011; Bacon *et al.*, 2015) and butterfly dispersal from Central

America to other Neotropical areas (Toussaint *et al.*, 2019) likely occurred throughout most of

the Miocene facilitated by land availability, with Uranium-lead dating confirming that segments

of the Panama arc had already emerged by 13–15 Mya (Montes *et al.*, 2015). However, only at

the end of the Pliocene ~3 to 4 Mya did shallow seawaters fully recede from the region (Coates

*et al.*, 2004; Montes *et al.*, 2012; O'Dea *et al.*, 2016; Jaramillo *et al.*, 2017b), thus increasing

opportunities for terrestrial species diversification in lowland areas that were sporadically

inundated by seawater during the Miocene (Hosner *et al.*, 2016). In contrast, speciation in the

Atlantic Forest remained low and dispersal out of this region decreased until the mid-Pleistocene.

The contraction of the Atlantic Forest since the mid-Miocene (Simon *et al.*, 2009; Edwards *et al.*,

2010; Strömberg, 2011; Werneck, 2011) may have restricted Brassolini speciation to montane

habitats in southeastern Brazil where most extant species occur, and old lineages having low

speciation rates through time survived in remnants of rainforest.

*FUTURE DIRECTIONS*

The evolutionary patterns inferred here indicate that geography has been an important

determinant of the diversity and distribution of extant Brassolini butterflies. Other unmeasured or

unknown traits might have driven the observed disparate trends in speciation through time, for

example larval hostplants. However, many Mesoamerican Brassolini feed on the same plant

families (e.g., Arecaceae), and even genera (e.g., *Euterpe*, *Bactris*), as do Atlantic Forest taxa

(Beccaloni *et al.*, 2008; Janzen *et al.*, 2009; Robinson *et al.*, 2010). Thus, despite the limited data

at hand, it seems that larval hostplants might have contributed little to the disparate

diversification dynamics of Brassolini.

We are aware that estimating diversification rates shifts through time is challenging (Moore *et*

*al.*, 2016). However, further evaluation via simulated data suggested that BAMM can infer

speciation rates accurately from extant-taxon phylogenies (Rabosky *et al.*, 2017). Although low

extinction rates might explain the museum of diversity pattern, we have conservatively focused

only on regional species origination rates to explain lineage accumulation within bioregions.

Furthermore, (Louca & Pennell, 2020) stressed that the identifiability of diversification processes

is problematic in a model-selection approach, but this remains to be evaluated in methods that

use rate shift models such as BAMM (Laudanno *et al.*, 2020). Reassuringly, we also used a

fundamentally distinct measure of within-area cladogenesis (Xing & Ree, 2017) that

corroborated the pattern of recent and rapid speciation in Mesoamerica + NW Andes, and an

older lineage accumulation and lower diversification in the Atlantic Forest. However, our

proposed checkered assemblage of Neotropical cradles and museums certainly needs to be

evaluated with other larger phylogenies of rainforest lineages, particularly in light of natural

history differences among various butterfly groups.

Our approach to disentangle regional speciation from dispersal to understand the origin of

within-area species diversity has two strengths. First, we defined bioregions that fit the

distribution and composition of Brassolini species by using geo-referenced occurrences, rather

than assuming any subjective criteria to adopt areas from hierarchical bioregion classifications

(Vilhena & Antonelli, 2015). This data-driven approach revealed a clustering of butterfly

communities from southern North America to Mesoamerica and the NW side of the Andes,

which conforms with the likely emergence of landmass in Central America, Chocó and

northeastern Colombia after the collision of the Panama Block and northwestern South America

by the late Oligocene (Coates & Stallard, 2013; Jaramillo, 2018). Furthermore, the inference of

large bioregions for Brassolini agrees with its apparent poor spatial structuring and high dispersal

capabilities. For instance, given a suitable environment, ancestral *Bia* had the capability to

disperse across pan-Amazonia in only 1,463–3,115 years (Penz *et al.*, 2015). The delimitation of

Mesoamerica + NW Andes and the Atlantic Forest as separate bioregions has likely been

determined by ecological and habitat suitability rather than geographical barriers to Brassolini

dispersal; this being a similar biogeographical pattern found in other Neotropical rainforest taxa

(Dexter *et al.*, 2017; Pérez-Escobar *et al.*, 2019).

**CONCLUSIONS**

We investigated whether the diversification and biogeographical history of butterflies classified

in the tribe Brassolini conforms to the premise that the Neotropics consists of a regional network

of museums and cradles of diversity (Fig. 5). We found that endemic species to Mesoamerica

plus the NW side of Andes have originated by old dispersal events into the region, and that

speciation rates increased only at about 2 Mya (cradle of diversity). In contrast, endemic species

to the Brazilian Atlantic Forest likely arose under low and continuous speciation rates since at

least 12 Mya (museum of diversity), and was likely intermittently connected to Amazonia since

the Miocene. The dynamic evolutionary history of Amazonian Brassolini lineages alternately

diversifying and accumulating old lineages appears to reflect paleoenvironmental changes,

including landscape reconfigurations in western Amazonia and paleoclimate fluctuations during

the Neogene and Quaternary. By focusing on a representative lineage of a major insect order, our

study paves the way for understanding why the Neotropics should be considered both a museum

and cradle of diversity.

**ACKNOWLEDGEMENTS**

We are grateful to Carlos Peña, Mirna Casagrande, Helena Romanowski, Augusto H. B. Rosa,

and Andrew Warren for contributing specimens used in this study; and to Diana Silva, Gerardo

Lamas, and SERFOR (Ministerio de Agricultura, Peru) for assistance with research permits (No

Autorización 223-2017-SERFOR/DGGSPFFS). We thank the anonymous reviewers for their

helpful comments. We are also grateful to the ICMBio for the research permits (SISBIO nº

10802-5, 10802-24 and 53016-10). Brazilian specimens are registered under the SISGEN

(A16771F). We acknowledge the computational resources provided by the CESNET

LM2015042 and the CERIT Scientific Cloud LM2015085, provided under the programme

"Projects of Large Research, Development, and Innovations Infrastructures". This publication is

part of the RedeLep 'Rede Nacional de Pesquisa e Conservação de Lepidópteros' SISBIOTA-

Brasil/CNPq (563332/2010-7). PMM was supported by a Marie Skłodowska-Curie fellowship

(European Commission, Grant No. MARIPOSAS-704035), by the Grant Agency of the Czech

Republic (GAČR grant: GJ20-18566Y), and by a fellowship from the PPLZ programme (Czech

Academy of Sciences, Fellowship No. L200961951). AVLF acknowledges support from

FAPESP (2011/50225-3, 2012/50260-6, 2013/50297-0), from the Brazilian Research Council -

CNPq (303834/2015-3), from the National Science Foundation (DEB-1256742) and from the

United States Agency for International Development - USAID / the U.S. National Academy of

Sciences (NAS), under the PEER program (Sponsor Grant Award Number: AID-OAA-A-11-

00012) (Mapping and Conserving Butterfly Biodiversity in the Brazilian Amazon). NW

acknowledges support from the Swedish Research Council. AA acknowledges support from the

Swedish Research Council, the Swedish Foundation for Strategic Research, the Knut and Alice

Wallenberg Foundation, and the Royal Botanic Gardens, Kew.

**REFERENCES**

**Albert JS, Val P & Hoorn C**. **2018**. The changing course of the Amazon River in the Neogene:

center stage for Neotropical diversification. *Neotropical Ichthyology* **16**: e180033.

**Antonelli A, Nylander J a a, Persson C & Sanmartín I**. **2009**. Tracing the impact of the

Andean uplift on Neotropical plant evolution. *Proceedings of the National Academy of Sciences*

**106**: 9749–9754.

**Antonelli A, Zizka A, Carvalho FA, Scharn R, Bacon CD, Silvestro D & Condamine FL**.

**2018a**. Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National*

*Academy of Sciences* **115**: 6034–6039.

**Antonelli A, Ariza M, Albert J, Andermann T, Azevedo J, Bacon C, Faurby S, Guedes T,**

**Hoorn C, Lohmann LG, Matos-Maraví P, Ritter CD, Sanmartín I, Silvestro D, Tejedor M,**

**ter Steege H, Tuomisto H, Werneck FP, Zizka A & Edwards S V. 2018b**. Conceptual and

empirical advances in Neotropical biodiversity research. *PeerJ* **6**: e5644.

**Antonelli A & Sanmartín I**. **2011**. Why are there so many plant species in the Neotropics?

*Taxon* **60**: 403–414.

**Austin GT, Warren AD, Penz CM, Llorente-Bousquets JE, Luis-Martínez A & Vargas-**

**Fernández I**. **2007**. A new species of *Opsiphanes* Doubleday, [1849] from western Mexico

(Nymphalidae: Morphinae: Brassolini). *Bulletin of the Allyn Museum* **150**: 1–20.

**Azevedo JAR, Collevatti RG, Jaramillo CA, Strömberg CAE, Guedes TB, Matos-Maraví**

**P, Bacon CD, Carillo JD, Faurby S & Antonelli A**. **2020**. On the young savannas in the land

of ancient forests. In: Rull V, Carnaval AC, eds. *Neotropical Diversification: Patterns and*

*Processes*. Cham, Switzerland: Springer Nature Switzerland AG 2020, 271–298.

**Bacon CD, Silvestro D, Jaramillo C, Smith BT, Chakrabarty P & Antonelli A**. **2015**.

Biological evidence supports an early and complex emergence of the Isthmus of Panama. *Proceedings of the National Academy of Sciences* **112**: 6110–6115.

**Basset Y, Barrios H, Segar S, Srygley RB, Aiello A, Warren AD, Delgado F, Coronado J, Lezcano J, ARizala S, Rivera M, Perez F, Bobadilla R, Lopez Y & Ramirez JA**. **2015**. The butterflies of Barro Colorado Island, Panama: local extinction since the 1930s. *PLoS ONE* **10**: e0136623.

**Batalha-Filho H, Fjeldså J, Fabre PH & Miyaki CY**. **2013**. Connections between the Atlantic and the Amazonian forest avifaunas represent distinct historical events. *Journal of Ornithology* **154**: 41–50.

**Beccaloni GW, Viloria ÁL, Hall SK & Robinson GS**. **2008**. *Catalogue of the hostplants of the Neotropical butterflies/Catálogo de las plantas huésped de las mariposas neotropicales.* Zaragoza, Spain: Sociedad de Entomología Aragonesa.

**Bermingham E & Dick C**. **2001**. The *Inga*—Newcomer or museum antiquity? *Science* **293**: 2214–2216.

**Bernal R, Bacon CD, Balslev H, Hoorn C, Bourlat SJ, Tuomisto H, Salamanca S, van Manen MT, Romero I, Sepulchre P & Antonelli A**. **2019**. Could coastal plants in western Amazonia be relicts of past marine incursions? *Journal of Biogeography* **46**: 1749–1759.

**Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, Suchard MA, Rambaut A & Drummond AJ**. **2014**. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Computational Biology* **10**: e1003537.

**Bouckaert RR & Drummond AJ**. **2017**. bModelTest: Bayesian phylogenetic site model averaging and model comparison. *BMC Evolutionary Biology* **17**: 42.

**Bristow CR**. **1981**. A revision of the brassoline genus *Catoblepia* (Lepidoptera: Rhopalocera).

*Zoological Journal of the Linnean Society* **72**: 117–163.

**Bristow CR**. **1982**. A revision of the brassoline genus *Selenophanes* (Lepidoptera: Rhopalocera).

*Zoological Journal of the Linnean Society* **76**: 273–291.

**Bristow CR**. **1991**. A revision of the brassoline genus *Opsiphanes* (Lepidoptera: Rhopalocera).

*Zoological Journal of the Linnean Society* **101**: 203–293.

**Capurucho JMG, Ashley M V., Ribas CC & Bates JM**. **2018**. Connecting Amazonian,

Cerrado, and Atlantic forest histories: Paraphyly, old divergences, and modern population

dynamics in tyrant-manakins (*Neopelma*/*Tyranneutes*, Aves: Pipridae). *Molecular Phylogenetics*

*and Evolution* **127**: 696–705.

**Casagrande MM**. **1982**. Quatro gêneros novos de Brassolinae (Lepidoptera-Nymphalidae).

*Revista Brasileira de Entomologia* **26**: 355–356.

**Casagrande MM**. **1995**. Notas sistemáticas sobre Brassolinae. I. Tribos (Lepidoptera,

Nymphalidae). *Revista Brasileira de Zoologia* **12**: 671–699.

**Casagrande MM**. **2002**. Naropini Stichel, taxonomia e imaturos (Lepidoptera, Nymphalidae,

Brassolinae). *Revista Brasileira de Zoologia* **19**: 467–569.

**Casagrande MM**. **2004**. Brassolini: Checklist. Part 4A. Hesperioidea - Papilionoidea. In:

Heppner JB, Lamas G, eds. *Atlas of Neotropical Lepidoptera*. Gainesville, Florida: Association

of Tropical Lepidoptera/Scientific publishers, 201–205.

**Chacón IA, Montero-Ramírez J, Janzen DH, Hallwachs W, Blandin P, Bristow CR &**

**Hajibabaei M**. **2012**. A new species of *Opsiphanes* Doubleday, [1849] from Costa Rica

(Nymphalidae: Morphinae: Brassolini), as revealed by its DNA barcodes and habitus. *Bulletin of*

*the Allyn Museum* **166**: 1–15.

**Chazot N, Willmott KR, Condamine FL, De-Silva DL, Freitas AVL, Lamas G, Morlon H,**

**Giraldo CE, Jiggins CD, Joron M, Mallet J, Uribe S & Elias M**. **2016**. Into the Andes:

multiple independent colonizations drive montane diversity in the Neotropical clearwing

butterflies Godyridina. *Molecular Ecology* **25**: 5765–5784.

**Chazot N, Wahlberg N, Freitas AVL, Mitter C, Labandeira C, Sohn JC, Sahoo RK,**

**Seraphim N, de Jong R & Heikkilä M**. **2019a**. Priors and posteriors in Bayesian timing of

divergence analyses: the age of butterflies revisited. *Systematic Biology* **68**: 797–813.

**Chazot N, Willmott KR, Lamas G, Freitas AVL, Piron-Prunier F, Arias CF, Mallet J, De-**

**Silva DL & Elias M**. **2019b**. Renewed diversification following Miocene landscape turnover in

a Neotropical butterfly radiation. *Global Ecology and Biogeography* **28**: 1118–1132.

**Coates AG, Collins LS, Aubury MP & Berggren WA**. **2004**. The geology of the Darien,

Panama, and the late Miocene-Pliocene collision of the Panama arc with northwestern South

America. *Bulletin of the Geological Society of America* **116**: 1327–1344.

**Coates AG & Stallard RF**. **2013**. How old is the Isthmus of Panama? *Bulletin of Marine*

*Science* **89**: 801–813.

**Condamine FL, Silva-Brandão KL, Kergoat GJ & Sperling FAH**. **2012**. Biogeographic and

diversification patterns of Neotropical Troidini butterflies (Papilionidae) support a museum

model of diversity dynamics for Amazonia. *BMC Evolutionary Biology* **12**: 82.

**Costa LP**. **2003**. The historical bridge between the Amazon and the Atlantic Forest of Brazil: a

study of molecular phylogeography with small mammals. *Journal of Biogeography* **30**: 71–86.

**De-Silva DL, Elias M, Willmott K, Mallet J & Day JJ**. **2016**. Diversification of clearwing

butterflies with the rise of the Andes. *Journal of Biogeography* **43**: 44–58.

**De-Silva LD, Mota LL, Chazot N, Mallarino R, Silva-Brandão KL, Piñerez LMG, Freitas**

**AVL, Lamas G, Joron M, Mallet J, Giraldo CE, Uribe S, Särkinen T, Knapp S, Jiggins CD,**

**Willmott KR & Elias M**. **2017**. North Andean origin and diversification of the largest ithomiine

butterfly genus. *Scientific Reports* **7**: 45966.

**DeVries PJ**. **1983**. Checklist of butterflies and natural history. In: Janzen DH, ed. *Costa Rican*

*Natural History*. Chicago, USA: University of Chicago Press, 654–678.

**DeVries PJ**. **1994**. Patterns of butterfly diversity and promising topics in natural history and

ecology. In: McDade LA, Bawa KS, Hespenheide HA, Hartshorn GS, eds. *La Selva: Ecology*

*and Natural History of a Neotropical Rain Forest*. Chicago, USA: The University of Chicago

Press, 187–194.

**Dexter KG, Lavin M, Torke BM, Twyford AD, Kursar TA, Coley PD, Drake C, Hollands R**

**& Pennington RT**. **2017**. Dispersal assembly of rain forest tree communities across the Amazon

basin. *Proceedings of the National Academy of Sciences* **114**: 2645–2650.

**Dickens JK, McMahon L & Binnie SE**. **2019**. The butterflies of a Cerrado–Atlantic Forest

ecotone at Laguna Blanca reveal underestimation of Paraguayan butterfly diversity and need for

conservation. *Journal of Insect Conservation* **23**: 707–728.

**Dupin J, Matzke NJ, Särkinen T, Knapp S, Olmstead RG, Bohs L & Smith SD**. **2017**.

Bayesian estimation of the global biogeographical history of the Solanaceae. *Journal of*

*Biogeography* **44**: 887–899.

**Edler D, Guedes T, Zizka A, Rosvall M & Antonelli A**. **2017**. Infomap Bioregions: interactive

mapping of biogeographical regions from species distributions. *Systematic Biology* **66**: 197–204.

**Edwards EJ, Osborne CP, Strömberg CAE, Smith SA, Bond WJ, Christin PA, Cousins**

**AB, Duvall MR, Fox DL, Freckleton RP, Ghannoum O, Hartwell J, Huang Y, Janis CM,**

**Keeley JE, Kellogg EA, Knapp AK, Leakey ADB, Nelson DM, Saarela JM, Sage RF, Sala**

**OE, Salamin N, Still CJ & Tipple B**. **2010**. The origins of $C_4$ grasslands: integrating

evolutionary and ecosystem science. *Science* **328**: 587–591.

**Elias M, Joron M, Willmott K, Silva-Brandão KL, Kaiser V, Arias CF, Gomez Piñerez**

**LM, Uribe S, Brower AVZ, Freitas AVL & Jiggins CD**. **2009**. Out of the Andes: patterns of

diversification in clearwing butterflies. *Molecular Ecology* **18**: 1716–1729.

**Emery E de O, Brown Jr KS & Pinheiro CEG**. **2006**. As borboletas (Lepidoptera,

Papilionoidea) do Distrito Federal, Brasil. *Revista Brasileira de Entomologia* **50**: 85–92.

**Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, Maunsell SC,**

**Aduse-Poku K, Talavera G, Eastwood R, Jarzyna MA, Guralnick R, Lohman DJ, Pierce**

**NE & Kawahara AY**. **2018**. A comprehensive and dated phylogenomic analysis of butterflies.

*Current Biology* **28**: 770–778.

**Faurby S, Eiserhardt WL & Svenning JC**. **2016**. Strong effects of variation in taxonomic

opinion on diversification analyses. *Methods in Ecology and Evolution* **7**: 4–13.

**FitzJohn RG, Maddison WP & Otto SP**. **2009**. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. *Systematic Biology* **58**: 595–611.

**Freitas AVL & Brown KSJ**. **2004**. Phylogeny of the Nymphalidae (Lepidoptera). *Systematic*

*Biology* **53**: 363–383.

**Furtado E & Campos-Neto FC**. **2004**. *Caligopsis seleucida* (Hewitson) e seus estágios

imaturos (Lepidoptera, Nymphalidae, Brassolinae). *Revista Brasileira de Zoologia* **21**: 593–597.

**Garzón-Orduña IJ, Benetti-Longhini JE & Brower AVZ**. **2014**. Timing the diversification of

the Amazonian biota: butterfly divergences are consistent with Pleistocene refugia (B Riddle,

Ed.). *Journal of Biogeography* **41**: 1631–1638.

**Garzón-Orduña IJ & Penz CM**. **2009**. Phylogeny of *Dynastor* and *Brassolis* butterflies

(Lepidoptera: Nymphalidae): a tough nut to crack. *Zootaxa* **2134**: 1–22.

**Gaston KJ & Blackburn TM**. **1996**. The tropics as a museum of biological diversity: an

analysis of the New World avifauna. *Proceedings of the Royal Society B: Biological Sciences*

**263**: 63–68.

**Goldberg EE, Roy K, Lande R & Jablonski D**. **2005**. Diversity, endemism, and age

distributions in macroevolutionary sources and sinks. *The American Naturalist* **165**: 623–633.

**Goloboff PA & Catalano SA**. **2016**. TNT version 1.5, including a full implementation of

phylogenetic morphometrics. *Cladistics* **32**: 221–238.

**Hoorn C, Wesselingh FP, ter Steege H, Bermudez MA, Mora A, Sevink J, Sanmartín I,**

**Sanchez-Meseguer A, Anderson CL, Figueiredo JP, Jaramillo C, Riff D, Negri FR,**

**Hooghiemstra H, Lundberg J, Stadler T, Sarkinen T & Antonelli A**. **2010**. Amazonia

through time: andean uplift, climate change, landscape evolution, and biodiversity. *Science* **330**:

927–931.

**Hosner PA, Braun EL & Kimball RT**. **2016**. Rapid and recent diversification of curassows,

guans, and chachalacas (Galliformes: Cracidae) out of Mesoamerica: phylogeny inferred from

mitochondrial, intron, and ultraconserved element sequences. *Molecular Phylogenetics and*

*Evolution* **102**: 320–330.

**Huelsenbeck JP, Larget B & Alfaro ME**. **2004**. Bayesian phylogenetic model selection using

reversible jump Markov chain Monte Carlo. *Molecular Biology and Evolution* **21**: 1123–1133.

**Igea J & Tanentzap AJ**. **2019**. Multiple macroevolutionary routes to becoming a biodiversity

hotspot. *Science Advances* **5**: eaau8067.

**Jablonski D, Roy K & Valentine JW**. **2006**. Out of the tropics: evolutionary diversity gradient.

*Science* **314**: 102–106.

**Janzen DH, Hallwachs W, Blandin P, Burns JM, Cadiou JM, Chacon I, Dapkey T, Deans**

**AR, Epstein ME, Espinoza B, Franclemont JG, Haber WA, Hajibabaei M, Hall JPW, Hebert PDN, Gauld ID, Harvey DJ, Hausmann A, Kitching IJ, Lafontaine D, Landry JFÇ, Lemaire C, Miller JY, Miller JS, Miller L, Miller SE, Montero J, Munroe E, Green SR, Ratnasingham S, Rawlins JE, Robbins RK, Rodriguez JJ, Rougerie R, Sharkey MJ, Smith MA, Solis MA, Sullivan JB, Thiaucourt P, Wahl DB, Weller SJ, Whitfield JB, Willmott KR, Wood DM, Woodley NE & Wilson JJ**. **2009**. Integration of DNA barcoding into an ongoing inventory of complex tropical biodiversity. *Molecular Ecology Resources* **9**: 1–26.

**Janzen DH & Hallwachs W**. **2009**. Dynamic database for an inventory of the macrocaterpillar fauna, and its food plants and parasitoids, of Area de Conservacion Guanacaste (ACG), northwestern Costa Rica. <http://janzen.sas.upenn.edu>.

**Jaramillo C, Romero I, D'Apolito C, Bayona G, Duarte E, Louwye S, Escobar J, Luque J, Carrillo-Briceño JD, Zapata V, Mora A, Schouten S, Zavada M, Harrington G, Ortiz J & Wesselingh FP**. **2017a**. Miocene flooding events of western Amazonia. *Sciences Advances* **3**: e1601693.

**Jaramillo C, Montes C, Cardona A, Silvestro D, Antonelli A & Bacon CD**. **2017b**. Comment (1) on "Formation of the Isthmus of Panama" by O'Dea *et al*. *Science Advances* **3**: e1602321.

**Jaramillo C**. **2018**. Evolution of the Isthmus of Panama: Biological, paleoceanographic and paleoclimatological implications. In: Hoorn C, Perrigo A, Antonelli A, eds. *Mountains, Climate and Biodiversity*. West Sussex, UK: John Wiley & Sons, Ltd, 323–338.

**Jaramillo C & Cárdenas A**. **2013**. Global warming and Neotropical rainforests: a historical perspective. *Annual Review of Earth and Planetary Sciences* **41**: 741–766.

**Laudanno G, Haegeman B, Rabosky DL & Etienne RS**. **2020**. Detecting lineage-specific shifts in diversification: A proper likelihood approach. *Systematic Biology*: syaa048.

**Ledo RMD & Colli GR**. **2017**. The historical connections between the Amazon and the Atlantic

Forest revisited. *Journal of Biogeography* **44**: 2551–2563.

**Lewis PO**. **2001**. A likelihood approach to estimating phylogeny from discrete morphological

character data. *Systematic Biology* **50**: 913–925.

**Louca S & Pennell MW**. **2020**. Extant timetrees are consistent with a myriad of diversification

histories. *Nature* **580**: 502–505.

**Löwenberg-Neto P**. **2014**. Neotropical region: a shapefile of Morrone's (2014) biogeographical

regionalisation. *Zootaxa* **3802**: 300.

**Machado AFP, Rønsted N, Bruun-Lund S, Pereira RAS & de Queiroz LP**. **2018**. Atlantic

forests to the all Americas: Biogeographical history and divergence times of Neotropical *Ficus*

(Moraceae). *Molecular Phylogenetics and Evolution* **122**: 46–58.

**Matos-Maraví P, Peña C, Willmott KR, Freitas AVL & Wahlberg N**. **2013**. Systematics and

evolutionary history of butterflies in the '*Taygetis* clade' (Nymphalidae: Satyrinae: Euptychiina):

Towards a better understanding of Neotropical biogeography. *Molecular Phylogenetics and*

*Evolution* **66**: 54–68.

**Matos-Maraví P**. **2016**. Investigating the timing of origin and evolutionary processes shaping

regional species diversity: Insights from simulated data and Neotropical butterfly diversification

rates. *Evolution* **70**: 1638–1650.

**Matzke NJ**. **2013**. Probabilistic historical biogeography: new models for founder-event

speciation, imperfect detection, and fossils allow improved accuracy and model-testing.

*Frontiers of Biogeography* **5**: 242–248.

**McKenna DD & Farrell BD**. **2006**. Tropical forests are both evolutionary cradles and museums

of leaf beetle diversity. *Proceedings of the National Academy of Sciences* **103**: 10947–10951.

**Melo DHA, Duarte M, Mielke OHH, Robbins RK & Freitas AVL**. **2019**. Butterflies

(Lepidoptera: Papilionoidea) of an urban park in northeastern Brazil. *Biota Neotropica* **19**:

e20180614.

**Miller MA, Pfeiffer W & Schwartz T**. **2010**. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. *Proceedings of the Gateway Computing Environments*

*Workshop (GCE), 14 Nov.*: 1–8.

**Montes C, Bayona G, Cardona A, Buchs DM, Silva CA, Morón S, Hoyos N, Ramírez DA,**

**Jaramillo CA & Valencia V**. **2012**. Arc-continent collision and orocline formation: closing of

the Central American seaway. *Journal of Geophysical Research* **117**: B04105.

**Montes C, Cardona A, Jaramillo C, Pardo A, Silva JC, Valencia V, Ayala C, Pérez-Angel**

**LC, Rodriguez-Parra LA, Ramirez V & Niño H**. **2015**. Middle Miocene closure of the Central

American Seaway. *Science* **348**: 226–229.

**Moore BR, Höhna S, May MR, Rannala B & Huelsenbeck JP**. **2016**. Critically evaluating the

theory and performance of Bayesian analysis of macroevolutionary mixtures. *Proceedings of the*

*National Academy of Sciences* **113**: 9569–9574.

**Moreau CS & Bell CD**. **2013**. Testing the museum versus cradle tropical biological diversity

hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of the ants.

*Evolution* **67**: 2240–2257.

**Moritz C, Patton JL, Schneider CJ & Smith TB**. **2000**. Diversification of rainforest faunas:

An integrated molecular approach. *Annual Review of Ecology and Systematics* **31**: 533–563.

**Morrone JJ**. **2014a**. Cladistic biogeography of the Neotropical region: identifying the main

events in the diversification of the terrestrial biota. *Cladistics* **30**: 202–214.

**Morrone JJ**. **2014b**. Biogeographical regionalisation of the Neotropical region. *Zootaxa* **3782**:

1–110.

**Mullen SP, Savage WK, Wahlberg N & Willmott KR**. **2011**. Rapid diversification and not

clade age explains high diversity in neotropical *Adelpha* butterflies. *Proceedings of the Royal*

*Society B: Biological Sciences* **278**: 1777–1785.

**Musher LJ, Ferreira M, Auerbach AL, McKay J & Cracraft J**. **2019**. Why is Amazonia a

'source' of biodiversity? Climate-mediated dispersal and synchronous speciation across the

Andes in an avian group (Tityrinae). *Proceedings of the Royal Society B: Biological Sciences*

**286**: 20182343.

**Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB & Kent J**. **2000**. Biodiversity

hotspots for conservation priorities. *Nature* **403**: 853–858.

**O'Dea A, Lessios HA, Coates AG, Eytan RI, Restrepo-Moreno SA, Cione AL, Collins LS,**

**de Queiroz A, Farris DW, Norris RD, Stallard RF, Woodburne MO, Aguilera O, Aubry**

**MP, Berggren WA, Budd AF, Cozzuol MA, Coppard SE, Duque-Caro H, Finnegan S,**

**Gasparini GM, Grossman EL, Johnson KG, Keigwin LD, Knowlton N, Leigh EG,**

**Leonard-Pingel JS, Marko PB, Pyenson ND, Rachello-Dolmen PG, Soibelzon E, Soibelzon**

**L, Todd JA, Vermeij GJ & Jackson JBC**. **2016**. Formation of the Isthmus of Panama. *Science*

*Advances* **2**: e1600883.

**Ogilvie HA, Bouckaert RR & Drummond AJ**. **2017**. StarBEAST2 brings faster species tree

inference and accurate estimates of substitution rates. *Molecular Biology and Evolution* **34**:

2101–2114.

**Peña C, Nylin S, Freitas AVL & Wahlberg N**. **2010**. Biogeographic history of the butterfly

subtribe Euptychiina (Lepidoptera, Nymphalidae, Satyrinae). *Zoologica Scripta* **39**: 243–258.

**Penz CM**. **2007**. Evaluating the monophyly and phylogenetic relationships of Brassolini genera

(Lepidoptera, Nymphalidae). *Systematic Entomology* **32**: 668–689.

**Penz CM**. **2008**. Phylogenetic revision of *Eryphanis* Boisduval, with a description of a new

species from Ecuador (Lepidoptera, Nymphalidae). *Insecta Mundi* **0035**: 1–25.

**Penz CM**. **2009a**. Phylogeny of *Dasyophthalma* butterflies (Lepidoptera, Nymphalidae,

Brassolini). *Insecta Mundi* **0069**: 1–12.

**Penz CM**. **2009b**. The phylogeny of *Opoptera* butterflies, and an assessment of the systematic

position of *O. staudingeri* (Lepidoptera, Nymphalidae). *Zootaxa* **1985**: 1–20.

**Penz CM, Freitas AVL, Kaminski LA, Casagrande MM & Devries PJ**. **2013**. Adult and

early-stage characters of Brassolini contain conflicting phylogenetic signal (Lepidoptera,

Nymphalidae). *Systematic Entomology* **38**: 316–333.

**Penz C, DeVries P, Tufto J & Lande R**. **2015**. Butterfly dispersal across Amazonia and its

implication for biogeography. *Ecography* **38**: 410–418.

**Penz CM, Casagrande MM, Devries P & Simonsen TJ**. **2017**. Documenting diversity in the

Amazonian butterfly genus *Bia* (Lepidoptera, Nymphalidae). *Zootaxa* **4258**: 201–237.

**Penz CM, Mohammadi N & Wahlberg N**. **2011a**. Neotropical *Blepolenis* butterflies: wing

pattern elements, phylogeny, and Pleistocene diversification (Lepidoptera, Nymphalidae).

*Zootaxa* **2897**: 1–17.

**Penz CM, Simonsen TJ & DeVries P**. **2011b**. A new *Orobrassolis* butterfly (Nymphalidae,

Brassolini): a casualty of habitat destruction? *Zootaxa* **2740**: 35–43.

**Pereira Martins L, da Costa Araujo Junior E, Pereira Martins AR, Santos Colins M,**

**Fonseca Almeida GC & Garcia Azevedo G**. **2017**. Butterflies of Amazon and Cerrado

remnants of Maranhão, Northeast Brazil. *Biota Neotropica* **17**: e20170335.

**Pérez-Escobar OA, Lucas E, Jaramillo C, Monro A, Morris SK, Bogarín D, Greer D,**

**Dodsworth S, Cano JA, Sanchez Meseguer A & Antonelli A**. **2019**. The origin and

diversification of the hyperdiverse flora in the Chocó biogeographic region. *Frontiers in Plant*

*Science* **10**: 1328.

**Pérez JH, Gaviria-Ortiz FG, Santos WIG, Carneiro E, Mielke OHH & Casagrande MM**.

**2017**. Long term survey of the butterfly fauna of Curitiba, Paraná, Brazil: how does a scientific

collection gather local biodiversity information? (Lepidoptera: Papilionoidea). *SHILAP Revista*

*de Lepidopterología* **45**: 433–446.

**Pinheiro CEG & Emery E de O**. **2006**. As borboletas (Lepidoptera: Papilionoidea e

Hesperioidea) da área de Proteção Ambiental do Gama e Cabeça de Veado (Distrito Federal,

Brasil). *Biota Neotropica* **6**: bn01506032006.

**Prates I, Rivera D, Rodrigues MT & Carnaval AC**. **2016**. A mid-Pleistocene rainforest

corridor enabled synchronous invasions of the Atlantic Forest by Amazonian anole lizards.

*Molecular Ecology* **25**: 5174–5186.

**R Core Team**. **2020**. R: A language and environment for statistical computing. *R Foundation for*

*Statistical Computing*: Version 4.0.0.

**Rabosky DL**. **2014**. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PLoS ONE* **9**: e89543.

**Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, Huang H & Larson JG**.

**2014**. BAMMtools: An R package for the analysis of evolutionary dynamics on phylogenetic

trees. *Methods in Ecology and Evolution* **5**: 701–707.

**Rabosky DL, Mitchell JS & Chang J**. **2017**. Is BAMM flawed? Theoretical and practical

concerns in the analysis of multi-rate diversification. *Systematic Biology* **66**: 477–498.

**Rambaut A, Drummond AJ, Xie D, Baele G & Suchard MA**. **2018**. Posterior summarization

in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**: 901–904.

**Ree RH, Moore BR, Webb CO & Donoghue MJ**. **2005**. A likelihood framework for inferring

the evolution of geographic range on phylogenetic trees. *Evolution* **59**: 2299–2311.

**Ribas CC, Aleixo A, Nogueira ACR, Miyaki CY & Cracraft J**. **2012**. A palaeobiogeographic

model for biotic diversification within Amazonia over the past three million years. *Proceedings*

*of the Royal Society B: Biological Sciences* **279**: 681–689.

**Richardson JE, Pennington RT, Pennington TD & Hollingsworth PM**. **2001**. Rapid

diversification of a species-rich genus of neotropical rain forest trees. *Science* **293**: 2242–2245.

**Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW & Hernandez LM**. **2010**. HOSTS - A

Database of the World's Lepidopteran Hostplants. Natural History Museum, London.

<http://www.nhm.ac.uk/hosts>. : http://www.nhm.ac.uk/hosts.

**Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L,**

**Suchard MA & Huelsenbeck JP**. **2012**. MrBayes 3.2: efficient Bayesian phylogenetic inference

and model choice across a large model space. *Systematic Biology* **61**: 539–542.

**Rosa BB, Melo GAR & Barbeitos MS**. **2019**. Homoplasy-based partitioning outperforms

alternatives in Bayesian analysis of discrete morphological data. *Systematic Biology* **68**: 657–

671.

**Roy K & Goldberg EE**. **2007**. Origination, extinction, and dispersal: integrative models for

understanding present-day diversity gradients. *The American Naturalist* **170**: S71–S85.

**Santos JP dos, Agra Iserhard C, Oliveira Teixeira M & Piccoli Romanowski H**. **2011**. Fruit-

feeding butterflies guide of subtropical Atlantic Forest and Araucaria Moist Forest in State of

Rio Grande do Sul, Brazil. *Biota Neotropica* **11**: 253–274.

**Santos JP dos, Freitas AVL, Brown KS, Carreira JYO, Gueratto PE, Rosa AHB, Lourenço**

**GM, Accacio GM, Uehara-Prado M, Iserhard CA, Richter A, Gawlinski K, Romanowski**

**HP, Mega NO, Teixeira MO, Moser A, Ribeiro DB, Araujo PF, Filgueiras BKC, Melo**

**DHA, Leal IR, Beirão M do V, Ribeiro SP, Cambuí ECB, Vasconcelos RN, Cardoso MZ,**

**Paluch M, Greve RR, Voltolini JC, Galetti M, Regolin AL, Sobral-Souza T & Ribeiro MC**.

**2018**. Atlantic butterflies: a data set of fruit-feeding butterfly communities from the Atlantic

forests. *Ecology* **99**: 2875.

**Shirai LT, Barbosa EP, Greve RR, Magaldi LM, Nascimento AR & Freitas AVL**. **2017**.

Natural history of *Selenophanes cassiope guarany* (Lepidoptera: Nymphalidae: Brassolini): an

integrative approach, from molecules to ecology. *Annals of the Entomological Society of*

*America* **110**: 145–159.

**Silva ARM, Castro CO de, Mafia PO, Mendonça MOC, Alves TCC & Beirão M do V**.

**2012**. Borboletas frugívoras (Lepidoptera: Nymphalidae) de uma área urbana (Área de Proteção

Especial Manancial Cercadinho) em Belo Horizonte, Minas Gerais, Brasil. *Biota Neotropica* **12**:

292–297.

**Simon MF, Grether R, de Queiroz LP, Skema C, Pennington RT & Hughes CE**. **2009**.

Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of

adaptations to fire. *Proceedings of the National Academy of Sciences* **106**: 20359–20364.

**Soldati D, Silveira FA da & Silva ARM**. **2019**. Butterfly fauna (Lepidoptera, Papilionoidea) in

a heterogeneous area between two biodiversity hotspots in Minas Gerais, Brazil. *Papéis Avulsos*

*de Zoologia* **59**: e20195902.

**Stebbins GL**. **1974**. *Flowering Plants: Evolution above the Species Level*. Cambridge, MA:

Harvard University Press.

**Stichel H**. **1916**. Neues über Genus *Narope* Westw. *Neue Beiträge zur Systematischen*

*Insektenkunde* **1**: 7–8.

**Strömberg CAE**. **2011**. Evolution of grasses and grassland ecosystems. *Annual Review of Earth*

*and Planetary Sciences* **39**: 517–544.

**Thomé MTC, Sequeira F, Brusquetti F, Carstens B, Haddad CFB, Rodrigues MT &**

**Alexandrino J**. **2016**. Recurrent connections between Amazon and Atlantic forests shaped

diversity in Caatinga four-eyed frogs. *Journal of Biogeography* **43**: 1045–1056.

**Toussaint EFA, Dias FMS, Mielke OHH, Casagrande MM, Sañudo-Restrepo CP, Lam A,**

**Morinière J, Balke M & Vila R**. **2019**. Flight over the Proto-Caribbean seaway: Phylogeny and

macroevolution of Neotropical Anaeini leafwing butterflies. *Molecular Phylogenetics and*

*Evolution* **137**: 86–103.

**Trujillo-Arias N, Dantas GPM, Arbeláez-Cortés E, Naoki K, Gómez MI, Santos FR,**

**Miyaki CY, Aleixo A, Tubaro PL & Cabanne GS**. **2017**. The niche and phylogeography of a

passerine reveal the history of biological diversification between the Andean and the Atlantic

forests. *Molecular Phylogenetics and Evolution* **112**: 107–121.

**Tufto J, Lande R, Ringsby TH, Engen S, Sæther BE, Walla TR & DeVries PJ**. **2012**.

Estimating Brownian motion dispersal rate, longevity and population density from spatially

explicit mark-recapture data on tropical butterflies. *Journal of Animal Ecology* **81**: 756–769.

**Vilhena DA & Antonelli A**. **2015**. A network approach for identifying and delimiting

biogeographical regions. *Nature Communications* **6**: 6848.

**Wahlberg N, Leneveu J, Kodandaramaiah U, Peña C, Nylin S, Freitas AVL & Brower**

**AVZ**. **2009**. Nymphalid butterflies diversify following near demise at the Cretaceous/Tertiary

boundary. *Proceedings of the Royal Society B: Biological Sciences* **276**: 4295–4302.

**Wahlberg N & Wheat CW**. **2008**. Genomic outposts serve the phylogenomic pioneers:

designing novel nuclear markers for genomic DNA extractions of Lepidoptera. *Systematic*

*Biology* **57**: 231–242.

**Werneck FP**. **2011**. The diversification of eastern South American open vegetation biomes:

historical biogeography and perspectives. *Quaternary Science Reviews* **30**: 1630–1648.

**Wesselingh FP & Salo JA**. **2006**. A Miocene perspective on the evolution of the Amazonian

biota. *Scripta Geologica* **133**: 439–458.

**Wiens JJ & Donoghue MJ**. **2004**. Historical biogeography, ecology and species richness.

*Trends in Ecology and Evolution* **19**: 639–644.

**Xing Y & Ree RH**. **2017**. Uplift-driven diversification in the Hengduan Mountains, a temperate

biodiversity hotspot. *Proceedings of the National Academy of Sciences* **114**: E3444–E3451.

**Zacca T & Bravo F**. **2012**. Borboletas (Lepidoptera: Papilionoidea e Hesperioidea) da porção

norte da Chapada Diamantina, Bahia, Brasil. *Biota Neotropica* **12**: 117–126.

**Zizka A, Silvestro D, Andermann T, Azevedo J, Duarte Ritter C, Edler D, Farooq H,**

**Herdean A, Ariza M, Scharn R, Svantesson S, Wengström N, Zizka V & Antonelli A**. **2019**.

CoordinateCleaner: Standardized cleaning of occurrence records from biological collection

databases. *Methods in Ecology and Evolution* **10**: 744–751.

**SUPPORTING INFORMATION**

Additional Supporting Information may be found in the online version of this article at the

publisher's website:

**Supplementary data analyses** (pp. 2–12).

**Figure S1.** Map with the clean and flagged Brassolini geo-referenced occurrences (pp. 13).

**Figure S2.** Consensus trees of molecular and morphological datasets (pp. 14–22).

**Figure S3.** Consensus tree of concatenated total-evidence dataset (pp. 23–24).

**Figure S4.** Parsimony-based partitioned Bremer support scores (pp. 25–26).

**Figure S5.** Time-calibrated tree using the multispecies coalescent (pp. 27–28).

**Figure S6.** Brassolini taxonomic resolution across the Neotropics (pp. 29–30).

**Figure S7.** Ancestral range probabilities plotted on species tree (pp. 31–35).

**Figure S8.** Dispersal rate through time between bioregions (pp. 36–38).

**Figure S9.** Within-area cladogenesis events through time (pp. 39–41).

**Table S1.** Voucher locality information and associated genetic data (pp. 42, separate file).

**Table S2.** Best-fit partitioning scheme for the molecular dataset (pp. 42).

**Table S3.** Best-fit partitioning scheme for the morphological dataset (pp. 43–44).

**Table S4.** Bayes factor between the strict and relaxed clock models (pp. 45).

**Table S5.** Bayes factor among clock partitions and tree models (pp. 46).

**Table S6.** Tree topology test of early divergent Brassolini lineages (pp. 47).

**Table S7.** Sampling fractions for taking into account missing species (pp. 48).

 **Table and figures**

1016   Table 1: Secondary calibration points for divergence time estimation of Brassolini using BEAST

1017   v.2.6.3. We constrained the monophyly of representative taxa within the butterfly subfamily

1018   Satyrinae and used uniform age distributions based on Chazot *et al*. (2019a).

1019

| MRCA (monophyletic) | Sampled taxa | Lower | Upper |
|---|---|---|---|
| Brassolini (Ingroup) - Morphini | Morphini: *Antirrhea*, *Morpho*, *Caerois* | 34 | 50 |
| Lethina - Parargina - Mycalesina | *Lethe*, *Pararge*, *Bicyclus* | 31 | 46 |
| Pronophilina - Euptychiina - Satyrina - Erebiina | *Calisto*, *Manerebia*, *Taygetis*, *Satyrus*, *Erebia* | 31 | 44 |
| Satyrini | Lethina, Parargina, Mycalesina, Pronophilina, Euptychiina, Satyrina, Erebiina, *Orsotriaena*, *Oressinoma* | 38 | 55 |
| Melanitini - Dirini | *Melanitis*, *Aeropetes* | 27 | 41 |
| Satyrinae (Root) | Brassolini, Morphini, Satyrini, Melanitini, Dirini, *Cithaerias*, *Pierella*, *Zeuxidia*, *Elymnias* | 46 | 65 |

1020

1021

1022   Table 2: Characteristics of the multi-locus dataset. Specimens, species, and the molecular

1023   composition of the data matrix include outgroup taxa. (bp): base pairs; (Pis): number of

1024   parsimony-informative sites.

1025

| Genes | Specimens | Species (%) | Length (bp) | Variable sites (Pis) | Missing data (%) | GC content (%) |
|---|---|---|---|---|---|---|
| CAD | 43 | 35 (42%) | 850 | 369 (291) | 22.1 | 32.7 |
| COI | 107 | 82 (99%) | 1475 | 668 (542) | 14.4 | 29.7 |
| EF1α | 87 | 72 (87%) | 1240 | 451 (355) | 12.9 | 48.5 |
| GAPDH | 51 | 47 (57%) | 691 | 271 (232) | 6.8 | 45.4 |

| | | | | | | |
|---|---|---|---|---|---|---|
| RpS5 | 86 | 70 (84%) | 617 | 245 (216) | 4.0 | 46.2 |
| *wingless* | 87 | 70 (84%) | 412 | 189 (149) | 7.8 | 55.3 |
| TOTAL | 109 | 83 (100%) | 5285 | 2193 (1785) | 35.0 | 40.5 |

1026

1027

Fig. 1: Map of the Neotropics and the defined bioregions based on occurrence data of Brassolini.

Inset map shows the most stable bioregions under different cluster cost values by the software

Infomap Bioregions using validated geo-coordinates from GBIF and ATLANTIC

BUTTERFLIES databases. Note that the delineated bioregions approximate the Dominions of

Morrone (2014b), encompassing the Mesoamerican plus Pacific dominions, except for three

provinces east of the northern Andes; the Paraná dominion (Brazilian Atlantic Forest); the

remaining dominions and provinces where Brassolini occurs. Shapefile from Löwenberg-Neto

(2014).

1036

Fig. 2: Time-calibrated species tree of the butterfly tribe Brassolini based on morphological and molecular data. The posterior probabilities for all genera and among genera are displayed next to the crown node. Ancestral range probability based on the DEC model and 10,000 biogeographical stochastic mappings is depicted as a pie chart following colors in legend. Images of representative Brassolini butterflies, all at the same scale: From top to down, *Bia actorion*

(DeVries collection, PJD), *Dynastor darius* (Milwaukee Public Museum, MPM),

*Dasyophthalma creusa* (MPM), *Opoptera fruhstorferi* (American Museum of the Natural

History, AMNH), *Narope panniculus* (AMNH), *Brassolis astyra* (MPM), *Caligopsis seleucida*

(PJD), *Eryphanis aesacus* (MPM), *Caligo martia* (MPM), *Penetes pamphanis* (MPM),

*Selenophanes cassiope* (PJD), *Catoblepia berecynthia* (MPM), *Mielkella singularis* (AMNH),

*Orobrassolis ornamentalis* (MPM), *Blepolenis bassus* (Museu de Zoologia - USP), *Opsiphanes*

*quiteria* (PJD).

1051

1052

1053

1054     Fig. 3: Colonization rates through-time based on 10,000 biogeographical stochastic mappings in

1055     BioGeoBEARS. Rates are displayed for a select pair of areas. Source areas (dispersal from)

1056     follow the color in the legend. Solid lines are the median values, colored ribbons are the lower

1057     and upper quartiles (0.25 and 0.75 quantiles), and dashed lines are the 0.1 and 0.9 quantiles.

1058

1059

1060

Fig. 4: Within-area speciation through-time based on 100 posterior phylogenies and 100

biogeographical stochastic mappings (BSMs) in BioGeoBEARS. A) The relative number of

within-area cladogenesis through time was calculated using a formula modified from Xing &

Ree (2017) based on cladogenesis events occurring in one area. Solid lines represent median

values, colored ribbons represent lower and upper quartiles (0.25 and 0.75 quantiles) and dashed

lines represent 0.1 and 0.9 quantiles. B) Per-lineage speciation rates through time estimated by

BAMM were assigned to a given bioregion based on the BSMs. Solid lines represent mean

speciation rates per lineage and the colored ribbons the 5% and 95% confidence intervals.

1069

1070

1071

Fig. 5: Summary of Brassolini colonization and speciation estimates in three time-windows, at

24, 10 and 2 Mya. Colored directional arrows represent dispersal from source areas, following

colors in legend. The width and shape of arrows represent the estimated relative colonization rate

between two areas in BioGeoBEARS, as depicted in Fig. 3. Bioregions were colored to

qualitatively represent the estimated relative cladogenesis/speciation rates, as either center of

speciation or area with comparatively lower speciation rates per time window.

1078

1079

1080