Title page

Title: Mesoamerica is a cradle and the Brazilian Atlantic Forest is a museum of Neotropical

butterfly diversity (Lepidoptera: Nymphalidae: Brassolini)

Authors: Pável Matos-Maraví[1,2,3]*, Niklas Wahlberg[4], André V. L. Freitas[5], Phil J. DeVries[6],

Alexandre Antonelli[1,2,7], Carla M. Penz[6]

Full correspondence addresses:

[1] Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg,

Sweden

[2] Gothenburg Global Biodiversity Centre, Gothenburg, Sweden

[3] Biology Centre of the Czech Academy of Sciences, Institute of Entomology, České

Budějovice, Czech Republic

[4] Department of Biology, Lund University, Lund, Sweden

[5] Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas,

Campinas, SP, Brazil

[6] Department of Biological Sciences, University of New Orleans, New Orleans, LA, USA

[7] Royal Botanical Gardens Kew, TW9 3AE Richmond, United Kingdom

* Corresponding author (E-mail: pavel.matos@entu.cas.cz, ORCiD id: https://orcid.org/0000-

0002-2885-4919)

Abstract

Regional species diversity is ultimately explained by speciation, extinction, and dispersal. Here

we estimate dispersal and speciation rates in Neotropical rainforest biomes to propose an

explanation for the distribution and diversity of extant butterfly species. We focus on the tribe

Brassolini (owl butterflies and allies): a Neotropical group that comprises 17 genera and 108

species, most of them endemic to rainforest biomes. We infer a total-evidence species tree using

the multispecies coalescent framework. By applying biogeographical stochastic mapping, we

infer ancestral ranges and estimate rates of dispersal and cladogenesis at the scale of millions of

years. We suggest that speciation in Mesoamerica and the northwestern flank of the Andes have

only increased within the past 2 million years. In contrast, speciation in the Brazilian Atlantic

Forest has been constant throughout the past 10 million years. The disparate species

diversification dynamics may be partly explained by the geological and environmental history of

each bioregion. Importantly, the dispersal rates into the Atlantic Forest and Mesoamerica plus

NW Andes increased simultaneously in the middle-Miocene, suggesting that lineages from such

regions have had comparable times for speciation despite their decoupled diversification

dynamics. Diversification of extant Amazonian lineages, on the other hand, has episodically

increased since the late Miocene, including a rise in speciation rate during the Pleistocene.

Altogether, our results reveal a mosaic of biome-specific evolutionary histories within the

Neotropics, where species have diversified rapidly (cradles: e.g., Mesoamerica), have

accumulated gradually (museums: e.g., Atlantic Forest), or have alternately diversified and

accumulated (e.g., Amazonia).

Keywords: Speciation; dispersal; multispecies coalescent; total-evidence analyses; ancestral

state inference; biogeographical stochastic mapping.

Introduction

Terrestrial biomes are heterogeneous, and some areas accumulate greater diversity and endemic species than others. An explanation for this biogeographical pattern includes three evolutionary mechanisms: species origination, extinction, and dispersal (Goldberg et al. 2005, Jablonski et al. 2006). For example, a species-rich region can attain its present diversity by exceptionally high rates of speciation, becoming a "cradle of diversity", or by the persistence of old lineages, e.g., via low species extinction rates, becoming a "museum of diversity" (Stebbins 1974). Furthermore, a species-rich region might be a "source of diversity" when dispersal out of the region is high, or a "sink of diversity" when regional species diversity accumulates from external sources by dispersal into the region. Characterizing the interplay among speciation, extinction, and dispersal is thus highly relevant for understanding the origin and evolution of total regional diversity and endemism (Wiens and Donoghue 2004, Roy and Goldberg 2007).

The Neotropical region contains a great deal of the world's biodiversity and endemism. Most of such diversity is concentrated in tropical rainforests in Central America, the Amazon basin, and the Atlantic Forest in southeastern Brazil. However, there is no consensus on whether the currently scattered rainforests were once connected, nor for how long (Jaramillo and Cárdenas 2013). It is also unclear whether rainforests represent centers of "ecological stability" that share a common evolutionary history (Moritz et al. 2000). Mammals and birds in Neotropical biodiversity hotspots, including the Brazilian Atlantic Forest, and rainforests in Mesoamerica and the neighboring area Chocó (Myers et al. 2000) apparently originated at faster rates and exported more lineages compared to surrounding areas, suggesting that such hotspots are both cradles and sources of diversity (Igea and Tanentzap 2019). Some non-hotspots regions are, in

contrast, museums of diversity given the old and gradual accumulation of species through time,

such as Amazonia for Troidini butterflies (Papilionidae) (Condamine et al. 2012). It thus seems

that the checkered spatial assemblage of rainforests acting as cradle and museum areas might

explain why the Neotropics is considered both museum and cradle of diversity (McKenna and

Farrell 2006, Moreau and Bell 2013, Antonelli et al. 2018a). However, the impacts of geological

and paleoenvironmental changes in the Neotropics together with the roles of speciation and

dispersal have seldom been jointly elucidated in individual rainforest hotspots.

Here we test whether there was significant biotic exchange among Neotropical rainforests during

the Neogene and Quaternary periods (i.e., the past 23 Ma), which might indicate physical

connectivity of biomes. If connection among biomes existed, we aim to quantify its timing and

magnitude in terms of dispersal and speciation of rainforest taxa. To do this, we estimate

dispersal rates and speciation of Neotropical butterflies that occur mainly in rainforest hotspots,

such as Mesoamerica, the northwestern slope (NW) of the Andes (including e.g., Chocó), the

Atlantic Forest, Amazonia, and the South American "dry diagonal" (Fig. 1). Butterflies in the

tribe Brassolini (Nymphalidae: Satyrinae) are particularly suited to address our question because

they are a monophyletic group (Freitas and Brown 2004, Wahlberg et al. 2009, Espeland et al.

2018, Chazot et al. 2019a), exclusively Neotropical, and have about one third of their species

diversity endemic to rainforest hotspots (see Penz 2007 for an overview).

Studies that focused on individual brassoline genera used adult morphology (Penz 2008, 2009a,

b, Garzón-Orduña and Penz 2009) and DNA data (Penz et al. 2011a, Shirai et al. 2017) to infer

species-level phylogenies, but several deep nodes had weak or even conflicting phylogenetic

support in tribal-level analyses (Penz et al. 2013). We infer a time-calibrated species phylogeny

of Brassolini by merging all previous datasets and generating new DNA and morphological data.

Specifically, we use this phylogeny to revisit and test the following predictions on diversification

and dispersal among three Neotropical rainforest biomes:

1. Amazonia is the main source of Brassolini diversity by means of dispersal, reflecting the

patterns from Neotropical vertebrates and plants during the past 60 million years (Antonelli et al.

2018b). Rapid species diversification of some butterfly groups occurred there since at least the

Oligocene (Matos-Maraví 2016), and old lineages might have gradually accumulated in

Amazonia since that time (Condamine et al. 2012). More recent diversification episodes

supporting Amazonia as alternating museum and cradle of diversity could have been triggered by

paleoenvironmental changes, such as the retreat of the Pebas wetland system (Antonelli et al.

2009, Antonelli and Sanmartín 2011, Chazot et al. 2019b).

2. Mesoamerica and the NW Andes are together a sink of Brassolini diversity due to dispersal

events into this region during the Miocene, and cradle of diversity due to more recent, rapid

speciation in the Pleistocene. This scenario conforms with the likely emergence of landmass in

Central America and Chocó after the collision of the Panama Block and northwestern South

America by the late Oligocene–early Miocene (Coates and Stallard 2013, Jaramillo 2018), and

with Uranium-lead dating confirming that segments of the Panama arc had already emerged by

13–15 Ma (Montes et al. 2015). Shallow seawaters probably existed in the region until ~3 to 4

Ma (O'Dea et al. 2016, Jaramillo et al. 2017a), and while dispersal might have significantly

increased since the early Miocene about 20 Ma, facilitated by land availability (Bacon et al.

2015), regional species diversification might have been hindered by lowland area water

inundations until the Plio-Pleistocene (Mullen et al. 2011, Hosner et al. 2016).

3. The Atlantic Forest might have been a sink of Brassolini diversity for old Amazonian lineages

(>10–15 Ma) and/or a recent source of diversity for present dry diagonal region (<10 Ma). The

onset of a period of global cooling at ~10 to 15 Ma likely drove the expansion of open

environments between Amazonia and the Atlantic Forest (Simon et al. 2009). Apparently,

intermittent forest expansions into the dry diagonal occurred throughout the Neogene and

Pleistocene (Werneck 2011, Trujillo-Arias et al. 2017), and Amazonia and the Atlantic Forest

might have repeatedly been in contact (Ledo and Colli 2017, Capurucho et al. 2018). Biotic

interchange from the Atlantic Forest toward the dry diagonal might have been a geologically

more recent event, since at least 5 Ma (Costa 2003, Batalha-Filho et al. 2013, Thomé et al.

2016).

Material and methods

Taxon sampling

Specimens were collected in Mexico, Costa Rica, Ecuador, Peru, and Brazil (Supplementary

material Appendix 1 Table S1), and identified to species based on various sources (Casagrande

2002, 2004, Austin et al. 2007, Penz 2008, 2009a, b, Garzón-Orduña and Penz 2009, Penz et al.

2017, 2011a, b, Chacón et al. 2012). To evaluate taxonomic bias in relation to geography, we

compiled the year of description of each Brassolini species and the year of the latest revision of

their current taxonomic status (e.g., when former subspecies were raised to species).

Molecular dataset

Based on protocols in Wahlberg and Wheat (2008), we Sanger-sequenced fragments of the

mitochondrial COI gene (1,475 bp) and the nuclear genes CAD (850 bp), EF1α (1,240 bp),

GAPDH (691 bp), RpS5 (617 bp), and *wingless* (400 bp) (outsourced to Macrogen, Seoul). We

obtained genetic data from 81 individuals classified in 57 out of 108 valid species in 15 out of 17

Brassolini genera (Penz 2007). The two missing genera (*Aponarope* and *Mielkella*) are

monotypic. We retrieved COI sequences available from the BOLD database

(http://boldsystems.org) for seven additional species; thus, our molecular dataset consisted of 15

genera and 64 species. To root the phylogeny, we included genetic data of 19 outgroup taxa in

the subfamily Satyrinae from public data deposited in GenBank. All DNA sequences from this

study are in GenBank (accession numbers MK551348–MK551551).

Morphological and total-evidence dataset

Our morphological dataset included 255 characters (201 from adults and 54 from early stages)

for 64 species from all 17 Brassolini genera. The morphological matrix was concatenated from

previous studies (Penz 2007, 2008, 2009a, b, Garzón-Orduña and Penz 2009, Penz et al. 2013)

by eliminating duplicate characters and scoring data for species missing entries. Specimens used

were the same as listed in the original publications. The combined molecular and morphological

dataset comprised all genera and 84 out of 108 Brassolini species. We call this the total-evidence

dataset, which we used to infer a time-calibrated phylogeny.

Distribution dataset

We delineated bioregions (*sensu* Vilhena and Antonelli 2015) using geo-referenced occurrence

points of Brassolini identified at the species level, of which 6,378 points came from GBIF

(https://gbif.org; retrieved on 21[st] January 2019) and 881 points from the ATLANTIC

BUTTERFLIES dataset (Santos et al. 2018). We used the R v3.5.3 (R Core Team 2019) package

CoordinateCleaner v2.0-11 (Zizka et al. 2019) for automated cleaning of species occurrences.

The software flagged 81 potential errors, such as occurrence records in the ocean, geographic

coordinates identical to country centroids, and occurrences in close proximity to biodiversity

institutions. We uploaded the 7,174 vetted geo-referenced occurrences to the web application

Infomap Bioregions (Edler et al. 2017) at http://bioregions.mapequation.org/. We set the

maximum and minimum values for cell sizes to 4 and 2, and for cell capacities to 40 and 4. We

used different values of cluster cost, from 0.1 to 3.0, in steps of 0.1. The number of inferred areas

remained stable between values of 1.8 and 2.4 and, thus, received the strongest support. The

bioregions identified correspond to previously identified biomes or combinations of these: 1)

Mesoamerica plus NW Andes, 2) Brazilian Atlantic Forest, and 3) the remaining tropical South

America, including the Amazon drainage basin and the seasonally dry diagonal biomes Cerrado

and Caatinga (Fig. 1).

To inform the biogeographical and within-area speciation rate analyses, we generated a species-

level distributional dataset. We surveyed the literature to score species occurrences in the defined

bioregions because the geo-referenced dataset did not include all Brassolini species. To

investigate dispersal patterns across the South American dry diagonal, we kept this area separate.

We used distributional maps and observations, not always with primary geographic coordinates,

from taxonomic revisions (Bristow 1981, 1982, 1991, Casagrande 2002, Furtado and Campos-

Neto 2004, Penz 2008, 2009a, b, Garzón-Orduña and Penz 2009, Penz et al. 2017) and published

butterfly inventories in Mesoamerica (DeVries 1983, 1994, Janzen and Hallwachs 2009, Basset

et al. 2015), Amazonia (Pereira Martins et al. 2017), Cerrado (Pinheiro and Emery 2006, Emery

et al. 2006, Silva et al. 2012, Pereira Martins et al. 2017, Dickens et al. 2019), Caatinga (Zacca

and Bravo 2012), and Atlantic Forest (Santos et al. 2011, 2018, Pérez et al. 2017, Melo et al.

2019, Soldati et al. 2019). We call this the presence/absence biogeographical dataset.

Total-evidence phylogenetic inference and divergence time calibration

We inferred time-calibrated species trees using the StarBEAST2 package (Ogilvie et al. 2017)

available in BEAST v2.5.2 (Bouckaert et al. 2014). We removed the species *Opsiphanes camena*

from the dataset because it had only 325 bp of the COI locus, and its phylogenetic position was

not stable in a preliminary analysis (Supplementary material Appendix 1). We used a gene-based

partitioning strategy, each having unlinked tree models in the multispecies coalescent

framework. We performed model averaging over 31 substitution models available in the

bModelTest v1.1.2 package (Bouckaert and Drummond 2017). We applied uncorrelated relaxed-

clock models for all partitions because strict clock models were rejected based on the stepping-

stone method (Supplementary material Appendix 1). We estimated clock rates relative to the

mitochondrial mean clock rate fixed to 1.0, and we used log normal distribution for the relative

clock mean priors (M = 0, S = 1). We used the Yule tree model because it received decisive

support over the birth-death tree model based on path-sampling steps (Supplementary material

Appendix 1). We added the morphological dataset onto the species tree and we conditioned the

characters to the Markov model (Lewis 2001).

Inasmuch as there are no described fossils of Brassolini, we relied on secondary calibrations to

time-calibrate the species tree. We conservatively used uniform distributions encompassing the

95% highest posterior density (HPD) from a large fossil-calibrated, genus-level butterfly

phylogeny (Chazot et al. 2019a) to constrain six outgroup nodes:

1) The crown age of the subfamily Satyrinae (46–65 Ma),

2) The divergence of the tribes Melaniti and Dirini (27–41 Ma),

3) The crown age of the tribe Satyrini (38–55 Ma),

4) The crown node of the subtribes Lethina, Parargina and Mycalesina to (31–46 Ma),

5) The crown node of the subtribes Pronophilina, Euptychiina, Satyrina and Erebiina (31–44

Ma),

6) The divergence of the tribes Brassolini and Morphini (34–50 Ma).

The analyses were run four independent times for 300 million generations, via the CIPRES

Science Gateway v3.3 (Miller et al. 2010). We sampled 20,000 trees from the posterior

distribution and discarded the first 25% as burnin. We checked that the estimated sample sizes

(ESS) were above 200 using Tracer v1.7.1 (Rambaut et al. 2018). We merged the independent

runs using LogCombiner (part of the BEAST v2.5.2 package) and we summarized the post-

burnin species trees as a maximum clade credibility (MCC) tree in TreeAnnotator (part of the

BEAST v2.5.2 package).

Within-area speciation and dispersal among bioregions through time

We estimated ancestral geographic ranges and colonization rates among bioregions using the

dispersal-extinction-cladogenesis (DEC) model (Ree et al. 2005), as implemented in the R

package BioGeoBEARS v1.1.2 (Matzke 2013). We used the presence/absence biogeographical

dataset and the four defined bioregions: 1) Mesoamerica plus NW Andes, 2) Atlantic Forest, 3)

Amazonia, and 4) Dry diagonal. We did not constrain any *a priori* dispersal multiplier, nor did

we stratify dispersal rates across the phylogeny. To account for ancestral state uncertainty, we

carried out 100 biogeographical stochastic mappings (BSM; Dupin et al. 2017). To account for

divergence time uncertainty, we used 100 random topologies from the BEAST2 posterior

distribution. To account for phylogenetic uncertainty of 25 missing species, we randomly added

such lineages to their currently assigned genera in the 100 posterior phylogenies. In this case, we

assumed that phylogenetic diversity per genus was maximized and we placed missing taxa in the

crown genera given our comprehensive taxonomic sampling. All genera are monophyletic, but

we conservatively joined *Selenophanes* and *Catoblepia* (posterior probability, PP = 1.0; Fig. 2)

when assigning missing species because the monophyly of *Catoblepia* was only moderately

supported (PP = 0.75).

We estimated the number of dispersal events and within-area-cladogenesis events by simulating

areas on nodes based on the 10,000 pseudoreplicated biogeographic histories (100 BSM × 100

posterior species trees). We calculated colonization rates through time as $c_{XtoY}(t_1) = d_{XtoY}(t_1)$ /

$L(t_1)$, where $d_{XtoY}(t_1)$ is the number of inferred dispersal events from area X to Y in a 1-million

year interval, and $L(t_1)$ is the total length of all branches in the 1-million-year interval (Antonelli

et al. 2018b). In addition, we calculated the relative number of *in situ* speciation events through

time as $\lambda(t_1) = s(t_1)$ / $s(t_0 / t_{0+1})$, where $s(t_1)$ is the number of within-area cladogenesis in a 1-

million-year interval, and $s(t_0 / t_{0+1})$ is the number of within-area cladogenesis in the previous 1-

million-year interval (t_0) relative to the number of within-area cladogenesis in time intervals t_0

and t_1 (modified from Xing and Ree 2017).

Speciation within bioregions through time

For a second, independent estimate of within-area speciation, we calculated speciation rates for

all tips and branches in the MCC species tree using BAMM v2.5.0 (Rabosky 2014). To account

for the 25 missing species, we generated clade-specific sampling fractions at the genus level. We

used the R package BAMMtools v2.1.6 (Rabosky et al. 2014) to estimate prior values for the

evolutionary rate parameters, and we assigned the expected number of shifts to 1.0 as

recommended for phylogenies including ~100 species (Rabosky 2014). We ran the analysis for

10 million generations, sampling event data every 1,000 generations. After discarding the first

20% samples, we checked that the ESS value was higher than 200 and the likelihood estimates

reached a stationary distribution.

First, we estimated speciation rates within bioregions at present using the function "getTipRates"

in BAMMtools. We computed tip-specific mean speciation rates for those species occurring only

in one bioregion, i.e., the endemic lineages. Second, we used the function "dtRates" in

BAMMtools to compute mean speciation rates along branches in ~1-million year intervals (tau

parameter set to 0.05). We assigned the most probable inferred area from the 10,000 BSMs to the

points where branches cross time intervals (Igea and Tanentzap 2019). When the most probable

area for a node and its parental one were the same, we assumed that the branch connecting both

nodes occurred in such an area. When inferred areas for a node and its parental one were

different, we assumed that the dispersal event occurred stochastically along the branch. We

computed mean speciation rates only for nodes that most probably occurred in a single

bioregion.

All the datasets from this study can be found in TreeBASE (ID 25040) and Dryad (DOI:

XXXXX).

Results

Phylogenetic and divergence time uncertainty

We found no disparate trend among Neotropical bioregions in taxon discovery, description and

taxonomic opinion (i.e., oversplitting species *vs*. multiple infraspecific taxa) (Supplementary

material Appendix 1). The multispecies coalescent model recovered low posterior probabilities

for deeper nodes (0.39–0.90; Fig. 2). In contrast, a concatenation-based analysis using only the

molecular dataset (Supplementary material Appendix 1) produced higher posterior probabilities

for the deep nodes (0.97–1.00). This suggests that the molecular dataset held strong phylogenetic

signal, but the multispecies coalescent model might have better reflected the phylogenetic

uncertainty in the morphological dataset. We estimated these nodes had ages between 15 and 17

Ma (entire highest posterior density HPD, 10.87–20.89 Ma). The total-evidence dataset

supported posterior probability of 1.0 for the crown node of almost all non-monotypic genera,

regardless of phylogenetic method.

Ancestral bioregions and colonization rates through time

The inferred ranges coming from the 10,000 pseudoreplicated biogeographic histories and

plotted against the MCC species tree in Fig. 2 suggested that Brassolini most probably ($Pr =$

0.36) originated in Amazonia at 21 Ma (entire HPD, 16.67–27.16 Ma). The early rapid radiation

of Brassolini between 15 and 17 Ma also probably took place in Amazonia ($Pr = 0.30$–0.52).

Dispersal out of Amazonia had the highest rates compared to other bioregions (Fig. 3).

Colonization rate from Amazonia to Mesoamerica showed a sharp increase by ~5 Ma and is

currently the highest rate among other dispersal types (Fig. 3A). Dispersal into Mesoamerica

from other bioregions occurred throughout most of the Miocene, and peaked at ~10 Ma

(Supplementary material Appendix 1). Dispersal out of Mesoamerica + NW Andes, however,

was very low through time, and only since the past ~2 Ma did it exponentially increase toward

Amazonia (Fig. 3A). Dispersal from Amazonia to the dry diagonal increased throughout the

Plio-Pleistocene and has since then surpassed dispersal to Atlantic Forest (Fig. 3B & 3C).

Dispersal from the Atlantic Forest into Amazonia was higher at ~10 Ma than from any other

region (Fig. 3C). Nonetheless, dispersal into or out of the Atlantic Forest from/to any other

bioregion declined from ~10 to 2 Ma, rendering the Atlantic Forest isolated for most of the late

Miocene and Pliocene; only since the Pleistocene ~2 Ma dispersal into and from the dry diagonal

has increased until the present (Fig. 3D).

Regional speciation through time

The relative rates of within-area cladogenesis estimated in BioGeoBEARS and speciation

estimated in BAMM were congruent (Fig. 4). Speciation rates increased through time in all areas

except in the dry diagonal, which has had near zero endemic cladogenesis. Diversification began

earlier in the Atlantic Forest, at ~12 Ma when the cladogenesis rate was slightly above zero (Fig.

4A) and speciation rate increased (Fig. 4B), specifically in the clade comprising *Penetes*,

Orobrassolis and *Blepolenis*. However, diversification in the Atlantic Forest decreased

throughout most of the Miocene until ~ 5 Ma, when it steadily rose again towards the present. In

contrast, speciation in Mesoamerica remained low during most of the Miocene and Pliocene, and

sharply increased only at ~2 to 3 Ma. Speciation in Amazonia increased since the late Miocene at

higher rate than in the Atlantic Forest and Mesoamerica, but it also dramatically increased during

the Pleistocene (<2 Ma). At present, the speciation rate in Mesoamerica + NW Andes is similar

to Amazonian lineages, both being much higher than in the Atlantic Forest (Fig. 4C).

Discussion

Here we investigated the role of regional speciation and dispersal to understand the origin of

species in the butterfly tribe Brassolini. One third of those species are endemic to rainforests in

Mesoamerica, NW Andes and the Brazilian Atlantic Forest. Recent and rapid butterfly speciation

occurred in Mesoamerica and NW Andes, suggesting that it acted as a cradle of diversity. In

contrast, low and constant diversification since at least the late Miocene has shaped Brassolini

endemic diversity in the Atlantic Forest. This finding, coupled with the survival of old lineages

found mainly in montane habitats (e.g., *Penetes*, *Dasyophthalma*) suggests that the Atlantic

Forest acts as a museum of diversity.

A longer time for speciation in the Atlantic Forest potentially explaining the older events of

cladogenesis is not supported because dispersal into Mesoamerica and NW Andes occurred near-

simultaneously. We argue that the uneven origin of endemic species among the Neotropical

rainforest biomes reflect the geological and environmental changes that took place throughout

the Neogene and Pleistocene, including the availability of substantial landmass in Mesoamerica

and Chocó by ~3 to 4 Ma and the spread of wet forest into what is now the South American Dry

Diagonal during the past 10 Ma.

Strengths and weaknesses of the approach

We are aware of two main limitations and took them into account in the interpretation of our

results. First, by using the DEC model, we assumed no species extinction in the phylogeny of

extant Brassolini. Simulation analyses comparing the DEC and state-dependent diversification

models suggested that the estimation of ancestral areas can be robust when extinction rates were

low, or not associated to particular geographic areas (Matzke 2014). The lack of Brassolini

fossils precludes assessing this directly, but a better fit of the Yule process over the birth-death

model suggests that extinction was probably low to moderate during the timeframe of their

diversification, and might not have substantially biased our inferences.

Second, BAMM's prior model used for estimating diversification rate shifts were contested by

Moore et al. (2016), but further evaluation via simulated data suggested that BAMM can infer

speciation rates accurately from extant-taxon phylogenies (Rabosky et al. 2017). Although low

extinction rates might explain the museum of diversity pattern, we have conservatively focused

only on regional species origination rates to explain lineage accumulation within bioregions. We

additionally used a fundamentally-distinct measure of within-area cladogenesis (Xing and Ree

2017) that corroborated the pattern of recent and rapid speciation in Mesoamerica + NW Andes,

and an older lineage accumulation and lower diversification in the Atlantic Forest.

Our approach to disentangle regional speciation from dispersal to understand the origin of

within-area species diversity has two strengths. First, we defined bioregions that fit the

distribution and composition of Brassolini species by using geo-referenced occurrences, rather

than expert opinion or another subjective criterion (Vilhena and Antonelli 2015). This approach

revealed a clustering of butterfly communities from southern North America to Mesoamerica and

the NW Andes, which are regions that otherwise belong to different biogeographic dominions

(Morrone 2014). Our inference of large bioregions for Brassolini agrees with its apparent poor

spatial structuring and high dispersal capabilities. For instance, given a suitable environment,

ancestral *Bia* had the capability to disperse across pan-Amazonia in only 1,463–3,115 years

(Penz et al. 2015). The delimitation of Mesoamerica + NW Andes and the Atlantic Forest as

separate bioregions has likely been determined by ecological and habitat suitability rather than

geographical barriers to Brassolini dispersal; this being a similar biogeographical pattern found

in other Neotropical rainforest taxa (Dexter et al. 2017, Pérez-Escobar et al. 2019).

Second, we used a multispecies coalescent framework which outperforms the concatenation-

based approach (Edwards et al. 2016), in the face of incomplete taxon sampling and incomplete

lineage sorting (Liu et al. 2015, Ogilvie et al. 2017). For additional gene-tree discordance tests,

see Supplementary material, Appendix 1. Our comprehensive dataset including molecular and

morphological characters allowed us to simultaneously use several lines of evidence for

recognizing and dating speciation events. The Bayesian probabilistic framework applied here

also considered three types of uncertainties: phylogenetic, temporal and taxonomic sampling.

This approach contributed toward a better assessment of confidence intervals when estimating

dispersal and diversification rates.

Species diversification and dispersal in bioregions

The calculated probability densities of dispersal and speciation rates supported three distinct

evolutionary processes: recent and fast speciation in Mesoamerica + NW Andes, old

diversification dynamics with low speciation rates in Atlantic Forest, and episodic increases in

speciation rate of Amazonian lineages. Our results suggest that Brassolini diversified mainly in

Amazonia and the Atlantic Forest from about 23 to 5 Ma (speciation rate remained above 0.4 in

Amazonia, and below 0.4 in the Atlantic Forest, Fig. 4B), and early Brassolini diversification

might have been accompanied by increasing dispersal out of these two bioregions.

In the case of Amazonia, the Pebas wetland system (Wesselingh and Salo 2006) may not have

drastically limited dispersal and diversification of Brassolini as it likely did for other butterfly

groups (Chazot et al. 2019b). These wetland system was occasionally affected by marine

flooding events, which were probably short-lived, and at least two dated events at ~17.8 Ma and

~13.7 (Jaramillo et al. 2017b) were concurrent with the early and rapid cladogenesis events of

Brassolini in Amazonia. The Pebas wetland might in fact have increased opportunities for

dispersal and diversification of some typical coastal plants (Bernal et al. 2019), including many

of the butterflies' hostplants in the region: Arecaceae (palms), Poaceae (bamboo) and

Zingiberales such as Marantaceae (Beccaloni et al. 2008, Janzen et al. 2009). Therefore, the

evolutionary consequences of the Miocene marine incursions in western Amazonia might have

been mixed, restricting dispersal and diversification in some lineages while providing ecological

opportunities for others, such as Brassolini butterflies.

The recent increased diversification in Amazonian lineages (< 3 Ma) might be related to younger

paleoenvironmental changes, such as fluctuating river dynamics mediated by Quaternary glacial

cycles (Ribas et al. 2012). For example, species in the genus *Bia* seem to have strong

biogeographical structure driven by major Amazon-basin rivers (Penz et al. 2017), suggesting

that riverine barriers might have been involved in allopatric differentiation. Indeed, in areas of

the Amazon River, the distance between river banks is larger than the estimated daily dispersal

of species such as *Bia actorion* (Tufto et al. 2012, Penz et al. 2015). However, a phylogeographic

evaluation including more samples per species would be necessary to estimate the influence of

river barriers in differentiating Brassolini populations, and potentially to explain the major

increase in diversification during the Pleistocene.

In the case of the Atlantic Forest, Miocene expansions of wet forests into Amazonia through the

dry diagonal might have been determinants of ancient butterfly diversity. Forest connections

have been documented mainly for the Quaternary based on palynological data and

phylogeographic studies (e.g., Werneck 2011, Prates et al. 2016, Trujillo-Arias et al. 2017).

Although evidence of Neogene connection between the Atlantic Forest and Amazonia is still

scarce, the origin of the modern transcontinental Amazon river by ~10 Ma might have facilitated

dispersal of rainforest taxa (Albert et al. 2018). Indeed, we found that late Miocene dispersal

rates between these wet forest biomes were higher than in the Plio-Pleistocene. We suggest that

episodic Miocene wet forest expansions might have represented ecological opportunities for

Brassolini to disperse across Central Brazil, either by increased sizes of wet forest galleries in the

dry diagonal biome, or by fully connecting Amazonia and Atlantic Forest.

By the Plio-Pleistocene, speciation in Mesoamerica and NW Andes seems to have increased and

surpassed the speciation rate in the Atlantic Forest. Biotic dispersal into Mesoamerica from

South America likely occurred throughout most of the Miocene (e.g., Mullen et al. 2011, Bacon

et al. 2015). However, only at the end of the Pliocene did shallow water fully recede from the

region (Coates et al. 2004, Montes et al. 2012), thus increasing opportunities for terrestrial

species diversification. In contrast, speciation in the Atlantic Forest remained low and dispersal

out of this region decreased until the mid-Pleistocene. These events coincided with the expansion

of open environments in the American continent and between Amazonia and Atlantic Forest

since the late Miocene (Simon et al. 2009, Edwards et al. 2010, Strömberg 2011, Werneck 2011).

The contraction of Atlantic Forest may have restricted speciation to montane habitats in

southeastern Brazil where most extant species occur, and old lineages having low speciation

rates through time survived in remnants of rainforest.

Consequences for Brassolini classification

Brassolini has been subdivided in three subtribes, the monotypic Biina and Naropina, and

polytypic Brassolina (Casagrande 1995, Penz 2007), but here *Narope* falls within Brassolina. At

the genus level, our analyses support sister taxa relationships between *Dynastor* and

Dasyophthalma, *Opoptera* and *Narope*, *Selenophanes* and *Catoblepia*, *Orobrassolis* and

Blepolenis, and *Brassolis* appears as sister to the *Opsiphanes*-group (sensu Penz 2007). We

propose the following changes to the Brassolini classification at the subtribe and species levels:

1) Naropina is subsumed within Brassolina. Brassolini is thus redefined to include the subtribes

Biina (monotypic) and Brassolina.

2) The monotypic *Aponarope* is subsumed within *Narope*. Stichel (1916) originally described

Narope sutor, but this species was later segregated in the monotypic *Aponarope* (Casagrande

1982). We revise the genus assignment for *Narope sutor*, REV. COMB., and synonymize

Aponarope, NEW SYN.

3) *Selenophanes orgetorix*, NEW COMB. This species grouped with members of *Selenophanes*

with high posterior probability (see also Shirai et al. 2017).

Conclusions

This study suggests that the Neotropics consists of a regional network of museums and cradles of

butterfly diversity. We found that endemic species to Mesoamerica plus NW Andes have

originated by old dispersal events into the region, and that speciation rates increased only at

about 2 Ma (cradle of diversity). In contrast, endemic species to the Brazilian Atlantic Forest

likely arose under low and continuous speciation rates since about 12 Ma (museum of diversity),

and was likely intermittently connected to Amazonia since the Miocene. This dynamic

evolutionary history of Neotropical rainforests appear to reflect paleoenvironmental changes,

including landscape reconfigurations in Central America and paleoclimate fluctuations during

the Neogene and Quaternary.

Acknowledgements

We are grateful to Carlos Peña, Mirna Casagrande, Helena Romanowski, Augusto H. B. Rosa and Andrew Warren for contributing specimens used in this study; and to Diana Silva, Gerardo Lamas and SERFOR (Ministerio de Agricultura, Peru) for assistance with research permits (No Autorización 223-2017-SERFOR/DGGSPFFS). We also are grateful to the ICMBio for the research permits (SISBIO nº 10802-5, 10802-24 and 53016-10). We acknowledge the computational resources provided by the CESNET LM2015042 and the CERIT Scientific Cloud LM2015085, provided under the programme "Projects of Large Research, Development, and Innovations Infrastructures". This publication is part of the RedeLep 'Rede Nacional de Pesquisa e Conservação de Lepidópteros' SISBIOTA-Brasil/CNPq (563332/2010-7).

Funding

PMM was supported by a Marie Skłodowska-Curie fellowship (European Commission, Grant No. MARIPOSAS-704035), and by a fellowship from the PPLZ programme (Czech Academy of Sciences, Fellowship No. L200961951). AVLF acknowledges support from FAPESP (2011/50225-3, 2012/50260-6, 2013/50297-0), from the Brazilian Research Council - CNPq (303834/2015-3), from the National Science Foundation (DEB-1256742) and from the United States Agency for International Development - USAID / the U.S. National Academy of Sciences (NAS), under the PEER program (Sponsor Grant Award Number: AID-OAA-A-11-00012) (Mapping and Conserving Butterfly Biodiversity in the Brazilian Amazon). AA acknowledges support from the Swedish Research Council, the Swedish Foundation for Strategic Research, the Knut and Alice Wallenberg Foundation, and the Royal Botanic Gardens, Kew.

Author contributions

PMM, NW, and CMP conceived the project, PMM, AVLF, PJD, and CMP conducted fieldwork,

PMM carried out the lab experiments and led the data analyses, PMM wrote the first draft of the

manuscript, and all authors discussed and contributed to the final product.

References

Albert, J. S. et al. 2018. The changing course of the Amazon River in the Neogene: center stage

for Neotropical diversification. - Neotrop. Ichthyol. 16: e180033.

Antonelli, A. and Sanmartín, I. 2011. Why are there so many plant species in the Neotropics? -

Taxon 60: 403–414.

Antonelli, A. et al. 2009. Tracing the impact of the Andean uplift on Neotropical plant evolution.

- Proc. Natl. Acad. Sci. 106: 9749–9754.

Antonelli, A. et al. 2018a. Conceptual and empirical advances in Neotropical biodiversity

research. - PeerJ 6: e5644.

Antonelli, A. et al. 2018b. Amazonia is the primary source of Neotropical biodiversity. - Proc.

Natl. Acad. Sci. 115: 6034–6039.

Austin, G. T. et al. 2007. A new species of *Opsiphanes* Doubleday, [1849] from western Mexico

(Nymphalidae: Morphinae: Brassolini). - Bull. Allyn Mus. 150: 1–20.

Bacon, C. D. et al. 2015. Biological evidence supports an early and complex emergence of the

Isthmus of Panama. - Proc. Natl. Acad. Sci. 112: 6110–6115.

Basset, Y. et al. 2015. The butterflies of Barro Colorado Island, Panama: local extinction since

the 1930s. - PLoS One 10: e0136623.

Batalha-Filho, H. et al. 2013. Connections between the Atlantic and the Amazonian forest

avifaunas represent distinct historical events. - J. Ornithol. 154: 41–50.

Beccaloni, G. W. et al. 2008. Catalogue of the hostplants of the Neotropical butterflies/Catálogo

de las plantas huésped de las mariposas neotropicales. - Sociedad de Entomología

Aragonesa.

Bernal, R. et al. 2019. Could coastal plants in western Amazonia be relicts of past marine

incursions? - J. Biogeogr. 46: 1749–1759.

Bouckaert, R. R. and Drummond, A. J. 2017. bModelTest: Bayesian phylogenetic site model

averaging and model comparison. - BMC Evol. Biol. 17: 42.

Bouckaert, R. et al. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. -

PLoS Comput. Biol. 10: e1003537.

Bristow, C. R. 1981. A revision of the brassoline genus *Catoblepia* (Lepidoptera: Rhopalocera). -

Zool. J. Linn. Soc. 72: 117–163.

Bristow, C. R. 1982. A revision of the brassoline genus *Selenophanes* (Lepidoptera:

Rhopalocera). - Zool. J. Linn. Soc. 76: 273–291.

Bristow, C. R. 1991. A revision of the brassoline genus *Opsiphanes* (Lepidoptera: Rhopalocera).

- Zool. J. Linn. Soc. 101: 203–293.

Capurucho, J. M. G. et al. 2018. Connecting Amazonian, Cerrado, and Atlantic forest histories:

Paraphyly, old divergences, and modern population dynamics in tyrant-manakins

(*Neopelma*/*Tyranneutes*, Aves: Pipridae). - Mol. Phylogenet. Evol. 127: 696–705.

Casagrande, M. M. 1982. Quatro gêneros novos de Brassolinae (Lepidoptera-Nymphalidae). -

Rev. Bras. Entomol. 26: 355–356.

Casagrande, M. M. 1995. Notas sistemáticas sobre Brassolinae. I. Tribos (Lepidoptera,

Nymphalidae). - Rev. Bras. Zool. 12: 671–699.

Casagrande, M. M. 2002. Naropini Stichel, taxonomia e imaturos (Lepidoptera, Nymphalidae,

Brassolinae). - Rev. Bras. Zool. 19: 467–569.

Casagrande, M. M. 2004. Brassolini: Checklist. Part 4A. Hesperioidea - Papilionoidea. - In:

Heppner, J. B. and Lamas, G. (eds), Atlas of Neotropical Lepidoptera. Association of

Tropical Lepidoptera/Scientific publishers, pp. 201–205.

Chacón, I. A. et al. 2012. A new species of *Opsiphanes* Doubleday, [1849] from Costa Rica

(Nymphalidae: Morphinae: Brassolini), as revealed by its DNA barcodes and habitus. -

Bull. Allyn Mus. 166: 1–15.

Chazot, N. et al. 2019a. Priors and posteriors in Bayesian timing of divergence analyses: the age

of butterflies revisited. - Syst. Biol. 68: 797–813.

Chazot, N. et al. 2019b. Renewed diversification following Miocene landscape turnover in a

Neotropical butterfly radiation. - Glob. Ecol. Biogeogr. 28: 1118–1132.

Coates, A. G. and Stallard, R. F. 2013. How old is the Isthmus of Panama? - Bull. Mar. Sci. 89:

801–813.

Coates, A. G. et al. 2004. The geology of the Darien, Panama, and the late Miocene-Pliocene

collision of the Panama arc with northwestern South America. - Bull. Geol. Soc. Am. 116:

1327–1344.

Condamine, F. L. et al. 2012. Biogeographic and diversification patterns of Neotropical Troidini

butterflies (Papilionidae) support a museum model of diversity dynamics for Amazonia. -

BMC Evol. Biol. 12: 82.

Costa, L. P. 2003. The historical bridge between the Amazon and the Atlantic Forest of Brazil: a

study of molecular phylogeography with small mammals. - J. Biogeogr. 30: 71–86.

DeVries, P. J. 1983. Checklist of butterflies and natural history. - In: Janzen, D. H. (ed), Costa

Rican Natural History. University of Chicago Press, pp. 654–678.

DeVries, P. J. 1994. Patterns of butterfly diversity and promising topics in natural history and

ecology. - In: McDade, L. A. et al. (eds), La Selva: Ecology and Natural History of a

Neotropical Rain Forest. The University of Chicago Press, pp. 187–194.

Dexter, K. G. et al. 2017. Dispersal assembly of rain forest tree communities across the Amazon

basin. - Proc. Natl. Acad. Sci. 114: 2645–2650.

Dickens, J. K. et al. 2019. The butterflies of a Cerrado–Atlantic Forest ecotone at Laguna Blanca

reveal underestimation of Paraguayan butterfly diversity and need for conservation. - J.

Insect Conserv. 23: 707–728.

Dupin, J. et al. 2017. Bayesian estimation of the global biogeographical history of the

Solanaceae. - J. Biogeogr. 44: 887–899.

Edler, D. et al. 2017. Infomap Bioregions: interactive mapping of biogeographical regions from

species distributions. - Syst. Biol. 66: 197–204.

Edwards, E. J. et al. 2010. The origins of C_4 grasslands: integrating evolutionary and ecosystem

science. - Science 328: 587–591.

Edwards, S. V. et al. 2016. Implementing and testing the multispecies coalescent model: a

valuable paradigm for phylogenomics. - Mol. Phylogenet. Evol. 94: 447–462.

Emery, E. de O. et al. 2006. As borboletas (Lepidoptera, Papilionoidea) do Distrito Federal,

Brasil. - Rev. Bras. Entomol. 50: 85–92.

Espeland, M. et al. 2018. A comprehensive and dated phylogenomic analysis of butterflies. -

Curr. Biol. 28: 770–778.

Freitas, A. V. L. and Brown, K. S. Jr. 2004. Phylogeny of the Nymphalidae (Lepidoptera). - Syst.

Biol. 53: 363–383.

Furtado, E. and Campos-Neto, F. C. 2004. *Caligopsis seleucida* (Hewitson) e seus estágios

imaturos (Lepidoptera, Nymphalidae, Brassolinae). - Rev. Bras. Zool. 21: 593–597.

Garzón-Orduña, I. J. and Penz, C. M. 2009. Phylogeny of *Dynastor* and *Brassolis* butterflies

(Lepidoptera: Nymphalidae): a tough nut to crack. - Zootaxa 2134: 1–22.

Goldberg, E. E. et al. 2005. Diversity, endemism, and age distributions in macroevolutionary

602 sources and sinks. - Am. Nat. 165: 623–633.

603 Hosner, P. A. et al. 2016. Rapid and recent diversification of curassows, guans, and chachalacas

604  (Galliformes: Cracidae) out of Mesoamerica: phylogeny inferred from mitochondrial,

605  intron, and ultraconserved element sequences. - Mol. Phylogenet. Evol. 102: 320–330.

606 Igea, J. and Tanentzap, A. J. 2019. Multiple macroevolutionary routes to becoming a biodiversity

607  hotspot. - Sci. Adv. 5: eaau8067.

608 Jablonski, D. et al. 2006. Out of the tropics: evolutionary diversity gradient. - Science 314: 102–

609  106.

610 Janzen, D. H. and Hallwachs, W. 2009. Dynamic database for an inventory of the

611  macrocaterpillar fauna, and its food plants and parasitoids, of Area de Conservacion

612  Guanacaste (ACG), northwestern Costa Rica. <http://janzen.sas.upenn.edu>.

613 Janzen, D. H. et al. 2009. Integration of DNA barcoding into an ongoing inventory of complex

614  tropical biodiversity. - Mol. Ecol. Resour. 9: 1–26.

615 Jaramillo, C. 2018. Evolution of the Isthmus of Panama: Biological, paleoceanographic and

616  paleoclimatological implications. - In: Hoorn, C. et al. (eds), Mountains, Climate and

617  Biodiversity. First. John Wiley & Sons, Ltd, pp. 323–338.

618 Jaramillo, C. and Cárdenas, A. 2013. Global warming and Neotropical rainforests: a historical

619  perspective. - Annu. Rev. Earth Planet. Sci. 41: 741–766.

620 Jaramillo, C. et al. 2017a. Comment (1) on "Formation of the Isthmus of Panama" by O'Dea *et*

621  *al*. - Sci. Adv. 3: e1602321.

622 Jaramillo, C. et al. 2017b. Miocene flooding events of western Amazonia. - Sci. Adv. 3:

623  e1601693.

624 Ledo, R. M. D. and Colli, G. R. 2017. The historical connections between the Amazon and the

Atlantic Forest revisited. - J. Biogeogr. 44: 2551–2563.

Lewis, P. O. 2001. A likelihood approach to estimating phylogeny from discrete morphological

character data. - Syst. Biol. 50: 913–925.

Liu, L. et al. 2015. Coalescent methods are robust to the simultaneous effects of long branches

and incomplete lineage sorting. - Mol. Biol. Evol. 32: 791–805.

Löwenberg-Neto, P. 2014. Neotropical region: a shapefile of Morrone's (2014) biogeographical

regionalisation. - Zootaxa 3802: 300.

Matos-Maraví, P. 2016. Investigating the timing of origin and evolutionary processes shaping

regional species diversity: Insights from simulated data and Neotropical butterfly

diversification rates. - Evolution 70: 1638–1650.

Matzke, N. J. 2013. Probabilistic historical biogeography: new models for founder-event

speciation, imperfect detection, and fossils allow improved accuracy and model-testing. -

Front. Biogeogr. 5: 242–248.

Matzke, N. J. 2014. Model selection in historical biogeography reveals that founder-event

speciation is a crucial process in island clades. - Syst. Biol. 63: 951–970.

McKenna, D. D. and Farrell, B. D. 2006. Tropical forests are both evolutionary cradles and

museums of leaf beetle diversity. - Proc. Natl. Acad. Sci. 103: 10947–10951.

Melo, D. H. A. et al. 2019. Butterflies (Lepidoptera: Papilionoidea) of an urban park in

northeastern Brazil. - Biota Neotrop. 19: e20180614.

Miller, M. A. et al. 2010. Creating the CIPRES Science Gateway for inference of large

phylogenetic trees. - Proc. Gatew. Comput. Environ. Work. (GCE), 14 Nov.: 1–8.

Montes, C. et al. 2012. Arc-continent collision and orocline formation: closing of the Central

American seaway. - J. Geophys. Res. 117: B04105.

Montes, C. et al. 2015. Middle Miocene closure of the Central American Seaway. - Science 348:

226–229.

Moore, B. R. et al. 2016. Critically evaluating the theory and performance of Bayesian analysis

of macroevolutionary mixtures. - Proc. Natl. Acad. Sci. 113: 9569–9574.

Moreau, C. S. and Bell, C. D. 2013. Testing the museum versus cradle tropical biological

diversity hypothesis: phylogeny, diversification, and ancestral biogeographic range

evolution of the ants. - Evolution 67: 2240–2257.

Moritz, C. et al. 2000. Diversification of rainforest faunas: An integrated molecular approach. -

Annu. Rev. Ecol. Syst. 31: 533–563.

Morrone, J. J. 2014. Biogeographical regionalisation of the Neotropical region. - Zootaxa 3782:

1–110.

Mullen, S. P. et al. 2011. Rapid diversification and not clade age explains high diversity in

neotropical *Adelpha* butterflies. - Proc. R. Soc. B Biol. Sci. 278: 1777–1785.

Myers, N. et al. 2000. Biodiversity hotspots for conservation priorities. - Nature 403: 853–858.

O'Dea, A. et al. 2016. Formation of the Isthmus of Panama. - Sci. Adv. 2: e1600883.

Ogilvie, H. A. et al. 2017. StarBEAST2 brings faster species tree inference and accurate

estimates of substitution rates. - Mol. Biol. Evol. 34: 2101–2114.

Penz, C. M. 2007. Evaluating the monophyly and phylogenetic relationships of Brassolini genera

(Lepidoptera, Nymphalidae). - Syst. Entomol. 32: 668–689.

Penz, C. M. 2008. Phylogenetic revision of *Eryphanis* Boisduval, with a description of a new

species from Ecuador (Lepidoptera, Nymphalidae). - Insecta mundi 0035: 1–25.

Penz, C. M. 2009a. Phylogeny of *Dasyophthalma* butterflies (Lepidoptera, Nymphalidae,

Brassolini). - Insecta mundi 0069: 1–12.

Penz, C. M. 2009b. The phylogeny of *Opoptera* butterflies, and an assessment of the systematic

position of *O. staudingeri* (Lepidoptera, Nymphalidae). - Zootaxa 1985: 1–20.

Penz, C. M. et al. 2011a. Neotropical *Blepolenis* butterflies: wing pattern elements, phylogeny,

and Pleistocene diversification (Lepidoptera, Nymphalidae). - Zootaxa 2897: 1–17.

Penz, C. M. et al. 2011b. A new *Orobrassolis* butterfly (Nymphalidae, Brassolini): a casualty of

habitat destruction? - Zootaxa 2740: 35–43.

Penz, C. M. et al. 2013. Adult and early-stage characters of Brassolini contain conflicting

phylogenetic signal (Lepidoptera, Nymphalidae). - Syst. Entomol. 38: 316–333.

Penz, C. et al. 2015. Butterfly dispersal across Amazonia and its implication for biogeography. -

Ecography 38: 410–418.

Penz, C. M. et al. 2017. Documenting diversity in the Amazonian butterfly genus *Bia*

(Lepidoptera, Nymphalidae). - Zootaxa 4258: 201–237.

Pereira Martins, L. et al. 2017. Butterflies of Amazon and Cerrado remnants of Maranhão,

Northeast Brazil. - Biota Neotrop. 17: e20170335.

Pérez-Escobar, O. A. et al. 2019. The origin and diversification of the hyperdiverse flora in the

Chocó biogeographic region. - Front. Plant Sci. 10: 1328.

Pérez, J. H. et al. 2017. Long term survey of the butterfly fauna of Curitiba, Paraná, Brazil: how

does a scientific collection gather local biodiversity information? (Lepidoptera:

Papilionoidea). - Shil. Rev. Lepidopterol. 45: 433–446.

Pinheiro, C. E. G. and Emery, E. de O. 2006. As borboletas (Lepidoptera: Papilionoidea e

Hesperioidea) da área de Proteção Ambiental do Gama e Cabeça de Veado (Distrito

Federal, Brasil). - Biota Neotrop. 6: bn01506032006.

Prates, I. et al. 2016. A mid-Pleistocene rainforest corridor enabled synchronous invasions of the

Atlantic Forest by Amazonian anole lizards. - Mol. Ecol. 25: 5174–5186.

R Core Team 2019. R: A language and environment for statistical computing. Version 3.5.3.:

http://www.r-project.org.

Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. - PLoS One 9: e89543.

Rabosky, D. L. et al. 2014. BAMMtools: An R package for the analysis of evolutionary

dynamics on phylogenetic trees. - Methods Ecol. Evol. 5: 701–707.

Rabosky, D. L. et al. 2017. Is BAMM flawed? Theoretical and practical concerns in the analysis

of multi-rate diversification. - Syst. Biol. 66: 477–498.

Rambaut, A. et al. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. -

Syst. Biol. 67: 901–904.

Ree, R. H. et al. 2005. A likelihood framework for inferring the evolution of geographic range on

phylogenetic trees. - Evolution 59: 2299–2311.

Ribas, C. C. et al. 2012. A palaeobiogeographic model for biotic diversification within Amazonia

over the past three million years. - Proc. R. Soc. B Biol. Sci. 279: 681–689.

Roy, K. and Goldberg, E. E. 2007. Origination, extinction, and dispersal: integrative models for

understanding present-day diversity gradients. - Am. Nat. 170: S71–S85.

Santos, J. P. dos et al. 2011. Fruit-feeding butterflies guide of subtropical Atlantic Forest and

Araucaria Moist Forest in State of Rio Grande do Sul, Brazil. - Biota Neotrop. 11: 253–274.

Santos, J. P. dos et al. 2018. Atlantic butterflies: a data set of fruit-feeding butterfly communities

from the Atlantic forests. - Ecology 99: 2875.

Shirai, L. T. et al. 2017. Natural history of *Selenophanes cassiope guarany* (Lepidoptera:

Nymphalidae: Brassolini): an integrative approach, from molecules to ecology. - Ann.

717  Entomol. Soc. Am. 110: 145–159.

718 Silva, A. R. M. et al. 2012. Borboletas frugívoras (Lepidoptera: Nymphalidae) de uma área

719  urbana (Área de Proteção Especial Manancial Cercadinho) em Belo Horizonte, Minas

720  Gerais, Brasil. - Biota Neotrop. 12: 292–297.

721 Simon, M. F. et al. 2009. Recent assembly of the Cerrado, a neotropical plant diversity hotspot,

722  by in situ evolution of adaptations to fire. - Proc. Natl. Acad. Sci. 106: 20359–20364.

723 Soldati, D. et al. 2019. Butterfly fauna (Lepidoptera, Papilionoidea) in a heterogeneous area

724  between two biodiversity hotspots in Minas Gerais, Brazil. - Pap. Avulsos Zool. 59:

725  e20195902.

726 Stebbins, G. L. 1974. Flowering Plants: Evolution above the Species Level. - Harvard University

727  Press.

728 Stichel, H. 1916. Neues über Genus *Narope* Westw. - Neue Beiträge zur Syst. Insektenkd. 1: 7–

729  8.

730 Strömberg, C. A. E. 2011. Evolution of grasses and grassland ecosystems. - Annu. Rev. Earth

731  Planet. Sci. 39: 517–544.

732 Thomé, M. T. C. et al. 2016. Recurrent connections between Amazon and Atlantic forests shaped

733  diversity in Caatinga four-eyed frogs. - J. Biogeogr. 43: 1045–1056.

734 Trujillo-Arias, N. et al. 2017. The niche and phylogeography of a passerine reveal the history of

735  biological diversification between the Andean and the Atlantic forests. - Mol. Phylogenet.

736  Evol. 112: 107–121.

737 Tufto, J. et al. 2012. Estimating Brownian motion dispersal rate, longevity and population

738  density from spatially explicit mark-recapture data on tropical butterflies. - J. Anim. Ecol.

739  81: 756–769.

Vilhena, D. A. and Antonelli, A. 2015. A network approach for identifying and delimiting

biogeographical regions. - Nat. Commun. 6: 6848.

Wahlberg, N. and Wheat, C. W. 2008. Genomic outposts serve the phylogenomic pioneers:

designing novel nuclear markers for genomic DNA extractions of Lepidoptera. - Syst. Biol.

57: 231–242.

Wahlberg, N. et al. 2009. Nymphalid butterflies diversify following near demise at the

Cretaceous/Tertiary boundary. - Proc. R. Soc. B Biol. Sci. 276: 4295–4302.

Werneck, F. P. 2011. The diversification of eastern South American open vegetation biomes:

historical biogeography and perspectives. - Quat. Sci. Rev. 30: 1630–1648.

Wesselingh, F. P. and Salo, J. A. 2006. A Miocene perspective on the evolution of the

Amazonian biota. - Scr. Geol. 133: 439–458.

Wiens, J. J. and Donoghue, M. J. 2004. Historical biogeography, ecology and species richness. -

Trends Ecol. Evol. 19: 639–644.

Xing, Y. and Ree, R. H. 2017. Uplift-driven diversification in the Hengduan Mountains, a

temperate biodiversity hotspot. - Proc. Natl. Acad. Sci. 114: E3444–E3451.

Zacca, T. and Bravo, F. 2012. Borboletas (Lepidoptera: Papilionoidea e Hesperioidea) da porção

norte da Chapada Diamantina, Bahia, Brasil. - Biota Neotrop. 12: 117–126.

Zizka, A. et al. 2019. CoordinateCleaner: Standardized cleaning of occurrence records from

biological collection databases. - Methods Ecol. Evol. 10: 744–751.

Supplementary material (available online as Appendix 1 at website). Appendix 1.

Figures

Fig. 1: Map of the Neotropics and the defined bioregions based on occurrence data of Brassolini. Inset map shows the most stable bioregions under different cluster cost values by the software Infomap Bioregions using validated geo-coordinates from GBIF and ATLANTIC BUTTERFLIES databases. Bioregions are delineated based on Morrone's (2014) Neotropical provinces, encompassing the Mesoamerican plus Pacific dominions, except for three provinces east of the northern Andes; the Paraná dominion (Brazilian Atlantic Forest); the remaining dominions and provinces where Brassolini occurs. Shapefile from Löwenberg-Neto (2014).

Fig. 2: Time-calibrated species tree of the butterfly tribe Brassolini. The posterior probabilities

for all genera and among genera are displayed next to the crown node. Ancestral range

probability based on the DEC model and 10,000 biogeographical stochastic mappings is depicted

as a pie chart following colors in legend. Images of representative butterflies of all Brassolini

genera, all at the same scale: From top to down, *Bia actorion* (DeVries collection, PJD),

Dynastor darius (Milwaukee Public Museum, MPM), *Dasyophthalma creusa* (MPM), *Opoptera*

fruhstorferi (American Museum of the Natural History, AMNH), *Narope panniculus* (AMNH),

Caligopsis seleucida (PJD), *Eryphanis aesacus* (MPM), *Caligo martia* (MPM), *Brassolis astyra*

(MPM), *Selenophanes cassiope* (PJD), *Catoblepia berecynthia* (MPM), *Penetes pamphanis*

(MPM), *Mielkella singularis* (AMNH), *Orobrassolis ornamentalis* (MPM), *Blepolenis bassus*

(Museu de Zoologia - USP), *Opsiphanes quiteria* (PJD).

Ancestral range

- 🔴 Pr. one area = 1
- 🔴 Most probable range = 1 area
- 🔴 Most probable range = 2 areas
- ⚪ Most probable range > 2 areas

Bioregion

- 🟩 Mesoamerica + NW Andes
- 🟥 Amazonia
- 🟧 Dry diagonal
- 🟦 Atlantic Forest

Bia actorion
Bia rebeli
Bia caelestis
Bia decaerulea
Opoptera sulcius
Opoptera syme
Opoptera fruhstorferi
Opoptera staudingeri
Opoptera aorsa
Opoptera hilaris
Opoptera arsippe
Opoptera bracteolata
Narope panniculus
Narope testacea
Narope nesope
Narope cyllene
Narope sutor **rev. comb.**
Narope cyllarus
Narope cyllabarus
Narope cyllastros
Dynastor macrosiris
Dynastor darius
Dynastor napoleon
Dasyophthalma creusa
Dasyophthalma vertebralis
Dasyophthalma rusina
Dasyophthalma geraensis
Caligopsis seleucida
Eryphanis gerhardi
Eryphanis reevesii
Eryphanis automedon
Eryphanis lycomedon
Eryphanis bubocula
Eryphanis aesacus
Eryphanis opimus
Eryphanis greeneyi
Eryphanis zolvizora
Caligo martia
Caligo arisbe
Caligo euphorbus
Caligo idomeneus
Caligo uranus
Caligo atreus
Caligo beltrao
Caligo placidianus
Caligo illioneus
Caligo teucer
Caligo memnon
Caligo oedipus
Caligo eurilochus
Caligo brasiliensis
Caligo telamonius
Brassolis haenschi
Brassolis isthmia
Brassolis dinizi
Brassolis granadensis
Brassolis sophorae
Brassolis astyra
Selenophanes orgetorix **new comb.**
Selenophanes josephus
Selenophanes cassiope
Catoblepia amphirhoe
Catoblepia xanthus
Catoblepia soranus
Catoblepia xanthicles
Catoblepia versitincta
Catoblepia berecynthia
Penetes pamphanis
Mielkella singularis
Orobrassolis ornamentalis
Blepolenis batea
Blepolenis bassus
Blepolenis catharinae
Opsiphanes bogotanus
Opsiphanes tamarindi
Opsiphanes cassiae
Opsiphanes zelotes
Opsiphanes blythekitzmillerae
Opsiphanes boisduvalii
Opsiphanes invirae
Opsiphanes quiteria
Opsiphanes cassina
Opsiphanes jacobsorum

25   20   15   10   5   0 **Ma**

Fig. 3: Colonization rates through-time based on 10,000 biogeographical stochastic mappings in

BioGeoBEARS. Rates are displayed for a select pair of areas. Source areas (dispersal from)

follow the color in the legend. Solid lines are the median values, colored ribbons are the lower

and upper quartiles (0.25 and 0.75 quantiles), and dashed lines are the 0.1 and 0.9 quantiles.

Fig. 4: Within-area speciation through-time based on 100 posterior phylogenies and 100

biogeographical stochastic mappings (BSMs) in BioGeoBEARS. A) The relative number of

within-area cladogenesis through time was calculated using a formula modified from Xing and

Ree (2017) based on cladogenesis events occurring only in one area. Solid lines represent

median values, colored ribbons represent lower and upper quartiles (0.25 and 0.75 quantiles) and

dashed lines represent 0.1 and 0.9 quantiles. The x-axis is truncated at the middle Miocene, as

most Brassolini cladogenesis events have occurred since then. B) Per-lineage speciation rates

through time estimated by BAMM were assigned to a given bioregion based on the BSMs. Solid

lines represent mean speciation rates per lineage and the colored ribbons the 5% and 95%

confidence intervals. The x-axis is truncated at the middle Miocene. C) Species (tips) specific

speciation rates as estimated by BAMM. Box plots per area show estimates from the posterior

distribution, and the cross within each box plot represents mean speciation rates per area.

Fig. 5: Summary of Brassolini colonization and speciation estimates in three time-windows, at

10, 5 and 1 Ma. Colored directional arrows represent dispersal from source areas, following

colors in legend. The width and shape of arrows represent the estimated relative colonization rate

between two areas in BioGeoBEARS, as depicted in Fig. 3. Bioregions were colored to

qualitatively represent the estimated relative cladogenesis/speciation rates, as either center of

speciation or area with comparatively lower speciation rates per time window.

Colonization and speciation through time