Exploring the Links between Specific Depression Symptoms and Brain Structure: A Network Study

Eva Hilland [1,2], **Nils Inge Landrø**[1,2], **Brage Kraft**[2], **Christian K. Tamnes**[1,3,4], **Eiko I. Fried**[5], **Luigi A. Maglanoc**[2,3], **& Rune Jonassen**[6]

[1] Department of Psychiatry, Diakonhjemmet Hospital, Oslo, Norway.

[2] Clinical Neuroscience Research Group, Department of Psychology, University of Oslo, Oslo, Norway.

[3] NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway.

[4] PROMENTA Research Center, Department of Psychology, University of Oslo, Oslo, Norway.

[5] Clinical Psychology Unit, Leiden University, Leiden, the Netherlands.

[6] Faculty of Health Sciences, Oslo Metropolitan University, Oslo, Norway.

Corresponding author: Eva Hilland, Diakonhjemmet Hospital, Department of Psychiatry, Postboks 23 Vinderen, 0319 Oslo, Norway. eva.hilland@psykologi.uio.no

Abstract

The network approach to psychopathology has recently received considerable attention, and is as a novel way of conceptualizing mental disorders as causally interacting symptoms. In this study, we modeled a joint network of depression symptoms and depression-related brain structures, using 21 symptoms and five regional brain measures. We used a mixed sample of 268 individuals previously treated for one or more major depressive episodes and never depressed individuals. The network revealed associations between brain structure and unique depressive symptoms, which may clarify relationships regarding symptomatic and biological heterogeneity in depression.

Introduction

Various patterns of structural brain abnormalities have been associated with depression, yet sensitive, specific and clinically predictive brain correlates have proven to be difficult to characterize[1]. The currently best available empirical evidence on neuroanatomical differences between patients with major depression (MDD) and healthy controls are two meta-analyses of approximately 10.000 individuals[2, 3]. These reports show widespread alterations in cortical regions and in hippocampal volume, but no associations between depression severity and brain structure. Inconsistencies in the neuroimaging literature may be explained by the fact that depression is highly heterogeneous, featuring over 50 symptoms[4], where symptom constellations may reflect different phenomena with distinct underlying biological causes[1].

Understanding the neural substrates of specific symptoms may provide important information about mechanisms underlying depression vulnerability. A growing body of research under the umbrella term 'network approach' has recently received considerable attention[5]; the approach understands and aims to model mental disorders as systems of causally interacting symptoms. So far, network studies have solely been based on symptoms, ignoring relevant neurobiological factors[6]. Here, we address this knowledge gap by modelling a joint network of depression-related brain structures and individual depression symptoms, using 21 symptoms and five regional brain measures. The sample is a mixed group of individuals that previously has been treated for one or more major depressive episodes (MDE), and never depressed individuals, with the goal to model a continuum of depression severity.

Methods

Depression symptoms were measured using the Beck Depression Inventory (BDI-II). Whole-brain volumetric segmentation and cortical surface reconstruction of MRI images were performed with FreeSurfer 5.3 (https://surfer.nmr.mgh.harvard.edu/). Five regional brain measures were selected based on the MDD case-control differences showing the largest bilateral effects in the studies from the ENIGMA MMD working group[2, 3]: hippocampal volume and cortical thickness in four regions - medial orbitofrontal cortex (mOFC), fusiform gyrus, insula and cingulate (weighted average of rostral anterior cingulate, caudal anterior cingulate and posterior cingulate). Brain structure measures were averaged across the left and right hemisphere for each participant, and z-residuals of hippocampal volume (controlling for sex and estimated intracranial volume) were calculated for further analyses. Network

structures of the 26 variables were computed using the R packages qgraph and bootnet; the graphical LASSO (least absolute shrinkage and selection operator) was used for regularization. Informed consent was obtained for all participants before enrolment and their anonymity was preserved.

Sample

This sample was drawn from two related clinical trials (Clin.gov ID NCT0265862 and NCT02931487), and a case-control research study conducted at the Department of Psychology, University of Oslo. The sample consists of 268 adult participants, 191 with at least one MDE (*M* age = 39.4 [*SD* = 13.2], 132 females, *M* education level *(*ISCED) level 6.0 [*SD* = 0.9], *M* BDI-II score 14.7 [*SD* = 10.4]) and 77 never depressed individuals (*M* age = 41.9 [*SD* = 12.9], *M* education level 5.7 [*SD* = 1.5], *M* BDI-II score 1.7 [*SD* = 2.9], 50 females). BDI-II sum score range was 0 - 49. A total of 172 subjects had experienced two or more MDE's.

Results

The symptom-brain network is depicted in *Figure 1*. All brain structures were positively inter-connected, with regularized partial correlations up to 0.41, see *Table 1.* Insula was negatively associated with *loss of interest in sex* and *sadness*. Hippocampus was negatively associated with *sadness* and *changes in appetite,* and a positively associated with *loss of interest*. Cingulate had a negative association with *sadness*, while fusiform gyrus had weak but positive associations with *self-criticism* and *crying*.

Figure 1. Depression symptom network including five brain areas. Blue lines represent positive associations, red lines negative associations, and the thickness and brightness of an edge indicate the association strength. Label descriptions: mOFC=Medial orbitofrontal cortex, CINGULATE=Rostral-,medial-, and anterior cingulate cortex, INSULA=Insula, FUSIFORM=Fusiform gyrus, HIPPOCAMP=Hippocampus, SAD=Sadness, PESS=Pessimism, FAIL=Past Failure, ANHED=Loss of Pleasure, GUILT=Guilty Feelings, PUNISH=Punishment Feelings, DISL=Self-Dislike, CRITIC=Self-Criticalness, SUIC=Suicidal Thoughts or Wishes, CRY=Crying, AGIT=Agitation, INTER=Loss of Interest, INDECISIVE=Indecisiveness, WORTH=Worthlessness, ENER=Loss of Energy, SLEEP=Changes in Sleep Pattern, IRRIT=Irritability, APPET=Changes in Appetite, CONC=Concentration Difficulty, FATIG=Tiredness or Fatigue, SEX= Loss of Interest in Sex.

Brain regions	Edge weight	Brain regions and symptoms	Edge weight
Cingulate – mOFC	0.41	Insula – Loss of interest in sex	-0.06
Cingulate – Insula	0.30	Hippocampus – Sadness	-0.04
Insula – Fusiform gyrus	0.30	Cingulate – Sadness	-0.04
Insula – Hippocampus	0.18	Hippocampus – Changes in appetite	-0.04
Fusiform gyrus – Cingulate	0.15	Hippocampus – Loss of interest	0.03
Insula – mOFC	0.11	Fusiform gyrus – Self criticism	0.02
Hippocampus – Cingulate	0.09	Fusiform gyrus – Crying	0.01
mOFC – Fusiform gyrus	0.01	Insula – Sadness	< -0.01

Table 1 Sparse partial correlations between brain structure measures, and between brain structure measures and depressive symptoms in the network model.

Discussion Here we establish the first link between individual depression symptoms and neuroanatomy using symptom network analysis. Our results broadly align with prior literature showing that depression symptoms differentially relate to important outcomes such as impairment and risk factors, and demonstrate the importance of studying specific features of depression over one heterogeneous category[5, 6]. The associations between symptoms and brain structure may reflect the heterogeneous nature of the disorder, and may offer important cues about underlying neural mechanisms in MDD. The results await replication in larger samples and other patient groups. In this study depression history was assessed retrospectively and previous MDE was classified independent of type of treatment, combination treatment, treatment response or time since the last episode. We hope the reported results can pave the way for future studies integrating neurobiological measures in network analyses, which represent a step towards validation of biomarkers.

Acknowledgements
We thank Department of Psychiatry, Diakonhjemmet Hospital for help and support with recruiting patients, and the Intervention Center, OUS for radiological assistance in MRI protocols, data acquisitions and screening for unexpected neuropathological findings. We thank Tor Endestad for establishing the infrastructure for MRI research at the Department of Psychology, University of Oslo. We want to thank our MRI research assistant Dani Beck. We also thank our external recruitment sites; Unicare, Coperiosenteret AS, Torgny Syrstad, MD, Synergi Helse AS and Lovisenberg Hospital.

Disclosure Statement

The project is supported by the South-Eastern Norway Regional Health Authority, grant number: 2015052 (to NIL), Research Council of Norway, grant number: 229135 (to NIL) and Department of Psychology, University of Oslo. CKT is funded by the Research Council of Norway, grant numbers: 223273; 288083; 230345 and the South-Eastern Norway Regional Health Authority, grant number: 2019069. NIL has received consultancy fees and travel expenses from Lundbeck. EH, BK, EF, LM CKT and RJ reports no biomedical financial interests or potential conflicts of interest.

References

Insel TR, Landis SC: Twenty-five years of progress: the view from NIMH and NINDS. Neuron 2013;80:561-567.
Schmaal L, Veltman DJ, van Erp TG, Sämann P, Frodl T, Jahanshad N, Loehrer E, Tiemeier H, Hofman A, Niessen W: Subcortical brain alterations in major depressive disorder: findings from the ENIGMA Major Depressive Disorder working group. Molecular psychiatry 2016;21:806.
Schmaal L, Hibar D, Sämann P, Hall G, Baune B, Jahanshad N, Cheung J, van Erp T, Bos D, Ikram M: Cortical abnormalities in adults and adolescents with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working Group. Molecular psychiatry 2017;22:900.
Fried EI: The 52 symptoms of major depression: Lack of content overlap among seven common depression scales. J Affect Disord 2017;208:191-197.
Borsboom D: A network theory of mental disorders. World psychiatry 2017;16:5-13.
Fried EI, Cramer AOJ: Moving Forward: Challenges and Directions for Psychopathological Network Theory and Methodology. Perspectives on Psychological Science 2017;12:999-1020.