Dendritic GABAB receptors control nonlinear information transfer along the dendro-somatic axis in layer 5 pyramidal neurons

Jan M. Schulz[1], Matthew E. Larkum[2]

[1]*Department of Biomedicine, University of Basel, Pestalozzistr. 20, 4056 Basel, Switzerland*

[2]*Institute for Biology, Humboldt-Universität zu Berlin, Charitéplatz 1, 10437 Berlin, Germany*

Correspondence:

Dr Jan M. Schulz[1,2]
Department of Biomedicine
University of Basel
Pestalozzistr. 20
4056 Basel
Switzerland
phone: +41-61-207 2727
Email: j.schulz@unibas.ch

Running title: GABAB inhibition of dendro-somatic synergy

Number of figures:	7
Number of tables:	0
Number of words:	Abstract – 199
	Introduction/Results/Discussion – 4973
	Methods - 1634
Number of Supplementary Files:	3

Key words: interhemispheric inhibition, neocortex, neuromodulation, GIRK channels, L-type calcium channels, HCN channels

Acknowledgements. We would like to thank Enrique Perez-Graci for helpful comments on the manuscript and Natalie Nevian for technical assistance. This work was supported by the Swiss National Science Foundation (SNSF, PP00A‑102721 and 31003A_130694).

Summary

Dendritic GABA$_B$ receptors (GABA$_B$Rs) mediate a slow form of interhemispheric inhibition. Surprisingly, this inhibition has no detectable effect on the somatic membrane potential of layer 5 pyramidal neurons, whereas the action potential (AP) output is robustly decreased even when the input is proximal to the cell body. To elucidate the underlying mechanisms, we systematically mapped the AP frequency-current (F-I) relationship during dual patch-clamp recordings from soma and apical dendrite. The AP output function was governed by the synergistic interaction between dendritic and somatic compartments as the local input and transfer resistance from dendrite to soma (K$_{ds}$) depended on the dendritic membrane potential. Thus, K$_{ds}$ doubled at an estimated rate of once per 28.7 mV depolarization due to HCN channel deactivation. In addition, dendritic L-type Ca^{2+} channels converted individual APs into dendritic Ca^{2+} spikes causing high-frequency bursts of APs (HFB) during large dendritic depolarization. Activation of dendritic GABA$_B$Rs greatly reduced both nonlinear mechanisms. While direct block of L-type Ca^{2+} channels reduced the number of HFBs, K$^+$ channel activation decreased voltage-dependent input and transfer resistances and decreased the AP rate under all conditions. These results highlight the powerful modulation of the input integration in pyramidal neurons by metabotropic receptor-activated K$^+$ channels.

INTRODUCTION

The mammalian cortex is a conspicuously layered structure that receives inputs from afferent subcortical and other cortical areas in a layer-specific manner (Douglas & Martin, 2004). The dendrites of pyramidal neurons typically span several layers and can therefore integrate inputs from different afferent sources. Together with the neurons' intrinsic physiological properties, this anatomical arrangement places pyramidal neurons in the ideal position to form associative computational elements of different input streams (Larkum, 2013). It has been suggested that pyramidal neurons, in particular those residing in layer 5 (L5), act as coincidence detectors of sensory feedforward inputs impinging onto the basal dendrites and higher-order feedback (e.g. attentional) inputs onto the apical tuft (Xu *et al.*, 2012; Takahashi *et al.*, 2016). One important mechanism for coincidence detection is the generation of dendritic Ca^{2+} spikes by the interaction of backpropagating APs and large dendritic postsynaptic potentials (PSP) resulting in a burst of axonal APs (Larkum *et al.*, 1999). This mechanism also contributes to the amplification of AP output during noisy somatic input currents by additional dendritic depolarization, i.e. it increases the gain of the input-AP output (F-I) relationship (Larkum *et al.*, 2004). Although other cell-intrinsic voltage-dependent mechanisms, such as the hyperpolarization and cyclic nucleotide activated (HCN) cation current (I_H), are known to shape synaptic integration (Williams & Stuart, 2000; Berger *et al.*, 2001; Harnett *et al.*, 2015), their contributions to the synergistic interaction between depolarization in the apical dendritic and perisomatic compartments are less well understood (Phillips *et al.*, 2016).

Active dendritic processing significantly contributes to sensory perception and behaviorally relevant neuronal computations (Xu *et al.*, 2012; Harnett *et al.*, 2013; Smith *et al.*, 2013; Bittner *et al.*, 2015; Takahashi *et al.*, 2016; Sheffield *et al.*, 2017). Not surprisingly, the integrative properties of pyramidal neurons are the target of multiple modulatory systems including serotonergic (Santello & Nevian, 2015), adrenergic (Liu *et al.*, 2017) and cholinergic signals (Williams & Fletcher, 2018). Thus, cholinergic inputs from the basal forebrain involved in attentional processes have recently been shown to enhance dendritic R-type Ca^{2+} channel activity via muscarinic receptor activation in L5

pyramidal neurons (Williams & Fletcher, 2018). In contrast, down-modulation of dendritic excitability by GABA$_B$ receptors (GABA$_B$Rs) involves both direct inhibition of dendritic L-type Ca^{2+} channels and activation of G protein-coupled inwardly-rectifying K$^+$ (GIRK) channels (Perez-Garci *et al.*, 2006; Palmer *et al.*, 2012; Perez-Garci *et al.*, 2013). GABA$_B$Rs mediate a slow form interhemispheric inhibition that decreases the *in vivo* AP rate specifically in L5 pyramidal neurons by about a third (Palmer *et al.*, 2012; Palmer *et al.*, 2013). This form of interhemispheric inhibition acts as silent inhibition that reduces L5 pyramidal neuron spiking without noticeable effects on the subthreshold voltage envelope at the cell body. While there is strong evidence that direct L-type Ca^{2+} channel inhibition fundamentally changes the integrative properties of pyramidal neurons (Perez-Garci *et al.*, 2006; Breton & Stuart, 2012; Palmer *et al.*, 2012; Perez-Garci *et al.*, 2013), the relative contribution of dendritic K$^+$ channels to the decreased somatic AP output has remained unclear.

In the current study, we sought to elucidate the precise contributions by both GABA$_B$R-activated effector pathways to the modulation of integrative properties of L5 pyramidal neurons.

RESULTS

We systematically mapped the F-I relationship and membrane depolarization during combined current injections into the somatic and apical dendritic compartments of thick-tufted L5b pyramidal neurons in rat (P28-40) somatosensory cortex using dual patch-clamp recordings. Somatic current injection into L5b pyramidal neurons resulted in classic type 1 action potential (AP) firing (Hodgkin, 1948). The frequency-input (F-I)-curve was well fitted by a square root function:

$$F = A * \sqrt{I - I_T} \qquad (1)$$

for all I>I$_T$; where I$_T$ denotes the current threshold and parameter *A* determines the slope, i.e. the gain of the input-output function (Fig. 1A-B). Next, we extended this spike rate model to predict what the effect of additional current injection into the apical dendrite would be on the somatic AP frequency:

$$F = A * \sqrt{I_{soma} + D * I_{dend} - I_T} \qquad (2)$$

where I$_{soma}$ and I$_{dend}$ stand for somatic and dendritic current amplitude, respectively; and

the factor D scales the relative impact of I_{dend}. We reasoned that, if the dendrite was just a passive cable, any dendritic current would depolarize the soma according to the transfer resistance from dendrite to soma ($K_{ds}= V_{soma}/I_{dend}$) measured at resting membrane potential. Thus, we used the ratio of transfer resistance to the somatic input resistance ($D=K_{ds}/R_{in}$) as an approximation for D (Fig. 1C). This spike rate model is henceforth called the passive dendrite spike rate (PDSR) model. To directly test the PDSR model's predictions, the F-I relationship were experimentally obtained during simultaneous somatic and dendritic patch-clamp recordings (Fig.1D). The experimentally observed F-I relationship exhibited considerably higher AP rates than predicted by the PDSR model (Fig. 1E&F). To determine the parameters that caused this deviation, we fitted equation (2) to the data and compared the obtained parameters to those of the PDSR model based on somatic current injections (Extended Data Fig. 1). The obtained gain and current threshold (I_T) were very similar between fits to purely somatic compared to combined somatic and dendritic current injections (Extended Data Fig. 1E). However, the experimentally established D (0.48 ± 0.03, n=25 neurons) was significantly larger than K_{ds}/R_{in} (0.29 ± 0.02, $P<0.0001$; Wilcoxon signed rank test). The strong nonlinear effects of dendritic current injections on spike patterns that included dendritic Ca^{2+} spikes and somatic burst firing (Fig. 1D) clearly indicated that active dendritic mechanisms significantly contributed to the F-I relationship.

Dendritic GABA_BRs minimize active dendritic contribution to the F-I relationship
Modulation via dendritic GABA_BRs is known to suppress dendritic Ca^{2+} spikes in L5 pyramidal neuron (Perez-Garci *et al.*, 2006; Palmer *et al.*, 2012; Perez-Garci *et al.*, 2013). We tested the effect of dendritic GABA_BR activation on the F-I relationship by puffing baclofen (50 µM) directly onto the apical dendrite (Fig. 1G). As previously documented (Perez-Garci *et al.*, 2006 ; Palmer *et al.*, 2012), baclofen prevented the activation of dendritic Ca^{2+} spikes and decreased the overall AP output (Fig. 1H). The F-I relationship during dendritic GABA_BR activation was surprisingly similar to the PDSR model's predictions (Fig. 1I). Thus, the summed squared error (SSE) of the experimentally observed F-I relationship to the PDSR model was reduced ($P=0.0031$, Wilcoxon signed rank test, n=23; Extended Data Fig. 1D left). Similarly, SSE of the data fit to equation (2)

was reduced indicating greater qualitative similarity of the actual F-I relationship to equation (*2*) (*P*<0.0001, Wilcoxon signed rank test, n=23; Extended Data Fig. 1D right). Furthermore, this fit showed a significant reduction of *D* (*P*=0.0023, Wilcoxon signed rank test, n=25) and increase of the current threshold during dendritic GABA$_B$R activation (*P*<0.0001; Extended Data Fig. 1E). Together, these results showed that dendritic GABA$_B$Rs removed active dendritic contributions and established an F-I relationship closely resembling the PDSR model.

Dendritic GABA$_B$ modulation reduces nonlinearities at subthreshold membrane potentials

We next tested the effect of dendritic GABA$_B$R activation on subthreshold membrane potential deflections in the absence of AP-associated dendritic Ca^{2+} spikes. Families of increasing current steps were injected into either dendrite or soma before and after puffing baclofen locally onto the apical dendrite (Fig.2A and B). When current steps of increasing amplitude were injected into the dendrite, the dendritic as well as the somatic membrane depolarizations increased in a supralinear fashion. Thus, the apparent local input resistance at the dendrite increased with increasing current step size (Fig. 2C, black). This effect was even stronger for the transfer resistance K$_{ds}$ (Fig. 2D, black). Dendritic GABA$_B$R activation strongly suppressed this nonlinear increase of both the local input resistance in the dendrites (*P*=0.0017; extra sum-of-squares F test, F(1,110)=10.38; Fig.2C, orange) and the transfer resistance (*P*<0.0001; F(1,236)=35.07; Fig.2D, orange). In addition, current steps of increasing amplitude into the soma induced supralinearly increasing membrane depolarizations in the soma. While dendritic GABA$_B$R activation had no effect on the somatic input resistance measured at resting membrane potential (Fig. 2E), it did reduce the nonlinear increase of the somatic input resistance significantly (*P*=0.0008; extra sum-of-squares F test, F(1,119)=11.82). These effects may explain the previously made observation that dendritic GABA$_B$R activation acts as silent inhibition without an apparent effect on the somatic input resistance (Palmer *et al.*, 2012). To better understand the nonlinearities in the integrative properties of L5 pyramidal neurons, we proceeded to map systematically the current-voltage relationship at subthreshold membrane potentials.

The transfer resistance is voltage-dependent

In linear systems, the transfer resistance is independent of the direction in which it is measured, i.e. it is transitive. In agreement with this prediction, K_{ds} closely matched K_{sd} (Fig. 3A and B). In addition, the transfer resistance is expected to depend on the distance between the two recordings sites and to decrease with increasing distance. Indeed, K_{ds} normalized to the somatic R_{in} showed a monoexponential decay with a length constant of 383 µm (Fig. 3C). However, larger current steps injected into the dendrite resulted in supralinear increases of the somatic membrane potential fluctuations (Fig. 3A). Plotting the observed K_{ds} normalized by the K_{ds} measured at the resting membrane potential versus the steady-state membrane potential locally induced by the current step showed that K_{ds} strongly depended on the membrane potential (Fig. 3D). On average, depolarization of 28.7 mV in the dendrite caused the K_{ds} to increase by a factor of two (fitted line). The local dendritic input resistance (K_{dd}) was also modulated by the dendritic membrane potential (Fig. 3E); however, this voltage-dependence consisted of a doubling every 59.5 mV which was a significantly lower rate than for K_{ds} ($P<0.001$; extra sum-of-squares F test, F(3,783)=90.35).

What are the mechanisms underlying the voltage-dependence of K_{ds}? A simple linear two-compartment model without any voltage-dependent conductances cannot reproduce the observed behavior (Fig. 3F). A good candidate for mediating the underlying nonlinear conductance are HCN channels, which are strongly expressed in dendrites of pyramidal neurons (Magee, 1998; Williams & Stuart, 2000; Berger *et al.*, 2001; Harnett *et al.*, 2015). The HCN-mediated conductance is highly voltage-dependent with about half of the channels activated at -80 to -95 mV and an *e*-fold current response per 7 to 10 mV (Solomon & Nerbonne, 1993; Berger *et al.*, 2001). In neocortical L5 and CA1 pyramidal, HCN channels significantly contribute to the resting conductance in dendrites (Magee, 1998; Williams & Stuart, 2000; Berger *et al.*, 2001). As the reversal potential of the HCN-mediated current, I_H, is close to -40 mV, I_H is the main driver for a more depolarized membrane potential in dendrites (Berger *et al.*, 2001). Including I_H in the two-compartmental model improved fits to the data (Fig. 3F).

As expected, the fits showed that there was a high density of I_H specifically in the dendrites (135.4 ± 26.1 nS, versus 52.5 ± 8.8 nS in the soma, P=0.003, n=11 neurons). These results clearly demonstrated that voltage-dependent conductances in the somato-dendritic compartments of L5 pyramidal neurons are the basis for the strong voltage-dependence of the transfer resistance.

The voltage-dependence of the transfer resistance could potentially explain the large difference between the parameter *D* derived from the fits of the F-I relationship and K_{ds} normalized by R_{in} (Extended Data Fig. 1E). K_{ds} had been measured on small dendritic current steps to minimize the influence of voltage-dependent nonlinearities. If K_{ds} was measured during large dendritic current injections inducing somatic depolarizations close to the AP threshold, K_{ds}/R_{in} provided a much better match of *D* derived from the actual F-I relationship (Fig. 3G). Taken together, these results highlighted the synergistic interaction between depolarization in the somatic and dendritic compartments even at subthreshold membrane potentials independent of Ca^{2+} channel activation.

GABA$_B$R activated K$^+$ channels diminish voltage-dependent increase of input and transfer resistance

GABA$_B$R activation is well known to increase conductance through GIRK channels (Newberry & Nicoll, 1984; Palmer *et al.*, 2012) and has been reported to recruit additionally other K$^+$ channels such as two-pore domain K$^+$ channels in L5 pyramidal neurons and entorhinal stellate cells (Deng *et al.*, 2009; Breton & Stuart, 2017). To test whether an increase of the K$^+$-mediated membrane conductance in the dendritic compartment was sufficient to cause the experimentally observed changes in dendritic input resistance and transfer resistance after the baclofen puff, we fitted the two-compartmental model to the membrane potential responses in the control condition for all experiments in Fig. 2C (Fig. 4A-B). We then refitted the models to the membrane potential responses during baclofen ejection with the dendritic membrane resistance and the resistance between both compartments as the only free parameters. While the inter-compartmental resistance remained unchanged (27.7 ± 5.7 versus 27.6 ± 5.1 MΩ, n=5 neurons), the fits resulted in a consistent increase in the local K$^+$-mediated

membrane conductance across all experiments (16.7 ± 3.6 versus 11.5 ± 2.2 nS) and were sufficient to induce the experimentally observed changes in the dendritic input resistance and transfer resistance (Fig. 4C-D). These results suggested that K^+ channel activation in the dendrite was a major contributor to reduced voltage-dependence of input and transfer resistances during dendritic $GABA_BR$ activation.

GIRK channel-activation in dendrites effectively inhibits action potential output

We tested experimentally the contribution of GIRK channels by applying the GIRK antagonist tertiapin (0.5 μM). Baclofen puffed onto the apical dendrite induced a small membrane potential hyperpolarization of -1.6 ± 0.2 mV and -0.7 ± 0.2 mV (n=15) in the apical dendrite and soma, respectively. Tertiapin reduced the baclofen-induced hyperpolarization by 54.1 ± 7.6% in the dendrite, consistent with a functionally significant contribution of GIRK channels to the baclofen-induced hyperpolarization (Extended Data Fig.4A). The incomplete block of the baclofen-induced hyperpolarization by tertiapin is in agreement with recent reports suggesting that other K^+ channels such as two-pore domain K^+ channels also contribute to $GABA_B$-mediated membrane potential hyperpolarizations in L5 pyramidal neurons and entorhinal stellate cells (Deng *et al.*, 2009; Breton & Stuart, 2017). Importantly, tertiapin partially blocked the effect of baclofen on the voltage-dependence of input and transfer resistances (Fig.4F-J), suggesting that GIRK channel activation in the dendrite was an important factor that mediated the effect of dendritic $GABA_BR$ activation. We proceeded to test the effect of GIRK channel activation on the F-I relationship.

Bath application of tertiapin increased the somatic R_{in} (Extended Data Fig.4B). This resulted in greater cellular excitability reflected by increased values of the F-I gain (Extended Data Fig.4C). These observations suggested that there was a baseline activity of GIRK channels in the control condition. In the presence of tertiapin, the F-I relationship showed a strong nonlinear deviation from the PDSR model similar to the control condition (Fig. 5C). We proceeded to test the effect of the baclofen puff in presence of tertiapin (Fig. 5D-F). In the presence of tertiapin, baclofen was less effective in reducing AP output as shown by ANOVA analyses of the baclofen-induced AP rate reduction (Fig. 5G). Thus, there was a main effect of tertiapin (P=0.0131,

F(1,6)=12.14, n =7), as well as a statistically significant interaction between tertiapin and stimulation-intensity (P=0.0083, F(3,18)=5.34). Post-hoc tests revealed that tertiapin reduced the effect of baclofen for all somatic current step sizes except 1000 pA, where the effect of baclofen was small anyway (Fig. 5G). Together, these results clearly demonstrated that dendritic $GABA_B$R-mediated GIRK channel activation significantly contributed to reduced somatic AP output. The underlying mechanism was the modulation of the nonlinear integrative properties of the L5 pyramidal neuron rather than a pronounced impact on the somatic input resistance or membrane potential.

The role of L-type Ca^{2+}-channels in $GABA_B$R-mediated inhibition

The previous results highlighted the importance of the nonlinear integrative properties of L5 pyramidal neurons at subthreshold membrane potentials. In how far was the F-I relationship shaped by dendritic Ca^{2+} spikes? To test this, we pharmacologically inhibited L-type Ca^{2+} channels by bath application of nimodipine (10 µM). Nimodipine inhibited dendritic Ca^{2+} spikes (cf. Fig. 6A vs. D) and hence reduced factor *D* of the F-I relationship (*P*=0.047, n=7, Wilcoxon signed rank test; Fig. 6L). However, the overall spike rate was not decreased (Fig. 6B vs. E) and the F-I relationship continued to significantly deviate from the PDSR model (Fig. 6C vs. F). Analysis of the inter-spike interval (ISI) distribution showed that nimodipine specifically reduced the number of fast ISIs of <15 ms emitted during Ca^{2+} spike-induced high frequency bursts (HFB) and resulted in an increased fraction of intermediate ISIs of 20 to 40 ms (Fig.6J). At the same time, the prominent slow afterhypolarzations (sAHP) following HFBs associated with particularly long ISIs (arrows in Fig.6A) also disappeared reflecting decreased activation of Ca^{2+} activated K^+ channels in agreement with previous reports (Moyer *et al.*, 1992).

Next, we tested whether the effect of puffed baclofen was decreased in the presence of nimodipine (Fig. 6G-I). Similar to the control situation, baclofen reduced the overall AP output (Fig. 6H) and rendered the F-I relationship more similar to the PDSR model (Fig. 6I). Thus, puffed baclofen not only prevented HFBs but also shifted the whole ISI distribution towards longer ISIs (Fig.6J). ANOVA analyses of the spike rate reduction after the baclofen puff in the absence and presence of nimodipine within the

same neurons did not show a main effect of nimodipine (P=0.12, F(1,6)=3.38), but revealed a significant interaction between nimodipine and stimulation intensity (P=0.0166, F(3,18)=4.45). Post-hoc tests revealed that the presence of nimodipine moderately decreased the impact of baclofen at intermediate somatic current levels (Fig. 6K).

Taken together, these results show that direct inhibition of L-type Ca^{2+}-channels by dendritic $GABA_BRs$ plays only a minor role in shaping the F-I relationship.

Dendritic $GABA_BR$ activation decouples dendritic input from somatic AP output

The conclusions so far were derived from experiments involving linear current steps. Is the direct inhibition of L-type Ca^{2+}-channels by dendritic $GABA_BRs$ more prominent during more realistic time-varying input patterns? To test this, we injected current into the soma and dendrite in the shape of responses recorded *in vivo* during contralateral hind limb stimulation (Fig. 7A; Palmer et al., 2012). Combining these dendritic and somatic current waves at increasing amplitudes resulted in a complex pattern of evoked APs that included HFBs of two or more APs with an ISI of <15 ms. High-frequency bursts occurred preferentially during episodes with large dendritic current amplitudes at well-defined time points (Fig. 7B and C, Extended Data Fig. 7B). Under control conditions, HFBs were commonly associated with dendritic Ca^{2+} spikes, while APs separated by ISIs>15 ms rarely coincided with dendritic Ca^{2+} spikes (Fig. 7H). Activation of dendritic $GABA_BRs$ by a puff of baclofen dramatically decreased HFB rate and eliminated Ca^{2+} spikes (Fig. 7D-E and I).

What determined whether HFBs rather than individual APs were emitted in the two conditions? We analyzed the spike-triggered average (STA) of the dendritic and somatic current. To obtain a more accurate estimate of the current that effectively drove the axonal AP output, the dendritic current was scaled by *D* (I_{eff}). *D* was obtained from the fit of *eq. 2* to the F-I relationship during linear current step injections in the same neurons under the same pharmacological conditions (Fig.1). The somatic I_{eff} was set equal to the injected somatic current. This analysis showed that, on average, a combination of constant intermediate dendritic I_{eff} of 271.3 ± 20.6 pA (n=7) and a peak somatic I_{eff} of 564.4 ± 42.4 pA drove single APs outside of bursts (ISI>15 ms; Extended

data Fig. 7B). By contrast, HFBs were evoked by significantly higher levels of dendritic I_{eff} (P=0.016) in combination with somatic I_{eff} that was in the rising phase at the time of the first AP of the burst. The contribution of somatic (462.3 ± 33.2 pA, n=7) and dendritic I_{eff} (452.2 ± 35.6 pA) at the time of the first AP of the HFB was approximately equal, indicating that HFBs were often the result of the synergistic interaction between membrane depolarizations in the dendritic and somatic compartments in agreement with their close association to dendritic Ca^{2+} spikes.

The activation of dendritic GABA$_B$Rs by baclofen altered the balance towards large somatic versus relatively small dendritic I_{eff} during initiation of single APs (somatic: 685.3 ± 48.2 pA vs. dendritic: 227.1 ± 27.5 pA). This effect was even more pronounced for APs at the start of the few remaining HFBs (somatic: 845.3 ± 33.4 pA vs. dendritic: 292.1 ± 23.3 pA), completely shifting the balance towards somatic I_{eff}. This indicated that APs were only evoked at times of large somatic current, as stronger somatic current was required to compensate for less effective dendritic current.

Next, we directly tested the contribution of dendritic Ca^{2+} spikes to the initiation of HFBs. Pharmacological blockade of L-type Ca^{2+} channels by bath application of nimodipine (10 µM) decreased the relative proportion of HFB ISI from 20.3 ± 2.8% to 9.2 ± 1.3% of all ISIs (P=0.016; Wilcoxon Signed Rank test, n=7 neurons; Fig.7G). The contribution of dendritic I_{eff} to the remaining HFBs was significantly smaller than in control condition (359.9 ± 31.2 pA vs. 452.2 ± 35.6 pA; P=0.016), while significantly greater somatic current was necessary to evoke HFBs (600.1 ± 40.0 pA vs. 462.3 ± 33.2 pA; P=0.016; Extended data Fig. 7C). Direct activation of dendritic GABA$_B$Rs by baclofen puffed onto the apical dendrite further reduced the HFB ISI proportion to 2.9 ± 0.8% comparable to the baclofen-induced effect in the absence of nimodipine (2.9 ± 0.8%). Thus, inhibition of L-type Ca^{2+} channels had qualitatively the same effect on HFBs as dendritic GABA$_B$R activation.

By contrast, the inhibition of L-type Ca^{2+} channels did not change the contribution of dendritic I_{eff} to the induction of single APs outside of HFBs (P=0.81; Extended data Fig. 7B and C). This suggested that the reduction of the AP frequency in the absence HFB by dendritic GABA$_B$Rs was independent of L-type Ca^{2+} channel inhibition. To directly test this hypothesis, we restricted the analysis to episodes with low AP

frequency (<7 Hz). For this analysis, only episodes with an intermediate dendritic current level (max. 750 pA) were included to avoid any HFBs due to dendritic Ca^{2+} spikes. In episodes with low AP frequency, activation of dendritic GABA$_B$Rs by puffed baclofen had the same effect in absence or presence of nimodipine (P=0.64, n=7; Fig.7F). This observation strongly supported the conclusion from experiments involving current step injections (Fig.4 and 5) that dendritic K$^+$ channels are the main effectors of dendritic GABA$_B$Rs to decrease the AP output.

Taken together, the results from both linear current steps and time-varying inputs converge at the conclusion that inhibition of L-type Ca^{2+} channels and activation of K$^+$ channels by dendritic GABA$_B$Rs synergistically alter the nonlinear integrative properties of L5 pyramidal neurons to suppress burst firing and decrease AP output under broadly varying conditions.

DISCUSSION

By systematically mapping the input-output relationship during dual patch-clamp recordings from soma and apical dendrite, we demonstrated here for the first time that voltage-dependent local input and transfer resistances mediate a synergistic interaction between dendritic and somatic membrane depolarization. This was the principal mechanism causing a significant deviation of the F-I relationship from the predictions based on a model with a passive dendrite (PDSR). Dendritic $GABA_BR$-activated K^+ channels greatly reduced the voltage-dependent transfer resistance, while having negligible effects on the somatic input resistance and membrane depolarization at membrane potentials well below action potential threshold. This explains the surprising impact of interhemispheric inhibition on AP output in the absence of a discernable membrane hyperpolarization at the soma of L5 pyramidal neurons observed *in vivo* (Palmer et al., 2012). In contrast, the $GABA_BR$-mediated direct block of L-type calcium channels had a more restricted effect and primarily prevented HFBs. These results highlight the powerful modulation of the input integration in pyramidal neurons by $GABA_BR$-activated K^+ channels.

The significance of the voltage dependence of K_{ds}

The contribution of dendritic K^+ channels to the $GABA_BR$-mediated response was much more pronounced than previously thought (Breton & Stuart, 2012; Palmer *et al.*, 2012). The main reason for the powerful impact of the seemingly small hyperpolarization by $GABA_BR$-activated K^+ channels (Newberry & Nicoll, 1984; Palmer *et al.*, 2012; Breton & Stuart, 2017) was due to its impact on other voltage-dependent conductances in the dendrites. While the local somatic and dendritic input resistances were weakly voltage dependent, the transfer resistance from dendrite to soma (K_{ds}) strongly depended on the dendritic membrane potential doubling at an estimated rate of every 28.7 mV depolarization step (Fig.3D). Consequently, the ratio of K_{ds} to R_{in} increased and the impact of dendritic inputs on the somatic membrane potential grew supralinearly with increasing dendritic membrane potential depolarization. Incorporating K_{ds} measured at depolarized membrane potential provided a much more accurate estimate of the dendritic impact (*D*) on the somatic AP output.

The voltage-dependent deactivation of I_H appeared to be an important factor mediating the nonlinear K_{ds}. At rest, I_H decreases the impact of small EPSPs evoked in the distal dendrite at the soma (Golding *et al.*, 2005; George *et al.*, 2009) and contributes to increased compartmentalization of synaptic inputs (Harnett *et al.*, 2015) that has also been observed in human L5 neurons (Beaulieu-Laroche *et al.*, 2018). However, larger depolarization deactivates I_H thereby increasing the local input resistance and decreasing the leakage for dendritic current flowing to the somatic compartment. This mechanism dramatically increases the impact of apical dendritic inputs on the soma when the dendrite is sufficiently depolarized.

Mechanisms of GABA_BR-mediated modulation of dendritic integration

Dendritic GABA_BR activation greatly lowers the dendritic membrane resistance. Although the membrane hyperpolarization induced by the dendritic puff of baclofen appeared to be small (Breton & Stuart, 2012; Palmer *et al.*, 2012), the strong impact on the local input resistance (Fig.2C) suggests that the conductance change was much larger. GIRK channel activation locally hyperpolarizes the membrane potential and thereby recruits additional shunt by the activation of I_H. In fact, the activation of I_H may have partially masked the actual hyperpolarization induced by GIRK channels. Interestingly, HCN channels appear to be structurally and functionally associated to GABA_BRs (Schwenk *et al.*, 2016). The combined effect of GIRK and HCN channel activation decreased the transfer resistance from dendrite to soma. Under these conditions, L5 pyramidal neurons behaved similar to the predictions of the simple PDSR model. In combination with the direct block of L-type Ca^{2+} channels (Perez-Garci *et al.*, 2013), GABA_BR-activated K^+ channels strongly down-modulate the impact of dendritic input currents on somatic AP discharge under all conditions.

The presynaptic interneuron population responsible for dendritic GABA_BR activation

GABA_BR activation mediates a slow acting form of interhemispheric inhibition (Palmer *et al.*, 2012). Activation of the contralateral sensory cortex by hind limb stimulation *in vivo* causes a GABA_BR-mediated inhibitory effect on spike activity in the ipsilateral cortex of

up to 400 ms (Palmer *et al.*, 2012). Interhemispheric callosal fibers preferentially activate interneurons in the upper cortical layers including L1 (Palmer *et al.*, 2012). GABA$_B$Rs are thought to be activated by spill-over of GABA from synaptic release sites (Scanziani, 2000; Kohl & Paulsen, 2010). In principle, any dendrite targeting interneuron population could activate dendritic GABA$_B$Rs in L5 pyramidal neurons given that they release sufficient amounts of GABA to overcome the effective perisynaptic GABA reuptake mechanisms (Destexhe & Sejnowski, 1995; Thomson & Destexhe, 1999). However, cortical neurogliaform cells (NGFCs) appear to be of particular relevance for the modulation of dendritic properties via GABA$_B$Rs. Thus, they are the only known interneurons that reliably activate postsynaptic GABA$_B$Rs after a single presynaptic AP (Tamas *et al.*, 2003; Price *et al.*, 2008). Two anatomical features may contribute to the efficient GABA$_B$Rs activation: the exceptional high density of about 1 bouton per 2.5 µm axon and the greater than usual distance of boutons from their target dendrites (Olah *et al.*, 2009; Overstreet-Wadiche & McBain, 2015). Hence, GABA released by NGFCs is thought to act via volume transmission potentially affecting many postsynaptic targets simultaneously rather than by 'point-to-point' synaptic transmission. Interestingly, NGFCs also engage a particularly slow form of ionotropic inhibition that reflects the slow GABA dynamics within the synaptic cleft as well as the slow receptor kinetics of high-affinity GABA$_A$R in the postsynaptic membrane (Tamas *et al.*, 2003; Karayannis *et al.*, 2010; Schulz *et al.*, 2018). A recent study demonstrated that a large fraction of L1 interneurons that express Neuron-Derived Neurotrophic Factor (NDNF) inhibits postsynaptic pyramidal neurons by a combination of slow GABA$_A$R and GABA$_B$R-mediated IPSCs as it is typical for NGFCs (Abs *et al.*, 2018). Furthermore, NDNF+ interneurons effectively inhibited Ca^{2+} spikes evoked in the apical dendrite of L5 pyramidal neurons (Abs *et al.*, 2018). Therefore, NDNF+ interneurons and other NGFCs in the superficial cortical layers are the prime candidates to activate dendritic GABA$_B$Rs in L5 pyramidal neurons.

Converging modulatory pathways

NGFCs receive inputs from many afferent cortical and subcortical areas that project to L1 of the cortex (Abs *et al.*, 2018). In general, these inputs signal top-down information

from higher order brain areas that control the processing of bottom-up information that is locally processed in the specific cortical area (Larkum, 2013). In addition, NGFCs also receive prominent neuromodulatory inputs such as cholinergic inputs from the basal forebrain (Lee *et al.*, 2010; Abs *et al.*, 2018; Poorthuis *et al.*, 2018). These observations show that modulation by NGFCs can be recruited via multiple converging associational pathways (D'Souza & Burkhalter, 2017; Schuman *et al.*, 2019).

Besides GABA, other neuromodulators including adenosine, serotonin, acetylcholine, endocannabinoids and somatostatin can activate GIRK channels in pyramidal neurons (Takigawa & Alzheimer, 1999; Seeger & Alzheimer, 2001; Chen & Johnston, 2005; Kim & Johnston, 2015; Lucas & Armstrong, 2015; Stumpf *et al.*, 2018). The results of the present study suggests that these neuromodulatory pathways will similarly affect dendritic integration. The functional relevance of dendritic GIRK channels has recently been highlighted by the observation that the higher induction threshold for LTP in dorsal versus ventral CA1 pyramidal neurons may be caused by tonically enhanced GIRK activation in dorsal CA1 pyramidal neurons (Malik & Johnston, 2017). In neurological disorders involving increased GIRK channel expression such as trisomy of chromosome 21 (Kleschevnikov *et al.*, 2012), increased dendritic GIRK channel activation is therefore expected to contribute to both deficits in dendritic integration and synaptic plasticity.

In conclusion, the results of the present study demonstrate that $GABA_BR$-mediated activation of dendritic K^+ channels effectively modifies the nonlinear integrative properties of L5 pyramidal neurons to decrease AP output under broadly varying conditions.

METHODS

In vitro preparation

Wistar rats (P30-P46) were anaesthetized with 95% CO_2 / 5% O_2 before decapitation. The brain was then rapidly transferred to ice-cold, oxygenated artificial cerebrospinal fluid (ACSF) containing (in mM) 125 NaCl, 25 $NaHCO_3$, 2.5 KCl, 1.25 NaH_2PO_4, 1 $MgCL_2$, 25 glucose and 2 $CaCl_2$ (pH 7.4). Parasagittal slices of the primary somatosensory cortex (300 μm thick) were cut with a vibrating microslicer (Leica) and incubated at 37 ºC for 30 minutes and subsequently maintained at room temperature (~22 ºC). Dual somatic and dendritic whole-cell patch recordings were made from visually identified neurons using infared Dodt gradient contrast or oblique illumination and a CCD camera (CoolSnap ES, Roper Scientific). During recordings, slices were bathed in ACSF maintained at 33-35 ºC.

Patch-clamp recordings

Whole-cell patch-clamp recordings were obtained from thick-tufted layer 5b (L5) pyramidal neurons using a patch pipette (resistance 6-9 MΩ) filled with an intracellular solution containing (in mM): 135 K gluconate, 4 KCl, 10 mM HEPES, 10 Na_2 phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP, 0.2% biocytin, adjusted to pH 7.3-7.4 with NaOH. Dual whole-cell voltage recordings were performed from the soma and dendrite (resistance 20-40 MΩ) using Axoclamp 2A (Molecular Devices) and Dagan BVC-700A amplifiers (Dagan Corporation). Voltage was filtered at 5 kHz and digitized at 10 kHz using a BNC900 (National Instruments, Austin, TX) or ITC-18 interface (InstruTech, Port Washington, NY). Custom written igor software was used for acquisition. No correction was made for the junction potential between the bath and pipette solutions. Dual recordings were made from the soma and dendrite in current clamp mode. Current of varying amplitude was injected through either one or both electrodes simultaneously. Step currents were 1 s long, and in other experiments, two different voltage responses to contralateral-HS recorded from the soma and dendrite *in vivo* (Palmer *et al.*, 2012) were injected into the soma and dendrite, respectively.

Drugs applications

Baclofen (50 µM) was pressure ejected from a glass pipette (tip diameter: 1 µm) placed 50–100 µm distal to the dendritic patch pipette (approx. 500-800 µm from the soma). The volume directly affected by the drugs pressure ejected from the puffing pipette was estimated to have a radius of approx. 100 µm, as measured by a test application of the fluorescent indicator Alexa 594 into a brain slice. Nimodipine (Sigma) was dissolved at 20 mM in DMSO on the day of the experiment. In experiments testing GIRK channel contribution, tertiapin (Sigma) was added to the bath ACSF (0.5 µM) and puff pipette (5.0 µM).

Neuronal reconstruction

After recordings, neurons were prepared for biocytin reconstruction. The slices were placed in 4% paraformaldehyde after the experiment for up to 4 days. Slices were processed for biocytin staining to reveal the morphology of the recorded neuron. Neuronal reconstruction was then performed with Neurolucida software.

Data analysis

Data were analyzed offline using MATLAB 7.13 with Signal Processing 6.16 and Statistics 7.6 Toolboxes. Action potentials (APs) were detected using a threshold criterion when the membrane potential crossed 0 mV. The time of the maximal depolarization was saved for each AP. The spike rate was defined as number of APs during the 1 s-long current step. Interspike intervals (ISI) were calculated as the difference in time between two subsequent APs. For experiments with fluctuating current stimuli, segments of the injected current waveform around the time of each AP were saved for visualization. APs were classified into three categories: single APs (preceding ISI_{n-1} and subsequent ISI_n>15 ms), APs at the start of a burst (ISI_{n-1}>15 ms and ISI_n<15 ms), and APs during a burst (ISI_{n-1}<15 ms). The spike-triggered average (STA) was calculated as the mean of these current segments over all spikes of a defined class. For the analysis of the baclofen-induced effect on trails with low spike rates (Fig.7F), only episodes with a maximal dendritic current amplitude of 750 pA and less than 7 spikes at baseline were included.

Input and transfer resistances were determined from the slope of a regression line fitted to four mean membrane potentials produced by a series of subthreshold current pulses around resting membrane potential (-100, 0, +100, +200 pA). For transfer resistances, current was injected into either somatic or dendritic electrode and membrane potentials were measured at the opposite electrode. For experiments involving tertiapin application, K_{ds} and R_{in} were directly derived from 250 pA current injections into dendrite and soma, respectively. To test for a voltage-dependence of the input (transfer) resistance, input (transfer) resistances were determined for increasing current step amplitudes according to Ohm's law and then normalized by the input (transfer) resistances measured at rest. For this analysis, only experiments with optimal compensation of series resistances were included.

Spike rate model. The AP frequency-input current (F-I) relationship was modeled by the following equation:

$$F = A * \sqrt{I_{soma} + D * I_{dend} - I_T} \qquad\qquad (2)$$

for all I>IT; where IT denotes the current threshold, parameter A determines the gain of the input-output function and the factor *D* scales the relative impact of the dendritic current injection (I_{dend}). This equation was fitted to AP rate data in response to increasing amplitudes of 1 s long current steps using the Matlab function fminsearch. For fits to somatic current injections only (Fig. 1B; 100 pA step size), *A* and I_T were free parameters. For fits to combined somatic and dendritic current steps (Fig. 1E; 250 pA step size), *A*, I_T and *D* were free parameters. Fits to somatic current injections only provided estimates of *A* and I_T that were combined with an estimation of *D*=K_{ds}/R_{in} to predict the AP rate of combined dendritic and somatic current injections according to the passive dendrite spike rate (PDSR) model. The parameters of the PDSR model were kept constant across pharmacological conditions and were compared to parameters derived from the fits of eq. (2) to the actual data. For experiments involving tertiapin application, *A* was directly derived from the actual fit to the AP rate data in response to combined dendritic and somatic current injections. The predicted spike rate from this model was compared to the measured spike rate in the presence and absence of an additional baclofen puff onto the apical dendrites.

Regression analyses. Values for *D* obtained from fits to the experimental F-I data were compared to theoretically predicted values using nonlinear regression analysis in GraphPad Prism 6. A regression line was fitted to the scatter plot of predicted versus measured values (Fig. 3G). The Y-axis intercept was constrained to zero. An extra sum-of-square F test was used to test for a significant deviation of the fitted slope from 1 indicating that theoretically predicted values systematically deviated from values derived from fits. Similarly, exponential functions were fitted to scatter plots of normalized input (transfer) resistances versus membrane potential or current step amplitude. Extra sum-of-square F tests were used to statistically compare growth rates between different data sets.

Two-compartment model. Two compartment models were simulated in the Python-based simulation environment Brian (Goodman & Brette, 2009). Membrane resistances of dendritic and somatic compartment as well as the resistance of the connecting resistor for linear models were derived from electrophysiological measurements of somatic R_{in} (K_{ss}), dendritic R_{in} (K_{dd}) and transfer resistance K_{ds} according to the following formulas. Dendritic membrane resistance:

$$R_{md} = (K_{ss} * K_{dd} - K_{ds}^2)/(K_{ss} - K_{ds}) \tag{3}$$

Axial resistance: $\qquad R_a = R_{md} * (K_{ss} - K_{ds})/K_{ds} \tag{4}$

Somatic membrane resistance: $\quad R_{ms} = (R_a * K_{ds})/(K_{dd} - K_{ds}) \tag{5}$

The capacitance for each compartment was estimated from the exponential fit to the decaying phase of small positive current steps at the soma and dendrite, respectively.

To model the voltage-dependence of the transfer resistance, persistent sodium channels were included in the somatic compartment, and HCN (hyperpolarization-activated cyclic nucleotide-gated cation) channels mediating I_H were included in both compartments. Persistent sodium current was modeled as:

$$I_{NaP} = (E_{Na} - v) * g_{NaP_{max}} * e^{-\frac{\theta_{NaP} - v}{\sigma_{NaP}}} \tag{6}$$

θ_{NaP} was set to -57.9 mV, and σ_{NaP} was set to 6.4 mV (Amarillo *et al.*, 2014).
I_H was modeled as:

$$I_H = (E_H - v) * g_H \tag{7}$$

$$dg_H = (g_{H\infty} - g_H)/\tau_H \qquad\qquad (8)$$

$$g_{H\infty} = g_{Hmax}/(1 + e^{-\frac{\theta_H - v}{\sigma_H}}) \qquad\qquad (9)$$

θ_H was set to -80 mV, σ_H to 12 mV, and τ_H to 40 ms, similar to published values in the literature (Solomon & Nerbonne, 1993; Berger *et al.*, 2001). The density of HCN and persistent sodium channels was derived from fits of experimentally observed steady-state membrane potential in soma and dendrites in response to somatic and dendritic current steps. The fitting procedure minimized the deviation of the model's steady state voltage responses to the applied current steps from the experimentally observed steady-state membrane potential responses by adjusting six conductance densities: the somatic and the dendritic leak membrane conductance, the axial conductance (1/ R_a), somatic and dendritic g_{Hmax} and somatic g_{NaPmax}. Initial values for the first three parameters were the inverse of R_{ms}, R_{md} and R_a calculated for the linear model (eqs. *3-5*). Initial values for dendritic and somatic g_{Hmax} and somatic g_{NaPmax} were set to 1/ R_{md}, 1/ 20*R_{ms} and 1/ 5*R_{ms}, respectively. Minimization was performed by the function minimize from the scipy.optimize package using the Sequential Least Squares Programming optimization algorithm (SLSQP). For fits of membrane potential responses during baclofen ejection, dendritic leak membrane conductance and axial conductance were the only free parameters, while the other four parameters were fixed at the values fitted to control responses.

Experimental Design and Statistical Analysis. Statistical analyses were performed in GraphPad Prism 6. All pharmacological tests were within experiment comparisons, i.e. the baclofen-induced effect in the presence of nimodipine/tertiapin were compared with the baclofen-induced effect alone in the same cell. The sample size was based on our experience with the reproducibility of similar experiments (Palmer *et al.*, 2012). Data sets were analyzed with the nonparametric Wilcoxon Signed Rank and the Mann-Whitney tests for paired and unpaired data, respectively. The effect of nimodipine/tertiapin on the baclofen-induced AP frequency reduction was tested by a two-way ANOVA on normalized AP rate data. The significance level was set to $P=0.05$. All data are shown as mean ± s.e.m.. Unless stated otherwise, the number (n) of

observations indicated reflects the number cells recorded from.

REFERENCES

Abs, E., Poorthuis, R.B., Apelblat, D., Muhammad, K., Pardi, M.B., Enke, L., Kushinsky, D., Pu, D.L., Eizinger, M.F., Conzelmann, K.K., Spiegel, I. & Letzkus, J.J. (2018) Learning-Related Plasticity in Dendrite-Targeting Layer 1 Interneurons. *Neuron*, **100**, 684-699.e686.

Amarillo, Y., Zagha, E., Mato, G., Rudy, B. & Nadal, M.S. (2014) The interplay of seven subthreshold conductances controls the resting membrane potential and the oscillatory behavior of thalamocortical neurons. *J Neurophysiol*, **112**, 393-410.

Beaulieu-Laroche, L., Toloza, E.H.S., van der Goes, M.S., Lafourcade, M., Barnagian, D., Williams, Z.M., Eskandar, E.N., Frosch, M.P., Cash, S.S. & Harnett, M.T. (2018) Enhanced Dendritic Compartmentalization in Human Cortical Neurons. *Cell*, **175**, 643-651.e614.

Berger, T., Larkum, M.E. & Lüscher, H.-R. (2001) High I h Channel Density in the Distal Apical Dendrite of Layer V Pyramidal Cells Increases Bidirectional Attenuation of EPSPs. *Journal of Neurophysiology*, **85**, 855-868.

Bittner, K.C., Grienberger, C., Vaidya, S.P., Milstein, A.D., Macklin, J.J., Suh, J., Tonegawa, S. & Magee, J.C. (2015) Conjunctive input processing drives feature selectivity in hippocampal CA1 neurons. *Nat Neurosci*, **18**, 1133-1142.

Breton, J.D. & Stuart, G.J. (2012) Somatic and dendritic GABA(B) receptors regulate neuronal excitability via different mechanisms. *J Neurophysiol*, **108**, 2810-2818.

Breton, J.D. & Stuart, G.J. (2017) GABAB receptors in neocortical and hippocampal pyramidal neurons are coupled to different potassium channels. *The European journal of neuroscience*, **46**, 2859-2866.

Chen, X. & Johnston, D. (2005) Constitutively active G-protein-gated inwardly rectifying K+ channels in dendrites of hippocampal CA1 pyramidal neurons. *J Neurosci*, **25**, 3787-3792.

D'Souza, R.D. & Burkhalter, A. (2017) A Laminar Organization for Selective Cortico-Cortical Communication. *Frontiers in neuroanatomy*, **11**, 71.

Deng, P.Y., Xiao, Z., Yang, C., Rojanathammanee, L., Grisanti, L., Watt, J., Geiger, J.D., Liu, R., Porter, J.E. & Lei, S. (2009) GABA(B) receptor activation inhibits neuronal excitability and spatial learning in the entorhinal cortex by activating TREK-2 K+ channels. *Neuron*, **63**, 230-243.

Destexhe, A. & Sejnowski, T.J. (1995) G-Protein Activation Kinetics And Spillover Of Gamma-Aminobutyric-Acid May Account For Differences Between Inhibitory Responses In The Hippocampus And Thalamus. *Proceedings of the National Academy of Sciences of the United States of America*, **92**, 9515-9519.

Douglas, R.J. & Martin, K.A.C. (2004) Neuronal circuits of the neocortex. *Annual review of neuroscience*, **27**, 419-451.

George, M.S., Abbott, L.F. & Siegelbaum, S.A. (2009) HCN hyperpolarization-activated cation channels inhibit EPSPs by interactions with M-type K+ channels. *Nature Neuroscience*, **12**, 577-584.

Golding, N.L., Mickus, T.J., Katz, Y., Kath, W.L. & Spruston, N. (2005) Factors mediating powerful voltage attenuation along CA1 pyramidal neuron dendrites. *Journal of Physiology-London*, **568**, 69-82.

Goodman, D.F. & Brette, R. (2009) The brian simulator. *Front Neurosci*, **3**, 192-197.

Harnett, M.T., Magee, J.C. & Williams, S.R. (2015) Distribution and function of HCN channels in the apical dendritic tuft of neocortical pyramidal neurons. *J Neurosci*, **35**, 1024-1037.

Harnett, M.T., Xu, N.L., Magee, J.C. & Williams, S.R. (2013) Potassium channels control the interaction between active dendritic integration compartments in layer 5 cortical pyramidal neurons. *Neuron*, **79**, 516-529.

Hodgkin, A.L. (1948) The local electric changes associated with repetitive action in a non-medullated axon. *The Journal of Physiology*, **107**, 165-181.

Karayannis, T., Elfant, D., Huerta-Ocampo, I., Teki, S., Scott, R.S., Rusakov, D.A., Jones, M.V. & Capogna, M. (2010) Slow GABA transient and receptor desensitization shape synaptic responses evoked by hippocampal neurogliaform cells. *J Neurosci*, **30**, 9898-9909.

Kim, C.S. & Johnston, D. (2015) A1 adenosine receptor-mediated GIRK channels contribute to the resting conductance of CA1 neurons in the dorsal hippocampus. *J Neurophysiol*, **113**, 2511-2523.

Kleschevnikov, A.M., Belichenko, P.V., Gall, J., George, L., Nosheny, R., Maloney, M.T., Salehi, A. & Mobley, W.C. (2012) Increased efficiency of the GABAA and GABAB receptor-mediated neurotransmission in the Ts65Dn mouse model of Down syndrome. *Neurobiol Dis*, **45**, 683-691.

Kohl, M.M. & Paulsen, O. (2010) The roles of GABAB receptors in cortical network activity. *Advances in pharmacology (San Diego, Calif.)*, **58**, 205-229.

Larkum, M. (2013) A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex. *Trends Neurosci*, **36**, 141-151.

Larkum, M.E., Senn, W. & Luscher, H.R. (2004) Top-down dendritic input increases the gain of layer 5 pyramidal neurons. *Cerebral Cortex*, **14**, 1059-1070.

Larkum, M.E., Zhu, J.J. & Sakmann, B. (1999) A new cellular mechanism for coupling inputs arriving at different cortical layers. *Nature*, **398**, 338-341.

Lee, S., Hjerling-Leffler, J., Zagha, E., Fishell, G. & Rudy, B. (2010) The largest group of superficial neocortical GABAergic interneurons expresses ionotropic serotonin receptors. *J Neurosci*, **30**, 16796-16808.

Liu, Y., Cui, L., Schwarz, M.K., Dong, Y. & Schlüter, O.M. (2017) Adrenergic Gate Release for Spike Timing-Dependent Synaptic Potentiation. *Neuron*, **93**, 394-408.

Lucas, S.J. & Armstrong, D.L. (2015) Protein phosphatase modulation of somatostatin receptor signaling in the mouse hippocampus. *Neuropharmacology*, **99**, 232-241.

Magee, J.C. (1998) Dendritic Hyperpolarization-Activated Currents Modify the Integrative Properties of Hippocampal CA1 Pyramidal Neurons. *The Journal of Neuroscience*, **18**, 7613-7624.

Malik, R. & Johnston, D. (2017) Dendritic GIRK Channels Gate the Integration Window, Plateau Potentials, and Induction of Synaptic Plasticity in Dorsal But Not Ventral CA1 Neurons. *J Neurosci*, **37**, 3940-3955.

Moyer, J.R., Jr., Thompson, L.T., Black, J.P. & Disterhoft, J.F. (1992) Nimodipine increases excitability of rabbit CA1 pyramidal neurons in an age- and concentration-dependent manner. *J Neurophysiol*, **68**, 2100-2109.

Newberry, N.R. & Nicoll, R.A. (1984) Direct hyperpolarizing action of baclofen on hippocampal pyramidal cells. *Nature*, **308**, 450-452.

Olah, S., Fule, M., Komlosi, G., Varga, C., Baldi, R., Barzo, P. & Tamas, G. (2009) Regulation of cortical microcircuits by unitary GABA-mediated volume transmission. *Nature*, **461**, 1278-1281.

Overstreet-Wadiche, L. & McBain, C.J. (2015) Neurogliaform cells in cortical circuits. *Nat Rev Neurosci*, **16**, 458-468.

Palmer, L.M., Schulz, J.M. & Larkum, M.E. (2013) Layer-specific regulation of cortical neurons by interhemispheric inhibition. *Communicative & integrative biology*, **6**, e23545.

Palmer, L.M., Schulz, J.M., Murphy, S.C., Ledergerber, D., Murayama, M. & Larkum, M.E. (2012) The cellular basis of GABA(B)-mediated interhemispheric inhibition. *Science*, **335**, 989-993.

Perez-Garci, E., Gassmann, M., Bettler, B. & Larkum, M.E. (2006) The GABA(B1b) isoform mediates long-lasting inhibition of dendritic Ca2+ spikes in layer 5 somatosensory pyramidal neurons. *Neuron*, **50**, 603-616.

Perez-Garci, E., Larkum, M.E. & Nevian, T. (2013) Inhibition of dendritic Ca2+ spikes by GABAB receptors in cortical pyramidal neurons is mediated by a direct Gi/o-beta-subunit interaction with Cav1 channels. *J Physiol*, **591**, 1599-1612.

Phillips, W.A., Larkum, M.E., Harley, C.W. & Silverstein, S.M. (2016) The effects of arousal on apical amplification and conscious state. *Neuroscience of consciousness*, **2016**, niw015.

Poorthuis, R.B., Muhammad, K., Wang, M., Verhoog, M.B., Junek, S., Wrana, A., Mansvelder, H.D. & Letzkus, J.J. (2018) Rapid Neuromodulation of Layer 1 Interneurons in Human Neocortex. *Cell Rep*, **23**, 951-958.

Price, C.J., Scott, R., Rusakov, D.A. & Capogna, M. (2008) GABA(B) receptor modulation of feedforward inhibition through hippocampal neurogliaform cells. *J Neurosci*, **28**, 6974-6982.

Santello, M. & Nevian, T. (2015) Dysfunction of cortical dendritic integration in neuropathic pain reversed by serotoninergic neuromodulation. *Neuron*, **86**, 233-246.

Scanziani, M. (2000) GABA spillover activates postsynaptic GABA(B) receptors to control rhythmic hippocampal activity. *Neuron*, **25**, 673-681.

Schulz, J.M., Knoflach, F., Hernandez, M.C. & Bischofberger, J. (2018) Dendrite-targeting interneurons control synaptic NMDA-receptor activation via nonlinear alpha5-GABAA receptors. *Nature communications*, **9**, 3576.

Schuman, B., Machold, R.P., Hashikawa, Y., Fuzik, J., Fishell, G.J. & Rudy, B. (2019) Four Unique Interneuron Populations Reside in Neocortical Layer 1. *J Neurosci*, **39**, 125-139.

Schwenk, J., Perez-Garci, E., Schneider, A., Kollewe, A., Gauthier-Kemper, A., Fritzius, T., Raveh, A., Dinamarca, M.C., Hanuschkin, A., Bildl, W., Klingauf, J., Gassmann, M., Schulte, U., Bettler, B. & Fakler, B. (2016) Modular composition and dynamics of native GABAB receptors identified by high-resolution proteomics. *Nat Neurosci*, **19**, 233-242.

Seeger, T. & Alzheimer, C. (2001) Muscarinic activation of inwardly rectifying K(+) conductance reduces EPSPs in rat hippocampal CA1 pyramidal cells. *J Physiol*, **535**, 383-396.

Sheffield, M.E.J., Adoff, M.D. & Dombeck, D.A. (2017) Increased Prevalence of Calcium Transients across the Dendritic Arbor during Place Field Formation. *Neuron*, **96**, 490-504.e495.

Smith, S.L., Smith, I.T., Branco, T. & Häusser, M. (2013) Dendritic spikes enhance stimulus selectivity in cortical

neurons in vivo. *Nature*, **503**, 115-120.

Solomon, J.S. & Nerbonne, J.M. (1993) Hyperpolarization-activated currents in isolated superior colliculus-projecting neurons from rat visual cortex. *J Physiol*, **462**, 393-420.

Stumpf, A., Parthier, D., Sammons, R.P., Stempel, A.V., Breustedt, J., Rost, B.R. & Schmitz, D. (2018) Cannabinoid type 2 receptors mediate a cell type-specific self-inhibition in cortical neurons. *Neuropharmacology*, **139**, 217-225.

Takahashi, N., Oertner, T.G., Hegemann, P. & Larkum, M.E. (2016) Active cortical dendrites modulate perception. *Science*, **354**, 1587-1590.

Takigawa, T. & Alzheimer, C. (1999) G protein-activated inwardly rectifying K+ (GIRK) currents in dendrites of rat neocortical pyramidal cells. *J Physiol*, **517 (Pt 2)**, 385-390.

Tamas, G., Lorincz, A., Simon, A. & Szabadics, J. (2003) Identified sources and targets of slow inhibition in the neocortex. *Science*, **299**, 1902-1905.

Thomson, A.M. & Destexhe, A. (1999) Dual intracellular recordings and computational models of slow inhibitory postsynaptic potentials in rat neocortical and hippocampal slices. *Neuroscience*, **92**, 1193-1215.

Williams, S.R. & Fletcher, L.N. (2018) A Dendritic Substrate for the Cholinergic Control of Neocortical Output Neurons. *Neuron*.

Williams, S.R. & Stuart, G.J. (2000) Site independence of EPSP time course is mediated by dendritic I_h in neocortical pyramidal neurons. *Journal of Neurophysiology*, **83**, 3177-3182.

Xu, N.L., Harnett, M.T., Williams, S.R., Huber, D., O'Connor, D.H., Svoboda, K. & Magee, J.C. (2012) Nonlinear dendritic integration of sensory and motor input during an active sensing task. *Nature*, **492**, 247-251.

FIGURE LEGENDS

Figure 1. The contribution of active dendritic processes to the F-I relationship is minimized by dendritic GABA$_B$R activation. A) Left, locations of dual dendritic and somatic patch-clamp recordings are indicated on a biocytin-filled layer 5 pyramidal neuron. Dendritic and somatic membrane potential responses to a current step in the somatic patch pipette. **B**) AP rate versus current (F-I) relationship for somatic current injections. Parameters of eq. (*1*) from the fit to the data of the same neuron are indicated. **C**) Extrapolation of the F-I relationship for combinations of dendritic and somatic current injections according to the passive dendrite spike rate (PDSR) model in eq. (*2*) with $D=K_{ds}/R_{in}$. **D**) Membrane potential responses of the same neuron to combined dendritic and somatic current injection. **E**) Experimentally observed F-I relationship. **F**) The average difference of the experimental F-I relationship from the neuron-specific PDSR model (n=23 neurons) shows a strong deviation from the passive case indicating a pronounced contribution of active dendritic processes to the F-I relationship. **G**) Membrane potential responses to combined dendritic and somatic current injection during a puff of baclofen onto the apical dendrite. **H**) Experimentally observed F-I relationship for the same neuron. **I**) During the baclofen puff, the average deviation of the F-I relationships from the PDSR model is small (n=23 neurons). Abbr.: Idend, dendritic current; Isoma, somatic current; Vdend, dendritic membrane potential; Vsoma, somatic membrane potential

Extended Data Figure 1. The contribution of active dendritic processes to the F-I relationship is minimized by dendritic GABA$_B$R activation. A) The F-I relationship of the PDSR model (eq. (*2*) with $D=K_{ds}/R_{in}$) for the neuron shown in Fig.1A. **B**) Fit of eq. (*2*) to the experimental data obtained from the same neuron during combined dendritic and somatic current injection (see Fig. 1E). Note the larger AP rates compared to the PDSR model in A. Fitted parameters are indicated. **C**) Fit to the F-I data recorded from the same neuron during a puff of baclofen onto the apical dendrite (see Fig.1H). **D**) Summed squared errors (SSE) of experimental F-I data to the PDSR model (left) and to the fit of eq. (*2*) (right). Data during baclofen puff (red) showed significantly lower SSE (both: *P*<0.01, Wilcoxon signed rank test, n=23). **E**) Group data (n=25) of parameters of

the PDSR model (light gray) and the fits of control data (gray) and baclofen data (red) to eq. (*2*). Statistically significant differences are indicated (*, P<0.01, Wilcoxon signed rank test).

Figure 2. GABA$_B$R activation reduces nonlinearities of input and transfer resistance. A) Schematic of recording arrangement during dual dendritic and somatic patch-clamp recordings. **B**) Families of current steps were injected into the dendrite (top) or soma (bottom) to evoke subthreshold membrane potential responses before (grey) and during the local application of baclofen (red) onto the apical dendrite. **C**) Local dendritic input resistance (K$_{dd}$) normalized to values at resting membrane potential are plotted versus dendritic current amplitude. Solid line indicates exponential fit (y=*Y0**e(*k**x)) to the data. Baclofen application significantly decreased *k* (*P*=0.0017, n=5 neurons, F(1,110)=10.38), thereby increasing the doubling interval from 1.4 to 3.3 nA. **D**) Dendro-somatic transfer resistance (K$_{ds}$) normalized to values at resting membrane potential are plotted versus dendritic current amplitude. Baclofen application significantly decreased *k* of the exponential fit indicating an increase of the doubling interval from 1.0 to 1.8 nA of K$_{ds}$ (*P*<0.0001, n=11, F(1,236)=35.07). **E**) Somatic input resistance (R$_{in}$) normalized to values at resting membrane potential are plotted versus somatic current amplitude. Baclofen application significantly increased the doubling interval of Rin from 1.1 to 1.9 nA (*P*=0.0008, n=11, F(1,119)=11.82).

Figure 3. The transfer and dendritic input resistances depend on the membrane potential. **A**) Left, schematic of recording arrangement during dual dendritic and somatic patch-clamp recordings. Families of current steps were injected into the soma (center) or the dendrite (right) to evoke subthreshold membrane potential responses. Labels indicate voltage responses recorded from the dendrite (Vd, grey) to somatic current (Is), and somatic responses (Vs, black) to dendritic current (Id). **B**) Scatter plot of normalized somato-dendritic transfer resistance (K$_{sd}$) versus dendro-somatic transfer resistance (K$_{ds}$) show that the transfer resistance is transitive. Transfer resistances were normalized to somatic input resistance (R$_{in}$). Inset shows the overlay of traces from soma (black) and dendrite (grey) for the same family of current injections into the

respective opposite compartment (accentuated in A). Note that they nicely match indicating transitivity. **C**) Relationship of normalized transfer resistance to distance of dendritic recording location from soma. The mono-exponential fit has a length constant of 383 µm. **D**) Scatter plot of K_{ds} normalized to K_{ds} measured at resting membrane potential versus the steady state depolarization at the dendritic recording site. K_{ds} values were estimated from the steady state depolarization in the soma to dendritic current steps of increasing amplitude in 38 neurons. Solid line indicates exponential fit (y=$Y0$*e(k*x)) to the data. **E**) Scatter plot of dendritic input resistance (K_{dd}) normalized to K_{dd} measured at the resting membrane potential versus the local steady state depolarization. **F**) A simple two-compartment model was fitted to the recording data from individual neurons. Inset shows a schematic with a dendritic compartment (green) and a somatic compartment (red) connected by a resistor. The model included I_H (blue channels) and persistent sodium (yellow channel) and captured the nonlinear voltage-dependence of the recording data. For comparison, voltage responses to dendritic current steps in a linear 2-compartmental model without voltage-dependent conductances are shown in the top right corner. **G**) Scatter plot of K_{ds} normalized to somatic Rin (light grey) versus D determined from fits of the F-I data shows a systematic deviation from the identity line ($P<0.0001$, F(1,17)=94.74). If K_{ds} was measured during large current steps just below AP threshold (@depol; green), the slope is not significantly different from the identity line ($P=0.27$, F(1,17)=1.31), indicating a close match. Lines were constrained to cross the origin.

Figure 4. GABA$_B$R-activated K$^+$ channels reduce nonlinearities of input and transfer resistance. A) Two-compartmental models were fitted to the data of all experiments from Fig. 2C (n=5 neurons). **B**) Fitted voltage responses to families of current steps injected into either the dendritic (top) or somatic compartment (bottom) before (grey) and during the local application of baclofen (red) onto the apical dendrite. **C-E**) Increased K$^+$-mediated conductance in the dendritic compartment was sufficient to capture the experimentally observed significant reductions of parameter k in the exponential fit (y=$Y0$*e(k*x)) of the voltage-dependence of K_{dd} ($P=0.0166$, n=5 models, F(1,116)=5.906) and K_{ds} ($P=0.0348$, F(1,116)=4.563) during the baclofen puff. **F**)

Schematic of recording arrangement during dual dendritic and somatic patch-clamp recordings testing the contribution of GIRK channels. **G**) Families of current steps were injected into the dendrite (top) or soma (bottom) to evoke subthreshold membrane potential responses in the presence of tertiapin (0.5 µM, blue) and during the local application of baclofen (red) onto the apical dendrite. **H-J**) Tertiapin bath-applied partially blocked GABA$_B$-mediated effects. Lighter shaded lines indicate the fitted curves in the absence of tertiapin from Fig. 2. The voltage-dependence of K_{dd} and K_{ds} was less affected by baclofen in the presence of tertiapin, as indicated by significantly larger *k*'s of the fitted lines (K_{dd}: *P*=0.033, n1=4, n2=5, $F_{(1,96)}$=4.678; K_{ds}: *P*=0.0354, n1=8, n2=11, $F_{(1,208)}$=4.484). In addition, the doubling interval of K_{dd} was no different after baclofen in the presence of tertiapin compared to tertiapin alone (2.2 nA vs. 1.9 nA; *P*=0.48, n=4 neurons, $F_{(1,71)}$=0.5063).

Extended Data Figure 4. GABA$_B$R-activated K$^+$ channels reduce nonlinearities of input and transfer resistance. A) Local hyperpolarization in the dendrite and soma induced by the dendritic baclofen puff and partial block by tertiapin. Significant effects are indicated ($P<0.05$, n=15, Wilcoxon signed rank test). **B**) The somatic input resistance was significantly increased in the presence of tertiapin (*P*=0.0156, n=7, Wilcoxon signed rank test). **C**) The F-I gain derived from the fit of eq. (*2*) to the data was also increased (*P*=0.0156, n=7).

Figure 5. GIRK channel-activation in dendrites effectively inhibits AP output. **A**) Left, locations of dual dendritic and somatic patch-clamp recordings on a biocytin-filled layer 5 pyramidal neuron. Right, membrane potential responses of the same neuron to combined dendritic and somatic current injection during the presence of tertiapin (0.5 µM). **B**) Experimentally observed AP rate versus current (F-I) relationship for the same neuron. **C**) Average deviation of the experimental F-I relationship from the PDSR model across all neurons (n=7). **D-F**) Membrane potential responses and F-I relationship during a puff of baclofen in the presence of tertiapin. **G**) AP rate reduction during baclofen application in percent of baseline AP rate. Episodes of different dendritic current amplitude were grouped according to somatic current amplitude. There was a

main effect of tertiapin (blue; P=0.0131, ANOVA, F(1,6)=12.14). Asterisks indicate significantly smaller reductions of the AP rate in the presence tertiapin (*P*<0.05, Bonferroni's multiple comparisons test), demonstrating a block of the baclofen effect.

Figure 6. GABA$_B$R-mediated inhibition of L-type Ca^{2+} channels prevents burst firing. **A**) Left, locations of dual dendritic and somatic patch-clamp recordings on a biocytin-filled layer 5 pyramidal neuron. Right, membrane potential responses of the same neuron to combined dendritic and somatic current injection. Dendritic Ca^{2+} spikes (arrowheads) and subsequent sAHPs (arrows) are indicated. **B**) Experimentally observed AP rate versus current (F-I) relationship for the same neuron. **C**) Average deviation of the experimental F-I relationship from the PDSR model across all neurons (n=7). **D**) Membrane potential responses of the same neuron to combined dendritic and somatic current injection after application of nimodipine (10 µM). **E**) Experimentally observed F-I relationship. **F**) Average deviation of the experimental F-I relationship during nimodipine application from the PDSR model across all neurons (n=7). Note the similarity to C. **G-I**) Membrane potential responses and F-I relationship during a puff of baclofen in the presence of nimodipine. Note the decreased deviation from the PDSR model. **J**) Grand mean of the cumulative ISI distribution (n=7 neurons). In control condition (grey), there was a high proportion of fast ISIs (<20 ms), which were lost after baclofen puff onto the apical dendrite (orange) or wash-in of nimodipine (green). Shaded area indicates the SEM. Inset shows the frequency distribution of all ISIs in control condition and after wash-in of nimodipine. Bin width was 5 ms. **K**) AP rate reduction during dendritic puff of baclofen in percent of baseline AP rate. Episodes of different dendritic current amplitude were grouped according to somatic current amplitude. Asterisks indicate significantly smaller reductions of the AP rate in the presence of nimodipine (green; *P*<0.01, Bonferroni's multiple comparisons test), indicating a partial occlusion of the baclofen effect. **L**) Parameters of eq. (*2*) from fits to F-I data in control condition (gray) and in the presence of nimodipine (green) alone or in combination with a baclofen puff (light red and red; n=7). Main effects of baclofen (#, P<0.05, ANOVA) and significant differences between groups (*, P<0.01, Wilcoxon signed rank test) are indicated.

Figure 7. Differential regulation of single APs and HFBs during *in vivo*-like input waveforms by dendritic GABA$_B$Rs. A) Injected current waveforms based on responses to sensory stimulation of the contralateral hind limb recorded *in vivo* from dendrite and soma, respectively (Palmer *et al.*, 2012). Dendritic current is shown in green, somatic in red. **B**) Raster plot of APs emitted in individual episodes during increasing levels of dendritic and somatic stimulation strength. The seven different levels of the injected current in 49 combinations are indicated by the right color bars (S=somatic, D=dendritic). The peak amplitude of the current waveform was increased from 0 pA (white) to 1500 pA (black). Step size was 250 pA. Red circles indicate APs that are part of a high-frequency burst (HFB, ISI<15ms). **C**) Peri-stimulus time histogram of all APs (black) and APs that are part of a HFB (red). **D**-**E**) Raster plot and PSTH of APs emitted in the same neuron while puffing baclofen onto the apical dendrite. **F**) Normalized AP rate during the dendritic baclofen puff in episodes of low AP probability (<7 Hz, and maximal dendritic current amplitude of 750 pA). The blockade of L-type Ca^{2+} channels by nimodipine had no effect on the effect of baclofen (P=0.64, n=7). **G**) The proportion of HFB ISIs (<15 ms) was significantly decreased by nimodipine (P=0.016; Wilcoxon Signed Rank test, n=7). Baclofen (orange) puffed onto the dendrite further decreased the proportion. **H**) Voltage traces recorded in dendrite (green) and soma (red) for all single APs outside of HFBs (ISI>15 ms, left) and APs at the start of a HFB (right). Three traces are highlighted to exemplify the variability of dendritic voltage traces during HFBs. **I**) Voltage traces recorded in dendrite and soma while puffing baclofen onto the apical dendrite.

Extended Data Figure 7. Differential regulation of single APs and HFBs during *in vivo*-like input waveforms by dendritic GABA$_B$Rs. A) Injected current waveforms based on responses to sensory stimulation of the contralateral hind limb recorded *in vivo*. Dendritic current is shown in green, somatic in red. **B**) Peri-stimulus time histograms (PSTH, left) of all APs (black) and APs that are part of a HFB (red) in a neuron during control condition (top) and while puffing baclofen onto the apical dendrite (bottom). Vertical dashed lines in the PSTH and the current waveforms (A) indicate time points of consistently reoccurring HFBs across episodes. On the right, the grand mean (n=7 neurons) of the dendritic (green) and the somatic (red) I$_{eff}$ is shown. Note the

different temporal modulation of somatic I_{eff} and the increased level of dendritic I_{eff} necessary for the induction of HFB compared to single APs during the control condition. Dendritic activation of $GABA_BRs$ by puffed baclofen strongly increased the contribution of somatic I_{eff}, suggesting that AP output was mainly driven by the somatic current injection. **C**) The effect of the L-type Ca^{2+} channel blocker nimodipine on AP output. Left, PSTH from the same neuron before and during dendritic baclofen application in the presence of the L-type Ca^{2+} channel blocker nimodipine. Right, grand means of I_{eff} from the same group of neurons before and during baclofen application. Note the reduced contribution of dendritic I_{eff} and the increased contribution of somatic I_{eff} to HFBs in the presence of nimodipine.

Figure 1. The contribution of active dendritic processes to the F-I relationship is minimized by dendritic GABAB receptor activation.

A PDSR Model
gain = 1.1   D-factor = 0.28
I_{thr} = 0.5 nA

B Fit to data
gain = 1.3   D-factor = 0.59
I_{thr} = 0.59 nA

C Fit to baclofen data
gain = 1.2   D-factor = 0.44
I_{thr} = 0.65 nA

Extended Data Fig. 1. The contribution of active dendritic processes to the F-I relationship is minimized by dendritic GABAB receptor activation.

Figure 2. GABABR activation reduces nonlinearities of input and transfer resistance.

Figure 3. The transfer and dendritic input resistances depend on the membrane potential.

Figure 4. GABABR-activated K+ channels reduce nonlinearities of input and transfer resistance.

Extended Figure 4. GABABR-activated K+ channels reduce nonlinearities of input and transfer resistance.

Figure 5. GIRK channel-activation in dendrites effectively inhibits action potential output.

Figure 6. GABABR-mediated inhibition of L-type Ca2+ channels prevents burst firing.

Figure 7. Differential regulation of single APs and HFBs during in vivo-like input waveforms by dendritic GABABRs.

Extended Data Figure 7. Differential regulation of single APs and HFBs by dendritic GABABRs.