Fitness functions, genetic and non-genetic inheritance, and why

ecological dynamics and evolution are inevitably linked

Tim Coulson[1,2], Tomos Potter[1], Anja Felmy[1]

[1] Department of Zoology, University of Oxford, Mansfield Road, Oxford, OX1 3SZ, UK

[2] tim.coulson@zoo.ox.ac.uk

ABSTRACT

Fitness functions are at the heart of ecology and evolution as they simultaneously determine both

population dynamics and the strength of selection. We link the Breeders and Price equations with

structured population models constructed from fitness functions to show how ecological and

evolutionary dynamics are intimately linked. Analyses of our models reveal i) how multi-

generational predictions of evolution are sensitive to assumptions about phenotypic plasticity and

non-genetic inheritance, ii) that factors that reduce the per time-step population growth rate such

as negative density-dependence and antagonistic species interactions are generally expected to

accelerate rates of evolution compared to cases when they are absent, iii) that evolution is

expected to occur fastest when it is cryptic (all other things being equal) and iv) that eco-

evolutionary dynamics are likely widespread and can be studied using evolutionarily explicit

structured population models.

1. INTRODUCTION

Biologists now appreciate that ecological and evolutionary dynamics can occur on the same time

scale, and this has spawned the field of eco-evolution (Hairston Jr et al. 2005). Research in this

field has flourished with a profusion of reviews (Post and Palkovacs 2009) (Hendry 2016), empirical

analyses of observational (Coulson et al. 2011, Travis et al. 2014) and experimental (Yoshida et al.

2003, Cameron et al. 2013) data, and theory papers (Cotto et al. 2017, Coulson et al. 2017). Yet it

is not always clear how work in eco-evolution links to existing ecological and evolutionary

understanding. In this paper, we show how key theory in ecology and evolutionary biology can be

linked by using a series of simple models constructed around fitness functions that can be

empirically estimated.

Fitness functions describe how phenotypic traits influence fitness or its survival and recruitment

components (Parker and Smith 1990). These functions are central to understanding both

ecological (Easterling et al. 2000) and evolutionary dynamics (Lande 1982) as they determine

population growth, and the strength of selection. Because of the key role that fitness functions

play in both ecology and evolution, they offer a logical starting point from which to develop eco-

evolutionary theory that is consistent with current understanding. Yet not all eco-evolutionary

theory is derived from such a starting point. Equations can be coupled to describe the dynamics of

population size and the mean of a phenotypic trait (Patel et al. 2018), but the coupling can be

phenomenological such that it does not capture known biological constraints.

Evolutionary theory has demonstrated that the mean phenotype and the population growth rate

can change simultaneously as a consequence of the shape of fitness functions (Fisher 1930). In

particular, for annual species, individual fitness is lifetime reproductive success (LRS) and mean

LRS is the population growth rate: selection on a phenotypic trait can lead to phenotypic evolution

as well as an increase in mean LRS and consequently the population growth rate (Fisher 1930). In

spite of this realisation, little evolutionary theory has investigated the simultaneous evolution of

phenotypic traits and population growth for non-annual life histories (but see (Lande 1982, van

Tienderen 2000)) for two reasons. First, weak selection is often assumed such that the dynamics of

populations occur on a shorter time scale to the dynamics of phenotypic trait means (Lande 1982).

Such an assumption simplifies model analysis by separating ecological and evolutionary dynamics,

but in cases where ecological and evolutionary change occurs on the same time scales, models of

weak selection can provide only limited insight. Second, evolutionary models frequently define

selection as a fixed per-generation parameter rather than an emergent consequence of the form

of the survival and reproductive fitness functions (e.g. (Charlesworth 1994)). Under this

assumption it is straightforward to model evolution but it is problematic to define the population

dynamics as a function of parameters describing selection.

Evolutionary models incorporate rules of genetic inheritance; ecological models do not. Instead,

ecological models assume either no covariance between parental and offspring phenotypic trait

values (Easterling et al. 2000), or incorporate a phenomenological association that is identified

from statistical analyses (Coulson 2012). However, many ecological models are constructed from

fitness functions (Ellner et al. 2016). A number of authors have used ecological models built from

fitness functions to simultaneously consider the selective, phenotypic and population dynamical

consequences of altering fitness functions (Coulson et al. 2010, Childs et al. 2016). These models

have demonstrated, for both annual and iteroparous species, that the population growth rate and

the strength of selection on a phenotypic trait vary simultaneously when fitness functions are

altered. However, unlike evolutionary models, these models cannot be used to study rates of

evolution (Chevin 2015). In spite of this, some evolutionary inference can be gained via sensitivity

analyses of ecological models (Caswell 2001) where a parameter in the model is altered in a way

that mimics evolution and the simultaneous impact of the change on population growth and

phenotypic trait means is investigated (Traill et al. 2014).

Ecological and evolutionary models constructed from fitness functions have consequently

revealed that selection, phenotypic traits and population dynamics can change simultaneously

when the form of fitness functions is altered. However, many empiricists verbally articulate a view

of eco-evolution whereby ecological change results in evolution, which in turn results in further

ecological change, and so on *ad infinitum* (Pelletier et al. 2009, Hendry 2016). Such a narrative is

not consistent with ecological and evolutionary understanding, but it is clear why it has arisen. The

majority of theoretical models linking the dynamics of populations and the mean phenotype

assume continuous time (Hairston Jr et al. 2005, Terhorst et al. 2018). One equation describes how

population size within a species changes as a function of the mean phenotypic trait value, while

another captures change in the mean phenotypic trait value as a function of population size.

Because time is continuous and population and phenotypic change is instantaneous, population

size and the mean phenotypic trait change together. However, some authors have interpreted

time as if it were discrete in these models, leading to a verbal narrative that phenotypic change

and population change happen in a stepwise linear sequence rather than together (Hendry 2016).

A better way to visualise simultaneous eco-evolutionary change is to ask how biotic or abiotic

change impacts fitness functions, and how change to these functions impacts ecological and

evolutionary dynamics (Caswell 2001).

Empiricists often work in discrete time because they collect and analyse data from population

censuses collected at regular time intervals (Clutton-Brock 1988). For example, capture-mark-

recapture analyses take a mix of individual-, population- and environment-level attributes and use

them to explain variation in survival rates from one time step to the next, while simultaneously

correcting for imperfect detection probabilities (White and Burnham 1999). Approaches such as

these can be used to estimate discrete time fitness functions, and these functions can be

incorporated into discrete time population dynamic models (Lefkovitch 1965, Easterling et al.

2000, Caswell 2001, Ellner and Rees 2006) to produce point estimates of the simulated processes

at fixed intervals of time.

Discrete time ecological models project over fixed time intervals such as a week, month, or year.

Discrete time evolutionary models typically provide projections on a per generation time scale

(Falconer 1960, Lynch and Walsh 1998). Models such as the Breeders' and Price equations are

consequently usually parameterised using LRS. Fisher (1930) demonstrated that LRS is individual

absolute fitness for species with an annual life history, and some empiricists have extended this

logic to iteroparous species (Morrissey et al. 2010). However, in iteroparous species, where

different individuals can live for different lengths of time and breed on a different number of

occasions and at different ages (Clutton-Brock 1988), mean LRS is not mean fitness or the

population growth rate (Grafen 1988). For iteroparous species it is consequently not

straightforward (and may be impossible) to link population dynamics and evolution using LRS as a

measure of fitness as can be done for annual life histories (Sæther and Engen 2015). It seems

improbable that a general theory of eco-evolution that is applicable to iteroparous species can be

derived from models of LRS. An alternative approach is needed. Discrete-time ecological models

provide a logical starting point (Childs et al. 2016).

Discrete time structured ecological models of iteroparous species based on fitness functions have

recently been extended to incorporate rules of genetic inheritance (Barfield et al. 2011, Coulson et

al. 2011, Childs et al. 2016, Coulson et al. 2017, Rees and Ellner 2019). These models can be shown

to be generalised, dynamic forms of the Breeders' and Price equations for iteroparous species that

can be easily parameterised with data collected from regular population censuses (Coulson et al.

2010, Childs et al. 2016). When viewed from this perspective the models reveal novel insights into

eco-evolution. We demonstrate this by first constructing and analysing simple models before

proceeding to more realistic models that are developed by extending existing models of Soay

sheep (*Ovis aries*) (Coulson 2012).

More specifically we develop eco-evolutionary models of strong selection. We start by considering

the Breeders' (Falconer 1960) and Price (Price 1970) equations of evolution of the mean

phenotype before extending these static models to form evolutionarily explicit dynamic

demographic models based on fitness functions that incorporate phenotypic development and

genetic and non-genetic inheritance. We do this by dividing the modelled phenotypic traits into

their additive genetic and environmental components (Falconer 1960). An additive function

determines how the two components combine to form the phenotypic trait, before fitness

functions operate on the entire trait, altering the bivariate distribution of both components (and

the conditional distributions of each) (Coulson et al. 2017). Additional functions then describe how

both the additive genetic and environmental components of the phenotypic trait are inherited

(the latter through non-genetic inheritance), and how the environmental component develops

throughout an individual's life, potentially as a function of the external biotic or abiotic

environment. Once constructed, models can be iterated and the dynamics of selection, population

size or growth rate, the phenotypic trait, and its additive genetic (i.e. evolution) and

environmental components (i.e. phenotypic plasticity) investigated.

2. METHODS

2.1 Selection

In seasonal breeders, absolute fitness of individual i over a time-step $w_{i,t}$ is always the sum of a

reproductive component $r_{i,t}$ and a survival component $s_{i,t}$: $w_{i,t} = r_{i,t} + s_{i,t}$. The reproductive

component $r_{i,t}$ is the number of offspring produced between time t and $t+1$ that survive to enter

the population at time $t+1$, scaled by the genetic relatedness between parent and offspring, while

the survival component $s_{i,t}$ equals one if the focal individual itself survives to time *t+1*, and zero

otherwise (Watson and Galton 1875). The definition holds for non-seasonal breeders as long as

all individuals born over the time-step can be considered as being within the same cohort.

In populations where immigration and emigration can be ignored, the mean of individual absolute

fitness is the per-time-step population growth rate $\bar{w}_t$. In addition, the mean of the reproductive

components of individual fitness is the mean reproductive rate $\bar{r}_t$, and the mean of the survival

component is the mean survival rate $\bar{s}_t$. Consequently $\bar{w}_t = \bar{r}_t + \bar{s}_t$ (Coulson et al. 2005).

Selection on a phenotypic trait z at time t is measured with a selection differential $\omega_{z,t}$. $\omega_{z,t}$

describes the difference between the mean phenotypic trait value of selected individuals at time t

$\bar{z}_{\omega,t}$ and the mean phenotypic trait value of the entire population prior to selection at time t $\bar{z}_t$

within a generation: $\omega_{z,t} = \bar{z}_{\omega,t} - \bar{z}_t$ (Price 1970). $\omega_{z,t}$ can be written as a quantity that includes

$w_{i,t}$, $\bar{w}_t$ and $z_{i,t}$ – the phenotypic trait value of individual i at time t:

152
$$\omega_{z,t} = \frac{\frac{1}{N_t}\int_{i=1}^{N_t}[w_{i,t} - \bar{w}_t][z_{i,t} - \bar{z}_t]}{\bar{w}_t} \tag{1}$$

153   where N_t is the population size at time t. Equation (1) can be expanded into its reproductive and

154   survival components such that

155
$$\omega_{z,t} = \frac{\bar{r}_t}{\bar{w}_t}\frac{\frac{1}{N_t}\int_{i=1}^{N_t}[r_{i,t} - \bar{r}_t][z_{i,t} - \bar{z}_t]}{\bar{r}_t} + \frac{\bar{s}_t}{\bar{w}_t}\frac{\frac{1}{N_t}\int_{i=1}^{N_t}[s_{i,t} - \bar{s}_t][z_{i,t} - \bar{z}_t]}{\bar{s}_t}. \tag{2}$$

The first term on the right-hand side of equation (2) is a fertility selection differential and the

second is a viability selection differential (Falconer 1960). We now refer to expressions such as

$\frac{1}{N_t}\int_{i=1}^{N_t}[w_{i,t} - \bar{w}_t][z_{i,t} - \bar{z}_t]$ as covariances, and express them as a function $cov(w_t, z_t)$. Note that

functions that take arguments are denoted with round parentheses. We now drop the subscripts i

to simplify notation.

It is clear from equations (*1*) and (*2*) that over a time-step the population growth rate $\bar{w}_t$ and the

strength of selection $\omega_{z,t}$ are related, and that the strength of selection is expected to change with

the population growth rate. However, will the population growth rate $\bar{w}_t$ change in response to

selection and evolutionary change? The answer is (usually) yes, but understanding why requires

formulating the Breeders' equation.

2.2 The Breeders' equation

We start by writing the Breeders' equation in terms of individual absolute fitness (Heywood 2005).

We assume an annual life history and an annual population census such that viability selection can

be ignored and $\bar{w}_t = \bar{r}_t$,

$$\Delta\bar{z}_t = \frac{cov(w_t, A_t + E_t)}{\bar{w}_t} \frac{var(A_t)}{var(A_t + E_t)}. \tag{3}$$

Where $\Delta\bar{z}_t$ is the change in the mean phenotypic trait value over a generation (=time-step for our

annual life history) (Lush 1943). We have now split the phenotypic trait z into its additive genetic

component A_t (the breeding value) and an environmental component E_t such that $z_t = A_t + E_t$.

We assume that z_t does not have any non-additive genetic components, such as dominance or

epistatic components or maternal effects (Falconer 1960). The variance in the breeding values at

time t is denoted $var(A_t)$ – the additive genetic variance.

Equation (*3*) states that a proportion $\frac{var(A_t)}{var(A_t + E_t)} = h^2$ (the narrow-sense heritability) of selection is

realised as evolution. This assumes that change in $\bar{A}_t$ caused by selection within the parental

generation is transmitted to the offspring generation, and that change in $\bar{E}_t$ is not. The equation

consequently assumes that $\Delta \bar{z}_t$ is entirely attributable to change in $\bar{A}_t$. We can rewrite equation

(*3*) as

182
$$\Delta \bar{z}_t = \frac{\beta_1 var(A_t)}{\overline{w}_t},$$ (4)

where β_1 is the slope of a linear regression between individual absolute fitness and the individual

phenotypic trait values (also the derivative $\frac{d\overline{w}_t}{dz_t}$). Note that this makes β_1 different from a selection

gradient (Lande and Arnold 1983), which is computed using relative fitness values (e.g. $\beta_1 /\overline{w}_t$ in

our notation). Mean individual fitness at time t, $\overline{w}_t$, is dependent upon the distribution of the

phenotypic trait and the form of the fitness function. For a linear fitness function $w_t = \beta_0 + \beta_1 z_t$

we can write

189
$$\overline{w}_t = \beta_0 + \beta_1 \bar{z}_t.$$ (5)

190   For non-linear fitness functions mean fitness will not necessarily be expected fitness at the mean

191   value of the phenotypic trait due to non-linear averaging (Ellner et al. 2016). However, as long as

192    $\overline{w}_t$ systematically changes with $\bar{z}_t$ as evolution alters $\bar{z}_t$, we can see that $\overline{w}_t$ (which is also the

193   population growth rate) will also change. For linear functions we can write:

194
$$\Delta \bar{z}_t = \frac{\beta_1 var(A_t)}{\beta_0 + \beta_1 [\bar{A}_t + \bar{E}_t]}.$$ (6)

The equations above assume that selection is the only process that alters $\bar{A}_t$ or $\bar{E}_t$ over a time-

step, and this is one reason why the Breeders' equation does not provide a complete description

of the dynamics of the mean of a phenotypic trait. What processes other than selection might

influence evolution? Non-genetic inheritance, ontogenetic development, and phenotypic plasticity

provide three possibilities. These can be captured, along with selection, with the Price equation.

2.3 The Price equation

The Price equation (Price 1970) provides a complete description of the change in the mean of a

phenotypic trait over a time-step. As before, we start by assuming an annual population census of

a species with an annual life history such that $\bar{w}_t = \bar{r}_{t,}$. The Price equation states that

204
$$\Delta \bar{z}_t = \frac{cov(w_t, z_t)}{\bar{w}_t} + [\bar{z}_{t+1} - \bar{z}_{\omega,t}]$$
(7)

where $\bar{z}_{\omega,t}$ is the mean phenotypic trait of individuals selected to produce offspring (i.e. the

parents), and $\bar{z}_{t+1}$ is the mean phenotypic trait of offspring. The first term on the right-hand side is

a selection differential, while the second term describes how imperfect inheritance (through

mutation, recombination and non-genetic processes) alters the mean phenotypic trait between

times t and $t + 1$.

As we go on to develop dynamic models, it becomes useful to separate out the Price equation into

terms that operate on the breeding value A_t, and on the environmental component E_t of the

phenotype. We start by rewriting equation (*6*) as,

213
$$\Delta \bar{z}_t = \frac{cov(w_t, A_t + E_t)}{\bar{w}_t} + [\bar{A}_{t+1} - \bar{A}_{\omega,t}] + [\bar{E}_{t+1} - \bar{E}_{\omega,t}].$$
(8)

The first term on the right-hand side describes how selection alters components of the mean

phenotype. The second term describes how imperfect genetic inheritance could alter the mean

breeding value between offspring and their parents. The third term describes how non-genetic

inheritance alters the mean environmental component of the phenotypic trait between offspring

and their parents. Note that we can separate $\frac{cov(w_t, A_t + E_t)}{\bar{w}_t} = \frac{cov(w_t, A_t)}{\bar{w}_t} + \frac{cov(w_t, E_t)}{\bar{w}_t}$. This is useful to

demonstrate linkages between the Price and Breeders' equation (below), but can erroneously

suggest that selection at a point in time can operate independently on different components of

the phenotype. It cannot if we assume $z_t = A_t + E_t$. Selection operates on the entire phenotype,

altering the distribution of the components that combine to produce it. For selection to operate in

contrasting ways on each component of the phenotype as sometimes argued (Morrissey et al.

2010, Bonnet et al. 2017) it is necessary to specify a genotype-phenotype maps where $z_t \neq A_t +$

E_t (Appendix 1).

How do the Price and Breeders' equation relate to one another? The Breeders' equation is a

special form of the Price equation (Eqn. 8) that assumes that $[\bar{A}_{t+1} - \bar{A}_{\omega,t}] = 0$ and $\frac{cov(w_t, E_t)}{\bar{w}_t} =$

$-[\bar{E}_{t+1} - \bar{E}_{\omega,t}]$. Biologically this means that genetic inheritance does not change $\bar{A}_t$ (i.e. there is

no transmission bias), and that $\bar{E}_t$ at birth remains identical in each generation. The first of these

assumptions is reasonable for large populations; the second is justified if the environmental

component of the phenotypic trait is determined by completely random developmental noise

(Falconer 1960). If the environmental component of the phenotypic trait is dependent upon non-

random aspects of the external environment, then the assumption will be violated (Kruuk et al.

2002). If this is the case, the Price equation is required. We can write down a form of the Price

equation based on the Breeders' equation by combining equations (4) and (8): $\Delta\bar{z}_t = \frac{\beta_1 var(A_t)}{\bar{w}_t} +$

$[\bar{E}_{t+1} - \bar{E}_{\omega,t}]$.

We now relax the annual life cycle assumption and expand the Price equation into its survival and

reproductive components:

$$\Delta\bar{z}_t = \frac{cov(s_t, A_t + E_t)}{\bar{w}_t} + \qquad (9a) \qquad \text{Effects of viability selection on components of the phenotype.}$$

$$+ \frac{cov(r_t, A_t + E_t)}{\bar{w}_t} + \qquad (9b) \qquad \text{Effect of fertility selection on components of the phenotype.}$$

$$+\frac{\bar{r}_t}{\bar{w}_t}[\bar{A}'_{t+1} - \bar{A}_{\omega_r,t}] + \qquad (9c)$$

Effect of genetic inheritance on the mean breeding value. $\bar{A}'_{t+1}$ is the mean breeding value amongst recruits; $\bar{A}_{\omega_r,t}$ is the mean breeding value among selected parents. Equals zero: inheritance does not change the mean breeding value between selected parents and their offspring.

$$+\frac{\bar{s}_t}{\bar{w}_t}[\bar{A}''_{t+1} - \bar{A}_{\omega_s,t}] + \qquad (9d)$$

Development of the breeding value within surviving individuals. $\bar{A}''_{t+1}$ is the mean breeding value amongst surviving individuals at time *t+1*; $\bar{A}_{\omega_s,t}$ is the mean breeding value amongst individuals that will survive from time *t* to time *t+1*. Equals zero: breeding values do not change within individuals as they age.

$$+\frac{\bar{r}_t}{\bar{w}_t}[\bar{E}'_{t+1} - \bar{E}_{\omega_r,t}] + \qquad (9e)$$

Effect of non-genetic inheritance on the mean environmental component of the phenotypic trait: non-genetic inheritance can change the mean of the environmental component of the phenotype between the selected parents and their offspring.

$$+\frac{\bar{s}_t}{\bar{w}_t}[\bar{E}''_{t+1} - \bar{E}_{\omega_s,t}]. \qquad (9f)$$

Effect of phenotypic plasticity on development of the mean environmental component of the phenotypic trait: the environmental component of the phenotype can change within individuals as they age.

In this formulation we assume that all processes remain constant with age. This assumption could

be relaxed using the approach in Coulson and Tuljapurkar (2008).

From equation (*9*) it is clear that change in the mean of a phenotypic trait over a time-step is

dependent on the strength of viability and fertility selection on the phenotypic trait, and on how

ontogenetic development and imperfect inheritance might alter its mean breeding value and

environmental component. Components 9(c) and 9(d) are usually assumed to be zero, as

inheritance is not assumed to change the mean breeding value between selected parents and

their offspring (9(c)), and as breeding values are not assumed to change within individuals as they

age (9(d)).

2.4 Making the Breeders' and Price equations dynamic

Each of the terms in the static Breeders' and Price equations are emergent properties of the

distribution of the phenotypic trait (and its components) at time t, fitness functions, and functions

that determine the terms in equations 9(c) to 9(f) (Coulson et al. 2010, Childs et al. 2016). In order

to make the Price and Breeders' equations dynamic, it is consequently necessary to specify fitness

functions that determine selection, as well as functions that determine development and

inheritance of the environmental component of the phenotypic trait, and inheritance of the

additive genetic component (Coulson et al. 2017). These functions then iterate forward the

dynamics of the distribution of $z_t = A_t + E_t$. We start by considering the dynamics of the

univariate distribution of z_t for an annual species before extending the model to track the

bivariate distribution of A_t and E_t for iteroparous life histories.

A reproductive fitness function, $R(z_t)$, can be used to map the distribution of the individual

phenotypic traits z_t at time t, $n(z_t)$, to a new distribution of the phenotypic trait at time $t + 1$,

$n(z'_{t+1})$ (Ellner et al. 2016),

$$n(z'_{t+1}) = \int H(z'_{t+1}|z_t)R(z_t)n(z_t)dz. \tag{9}$$

$H(z'_{t+1}|z_t)$ is a transition function that describes the probability of transitioning from trait value z

at time t to all possible trait values z'_{t+1} at time $t + 1$ (Easterling et al. 2000). Trait value z_t is

expressed in the focal individual, while trait values z'_{t+1} are expressed in offspring in annual

species where projections are made over a time step. The transition function is needed when

predicting evolutionary change over multiple time-steps if offspring that recruit to the population

at time $t + 1$ and their parents at time t do not perfectly resemble one another.

To produce a model that projects terms in the Breeders and Price equation over a time step, z_t

can be replaced with $A_t + E_t$ (Coulson et al. 2017). First, write,

271
$$n(A'_{t+1}, E'_{t+1}) = \int H(A'_{t+1}, E'_{t+1} | A_t, E_t) R(A_t + E_t) n(A_t, E_t) dA dE. \qquad (10)$$

The transition function $H(A'_{t+1}, E'_{t+1} | A_t, E_t)$ can be formulated to capture the second term on

the right-hand side of the Breeders' equation (3) or the last two terms on the right-hand side of

the Price equation (8), and so can describe genetic or non-genetic inheritance rules. A dynamic

form of the Breeders' equation (3) requires $H(A'_{t+1}, E'_{t+1} | A_t, E_t)$ to be specified such that it

produces a distribution of $n(A'_{t+1}, E'_{t+1})$ that has means of $\bar{A}'_{t+1} = \bar{A}_{\omega,t}$ and $\bar{E}_t = \bar{E}_{t+1}$ and a

variance-covariance matrix $\begin{pmatrix} var(A_t) & 0 \\ 0 & var(E_t) \end{pmatrix}$ where $var(A_t)$ can be selected to be a constant

or to erode with time as a function of selection, and $var(E_t)$ is constant (Childs et al. 2016,

Coulson et al. 2017, Rees and Ellner 2019). There are a number of ways this can be achieved

(Appendix 2). Any formulation of $H(A'_{t+1}, E'_{t+1} | A_t, E_t)$ that results in $\bar{A}'_{t+1} \neq \bar{A}_{\omega,t}$ and/or $\bar{E}_t \neq$

$\bar{E}_{t+1}$ provides a dynamic form of the Price equation (8). Once the transition and fitness functions

are parameterised, derivations from Coulson *et al.* (2010) can then be used to create time series of

$\Delta \bar{z}_t$, β_0, β_1, $var(A_t)$, $\bar{A}_t$ and $\bar{E}_t$ for different parameterisations of $H(A'_{t+1}, E'_{t+1} | A_t, E_t)$ and

$R(A_t + E_t)$ and different starting values of $n(A_t, E_t)$.

Equations (*9*) and (*10*) can be extended to incorporate per-time step survival and recruitment

functions to allow models to be constructed for iteroparous species. Models now include two

transition functions: an ontogenetic development function $G(z'_{t+1} | z_t)$ operating on surviving

individuals, and an inheritance function $D(z'_{t+1} | z_t)$ that captures the similarity between parental

and offspring traits (Easterling et al. 2000, Ellner and Rees 2006, Coulson 2012). The development

function operates with a survival fitness function $S(z_t)$, while the inheritance function operates

with the recruitment fitness function $R(z_t)$:

292
$$n(z'_{t+1}) = \int [D(z'_{t+1} | z_t) R(z_t) + G(z'_{t+1} | z_t) S(z_t)] n(z_t) dz. \qquad (11)$$

Equation (*11*) can be extended to project forwards the dynamics of the phenotypic trait

partitioned into A_t and E_t. Selection operates on the phenotypic trait $z_t = A_t + E_t$, potentially

altering the size, location and shape of the conditional distributions of both A_t and E_t, before

inheritance and development functions then operate on the post-selection distribution of A_t and

E_t:

$$n(A'_{t+1}, E'_{t+1}) = \int [D(A'_t, E'_t | A_t, E_t) R(A_t + E_t) + G(E'_t | A_t, E_t) S(A_t + E_t)] n(A_t, E_t) dA_t dE_t. \quad (12)$$

How does this equation relate to equation (9)? Term 9(a) is determined by $S(A_t + E_t) n(A_t, E_t)$,

term 9(b) by $R(A_t + E_t) n(A_t, E_t)$, terms 9(c) and 9(e) by $\int D(E'_t | A_t, E_t) R(A_t + $

$E_t) n(A_t, E_t) dA_t dE_t$ and terms 9(d) and 9(f) by $\int G(E'_t | A_t, E_t) S(A_t + E_t) n(A_t, E_t) dA_t dE_t$ (Childs

et al. 2016).

2.4 A simple evolutionarily explicit IPM

We start with a model of the form of equation (*10*) with the phenotypic trait separated into its

additive genetic and environmental components $z_t = A_t + E_t$, but where fitness is not separated

into its recruitment and survival components. We assume a linear fitness function and iterate

$n(A_t, E_t)$ given the following parameterisation,

$$w(A_t + E_t) = -0.2 + 0.05(A_t + E_t) \quad (13)$$

and a starting bivariate normal distribution $n(A_{t=1}, E_{t=1})$ at as having means $\bar{A}_t = \bar{E}_t = 10$ and a

variance-covariance matrix $\sigma^2 = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$. The choice of parameter values is for illustration rather

than representing any particular system. The mean of the breeding value distribution does not

change between parents and their offspring. We run four model scenarios. In scenarios 1) and 3)

selection erodes additive genetic variation, while in scenarios 2) and 4) additive genetic variation is

not eroded by selection. In scenarios 1) and 2) the mean and variance of the environmental

component of the phenotype remains constant across generations, while in scenarios 3) and 4)

the mean of the environmental component declines with time such that $\bar{E}_t = 10 - 0.05t$.

Scenarios 1) and 2) are dynamic forms of the Breeders' equation, while scenarios 3) and 4) are

dynamic forms of the Price equation. The last two scenarios capture a process where some aspect

of the environment can determine the dynamics of the environmental component of the

phenotype (Kruuk et al. 2002).

Each simulation is run for 50 generations. For each simulation we report the rate of evolution

(defined as $\Delta \bar{A}_t$), the rate of change in the mean phenotype $\Delta \bar{z}_t$, the population growth rate

(mean fitness $\bar{w}_t$), trends in the additive genetic variance, covariances between $\Delta \bar{A}_t$ and $\Delta \bar{z}_t$ with

$\bar{w}_t$ and all terms in equations (1)-(9).

2.5 A more realistic model using data from Soay sheep

The assumption of linear fitness (equation (13)) is unrealistic because litter size and survival rates

are bounded. In addition, in variable environments, survival or transition functions for the

environmental component of the phenotype depend on environmental factors such as the

population size or phenotypic or age structure of the focal species or of an interacting one (Kruuk

et al. 2002, Bassar et al. 2017). Moreover, although breeding values remain constant with age, the

environmental component of the phenotypic trait can change within individuals (Falconer 1960).

We consequently develop a density-dependent model, where fitness is separated into its survival

and recruitment components, and where the environmental component of the phenotype can

develop within surviving individuals, and can vary between parents and their offspring.

We took a published density-dependent IPM of the dynamics of body size from the female

component of an individually marked population of Soay sheep (*Ovis aries*) living on the island of

Hirta in the St. Kilda archipelago in Scotland (Coulson 2012). The published IPM was of the form of

equation (*11*), and was modified to be of the form of equation (*12*). Functions and parameter

values are provided in Appendix 3.

The parameterisation in Appendix 3 describes the baseline model. We ran four simulations where

we set the density-dependence parameter to 0 in either the survival fitness function, the

recruitment fitness function, the development or inheritance functions for E_t. Although this is a

large perturbation, it reveals some interesting dynamics. We then compared the dynamics of the

breeding value, the mean phenotypic trait, and mean fitness across the simulations.

None of the simulations reached an equilibrium because of the positive slopes for the phenotypic

trait in the fitness functions and some additive genetic variance is always maintained.

3. RESULTS

3.1 Theoretical insight

From equation (*6*) it is clear that the rate of evolution of a trait over a time step will i) scale linearly

with the additive genetic variance for the trait (Falconer 1960, Lynch and Walsh 1998), ii) will

reduce as the elevation (intercept) of the fitness function (β_0) is increased, iii) will slow as the

mean phenotype is increased (assuming $\bar{A}_t$ and $\bar{E}_t$ are both positive), and iv) has a complex

association with the linear slope of the phenotype with fitness (β_1) as this term appears in both

the numerator and denominator.

The transition terms in the Price equation provide a way to capture processes other than selection

that influence either $\bar{E}_t$ or $\bar{A}_t$ (Price 1970, Coulson and Tuljapurkar 2008). For example, if $\Delta\bar{A}_t -$

$\Delta\bar{E}_t = 0$ then $\Delta\bar{Z}_t = 0$ and phenotypic change due to evolution is countered by change in the

environmental component of the phenotype, a phenomenon termed cryptic evolution (Merilä et

al. 2001a).

3.2 Results from the simple model

Simulating the simple model provides graphical representation to the theoretical insights.

Evolution, defined as a change in the mean breeding value, was fastest when additive genetic

variance was not eroded by selection, and when there was a negative temporal trend in $\bar{E}_t$ (Figure

1(A)). The negative temporal trend in $\bar{E}_t$ accelerated the rate of evolution via its effect on reducing

mean fitness. In spite of accelerating the rate of evolution, the trending values of $\bar{E}_t$ with time

slowed the rate of phenotypic change as positive change in $\bar{A}_t$ was countered by negative change

in $\bar{E}_t$ (Figure 1(B)). Mean fitness evolved fastest when additive genetic variance was not eroded,

and $\bar{E}_t$ did not decline with time (Figure 1(C)). Increased levels of the additive genetic variance in

the phenotypic trait increased the rate of change in $\bar{w}_t$. Linear selection reduced the additive

genetic variance if it was not in each generation (Figure 1(D)). When there was no temporal trend

in $\bar{E}_t$, the association between the mean breeding value and mean fitness was linear (Figure 1(E)),

while it was slightly curvilinear when $\bar{E}_t$ trends with time. As the fitness function was linear, there

was by definition a linear association between the mean phenotypic trait and mean fitness, with a

slope of β_1 (Figure 1(F)).

We can graphically illustrate the evolutionary, phenotypic trait, and fitness dynamics of this simple

model with a plot of A_t against E_t that includes fitness clines (Figure 2(a)). Each diagonal dashed

line represents equivalent values of the phenotypic trait made from different combinations of A_t

and E_t. Because the fitness function is linear and includes no terms additional terms beyond β_0

and β_1, these dashed lines also represent fitness. The darker the colour of the line, the larger the

value of z_t and fitness. The central solid line contours represent the bivariate distribution of A_t

and E_t at time $t = 0$ when simulations are begun. The coloured arrows represent phenotypic

change. When evolution is cryptic and $\Delta \bar{A}_t - \Delta \bar{E}_t = 0$, the A_t increases at the same rate as E_t

decreases and neither z_t, nor $\bar{w}_t$ change. Because $\bar{w}_t$ does not change (blue arrow), evolution

proceeds rapidly as the denominator in the selection differential remains constant with time.

Dynamics forms of the Price equation are required to capture such cryptic evolution. The green

line represents dynamics predicted by dynamic forms of the Breeders' equation with $\bar{A}_t$ and $\bar{w}_t$,

evolving, but the $\bar{E}_t$ remaining constant with time. Evolution occurs more slowly than when it is

cryptic because $\bar{w}_t$ increases with time, increasing the denominator of the selection differential.

The red line captures evolutionary dynamics where change in the mean breeding value and the

mean environmental component of the phenotype act in the same direction, increasing both $\bar{z}_t$

and $\bar{w}_t$. A dynamic form of the Price equation is required to capture such dynamics, and evolution

proceeds even more slowly than in the previous two examples, because $\bar{w}_t$ increases even faster.

Finally, the purple arrow represents no evolution, but phenotypic change due solely to change in

$\bar{E}_t$ – i.e. via non-genetic inheritance (or phenotypic plasticity in iteroparous species). A dynamic

form of the Price equation is required to capture such dynamics.

3.2 Applied results from a more realistic model using data from Soay sheep

The more complex Soay sheep model extends insights from the simple model in three ways. First,

we consider an iteroparous life history (equation (*12*)), second, the incorporation of density-

dependence in the survival and recruitment functions has the effect of reducing the intercepts of

the fitness functions with time as $\bar{w}_t$ evolves and population size increases, and third, the non-

linear fitness functions mean that the derivative of mean fitness to the phenotype (i.e. the slope

β_1 in equation (4)) can vary as evolution proceeds. Figure (2(b)) shows the joint dynamics of $\bar{A}_t$

and $\bar{E}_t$ for each simulation. We do not include fitness clines such as those in Figure (2(a)) because

the incorporation of density-dependence complicates the association between $\bar{Z}_t$ and $\bar{w}_t$, with the

different models exhibiting different associations (Figure 2(c)). The blue points demonstrate that

as evolution proceeds $\bar{Z}_t\bar{Z}_t$ can change, but the mean of a fitness component can remain constant

if the fitness function changes with time. However, if $\bar{w}_t$ does evolve at some rate with z_t, the

derivative β_1 (red lines) can change with time if fitness functions are non-linear. Both of the

processes operate in the modified Soay sheep model.

Removing density-dependence from the recruitment and inheritance functions in the modified

sheep model increased the rate of evolution of body size, while removing it from the survival and

development function decreased it (Figure 2(b)). By far the largest effect was caused by perturbing

the survival function.

Temporal changes in $\bar{E}_t$ were negative in all models, with the greatest rate of change observed

when density-dependence was removed from the recruitment function (Figure 2(b)). In all models,

$\bar{E}_t$ trended to smaller values. This negative trend partially countered the positive trends in $\bar{A}_t$,

such that $\bar{z}_t$ increased with time in all models (Figure 3(b)), but to a lesser extent than change in

$\bar{A}_t$.

$\bar{w}_t$ evolved at similar rates in the baseline model and models where density-dependence was

removed from the recruitment, inheritance and development functions. In contrast, $\bar{w}_t$ evolved

much more slowly when density-dependence was removed from the survival function (Figure

3(c)), although initial population size was much larger in the density-independent survival model

than in any of the other models (Figure 2(b)). The additive genetic variance increased slightly with

time when density-dependence was removed from the survival and recruitment functions, but

remained constant when development and inheritance were density-independent (Figure 3(d)).

Removing density-dependence from each fitness functions impacted evolutionary, phenotypic and

population size change via a number of different routes. First, removing density-dependence

resulted in different starting population sizes and structures (Figure 4(a)-(e)). In the baseline

model and the model with density-dependence removed from the inheritance function, there was

clear bimodality in the distributions z_t throughout the simulation (Figures 3(a,e)). The bimodality

reflects a population with similar proportions of smaller juveniles and larger adults. There is

weaker evidence of a bimodal distribution at time $t = 1$ for the other three models, with

bimodality weakening as the simulations proceeded (Figures (3(b-d)). In the models with density-

independent survival and inheritance, small juveniles are comparatively rare, and their frequency

decreases as the simulations proceed. In contrast, in the density-independent recruitment model,

the population is dominated by small juveniles and becomes increasingly so as the simulation

proceeds.

The rates of evolution of the phenotypic trait and mean fitness are determined by i) β_1, ii) $\overline{w}_t$

(which depends upon $n(z_t)$ and the fitness functions), and iii) $var(A_t)$. Because the models are

density-dependent, the mean population growth rate tends towards unity, and differs little

between the simulations (Figure 3(c)). Relatively little of the dynamical difference between the

models is consequently due to differences in $\overline{w}_t$ across simulations. Similarly, because the additive

genetic variance remains approximately constant in all simulations (Figure 3(d)) this too is not a

major driver of the dynamics we report from the sheep model.

Most of the dynamical differences between the simulations are consequently caused by processes

generating variation in β_1: the covariance between individual phenotypic traits and individual

absolute fitness (equation 1) and the phenotypic variance. These two terms tend to show similar

temporal dynamics (Figure 4(f, g, h)) resulting in the derivative between the mean survival and the

phenotype (the strength and direction of viability selection) remaining approximately constant in

the baseline model and the density-independent growth and inheritance function models (Figure

3(i)). In contrast, the derivative trended downwards as the simulation proceeded in the density-

independent survival and recruitment models (Figure 3(i)). Similar patterns were observed for the

derivative of mean recruitment to the mean phenotype (not shown).

These temporal changes in the derivatives of the mean fitness components to the phenotypic trait

value resulted in approximately constant survival and recruitment rates in the density-

independent survival model, weakly trending decreases in survival rates and weakly trending

increases in recruitment rates in the baseline, density-independent development and density-

independent inheritance models, and stronger temporal trends for decreased survival and

increased recruitment rates in the density-independent recruitment model (Figure 4(j)). In other

words, the patterns we report in mean fitness are caused by opposing trends in fitness

components in these simulations.

The dynamics of β_1 that generated these contrasting patterns across the different simulations are

determined by the shape of the fitness function (Figure 2(c)). As density increased, the intercept

of the logistic fitness functions became increasingly more negative due to the density-dependence

terms in the models. In contrast, as mean body size increased, the body size distribution moved to

the right on the x-axis of figure 2(c). To some extent, these two processes tended to cancel one

another out, such that mean survival and mean recruitment tended to change relatively slowly.

The one exception to this pattern is in the density-independent recruitment model where

increasing body size increased recruitment rates but did not lower the intercept of the function.

Instead, mean survival rates decreased as density increased as mean fitness evolved.

The Soay sheep population is primarily regulated via density-dependent survival. When density is

removed from the survival function, the population consequently increases to a large size, in turn

reducing recruitment rates, development rates, and the mean value of the environmental

component of the phenotype. Large individuals, however, have very high survival, at a point when

the derivative of mean survival to body weight is very shallow – i.e. β_1 is close to zero and

evolution of both the mean phenotypic trait and mean fitness is slow.

4. DISCUSSION

The dynamic forms of the Breeders' and Price equations we develop from demographic models

build on existing approaches by Lande (Lande 1982), Barfield et al. (Barfield et al. 2011), Childs et

al.(Childs et al. 2016), Coulson et al. (Coulson et al. 2011, Coulson et al. 2017) and Rees and Ellner

(Rees and Ellner 2019). The approach is powerful as it allows the simultaneous study of population

dynamics with the evolution of phenotypic traits, life histories, and mean fitness (Coulson et al.

2011). In this paper we use simple dynamic models to demonstrate how our approach generate

dynamics of terms in the Breeders' and Price equations.

The first insight our work provides concerns eco-evolution. A widely championed view of eco-

evolution is that ecological change generates evolutionary change and so on *ad infinitum* (Hendry

2016). The logic behind this interpretation is that ecology and evolution are two separate

processes that can influence one another (Pelletier et al. 2009). Our approach reveals that

ecological change and evolution occur simultaneously when an environmental process alters

either the fitness, development or inheritance functions in demographic models (see also (Coulson

et al. 2011, Childs et al. 2016, Coulson et al. 2017)). For example, the reduction in size of a

predator population may increase the elevation of fitness functions, simultaneously increasing the

population growth rate and reducing the strength of selection and the rate of evolution.

Our view of eco-evolution is based on the logic that a population can be represented by a

multivariate distribution of genotypes and phenotypes (Easterling et al. 2000, Ellner and Rees

2006). The dynamics of this distribution are then determined by fitness, development and

inheritance functions. The dynamics of the distribution's volume (size) are the population

dynamics, while the dynamics of the mean genotype (and sometimes phenotypic trait) capture

evolution (Smallegange and Coulson 2013). These models can also be used to study the dynamics

of life history descriptors such as generation length and LRS (Ellner and Rees 2006, Steiner et al.

2012, Steiner et al. 2014), and examine life history evolution (Childs et al. 2003, 2004). The fitness,

development, and inheritance functions that underpin these models can vary as a function of age,

sex, location, and genotype, and they can include aspects of the biotic and abiotic environment

such as the size and structure of the focal species or of other species in the community, or climatic

variation (Ellner and Rees 2006, Coulson 2012, Schindler et al. 2015, Ellner et al. 2016).

There are two criticisms that have been levied at the demographic modelling approach that we

use. First, that models tend to be data driven rather than mechanistic (Chevin 2015, Janeiro et al.

2017), and second, that the data required to parameterise models for natural systems can be

prohibitive. It is correct that most models of real systems have been phenomenological rather

than mechanistic. However, following observations that particular mechanisms were not included

in early models, the structured modelling approach has been extended to incorporate a number of

mechanisms. Genetic inheritance (Coulson et al. 2011, Childs et al. 2016), dynamic energy budgets

(Smallegange et al. 2017), and resource dependence (Bassar et al. 2017) have now been explicitly

included into models, with ongoing work including additional mechanisms.

The more complex a model is, the harder it will be to parameterise, and analyse. However, it is

possible to collect the required data from many systems, and complex models have been

parameterised, and analysed, for many natural systems (Merow et al. 2014, Rees et al. 2014,

Ellner et al. 2016). We are not advocating that simpler models (or even individual-based models

that are often even more parameter-rich than the ones we develop) be abandoned. Instead, we

are demonstrating how structured models can be remarkably powerful at linking ecology and

evolution. The choice of model to construct always depends upon the question, the data available,

and the desired degree of analytical, or numerical, tractability. All models are wrong, and

criticising any modelling approach for what it does not include is easy; the challenge is selecting

the approach that is least wrong for the question at hand.

The question we are interested in is how might perturbations impact the fitness, development,

and inheritance functions that constitute demographic models, and how do such impacts

simultaneously influence population dynamics, phenotypic trait dynamics, and evolution? To study

evolution, we examine how environmental variation can impact terms in the Breeders' and Price

equations.

Our results reveal that there are multiple ways that the environment can impact evolution. The

first route is via its impact on mean fitness that can influence the strength of selection. We

explicitly separate the covariance between a phenotypic trait and relative individual fitness – a

quantity central to evolution – into the covariance between the phenotypic trait and absolute

individual fitness, divided by mean absolute individual fitness (Heywood 2005). This is a

mathematically trivial alteration to standard derivations, but is biologically illuminating, as over a

time step, mean individual absolute fitness is the population growth rate (Watson and Galton

1875, Engen et al. 2009). Population ecologists have long studied the population growth rate

(Caswell 2001), and by separating it out of the selection differential, the understanding they have

garnered can be used to gain insight into the dynamics of selection. In particular, population

ecologists have revealed how intraspecific competition for resources, interspecific interactions,

and the abiotic environment can impact the distribution of per-time step population growth rates

(Turchin 2003): these drivers have the potential to impact the temporal dynamics of selection via

the temporal dynamics of mean fitness.

Writing relative fitness in terms of absolute fitness also allows us to reformulate the Breeders' and

Price equations so that the numerator of the equation now contains the derivative of mean fitness

to the phenotypic trait value. Much ecological research has revealed how phenotypic traits such

as body size are associated with demographic rates (components of absolute fitness), and how

these associations can vary with aspects of the biotic and abiotic environment (Merow et al. 2014,

Ellner et al. 2016). In general, recruitment and survival rates tend to be higher in good

environments. However, because these fitness components are bounded, as their means increase,

the derivative of fitness to phenotypic trait values is (often) expected to decrease as favourable

environments act to increase the intercept of fitness functions. Because favourable environments

also result in higher per-time step population growth rates, we may expect selection to be weaker

in favourable rather than in harsh environments. Such a pattern has been observed in some cases

(Wilson et al. 2006) but is not universal (Siepielski et al. 2017). Examining how the derivative of

components of fitness to phenotypic traits changes with aspects of the environment across a

range of species will help to understand why.

The third route via which the environment can impact selection is via change in the mean of the

environmental component of the phenotype (Nussey et al. 2007). Non-genetic change in the mean

phenotype via phenotypic plasticity or non-genetic inheritance can impact the rate of phenotypic

change (Reed et al. 2011, Coulson et al. 2017), which in turn can influence both mean fitness

(Figure 2(a)), or the derivative of mean fitness to the phenotype (Figure 2(c)).

We only consider additive genotype-phenotype maps such that $z_t = A_t + E_t$ where A_t and E_t are

on the same scale of measurement (Fisher 1930). When genotype-by-environment interactions

occur, the environment can influence evolution (and consequently ecological dynamics) via an

alternative route (Via and Lande 1985). Some authors have argued that selection acts on the

environmental and additive genetic components of the phenotype in contrasting ways (Morrissey

et al. 2010, Bonnet et al. 2017), thus invoking a genotype-by-environment interaction. Indeed, it

seems plausible that genotype-by-environment interactions are widespread in natural settings.

Our approach can be extended to include non-additive genotype-phenotype maps (Coulson et al.

2017).

Given the multiple routes via which environmental variation can impact ecology and evolution it is

perhaps unsurprising how difficult it has proven to predict evolution in variable environments

(Merilä et al. 2001b), yet understandable why models such as the Breeders' equation do tend to

make more accurate predictions in the more constant environments under which most artificial

selection regimes are imposed (Lynch and Walsh 1998). Our approach does provide a way to

explore how different forms of environmental variation operating via different components of the

life history might simultaneously impact ecological dynamics and evolution.

The modelling framework we work within allows the effects of phenotypic plasticity and non-

genetic inheritance on evolution, phenotypic change, and ecology to be explored. Results are

consistent with current quantitative genetic understanding, but our approach significantly

advances our ability to understand how environmental variation, and processes other than

selection such as phenotypic plasticity and non-genetic inheritance, can impact evolutionary

outcomes over multiple time-steps. Because life-history descriptors like generation length and life

expectancy can be calculated from integral projection models (Steiner et al. 2012, Steiner et al.

2014), it is also possible to explore how phenotypic evolution impacts life history. The next steps

involve constructing models for interacting species, incorporating multiple traits, and

parameterising models for a range of systems.

Figure 1. Evolution of (A) the mean breeding value, (B) the mean phenotypic trait, (C) mean

fitness, and (D) the additive genetic variance in simple models over 50 generations. The red lines

represent sim 2), where additive genetic variance is not eroded by selection, and the mean and

variance in the environmental component of the phenotypic trait remain constant with time. The

black line represents sim 1), where the additive genetic variance is eroded and the mean and

variance in the environmental component of the phenotypic trait remain constant with time. The

green (sim 4)) and blue line (sim 3)) are the same as sim 2) and sim 1), except we impose a weak

temporal trend in the mean of the environmental component of the phenotypic trait. (E) In

models with no temporal trend in the environmental component of the phenotypic trait, the

relationship between the mean breeding value and mean fitness is linear, while it is slightly

curvilinear in models with temporal trends in E. In contrast, there is (F) a linear association

between the mean phenotypic trait and mean fitness in all four models.

Figure 2. The dynamics of phenotypic traits and mean fitness. (A) Because the breeding value and

the value of the environmental component of a phenotypic trait sum to produce the phenotypic

trait, there are multiple ways the same phenotypic trait value can be produced (dashed grey lines).

If the phenotypic trait linearly influences fitness, these phenotypic trait clines also represent

fitness clines, generating linear fitness contours. Larger values of the trait, and of mean fitness, are

depicted with darker dashed lines. Depending upon the dynamics of the environmental

component of the phenotype, selection can fail to climb across contours, or can climb at a range

of rates even for the same, constant, derivative of mean fitness to the mean phenotype (i.e.

phenotypic slope in the linear fitness function). Because mean fitness remains lowest when

evolution is cryptic, all other things being equal, selection will also be strongest, and evolution

fastest, when non-genetic inheritance or phenotypic plasticity, impacts the environmental

component of the phenotype in a manner that exactly counters gains to the additive genetic

component of the phenotype attributable to selection. (B) The bivariate dynamics of the mean

breeding value and the mean environmental component of the phenotypic trait over time in each

simulation for the modified sheep model. The starting point of each simulation is towards the

bottom of the figure. The increasing size of the thick lines represent how population size (as a

consequence of the dynamics of mean fitness) changes as the simulation progresses. Bivariate

distributions of the components of the phenotypic trait are drawn as contour plots at time $t=1$,

$t=50$ and $t=100$ for the simulation where density was removed from the recruitment (fitness)

function. Evolution is captured by the length of the line in the y-axis, while change in the

environmental component of the phenotypic trait is captured by change in the x-axis. (C) The

derivative of mean fitness to the mean phenotype (red line) can vary as either the mean

phenotypic trait increases when linear functions are non-linear, or when environment drivers such

as an increase in density act to lower the intercept of the fitness function (black lines).

Figure 3. Evolution of (A) the mean breeding value, (B) the mean phenotypic trait, (C) mean

fitness, and (D) the additive genetic variance in the modified sheep model. The baseline model

(black lines) has negative density-dependence in each function, while the coloured lines

represents simulations where density-dependence has been removed from one particular

function.

Figure 4. Dynamics of the phenotypic trait distribution in the baseline model (A) and in models

with negative density-dependence removed from each function (B-E). Distributions are drawn at

times $t=1$, $t=50$ and $t=100$. These dynamics are determined in part by the selection (the derivative

of mean fitness to the mean phenotypic trait value) that consists of (F, G) covariances between the

trait and components of fitness, (H) the phenotypic variance in the trait. (i) derivative of mean

survival to the mean phenotype (similar patterns for recruitment not shown) and (J) the temporal

dynamics of mean survival and recruitment rates.

Figure 1

Figure 2.

Figure 3.

Figure 4.

Barfield, M., R. D. Holt, and R. Gomulkiewicz. 2011. Evolution in stage-structured populations. The
American Naturalist **177**:397-409.
Bassar, R. D., J. Travis, and T. Coulson. 2017. Predicting coexistence in species with continuous
ontogenetic niche shifts and competitive asymmetry. Ecology **98**:2823-2836.
Bonnet, T., P. Wandeler, G. Camenisch, and E. Postma. 2017. Bigger is fitter? Quantitative genetic
decomposition of selection reveals an adaptive evolutionary decline of body mass in a wild
rodent population. PLoS biology **15**:e1002592.
Cameron, T. C., D. O'Sullivan, A. Reynolds, S. B. Piertney, and T. G. Benton. 2013. Eco-evolutionary
dynamics in response to selection on life-history. Ecology Letters **16**:754-763.
Caswell, H. 2001. Matrix population models: Construction, Analysis and Interpretation. Sinauer
Associates.
Charlesworth, B. 1994. Evolution in age-structured populations. Cambridge University Press
Cambridge.
Chevin, L. M. 2015. Evolution of adult size depends on genetic variance in growth trajectories: a
comment on analyses of evolutionary dynamics using integral projection models. Methods
in Ecology and Evolution **6**:981-986.
Childs, D. Z., M. Rees, K. E. Rose, P. J. Grubb, and S. P. Ellner. 2003. Evolution of complex flowering
strategies: an age–and size–structured integral projection model. Proceedings of the Royal
Society of London. Series B: Biological Sciences **270**:1829-1838.
Childs, D. Z., M. Rees, K. E. Rose, P. J. Grubb, and S. P. Ellner. 2004. Evolution of size–dependent
flowering in a variable environment: construction and analysis of a stochastic integral
projection model. Proceedings of the Royal Society of London. Series B: Biological Sciences
271:425-434.
Childs, D. Z., B. C. Sheldon, and M. Rees. 2016. The evolution of labile traits in sex-and age-
structured populations. Journal of Animal Ecology **85**:329-342.
Clutton-Brock, T. H. 1988. Reproductive success: studies of individual variation in contrasting
breeding systems. University of Chicago Press.
Cotto, O., J. Wessely, D. Georges, G. Klonner, M. Schmid, S. Dullinger, W. Thuiller, and F.
Guillaume. 2017. A dynamic eco-evolutionary model predicts slow response of alpine
plants to climate warming. Nature Communications **8**:15399.
Coulson, T. 2012. Integral projections models, their construction and use in posing hypotheses in
ecology. Oikos **121**:1337-1350.
Coulson, T., T. Benton, P. Lundberg, S. Dall, B. Kendall, and J.-M. Gaillard. 2005. Estimating
individual contributions to population growth: evolutionary fitness in ecological time.
Proceedings of the Royal Society B: Biological Sciences **273**:547-555.
Coulson, T., B. E. Kendall, J. Barthold, F. Plard, S. Schindler, A. Ozgul, and J.-M. Gaillard. 2017.
Modeling adaptive and nonadaptive responses of populations to environmental change.
The American Naturalist **190**:313-336.
Coulson, T., D. R. MacNulty, D. R. Stahler, R. K. Wayne, and D. W. Smith. 2011. Modeling effects of
environmental change on wolf population dynamics, trait evolution, and life history.
Science **334**:1275-1278.
Coulson, T., and S. Tuljapurkar. 2008. The dynamics of a quantitative trait in an age-structured
population living in a variable environment. The American Naturalist **172**:599-612.
Coulson, T., S. Tuljapurkar, and D. Z. Childs. 2010. Using evolutionary demography to link life
history theory, quantitative genetics and population ecology. Journal of Animal Ecology
79:1226-1240.
Easterling, M. R., S. P. Ellner, and P. M. Dixon. 2000. Size-specific sensitivity: applying a new
structured population model. Ecology **81**:694-708.

Ellner, S. P., D. Z. Childs, and M. Rees. 2016. Data-driven modelling of structured populations.
Springer.
Ellner, S. P., and M. Rees. 2006. Integral projection models for species with complex demography.
The American Naturalist **167**:410-428.
Engen, S., R. Lande, B.-E. Sæther, and F. S. Dobson. 2009. Reproductive value and the stochastic
demography of age-structured populations. The American Naturalist **174**:795-804.
Falconer, D. S. 1960. Introduction to quantitative genetics. Oliver & Boyd.
Fisher, R. A. 1930. The genetical theory of natural selection. Clarendon Press.
Grafen, A. 1988. 28 On the Uses of Data on Lifetime Reproductive Success.*in* T. H. Clutton-Brock,
editor. Reproductive success: studies of individual variation in contrasting breeding
systems. University of Chicago Press.
Hairston Jr, N. G., S. P. Ellner, M. A. Geber, T. Yoshida, and J. A. Fox. 2005. Rapid evolution and the
convergence of ecological and evolutionary time. Ecology letters **8**:1114-1127.
Hendry, A. P. 2016. Eco-evolutionary dynamics. Princeton university press.
Heywood, J. S. 2005. An exact form of the breeder's equation for the evolution of a quantitative
trait under natural selection. Evolution **59**:2287-2298.
Janeiro, M. J., D. W. Coltman, M. Festa-Bianchet, F. Pelletier, and M. B. Morrissey. 2017. Towards
robust evolutionary inference with integral projection models. Journal of Evolutionary
Biology **30**:270-288.
Kruuk, L. E. B. 2004. Estimating genetic parameters in natural populations using the 'animal
model'. Philosophical Transactions of the Royal Society of London. Series B: Biological
Sciences **359**:873-890.
Kruuk, L. E. B., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, and T. Clutton-Brock. 2002.
Antler size in red deer: heritability and selection but no evolution. Evolution **56**:1683-1695.
Lande, R. 1982. A quantitative genetic theory of life history evolution. Ecology **63**:607-615.
Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.
Evolution **37**:1210-1226.
Lefkovitch, L. 1965. The study of population growth in organisms grouped by stages. Biometrics:1-
18.
Lush, J. L. 1943. Animal breeding plans. Iowa State College Press.
Lynch, M., and B. Walsh. 1998. Genetics and analysis of quantitative traits. Sinauer Sunderland,
MA.
Merilä, J., L. Kruuk, and B. Sheldon. 2001a. Cryptic evolution in a wild bird population. Nature
412:76.
Merilä, J., B. Sheldon, and L. Kruuk. 2001b. Explaining stasis: microevolutionary studies in natural
populations. Genetica **112**:199-222.
Merow, C., J. P. Dahlgren, C. J. E. Metcalf, D. Z. Childs, M. E. Evans, E. Jongejans, S. Record, M.
Rees, R. Salguero-Gómez, and S. M. McMahon. 2014. Advancing population ecology with
integral projection models: a practical guide. Methods in Ecology and Evolution **5**:99-110.
Morrissey, M., L. Kruuk, and A. J. Wilson. 2010. The danger of applying the breeder's equation in
observational studies of natural populations. Journal of evolutionary biology **23**:2277-2288.
Nussey, D., A. Wilson, and J. Brommer. 2007. The evolutionary ecology of individual phenotypic
plasticity in wild populations. Journal of Evolutionary Biology **20**:831-844.
Parker, G. A., and J. M. Smith. 1990. Optimality theory in evolutionary biology. Nature **348**:27.
Pelletier, F., D. Garant, and A. Hendry. 2009. Eco-evolutionary dynamics. Philosophical
Transactions of the Royal Society B: Biological Sciences **364**:1483-1489.
Post, D. M., and E. P. Palkovacs. 2009. Eco-evolutionary feedbacks in community and ecosystem
ecology: interactions between the ecological theatre and the evolutionary play.
Philosophical Transactions of the Royal Society B: Biological Sciences **364**:1629-1640.

Price, G. R. 1970. Selection and covariance. Nature **227**:520-521.
Reed, T. E., D. E. Schindler, and R. S. Waples. 2011. Interacting effects of phenotypic plasticity and
evolution on population persistence in a changing climate. Conservation Biology **25**:56-63.
Rees, M., D. Z. Childs, and S. P. Ellner. 2014. Building integral projection models: a user's guide.
Journal of Animal Ecology **83**:528-545.
Rees, M., and S. P. Ellner. 2019. Why so variable: can genetic variance in flowering thresholds be
maintained by fluctuating selection? The American Naturalist **194**:E13-E29.
Sæther, B.-E., and S. Engen. 2015. The concept of fitness in fluctuating environments. Trends in
ecology & evolution **30**:273-281.
Schindler, S., J. M. Gaillard, A. Grüning, P. Neuhaus, L. W. Traill, S. Tuljapurkar, and T. Coulson.
2015. Sex-specific demography and generalization of the Trivers–Willard theory. Nature
526:249.
Siepielski, A. M., M. B. Morrissey, M. Buoro, S. M. Carlson, C. M. Caruso, S. M. Clegg, T. Coulson, J.
DiBattista, K. M. Gotanda, and C. D. Francis. 2017. Precipitation drives global variation in
natural selection. Science **355**:959-962.
Smallegange, I. M., H. Caswell, M. E. Toorians, and A. M. Roos. 2017. Mechanistic description of
population dynamics using dynamic energy budget theory incorporated into integral
projection models. Methods in Ecology and Evolution **8**:146-154.
Smallegange, I. M., and T. Coulson. 2013. Towards a general, population-level understanding of
eco-evolutionary change. Trends in ecology & evolution **28**:143-148.
Steiner, U. K., S. Tuljapurkar, and T. Coulson. 2014. Generation time, net reproductive rate, and
growth in stage-age-structured populations. The American Naturalist **183**:771-783.
Steiner, U. K., S. Tuljapurkar, T. Coulson, and C. Horvitz. 2012. Trading stages: life expectancies in
structured populations. Experimental gerontology **47**:773-781.
Terhorst, C. P., P. C. Zee, K. D. Heath, T. E. Miller, A. I. Pastore, S. Patel, S. J. Schreiber, M. J. Wade,
and M. R. Walsh. 2018. Evolution in a community context: trait responses to multiple
species interactions. The American Naturalist **191**:368-380.
Traill, L. W., S. Schindler, and T. Coulson. 2014. Demography, not inheritance, drives phenotypic
change in hunted bighorn sheep. Proceedings of the National Academy of Sciences
111:13223-13228.
Travis, J., D. Reznick, R. D. Bassar, A. López-Sepulcre, R. Ferriere, and T. Coulson. 2014. Do eco-evo
feedbacks help us understand nature? Answers from studies of the Trinidadian guppy.
Pages 1-40 Advances in Ecological Research. Elsevier.
Turchin, P. 2003. Complex population dynamics: a theoretical/empirical synthesis. Princeton
university press.
van Tienderen, P. H. 2000. Elasticities and the link between demographic and evolutionary
dynamics. Ecology **81**:666-679.
Via, S., and R. Lande. 1985. Genotype-environment interaction and the evolution of phenotypic
plasticity. Evolution **39**:505-522.
Watson, H. W., and F. Galton. 1875. On the probability of the extinction of families. The Journal of
the Anthropological Institute of Great Britain and Ireland **4**:138-144.
White, G. C., and K. P. Burnham. 1999. Program MARK: survival estimation from populations of
marked animals. Bird study **46**:S120-S139.
Wilson, A. J., J. M. Pemberton, J. Pilkington, D. W. Coltman, D. Mifsud, T. H. Clutton-Brock, and L.
B. Kruuk. 2006. Environmental coupling of selection and heritability limits evolution. PLoS
Biology **4**:e216.
Yoshida, T., L. E. Jones, S. P. Ellner, G. F. Fussmann, and N. G. Hairston Jr. 2003. Rapid evolution
drives ecological dynamics in a predator–prey system. Nature **424**:303.
797

Appendix 1

A key premise of the standard quantitative genetic approach is that the genotype-phenotype map

can be described as $Z = A + E$ (Falconer 1960). If a phenotypic trait is measured in a particular

unit, say length in millimetres, then A and E also take these units. The second premise is that

selection operates on the entire phenotype via a selection differential. If we consider a linear

fitness function $E(w) = \beta_0 + \beta_1 Z$ we can also write $E(w) = \beta_0 + \beta_1(A + E) = \beta_0 + \beta_1 A + \beta_1 E$.

This equation says the effect of a one unit change on fitness in the additive genetic and

environmental components of the phenotype is identical.

A linear regression slope is defined as the covariance between the response and predictor variable

divided by the variance in the predictor variable. Consequently,

$$\bar{w}_t = \beta_0 + \frac{cov(w, A)}{var(A)} A + \frac{cov(w, E)}{var(E)} E,$$

where $\frac{cov(w,A)}{var(A)} = \frac{cov(w,E)}{var(E)}$. If $\frac{cov(w,A)}{var(A)} \neq \frac{cov(w,E)}{var(E)}$ then $Z \neq A + E$ but will be of the form $Z = A + \gamma E$

where $\gamma \neq 1$.

Some authors have argued that selection can operate in contrasting ways on the environmental

and additive genetic components of the phenotype while assuming $z_t = A_t + E_t$ (Morrissey et al.

2010). In the most extreme cases they argue that selection on the breeding value is in one

direction, while selection on the environmental component of the phenotype is in a contrasting

direction such that $\gamma < 0$ while A_t and E_t do not covary (Bonnet et al. 2017). Although

mathematically possible, such a genotype-phenotype map seems biologically implausible. We

have been unable to identify a biologically plausible genotype-phenotype map where 1 unit

increase in A_t will result in an increase in fitness while a 1 unit increase in E_t will result in a

decrease without invoking some dependence of A_t and E_t on one another.

A more plausible genotype-phenotype map would involve phenotypic plasticity where a breeding

value A_t produces a different phenotypic value z_t in different environments. If $z_t = A_t + E_t$ this

requires E_t to vary with the environment θ, such that E can be made a function of the

environment $E(\theta_t)$. Phenotypic plasticity can then be captured with a number of formulations

such as $z_t = A_t + E(\theta_t)$ or $z_t = A_t + E(\theta_t) + \alpha A_t E(\theta_t)$ (Coulson et al. 2017). In the first

formulation, the mean and variance of E_t can vary with the environment, but the genotype-

phenotype map remains $z_t = A_t + E_t$ such that the slope between the environmental component

of the phenotype and fitness equals the slope between the additive genetic component of the

phenotype and fitness. In contrast, in the second formulation, the breeding value-by-environment

interaction means that the slope between fitness and each component of the phenotype differs.

Empirical evidence of so-called "selection on the environmental component of the phenotype"

(Bonnet et al. 2017) is consequently likely evidence of a genotype-by-environment interaction:

something that cannot be detected with widely used statistical analyses of observational data

(Kruuk 2004).

Appendix 2

Four approaches can be used to specify how inheritance alters the distribution of breeding values

between selected parents and their offspring. Approach 1: A convolution of the distribution of

breeding values in mothers and fathers is taken to construct a distribution of mid-parental values

assuming random mating. The distribution of mid-parental values is then convolved with a normal

distribution with a mean of 0 and a constant variance to construct the distribution of breeding

values in offspring. Approach 2: Breeding values in the offspring generation are assumed to be

normally distributed, with a mean equal to that of the mid-point parental breeding value

distribution, and a constant variance that does not change across generations. Approach 3: The

distribution of breeding values of parents is operated on by an inheritance function which

maintains the parental mid-parental mean value and injects Gaussian distributed segregation

variance around each mid-point breeding value. Approach 4: The distribution of breeding values in

parents and offspring is identical. As long as selection does not generate distributions with

substantial skew or kurtosis, the first three of these approaches give very similar predictions.

Various properties of these four approaches are summarised in Table A2.1

Table A2.1. Properties of transition functions for breeding values.

	Approach 1	Approach 2	Approach 3	Approach 4
Are the breeding values in offspring normally distributed?	Not necessarily	yes	Not necessarily	Not necessarily

Does the distribution of breeding values in mid-parents and offspring have the same mean?	yes	yes	yes	yes
Does the distribution of breeding values in parents and offspring have the same shape?	No	No	No	yes

852　Appendix 3

853　The Soay sheep model consisted of the following parameterisations:

854
$$S(A_t + E_t) = \frac{1}{1 + e^{-(-0.2392 + 1.1175(A_t + E_t) - 0.0043N_t)}}$$

855
$$R(A_t + E_t) = \frac{1}{1 + e^{-(-1.7839 + 0.1090(A_t + E_t) - 0.0032N_t)}}$$

856
$$G(A'_{t+1} | A_t) = A_t$$

857
$$G(E'_{t+1} | E_t) = norm(\mu_G(E_t), \sigma_G^2(E_t)),$$

858　where $\mu_G(E_t) = 10.603 + 0.6049E_t - 0.0026N_t$ and $\sigma_G^2(E_t) = 2.874 - 0.0124E_t$

859
$$D(A'_{t+1} | A_t) = norm(\mu_D^\omega(A_t), \sigma_D^2(A_t)),$$

860　where $\mu_D^\omega(A_t)$ is the mean breeding value post-fertility selection, and , $\sigma_D^2(A_t)$ is the additive

861　genetic variance in offspring at birth, which we assumed to be constant and Gaussian distributed.

862　Finally,

863
$$D(E'_{t+1} | E_t) = norm(\mu_D(E_t), \sigma_D^2(E_t)),$$

where $\mu_D(E_t) = 9.8065 + 0.2544E_t - 0.0069N_t$ and , $\sigma_D^2(E_t) = 4.581 - 0.1196E_t$.

We started each simulation with mean values of $\bar{A}_t = 5.5$ and $\bar{E}_t = 19.5$, an additive genetic

variance of 4 and a variance in the environmental component of 6.

We selected the starting population structure of each simulation by using the equilibrium

population structure from an ecological simulation where the additive genetic variance was set to

0 so evolution cannot occur. Each simulation was run for 100 time-steps.