1 Dynamic Price and Breeder's equations for variable environments

Tim Coulson[1,2], Tomos Potter[1], Anja Felmy[1]

[1] Department of Zoology, University of Oxford, Mansfield Road, Oxford, OX1 3SZ, UK

[2] tim.coulson@zoo.ox.ac.uk

ABSTRACT

The Breeder's and Price equations are static models of evolution. Recent work has demonstrated

how components of these equations, such as selection differentials and heritabilities, can be

calculated for each time-step from dynamic, evolutionarily-explicit, structured population models.

These dynamic models consist of functions that describe how environmental factors impact

associations of genotypes and phenotypic traits with survival, reproduction, development, and

non-genetic inheritance. Because evolutionarily explicit versions of these structured models can

include feedbacks and can capture all the routes via which environmental variation can impact

evolution, they i) provide a more powerful predictive tool for modelling eco-evolution in natural

settings than existing approaches, and ii) reveal how key evolutionary parameters incorporated

into static models change with time as evolution, or environmental change, proceeds.

INTRODUCTION

Eco-evolutionary dynamics occur when selection on heritable phenotypic traits and the population

growth rate within a focal species change together (Hairston Jr et al. 2005). Eco-evolutionary

feedbacks then arise when a change in the population growth rate alters the strength of selection

within a species, and vice versa (Patel et al. 2018). By using dynamic, evolutionarily-explicit,

structured models (Coulson et al. 2017) to examine the dynamics of both population growth and

terms in the Breeder's and Price equations (Coulson et al. 2010, Childs et al. 2016), we show how

eco-evolutionary dynamics and feedbacks influence evolution, phenotypic trait dynamics, and

population growth. Our approach is based on understanding the interplay of fitness functions with

functions that describe development, and genetic and non-genetic inheritance (see also (Wallace

et al. 2013)). Fitness functions define how the association between phenotypic trait values and

individual absolute fitness (or survival and reproductive components of fitness) varies with

environmental drivers; development functions characterise the phenotypic development of

surviving individuals over a time-step as a function of their environment; inheritance functions

capture both genetic inheritance and environmentally-induced, non-genetic inheritance (Childs et

al. 2016, Coulson et al. 2017).

Adaptive evolution requires directional selection on a heritable phenotypic trait (Fisher 1930). The

strength of selection is determined by the phenotypic variance and the association between the

phenotypic trait and absolute fitness, and is measured with a selection differential (Price 1970,

Lande and Arnold 1983). Selection alters the distribution of the trait within a population within a

generation. For evolution to occur, this change must, at least partially, be reflected in the

distributions of alleles that determine trait values within the population (Falconer 1960). Rules of

genetic inheritance mean that allele frequencies among selected parents are expected to be

observed in offspring, although processes such as mutation and genetic drift can lead to

departures from this expectation (Crow and Kimura 1970). For eco-evolutionary dynamics and

feedbacks to occur, it is necessary, but not sufficient, that the strength of selection is influenced by

the population growth rate. The selection differential can be written as the covariance between

phenotypic trait values and individual absolute fitness, divided by the mean individual absolute

fitness (Heywood 2005). When the mean individual absolute fitness over a time-step is shorter

than a generation, it is the population growth rate (Fisher 1930). The strength of selection is

consequently influenced by the population growth rate, as long as the covariance between

phenotypic trait values and individual absolute fitness does not vary systematically with the

population growth rate.

A dynamical consequence of selection being partially determined by the covariance between

phenotypic trait values and individual absolute fitness, is that if the mean of one quantity changes,

the mean of the other will also change. This is true as long as i) the function linking the two

quantities does not change with time, and ii) there is variation in each quantity within the

population. For example, evolution of a phenotypic trait will cause an increase in mean absolute

fitness (i.e. the population growth rate) (Fisher 1930). Under these scenarios, both eco-

evolutionary dynamics and feedbacks will occur (Coulson et al. 2017). This is because i) the mean

of the phenotypic trait and the population growth rate will evolve together, ii) an increase in the

population growth rate will increase the size of the denominator in the selection differential, iii)

this will reduce the strength of selection which will, in turn, iv) reduce the rate of evolution of the

population growth rate. The speed of the eco-evolutionary dynamics, and the magnitude of the

eco-evolutionary feedback, both depend upon the association between phenotypic trait values

and individual absolute fitness, the phenotypic variance, and the form of the map between

genotype and phenotype.

In artificial settings it is possible to control i) the association between phenotypic trait values and

individual absolute fitness, ii) the phenotypic variance, and iii), to some extent, the way that

environmental variation might influence the genotype-phenotype map (Falconer 1960). Such

control does not exist in the natural world, where density-dependent feedbacks and

environmental variation can generate temporal change in each of these three quantities. For

example, if evolution occurs while the association between phenotypic trait values and individual

absolute fitness remains constant in time, populations grow hyper-exponentially (Witting 2002).

This is not observed in nature because density-dependent processes such as food limitation and

predation prevent long-term hyper-exponential (or even exponential) population growth.

Negative density-dependence operates to lower the elevation of the association between

phenotypic trait values and individual absolute fitness, which in turns reduces mean individual

absolute fitness (the population growth rate) (Coulson et al. 2017).

We use evolutionarily-explicit, density-dependent, structured population models to examine how

environmental variation can alter the form of the association between phenotypic trait values and

individual absolute fitness, the genotype-phenotype map, and the phenotypic variance, thereby

impacting eco-evolution and potentially leading to eco-evolutionary feedbacks.

THEORETICAL UNDERPINNINGS

Understanding eco-evolutionary dynamics and feedbacks requires dynamic models. However,

standard evolutionary theory widely used by empiricists is static, in that it only provides

predictions over a single time-step (Falconer 1960, Morrissey et al. 2010). Nonetheless, this static

theory is useful as selection differentials on which it is based provide exact descriptions of the

strength of observed selection on phenotypic traits (Lande and Arnold 1983), and the Price

equations builds on this to provide a complete description of evolution (Price 1970). If models of

eco-evolution are to link with the understanding these static models have provided, they must

predict time series of components of these static models (Coulson et al. 2010, Coulson et al. 2017,

Coulson et al. 2018). We consequently start by considering these models.

We start with the Breeder's equation. Following standard quantitative genetic theory, we assume

that an individual's phenotypic trait value z_t can be decomposed into a breeding value A_t and an

environmental component E_t, so that $z_t = A_t + E_t$. Next, we write the Breeder's equation in

terms of individual absolute fitness w_t (Heywood 2005) and assume an annual life history and an

annual population census such that viability selection can be ignored and mean absolute fitness is

equal to the mean recruitment rate $\bar{w}_t = \bar{r}_t$, where $\bar{r}_t$ is defined as the per capita production of

offspring between times t and $t + 1$ that survive to enter the population at time $t + 1$

$$\Delta \bar{z}_t = \frac{cov(w_t, A_t + E_t)}{\bar{w}_t} \frac{var(A_t)}{var(A_t + E_t)}. \tag{1}$$

$\Delta \bar{z}_t$ is the change in the mean phenotypic trait value over a generation (=time-step for our annual

life history) (Lush 1943), the first term on the right-hand side is a phenotypic selection differential,

and the second is the narrow-sense heritability h^2. The variance in the breeding values at time t is

denoted $var(A_t)$ – the additive genetic variance. Equation (1) assumes that change in $\bar{A}_t$ caused

by selection within the parental generation is transmitted to the offspring generation, and that

change in $\bar{E}_t$ is not.

We can rewrite equation (1) as $\Delta \bar{z}_t = \frac{\beta_1 var(A_t)}{\bar{w}_t}$ where β_1 is the slope of a linear regression

between individual absolute fitness and individual phenotypic trait values (the derivative $\frac{d\bar{w}_t}{d\bar{z}_t}$).

Note that this makes β_1 different from a selection gradient (Lande and Arnold 1983), which is

computed using relative fitness values (e.g. $\beta_1 / \bar{w}_t$ in our notation).

In this formulation, $\bar{w}_t = \bar{r}_t$ is mean fitness and the population growth rate. Assuming that β_1 and

$var(A_t)$ do not systematically vary with $\bar{w}_t$, it is apparent that the rate of evolution $\Delta \bar{z}_t$ and the

population growth rate $\bar{w}_t$ are intimately linked. For example, if we assume a linear fitness

function $w_t = \beta_0 + \beta_1 z_t$ we can write:

$$\Delta \bar{z}_t = \frac{\beta_1 var(A_t)}{\beta_0 + \beta_1 [\bar{A}_t + \bar{E}_t]}. \tag{2}$$

The equations above assume that selection is the only process that alters $\bar{A}_t$ or $\bar{E}_t$ from one

generation to the next. This is one reason why the Breeder's equation does not provide a

complete description of evolution of the mean of a phenotypic trait. Non-genetic inheritance and

phenotypic plasticity, that can influence the value of $\bar{E}_t$ and contribute to evolutionary dynamics,

are processes that are not included in the Breeder's equation. Put another way, the Breeder's

equation assumes that the genotype-phenotype map does not vary with time due to processes

that can alter $\bar{E}_t$. These processes can be captured, along with selection, with the Price equation

(Price 1970).

The Price equation provides a complete description of the change in the mean of a phenotypic

trait over a generation. Continuing to assume $z_t = A_t + E_t$,

$$\Delta \bar{z}_t = \frac{cov(w_t, A_t + E_t)}{\bar{w}_t} + [\bar{A}_{t+1} - \bar{A}_{\omega,t}] + [\bar{E}_{t+1} - \bar{E}_{\omega,t}]. \tag{3}$$

$\bar{A}_{\omega,t}$ and $\bar{E}_{\omega,t}$ respectively describe the mean of the breeding value and environmental component

of the phenotype among selected parents.

The first term on the right-hand side is a selection differential. The second term describes how

imperfect genetic inheritance (e.g., mutation, recombination) could alter the mean breeding value

between offspring and their parents, and the third how non-genetic inheritance (e.g., epigenetics)

alters the mean environmental component of the phenotypic trait between offspring and their

parents.

The Breeder's equation is a special form of the Price equation that assumes that $[\bar{A}_{t+1} - \bar{A}_{\omega,t}] =$

0 and $\frac{cov(w_t, E_t)}{\bar{w}_t} = -[\bar{E}_{t+1} - \bar{E}_{\omega,t}]$. Biologically this means that genetic inheritance does not change

$\bar{A}_t$ (i.e. there is no transmission bias), and that $\bar{E}_t$ at birth remains identical in each generation.

The first of these assumptions is reasonable for large populations; the second is justified if the

environmental component of the phenotypic trait is determined by completely random

developmental noise (Falconer 1960) and not by any variation in the environment. If the

environmental component of the phenotypic trait is dependent upon non-random aspects of the

external environment (including non-genetic inheritance), then the assumption will be violated

(Kruuk et al. 2002). If this is the case, the Price equation is required to describe evolution.

In an annual life history $\bar{w}_t = \bar{r}_t$; in a non-annual life history, the population growth rate and mean

fitness over a time-step are the sum of mean survival and mean recruitment, $\bar{w}_t = \bar{s}_t +$

$\bar{r}_t$. Replacing $\bar{w}_t$ with $\bar{r}_t + \bar{s}_t$ and expanding (3) we can write the Price equation in terms of fitness

and transition functions acting through survival, recruitment, development and genetic and non-

genetic inheritance on the distribution of z. In this form, the Price equation applies to iteroparous

species:

$$\Delta \bar{z}_t = \frac{cov(s_t, A_t + E_t)}{\bar{w}_t} + \qquad (4a)$$

Effect of viability selection on components of the phenotype.

$$+ \frac{cov(r_t, A_t + E_t)}{\bar{w}_t} + \qquad (4b)$$

Effect of fertility selection on components of the phenotype.

$$+ \frac{\bar{r}_t}{\bar{w}_t}[\bar{A}'_{t+1} - \bar{A}_{\omega r,t}] + \qquad (4c)$$

Effect of genetic inheritance on the mean breeding value. $\bar{A}'_{t+1}$ is the mean breeding value amongst recruits; $\bar{A}_{\omega r,t}$ is the mean breeding value among

selected parents. Equals zero: inheritance does not change the mean breeding value between selected parents and their offspring.

$$+\frac{\bar{s}_t}{\overline{w}_t}[\bar{A}''_{t+1} - \bar{A}_{\omega_s,t}] + \qquad (4d)$$

Development of the breeding value within surviving individuals. $\bar{A}''_{t+1}$ is the mean breeding value amongst surviving individuals at time $t+1$; $\bar{A}_{\omega_s,t}$ is the mean breeding value at time t amongst individuals that will survive from time t to time $t+1$. Equals zero: breeding values do not change within individuals as they age.

$$+\frac{\bar{r}_t}{\overline{w}_t}[\bar{E}'_{t+1} - \bar{E}_{\omega_r,t}] + \qquad (4e)$$

Effect of non-genetic inheritance on the mean environmental component of the phenotypic trait: non-genetic inheritance can change the mean of the environmental component of the phenotype between selected parents and their offspring.

$$+\frac{\bar{s}_t}{\overline{w}_t}[\bar{E}''_{t+1} - \bar{E}_{\omega_s,t}]. \qquad (4f)$$

Effect of phenotypic plasticity on the development of the mean environmental component of the phenotypic trait: the environmental component of the phenotype can change within individuals as they age.

This provides a complete description of all of the processes that can influence change in the mean

phenotype in an iteroparous species over a time-step. The formulation does assume $z_t = A_t + E_t$

and that all processes remain constant with age. The latter assumption could be relaxed using the

approach in (Coulson and Tuljapurkar 2008). Note that some authors have further decomposed

$cov(r_t, A_t + E_t)$ into $cov(r_t, A_t) + cov(r_t, E_t)$ and assumed that these two quantities can change

independently of one another (Bonnet et al. 2017). We do not do this here as this violates the

additivity assumption of $z_t = A_t + E_t$ of quantitative genetics (Appendix 1) that we are currently

employing.

Each of the terms in the static Breeder's and Price equations are emergent properties of the

distribution of the phenotypic trait (and its components) at time t, fitness functions determining

terms in equation (*4a*) and (*4b*), and transition functions determining terms in equations (*4c*) to

(*4f*) (Coulson et al. 2010, Childs et al. 2016). Integral projection models (IPMs), which calculate

changes in the distribution of a continuous trait by integrating over a series of fitness and

transition functions (Coulson et al. 2010, Childs et al. 2016) provide a framework with which to

make the Price and Breeder's equations dynamic. To do this, it is consequently necessary to

specify these functions (Coulson et al. 2017) that can then be combined into a model to iterate

forward the dynamics of the distribution of $z_t = A_t + E_t$.

In the case of an annual life history, a reproductive fitness function, $R(A_t + E_t)$, is used to map

the bivariate distribution of breeding values and environmental components of the phenotype at

time t, $n(A_t, E_t)$, to a new distribution at time $t + 1$, $n(A'_{t+1}, E'_{t+1})$ (Ellner et al. 2016),

$$n(A'_{t+1}, E'_{t+1}) = \int [H(A'_{t+1}, E'_{t+1}|A_t, E_t) R(A_t + E_t) n(A_t, E_t)] dA_t dE_t. \qquad (5)$$

$H(A'_{t+1}, E'_{t+1}|A_t, E_t)$ is a transition function that describes the probability of transitioning from

trait value $A_t + E_t$ at time t to all possible trait values $A'_{t+1} + E'_{t+1}$ at time $t + 1$ (Easterling et al.

2000). Trait value $z_t = A_t + E_t$ is expressed in the focal individual, while trait values $z'_{t+1} =$

$A'_{t+1} + E'_{t+1}$ are expressed in offspring. The transition function is needed when predicting

evolutionary change over multiple time-steps if offspring that recruit to the population at time $t +$

1 and their parents at time t do not perfectly resemble one another.

$H(A'_{t+1}, E'_{t+1}|A_t, E_t)$ can be formulated to capture the second term on the right-hand side of the

Breeder's equation (*1*) or the last two terms on the right-hand side of the Price equation (*3*), and

so can describe genetic and non-genetic inheritance. A dynamic form of the Breeder's equation (*1*)

requires $H(A'_{t+1}, E'_{t+1}|A_t, E_t)$ to be specified such that it produces a distribution $n(A'_{t+1}, E'_{t+1})$

that has means of $\bar{A}'_{t+1} = \bar{A}_{\omega,t}$ and $\bar{E}'_{t+1} = \bar{E}_{\omega,t}$ and a variance-covariance matrix

$\begin{pmatrix} var(A_t) & 0 \\ 0 & var(E_t) \end{pmatrix}$ where $var(A_t)$ can be selected to be a constant, or to change with time as a

function of selection, and $var(E_t)$ is constant (Childs et al. 2016, Coulson et al. 2017, Rees and

Ellner 2019). There are a number of ways this can be achieved (Appendix 2). Any formulation of

$H(A'_{t+1}, E'_{t+1} | A_t, E_t)$ that results in $\bar{A}'_{t+1} \neq \bar{A}_{\omega,t}$ and/or $\bar{E}'_{t+1} \neq \bar{E}_{\omega,t}$ provides a dynamic form of

the Price equation (*3*, 4). Once the transition and fitness functions are parameterised, derivations

from Coulson *et al.* (2010) can then be used to create time series of each of the components in the

various forms of the Price and Breeder's equations described above.

Equation (*5*) can be extended to incorporate per-time-step survival and recruitment functions to

allow models of iteroparous species. Models now include two transition functions: a development

function $G(E'_{t+1} | A_t, E_t)$ operating on surviving individuals, and an inheritance function

$D(E'_{t+1} | A_t, E_t)$ that captures the similarity between parental and offspring traits (Easterling et al.

2000, Ellner and Rees 2006, Coulson 2012). The development function operates with a survival

fitness function $S(A_t + E_t)$ that determines the strength of viability selection, while the

inheritance function operates with the recruitment fitness function $R(A_t + E_t)$ that determines

the strength of fertility selection:

193
$$n(A'_{t+1}, E'_{t+1}) = \int \begin{bmatrix} D(E'_{t+1}|A_t, E_t)R(A_t + E_t) + \cdots \\ \cdots + G(E'_{t+1}|A_t, E_t)S(A_t + E_t) \end{bmatrix} n(A_t, E_t) dA_t dE_t. \qquad (6)$$

Each process that can contribute to the change in the mean phenotype in an iteroparous species

(i.e. Equation(4)) is captured in this IPM: term (*4a*) is determined by $S(A_t + E_t)n(A_t, E_t)$, term

(*4b*) by $R(A_t + E_t)n(A_t, E_t)$, terms (*4c*) and (*4e*) by $\int D(E'_t|A_t, E_t)R(A_t + E_t)n(A_t, E_t)dA_t dE_t$

and terms (*4d*) and (*4f*) by $\int G(E'_t|A_t, E_t)S(A_t + E_t)n(A_t, E_t)dA_t dE_t$ (Childs et al. 2016).

METHODS

Our aim here is to incorporate environmental trends and feedbacks caused by density-

dependence into the functions in equations (*5*) and (*6*). We will examine how this impacts the

association between phenotypic trait values and individual absolute fitness, the phenotypic

variance, and the map between genotype and phenotype, and how, ultimately, this generates

temporal dynamics in the components of the Breeder's and Price equations, rates of evolution,

phenotypic trait dynamics, and population growth.

We start with a model of the form of equation (5) with the phenotypic trait separated into its

additive genetic and environmental components, $z_t = A_t + E_t$, but where fitness is not separated

into its survival and recruitment components. We assume a linear association between phenotypic

trait values and individual absolute fitness and iterate $n(A_t, E_t)$ given the following

parameterisation,

$$w(A_t + E_t) = -0.2 + 0.05(A_t + E_t), \qquad (7)$$

and a starting bivariate normal distribution $n(A_{t=1}, E_{t=1})$ with means $\bar{A}_t = \bar{E}_t = 10$ and a

variance-covariance matrix $\sigma^2 = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$. The choice of parameter values is for illustration rather

than representing any particular system. The mean of the breeding value distribution does not

change between parents and their offspring.

We ran four model scenarios. In scenarios 1) and 2) the mean and variance of the environmental

component of the phenotype remained constant across generations, while in scenarios 3) and 4)

the mean of the environmental component declined with time such that $\bar{E}_t = 10 - 0.05t$. In

scenarios 1) and 3) selection eroded additive genetic variation, while in scenarios 2) and 4) it did

not. Scenarios 1) and 2) are dynamic forms of the Breeder's equation, while scenarios 3) and 4)

are dynamic forms of the Price equation. The last two scenarios capture a process where some

aspect of the environment can determine the dynamics of the environmental component of the

phenotype (Kruuk et al. 2002).

The linear assumption of the simple model is unrealistic because litter size and survival rates are

bounded. In addition, in variable environments, survival, recruitment, development and

inheritance functions for the environmental component of the phenotype depend on

environmental factors such as the population size (Kruuk et al. 2002, Bassar et al. 2017).

Moreover, although breeding values remain constant with age, the environmental component of

the phenotypic trait can change within individuals – phenotypic plasticity (Falconer 1960). We

consequently developed a more complex, density-dependent model, where fitness is separated

into its survival and recruitment components, and where the environmental component of the

phenotype can develop within surviving individuals, and can vary between parents and their

offspring. We took a published density-dependent Integral Projection Model (IPM) of the

dynamics of body size from females in an individually marked population of Soay sheep (*Ovis*

aries) living on the island of Hirta in the St. Kilda archipelago in Scotland (Coulson 2012). The

published IPM was modified to be of the form of equation (*6*). Functions and parameter values are

provided in Appendix 3.

We selected the starting population structure of each evolutionarily explicit simulation by using

the equilibrium population structure from an ecological simulation where the additive genetic

variance was set to 0 so evolution cannot occur. Each simulation was run for 100 time-steps.

As well as the baseline model, in which the survival, recruitment, development and inheritance

functions were all density-dependent, we ran four simulations where we set the density-

dependence parameter to 0 in either the survival fitness function, the recruitment fitness

function, the development function, or the inheritance function for E_t. Although this is a large

perturbation, it reveals interesting dynamics whereby perturbing one function influences the role

of other functions on eco-evolution.

RESULTS

In our evolutionarily explicit IPM with a linear fitness function, evolution, defined as a change in

the mean breeding value $\Delta \bar{A}_t$, was fastest when additive genetic variance was not eroded by

selection, and when, simultaneously, there was a negative temporal trend in the mean

environmental component of the phenotype $\bar{E}_t$ (Figure 1(A)). The negative temporal trend in $\bar{E}_t$

accelerated the rate of evolution by reducing the population growth rate $\bar{w}_t$ (Figure 1(B)). In spite

of accelerating the rate of evolution, the trending values of $\bar{E}_t$ with time slowed the rate of

phenotypic change, as positive change in $\bar{A}_t$ was countered by negative change in $\bar{E}_t$ (Figure 1(C)).

The population growth rate $\bar{w}_t$, which is also mean fitness, increased fastest when additive genetic

variance was not eroded, and $\bar{E}_t$ did not decline with time (Figure 1(C)). As expected, linear

selection reduced the additive genetic variance if it was not replaced in each generation (Figure

1(D)). When there was no temporal trend in $\bar{E}_t$, the association between the mean breeding value

and the population growth rate was linear (Figure 1(E)), while it was slightly curvilinear when $\bar{E}_t$

declined with time. As the association between phenotypic trait values and individual absolute

fitness was linear, there was, by definition, a linear association between the mean phenotypic trait

and the population growth rate, with a slope of β_1 (Figure 1(F)).

We can combine these insights to graphically illustrate the evolutionary, phenotypic, and

population-growth dynamics our results reveal with a plot of A_t against E_t that includes fitness

clines (Figure 2(A)). Each diagonal dashed line represents equivalent values of the phenotypic trait

z_t made from different combinations of A_t and E_t. Because the association between phenotypic

trait values and individual absolute fitness was linear and included no additional terms beyond β_0

and β_1, these dashed lines also represent the population growth rate. The darker the colour of the

line, the larger the value of z_t and the population growth rate. The central solid line contours

represent the bivariate distribution of A_t and E_t at time $t = 0$ when simulations are begun. The

green, red and purple arrows represent phenotypic change. When evolution was cryptic (blue

arrow), meaning that neither $\bar{z}_t$, nor $\bar{w}_t$, changed, $\bar{A}_t$ increased at the same rate as $\bar{E}_t$ decreased

such that $\Delta\bar{A}_t - \Delta\bar{E}_t = 0$. Because $\bar{w}_t$ did not change in this situation, evolution proceeded rapidly

as the denominator in the selection differential remained constant with time. Dynamic forms of

the Price equation are required to capture such cryptic evolution. The green line represents

dynamics predicted by dynamic forms of the Breeder's equation, with $\bar{A}_t$ and $\bar{w}_t$ evolving, thus

resulting in phenotypic change, but $\bar{E}_t$ remaining constant with time. Evolution occurred more

slowly than when it was cryptic because $\bar{w}_t$ increased with time, increasing the denominator of

the selection differential. The red line captures evolutionary dynamics where the mean breeding

value and the mean environmental component of the phenotype change in the same direction,

increasing both $\bar{z}_t$ and $\bar{w}_t$. A dynamic form of the Price equation is required to capture such

dynamics, and evolution proceeded even more slowly than in the previous two examples, because

$\bar{w}_t$ increased even faster. Finally, the purple arrow represents no evolution, but phenotypic

change due solely to change in $\bar{E}_t$ – i.e. via non-genetic inheritance (or phenotypic plasticity in

iteroparous species). A dynamic form of the Price equation is required to capture such dynamics.

We now turn to the more complex Soay sheep model that extends insights from the linear model

in three ways: i) it is for an iteroparous life history, ii) it incorporates density-dependence in the

survival and recruitment functions (thus reducing the intercepts of the fitness functions with time

as evolution proceeds), and iii) it has non-linear fitness functions (meaning that the slope of the

association between phenotypic trait values and individual absolute fitness will vary as evolution

proceeds).

Figure (2(B)) shows the joint dynamics of $\bar{A}_t$ and $\bar{E}_t$ for each of the simulations of body size in

female Soay sheep. We do not include fitness clines such as those in Figure (2(A)) because the

incorporation of density-dependence complicates the association between $\bar{z}_t$ and $\bar{w}_t$. Before we

explore these results in further detail, it is helpful to appreciate that change in $\bar{z}_t$ can alter both

the elevation (blue dots) and slope (red lines) of the expected association between phenotypic

trait values and individual absolute fitness (red lines; Figure (2(C))). The way a system evolves

depends on both the rate that phenotypic change alters the location of the distribution of

phenotypic trait values on a fitness function, and the rate at which evolution of the population

growth rate alters the form of the fitness function itself.

Compared to the baseline model where all four functions were density-dependent, removing

density-dependence from the recruitment and inheritance functions in the modified sheep model

increased the rate of evolution of body size, while removing it from the survival and development

function decreased it (Figure 2(B)). By far the largest effect was caused by perturbing the survival

function.

Temporal changes in $\bar{E}_t$ were negative in all models, with the greatest change observed when

density-dependence was removed from the recruitment function (Figure 2(B)). This negative trend

only partially countered the positive trends in $\bar{A}_t$ (Figure 3(A, B)), such that $\bar{z}_t$ increased with time

in all models (Figure 3(B)), but to a lesser extent than $\bar{A}_t$ (Figure 3(A)).

$\bar{w}_t$ evolved at similar rates in the baseline model and models where density-dependence was

removed from the recruitment, inheritance and development functions (Figure 3(C)). In contrast,

$\bar{w}_t$ evolved much more slowly when density-dependence was removed from the survival function

(Figure 3(C)), although initial population size was much larger in the density-independent survival

model than in any of the other models (Figure 2(B)). The additive genetic variance increased

slightly with time when density-dependence was removed from the survival, recruitment and

development functions, but remained constant when inheritance was density-independent, as

well as in the baseline model (Figure 3(D)).

Removing density-dependence from each fitness function impacted evolutionary, phenotypic and

population-size change via a number of different routes. First, removing density-dependence

resulted in different starting population sizes and structures (Figures 4(A)-(E)). In the baseline

model and the model with density-dependence removed from the development function, there

was clear bimodality in the distribution of z_t throughout the simulation (Figures 4(A, E)). The

bimodality reflects a population with similar proportions of smaller juveniles and larger adults.

There was weaker evidence of a bimodal distribution at time $t = 1$ for the other three models,

with bimodality weakening as the simulations proceeded (Figures 4(B-D)). In models with density-

independent survival and inheritance, small juveniles were comparatively rare from the start, and

became rarer still over time. In contrast, in the density-independent recruitment model, the

population was initially dominated by small juveniles, but then also became increasingly

dominated by larger adults as the simulation proceeded.

The rates of evolution of the phenotypic trait and mean fitness are determined by i) β_1, ii) $\overline{w}_t$

(which depends upon $n(z_t)$ and the fitness functions), and iii) $var(A_t)$. Because the models were

density-dependent, the mean population growth rate tended towards unity, and differed little

between simulations (Figure 3(C)). Relatively little of the dynamical difference between the

models was consequently due to differences in $\overline{w}_t$ across simulations. Similarly, because the

additive genetic variance remained approximately constant in all simulations (Figure 3(D)), this too

was not a major driver of the dynamics we report from the sheep model.

Most of the dynamical differences between the simulations were consequently caused by

processes generating variation in β_1: the covariance between individual phenotypic trait values

and individual absolute fitness and the phenotypic variance. These two terms tended to show

similar temporal dynamics (Figure 4(F-H)). Consequently, the derivative between mean survival

and the phenotype (the strength and direction of viability selection) remained approximately

constant in the baseline model and the density-independent development and inheritance

models, while it trended downwards as the simulation proceeded in the density-independent

survival and recruitment models (Figure 4(I)). Similar patterns were observed for the derivative of

mean recruitment to the mean phenotype (not shown).

These temporal changes in the derivatives of the mean fitness components to the phenotypic trait

value resulted in approximately constant survival and recruitment rates in the density-

independent survival model, weak decreases in survival rates and weak increases in recruitment

rates in the baseline, density-independent development and density-independent inheritance

models, and stronger temporal trends for decreased survival and increased recruitment rates in

the density-independent recruitment model (Figure 4(J)). In other words, the patterns we report

in mean fitness were caused by opposing trends in fitness components in these simulations.

The dynamics of β_1 that generated these contrasting patterns across the different simulations

were determined by the shape of the fitness function (Figure 2(C)). As density increased, the

intercept of the logistic fitness functions became smaller due to the density-dependence terms in

the models. In contrast, as mean body size increased, the body size distribution moved to the right

on the x-axis of figure 2(C). To some extent, these two processes tended to cancel one another

out, such that mean survival and mean recruitment tended to change relatively slowly. The one

exception to this pattern is in the density-independent recruitment model, where increasing body

size increased recruitment rates but did not lower the intercept of the function. Instead, mean

survival rates decreased as density increased because mean fitness evolved.

To sum up, our results show that the Soay sheep population is primarily regulated via density-

dependent survival. When density was removed from the survival function, the population

consequently increased to a large size, in turn reducing recruitment rates, development rates, and

the mean value of the environmental component of the phenotype. Large individuals, however,

had very high survival, at a point when the derivative of mean survival to body weight was very

shallow – i.e. β_1 was close to zero and evolution of both the mean phenotypic trait and mean

fitness was slow.

DISCUSSION

The structured, evolutionarily-explicit models we developed to examine the dynamics of terms in

the Breeder's and Price equations build on existing approaches by Lande (Lande 1982), Barfield et

al. (Barfield et al. 2011), Childs et al.(Childs et al. 2016), Coulson et al. (Coulson et al. 2011,

Coulson et al. 2017) and Rees and Ellner (Rees and Ellner 2019). The approach is powerful as it

allows the simultaneous study of population dynamics and the evolution of phenotypic traits, life

histories, and mean fitness (Coulson et al. 2011). In this paper we use dynamic models to generate

dynamics of components of the Breeder's and Price equations.

Population dynamics and phenotypic evolution are both determined by three drivers: i) the

distribution of phenotypic trait values, ii) the fitness functions that determine how expected

survival and reproductive rates vary with phenotypic trait values and the environment, and iii)

details of the genotype-phenotype map (Lande 1982, Lynch and Walsh 1998). The latter can

include phenotypic plasticity, and genetic and non-genetic inheritance (in all its forms). Our

models reveal how rates of evolution (defined as $\Delta\bar{A}_t$), of phenotypic change, and of change in the

population growth rate, are determined by factors that influence each of these three drivers.

In the absence of any processes that alter these drivers with time, genetic change, phenotypic

change, and population growth rate change will occur simultaneously (Witting 2002). In the case

of a linear association, the rate of change of each will slow with time. This is an example of eco-

evolutionary feedbacks. If the association between the phenotypic trait and fitness is exponential,

then the rate of change is constant. This is because the slope of the association between the

phenotypic trait and fitness changes with time as the mean phenotype evolves.

In natural systems, because populations cannot increase in size indefinitely, the association

between the phenotypic trait and fitness must be influenced by processes such as density-

dependence or environmental change, to prevent the population growing hyper-exponentially.

These processes are typically modelled as being additive (Ellner et al. 2016), and act to reduce the

intercept of the association between the phenotypic trait and fitness, as evolution increases the

mean value of the phenotypic trait (Smallegange and Coulson 2013). However, these processes

can also interact with the phenotypic trait to alter the slope of the association. In our models, the

survival and fertility (fitness) functions can capture all these processes (Coulson et al. 2017).

Environmental drivers such as density-dependence and climatic variation can influence the rate of

evolution, phenotypic change, and the population growth rate, by directly impacting the

phenotypic trait via processes other than selection – e.g. phenotypic plasticity and non-genetic

inheritance (Nussey et al. 2007, Reed et al. 2011). These processes can affect the rate of evolution,

phenotypic trait dynamics, and the ecology of a system. In particular, these drivers can alter the

mean or variance of the trait, by changing the environmental component of the trait (Coulson et

al. 2017). In our models, these effects are captured by the development and inheritance functions

– where the latter also captures genetic inheritance.

In natural systems, environmental drivers typically impact fitness, development and inheritance

functions simultaneously, and potentially in different ways in different ages and sexes (Simmonds

et al. 2019). A consequence of this is that the environment can simultaneously influence evolution,

phenotypic trait dynamics, and the population growth rate (e.g. ecology) in multiple, interacting

ways (Coulson et al. 2011). For example, density-dependence can alter the association between a

phenotypic trait and fitness, and alter the mean and variance of the environmental component of

the phenotype.

The Breeder's equation was developed as a model to predict evolution over a single time-step in

artificial environments when humans specify the association between the phenotypic trait and

fitness to produce a desired number of offspring from those individuals they allow to reproduce

(Lush 1943, Falconer 1960, Lynch and Walsh 1998). In addition, they minimise the effect of

environmental drivers on the phenotypic trait by typically providing *ad libitum* food and removing

predation- and disease-related causes of death. Under these circumstances, the Breeder's

equation performs well. Yet, when it has been used as a retrospective tool to explain evolution in

natural systems it has typically failed (Merilä et al. 2001). This is because processes that can alter

the rate of evolution in natural settings are not included in the model, and are treated as nuisance

variables to be corrected for in statistical analyses (Kruuk 2004). However, these processes can

influence evolution (and ecology) and should consequently be included in predictive models if

they are found to explain significant variation in statistical analyses used to estimate additive

genetic variances and the strength of selection. Our approach allows for these processes to be

explicitly incorporated into predictive models (Childs et al. 2016, Coulson et al. 2017, Simmonds et

al. 2019).

The Breeder's equation does not incorporate all the processes that can generate phenotypic

change (Morrissey et al. 2010). This observation limits the utility of estimates of additive genetic

variances and heritabilities from free-living populations (Kruuk 2004), because, if these processes

are important, they should be incorporated into predictive models. The Price equation, on the

other hand, does provide an exact description of phenotypic evolution (Price 1970). However, it is

a tautology (Heywood 2005), and simply describes the difference between two means. It is

consequently a very useful tool to retrospectively decompose phenotypic change into all factors

that contributed to it (Coulson and Tuljapurkar 2008), but it is not dynamically sufficient, and

consequently cannot be used to make predictions of future dynamics. The structured population

models we develop are dynamically sufficient, and provide a dynamic form of the Price equation

(Childs et al. 2004, Coulson et al. 2010). In previous work, we have demonstrated how these

models can be parameterised for real systems (Simmonds et al. 2019).

The class of structured models we have developed to predict evolutionary change in natural

systems is widely used in ecology (Caswell 2001). A consequence of this is that our models can be

used to predict simultaneous ecological and evolutionary change (Coulson et al. 2011). Eco-

evolution is frequently described as ecological change generating evolutionary change, which in

turn generates ecological change, and so on *ad infinitum* (Hendry 2016). The logic behind this

interpretation is that ecology and evolution are two separate processes that can influence one

another (Pelletier et al. 2009). Our work challenges this interpretation, as we show that ecological

change and evolution occur simultaneously, as a consequence of ecological change or evolution

altering fitness, development and inheritance functions (see also (Coulson et al. 2011, Childs et al.

2016, Coulson et al. 2017)).

Our approach is based on the logic that a population can be represented by a multivariate

distribution of genotypes and phenotypes (Easterling et al. 2000, Ellner and Rees 2006). The

dynamics of this distribution are then determined by survival and reproduction, development and

inheritance functions. We consider these four processes as fundamental. The dynamics of the

distribution's volume (size) are the population dynamics, while the dynamics of the mean

genotype (and sometimes phenotypic trait) capture evolution (Smallegange and Coulson 2013).

These models can also be used to study the dynamics of life history descriptors such as generation

length and lifetime reproductive success (Ellner and Rees 2006, Steiner et al. 2012, Steiner et al.

2014), and to examine life history evolution (Childs et al. 2003, 2004). The fundamental processes

that these models capture can vary as a function of age, sex, location, and genotype, and they can

include aspects of the biotic and abiotic environment such as the size and structure of the focal

species or of other species in the community, climatic variation, or variation in resource

availability (Ellner and Rees 2006, Coulson 2012, Schindler et al. 2015, Ellner et al. 2016, Bassar et

al. 2017). These models consequently offer a flexible framework to explore how populations will

ecologically and evolutionarily respond to environmental change.

In this paper we provide two simple models to demonstrate the utility of our approach. However,

more complex models can be developed by extending such models. For example, we can relax the

assumption of an additive genotype-phenotype map, and extend models for multiple traits, age-

and sex-structure (Schindler et al. 2015, Coulson et al. 2017). It is also possible to construct models

of interacting species (Bassar et al. 2017). Although there will doubtless be challenges in

parameterising some versions of these models, the approach we describe here offers enormous

potential to improve predictions of eco-evolution in natural settings.

ACKNOWLEDGEMENTS

Tom Potter is joint funded by NERC DTP and Lamb and Flag studentships at Oxford University, and

Anja Felmy is funded by an Early Postdoc Mobility Fellowship from the Swiss National Science

Foundation (P2EZP3_181775). We thank Dylan Childs and Joe Travis for helpful comments on an

earlier version of the manuscript.

Barfield, M., R. D. Holt, and R. Gomulkiewicz. 2011. Evolution in stage-structured populations. The
American Naturalist **177**:397-409.
Bassar, R. D., J. Travis, and T. Coulson. 2017. Predicting coexistence in species with continuous
ontogenetic niche shifts and competitive asymmetry. Ecology **98**:2823-2836.
Bonnet, T., P. Wandeler, G. Camenisch, and E. Postma. 2017. Bigger is fitter? Quantitative genetic
decomposition of selection reveals an adaptive evolutionary decline of body mass in a wild
rodent population. PLoS biology **15**:e1002592.
Caswell, H. 2001. Matrix population models: Construction, Analysis and Interpretation. Sinauer
Associates.
Childs, D. Z., M. Rees, K. E. Rose, P. J. Grubb, and S. P. Ellner. 2003. Evolution of complex flowering
strategies: an age–and size–structured integral projection model. Proceedings of the Royal
Society of London. Series B: Biological Sciences **270**:1829-1838.
Childs, D. Z., M. Rees, K. E. Rose, P. J. Grubb, and S. P. Ellner. 2004. Evolution of size–dependent
flowering in a variable environment: construction and analysis of a stochastic integral
projection model. Proceedings of the Royal Society of London. Series B: Biological Sciences
271:425-434.
Childs, D. Z., B. C. Sheldon, and M. Rees. 2016. The evolution of labile traits in sex-and age-
structured populations. Journal of Animal Ecology **85**:329-342.
Coulson, T. 2012. Integral projections models, their construction and use in posing hypotheses in
ecology. Oikos **121**:1337-1350.
Coulson, T., B. E. Kendall, J. Barthold, F. Plard, S. Schindler, A. Ozgul, and J.-M. Gaillard. 2017.
Modeling adaptive and nonadaptive responses of populations to environmental change.
The American Naturalist **190**:313-336.
Coulson, T., D. R. MacNulty, D. R. Stahler, R. K. Wayne, and D. W. Smith. 2011. Modeling effects of
environmental change on wolf population dynamics, trait evolution, and life history.
Science **334**:1275-1278.
Coulson, T., S. Schindler, L. Traill, and B. E. Kendall. 2018. Predicting the evolutionary
consequences of trophy hunting on a quantitative trait. The Journal of Wildlife
Management **82**:46-56.
Coulson, T., and S. Tuljapurkar. 2008. The dynamics of a quantitative trait in an age-structured
population living in a variable environment. The American Naturalist **172**:599-612.
Coulson, T., S. Tuljapurkar, and D. Z. Childs. 2010. Using evolutionary demography to link life
history theory, quantitative genetics and population ecology. Journal of Animal Ecology
79:1226-1240.
Crow, J. F., and M. Kimura. 1970. An introduction to population genetics theory. An introduction
to population genetics theory.
Easterling, M. R., S. P. Ellner, and P. M. Dixon. 2000. Size-specific sensitivity: applying a new
structured population model. Ecology **81**:694-708.
Ellner, S. P., D. Z. Childs, and M. Rees. 2016. Data-driven modelling of structured populations.
Springer.
Ellner, S. P., and M. Rees. 2006. Integral projection models for species with complex demography.
The American Naturalist **167**:410-428.
Falconer, D. S. 1960. Introduction to quantitative genetics. Oliver & Boyd.
Fisher, R. A. 1930. The genetical theory of natural selection. Clarendon Press.
Hairston Jr, N. G., S. P. Ellner, M. A. Geber, T. Yoshida, and J. A. Fox. 2005. Rapid evolution and the
convergence of ecological and evolutionary time. Ecology letters **8**:1114-1127.
Hendry, A. P. 2016. Eco-evolutionary dynamics. Princeton university press.
Heywood, J. S. 2005. An exact form of the breeder's equation for the evolution of a quantitative
trait under natural selection. Evolution **59**:2287-2298.

Kruuk, L. E. B. 2004. Estimating genetic parameters in natural populations using the 'animal
526   model'. Philosophical Transactions of the Royal Society of London. Series B: Biological
527   Sciences **359**:873-890.
528   Kruuk, L. E. B., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, and T. Clutton-Brock. 2002.
529   Antler size in red deer: heritability and selection but no evolution. Evolution **56**:1683-1695.
530   Lande, R. 1982. A quantitative genetic theory of life history evolution. Ecology **63**:607-615.
531   Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.
532   Evolution **37**:1210-1226.
533   Lush, J. L. 1943. Animal breeding plans. Iowa State College Press.
534   Lynch, M., and B. Walsh. 1998. Genetics and analysis of quantitative traits. Sinauer Sunderland,
535   MA.
536   Merilä, J., B. Sheldon, and L. Kruuk. 2001. Explaining stasis: microevolutionary studies in natural
537   populations. Genetica **112**:199-222.
538   Morrissey, M., L. Kruuk, and A. J. Wilson. 2010. The danger of applying the breeder's equation in
539   observational studies of natural populations. Journal of evolutionary biology **23**:2277-2288.
540   Nussey, D., A. Wilson, and J. Brommer. 2007. The evolutionary ecology of individual phenotypic
541   plasticity in wild populations. Journal of evolutionary biology **20**:831-844.
542   Patel, S., M. H. Cortez, and S. J. Schreiber. 2018. Partitioning the effects of eco-evolutionary
543   feedbacks on community stability. The American Naturalist **191**:381-394.
544   Pelletier, F., D. Garant, and A. Hendry. 2009. Eco-evolutionary dynamics. Philosophical
545   Transactions of the Royal Society B: Biological Sciences **364**:1483-1489.
546   Price, G. R. 1970. Selection and covariance. Nature **227**:520-521.
547   Reed, T. E., D. E. Schindler, and R. S. Waples. 2011. Interacting effects of phenotypic plasticity and
548   evolution on population persistence in a changing climate. Conservation Biology **25**:56-63.
549   Rees, M., and S. P. Ellner. 2019. Why so variable: can genetic variance in flowering thresholds be
550   maintained by fluctuating selection? The American Naturalist **194**:E13-E29.
551   Schindler, S., J. M. Gaillard, A. Grüning, P. Neuhaus, L. W. Traill, S. Tuljapurkar, and T. Coulson.
552   2015. Sex-specific demography and generalization of the Trivers–Willard theory. Nature
553   **526**:249.
554   Simmonds, E. G., E. F. Cole, B. C. Sheldon, and T. Coulson. 2019. Testing the effect of quantitative
555   genetic inheritance in structured models on projections of population dynamics. Oikos.
556   Smallegange, I. M., and T. Coulson. 2013. Towards a general, population-level understanding of
557   eco-evolutionary change. Trends in ecology & evolution **28**:143-148.
558   Steiner, U. K., S. Tuljapurkar, and T. Coulson. 2014. Generation time, net reproductive rate, and
559   growth in stage-age-structured populations. The American Naturalist **183**:771-783.
560   Steiner, U. K., S. Tuljapurkar, T. Coulson, and C. Horvitz. 2012. Trading stages: life expectancies in
561   structured populations. Experimental gerontology **47**:773-781.
562   Wallace, K., A. Leslie, and T. Coulson. 2013. Re-evaluating the effect of harvesting regimes on N ile
563   crocodiles using an integral projection model. Journal of Animal Ecology **82**:155-165.
564   Witting, L. 2002. Evolutionary dynamics of exploited populations selected by density dependent
565   competitive interactions. Ecological Modelling **157**:51-68.
566

Figure 1. Evolution of (A) the mean breeding value, (B) the mean phenotypic trait, (C) mean

fitness, and (D) the additive genetic variance in simple models over 50 generations. The black line

represents sim 1), where the additive genetic variance is eroded and the mean and variance of the

environmental component of the phenotypic trait remain constant with time. The red lines

represent sim 2), where additive genetic variance is not eroded by selection, and the mean and

variance of the environmental component of the phenotypic trait remain constant with time. The

blue (sim 3)) and green line (sim 4)) are the same as sim 1) and sim 2), except we impose a weak

temporal downward trend on the mean environmental component of the phenotypic trait. (E) In

models with no temporal trend in the environmental component of the phenotypic trait, the

relationship between the mean breeding value and mean fitness is linear, while it is slightly

curvilinear in models with temporal trends in E. In contrast, there is (F) a linear association

between the mean phenotypic trait and mean fitness in all four models.

Figure 2. The dynamics of phenotypic traits and mean fitness. (A) Because the breeding value and

the value of the environmental component of a phenotypic trait sum to produce the phenotypic

trait, there are multiple ways the same phenotypic trait value can be produced (dashed grey lines).

If the phenotypic trait linearly influences fitness, these phenotypic trait clines also represent

fitness clines, generating linear fitness contours. Larger values of the trait, and of mean fitness, are

depicted with darker dashed lines. Depending upon the dynamics of the environmental

component of the phenotype, selection can fail to climb across contours, or can climb at a range

of rates even for the same, constant, derivative of mean fitness to the mean phenotype (i.e.

phenotypic slope in the linear fitness function). Because mean fitness remains lowest when

evolution is cryptic, all other things being equal, selection will also be strongest, and evolution

fastest. Cryptic evolution occurs when non-genetic inheritance or phenotypic plasticity impact the

environmental component of the phenotype in a manner that exactly counters gains to the

additive genetic component of the phenotype attributable to selection. (B) The bivariate dynamics

of the mean breeding value and the mean environmental component of the phenotypic trait over

time in each simulation for the modified sheep model. The starting point of each simulation is

towards the bottom of the figure. The increasing width of the thick lines represent how population

size (as a consequence of the dynamics of mean fitness) changes as the simulation progresses.

Bivariate distributions of the components of the phenotypic trait are drawn as contour plots at

time $t=1$, $t=50$ and $t=100$ for the simulation where density-dependence was removed from the

recruitment (fitness) function. Evolution is captured by the length of the thick line in the y-axis,

while change in the environmental component of the phenotypic trait is captured by change in the

x-axis. (C) The derivative of mean fitness to the mean phenotype (red line) can vary as either the

mean phenotypic trait increases when fitness functions are non-linear (blue dots), or when

environmental drivers such as increasing population density lower the intercept of the fitness

function (dashed black lines).

Figure 3. Evolution of (A) the mean breeding value, (B) the mean phenotypic trait value, (C) mean

fitness, and (D) the additive genetic variance in the modified sheep model. The baseline model

(black lines) has negative density-dependence in each function, while the coloured lines represent

simulations where density-dependence has been removed from one particular function.

Figure 4. Dynamics of the phenotypic trait distribution in the baseline model (A) and in models

with negative density-dependence removed from each function (B-E). Distributions are drawn at

times $t=1$, $t=50$ and $t=100$. These dynamics are determined in part by selection (the derivative of

mean fitness to the mean phenotypic trait value) that consists of (F, G) covariances between the

trait and components of fitness, and (H) the phenotypic variance in the trait. (I) derivative of mean

survival to the mean phenotype (similar patterns for recruitment not shown) and (J) the temporal

dynamics of mean survival and recruitment rates.

618 Figure 1

Figure 2.

Figure 3.

Figure 4.

Appendix 1

A key premise of the standard quantitative genetic approach is that the genotype-phenotype map

can be described as $z = A + E$ (Falconer 1960). If a phenotypic trait is measured in a particular

unit, say length in millimetres, then A and E also take these units. The second premise is that

selection operates on the entire phenotype via a selection differential. If we consider a linear

fitness function $E(w) = \beta_0 + \beta_1 z$ we can also write $E(w) = \beta_0 + \beta_1(A + E) = \beta_0 + \beta_1 A + \beta_1 E$.

This equation says the effect of a one-unit change on fitness in the additive genetic and

environmental components of the phenotype is identical.

A linear regression slope is defined as the covariance between the response and predictor

variable, divided by the variance in the predictor variable. Consequently,

$$\bar{w}_t = \beta_0 + \frac{cov(w,A)}{var(A)}A + \frac{cov(w,E)}{var(E)}E,$$

where $\frac{cov(w,A)}{var(A)} = \frac{cov(w,E)}{var(E)}$. If $\frac{cov(w,A)}{var(A)} \neq \frac{cov(w,E)}{var(E)}$ then $z \neq A + E$. Instead, the genotype-phenotype

map will be of the form $z = A + \gamma E$, where $\gamma \neq 1$. In general, this means that any difference in

slopes of the breeding values and the environment components of the phenotype with fitness can

only occur if the genotype-phenotype map has changed with time during the period over which

data were collected to parameterise models.

Some authors have argued that selection can operate in contrasting ways on the environmental

and additive genetic components of the phenotype while assuming $z_t = A_t + E_t$ (Morrissey et al.

2010). In the most extreme cases they argue that selection on the breeding value is in one

direction, while selection on the environmental component of the phenotype is in a contrasting

direction such that $\gamma < 0$ while A_t and E_t do not covary (Bonnet et al. 2017). Although

mathematically possible, such a genotype-phenotype map seems biologically implausible. We

have been unable to identify a biologically plausible genotype-phenotype map where a one-unit

increase in A_t will result in an increase in fitness, while a one-unit increase in E_t will result in a

decrease, without invoking some dependence of A_t and E_t on one another.

A more plausible genotype-phenotype map would involve phenotypic plasticity, where a breeding

value A_t produces a different phenotypic value z_t in different environments. If $z_t = A_t + E_t$ this

requires E_t to vary with the environment θ, such that E can be made a function of the

environment $E(\theta_t)$. Phenotypic plasticity can then be captured with a number of formulations

such as $z_t = A_t + E(\theta_t)$ or $z_t = A_t + E(\theta_t) + \alpha A_t E(\theta_t)$ (Coulson et al. 2017). In the first

formulation, the mean and variance of E_t can vary with the environment, but the genotype-

phenotype map remains $z_t = A_t + E_t$ such that the slope between the environmental component

of the phenotype and fitness equals the slope between the additive genetic component of the

phenotype and fitness, but $\bar{E}_t$ change with time, altering the numerical form of the genotype-

phenotype map. In contrast, in the second formulation, the breeding value-by-environment

interaction means that the slope between fitness and each component of the phenotype differs.

Empirical evidence of so-called "selection on the environmental component of the phenotype"

(Bonnet et al. 2017) is consequently likely evidence of a genotype-by-environment interaction:

something that cannot be detected with widely used statistical analyses of observational data

(Kruuk 2004).

Appendix 2

Four approaches can be used to specify how inheritance alters the distribution of breeding values

between selected parents and their offspring. Approach 1: A convolution of the distribution of

breeding values in mothers and fathers is taken to construct a distribution of mid-parental values

assuming random mating. The distribution of mid-parental values is then convolved with a normal

distribution with a mean of 0 and a constant variance to construct the distribution of breeding

values in offspring. Approach 2: Breeding values in the offspring generation are assumed to be

normally distributed, with a mean equal to that of the mid-point parental breeding value

distribution, and a constant variance that does not change across generations. Approach 3: The

distribution of breeding values of parents is operated on by an inheritance function which

maintains the mid-parental mean value and injects Gaussian distributed segregation variance

around each mid-point breeding value. Approach 4: The distribution of breeding values in parents

and offspring is identical. As long as selection does not generate distributions with substantial

skew or kurtosis, the first three of these approaches give very similar predictions. Various

properties of these four approaches are summarised in Table A2.1

Table A2.1. Properties of transition functions for breeding values.

	Approach 1	Approach 2	Approach 3	Approach 4
Are the breeding values in offspring normally distributed?	not necessarily	yes	not necessarily	not necessarily

Does the distribution of breeding values in mid-parents and offspring have the same mean?	yes	yes	yes	yes
Does the distribution of breeding values in parents and offspring have the same shape?	no	no	no	yes

687   **Appendix 3**

688   The Soay sheep model consisted of the following parameterisations:

689
$$S(A_t + E_t) = \frac{1}{1 + e^{-(-0.2392 + 1.1175(A_t + E_t) - 0.0043N_t)}}$$

690
$$R(A_t + E_t) = \frac{1}{1 + e^{-(-1.7839 + 0.1090(A_t + E_t) - 0.0032N_t)}}$$

691
$$G(A'_{t+1}|A_t) = A_t$$

692
$$G(E'_{t+1}|E_t) = norm(\mu_G(E_t), \sigma_G^2(E_t)),$$

693   where $\mu_G(E_t) = 10.603 + 0.6049E_t - 0.0026N_t$ and $\sigma_G^2(E_t) = 2.874 - 0.0124E_t$

694
$$D(A'_{t+1}|A_t) = norm(\mu_D^\omega(A_t), \sigma_D^2(A_t)),$$

695   where $\mu_D^\omega(A_t)$ is the mean breeding value post-fertility selection, and $\sigma_D^2(A_t)$ is the additive

696   genetic variance in offspring at birth, which we assumed to be constant and Gaussian distributed.

697   Finally,

698
$$D(E'_{t+1}|E_t) = norm(\mu_D(E_t), \sigma_D^2(E_t)),$$

where $\mu_D(E_t) = 9.8065 + 0.2544E_t - 0.0069N_t$ and $, \sigma_D^2(E_t) = 4.581 - 0.1196E_t$.

We started each simulation with mean values of $\bar{A}_t = 5.5$ and $\bar{E}_t = 19.5$, an additive genetic

variance of 4 and a variance in the environmental component of 6. Because the genetic variances

influence the rate of evolution, and selection operate on the entire phenotype, the model results

are insensitive to the starting values of $\bar{A}_t$ and $\bar{E}_t$. Rather it is their sum that determines dynamics.

In an initial (unreported) model we tracked a multivariate distribution of age-related breeding

values with genetic correlations close to unity. We simplified the model to so each age-related

trait has the same breeding value – i.e. genetic correlations are equal to unity. The simplification

made no material difference to the results we report.