The role of the diencephalon in the guidance of thalamocortical

axons in mice

Idoia Quintana-Urzainqui[1], Pablo Hernández-Malmierca P[2,3], James M. Clegg[1],

Ziwen Li[4], Zrinko Kozić[1], David J Price[1]

[1] Centre for Discovery Brain Sciences, Hugh Robson Building, George Square,

Edinburgh EH8 9XD, United Kingdom

[2] Heidelberg Institute for Stem Cell Technology and Experimental Medicine (HI-

STEM gGmbH), 69120 Heidelberg, Germany

[3] Division of Stem Cells and Cancer, Deutsches Krebsforschungszentrum (DKFZ)

and DKFZ-ZMBH Alliance, 69120 Heidelberg, Germany

[4] Centre for Cardiovascular Science, Queen's Medical Research Institute, 47 Little

France Crescent, Edinburgh EH16 4TJ, United Kingdom

Corresponding author address: idoia.quintana@ed.ac.uk

Abstract

Thalamocortical axons (TCAs) cross several tissues on their journey to the cortex.

Mechanisms must be in place along the route to ensure they connect with their

targets in an orderly fashion. The ventral telencephalon acts as an instructive tissue,

but the importance of the diencephalon in TCA mapping is unknown. We report that

disruption of diencephalic development by Pax6 deletion results in a thalamocortical

projection containing mapping errors. We used conditional mutagenesis to test

whether these errors are due to the disruption of pioneer projections from

prethalamus to thalamus and found that, while this correlates with abnormal TCA

fasciculation, it does not induce topographical errors. To test whether the thalamus

contains navigational cues for TCAs, we used slice culture transplants and gene

expression studies. We found the thalamic environment is instructive for TCA

navigation and that the molecular cues Netrin1 and Semaphorin3a are likely to be

involved. Our findings indicate that the correct topographic mapping of TCAs onto

the cortex requires the order to be established from the earliest stages of their

growth by molecular cues in the thalamus itself.

Introduction

A striking feature of the axonal tracts that interlink the nervous system's component parts is the high degree of order with which they map the array of neurons in one structure onto the array of neurons in their target. Often, the order of axons at the target closely mirrors that at the source. An excellent example is the mapping of thalamic neurons onto their cerebral cortical targets via the thalamocortical pathway (Fig. 1A). Thalamic neurons located at one end of the thalamus in a dorsolateral region called the dorsal lateral geniculate nucleus (dLGN) innervate the caudal (visual) part of cortex; neurons located at the other end of the thalamus in more rostral-medial regions known as the ventrolateral (VL) and ventromedial (VM) nuclei innervate more rostral cortical regions, including motor and frontal cortex; neurons located in between - in the ventromedial posterior (VMP) nuclei - innervate central (somatosensory) cortex (Fig. 1A) (Amassian and Weiner, 1966; Bosch-Bouju et al., 2013; Jones, 2007; Tlamsa and Brumberg, 2010). The mechanisms that generate this orderly topographic mapping remain poorly understood.

The maintenance of order among thalamic axons as they grow is likely to contribute to the generation of orderly topographic mapping in the mature thalamocortical tract. During embryogenesis, thalamic axons exit the thalamus from about E12.5 onwards (Auladell and Hans, 2000; Braisted et al., 1999; Tuttle et al., 1999), approximately coincident with the cessation of neurogenesis in this structure (Angevine, 1970; Li et al., 2018). They then cross the adjacent prethalamus and turn laterally out of the diencephalon and into the ventral telencephalon where they traverse two consecutive instructive regions - the corridor (Lopez-Bendito et al., 2006) and the striatum - before entering the cortex. There is evidence that the maintenance of spatial order among thalamocortical axons (TCAs) crossing the ventral

telencephalon requires interactions between the axons and signals released by cells they encounter in this region (Bielle et al., 2011; Bonnin et al., 2007; Braisted et al., 1999; Dufour et al., 2003; Molnár et al., 2012; Powell et al., 2008). The importance of earlier interactions within the diencephalon remains unclear.

Here, we tested the effects of mutating the gene for the Pax6 transcription factor, which is essential for normal diencephalic patterning (Caballero et al., 2014; Clegg et al., 2015; Parish et al., 2016; Pratt et al., 2000; Stoykova et al., 1996; Warren and Price, 1997), on the topographic mapping of TCAs onto the cortex. Pax6 starts to be expressed in the anterior neural plate well before TCAs start to form (Walther and Gruss, 1991). As the forebrain develops from the anterior neural plate, Pax6 expression becomes localized in (i) cortical progenitors that generate the target neurons for TCAs, (ii) diencephalic (thalamic and prethalamic) progenitors and (iii) prethalamic (but not thalamic) postmitotic neurons (Quintana-Urzainqui et al., 2018; Stoykova et al., 1996; Warren and Price, 1997). We discovered that deletion of Pax6 from mouse embryos at the time when thalamic axons are starting to grow results in the development of a thalamocortical projection containing mapping errors. Axons from dorsolateral thalamus are misrouted medially and end up projecting abnormally rostrally in the cortex. We went on to explore the reasons for this defect.

We first used conditional mutagenesis to test whether misrouting is due to the loss of Pax6 from prethalamic neurons, since previous work has shown that (i) Pax6 is not required in the cortex for normal TCA topography (Piñon et al., 2008) and (ii) Pax6 is neither expressed nor required autonomously by thalamic neurons for them to acquire the ability to extend axons to the cortex (Clegg et al., 2015). We found that while loss of Pax6 from a specific set of prethalamic neurons prevented them developing their normal axonal projections to thalamus and resulted in the abnormal

fasciculation of thalamic axons, it did not cause TCAs to misroute. This suggested

that the thalamus itself contains important navigational cues for TCAs. We used slice

culture transplants and gene expression studies to show (i) that the thalamic

environment is indeed instructive for TCA navigation and (ii) to identify molecular

changes within the thalamus that likely cause the disruption in TCA topography

observed upon Pax6 deletion. Our findings indicate that the normal topographic

mapping of TCAs requires that order be established and maintained from the earliest

stages of their growth by molecular cues in the thalamus itself.

Results

Thalamocortical topography is disrupted in CAG^{CreER} but not in Emx^{CreER} Pax6

conditional knockouts

Previous studies have shown that constitutive loss of Pax6 function causes a total

failure of TCA development, which is hypothesized to be a secondary consequence

of anatomical disruption at the interface between the diencephalon and the

telencephalon (Clegg et al., 2015; Georgala et al., 2011; Jones et al., 2002). No such

failure occurs if Pax6 is deleted conditionally after this anatomical link is formed

(Clegg et al., 2015). We first assessed whether delayed ubiquitous Pax6 deletion,

induced in $CAG^{CreER-TM} Pax6^{fl/fl}$ embryos (referred to here as CAG^{CreER} Pax6 cKOs),

disrupts the topography of TCA connections. We induced Cre recombinase

activation by tamoxifen administration at E9.5 which caused Pax6 protein loss in

CAG^{CreER} Pax6 cKOs from E11.5 onwards (Quintana-Urzainqui et al., 2018), which is

when the generation of most thalamic neurons is starting (Li et al., 2018) and before

many TCAs have begun to grow (Auladell and Hans, 2000; López-Bendito and

Molnár, 2003). Diencephalic progenitor domains in CAG^{CreER} Pax6 cKOs are fully

recognizable (Quintana-Urzainqui et al., 2018) and patterning seems largely

unaffected in the thalamus (Fig. S1). We used both wild type and $CAG^{CreER}Pax6^{fl/+}$

littermate embryos as controls since the latter express normal levels of Pax6 protein

(see Methods; Caballero et al., 2014; Manuel et al., 2015).

We inserted two different axonal tracers in two cortical areas in E15.5 fixed brains.

DiA was placed in the visual (caudal) cortex while DiI was placed in the

somatosensory (more rostral) cortex (Fig. 1B). In controls (both wild type and

$Pax6^{fl/+}$), DiA retrogradely labelled cells in dorsolateral thalamic areas (dLGN; green

labelling in Fig. 1C-E), while DiI labelled cells in ventromedially-located thalamic

regions, identified as ventral-posterior thalamic nucleus (VP) (red labelling in Fig. 1C-

E). Labelling of these two thalamic regions was clearly separated in all cases

(indicated by dotted line in Fig. 1C-E). In CAG^{CreER} Pax6 cKOs, however, the two

labelled populations overlapped (Fig. 1F-H). In these mutants, the distribution of the

DiA-labelled thalamic cells (from caudal cortical injections) was not obviously

changed with respect to controls. However, the DiI-labelled thalamic cells (projecting

to more rostral cortical areas) showed a much wider distribution than in controls and

expanded to lateral thalamic areas (compare Fig.1 C-E,C'-E' versus F-H,F'-H'), even

overlapping with DiA stained cells at the dLGN (Fig. 1 G,H,G',H'). To quantitate this,

we measured the area occupied by DiI and DiA within each nucleus of interest in

controls versus CAG^{CreER} Pax6 cKOs (in transverse E15.5 sections from three

different litters: four controls, three cKOs, five sections per embryo). We defined the

dLGN and VP nucleus as regions of interest (ROI) blind to DiI/DiA labelling and

measured the percentage of each ROI occupied by DiI and DiA (for details see

Methods). We found highly significant increase (from virtualy 0% to 7.7%) of the area

occupied by DiI in the dLGN in mutants compared to controls (Fig. 1I). We also

found a significant increase in DiA in VP nucleus, although its magnitude was very

small (0.26%) (Fig. 1I). Areas occupied by DiI in VP or DiA in dLGN were not

significantly altered. This analysis shows that *CAG^CreER Pax6* cKOs display

topographic errors, the main one being the misrouting of axons from dLGN neurons

towards abnormally rostral cortical areas in the *CAG^CreER Pax6* cKOs.

Since Pax6 is expressed both in the cortex and diencephalon during TCA

development, the mapping defects described above might have been due to the loss

of Pax6 from the cortex. This was unlikely because a previous study showed that

Pax6 is not required in the cortex for the establishment of proper topographical

thalamocortical connections (Piñon et al., 2008). To confirm this, we used a cortex-

specific, tamoxifen-inducible Cre line (*Emx1^CreER*). We administered tamoxifen at

E9.5, which results in a near-complete loss of cortical Pax6 between E11.5-12.5

(Georgala et al., 2011; Mi et al., 2013), and performed DiI/DiA labelling at E15.5,

following the same experimental design described above for the *CAG^Cre* line. We

found that the two retrogradely-labelled populations did not overlap in controls

(*Emx1^CreER Pax6^fl/+*) or in mutants (*Emx1^CreER Pax6^fl/fl*) (Fig. S2A,B), suggesting that

the defects of TCA mapping found in the *CAG^CreER Pax6* cKOs were not attributable

to cortical abnormalities.

To define the anatomical region where thalamic axons probably deviated from

ordered growth, we examined the TCA bundle in *CAG^CreER Pax6* cKOs. This bundle

was ordered and segregated into rostral/somatosensory (DiI) and caudal/visual (DiA)

halves at its point of exit from the prethalamus and entry into the ventral

telencephalon (arrows in Fig. 1 F-H), which indicated that the misrouting of lateral

TCAs from dorsolateral thalamus might happen before this point, i.e. within the

diencephalon.

TCAs fasciculate prematurely as they cross the prethalamus in *Pax6* conditional mutants

Within the diencephalon, the first structure that thalamic axons encounter as they leave the thalamus is the prethalamus, and its neurons normally express high levels of Pax6. Therefore, we investigated whether the defects of TCA mapping in *CAG^CreER Pax6* cKOs might arise from a disordered growth of thalamic axons through the prethalamus. As a first step, we examined the effects of Pax6 deletion on the behaviour of thalamic axons as they cross the prethalamus.

Since the neural cell adhesion molecule L1CAM (L1) is expressed in TCAs (Fukuda et al., 1997; Ohyama et al., 2004), we examined the distribution of L1-positive thalamic axons at E13.5 in transverse and sagittal sections through the prethalamus. In controls, axons emerging from the thalamus converge progressively as they cross the prethalamus (Fig. 2A-E) to subsequently form a single thalamocortical bundle that turns laterally and exits the prethalamus (arrows in Fig. 2A,B,E). We found that in *CAG^CreER Pax6* cKOs (Fig. 2F-L), thalamic axons prematurely converge into larger bundles as soon as they cross the thalamic-prethalamic boundary (empty arrows in Fig. 2G-I,L).

To obtain a quantitative measurement of this observation we positioned three equally-spaced lines across different diencephalic levels in sagittal sections (see Methods; lines represented in Fig. 2D,E,K,L). We used Fiji software (Schindelin et al., 2012) to quantify the number of axon bundles crossing each line and the width of each individual bundle. We recognized individual bundles as each single L1-positive structure above a consistent intensity threshold (red lines in Fig. 2M,N). We found a significant decrease in the number of bundles crossing all three checkpoints (Fig.

2O). Axon bundle width strikingly increased at the Th-PTh border and the mid-PTh,

with no significant change at the low-PTh checkpoint line (Fig. 2P) (see figure legend

for statistical details). These data indicate that, in the absence of Pax6, TCAs begin

to fasciculate prematurely in their route, forming bigger and fewer bundles as they

cross the prethalamus (Fig. 2Q). We next tested the role of the prethalamus in TCA

formation and the potential establishment of their topography.

Loss of prethalamic pioneer axons correlates with abnormal TCA fasciculation

but not with changes in topography

The prethalamus has been proposed to host a population of neurons that extend

axons to the thalamus which act as "pioneer guides" for TCA navigation (Price et al.,

2012; Tuttle et al., 1999). We assessed whether this population is disrupted by Pax6

loss from the prethalamus since, if it is, this might provide an explanation for

phenotypes described above.

From E9.5 on, most cells in the prethalamus express, or are derived from cells that

expressed, *Gsx2*. We used a *Gsx2^{Cre}* line (Kessaris et al., 2006) carrying an EGFP

Cre reporter (Sousa et al., 2009) to visualize neurons and axons belonging to the

Gsx2 lineage and we observed that prethalamic Pax6-expressing cells are included

within the location of the *Gsx2* lineage prethalamic population (Fig. 3A,B). *Zic4* is

also expressed by some prethalamic cells, with an onset of expression similar to that

of *Gsx2* (about E9.5; Gaston-massuet et al., 2005), and most diencephalic *Zic4*

lineage cells express and require Pax6 for their normal development (Li et al., 2018).

Using a *Zic4^{Cre}* line (Rubin et al., 2011) we observed that prethalamic neurons

derived from *Zic4* lineage were located in a narrow band close to the thalamic-

prethalamic border (Fig. 3C-D). We assessed whether these prethalamic populations

normally send axons to the thalamus.

Gsx2-lineage GFP-positive axons extended throughout the thalamus forming

ordered and parallel projections (Fig. 3E) and running in close apposition to L1-

positive TCAs (Fig. 3E',E'') from E12.5 onwards (Fig. S3). By contrast, *Zic4*-lineage

prethalamic cells did not project axons to the thalamus (Fig. 3D), indicating that

prethalamic pioneer axons arise from *Gsx2*-lineage and not from *Zic4*-lineage

prethalamic cells.

Since *Gsx2* is also expressed in the ventral telencephalon (Fig. 3A), and ventral

telencephalic neurons are known to project to the thalamus (López-Bendito and

Molnár, 2003; Métin and Godement, 1996; Molnár et al., 1998), there was a

possibility that *Gsx2*^{Cre} lineage axons innervating the thalamus actually originated

from ventral telencephalic neurons. To confirm the existence of *Gsx2*-lineage

prethalamic neurons projecting to the thalamus we injected the neuronal tracer

Neurobiotin^{TM} in the thalamus of E13.5 *Gsx2*^{Cre} embryos and successfully labelled

prethalamic neurons (arrow in Fig. 3F). Neurobiotin^{TM}-positive cells were GFP-

expressing *Gsx2* lineage (Fig. 3F-F'',G) and most of them also expressed Pax6

(arrows in Fig. 3H,H'; see summary in Fig. 3I). (Note that individual injections each

involved only subregions of the thalamus, explaining why each one only labelled a

discrete subset of the prethalamic neurons projecting to the thalamus). This

experiment confirmed that *Gsx2*-lineage cells in the prethalamus project to the

thalamus.

Having established that pioneer prethalamic axons belong to the *Gsx2* lineage and

express Pax6 we next aimed at disrupting their formation by conditionally deleting

Pax6 in *Gsx2*-lineage cells. We crossed mice carrying the floxed *Pax6* allele and

EGFP Cre reporter with the *Gsx2^Cre* line. Pax6 conditional deletion in *Gsx2*-lineage

cells (*Gsx2^Cre Pax6* cKOs) caused a visible reduction in the number of GFP-positive

axons projecting from prethalamus to thalamus in E12.5, E13.5 and E14.5 embryos

(Fig. 4A-F). To confirm that the prethalamic axons that were lost in *Gsx2^Cre*;

Pax6^loxP/loxP embryos were Pax6-expressing, we used the DTy54 YAC reporter allele

to express tauGFP in cells in which the *Pax6* gene is active, irrespective of whether it

is mutant or not (Tyas et al., 2006). Whereas there were many tauGFP-labelled

axons running from prethalamus to thalamus in controls, there were very few in

experimental embryos (Fig. 4G-J).

DiI placed in the thalamus of E13.5 *CAG^CreER* controls (*CAG^CreER Pax6 ^fl/+*)

retrogradely labelled a prethalamic population (arrow in Fig. 4K). In the absence of

Pax6 (*CAG^CreER Pax6* cKOs), no prethalamic cell bodies were labelled by thalamic

DiI injection (Fig. 4L), providing further evidence that the prethalamic pioneer

population does not form correctly when Pax6 is deleted. Overall, our results show

that prethalamic pioneer axons originating from *Gsx2*-lineage cells both express and

require Pax6 to develop normal connections with the thalamus (Fig. 4T).

We then studied the TCAs of *Gsx2^Cre Pax6* cKOs. Similar to the phenotype

described in *CAG^CreER Pax6* cKOs, Pax6 deletion in *Gsx2* lineage caused abnormal

premature fasciculation of axons crossing the thalamic-prethalamic border, as

evidenced by L1 immunohistochemistry (Fig. 4M-R). However, unlike *CAG^CreER Pax6*

cKOs, *Gsx2^Cre Pax6* cKOs did not show abnormal topographical projections, with no

obvious overlap between thalamic retrogradely-labelled populations after cortical DiA

and DiI placement in caudal and more rostral cortex respectively (Fig. 1B; Fig. 4S).

We conclude that, while prethalamic pioneer axons may play a role in avoiding

premature TCA fasciculation, they are not required for the establishment of accurate

thalamocortical topographic mapping (Fig. 4T).

Evidence for the importance of navigational cues in the thalamus itself

We next considered the potential importance of thalamic factors in the establishment

of thalamocortical topographic order. Our results above indicate that thalamic axons

might have deviated from their normal trajectories before they exited the thalamus in

CAG^CreER Pax6 cKOs (arrows mark deviant axons in Fig. 1 G',H'; no such axons

were observed in the controls, Fig. 1C',D',E'). A misrouting of TCAs in the thalamus

was also evident with L1 staining (Fig. 5 A,B). The largest collections of deviant

axons were observed projecting from lateral to medial thalamic regions (arrow in Fig.

5B), suggesting that the loss of Pax6 had disrupted normal navigational mechanisms

operating within the thalamus.

We looked for evidence that thalamic axons are actively guided through the normal

thalamus by using *in vitro* slice culture transplants to assess the effects of

repositioning lateral or medial thalamic neurons on the routes taken by their axons.

We grafted thalamic slice explants from E13.5 GFP-positive donor embryos into

GFP-negative host slices and cultured them for 72 hours to allow thalamic axons to

regrow and navigate through the host environment. The donor grafts were positioned

so that their axons had to traverse at least 200µm of host thalamic tissue before

encountering the Th-PTh boundary, allowing us to assess how the host thalamic

tissue affected the trajectory of the axons emerging from the donor tissue. We

isolated donor explants from either lateral or medial thalamus, and grafted them

either medially or laterally into host thalamus (Fig. 5C,E,G,I).

We found that axons from lateral thalamic explants showed a strong preference to

follow a lateral trajectory, while axons from medial explants extended much broadly

throughout the thalamus. In both cases, this effect was irrespective of whether they

were grafted laterally or medially (Fig. 5C-J). To quantitate this, we divided the

thalamus in three equal sectors (lateral, intermediate, medial; Fig 5K) and measured

the area occupied by GFP-positive elements in each of the sectors. We quantified

six explants (three lateral and three medial, 5 sections per explant) from three

different litters. In average, 95.7% of axons from lateral explants navigated through

lateral-most area (Fig 5.K), while medial explants extended axons more evenly

across the three regions (45.8% lateral, 31.8% intermediate and 22.4% medial; Fig.

5K). Moreover, we observed that when lateral axons were confronted with medial

host thalamus, most made a sharp turn towards lateral positions before heading

towards the prethalamus (Fig. 5E,F). When medial explants were grafted laterally

many of their axons turned medially (arrows in Fig. 5J).

These results indicated that different subsets of thalamic axons exhibit different

chemotactic responses to the thalamic environment and therefore that thalamic

axons are actively guided by mechanisms operating within the thalamus itself. To

gain further insight into what these mechanisms might be, we went on to examine

the expression of guidance molecules in the normal thalamus and in thalamus from

which Pax6 has been removed.

Axon guidance molecule expression in normal and Pax6 deficient thalamus

Semaphorin 3a (Sema3a) and Netrin 1 (Ntn1) are secreted guidance molecules

whose complementary gradients of expression in the ventral telencephalon are key

for the correct establishment of topographical connections between thalamus and

cortex (Bielle et al., 2011; Braisted et al., 1999; Molnár et al., 2012; Powell et al.,

2008; Wright et al., 2007). Interestingly, we found that their transcripts are also

distributed in opposing gradients in the thalamus (Fig. S4A,B). *Ntn1* is most highly

expressed at rostral-medial levels (Fig. S4A) while *Sema3a* is most highly expressed

in a more caudal-lateral aspect of the thalamus and in the lateral prethalamus (Fig.

S4B). In transverse *in situ* hybridization (ISH) of E13.5 controls we observed that

Ntn1 is expressed in a narrow rostral-medial thalamic population of neurons (arrow

in Fig. 6A) while *Sema3a* is expressed in caudal-lateral thalamic neurons (arrows in

Fig. 6B) as well as flanking the TCA bundles in the prethalamus (empty arrow in Fig.

6B).

To obtain a clearer three-dimensional view of these expression patterns we

reconstructed them from serial, adjacent sections stained for *Sema3a*, *Ntn1*, Pax6

and L1 in controls (Fig. 6C, see Methods). The 3D reconstruction confirmed that

Sema3a and *Ntn1* form opposing gradients in the normal embryonic thalamus (Fig.

6C), with *Sema3a* highest at caudal-lateral thalamic levels while *Ntn1* is highest at

rostral-medial thalamic levels.

We next investigated the expression patterns of the main receptors for *Ntn1* and

Sema3a in the thalamus of control embryos. The most interesting finding was that

Unc5c, encoding a Ntn1 receptor mediating axonal repulsion (Leonardo et al., 1997),

was expressed differentially from lateral to medial across the thalamus (Fig. 6D).

Laterally, almost all cells expressed high levels of *Unc5c* whereas medially many

cells did not (Fig. 6D'). *Unc5c* was largely absent from a narrow strip of cells close to

and parallel with the ventricular zone. This strip coincided with the region that

contained *Ntn1*-positive cells (compare Fig. 6D and A). *Plxna1*, encoding a Sema3a

receptor that mediates repulsion (Rohm et al., 2000; Takahashi et al., 1999;

Tamagnone et al., 1999) was found to be distributed relatively homogenously across

the thalamus (Fig. 6E, S4C).

These expression patterns suggest that, whereas all thalamic axons might be

repelled by Sema3a (due to their expression of *Plxna1*), only some axons might be

repelled by Ntn1 (i.e. those originating laterally, which express *Unc5c*, and those

Unc5c-expressing axons that originate medially) (Fig. 6K). This could explain why, in

the grafting experiments described above, axons from lateral explants invariably

navigated laterally, which would be away from medially located high levels of Ntn1. It

could also explain why medial explants generated axons able to navigate on a

broader front: some axons (those that express *Unc5c*) would be pushed relatively

laterally by repulsion from medially expressed *Ntn1*; others (those that do not

express *Unc5c*) would be able to maintain a medial trajectory through *Ntn1*-

expressing territory, thereby avoiding the high levels of *Sema3a* expressed in lateral

thalamus (Fig. 6K).

Other receptor-coding genes analysed (*Dcc*, *Unc5a*, *Unc5d*) showed little or no

expression within the main body of the thalamus and are therefore unlikely to

contribute to the navigation of thalamic axons within the thalamus (Fig. S4E,G,I).

We next asked whether the thalamic expression of *Ntn1* and *Sema3a* and their

receptors change in a way that might explain the medially-directed deviation of

lateral axons that we observed in the thalamus of *CAG^CreER Pax6* cKOs. In these

embryos, we found that the medial domain of *Ntn1*-expression was retained and

appeared enlarged. *Sema3a* was still expressed higher laterally, although overall

levels seemed reduced (Fig. 6F,G). These patterns are reconstructed in 3D in Fig.

6H. *Ntn1* and *Sema3a* expression in the subpallium of *CAG^CreER Pax6* cKOs

appeared to be unaffected (Fig. S4K-N). The significance of changes in the

expression levels of Sema3a and Ntn1 in the thalamus of CAG^{CreER} $Pax6$ cKOs was

confirmed by analysing a previously published RNAseq dataset (Fig. S4C, dataset

from Quintana-Urzainqui et al., 2018).

Regarding the expression of guidance receptors, fewer laterally-located neurons

expressed $Unc5c$ in CAG^{CreER} $Pax6$ cKOs than in controls (compare Fig. 6I' and D').

Significant numbers of $Unc5c$-negative neurons were now intermingled with $Unc5c$-

positive neurons even in the most lateral thalamic tissue (Fig. 6I'). $Plxna1$'s thalamic

expression pattern did not change in the absence of Pax6 (Fig. 6J, S4D), nor did that

of any of the other receptor-coding genes studied (Fig. S4E-J).

As reported above, we discovered that CAG^{CreER} $Pax6$ cKOs show a misrouting in a

medial direction of axons from the lateral thalamus (Fig. 1), and our finding that

many laterally-located thalamic neurons lose their expression of $Unc5c$ in these

mutants, provides a likely explanation, summarized in Fig. 6K. We propose that

$Unc5c$-negative laterally-located thalamic neurons in CAG^{CreER} $Pax6$ cKO thalamus

would no longer be repelled from the medial thalamus by its high levels of $Ntn1$.

Consequently, they would be more likely to stray, or perhaps to be pushed by

relatively high lateral levels of $Sema3a$, towards a medial direction (Fig. 6K).

Subsets of Pax6-deficient lateral thalamic axons deviate medially when

confronted with control thalamus

Finally, we grafted GFP-positive lateral thalamic explants from CAG^{CreER} $Pax6$ cKO

embryos into the lateral thalamus of GFP-negative slices from littermate control

embryos, following the same experimental paradigm as before (Fig. 7A, see Figure 5

and Methods). Our model would predict that subsets of Pax6-deficient lateral

thalamic axons, presumably those that have lost their repulsion to Ntn1, would

deviate towards medial areas when confronted with control thalamus. We used 11

embryos from four different litters and performed a total of 22 (bilateral) graft

experiments. In all cases we observed subsets of GFP-positive axons deviating

medially (Figure 7B, red arrows). To measure the navigation pattern of these axons

we divided the thalamus in three equal medial-lateral sectors and quantified the

percentage of GFP on each of them (Fig. 7C). For this analysis we used four

explants from four different litters (5 slices per explant). We found that axons spread

throughout the three areas of the thalamus (45.8% laterally, 31.8% intermedially,

22.44% medially; Fig 7C). This is in striking contrast to the almost invariably (95.7%)

lateral trajectories of axons from control lateral thalamic explants grafted into control

thalamic tissue (Fig. 5C-F, K). This evidence supports our model and the idea that

thalamic neurons hold intrinsic information that determines the medial-lateral position

of their axons when crossing the thalamus.

Overall, our findings indicate that mechanisms exist within the thalamus itself to

ensure that its TCAs exit in an orderly manner and that these mechanisms may play

an important part in the correct topographic mapping of TCAs onto the cortex.

DISCUSSION

During embryonic development, thalamic axons undertake a long journey, navigating

through several tissues before they arrive at the cortex. It is therefore crucial that

their guidance is tightly regulated by mechanisms placed along the route. Previous

studies have demonstrated the importance of the ventral telencephalon as an

intermediate target for establishing correct topographical thalamocortical

connections. Here, gradients of signalling molecules sort different subsets of TCAs

towards different areas of the embryonic cortex (Antón-Bolaños et al., 2018; Bielle et

al., 2011; Braisted et al., 1999; Dufour et al., 2003; Métin and Godement, 1996;

Molnár et al., 2012; Vanderhaeghen and Polleux, 2004). The importance of other

tissues along the route in the establishment and/or maintenance of axonal

topography remained unexplored. We now show that if thalamic axons do not

emerge in order from the thalamus they will connect with the wrong areas of the

cortex, resulting in topographical defects. This highlights the importance of

maintaining axonal order throughout the route and suggests the existence of

guidance mechanisms within the diencephalon to guarantee this happens.

Our *in vitro* graft experiments demonstrated that embryonic thalamic tissue is

instructive for TCAs and sorts axons according to their original medio-lateral position.

We went on to find a possible guidance mechanism acting within the thalamus. The

thalamus expresses *Ntn1* and *Sema3a,* some of the same guidance molecules

known to guide TCAs in the ventral telencephalon (Bielle et al., 2011; Powell et al.,

2008; Wright et al., 2007). What is more, there is an interesting correspondence

between the regions expressing each of those molecules in the thalamus and in the

ventral telencephalon. TCAs that emerge and navigate through the *Sema3a*-high

region of the thalamus (lateral-caudal thalamus, dLGN) are steered towards the

Sema3a-high region in the ventral telencephalon, while axons that emerge and

navigate through the *Ntn1*-high region of the thalamus (ventral-medial

thalamus,VMP) are sorted towards *Ntn1*-high regions in the ventral telencephalon.

This suggests that each subset of thalamic axons maintains the expression of the

same combination of axon guidance receptors along the route and therefore show

the same chemotactic response when confronting gradients of signalling cues.

Likewise, it indicates that the same gradients of guidance molecules are re-used at different levels of the thalamocortical pathway to maintain topographic order.

The chemotactic behaviour of TCAs with respect to *Sema3a* and *Ntn1* gradients in the thalamus and ventral telencephalon can be explained by our observations of the expression of Sema3a and Ntn1 receptors in developing thalamus. We show that all thalamic neurons seem to express homogeneous levels of *Plxna1*, a receptor mediating repulsion to Sema3a (Rohm et al., 2000; Takahashi et al., 1999; Tamagnone et al., 1999), while Unc5c, a receptor mediating repulsion to Ntn1 (Leonardo et al., 1997), was found to be expressed in a lateral-high medial-low gradient. According to these observations, we propose a model in which complementary expression patterns of Sema3a and Ntn1 can establish topographical order on TCAs by a mechanism of double repulsion, in which all thalamic axons have the potential to be repelled by Sema3a but only lateral axons are additionally repelled by Ntn1. It is possible that laterally-derived axons experience stronger Ntn1 repulsion the more lateral they are. Thus, lateral thalamic axons prefer to navigate through Sema3a-high, Ntn1-low regions because they might be more strongly repelled by Ntn1 than by Sema3a. Axons located in intermediate regions of the thalamus express lower levels of Unc5c; thus they might be equally repelled by Sema3a and Ntn1 and chose to navigate across regions with moderate levels of both signalling cues. Likewise, medial axons are only repelled by Sema3a and neutral to Ntn1: therefore they chose to navigate through Sema3a-low, Ntn1-high areas.

Supporting this model are the experiments showing that TCAs are repelled by Nnt1 (Bielle et al., 2011; Bonnin et al., 2007; Powell et al., 2008) and thalamic growth cones show retraction in the presence of Sema3a (Bagnard et al., 2001). Moreover,

Wright and colleagues reported that in mice harbouring a mutation that makes the

axons non-responsive to Sema3a, axons from the ventrobasal (VB) thalamic nucleus

were caudally shifted and target the visual cortex instead of the somatosensory

cortex (Wright et al., 2007). Our double repulsion model satisfactorily explains this

phenotype. The VB nucleus is located in an intermediate thalamic region that would

contain substantial number of Unc5c-positive neurons. In those mutants, VB axons

lose their repulsion to Sema3a but many would still be repelled by Ntn1, and

therefore would steer towards Sema3a-high, Ntn1-low regions both in the thalamus

and the ventral telencephalon.

The behaviour of thalamic axons in our explant experiments using donor tissue from

controls and CAG^{CreER} Pax6 cKOs, also support the model.

Other molecules known to form gradients and guide TCAs in the ventral

telencephalon, like Slit1 or Ephrin A5 (Bielle et al., 2011; Dufour et al., 2003; Molnár

et al., 2012; Seibt et al., 2003; Vanderhaeghen and Polleux, 2004) were not

analysed in this study. It remains to be tested whether these molecules and their

receptors are expressed in the thalamus in a gradient fashion and if they follow the

same rules proposed in our model.

It is important to highlight that in this study we only considered the medio-lateral axis

of the main thalamic body, but the same or other guidance cues and receptors

probably function in other directions. For example, work in mice showed that Unc5c

(Bonnin et al., 2007) and DCC (Powel et al., 2008) are also highly expressed in the

rostral thalamus, at a level we did not cover in our expression and tracer analyses.

Other important question is how Pax6 inactivation leads to deficits in thalamic

organization, since Pax6 is only expressed in progenitors. We show here and in a

previous paper that thalamic patterning does not seem to be largely affected in

CAG^CreER Pax6 cKOs (Fig. S1 and Quintana-Urzainqui et al., 2018) and that the

main changes in gene expression in postmitotic neurons are related with axon

guidance molecules. One possibility is that Pax6 expression in thalamic progenitors

affects transcriptional programmes that indirectly translates into actions on the

postmitotic expression of certain axon guidance molecules.

Finally, our results have given interesting new insights into the development and the

importance of the pioneer axons from the prethalamus to the thalamus. First, we

found that the prethalamic neurons extending pioneer axons to the thalamus belong

to a particular lineage, the Gsx2-lineage and not the Zic4-lineage. Second, although

disturbing the prethalamus-to-thalamus pioneers did not stop TCAs reaching the

cortex without any topographic error, it did cause them to fasciculate prematurely as

they crossed the prethalamus. Growing axons often increase their fasciculation when

they cross regions that are hostile to their growth. Our and other studies have shown

that the prethalamus also expresses guidance cues with potential to exert a

repulsive response of TCAs (Ono et al., 2014), thus it is possible that interactions

between the developing TCAs and prethalamic pioneers somehow helps the

passage of the TCAs across this region. Further work is needed to discover what the

consequences are if this help is unavailable.

Material and Methods

Mice

All animals (*Mus musculus*) were bred according to the guidelines of the UK Animals

(Scientific Procedures) Act 1986 and all procedures were approved by Edinburgh

University's Animal Ethics Committee.

The day the vaginal plug was detected was considered E0.5

For conditional inactivation of Pax6, we used a tamoxifen-inducible Pax6^{loxP} allele

(Simpson et al., 2009) and a RCE:LoxP EGFP Cre reporter allele (Sousa et al.,

2009) and we combined them with different Cre lines. To generate a deletion of Pax6

throughout the embryo, we used lines carrying a CAGGCre-ER^{TM} allele (Hayashi

and McMahon, 2002; Quintana-Urzainqui et al., 2018). As controls, we used both

wild type and *CAG^{CreER}Pax6$^{fl/+}$* littermate embryos since the latter express normal

levels of Pax6 protein, almost certainly because of a feedback loop that

compensates for a deletion in one allele by increasing the activity of the other

(Caballero et al., 2014; Manuel et al., 2015).

To inactivate Pax6 in different parts of the prethalamus we used either a Gsx2-Cre

(Kessaris et al., 2006) or the Zic4-Cre allele (Rubin et al., 2011). For cortex-specific

deletion of Pax6, we used Emx1Cre-ER^{T2} (Kessaris et al., 2006).

The DTy54 YAC reporter allele (Tyas et al., 2006) was combined with the Gsx2-Cre

allele to generate Gsx2Cre; Pax6$^{loxP/loxP}$ embryos expressing tauGFP in cells in which

the Pax6 gene is active.

Embryos heterozygous for the Pax6^{loxP} allele (Pax6$^{fl/+}$) were used as controls since

previous studies have shown no detectable defects in the forebrain of Pax6$^{fl/+}$

embryos (Simpson et al., 2009). Embryos carrying two copies of the floxed Pax6

allele (Pax6$^{fl/fl}$) were the experimental conditional knock-out (cKO) groups.

For thalamic explant experiments, we generated litters containing GFP-positive and

negative embryos by crossing a line of heterozygous studs for a constitutively active

form of CAGGCre-ER^{TM} allele and the and a RCE:LoxP EGFP Cre reporter allele

with wild type females. In the second set of explant experiments, we performed

crossings to generate litters containing GFP-positive Pax6 cKO embryos

(CAGCRE$^{ER\ TM}$; Pax6$^{fl/fl}$), and GFP-negative control embryos (homozygous for the

Pax6 floxed allele but not expressing the CRE allele). Both embryos carried a

RCE:LoxP EGFP allele reporting Cre activity and allowing us to select the embryos

within the same litter and visualizing Pax6 cKO axons in the explants

The day the vaginal plug was detected was considered E0.5. Pregnant mice were

given 10mg of tamoxifen (Sigma) by oral gavage on embryonic day 9.5 (E9.5) and

embryos were collected on E12.5, E13.5, E14.5, E15.5, E16.5 or E18.5 For the DiI

and DiA tracing experiments, wild type embryos (CD1 background) were additionally

used as controls.

Immunohistochemistry

Embryos were decapitated and fixed in 4% paraformaldehyde (PFA) in phosphate

buffered saline (PBS) overnight at 4^{o}C. After washes in PBS, heads were

cryoprotected by immersion in 30% sucrose in PBS, embedded in OCT Compound

and sectioned using a cryostat at 10μm.

Cryo-sections were let to stabilize at room temperature for at least 2 hours and then

washed three times in PBST (1X PBS with 0.1% Triton X-100, Sigma). To block

endogenous peroxidase, sections were treated with 3% H_2O_2 for 10 minutes. After

PBS washes, antigen retrieval was performed by immersing the sections in Sodium

Citrate buffer (10mM, pH6) heated at approximate 90°C using a microwave for 20

minutes. Sections were then incubated with the rabbit polyclonal anti-Pax6 (1:200,

BioLegend Cat # 901302) overnight at 4°C. The secondary antibody (goat anti-rabbit

bioninylated, 1:200, Vector laboratories Cat # BA-1000) was incubated for 1 hour at

room temperature followed by a 30-minute incubation with Avidin-Biotin complex

(ABC kit, Vector laboratories Cat # PK6100). Finally, diaminobenzidene (DAB,

Vector Laboratories, Cat # SK4100) reaction was used to obtain a brown precipitate

and sections were mounted in DPX media (Sigma-Aldrich, Cat # 06522).

For immunofluorescence, cryosections were incubated overnight at 4°C with the

following primary antibodies: rat monoclonal anti-Neural Cell Adhesion Molecule L1

(1:500 Millipore, Cat # MAB5272, clone 324, RRID:AB_2133200), rabbit polyclonal

anti-Pax6 (1:200, BioLegend Cat # 901302; RRID:AB_2565003), goat polyclonal anti-

GFP (1:200, Abcam Cat # ab6673, RRID:AB_305643), rabbit polyclonal anti-GFP

(1:200, Abcam Cat # ab290, RRID:AB_303395). The following secondary antibodies

from Thermo Fisher Scientific were incubated at room temperature for one hour:

Donkey anti-rat Alexa[488] (1:100, Thermo Fisher, Cat # A-21208), Donkey anti-rat

Alexa[594] (1:100, Thermo Fisher, Cat # A-21209), Donkey anti-rabbit Alexa[568] (1:100,

Thermo Fisher, A10042), Donkey anti-rabbit Alexa[488] (1:100, Thermo Fisher Cat #

R37118), Donkey anti-goat Alexa[488] (1:100, Invitrogen, Cat # A11055). Sections

were counterstained with DAPI (Thermo Fisher Scientific, Cat # D1306) and

mounted in ProLong Gold Antifade Mountant (Thermo Fisher Scientific, Cat #

P36930).

In situ hybridization

In vitro transcription of digoxigenin-labelled probes was done with DIG RNA-labeling

kit (Sigma-Aldrich, Cat # 11175025910). The following digoxigenin-labelled probes

were synthetized in the lab from cDNA: Ntn1 (kindly donated by Dr Thomas Theil;

forward primer: CTTCCTCACCGACCTCAATAAC, reverse primer:

GCGATTTAGGTGACACTATAGTTGTGCC TACAGTCACACACC), Sema3a

(forward primer: ACTGCTCTGACTTGGAGGAC, reverse primer:

ACAAACACGAGTGCTGGTAG), Plxna1 (forward primer:

GACGAGATTCTGGTGGCTCT, reverse primer: CATGGCAGGGAGAGGAAGG),

DCC (forward primer: AACAGAAGGTCAAGCACGTG, reverse primer:

CAATCACCACGACCAACACA), Unc5a (forward primer:

CTGTCAGACCCTGCTGAGT, reverse primer: GGGCTAGAGTTCGCCAGTC),

Unc5d (forward primer: GGACAGAGCTGAGGACAACT, reverse primer:

GTATCAAACGTGGCGCAGAT). Unc5c probe was kindly donated by Dr. Vassiliki

Fotaki, University of Edinburgh, UK and Dr Suran Ackerman, UCSanDiego, USA).

Cryosections were processed for *in situ* hybridization (ISH) using standard protocols.

Some slides were counterstained for nuclear fast red (Vector Laboratories, Cat# LS-

J1044-500).

Axon Tract Tracing

For cortical injections, brains were dissected between E15.5 and E18.5 and fixed in

4% PFA in PBS at 4°C for at least 48 hours. After washes in PBS, filter paper

impregnated in DiI (NeuroVue Red, Molecular Targeting Technologies, Cat # FS-

1002) and DiA (NeuroVue Jade, molecular Targeting Technologies, Cat # FS-1006)

was inserted approximately in the somatosensory and visual areas of the cortex,

respectively. Brains were incubated at 37°C in PBS for 4 weeks to allow the diffusion

of the tracers.

For thalamic injections in fixed tissue, embryos were dissected at E13.5 and fixed

overnight in 4%PFA in PBS at 4°C. After PBS washes, brains were cut in half at the

midline and DiI was inserted in the thalamus using a fine probe. Brains were

incubated for 1 week in PBS at 37°C.

Brains were then cryoprotected in 30% sucrose, embedded in OCT Compound and

sectioned in a cryostat at 30μm. Sections were counterstained with DAPI diluted

1:1000 in distilled water.

For thalamic injections in non-fixed tissue, we applied neurobiotin (Vector

Laboratories, Cat # SP-1120), and amino derivative of biotin used as an intracellular

label for neurons. The tracer in powder was held at the tip of an entomological

needle (00) and recrystallized using vapour from distilled water. Brains were cut in

half and the crystal was inserted in the thalamus. Brains were then immersed in

continuously oxygenated Ringer (124mM NaCl, 5mM KCl, 1.2mM KH_2P0_4, 1.3mM

$MgSO_4$ $7H_2O$, 26mM $NaHCO_3$, 2.4mM $CaCl_2$ $2H_2O$, 10mM glucose) and incubated

overnight at RT.The tissue was fixed in 4% PFA in PBS overnight at 4°C, washed in

PBS, cryoprotected in 30% sucrose and sectioned in a cryostat at 10μm. Neurobiotin

was visualized by incubating the sections with either Strep[488] or Strep[546].

Thalamic explants and slice culture

E13.5 embryos were dissected, embedded in 4% low melting temperature agarose

(Lonza, Cat # 50100) and sectioned in a vibratome to produce 300μm-thick coronal

slices. Lateral or medial thalamic explants were dissected from slices belonging to

GFP-positive embryos and transplanted into equivalent rostral/caudal slices

belonging to GFP-negative embryos (see schemas in Fig. 5). The thalamus and its

different medio-lateral regions were recognized under the dissecting scope by

anatomical landmarks. Slices were then cultured for 72 hours in floating membranes

(Whatman nuclepore track-etched membranes, Cat# WHA110414) over serum-free

Neurobasal medium (Thermo Fisher Scientific, Cat# 21103049) in 60mm center well

organ culture dishes (Falcon, Cat# 353037). Cultures were fixed in 4% PFA

overnight at 4°C, cryoprotected in 30% sucrose and cryosectioned at 10µm to be

processed for immunofluorescence. All grafts were positioned in direct contact with

presectioned host tissue to avoid pial growth.

We performed a total of 31 transplant experiments. In the first set of experiments, we

grafted control thalamus into control tissue. We used embryos from four different

litters to a total of nine different transplants (four using lateral thalamus and five

using medial thalamus as donor tissue). In the second set of experiments, we grafted

Pax6 cKO lateral thalamus into control thalamus, using 11 embryos from four

different litters and a total of 22 individual (bilateral) transplants.

Image analysis and quantification of thalamic area occupied by DiI and DiA

Images of transverse sections were analysed blind using Fiji Software (Schindelin et

al., 2012) for seven E15.5 embryos belonging to three different litters (four controls

and three *CAG^{CreER} Pax6* cKOs). For each embryo, we analysed at least 5 sections

at different rostral-caudal levels. For each section, we isolated the DAPI channel and

defined the dLGN and VP nucleus as regions of interest (ROIs) blind to the other

channels (DiI and DiA). We next measured the area occupied by DiI and DiA within

each ROI, defining positive label using the automated thresholding function in Fiji

software set for "MaxEntropy". Data from all sections and genotypes was statistically

assessed to test the effects of Pax6 inactivation on the area occupied by each dye

on each nucleus. Data was fitted to a mixed linear model using the lmer() function

from lme4 R package (Bates et al., 2015). "DiI in dLG", "DiI in VP", "DiA in dLG" and

"DiA in VP" were set as dependant variables, with "genotype" as a fixed effect and

"litter:embryo" factors as nested random effects. P-values of fixed effects were

obtained using the Anova() function from car package (Fox and Weisberg, 2011).

Quantification of thalamic explant experiments

Images of transverse sections of our explants were analysed using Fiji Software

(Schindelin et al., 2012). We quantified a total of ten culture thalamic explants (three

from lateral control donor tissue, three from medial control donnor tissue and four

from lateral Pax6 cKO tissue) from seven different litters, using five slices per explant

corresponding to different rostral-caudal levels of the explants. For each section we

first divided the thalamus into three equal medial-lateral sectors (defined as ROIs,

see Figure 5), avoiding the area occupied by the explant itself. We then isolated the

GFP channel and measured the area occupied by GFP-positive elements within

each ROI, defining positive label using the automated thresholding function in Fiji

software set for "MaxEntropy". To test for the differences of GFP occupancy in the

three different areas (lateral, intermedial, medial), the data was fitted to a mixed

linear model using the lmer() function from lme4 R package (Bates et al., 2015).

For the DiI and thalamic explant quantifications, data was presented as box plots

representing the median value and the distribution of the quartiles. Individual

datapoints representing measurements of each slice were plotted overlying the box

plots.

Quantifications of numbers of axons and bundle width

Images were blinded analysed for at least three E13.5 embryos for each condition

belonging to three different litters. We positioned three lines across the prethalamus:

(1) at the thalamic-prethalamic border (Th-PTh), guided by prethalamic expression of

Pax6; (2) at a lower prethalamic position (low-PTh), guided by the end of Pax6

prethalamic expression; and (3) at the midpoint position between the two other lines

(mid-PTh). We generated a L1 intensity profile using Fiji Software (Schindelin et al.,

2012). Intensity profiles were then processed by tracing a line at an arbitrary (but

constant for all quantifications) intensity level and quantifying the number and width

of bundles crossing the line. Statistical significance was assessed applying two-tailed

unpaired Student's t-test and N=3.

3D reconstruction

We used Free-D software (Andrey and Maurin, 2005) to reconstruct the structure of

thalamus and prethalamus from transverse slices stained with DAPI and antibodies

against L1 and Pax6 to reveal the thalamocortical tract and the limits of the

diencephalic structures, respectively. The thalamus territory was recognisable by an

intense DAPI staining and Pax6-negative mantle zone, contrasting with prethalamus

and pretectum, which express high levels of Pax6 in the postmitotic neurons. The

location of the signalling molecules was included in the model by comparison of

transversal and sagittal sections stained for Ntn1 and Sema3a and their adjacent

sections processed for Pax6 and L1 with the sections used to build the model

scaffold. Dots are representation of staining density. We used sagittal and transverse

sections from four embryos from three different litters.

Microscopy and imaging

ISH and IHQ images were taken with a Leica DMNB microscope coupled to a Leica

DFC480 camera. Fluorescence images were taken using a Leica DM5500B

automated epifluorescence microscope connected to a DFC360FX camera. Image

panels were created with Adobe Photoshop CS6.

Acknowledgements

We thank Dr Martine Manuel for her invaluable help with the generation of mice lines

for the explant experiments and Dr Vassiliki Fotaki for providing the Unc5c probe.

Competing interests

No competing interests.

Funding

This work was supported by a Marie Curie Fellowship from the European

Commission [624441]; a Medical Research Council UK Research Grant [N012291]

and a Biotechnology and Biological Sciences Research Council UK Research Grant

[N006542].

References

Amassian, V. and Weiner, H. (1966) Monosynaptic and polysynaptic activation of pyramidal

tract neurons by thalamic stimulation. In *The thalamus* (ed. D.P. Purpura and M.D. Yahr

editors), pp 255–282. New York: Columbia University Press.

Andrey, P. and Maurin, Y. (2005). Free-D: an integrated environment for three-dimensional

reconstruction from serial sections. *J. Neurosci. Methods* **145**, 233–244.

Angevine, J. A. Y. B. (1970). Time of Neuron Origin in the Diencephalon of the Mouse . An

Autoradiographic Study '.

Antón-Bolaños, N., Espinosa, A. and López-Bendito, G. (2018). Developmental interactions

between thalamus and cortex: a true love reciprocal story. *Curr. Opin. Neurobiol.* **52**,

33-41.

Auladell, C. and Hans, P. P. (2000). The early development of thalamocortical and

corticothalamic projections in the mouse. *Anat Embryol* **201**, 169–179.

Bagnard, D., Chounlamountri, N., Püschel, A.W., Bolz, J. (2001). Axonal Surface Molecules

Act in Combination with Semaphorin 3A during the Establishment of Corticothalamic

Projections. *Cereb. Cortex* **11**, 278–285.

Bates, D., Maechler, M., Bolker, B., Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *J. Stat. Softw.* **67**, 1-48.

Bielle, F., Marcos-Mondéjar, P., Leyva-Díaz, E., Lokmane, L., Mire, E., Mailhes, C., Keita,

M., García, N., Tessier-Lavigne, M., Garel, S., et al. (2011). Emergent growth cone

responses to combinations of Slit1 and Netrin 1 in thalamocortical axon topography.

Curr. Biol. **21**, 1748–1755.

Bonnin, A., Torii, M., Wang, L., Rakic, P. and Levitt, P. (2007). Serotonin modulates the

response of embryonic thalamocortical axons to netrin-1. **10**, 588–597.

Bosch-Bouju, C., Hyland, B. I. and Parr-Brownlie, L. C. (2013). Motor thalamus integration

of cortical, cerebellar and basal ganglia information: implications for normal and

parkinsonian conditions. *Front. Comput. Neurosci.* **7**, 1–21.

Braisted, J. E., Tuttle, R. and Leary, D. D. M. O. (1999). Thalamocortical Axons Are

Influenced by Chemorepellent and Chemoattractant Activities Localized to Decision

Points along Their Path. **440**, 430–440.

Caballero, I. M., Manuel, M. N., Molinek, M., Quintana-Urzainqui, I., Mi, D., Shimogori, T.

and Price, D. J. (2014). Cell-Autonomous Repression of Shh by Transcription Factor

Pax6 Regulates Diencephalic Patterning by Controlling the Central Diencephalic

Organizer. *Cell Rep.* **8**, 1405–1418.

Clegg, J. M., Li, Z., Molinek, M., Caballero, I. M., Manuel, M. N. and Price, D. J. (2015). Pax6

is required intrinsically by thalamic progenitors for the normal molecular patterning of

thalamic neurons but not the growth and guidance of their axons. *Neural Dev.* 1–12.

Dufour, A., Seibt, J., Passante, L., Depaepe, V., Ciossek, T., Frisén, J., Kullander, K.,

Flanagan, J. G., Polleux, F. and Vanderhaeghen, P. (2003). Area specificity and

topography of thalamocortical projections are controlled by ephrin/Eph genes. *Neuron*

39, 453–465.

Fox, J., Weisberg, S. (2019). An R Companion to Applied Regression, Third edition. Sage,

Thousand Oaks CA

Fukuda, T., Kawano, H., Ohyama, K. and Li, H. (1997). Immunohistochemical localization of

Neurocan and L1 in the Formation of Thalamocortical Pathway. **152**, 141–152.

Gaston-massuet, C., Henderson, D. J., Greene, N. D. E. and Copp, A. J. (2005). Zic4 , a Zinc-

finger transcription factor, is expressed in the developing mouse nervous system. **233**,

1110–1115.

Georgala, P. A., Carr, C. B. and Price, D. J. (2011). The role of Pax6 in forebrain

development. *Dev. Neurobiol.* **71**, 690–709.

Hayashi, S. and McMahon, A. P. (2002). Efficient Recombination in Diverse Tissues by a

Tamoxifen-Inducible Form of Cre: A Tool for Temporally Regulated Gene

Activation/Inactivation in the Mouse. *Dev. Biol.* **244**, 305–318.

Jones, L., López-bendito, G., Gruss, P., Stoykova, A. and Molnár, Z. (2002). Pax6 is required

for the normal development of the forebrain axonal connections. **5052**, 5041–5052.

Jones, E. G. (2007). The Thalamus. 2nd Edn. Cambridge: Cambridge University Press.

Kessaris, N., Fogarty, M., Iannarelli, P., Grist, M., Wegner, M. and Richardson, W. D.

(2006). Competing waves of oligodendrocytes in the forebrain and postnatal

elimination of an embryonic lineage. *Nat. Neurosci.* **9**, 173–179.

Leonardo, E. D., Hinck, L., Masu, M., Keino-Masu, K., Ackerman, S. L. and Tessier-Lavigne,

M. (1997). Vertebrate homologues of C. elegans UNC-5 are candidate netrin receptors.

Nature **386**, 833–838.

Li, Z., Pratt, T. and Price, D. J. (2018). Zic4 -Lineage Cells Increase Their Contribution to

Visual Thalamic Nuclei during Murine Embryogenesis If They Are Homozygous or

Heterozygous for Loss of Pax6 Function. *eNeuro* **1**, 1–19.

Lopez-Bendito, G., Cautinat, A., Sanchez, J. A., Bielle, F., Flames, N., Garratt, A. N.,

Talmage, D. A., Role, L. W., Charnay, P., Marin, O., et al. (2006). Tangential neuronal

migration controls axon guidance: a role for neuregulin-1 in thalamocortical axon

navigation. *Cell* **125**, 127–142.

López-Bendito, G. and Molnár, Z. (2003). Thalamocortical development: How are we going

to get there? *Nat. Rev. Neurosci.* **4**, 276–289.

Manuel, M. N., Mi, D., Mason, J. O. and Price, D. J. (2015). Regulation of cerebral cortical

neurogenesis by the Pax6 transcription factor. *Front. Cell. Neurosci.* **9**, 1–21.

Métin, C. and Godement, P. (1996). The Ganglionic Eminence May Be an Intermediate

Target for Corticofugal and Thalamocortical Axons. *J. Neurosci.* **16**, 3219–3235.

Mi, D., Carr, C. B., Georgala, P. A., Huang, Y. T., Manuel, M. N., Jeanes, E., Niisato, E.,

Sansom, S. N., Livesey, F. J., Theil, T., et al. (2013). Pax6 Exerts regional control of

cortical progenitor proliferation via direct repression of Cdk6 and Hypophosphorylation

of pRb. *Neuron* **78**, 269–284.

Molnár, Z., Adams, R. and Blakemore, C. (1998). Mechanisms Underlying the Early

Establishment of Thalamocortical Connections in the Rat. *J. Neurosci.* **18**, 5723–5745.

Molnár, Z., Garel, S., López-bendito, G., Maness, P. and Price, J. (2012). Mechanisms

Controlling the Guidance of Thalamocortical Axons Through the Embryonic Forebrain.

35, 1573–1585.

Ohyama, K., Tan-takeuchi, K., Kutsche, M., Schachner, M., Uyemura, K. and Kawamura, K.

(2004). Neural cell adhesion molecule L1 is required for fasciculation and routing of

thalamocortical fibres and corticothalamic fibres. **48**, 471–475.

Ono, K., Clavairoly, A., Nomura, T., Gotoh, H., Uno, A., Armant, O., Takebayashi, H., Zhang,

Q., Shimamura, K., Itohara, S., et al. (2014). Development of the prethalamus is crucial

for thalamocortical projection formation and is regulated by Olig2. **5**, 2075–2084.

Parish, E. V., Mason, J. O. and Price, D. J. (2016). Expression of Barhl2 and its relationship

with Pax6 expression in the forebrain of the mouse embryo. *BMC Neurosci.* **17**, 1–16.

Piñon, M. C., Tuoc, T. C., Ashery-padan, R. and Stoykova, A. (2008). Altered Molecular

Regionalization and Normal Thalamocortical Connections in Cortex-Specific Pax6

Knock-Out Mice. **28**, 8724–8734.

Powell, A. W., Sassa, T., Wu, Y., Tessier-lavigne, M. and Polleux, F. (2008). Topography of

Thalamic Projections Requires Attractive and Repulsive Functions of Netrin-1 in the

Ventral Telencephalon. **6**, e116.

Pratt, T., Vitalis, T., Warren, N., Edgar, J. M., Mason, J. O. and Price, D. J. (2000). A role for

Pax6 in the normal development of dorsal thalamus and its cortical connections.

Development **127**, 5167-5178.

Price, D. J., Clegg, J., Duocastella, X. O., Willshaw, D. and Pratt, T. (2012). The importance

of combinatorial gene expression in early mammalian thalamic patterning and

thalamocortical axonal guidance. **6**, 1–15.

Quintana-Urzainqui, I., Kozic, Z., Tian, T., Manuel, M., Mason, J. O., Price, D. J., Mitra, S.,

Tian, T., Manuel, M. and Mason, J. O. (2018). Tissue-Specific Actions of Pax6 on

Proliferation and Differentiation Balance in Developing Forebrain Are Foxg1

Dependent. **10**, 171–191.

Rohm, B., Ottemeyer, A., Lohrum, M. and Puschel, A. W. (2000). Plexin/neuropilin

complexes mediate repulsion by the axonal guidance signal semaphorin 3A. *Mech. Dev.*

93, 95–104.

Rubin, A. N., Alfonsi, F., Humphreys, M. P., Choi, C. K. P. and Susana, F. (2011). The

germinal zones of the basal ganglia but not the septum generate GABAergic

interneurons for the cortex. **30**, 12050–12062.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for

biological-image analysis. *Nat. Methods* **9**, 676–682.

Seibt, J., Schuurmans, C., Dehay, C., Vanderhaeghen, P., Hill, C. and Carolina, N. (2003).

Neurogenin2 Specifies the Connectivity of Thalamic Neurons by Controlling Axon

Responsiveness to Intermediate Target Cues. **39**, 439–452.

Simpson, T. I., Pratt, T., Mason, J. O. and Price, D. J. (2009). Normal ventral telencephalic

expression of Pax6 is required for normal development of thalamocortical axons in

embryonic mice. **5**, 1–18.

Sousa, V. H., Miyoshi, G., Hjerling-Leffler, J., Karayannis, T. and Fishell, G. (2009).

Characterization of Nkx6-2-derived neocortical interneuron lineages. *Cereb. Cortex* **19**.

Stoykova, A., Fritsch, R., Walther, C. and Gruss, P. (1996). Forebrain patterning defects in

Small eye mutant mice. *Development* **122**, 3453-3465.

Takahashi, T., Fournier, A., Nakamura, F., Wang, L. H., Murakami, Y., Kalb, R. G., Fujisawa,

H. and Strittmatter, S. M. (1999). Plexin-neuropilin-1 complexes form functional

semaphorin-3A receptors. *Cell* **99**, 59–69.

Tamagnone, L., Artigiani, S., Chen, H., He, Z., Ming, G. I., Song, H., Chedotal, A., Winberg,

M. L., Goodman, C. S., Poo, M., et al. (1999). Plexins are a large family of receptors for

transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. *Cell* **99**, 71–

80.

Tlamsa, A. P. and Brumberg, J. C. (2010). Organization and morphology of thalamocortical

neurons of mouse ventral lateral thalamus. *Somatosens. Mot. Res.* **27**, 34–43.

Tuttle, R., Nakagawa, Y., Johnson, J. E. and Leary, D. D. M. O. (1999). Defects in

thalamocortical axon pathfinding correlate with altered cell domains in Mash-1-

deficient mice. **1916**, 1903–1916.

Tyas, D. A., Simpson, T. I., Carr, C. B., Kleinjan, D. A., van Heyningen, V., Mason, J. O. and

Price, D. J. (2006). Functional conservation of Pax6 regulatory elements in humans and

mice demonstrated with a novel transgenic reporter mouse. *BMC Dev. Biol.* **6**, 1–11.

Vanderhaeghen, P. and Polleux, F. (2004). Developmental mechanisms patterning

thalamocortical projections: Intrinsic, extrinsic and in between. *Trends Neurosci.* **27**,

384–391.

Visel, A., Thaller, C. and Eichele, G. (2004). GenePaint.org: an atlas of gene expression

patterns in the mouse embryo. *Nucleic Acids Res.* **32**, D552-6.

Walther, C. and Gruss, P. (1991). Pax-6, a murine paired box gene, is expressed in the

developing CNS. *Development* **113**, 1435-1449.

Warren, N. and Price, D. J. (1997). Roles of Pax-6 in murine diencephalic development.

Development **124**, 1573-1582.

Wright, A. G., Demyanenko, G. P., Powell, A., Schachner, M., Enriquez-barreto, L., Tran, T.

S., Polleux, F. and Maness, P. F. (2007). Close Homolog of L1 and Neuropilin 1 Mediate

Guidance of Thalamocortical Axons at the Ventral Telencephalon. **27**, 13667–13679.

Figure 1

Figure 1. Ubiquitous conditional Pax6 deletion at E9.5 causes mapping errors in the thalamocortical connection. A) Schema showing how thalamic axons map to specific areas of the cortex via the thalamocortical pathway. **B)** Schematic drawing representing the cortical location where the tracers were placed at E15.5. **C-H)** Fluorescent microphotographs showing cells and axons retrogradely labelled by DiI and DiA in transverse sections. While in controls (C,E) the areas labelled in the thalamus were clearly separated (dotted lines), in CAG^{CreER} Pax6 cKOs (F-H) they visibly overlapped, indicating the existence of mapping errors in these mutants. Arrows in G,H and I indicate that the thalamocortical bundle was segregated in visual (green) and somatosensory (red) halves at the level of their exit from the diencephalon. **C', D', E',F',G',H')** High power images from insets in C,D,E,F,G and H, respectively, where only the DiI channel is shown. Arrows in G' and H' point to DiI-labeled axons showing abnormal trajectories towards medial thalamic regions. **I)** Box plots showing the area occupied by each tracer (DiI or DiA) in each defined area (dLGN or VP) in controls versus Pax6 cKOs. Seven E15.5 embryos (four controls and three Pax6 cKOs), from three different litters were analysed. For each embryo, at least five rostrocaudal sections were quantified. Each dot represents an area value of a single section. P values were calculated by fitting the data to a generalized mixed linear model. * indicates $p<0.001$, *** indicates $p< 1.0 e-15$ **J)** Schematic summary of the tract tracing results. Scale bars: 200µm (C,D,E-H); 50µm (C',D',E',F',G',H'). dLGN= dorsal lateral geniculate nucleus, FC= Fold Change, OB= Olfactory Bulb, VL=ventral lateral nucleus, VM= ventral medial nucleus, VMP= ventral medial posterior nucleus, VP= ventral posterior, vTel= ventral telencephalon, WT= wild type.

Figure 2

Figure 2. Thalamic axons exhibit abnormal fasciculation as they cross the

prethalamus in CAG^{CreER} Pax6 cKOs. A-L) Transverse (A,B, F-I) and sagittal (C-D,

J-L) sections showing immunofluorescence for L1 and Pax6 in controls (A-E) and

CAG^{CreER} Pax6 cKOs (F-L) at E13.5. White arrows in A,B and E show the point at

which the thalamocortical tract forms a single bundle and exits the diencephalon in controls. Empty arrows in G-I and L point to thalamic axons converging prematurely into big bundles as soon as they cross the thalamic-prethalamic boundary in Pax6 cKOs. Discontinuous lines in A, F-I mark the thalamic-prethalamic boundary. Discontinuous lines in B mark the level of sections in C, D and E. Discontinuous lines in D,E,K and L show the position of the lines used to quantify the number and width of bundles crossing the prethalamus. **M,N)** Examples of the measurements taken for the quantifications of the number and width of axon bundles crossing each line. Individual bundles were identified as single L1-positive structure above a consistent intensity threshold (red lines). **O,P)** Graphs showing the quantification of the number (O) and width (P) of axon bundles crossing each reference line and the statistical significance. Representation of means +/- SEM, n=3 (three embryos analysed from three different litters, five slices per embryo), where *, ** and *** stand for a p-value ≤ 0.05, 0.01 and 0.001 respectively after two-tailed unpaired Student's t-test and N=3. O) We found a significant decrease in the number of bundles crossing all three checkpoints (Th-PTh boundary: p value=0.046, t=2.86, df=4; Mid Pth: p value=0.0085, t=4.82, df=4; low-PTh: p-value=0.012, t=4.39, df=4). P) We detected a big increase in axon bundle width at the Th-PTh border (p-value<<0.001, t=9.12, df=443) and the mid-PTh (p-value<< 0.001, t=7.21, df=350), but no significant change at the low-PTh. **Q)** Schematic summary of the results showing how thalamic axons undergo abnormal and premature fasciculation when crossing the prethalamus in CAG^{CreER} Pax6 cKOs. Scale bars: 200µm. Ctx= cortex, Pth= prethalamus, Th= thalamus, vTel= ventral telencephalon.

Figure 3

Figure 3. Prethalamic pioneer axons belong to the Gsx2 lineage and express

Pax6. (A) EGFP reporter of Cre recombinase activity and L1 staining in Gsx2^{Cre}

embryos at E14.5. **B)** Immunohistochemistry showing Pax6 expression **C)** EGFP

reporter of Cre activity in Zic4^{Cre} embryos at E13.5 and thalamocortical axons

expressing L1. **D)** High power image showing the absence of axonal projections from

Zic4-lineage prethalamic cells towards the thalamus. **E-E'')** Neurons belonging to the

Gsx2 lineage extend parallel projections across the thalamus in close apposition to

L1-positive thalamocortical axons. E' and E'' are a higher magnification of the area

framed in E. **F-H)** Injection of the tracer Neurobiotin in the thalamus at E13.5 resulted

in the retrograde labelling of neurons in the prethalamus (arrow in F). F-G: Double

labelling of GFP and Neurobiotin shows that prethalamic labelled neurons belong to

the Gsx2. H,H': Parallel section of F'' processed for Pax6 immunoshistochemistry

combined with Neurobiotin visualization showing that most prethalamic neurobiotin-

positive neurons are also Pax6 positive (arrows). **I)** Schematic summary. Scale bars:

500μm (A,B,C,E), 250μm (F), 100μm (D, E',G,H). Ctx= cortex, Nb= Neurobiotin,

Pth= prethalamus, Th= thalamus, vTel= ventral telencephalon. All descriptions where

observed in at least three embryos from three different litters.

Figure 4

Figure 4. Disruption of prethalamic pioneer axons in Gsx2^{Cre} Pax6 cKOs

causes abnormal fasciculation but not changes in topography. A-F)

Prethalamic pioneer axons visualized by EGFP reporter in Gsx2^{Cre} embryos are
reduced in Gsx2^{Cre} Pax6 cKOs at E12.5 (A,B), E13.5 (C,D) and E14.5 (E,F) **G-J)**
Use of the DTy54 YAC reporter allele, which labels cells in which the *Pax6* gene is
active, also reveals a reduction in prethalamic pioneer axons crossing the thalamus.
K,L) Injection of the tracer DiI in the thalamus of E13.5 retrogradelly labels cell
bodies in the prethalamus of controls (arrow in K) but not in CAG^{CreER} Pax6 cKOS
(L). Inset in K is a fluorescent immunohistochemistry for Pax6 combined with DiI
visualization showing that the DiI-labelled prethalamic neurons are expressed in the
Pax6 positive area. **M-R)** Immunofluorescence for L1 shows formation of abnormal
big bundles of thalamocortical axons crossing the prethalamus in Gsx2^{Cre} Pax6
cKOs (arrow in N,P,R) with respect to controls (M,O,Q) at E14.5 (M,N), E16.5 (O,P)
and E18.5 (Q,R). **S)** DiI and DiA injection in the cortex of Gsx2^{Cre} Pax6 cKO embryos
shows that the topography of TCAs is not affected in these mutants. **T)** Schematic
summary. Scale bars: 250µm (G,H,S), 100 µm (A-F, I-P). Ctx= cortex, Pth=
prethalamus, Th= thalamus, vTel= ventral telencephalon. All phenotypes where
observed in at least three embryos from three different litters, unless otherwise
stated.

Figure 5

Figure 5. Subsets of thalamocortical axons show a preference to navigate

through different parts of the thalamus. A,B) Immunohistochemistry for L1 shows

that thalamocortical axons exhibit disorganized and erroneous trajectories in the

thalamus of E13.5 CAG^{CreER} Pax6 cKOs. Arrow in B points to one big bundle

deviating from lateral to medial thalamus. **C-J)** Schemas and images of the

transplants using lateral (C-F) or medial (G-J) donor thalamus and grafted into lateral

(C,D,G,H) or medial (E,F,I,J) host tissue. Images show immunofluorescence for GFP

and Pax6. White arrows in J show medial axons turning from lateral to medial

regions of the thalamus.. **K)** Quantification of the navigation of GFP-positive axons through three sectors of the thalamus shows that lateral axons have a strong preference to navigate through lateral thalamus while medial axons show a more variable response. We quantified six thalamic graft experiments with embryos from three different litters (three using lateral explants and three using medial explants). Box plots show the percentage of GFP-positive axons detected on each sector of the thalamus. Five rostrocaudal sections were quantified per explant. Each dot represents an area value of a single section. Scale bars: 100µm. Ctx= cortex, Pth= prethalamus, Th= thalamus, vTel= ventral telencephalon.

Figure 6

Figure 6. Axon guidance molecules and their receptors are altered in the diencephalon of CAG^{CreER} Pax6 cKOs. A-I') Expression pattern of *Ntn1* and *Sema3a* mRNA and their receptors *Unc5c* and *Plxna1* in the E13.5 diencephalon of controls (A-E') and CAG^{CreER} Pax6 cKOs (F-I'). C and H show two views of 3D reconstructions of the expression pattern of *Ntn1* and *Sema3a* from transverse ISH sections. **J,K)** Schematic model of the changes in *Ntn1* and *Sema3a* expression in the thalamus of controls (J) versus CAG^{CreER} Pax6 cKOs (K) and how we hypothesize might affect the guidance of different subsets of thalamic axons. In controls (J), lateral thalamic neurons express Unc5c and Plxna1, being repelled by both guidance cues, which direct their axons out from the thalamus and towards the prethalamus in a straight trajectory. In Pax6 cKOs (K) some lateral thalamic neurons lose their *Unc5c* expression but maintain their *Plxna1* expression, losing their repulsion to *Ntn1* but maintaining it for *Sema3a* and making their axons deviate towards medial thalamic regions. Scale bars: 500µm (A,B,D-E,F,G,I,J), 100µm (D',I'). Ctx= cortex, Pth= prethalamus, Th= thalamus, vTel= ventral telencephalon. All described patterns were observed at least in three embryos belonging to three different litters for each genotype, unless otherwise stated.

1014 **Figure 7**

1015

Figure 7. Subsets of lateral thalamocortical axons from CAG^{CreER} Pax6 cKOs

deviate medially when confronted with control thalamus. A) Schema of the slice

culture transplant experiment in which lateral thalamus from GFP-positive Pax6

cKOs embryos (CAG$^{CreER\,+}$; Pax6 $^{fl/fl}$) was grafted into lateral thalamus of control

GFP-negative littermates (CAG$^{CreER\,-}$; Pax6 $^{fl/fl}$). **B)** Subsets of axons from Pax6 cKO

lateral thalamus deviate from their normal trajectory towards medial thalamic regions

(red arrows). 11 embryos from four different litters were used. A total of 22 (bilateral)

graft experiments were performed. We observed medially deviated axons in all

cases. **C)** Quantification (four grafts from embryos from three different litters) showed

that Pax6 deficient lateral thalamic axons navigate across three delimited sectors,

unlike their control counterparts which showed a strong preference for the lateral

sector (Fig. 5K). Box plots show the percentage of GFP-positive axons detected on

each sector of the thalamus. Five rostrocaudal sections were quantified per explant.

Each dot represents an area value of a single section. Scale bar: 100µm. Lat= lateral

aspect of diencephalon, Med= medial aspect of diencephalon,Th=thalamus,

PTh=prethalamus

1032