A versatile bulk electrotransfection protocol for mouse embryonic fibroblast and iPS

cells

Shahin Eghbalsaied[1,2*], Iqbal Hyder[1], Wilfried A. Kues[1]

[1]Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Mariensee, Neustadt am

Rübenberge, Germany

[2]Department of Animal Science, Isfahan (Khorasgan) branch, Islamic Azad University, Iran.

[*]**Corresponding author: Shahin.eghbalsaied@fli.de , Shahin.eghbal@khuisf.ac.ir**

Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Mariensee, Neustadt am

Rübenberge, Germany.

Postal address: Höltystr. 10, 31535 Neustadt am Rübenberge, Germany.

Telephone: +49 5034 871 5120

Fax: +49 5034 871 5101

Abstract

A square-wave pulsing protocol was developed using OptiMEM-GlutaMAX for high efficient transfection of mouse embryonic fibroblast (MEF) and induced pluripotency stem (iPS) cells. An electrotransfection efficiency of > 95% was repeated for both MEF and iPS cells using reporter-encoding plasmids. The protocol was very efficient for plasmid size ranging from 6.2 to 13.5 kb. A high rate of targeted gene knockout (> 95 %) was produced in Venus transgenic cells using indels formation. Targeted deletions in the Venus transgene were performed by co-electroporation of two gRNA-encoding plasmids. In conclusion, this plasmid electrotransfection protocol is straight-forward, cost-effective, and efficient for CRISPRing mouse primary cells.

Keywords: CRISPR/Cas9, Electrotransfection, Targeted deletion, Large plasmids, OptiMEM-GlutaMAX

Background

The CRISPR (clustered regularly interspaced short palindromic repeats)–Cas9 (CRISPR-associated 9) protein nuclease system is a straight-forward, versatile, highly efficient, and nascent tool for genome editing of various organisms. Using 'all-in-one' expression vectors containing expression cassettes for guide RNAs and Cas9 nuclease/nickase it is possible to conduct CRISPR/Cas9 studies through a low-cost and straight-forward approach (1). However, the efficiency of gene editing using plasmid-based delivery methods remains relatively low, which subsequently increased attentions toward using alternate methods of employing the Cas9 protein and/or guideRNAs and using ribonucleoproteins (RNPs) (2, 3).

Basically, gene transfer into cells can be achieved via electroporation, lipofection, or viral transduction. Although viral gene transfer is very efficient, it requires time, skilled staff, and high levels of safety issues, whereas it has a limitation in the insert size and more importantly enhanced risks for clinical researches (4). On the other side, lipofection suffers from a low efficiency. Electroporation is an approach to instantly create several pores in the cell membrane using a burst electric pulse and to mediate the transfer of micro- and macro-molecules into cells, embryos, tissues, and organs (5). From the early papers on electroporation (6), it has been evident that gene transfer via electroporation is simple, easily applicable, and also efficient compared to lipofection (6). Although electroporation has been widely accepted as the main gene transfer tool in various laboratories around the world, the underlying mechanism has not been completely understood (7). Therefore, optimization of various factors, such as the electroporation medium, cuvette type (path width, length, and high), as well as the pulsing method which includes the amount, number, duration, and interval of pulses, is needed to have a high electrotransfection efficiency particularly in primary cells. Based on the above-mentioned electroporation parameters various types of electroporation-based devices, such as nucleofection (8), nano-electroporation (9),

mechanical-electrical approach (7), and microfluidic membrane deformation (2) as well as

new generation of electroporator devices (10), have been invented. However, still there is a

huge room to improve the electroporation efficiency to transfer exogenous DNAs into cells,

specifically for large plasmids (7, 11).

Here, we introduced a highly efficient electrotransfection method for both mouse embryonic

fibroblast (MEF) and induced pluripotent stem (iPS) cells based on the square-wave pulsing

method using a Bio-Rad electroporation device. We developed the technology using different

types of reporter plasmids and confirmed its very high efficiency for making gene knockout

(KO) which was induced by both indels and targeted deletions using single or double gRNAs

through CRISPR/Cas9 plasmids, respectively.

Results

Optimization of electrotransfection protocol for mouse iPS and EF cells

Three media were compared for electrotransfection efficiency. Initial results of transfection of

mouse iPS cells with Bio-Rad electroporation buffer, PBS, and OptiMEM-GlutaMAX using a

10 ms single pulse of 200 V resulted in 84.7 % transfection rate in OptiMEM-GlutaMAX

medium compared to 17.9 % in Bio-Rad buffer (Figure S1). More importantly, the number of

cells and colonies 24 h after the electroporation was considerably higher in OptiMEM-

GlutaMAX compared to both Bio-Rad and PBS buffers ($p<0.05$).

Then, we optimized pulse voltage, number, and duration as well as the medium temperature

for electrotransfection of iPS cells based on the OptiMEM-GlutaMAX medium (Figures S2).

We developed an electroporation protocol using the square-wave pulsing program of 250 V, 2

pulses, each 10 ms length, and 10 s interval in 4 mm cuvettes. Electroporations were carried

out at room temperature with 20 µg DNA in 250 µl of OptiMEM-GlutaMAX. Applying this

protocol for transient expression of mCherry and Venus reporters in mouse iPS cells showed a

95.5 and 92.8 % electrotransfection efficiency, respectively (Figure 1). Then, we implemented

the same protocol with a higher voltage (300 V instead of 250 V) for MEF cells and reached

96.0 and 95.1 % expression of mCherry and Venus transgenes, respectively, 24 h after the

electroporation (Figure 1). The electrotransfection efficiency was reproducible using mCherry

reporter in three MEF and three iPS cell lines.

Finally, using the optimized electrotransfection protocol the transient transfection efficiency

of mouse iPS and EF cells was assessed in 250 µl of either of Bio-Rad, OptiMEM-

GlutaMAX, and PBS media (Figure 2). Viability of iPS cells was significantly lower in Bio-

Rad (22.0 %) and PBS (3.2 %) compared to the OptiMEM-GlutaMAX medium (78.2 %).

Also, the electrotransfection efficiency of iPS cells was more than two-fold higher in

OptiMEM-GlutaMAX (99.3 %) than that of Bio-Rad (42.7 %) and PBS media (47.2 %). The

electrotransfection efficiency of mouse EF cells was 30 % higher in OptiMEM-GlutaMAX

compared to the Bio-Rad medium (p-value <0.05). In addition, the viability of mouse EF cells

averaged 78 % in the OptiMEM-GlutaMAX group.

Efficient knockout of Venus transgene using indels by Cas9/gRNA encoding plasmids

MEF cells carrying a single-copy of the Venus transgene were transfected by plasmids

encoding a gRNA and Cas9 protein using the optimized electroporation method (Figure S3).

Venus expression was not reduced by three gRNAs which targeted the upstream region of the

Venus transgene although they induced indels in the targeted sites (Figure 3). However,

targeting the beginning and ending parts of the transgene, ranging from +36 to +554 bp of the

cDNA, could knockout the transgene in > 90 % of the electroporated cells (Figure S3). The

Venus KO efficiency was maximized using gRNA+100 which was complimentary to the

beginning part of the cDNA, so that only 2 % of cells maintained the functional Venus. The

Venus knockout results were confirmed by fluorescent microscopy, as well as FACS analysis

and DNA sequencing. The knockout efficiency of Venus transgene was not affected by the

puromycin treatment; 99 % vs. 93 % for 100 gRNA with and without puromycin selection,

respectively (Figure S4). Although electrotransfection results were very promising using

OptiMEM-GlutaMAX medium, substituting the medium with the standard OptiMEM

supplemented with glutamine was completely inefficient for making Venus KO (Figure S5).

Moreover, electrotransfection of MEF cells with pSGD-Lys-72 plasmid (13.5 kb) encoding

human lysozyme, puromycin, and gRNA-72 compared to the modified pX459-gRNA-72

showed a similar cell viability following a one-day selection against puromycin antibiotic

(Figure S6).

High efficient deletion of Venus fragments using dual gRNAs

Following making Venus knockouts using indels, we were interested in testing the efficiency

of the electrotransfection protocol for making deletions inside of the target gene using two

gRNAs. Simultaneously, MEF cells carrying a single-copy of Venus were co-electroporated

using 15 combination of two gRNAs. The result of end-point PCR showed deletion of

expected fragments ranging from 398 to 748 bp in the Venus transgene using all of gRNAs

pairs (Figure 4). Only shortened Venus amplicons were amplified by a primer set (Figure S7),

while using another primer set the original transgene with no deletion was also detectable

(Figure 4). In overall, the deletion efficiency was more evident using the combinations of

gRNA+676, although gRNA+554 and +518 were also efficient for making deletions. Results

of the DNA sequencing confirmed the fragment deletion in co-transfected groups (Figure S8).

Discussion

Nowadays, implementation of the nascent CRISPR/Cas9 technology is a straightforward

approach to make genetic engineering in any organism. Electroporation of large plasmids (> 6

kb) is almost inefficient and associating with a low viability in primary cells (11). In the

current study, we identified OptiMEM-GlutaMAX as the best medium for electroporation of

MEF and iPS cells. The electrotransfection efficiencies were quite high throughout the initial

setup condition with various pulsing conditions. A > 95 % electrotransfection rate was

achieved with the optimized pulsing condition. Even the rate of transient expression of

reporter genes, as well as the cell viability using the OptiMEM-GlutaMAX in the current

study were as high as that of newly reported mechanical-electrical (7) and nanopore-electrical

approaches (12). We used plasmids with various sizes, from 6.2 to 13.5 kb, and found a high

transfection rate and cell viability irrespective of the plasmid size. It has been evident that

electrotransfer of large plasmids (> 6 kb) associates with a very high cell toxicity, while the

cell transfection rate was also low (11). However, in our study, electrotransfection of MEF

cells using a 13.5 kb was as efficient as a 9.2 kb plasmid, whereas the cell viability remained

the same. Therefore, the electrotransfection protocol for primary cells in the current study

could overcome impediments of hard to transfect cells, as well as high toxicity, and low

transfection rate using large plasmids (11). So far, invention of minicircles was considered as

the main solution to overcome this problem (13). However, converting a plasmid to a

minicircle is time- and cost-consuming and it does not solve the problem of co-transfection of

plasmids (11). Nonetheless, our results showed that electrotransfection of large plasmids is

possible with a high efficiency and cell viability in OptiMEM-GlutaMAX medium with no

need for a long post-pulse recovery time, as suggested by Lesueur et al. (11). It has been

well-documented that iPS cells are resistant and vulnerable to electroporation (14). The

current study showed that comparing to the Bio-Rad electroporation buffer, the OptiMEM-

GlutaMAX was compatible with iPS cells in terms of transfection rate and cell viability.

However, unlike reported by Potter et al. (15), conducting the electroporation in ice-cold

medium did not improve the transfection efficiency in the current study, although the

transfection efficiency of ice-cold medium was almost independent of pulsing condition.

Then, we were interested to know if the transient expression is sufficient for genome

engineering using the CRISPR/Cas9 technology. Following optimization of the

electrotransfection protocol for reporter-encoding transgenes, we implemented the same

protocol for transfer of plasmids encoding Cas9 and gRNA aiming to knockout the Venus

transgene in MEF cells carrying a single-copy of Venus. The highest rate of Venus knockout

through indels formation (> 95 % using gRNA+100) was as high as the electrotransfection

efficiency of reporter-encoding plasmids. The indels rate using plasmids in the current study

was higher than reported results by using both plasmid and RNP electrotransfection(16-18).

Following making Venus knockouts using indels, we were interested to test the efficiency of

the electrotransfection protocol for making deletions inside of the target gene using double

gRNAs. Deletion of expected fragments in the Venus transgene by co-electroporation of two

plasmids encoding two different gRNAs was confirmed using endpoint-PCR as well DNA

sequencing. The deletion efficiency obtained in the current study is higher than previous

results using dual gRNAs and Cas9 protein (19) as well as Cas9-encoding plasmids using the

BTX electroporation buffer in mouse cell lines (20).

Conclusion

In this study, we introduced a highly efficient electrotransfection method for both MEF and

iPS cells based on the square-wave pulsing method using OptiMEM-GlutaMAX medium.

This plasmid-based delivery method could induce > 95 % transfection efficiency for

fluorescent reporter genes. Electrotransfection of Cas9/gRNA encoding plasmids caused in

approximately 98 % knockouts of Venus transgene. Apart from indels creation, targeted

deletions in the Venus transgene were achieved using the co-electroporation of two gRNA-

encoding plasmids. Therefore, the developed protocol in the current study can be an alternate

to using Cas9 protein and RNPs.

Materials and methods

Materials

All plastic consumables including cell culture flasks and plates, tubes, and filter tips have been purchased from SARSTEDT AG & Co. (Germany). Chemical reagents were purchased with the following information: Dulbecco's phosphate buffered saline without calcium chloride and magnesium chloride (#D6662-10X1L, SIGMA-ALDRICH, Germany); Opti-MEM 1X + GlutaMAX, reduced serum medium (#1854076, Gibco, Life Technologies, Germany); DMEM high glucose (4.5 g/l) w/o L-glutamine (#DMEM-HXA, Capricorn Scientific GmbH, Germany); L-glutamine for cell culture (#A3704,0100, Applichem, Germany); MEM nonessential amino acids solution (100X) (#NEAA-B, Capricorn Scientific GmbH, Germany); 2-mercaptoethanol (#BCBS5481, SIGMA-ALDRICH, Germany); penicillin/streptomycin solution (100x) (#PS-B, Capricorn Scientific GmbH, Germany); trypsin-EDTA (10X) (#L11-003, GE Healthcare, PPA Laboratories GmbH, Austria); fetal bovine serum (#10270-106, Gibco, ThermoFisherScientific, Germany); dimethyl sulfoxide (#D4540-500ML, SIGMA-ALDRICH, Germany); LIF (hBA-FL) (#sc-4377, Santa Cruz Biotechnology, Germany); sodium pyruvate (#P2256, SIGMA-ALDRICH, Germany); gelatin from bovine skin (#G939-100G, SIGMA-ALDRICH, Germany); and Hoechst 33342 (#62249, Thermo Scientific, Germany). The Gene Pulser Xcell™ system with CE Module from BIO-RAD (Germany) was used with 4 mm electroporation cuvettes (#748052, Biozym, Germany).

Plasmids

Plasmids carrying *Sleeping Beauty* (SB) system

The *Sleeping Beauty* transposase was encoded in pCMV-T7_SB100X (4756 bp) which contains the hyperactive variant 100 of SB under the CMV promoter (21). Two reporter plasmids, pT2-Venus (6301 bp) encoding the Venus fluorescent marker under CAGGS

promoter and pT2-mCherry (7756 bp) encoding mCherry, were used (21). In addition, a 13.5

kb plasmid, pSGD-Lys-72, was also used as a large plasmid which encodes human lysozyme

under CAGGS promoter followed by Cas9 protein which was separated from puromycin

resistance by a T2A peptide and expressed under CMV promoter, and gRNA-72 under U6

promoter. All pT2-Venus, pT2-mCherry, and pSGD-Lys-72 plasmids contained inverted

terminal repeats (ITRs) of the SB transposase. The SB reprogramming transposon involved a

CAGGS promoter driven cassette containing Oct4, Sox2, Klf4, and c-Myc, which were

separated by sequences for coding the self-cleaving 2A peptides, and flanked by SB-ITRs

(22).

Plasmid carrying the CRISPR/Cas9 system

For the CRISPR/Cas9 study, we modified pX459 plasmid (9151 bp) which encodes a Cas9

protein followed by puromycin resistance under CAGGS promoter/enhancer, as well as the

gRNA scaffold under the U6 promoter. All gRNAs were synthetized and cloned into the

backbone vector based on the Franham protocol (23). Briefly, gRNAs were selected to target

Venus promoter or open reading frame (ORF) via CRISPOR software, available online

(http://crispor.tefor.net/) (24). The list of gRNAs is available in Table 1. The 100 µM forward

and reverse oligos were annealed in 10 µl reaction volume by incubation in a water bath

containing pre-boiled water and letting it to cool down to the room temperature. The pX459

plasmid was digested with High Fidelity BbsI-HF (NEB #R3539) at 37°C for 10 min

followed by gel purification using NucleoSpin Gel and PCR Clean-up Midi kit (#740986.20,

MACHEREY/NAGEL). The purified fragment was kept at -20°C for further applications.

Ligation of the annealed oligo-duplex with the digested pX459 was carried out as follow:

diluted oligo-duplex (1:20 ratio from the 10 µM source) (1 µl), digested pX459 vector (50

ng), 10× T4 DNA ligase buffer (2 µl), and T4 DNA ligase (1 µl) in a 20 µl final reaction. The

ligation reaction was incubated at room temperature for 1 hour. Transfection of the ligation

mixture was carried out into NEB 5-alpha Competent E. coli (Cat. # C2987I) following

incubation of 10 µl of the ligation reaction with the thawed competent cells on ice for 20 min

and then at 37°C for 5 min. Then, 400 µl of SOC medium was added into the transformation

tube, incubated at 37° C for 30 min, plated on Agar plates supplemented with 100 µg/ml

ampicillin, and incubated at 37°C overnight. From the cultured plate, 10 colonies were picked

and each was cultured in 3 ml LB medium followed by miniprep plasmid extraction (Genejet

Plasmid miniPrep kit, #K0502). The extracted plasmids were simultaneously digested by BbsI

and EcoRV FastDigest restriction enzymes (Fermentas). Plasmids with the expected bands of

6244 and 2928 bp were sent for sequencing.

Cell culture and electroporation

Gelatinization of cell culture plates

A 0.1 % gelatin solution was prepared by stirring 250 ml PBS and 2.5 g gelatin for one hour

without heating, and the solution was then autoclaved. Five milliliter of the 0.1 % gelatin was

poured into a 6-well plate and aspirated after 1 min. Plates were left to be dried under the

laminar flow hood for 30 min.

Cell lines

We used mouse embryonic fibroblast cells (MEFs), carrying either no- or a single-copy of the

Venus transgene (Venus +/-). Mouse embryonic fibroblast cells were isolated from day 11

embryos of the Venus -/- (wild type) or Venus +/- embryos . Embryos were beheaded,

eviscerated, sliced, and cultivated with 5 ml fibroblast medium (DMEM with 10% FBS, L-

glutamine, penicillin-streptomycin, nonessential amino acids, and β-mercaptoethanol) in T75

flasks. After 3 days of the incubation, propagated cells were trypsinized, passaged, and used

for cell culture and transfection. Moreover, mouse iPS cells were derived from MEFs with no-

or a single-copy of the Venus, respectively (22). Briefly, MEFs were electroporated with

plasmids containing the 4-factor reprogramming transposon and the helper plasmid containing

the hyperactive SB transposase (22). Electroporated cells were cultured in gelatinized 6-well

plates and iPS medium. Presumptive iPS colonies were picked under microscope, and plated

into individual wells of 96-well plates containing trypsin. Trypsin was neutralized with

DMEM and 10% FCS, and the cells suspension was dispensed into individual wells of

gelatinized 96-well plates (22).

284
Cell Electroporation

To assess the transfection efficiency using reporter plasmids, MEFs and iPS cells with no-

copy of Venus reporter were used. Both MEFs and iPS cells were cultured in 6-well plates.

From the highly confluent culture, cells from each well of a 6-well plate were used for each

electroporation reaction using both iPS and MEFs cells. Cells were washed once with 2 ml

PBS and then trypsinized with 200 µl trypsin-EDTA. Trypsin activity was inhibited by 4 ml

PBS per well, and then was removed through centrifugation at 1,000 rpm for 3 min. The cell

pellet was resuspended into 250 µl of the electroporation medium. In an initial experiment,

the highest transfection efficiency of iPS cells were reached by OptiMEM-GlutaMAX

medium (Figure S1). Then, the effect of voltage and medium temperature was optimized for

iPS cells (Figures S2 and S3). Finally, we reached the following optimized condition for

electroporation of mouse iPS cells. Twenty micrograms of plasmid DNA were mixed with

cells, and the cell-plasmid mixture was transferred into the 4 mm cuvettes, and underwent the

following electroporation program: The square-wave protocol with 250 V, each 10 ms pulse

length, 2 pulses, and 10 s pulse interval. MEFs underwent the same electroporation program

which was optimized for iPS cells, except conduction of a 300 V. After the electroporation

process, cells were transferred into the culture medium in 6-well plates and incubated at 37°C

and 5 % CO_2.

Venus knockout using one and two gRNAs

For the CRISPR/Cas9 experiment, we used MEFs cells with a single-copy of Venus for

making KO-Venus cells. The above-mentioned optimized protocol for cell electrotransfection

was also used for Cas9/gRNA encoding plasmids. Knockout of Venus was carried out either

by indels using 20 µg of a modified pX459 plasmid or by deletions via co-electroporation of

two modified pX459 plasmids (each 20 µg per reaction). To consider the deletion efficiency

of various sizes, three gRNAs which target the end part of Venus transgene (gRNA +518,

+554, and +676) were pair-wisely co-electroporated with five gRNAs which targeted the

beginning part of Venus (gRNA -72, -69, +36, +100, and +121). A pair of primers which

cover all the gRNA complementary sites were used for the end-point PCR. DNA was

extracted from each treated group using salting-out method. Briefly, confluent cells in each

well of 6-well plates were trypsinized with 250 µl Trypsin-EDTA, directly transferred into a

1.5 ml tube containing 1 ml of cell lysis buffer (0.2 mg/ml proteinase K, 150 mM NaCl, 10

mM Tris, 10 mM EDTA, and 0.1% sodium dodecyl sulfate (SDS)) (25), and were incubated

overnight at 55 °C in a shaker-incubator. Then, 500 µl saturated NaCl was added into each

tube, converted for 5-10 times, and centrifuged with the high speed at room temperature.

Supernatants were transferred into two new tubes and underwent ethanol precipitation with

absolute ethanol followed by twice washing with 70 % ethanol using centrifugation with the

high speed at 4°C. The DNA pellet was dissolved in 200 µl distilled water and kept at -20 °C

till further usages.

Fluorescent microscope imaging and flow cytometry analysis

For the fluorescent imaging, cells in each well of 6-wll plate were washed once with PBS, the

medium was replaced with the transparent OptiMEM (without phenol) supplemented with

Hoechst 33342 (2: 10,000 ratio from 1 mg/ml stock solution), and incubated at 37°C for 30

min. Transfection efficiency for mCherry and Venus reporters were assessed either by FACS

machine or fluorescent microscopy. Cell viability was defined as cell numbers in the

electroporated group divided by the cell number in the 0 voltage group (negative control).

Knockout efficiency of various gRNA sequences targeting the Venus reporter was assessed

10 days after the electroporation. A flow cytometer, MACSQuant® Analyzer, was also used

to assess cell transfection rate. We used Blue 488 nm in B1 channel with 525/550 nm filter

and Yellow 461 nm Y2 channel with 615/20 nm filter for detection of Venus and mCherry

proteins, respectively.

Statistical analysis

Means comparison was carried out using the least significant difference (LSD) test (p-values

0.05). All experiments were carried out at least three times.

Declarations

Ethics approval and consent to participate

The German animal welfare law does not require an approval for humanely sacrifice of

animals for the purpose of organ biopsies. The number of sacrificed animals are annually

reported to the regulating authorities.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Competing interests

Authors have no conflict of interest to declare.

Funding

This project was funded by the grant number: Ref 3.4 - IRN - 1191261 - GF-E from the Alexander von Humboldt Foundation, Germany.

Author's contributions

Conceived the idea and designed the experiment: SE and WK; Carried out the experiments: SE, IH, and WK; Wrote the manuscript: SE; all authors read and confirmed the manuscript content.

Acknowledgements

Authors would like to thank Antje Frenzel for conducting the FACS analysis.

References

1.   Sakuma T, Nishikawa A, Kume S, Chayama K, Yamamoto T. Multiplex genome engineering in human cells using all-in-one CRISPR/Cas9 vector system. Scientific reports. 2014;4:5400.
2.   Han X, Liu Z, chan Jo M, Zhang K, Li Y, Zeng Z, et al. CRISPR-Cas9 delivery to hard-to-transfect cells via membrane deformation. Science advances. 2015;1(7):e1500454.
3.   Liu C, Zhang L, Liu H, Cheng K. Delivery strategies of the CRISPR-Cas9 gene-editing system for therapeutic applications. Journal of controlled release. 2017;266:17-26.
4.   Gresch O, Engel FB, Nesic D, Tran TT, England HM, Hickman ES, et al. New non-viral method for gene transfer into primary cells. Methods. 2004;33(2):151-63.
5.   Weaver JC, Smith KC, Esser AT, Son RS, Gowrishankar T. A brief overview of electroporation pulse strength–duration space: A region where additional intracellular effects are expected. Bioelectrochemistry. 2012;87:236-43.
6.   Neumann E, Schaefer-Ridder M, Wang Y, Hofschneider P. Gene transfer into mouse lyoma cells by electroporation in high electric fields. The EMBO journal. 1982;1(7):841-5.
7.   Ding X, Stewart MP, Sharei A, Weaver JC, Langer RS, Jensen KF. High-throughput nuclear delivery and rapid expression of DNA via mechanical and electrical cell-membrane disruption. Nature biomedical engineering. 2017;1(3):0039.
8.   Lakshmipathy U, Pelacho B, Sudo K, Linehan JL, Coucouvanis E, Kaufman DS, et al. Efficient transfection of embryonic and adult stem cells. Stem cells. 2004;22(4):531-43.
9.   Yang Z, Chang L, Chiang C-l, James Lee L. Micro-/nano-electroporation for active gene delivery. Current pharmaceutical design. 2015;21(42):6081-8.
10.   Kaneko T, Sakuma T, Yamamoto T, Mashimo T. Simple knockout by electroporation of engineered endonucleases into intact rat embryos. Scientific reports. 2014;4:6382.
11.   Lesueur LL, Mir LM, André FM. Overcoming the specific toxicity of large plasmids electrotransfer in primary cells in vitro. Molecular Therapy-Nucleic Acids. 2016;5:e291.
12.   Cao Y, Ma E, Cestellos-Blanco S, Zhang B, Qiu R, Su Y, et al. Nontoxic nanopore electroporation for effective intracellular delivery of biological macromolecules. Proceedings of the National Academy of Sciences. 2019;116(16):7899-904.

13.   Joubert V, André FM, Schmeer M, Schleef M, Mir LM. Increased efficiency of minicircles
versus plasmids under gene electrotransfer suboptimal conditions: an influence of the extracellular
matrix. Minicircle and Miniplasmid DNA Vectors: The Future of Nonviral and Viral Gene Transfer.
2013:215-25.
14.   Qian K, Huang CL, Chen H, Blackbourn IV LW, Chen Y, Cao J, et al. A simple and efficient
system for regulating gene expression in human pluripotent stem cells and derivatives. Stem Cells.
2014;32(5):1230-8.
15.   Potter H, Heller R. Transfection by electroporation. Current protocols in immunology.
2017;117(1):10.5. 1-.5. 9.
16.   Liang X, Potter J, Kumar S, Zou Y, Quintanilla R, Sridharan M, et al. Rapid and highly efficient
mammalian cell engineering via Cas9 protein transfection. Journal of biotechnology. 2015;208:44-53.
17.   Jacobi AM, Rettig GR, Turk R, Collingwood MA, Zeiner SA, Quadros RM, et al. Simplified
CRISPR tools for efficient genome editing and streamlined protocols for their delivery into
mammalian cells and mouse zygotes. Methods. 2017;121:16-28.
18.   Liang X, Potter J, Kumar S, Ravinder N, Chesnut JD. Enhanced CRISPR/Cas9-mediated precise
genome editing by improved design and delivery of gRNA, Cas9 nuclease, and donor DNA. Journal of
biotechnology. 2017;241:136-46.
19.   Kraft K, Geuer S, Will AJ, Chan WL, Paliou C, Borschiwer M, et al. Deletions, inversions,
duplications: engineering of structural variants using CRISPR/Cas in mice. Cell reports.
2015;10(5):833-9.
20.   Canver MC, Bauer DE, Dass A, Yien YY, Chung J, Masuda T, et al. Characterization of genomic
deletion efficiency mediated by clustered regularly interspaced short palindromic repeats
(CRISPR)/Cas9 nuclease system in mammalian cells. The Journal of biological chemistry.
2017;292(6):2556.
21.   Ivics Z, Mátés L, Yau TY, Landa V, Zidek V, Bashir S, et al. Germline transgenesis in rodents by
pronuclear microinjection of Sleeping Beauty transposons. Nature protocols. 2014;9(4):773.
22.   Talluri TR, Kumar D, Glage S, Garrels W, Ivics Z, Debowski K, et al. Non-viral reprogramming of
fibroblasts into induced pluripotent stem cells by Sleeping Beauty and piggyBac transposons.
Biochemical and biophysical research communications. 2014;450(1):581-7.
23.   Guo Y, Perez AA, Hazelett DJ, Coetzee GA, Rhie SK, Farnham PJ. CRISPR-mediated deletion of
prostate cancer risk-associated CTCF loop anchors identifies repressive chromatin loops. Genome
biology. 2018;19(1):160.
24.   Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, et al. Evaluation of off-
target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR.
Genome biology. 2016;17(1):148.
25.   Minematsu T, Sugiyama M, Tohma Y, Tajima A, Kanai Y. Simplified DNA extraction methods
for sexing chick embryos. The Journal of Poultry Science. 2004;41(2):147-54.

Figure Legends:

Figure 1. Electrotransfection efficiency of mouse embryonic fibroblast (EF) and iPS

cells. In each electroporation reaction, 20 μg of either pT2-Venus and pT2-mCherry which

encode Venus and mCherry proteins, respectively, was pre-mixed with cells and underwent

electroporation. A) transfection efficiency and B) fluorescent expression of Venus and

mCherry 36 h after electroporation. The following electroporation program was used: square-

wave protocol with 300 V voltage, each 10 ms pulse length, 3 number of pulses, 10 s pulse

interval, and 4 mm cuvette. Transfected cells for mCherry and Venus are depicted by red and

green bars, respectively. Scale bars equal 100 µm. Results are means and standard deviation

(SD).

Figure 2. Cell electrotransfection and viability rates using Bio-Rad buffer, OptiMEM-

GlutaMAX, and PBS. Electrotransfection efficiency of mouse iPS (A and B) and MEF cells

(C and D) by BioRad electroporator. In each electroporation reaction, 20 µg of the reporter

plasmid (encoding either Venus or mCherry) was pre-mixed with cells and underwent

electroporation using the square-wave protocol with 250 and 300 V (for iPS and MEF cells,

respectively), 2 pulses, each 10 ms length, 10 s interval, and 4 mm cuvette. The reporter

expression was assessed 36 h after electroporation under a fluorescent microscope. White and

black bars in A and C are electrotransfection efficiency and cell viability, respectively. Bars

with different letters with are significantly different (p-value < 0.05). Scale bars equal 100

µm. Results are means and standard deviation (SD).

Figure 3. Knockout of Venus transgene by making indels using gRNA-encoding

plasmids. MEF cells carrying a single-copy of the Venus transgene were electrotransfected

with different gRNA-encoding plasmids (9.1 kb length). Induction of indels using gRNA-72

had no effect on the Venus expression, whereas gRNA+100 could completely remove the

Venus signal. Electroporated cells were selected against puromycin and screened 10 days

after the electroporation. Cells were stained with Hoechst 33342 and the efficiency of Venus

knockout was assessed (A and B). Histoplots of the FACS results are shown with the

percentage of Venus positive cells is embedded for each group (C). Results were confirmed

by sequencing of the amplified Venus transgene (D). NK and PK are negative and positive

controls, respectively. The following electroporation program was used: square-wave protocol

with 300 V voltage, each 10 ms pulse length, 3 number of pulses, 10 s pulse interval, and 4

mm cuvette. Scale bar equals 100 μm.

Figure 4. Targeted deletions of Venus transgene by co-electroporation of two gRNA-

encoding plasmids. End-point PCR showed deletion of 15 different fragments from the

original amplicon (1204 bp) using pairwise combination of gRNAs targeting the beginning (-

72, -69, +36, +100, and +121) and the ending part (+518, +554, and +676) of Venus. Primer

Venus Forward and Reverse1 amplified both the shortened and original fragments in the co-

electrotransfected groups.

100 bp	gRNA +676					gRNA +554					gRNA +518					Venus+/-	pVenus	-Ve	1 kb
Ladder	-72	-69	+36	+100	+121	-72	-69	+36	+100	+121	-72	-69	+36	+100	+121	Cells		Control	Ladder