Original Research Article

Lipids in xylem sap of woody plants across the angiosperm phylogeny

H. Jochen Schenk[1],[*], Joseph M. Michaud[1], Kerri Mocko[1], Susana Espino[1], Tatiana

Melendres[1], Mary R. Roth[2], Ruth Welti[2], Lucian Kaack[3], and Steven Jansen[3]

[1]*Department of Biological Science, California State University Fullerton, 800 N. State College*

Boulevard, Fullerton, CA 92831, USA; [2]*Kansas Lipidomics Research Center; Division of*

Biology; Kansas State University; Manhattan, KS 66506, USA; [3]*Institute of Systematic Botany*

and Ecology, Ulm University, Albert-Einstein-Allee 11, D–89081, Ulm, Germany.

[*]*For correspondence. E-mail jschenk@fullerton.edu*

Short-running title: Lipids in xylem sap of angiosperms

Keywords: angiosperms, apoplast, cohesion-tension theory, galactolipids, lipidomics,

phospholipids, vessel volume, xylem, xylem sap, *Laurus nobilis*

1
2   **Abstract**
3
4   16   Lipids have been observed attached to lumen-facing surfaces of mature xylem conduits of
5
6   17   several plant species, but there has been little research on their functions or effects on water
7
8   18   transport, and only one lipidomic study of the xylem apoplast. Therefore, we conducted
9
10 19   lipidomic analyses of xylem sap from woody stems of seven plants representing six major
11
12 20   angiosperm clades, including basal magnoliids, monocots, and eudicots, to characterize and
13
14 21   quantify phospholipids, galactolipids, and sulfolipids in sap using mass spectrometry. Locations
15
16 22   of lipids in vessels of *Laurus nobilis* were imaged using TEM and confocal microscopy. Xylem
17
18 23   sap contained the galactolipids di- and mono-galactosyldiacylglycerol (DGDG and MGDG), as
19
20 24   well as all common plant phospholipids, but only traces of sulfolipids, with total lipid
21
22 25   concentrations in extracted sap ranging from 0.18 to 0.63 nmol / mL across all seven species.
23
24 26   Contamination of extracted sap from lipids in cut living cells was found to be negligible. Lipid
25
26 27   composition of sap was compared to wood in two species and was largely similar, suggesting
27
28 28   that sap lipids, including galactolipids, originate from cell content of living vessels. Seasonal
29
30 29   changes in lipid composition of sap were observed for one species. Lipid layers coated all lumen-
31
32 30   facing vessel surfaces of *Laurus nobilis*, and lipids were highly concentrated in inter-vessel pits.
33
34 31   The findings suggest that apoplastic, amphiphilic xylem lipids are a universal feature of
35
36 32   angiosperms. The findings require a reinterpretation of the cohesion-tension theory of water
37
38 33   transport to account for the effects of apoplastic lipids on dynamic surface tension and hydraulic
39
40 34   conductance in xylem.
41
42
43 35
44
45 36
58   2
59
60

INTRODUCTION

The lipidome of the plant apoplast has been characterized as a black box (Misra, 2016), because

there have been almost no lipidomic analyses of cell walls, intercellular spaces, and xylem

conduits and fibers, except for a few studies of surface lipids involved in the making of suberin,

cutin, and waxes. Studies of lipids in xylem conduits, including vessels, are especially limited.

Wagner *et al*. (2000) observed under transmission electron microscopy (TEM) a dark layer lining

vessel and tracheid surfaces of the resurrection plant *Myrothamnus flabellifolia* Welw.

(Myrothamnaceae) and after further studies concluded that it was composed of phospholipids

(Schneider *et al.*, 2003). Dark layers on vessel walls that appeared under TEM after OsO_4

fixation were also found in a few other species (Fineran, 1997, Zimmermann *et al.*, 2004,

Westhoff *et al.*, 2008). In fact, it had been shown much earlier that a thin coat of lipids remains

on the lumen walls of xylem conduits from live cell content after conduit maturation (Scott *et al.*,

1960, Esau, 1965, Esau *et al.*, 1966).

The presence of lipids in xylem conduits may seem difficult to reconcile with the cohesion

tension (CT) theory of water transport (Askenasy, 1895, Dixon and Joly, 1895), which posits the

existence of negative pressure (i.e., tension) in xylem sap. Lipids are either hydrophobic or

amphiphilic, either of which properties would appear to pose a problem for maintaining negative

pressure in the sap without forming gas bubbles on hydrophobic or amphiphilic surfaces.

Accordingly, Zimmermann *et al*. (2004) cited evidence for lipids in the xylem apoplast to bolster

their arguments against the CT theory. However, the CT theory is strongly supported by multiple

lines of evidence, both indirect and direct (Angeles *et al.*, 2004, Wheeler and Stroock, 2008,

Jansen and Schenk, 2015, Venturas *et al.*, 2017), so the existence of apoplastic xylem lipids

would require an explanation within the context of this theory (Schenk *et al.*, 2017).

Any idea that there are no amphiphilic lipids in the xylem apoplast was put to rest when

Gonorazky *et al*. (2012) published an analysis of phospholipids in intercellular fluids and xylem

fluids of tomato plants. This was followed by reports of phospholipids in xylem sap and on

conduit surfaces of five woody angiosperm species from diverse phylogenetic backgrounds

(Schenk *et al.*, 2017, Schenk *et al.*, 2018). These findings raise questions about the origins of

these lipids, which could be remains from previous living vessel cell content (Scott *et al.*, 1960,

Esau, 1965, Esau *et al.*, 1966) or potentially be transported into vessels from conduit-associated

parenchyma cells (Morris *et al.*, 2018b). Even more important are questions about their functions

in xylem (Schenk *et al.*, 2015, Schenk *et al.*, 2017). However, surely the first question about

these lipids regards their chemical nature. What are they? There is clear evidence for

phospholipids in the xylem apoplast, but what are their characteristics, and how about galacto-

and sulfolipids?

To answer these questions, we conducted a lipidomic analysis of xylem sap extracted from seven

species sampled across the angiosperm phylogeny, including five woody angiosperm species

included in previous studies (Schenk *et al.*, 2017, Schenk *et al.*, 2018), to quantify and

characterize all phospho- and galactolipids in xylem sap and also test for the presence of

sulfolipids in a subset of species. Because contamination from cut surfaces is a notorious

problem with all xylem sap analyses (Schurr, 1998), we analyzed contamination controls from

cut and cleaned xylem surfaces at the sap collection end of stems and compared their lipid

conentrations and compositions to xylem sap samples to test if lipids in extracted sap originate

from damaged living cells during extraction. We also used a lipid tracer to test for lipid

contamination from successive cuts of stems during xylem sap extractions. Because lipids in

xylem vessels have been reported to originate from lipid bilayer membranes in the cell content of

living vessels (Scott *et al.*, 1960, Esau, 1965, Esau *et al.*, 1966), we compared the lipid

composition of xylem sap for two species to that of wood from the same stems to test the

prediction that there would be no differences in composition. Lipids in sap sampled for one

species were compared in July and March to test for seasonal differences in lipid composition

that could indicate developmental effects, lipid transport into sap, or apoplastic enzyme activity.

In addition, locations of lipids in xylem were visualized via confocal and transmission electron

microscopy in bay laurel, *Laurus nobilis*, a species much studied for its hydraulic properties

(e.g., Tyree *et al.*, 1999, Zwieniecki *et al.*, 2001, Hacke and Sperry, 2003, Gascó *et al.*, 2006,

Espino and Schenk, 2011, Nardini *et al.*, 2017).

MATERIAL AND METHODS

Plant species

The study was conducted with seven plant species from six major angiosperm clades (APG III,

2009) and different growth forms: *Liriodendron tulipifera* L. (winter-deciduous tree,

Magnoliaceae, Magnoliales, magnoliid clade), *Laurus nobilis* L., (evergreen tree, Lauraceae,

Laurales, magnoliid clade), *Bambusa oldhamii* Munro (bamboo, Poaceae, Poales, commelinid

clade), *Triadica sebifera* Small (syn. *Sapium sebiferum*, winter-deciduous tree, Euphorbiaceae,

Malpighiales, fabid clade), *Geijera parviflora* Lindl. (evergreen tree, Rutaceae, Sapindales,

malvid clade), *Distictis buccinatoria* (DC.) A.H.Gentry (syn. *Amphilophium buccinatorium*

(DC.) L.G. Lohmann, evergreen liana, Bignoniaceae, Lamiales, lamiid clade), and *Encelia*

farinosa Torr. & A.Gray (drought-deciduous desert shrub, Asteraceae, Asterales, campanulid

clade). All grow in the Fullerton Arboretum or on the California State University Fullerton

campus in Fullerton, California, United States, except for *Laurus nobilis*, which was collected

from the Los Angeles County Arboretum in Arcadia, California, United States. Specimens of

Laurus nobilis for imaging studies were collected from the Botanical Garden at Ulm University,

Germany. All species will be referred to by their generic names in this paper.

Measurements were conducted in two stages: 1. Lipid composition of xylem sap and

contamination controls from all seven species: Stems for xylem sap extraction for the first stage

of measurements (n = 3 for each species) were collected in August 2017 from *Liriodendron*,

Triadica, *Geijera*, *Distictis*, and *Encelia*, and in May 2018 from *Bambusa* and *Laurus*. 2.

Comparisons of lipid composition in xylem sap and wood in a subset of three species, and

additional tests for potential contamination of xylem sap from cut surfaces using lipid tracers in

two species (*Geijera* and *Distictis*). Collections for the second stage of measurements (n = 4 for

each species) were done in October 2019 for *Liriodendron*, January 2020 for Distictis, and

March 2020 for *Geijera*. Some methods were common to both sets of measurements while others

differed, as indicated below.

Xylem sap extraction (all measurements)

Most collections were done at dawn to ensure full hydration of stems. The previously published

method for xylem sap extraction (Schenk *et al.*, 2017) was modified to completely eliminate

contact of xylem sap with hydrophobic surfaces in order to avoid loss of lipids that could cling to

such surfaces (Fig. 1). Samples were only in contact with glass or, during stage 1 measurements,

with LoBind surfaces of microcentrifuge tubes (Eppendorf AG, Hamburg Germany), which

consist of a two-component polymer mix that creates a hydrophilic surface. Branches were cut

from the plants at a length exceeding the longest vessel for each species, which had been

previously measured via air injection (Greenidge, 1952). Stems were transported immediately to

the lab, and cut under water, with the final cuts made with a fresh razor blade. The bark was

removed from the proximal end for about 4 cm length to expose the xylem cylinder (Fig. 1). The

cut surface was thoroughly cleaned with deionized water using a high-pressure dental flosser

(WP-100 Ultra Water Flosser, Waterpik Inc., Fort Collins, CO, USA) for two minutes to remove

cell debris and cytoplasmic content from the surface. A control sample to determine the amount

of contamination from living cell remnants from the cleaned, cut surface was taken by inserting

the cut end into a glass vial containing 1 mL of nanopure water and leaving it there for one

minute. The liquid was then moved in a glass pipette into a pre-weighed 1.5 mL LoBind

Eppendorf microcentrifuge tube (for stage 1 extractions using partial freeze drying) or 2 mL

glass vials (for stage 2 extractions using a SpeedVac), immediately flash-frozen in liquid

nitrogen, and stored in a -20°C freezer until further processing, usually with 1-2 days.

Xylem sap was extracted under vacuum (Fig. 1). Tight-fitting latex tubing, about 2 cm long, was

fitted over about half of the exposed xylem cylinder next to the bark, leaving about 2 cm of

xylem cylinder freely exposed. The stem was then inserted from the top into a rubber flanged

stopper, avoiding any contact of the xylem with the stopper during insertion and creating a tight

seal between latex tubing and the stopper. The exposed xylem cylinder was then cleaned again

with deionized water using a high-pressure dental flosser and excess water removed with a

Kimwipe. Xylem sap was collected in a glass test tube embedded in ice inside a 1 L Buchner

flask, which was also embedded in ice. The flask was subjected to lab vacuum for about 30

seconds. The distal end of the branch was then cut back by about 4 cm, followed by successive 2

cm cuts, which were made to all the side branches of the stem until sap was observed dripping

into the test tube. Once dripping sap was observed, further 1 cm cuts were made every minute to

allow for slow, continuous removal of xylem sap. Depending on stem size, 1 to 2 mL of sap were

extracted from each stem, moved with a glass pipette into pre-weighed 1.5 mL LoBind

Eppendorf microcentrifuge tubes, (for stage 1 extractions using partial freeze drying) or 2 mL

glass vials (for stage 2 extractions using a SpeedVac), flash-frozen in liquid nitrogen, and stored

in the freezer until further processing.

Lipids were extracted using two different methods, because a SpeedVac evaporator instrument

had become available by the time stage 2 measurements were conducted, which allowed us to

simplify procedures. The two methods were compared and validated against each other

[**Supplementary Information**].

Lipid extraction in the stage 1 measurements of seven species: Partial freeze-drying

In preliminary experiments we had found that lipid extraction from completely freeze-dried sap

samples was incomplete, most likely due to formation of insoluble aggregates. We therefore

developed a new protocol to partially freeze-dry samples for lipid extraction and thereby avoided

aggregate formation, which yielded about 8 times more lipids than complete freeze-drying (data

not shown). For this protocol, weighed cell contamination controls and xylem sap samples of

about 1 mL in pre-weighed LoBind Eppendorf microcentrifuge tunes were partially lyophilized

in a freeze-dryer (FreeZone 1 Liter Benchtop Freeze Dry System, Labconco, Kansas City, MO,

USA) and observed until only about 100 µL of sample remained. They were then taken from the

freeze-dryer and weighed again to determine the remaining volume of aqueous sample. The

remaining aqueous samples in two (for *Encelia*, *Bambusa*, and *Laurus* xylem sap and *Bambusa*

and *Laurus* controls) or four (all xylem sap and controls from all other species) LoBind

microcentrifuge tubes were combined into one sample originating from 2 mL or 4 mL (see

breakdown of species and controls above) of xylem sap or control using a glass pipette.

Methanol:chloroform 1:1 (both HPLC grader, Fisher Scientific) was added to the combined

aqueous sample to create an approximate 5:5:1 methanol:chloroform:water one-phase mixture.

The mixture was then vortexed, centrifuged, and the supernatant collected with a 1 mL glass

syringe with PrecisionGlide Needle. A new one-phase 5:5:1 mixture of

methanol:chloroform:water was then added to the residue, the mixture vortexed, centrifuged, and

the supernatant again collected and combined with the previously collected supernatant. Samples

were then dried in a desiccator outfitted with an in-line carbon filter capsule (model 6704-7500,

Whatman, GE Healthcare Life Sciences, UK) and sent to the Kansas Lipidomics Research

Center at Kansas State University for mass spectrometry analysis.

Lipid extraction in stage 2 measurements of three species: SpeedVac evaporation

The lipid extraction protocol at this stage was changed, because a SpeedVac evaporator designed

for chloroform extraction had become available, which allowed us to simplify the procedure.

Following xylem sap extraction, samples were immediately freeze-dried using a SpeedVac

Concentrator unit (model Savant SPD121P, Thermo Scientific, Waltham, MA, USA). Samples

were freeze-dried for 7 hours to permit complete drying of xylem sap in 2 mL glass vials.

Freeze-dried samples were then stored at -15 °C.

To isolate lipids from freeze-dried xylem sap samples, a 5:5:1 methanol:chloroform:water one-

phase mixture (341 µL methanol, 341 µL chloroform, 68 µL nanopure water) was added to

previously fully freeze-dried vials containing xylem sap. Following vortexing for 60 s, the

solution was transferred to new 5 mL glass centrifuge tubes via glass pipette and centrifuged at

3500 rpm for 8 min. The supernatant of the centrifuged sample containing lipids was then

collected using a glass pipette into a new 2 mL glass vial. This process was repeated to ensure

maximal collection of lipids from all glass surfaces, with supernatant of the repeated steps added

into the final 2 mL glass vial. Samples were then freeze-dried using the SpeedVac Concentrator

unit for 7 h to permit complete removal of all liquid in 2 mL glass vials. Freeze-dried lipid-

isolate samples were then stored on dry ice and sent to the Kansas Lipidomics Research Center at

Kansas State University for mass spectrometry analysis.

The partial freeze-drying and SpeedVac extraction methods were subjected to a methods

comparison and were found to yield similar results (Methods S1) [**Supplementary**

Information].

Seasonal changes in lipid composition of xylem sap in Geijera

Xylem sap was extracted from *Geijera* in mid July 2019 (n = 8), and early March 2020 (n = 8).

At both times, half of the 8 samples were from lipid tracer experiments (see below), and because

no tracer was found in collected xylem sap for *Geijera* (see Results), samples from tracer

treatments and controls were combined to increase the statistical power of comparisons. Lipids

were extracted as described above, and lipid concentrations and composition compared between

sampling times to determine if there were changes between summer and winter collections.

Lipid extraction from wood in stage 2 measurements of three species

Lipids were extracted from wood by using a modification of methods developed originally for

lipid extractions from leaves (Shiva *et al.*, 2018). Following xylem sap extractions (see above)

from the same stems, four thin wood sections of exposed xylem (~200 mg) were cut with a

sledge microtome (model GSL 1, Schweingruber, Switzerland) and immediately submerged in

0.4 mL of hot isopropanol (75°C) for 15 min in 2 mL glass vials to prevent enzymatic

degradation of lipids from cut cells. This procedure made it impossible to determine the precise

weight of cut sections, so it allows only determination of lipid composition, not lipid

concentrations. To approximate the average fresh weight of wood samples, separate sections

from the same stem were cut and weighed. To isolate lipids from wood samples, 0.54 mL

chloroform, 0.54 mL methanol, and 0.104 mL nanopure water were added to the vial and softly

shaken for 24 h on a compact orbital shaker (Cole-Parmer). The 5/5/1

chloroform/methanol/water solution was then transferred with a glass pipette into a new 2 mL

glass vial and freeze-dried using a Speedvac.

Lipid tracer studies to quantify lipid contamination from cut stem surfaces (stage 2

measurements)

To determine if cutting stems during vacuum extraction of xylem sap introduces lipids into

extracted sap, lipid tracer studies were conducted with *Geijera* (short vessels) and *Distictis* (long

vessels). The tracer was di17:0 phosphatidylethanolamine (di17:0 PE) (Avanti Polar Lipids,

Alabaster, AL, USA), because it was absent in previous measurements of xylem sap. All cutting

blades were cleaned vigorously with 1% Alconox solution (product No. 1104, Alconox, White

Plains, NY, USA) before each cut to prevent surface contamination with tracer from blades. Sap

was extracted from stems as described above until the remaining stems were 50 cm long. At this

point, vacuum was relaxed, and 0.1 mL of 0.07 mg/ml di17:0 PE lipid tracer in chloroform was

added to the freshly-cut distal stem surface (see Fig. 1). The chloroform evaporated immediately,

and the tracer was allowed to absorb into vessels for 15 s. Vacuum was reapplied to the stem

until approximately 0.5 mL xylem sap was collected. Two more cuts were done using the same

tracer application at 45 cm and 40 cm length, at which point the extraction was concluded. Lipids

were extracted from xylem sap as described above for stage 2 measurements.

Vessel volume analysis

1
2
3
4 239   Because lipid micelles are mostly too large to pass through pit membranes (Zhang *et al.*, 2020),
5
6 240   we hypothesized that lipid concentrations in extracted xylem sap would be related to the volume
7
8 241   of vessels opened when a stem is cut for sap extraction. To determine the average opened vessel
9
10 242   volume for each species, we used silicone injections (Sperry *et al.*, 2005, Wheeler *et al.*, 2005).
11
12 243   Silicone compound (Rhodorsil RTV-141; Rhodia USA, Cranbury, NJ, USA) mixed with 0.41%
13
14 244   blue silicone dye (Blue Silc Pig; Smooth On, Easton, PA, USA) was injected from the proximal
15
16 245   end into flushed stems ($n = 4$ per species) that were between 0.9 and 1.7 cm in diameter and 25
17
18 246   cm in length, except 50 cm in length for *Distictis*. Injection took place at approximately 50 kPa
19
20 247   for 24 h, after which the silicone was allowed to dry for 72 h prior to sectioning. Cross sections
21
22 248   (cut at 0.7, 1.4, 2.9, 5.8, 8.3, 11.8, and 24 cm from the cut surface for all species, except at 0.7,
23
24 249   1.4, 2.8, 5.8, 11.6, 16.6, 23.6, and 48 cm for *Distictis*) were thin-sectioned using a sledge
25
26 250   microtome (model GSL 1, Schweingruber, Switzerland), dry-mounted on glass slides, and
27
28 251   imaged using a Leica MZ16 stereomicroscope with digital camera (model INFINITY 2-1C-IQ,
29
30 252   Lumenera, Ottawa, ONT, Canada) at 20× zoom and backlit using a transmitted light stage
31
32 253   connected to a halogen light source (model KL 1500 LCD, Schott, Germany). Adobe Photoshop
33
34 254   (version CS6) was used to blend images into one combined image for each cross-section (Fig.
35
36 255   S10) [**Supplementary Information**], and an analysis of filled vessels, vessels diameters, vessel
37
38 256   wall perimeters, and vessel density was completed using ImageJ (Schneider *et al.*, 2012).
39
40 257   Silicone-filled vessels were counted in each image, and the area of silicone-filled vessels in each
41
42 258   image was quantified both in mm^2 and as percent of the total xylem area. The silicone-filled
43
44 259   vessel volume was calculated from these data by linear interpolation between each cross-section
45
46 260   (Fig. S11) [**Supplementary Information**]. Counts of filled vessels (*C*) at each distance (*d*) from
47
48 261   the injected surface were used to estimate median vessel lengths, based on fitting the equation
49
50 262 $\ln C = a + b \times d$ using the software TableCurve 2D (version 5.01, Systat, San Jose, CA, USA).
58   11
59
60

Mass spectrometry

Direct-infusion electrospray ionization triple-quadrupole mass spectrometry was performed

using a Waters Xevo TQS mass spectrometer or Applied Biosystems 4000 (Shiva *et al.*, 2013).

Instrument-specific details can be found in Supplemental Tables S1 and S2. The dried samples,

each originating from 2 mL or 4 mL of xylem sap or cell contamination control, or from wood

samples (see breakdown of species above), were dissolved in 1 mL chloroform in a 2 mL glass

vial. After a test to determine the optimal amount for analysis, 50-1000 µL of each sample was

transferred to a 2-mL vial containing internal standards (Supplemental Table S2). 1.2 mL of

chloroform:methanol:300 mM ammonium acetate in water (30:66.5:3.5) was added and the

sample mixed. The sample was continuously infused to the mass spectrometer from a 1-mL

loop.

In addition to the scans described in Shiva *et al*. (2013), a scan for SQDG (Pre 261.1) in positive

ion mode was also acquired (except for samples from *Laurus* and *Bambusa*) (details in

Supplemental Table S2). Data were processed as described in Shiva *et al*. (2013) using the

LipidomeDB Data Calculation Environment at http://lipidome.bcf.ku.edu:8080/Lipidomics/

(Zhou *et al.*, 2011). Appropriate response factor for the biological galactolipid molecular species

in comparison to the saturated galactolipid internal standards were applied. The limit of

detection in the samples analyzed was 0.002 nmol for the stage 1 measurements of seven species.

And 0.0005 nmol for stage 2 measurements.

Confocal laser scanning microscopy

To test for lipids in xylem of *Laurus*, fluorescent FM1-43 dye (Molecular Probes, Life

Technologies, Eugene, OR, USA) was infiltrated into living xylem and imaged with laser

scanning confocal microscopy, as described in Schenk *et al*. (2018). FM1-43 is an amphiphilic

fluorophore that is virtually non-fluorescent in water and strongly fluorescent under cyan light

excitation when bound to lipids, including amphiphilic lipids, and biological membranes

(Jelínková *et al.*, 2010). It was used as a 5 µg/mL solution in nanopure water with 488 nm

excitation and an emission window of 578 to 618 nm. Woody stems, approximately 5-8 mm in

diameter and about 10 cm long, were cut from the plants and submerged in DI water. About 2.5

cm were cut under water from each end of the stem to remove most bubbles in the xylem that

may have been induced by the cutting. The cut ends were then recut with new razor blades to

create clean and smooth surfaces. Bark was removed from about 1 cm length at the distal end to

expose the xylem. The proximal end of each stem was connected to a vacuum flask using tight-

fitting latex tubing around the stem inside Tygon tubing. A short piece of Tygon tubing was

fitted to the exposed xylem cylinder at the distal end, using latex tubing around the cylinder to

create a tight fit. The distal cut end was cleaned by washing it three times with 500 µL nanopure

water using a pipette. 1 mL of fluorophore solution was then added into the tubing and onto the

exposed xylem surface. Lab vacuum was turned on and turned off once the solution was

absorbed into the xylem, but no air was aspirated. The control was treated the same way but,

instead of the fluorophore solution, 1 mL of nanopure water was sucked through the stem. A 2

mm long stem piece was then cut along its transverse plane from the center of the stem segment

and the distal surface recut using a sledge microtome (Sledge microtome GSL 1, Schweingruber,

Switzerland). The stem piece was then placed onto a cell culture dish with a glass bottom of 175

µm (CELLview™ dish, Greiner Bio-One, Germany) in a drop of water and observed using the

inverted microscope of a confocal laser scanning microscope (model TCS SP8 with a

HyVolution super-resolution detector, Leica Microsystems, Wetzlar, Germany). Lignin was

detected with 405 nm excitation and an emission window of 467 to 509 nm. Imaging was done

with a 63.0×1.20 objective and $3\times$ digital zoom in water, using four scans per image at $1760 \times$

1760 resolution.

Transmission electron microscopy (TEM)

Small sectioning blocks of xylem from fresh *Laurus* stems were given two different treatments

as described in Schenk *et al*. (2017): (1) fixation with a solution containing 2.5% glutaraldehyde,

1% sucrose, and 0.1 M phosphate buffer at pH 7.3, and (2) fixation with glutaraldehyde and then

postfixation with a 2% aqueous osmium tetroxide (OsO_4) solution for 2 hours at room

temperature. The samples were dehydrated through a gradual ethanol series and embedded in

Epon resin. Moreover, samples treated with glutaraldehyde only were observed with and without

post staining of TEM grids with aqueous uranyl acetate and lead citrate to test the potential

staining effect on vessel-vessel pit membranes (Ellis, 2014). The duration of the staining was 5

minutes for uranyl acetate and 1 minute for lead citrate. Transverse semi-thin sections were cut

with an ultramicrotome (Leica Ultracut UCT, Leica Microsystems, Vienna, Austria), stained

with 0.5% toluidine blue in 0.1 M phosphate buffer, and mounted on microscope slides using

Eukitt. Ultra-thin sections between 60 nm and 90 nm were mounted on copper grids (Athena,

Plano GmbH, Wetzlar, Germany) and observed with a JEM-1210 TEM (Jeol, Tokyo, Japan) at

120 kV. Digital images were taken using a MegaView III camera (Soft Imaging System,

Münster, Germany).

Data analysis

To test for the presence of di17:0 PE tracer in tracer experiments, concentrations of di17:0 PE in

xylem sap from stems treated with di17:0 PE tracer at cut stem surfaces were compared to

controls without tracer applications using two-tailed t-tests (n = 4) assuming equal variance in

Microsoft Excel. Mean lipid concentrations were compared to various anatomical parameters of

xylem for each species (incl. opened vessel volume, opened vessel volume relative to surface

area, mean vessel diameter and wall perimeter, both calculated from the diameter of a circle with

the same area as the vessel, and mean vessel density) via reduced major axis regression in the

statistical software package PAST (ver. 4.02) (Hammer *et al.*, 2001).

To compare lipid compositions between samples, mass spectrometry data were analyzed using

the online MetaboAnalyst software package (https://www.metaboanalyst.ca). Any concentrations

below the detection limit were set to zero, and data were log-transformed to normalize

distributions. Compositions of polar lipids in xylem sap were compared in MetaboAnalyst as

normalized concentrations between seven species and between two sampling periods for *Geijera*

using principal component analysis (PCA) and heatmaps, based on Euclidean distance measures,

and a Ward clustering algorithm to visualize differences among the species or sampling times.

Unpaired t-tests, adjusted for false discovery rates (FDR) were used to identify lipids that

differed between sampling times.

Compositions of polar lipids in xylem sap from six species (one sample for *Encelia* was lost,

causing the species to be excluded from this analysis) were compared to those in paired

contamination controls to test the hypothesis that xylem sap lipids originate from damaged living

cells and therefore do not differ in lipid composition to contamination controls. For these

comparisons, concentrations were converted into percentages, data were log-transformed and

analyzed using PCA in MetaboAnalyst.

Polar lipid compositions were compared between wood and xylem sap for *Geijera* and *Distictis*

by converting concentrations to percentages, then excluding any measurements that were below

the detection limit, and using PCA (95% CI). Paired t-tests adjusted for FDR were used to

identify lipids that differed between sap and wood. Because wood samples included xylem sap

lipids, the two data sets were not independent, so PCA and t-tests were used qualitatively to

determine if there were differences between sap and wood and identified lipids that differed the

most.

RESULTS

Comparisons of xylem sap and contamination controls (seven species)

Polar lipid concentrations in cell contamination controls ranged from 1.1% of the amount

detected in xylem sap in *Distictis* to 9.0% in *Laurus*, averaging 4.49 (± 2.80 SD) percent across

species (Table 1). The complete lipidomics data set may be found in Table S2 for xylem sap and

Table S3 for cell contamination controls [**Supplementary Information**]. All contamination

controls were far more variable and distinctly different in polar lipid composition compared to

xylem sap, as shown in principal component analyses (PCA; Fig. S12), clearly demonstrating

that xylem sap lipids do not originate from cells cut open during sap extraction [**Supplementary**

Information].

Lipid tracer studies to detect lipid contamination from cut surfaces (two species)

Concentrations of di17:0 PE in xylem sap of *Geijera* stems treated with di17:0 PE tracer at cut

distal stem surfaces and in controls without tracer applications were below 1 pmol/mL in both

treatments and not significantly different ($p = 0.738$). Concentrations of di17:0 PE in xylem sap

of *Distictis* stems treated with di17:0 PE tracer at cut distal stem surfaces were 0.9 pmol/mL (±

0.8 SE), 0.2 pmol/mL (± 0.1 SE) in controls without tracer applications, and not significantly

different ($p = 0.131$).

Xylem sap lipidomics (seven species)

Polar lipids extracted from xylem sap and cell contamination controls included the galactolipids

digalactosyldiacylglycerol (DGDG, Fig. S1) and monogalactosyldiacylglycerol (MGDG, Fig.

S2), as well as all common plant phospholipids. Total lipid concentrations for xylem sap varied

between 0.18 nmol / mL for *Laurus* and *Liriodendron* and 0.63 nmol / mL for *Bambusa*,

averaging 0.41 (± 0.17 SD) nmol / mL across species (Table 1; Fig. 2 A). Lipid concentrations in

sap were not statistically related to the vessel volume opened by cutting stems, but there was a

significant negative relationship to the total vessel perimeter at the xylem surface (Fig. 2 C).

The most common lipids grouped by their head groups (Fig. 2 A) were DGDG, MGDG,

phosphatidylcholine (PC, Fig. S4), phosphatidic acid (PA, Fig. S8), phosphatidylethanolamine

(PE, Fig. S5), phosphatidylinositol (PI, Fig. S6), with other phospholipids present in much lower

amounts, including phosphatidylserine (PS, Fig. S7), phosphatidylglycerol (PG, Fig. S3),

lysophosphatidylethanolamine (LPE, Fig. S9), lysophosphatidylcholine (LPC, Fig. S9), and

lysophosphatidylglycerol (LPG). Trace amounts of a contaminant that was also found in solvent

interfered with the detection of LPG, which was therefore excluded from all tables, figures, and

analyses for stage 1 measurements. (The problem was solved for stage 2 measurements by using

glass vials instead of LoBind Eppendorf tubes in all subsequent experiments.)

The most pronounced differences in chemical compositions between species were in the

percentage of galactolipids (*Laurus* 75.4%, *Liriodendron* 35.9%, *Bambusa* 57.0%, *Triadica*

18.8%, *Geijera* 10.6%, *Distictis* 20.4%, *Encelia* 7.1%). Sulfolipids were detected only in trace

amounts. In the PCA comparing the seven species, no species stood out as different in polar lipid

compositions from the other six species on the PC1 axis, which explained 46.3% of the variation,

while *Laurus* and *Bambusa* stood out on the PC2 axis, explaining 16.7% of the variation, as

distinctly different (Fig. 2 D). The most pronounced differences between species were in the

percentages contributed by DGDG, MGDG, PC, and PA (Table 1; Fig. 2 A). DGDG varied from

about 3.0% of all lipids in *Encelia* to 32.0% in *Bambusa*, with a mean across species of 12.4 (±

10.5 SD)%; MGDG varied from 4.1% in *Encelia* to 54.6% in *Laurus*, with a mean across species

of 19.8 (±17.8 SD)%; PC varied from 12.3% in *Laurus* to 30.1% in *Triadica*, with a mean across

species of 18.2 (±6.3 SD)%; PA varied from 1.9% in *Laurus* to 50.6% in *Encelia*, with a mean

across species of 28.7 (±17.6 SD)%.

The most common number of acyl chain carbon atoms were 34 and 36, with galactolipids mostly

having more 36-carbon combinations than phospholipids (Fig. 2 B, Fig. S1-2). Only on average

1.56 (± 0.70 SD)% of all lipids were fully saturated phospholipids, and the most common

numbers of double bonds were 1-3 for the 34 carbon combination of 2 acyl chains and 3-5 for the

36 carbon combination (Fig. 2 B, Fig. S3-9). Averaged across species, lipids with 34:1-3 (carbon

atoms:number of double bonds) and 36:3-5 accounted for 84.0 (± 4.9 SD)% of all apoplastic

xylem lipids. PA 34:2 was especially abundant in all species, except for *Laurus*, making up about

23% of all polar lipids in *Encelia* and 25% in *Geijera*. The lipid composition of the galactolipids,

with a total of four double bonds prominent, suggests that 18:2 fatty acids may be common.

Seasonal changes in lipid composition of xylem sap in Geijera

Total concentrations of lipids in xylem sap were not different between samples collected in July

and March, but concentrations of the galactolipids DGDG and MGDG were significantly higher

in sap collected in March (Table 2). Concentrations of PA tended to be lower in March, but

differences were not significant because of high variability among samples. Overall, 18

individual lipids differed significantly in their concentrations between July and March (Fig. 3 B),

including seven types of DGDG, four of MGDG, and the remaining ones phospholipids. In PCA

(Fig. 3 A), March samples were slightly more variable than July samples, and the 95%

confidence regions for the two periods were clearly separated on the PC1 axis, which explained

27% of the variation. Overall, there were clear seasonal differences in lipid composition.

Comparisons of lipid compositions in wood and xylem sap (two species)

Wood samples include xylem sap lipids, so the samples are not independent and can be

compared only qualitatively (Table 3). However, PCAs revealed some differences between wood

and sap, especially for *Distictis* (Fig. 4 A), where FDR-adjusted t-tests identified five

phospholipids (out of 125) as different between wood and sap. No headgroups, chain lengths, or

unsaturation levels were consistently different between wood and sap for *Distictis*, but several

types of PC stood out as being more abundant in sap than in wood, while several types of PS and

lysophospholipids showed the opposite trend (Fig. 4 B). For *Geijera*, only one polar lipid, PI

34:3, was identified by t-tests as different between wood and sap (Fig. 4 B), and PE was slightly

more abundant (by 3.3%) in wood than in sap, and lysoPC was slightly less abundant in wood

than in sap (by -0.14%). PA was just as abundant in wood as in sap (Table 3), even though wood

samples were treated immediately with hot isopropanol to inactive PLD enzyme activity. No

galactolipids were flagged as being different between wood and sap of either species. Overall,

differences in lipid composition between wood and sap were not pronounced.

Visualization of lipids in xylem vessels of Laurus

Treatment with FM1-43 showed continuous lipid layers on vessel surfaces of *Laurus nobilis*,

with high concentrations in inter-vessel pits (Fig. 5 A-B). Pit membranes were clearly visible

under confocal laser scanning microscopy as thick, pillow-shaped, black structures with lipid

coatings on either side and some faint labelling inside the membranes. Never-dried pit

membranes of *Laurus* were 1,252 nm (± 238 nm, standard deviation; n = 36 pit membranes)

thick under confocal microscopy (Fig. 5 A-B), but 601 nm (± 150 nm; n = 19) under TEM (Fig.

5 D-G), showing that the dehydration required for TEM by propanol could result in shrinkage of

the pit membrane. These observations were very similar to those published previously for five of

the other study species, including *Liriodendron*, *Triadica*, *Geijera*, *Distictis*, and *Encelia*

(Schenk *et al.*, 2018). TEM samples not treated with OsO_4 showed a dark and homogeneous

appearance (Fig. 5 D-E). After treatment with OSO_4, the lipid layer appeared to be broken up

into aggregates under TEM (Fig. 5 F-G) but was found in a continuous layer in the same

locations observed under confocal microscopy on vessel and pit surfaces and on the pit

membrane (Fig. 5 A-B).

DISCUSSION

The most important take-home message of this study is that amphiphilic lipids exist in xylem

conduits across the angiosperm phylogeny, including magnoliids, monocots, and eudicots. The

available evidence based on our results indicate clearly that apoplastic lipids do not originate

from cells cut open during sap extraction, as shown by contamination controls, lipid tracer

experiments, and imaging. Phospholipids have been reported to occur in xylem sap (Gonorazky

et al., 2012, Schenk *et al.*, 2017), but not galactolipids. These findings raise questions about the

chemical composition of xylem sap lipids, their origins, and functions of these lipids in

angiosperm vessels.

Chemical composition of xylem sap lipids

The concentrations of lipids in xylem sap were surprisingly consistent across species, with the

two species from the magnoliid clade, *Laurus* and *Liriodendron*, having about a third of the

concentrations of the other five species (Table 1; Fig. 2 A). Because lipid micelles or lipid-

coated gas bubbles in xylem sap are mostly larger than 50 nm in diameter (Schenk *et al.*, 2017)

and therefore too large to pass through pit membrane pores, which do not exceed 20 nm in

hydrated pit membranes (Choat *et al.*, 2003, Zhang *et al.*, 2017, Zhang *et al.*, 2020), it is likely

that the observed xylem sap lipids originate largely from vessels cut open for the sap extraction.

To account for this, we analysed the opened vessel volumes of the seven species and found no

significant relationship between lipid concentrations in sap and vessel volumes, but there was a

significant negative relationship with the total vessel wall perimeter at the cut xylem surface

(Fig. 2 C). It appears that the xylem functions like a filter during lipid extraction, so that xylem

with higher wall surface area, i.e., more and smaller vessels, traps more surface-active lipids,

leading to lower lipid yields in extracted sap. Because the open vessel volumes were typically

smaller than the 1 to 2 mL of xylem sap extracted (except for the liana species *Distictis*), it is

likely that the total amount of lipids in sap is underestimated due to a filtering and dilution effect.

Lipids are insoluble and in xylem conduits most are attached to surfaces (Fig. 4; Schenk *et al.*,

2018), so the amounts extracted in sap may only be a small fraction of all apoplastic xylem

lipids. Quantification of all apoplastic lipids in xylem would require either complete extraction

from individual conduits with organic solvents and/or detergents, which could cause artifacts if

these substances penetrated into living vessel-associated cells (Morris *et al.*, 2018b), or

quantification of lipids in vessel images via mass spectrometry imaging (Ellis *et al.*, 2013).

The very low concentrations of lipids found in cell contamination controls from the basal xylem

surface show that on average at least 95% of the lipids in xylem sap samples did not originate

from cut parenchyma cells at basal xylem surfaces. *Bambusa* and *Distictis*, both of which have

phloem embedded with xylem (*Distictis* has intraxylary phloem) had the lowest percentages of

lipid concentrations in their control samples (4.3% in *Bambusa* and 1.1% in *Distictis*),

demonstrating that hardly any lipids in xylem sap samples originated from the phloem. Lipid

compositions of cell contamination controls were far more variable and different from those of

sap (Fig. S12), which also confirms that sap lipids, except for traces, do not originate from cut

living cells. Lipids in sap also clearly do not originate from cells cut at the distal end of stems, as

lipid tracer applied to distal stem surfaces during xylem sap extraction did not show up in sap

collections. Minute tracer amounts may have made their way through the exceptionally wide and

long vessels of *Distictis*, but not enough to raise tracer levels significantly. Lipids from cut cells

at the distal stem end are likely to attach to vessel surfaces as sap is extracted, or will be held up

by inter-vessel pit membranes, instead of moving through >40 cm of vessel length and showing

up in extracted sap.

The only other lipidomic analysis of xylem sap, for tomato (Gonorazky *et al.*, 2012), did not

include analyses of galactolipids, but the chemical composition of phospholipids was roughly

comparable to our findings, with PC and PA being the most abundant phospholipids. The xylem

sap concentration of phospholipids in tomato (0.037 μmol kg^{-1}) was similar to concentrations in

Laurus (0.044 µmol L^{-1}), but much lower than the average phospholipid concentrations across all

species studied (0.296 ± 0.161 SD µmol L^{-1}).

The presence of the galactolipids DGDG and MGDG in the xylem apoplast is especially notable,

because galactolipids are synthesized exclusively in plastids (Dörmann and Benning, 2002, Botté

and Maréchal, 2014) and are the most abundant lipids in chloroplasts, but they have never been

found in any apoplastic compartment, presumably because there have been so few studies of "the

black box of plant apoplast lipidomes" (Misra, 2016). Moreover, their physical properties, such

as interactions of the head groups with water molecules, are different from phospholipids

(Kanduč *et al.*, 2017). MGDG have very small head groups and form bilayers only in mixture

with other lipids (Dörmann and Benning, 2002), where interactions between MGDG and DGDG

head groups affect the physical properties of galactolipids at gas-water interfaces (Bottier *et al.*,

2007). It is also notable that sulfolipids, which are also abundant in plastids (Shimojima, 2011),

were only found in trace amounts in xylem sap.

Regarding acyl chains, it was observed that galactolipids mostly had 36-carbon combinations,

most likely consisting of two 18-carbon chains (Fig. 2 B). Some plants, termed "16:3 plants" can

incorporate 16C-fatty acids into galactolipids and desaturate them to 16:3 (via the "prokaryotic

pathway"), whereas others, termed "18:3 plants", assemble the diacylglycerol backbone of

galactolipids only in the E.R. 16:3 plants contain a mixture of 34C and 36C galactolipids,

whether 18:3 plants contain only 36C galactolipids. Leaves of both *Laurus nobilis* and

Liriodendron tulipifera contain small amounts (slightly over 1%) of plastid-synthesized 16:3

(Mongrand *et al.*, 1998). *Triadica*, *Geijera*, and *Distictis* leaves also may contain small amounts

of 16:3, but *Bambusa* and *Encelia* are likely to be 18:3 plants. Still, the majority of xylem sap

galactolipid backbones from all seven species are di-18C molecular species, consistent with

synthesis in the ER (Mongrand *et al.*, 1998). Additionally, it has previously been shown that a

16:3 plant, such as *Arabidopsis thaliana*, favors the eukaryotic pathway in non-photosynthetic

tissue (Devaiah *et al.*, 2006, Huynh *et al.*, 2012).

Along the same lines, the bulk of the observed lipid species appear to be less unsaturated than

most leaf tissues, containing only minor amounts of 18:3 chains, which are the most prominent

fatty acids in leaves. Whole tissue data demonstrate that generally *Arabidopsis* shoots are very

high in 18:3, while roots and seeds are less unsaturated with more 18:2 and 18:1-containing

galactolipids (Devaiah *et al.*, 2006, Huynh *et al.*, 2012). Indeed, Gonorazky *et al*. (2011)

observed that 18:3-containing phospholipids in leaves were less likely to be found in the

extracellular fluid than in the whole leaf. The lipid composition of the xylem galactolipids, with

36:4 species prominent, suggests that 18:2 fatty acids may be common. Similarly, 34:2 species,

which are likely 16:0/18:2 combinations are very common in phospholipids, suggesting an

abundance of 18:2 in xylem sap.

Besides finding galactolipids in xylem sap, the presence of phosphatidic acid (PA) at high

concentrations in some species is also notable (Fig. 2 A). High concentrations of PA in plant

samples are usually attributed to phospholipase D (PLD) activity, which cleaves head groups

such as the choline head-group of phosphatidylcholine (PC) off phospholipids to create PA

(Christie, 1993, Welti *et al.*, 2002). PLD is active when bound to cell membranes (Kolesnikov *et*

al., 2012), and cutting living cells causes PLD activation as part of a wounding response

(Bargmann *et al.*, 2009). High PA concentrations can be artifacts of wounding, and thorough

cleaning of cut xylem surfaces is therefore essential for keeping fragments of cell membranes

containing PLD out of xylem sap collections. PLD can remain active even in organic solvents

such as chloroform:methanol (Christie, 1993), possibly because membrane fragments form

inverted vesicles in organic solvents that allow for continued PLD activity in the hydrophilic

interior of the vesicle, as has been suggested for phospholipase C (PLC) (Kates, 1957).

The high PA concentrations found in xylem sap of some species, such as *Encelia*, *Geijera*, and

Distictis (Fig. 2 A) were initially a concern, because it was suspected that they could be sampling

artifacts. However, treatments of cut xylem surfaces with the PLD inhibitors 1-butanol, 5-fluoro-

2-indolyl des-chlorohalopemide (FIPI) (Su *et al.*, 2009), and a lipase inhibitor cocktail (Furse *et*

al., 2013) had no effect on PA concentrations in xylem sap of *Distictis* (data not shown).

Moreover, PLD or other phospholipases have never been detected in proteomic analyses of

xylem sap (Alvarez *et al.*, 2006, Djordjevic *et al.*, 2007, Aki *et al.*, 2008, Krishnan *et al.*, 2011,

Ligat *et al.*, 2011, Dugé de Bernonville *et al.*, 2014), which may not be surprising, because PLD

is often bound to membranes and therefore unlikely to occur naturally in xylem sap. Moreover,

thin wood samples also contained high amounts of PA (Table 3), even though the slices were

immediately dropped into hot isopropanol to inactivate PLD.

These lines of argument all support PA as a major natural ingredient of xylem saps, except for

that of *Laurus* (Fig. 2 A; Table 1). PA has also been found in relatively high concentrations in

xylem sap of tomato plants (Gonorazky *et al.*, 2012). PA has many functions in plants, interacts

with many proteins, and has important roles in lipid metabolism and signalling (Testerink and

Munnik, 2011), wounding responses (Bargmann *et al.*, 2009), and in responses to stress, such as

drought and salinity (McLoughlin and Testerink, 2013). The physical properties of PA are also

unique when compared to other amphiphilic lipids in having the smallest and negatively charged

head-group, binding of cations such as Ca^{2+} and Mg^{2+} (Ohki and Ohshima, 1985), and causing

PA to have lower surface tension in monolayers than other phospholipids (Weschayanwiwat *et*

al., 2005). Of all the various roles of PA, its interactions with cations and effects on surface

tension at gas interfaces are likely to be most important in the xylem apoplast environment.

The amounts of choline-containing phospholipids (PC and LPC) in xylem sap reported here were

much lower than estimated in previous research using an enzymatic assay (Schenk *et al.*, 2017),

where all choline quantified in xylem sap was assumed to originate via PLD activity from

choline-containing phospholipids. However, that assumption appears to have been incorrect.

Choline has been found in xylem sap at molar concentrations far exceeding those of lipids (Lima

et al., 2017) and is likely to originate from sources other than phospholipids in sap.

Where do apoplastic xylem lipids come from?

There are two possible and not mutually exclusive origins for apoplastic xylem lipids: Lipids

have been shown to remain from the living cell content of conduits during conduit development

(Scott *et al.*, 1960, Esau *et al.*, 1966). If this includes lipids from plastids, then this would easily

explain the presence of galactolipids in xylem sap. Lipids could also be transported into conduits

from conduit-associated parenchyma cells (Czaninski, 1977, Morris *et al.*, 2018b), but it seems

unlikely that plastid lipids would be moved into vessels. We compared the lipid composition in

sap to that of the wood that the sap was extracted from (Fig. 4) and found only minor differences

in composition and none in galactolipids. To some degree, this is expected, as wood includes

sap, so it is impossible for sap to include any lipids not found in wood samples. Out of 157 polar

lipids tested, only one for *Geijera* and five for *Distictis* stood out as different in relative amounts

between sap and wood (Table 2). Overall, the differences between sap and wood were small, so

it seems reasonable to conclude that sap and wood lipids are essentially the same, supporting the

origin of apoplastic xylem lipids, including galactolipids, from living vessel contents, as shown

previously in microscopic studies of developing vessels (Scott *et al.*, 1960, Esau *et al.*, 1966).

That said, there were clear seasonal differences in the polar lipid composition of *Geijera* xylem

sap between July and March (Fig. 3), with elevated galactolipid concentrations in March.

Seasonal changes could be a reflection of wood development, with newer vessels containing

different lipids than older ones, could be caused by apoplastic enzyme activities, or by lipid

transport from vessel-associated cells. This would require transport of lipids across a cell

membrane, the protective layer surrounding such cells, which consists mainly of polysaccharides

(Chafe and Chauret, 1974, Fujii *et al.*, 1981, Wisniewski *et al.*, 1991a, Wisniewski *et al.*, 1991b,

Wisniewski and Davis, 1995), as well as across the pit membrane, which mainly consists of

cellulose. The transport could involve non-specific lipid transfer proteins (nsLTPs), which can

transport lipids across cell walls (Domínguez *et al.*, 2015, Fich *et al.*, 2016, Li *et al.*, 2016,

Misra, 2016) and which have been found in xylem cell walls and xylem sap (Buhtz *et al.*, 2004,

Djordjevic *et al.*, 2007, Krasikov *et al.*, 2011, Ligat *et al.*, 2011). However, it is exceedingly

unlikely that plastid lipids would be transported into vessels, so seasonal differences in

galactolipids in xylem sap are more likely to be caused either by apoplastic enzyme activities or

by immobilization of galactolipids on vessel surfaces and in pit membranes, which would

prevent them from being extracted in sap. Further research will be required to understand these

seasonal changes.

Lipid transport or secretion of apoplastic lipid-altering enzymes from conduit-associated cells

would allow for changes in lipid concentration and composition in conduits seasonally or in

response to environmental conditions, which could possibly affect hydraulic functions and

embolism resistance (Charrier *et al.*, 2018). Plants species vary widely in the amount and degree

of contact between conduits and living parenchyma cells in secondary xylem (Martínez-Cabrera

et al., 2009, Morris and Jansen, 2016, Morris *et al.*, 2016, Morris *et al.*, 2018a), and it could be

that some species undergo seasonal changes in apoplastic lipids while others do not. Seasonal

changes in the electron-density of pit membranes have been observed (Schmid and Machado,

1968), and this could potentially indicate lipid accumulation in conduits over time.

Apoplastic xylem lipids and functional implications

The presence of lipids on xylem conduit surfaces and pit membranes is incompatible with the

widely accepted assumption that gas bubble formation in xylem is prevented by the high surface

tension of water (Oertli, 1971, Sperry and Tyree, 1988, Tyree and Zimmermann, 2002, Stroock

et al., 2014). There is unequivocal evidence for negative pressure in xylem, and this aspect of the

CT theory is not in question, although many open questions remain (Jansen and Schenk, 2015,

Schenk *et al.*, 2015, Schenk *et al.*, 2017, Venturas *et al.*, 2017). Molecular dynamics modelling

of lipid bilayers under negative pressure confirmed that cavitation in lipid bilayers and micelles

is unlikely to occur within the normal range of pressures (above -10 MPa) experienced in plants

(Kanduč *et al.*, 2020).

However, layers of phospho- and galactolipids on conduit surfaces and in pit membranes (Fig. 5)

will reduce the surface tension at gas-water interfaces substantially below that of pure water,

with the dynamic surface tension depending on the local concentration of lipids at the surface

(Zuo *et al.*, 2004, Kwan and Borden, 2010). Measurements of xylem lipids extracted from four

of the species studied in this paper found average equilibrium surface tensions to be about 25

mN/m, roughly a third of the surface tension of pure water (Yang *et al.*, 2020). These findings

are entirely compatible with the air seeding pressure required to force air through pore

constrictions in angiosperm pit membranes. Pore constrictions rarely exceed 20 nm in diameter

(Zhang *et al.*, 2020), and it would take about 7 MPa of pressure to force air through such a

constriction if surface tension were that of pure water (Kaack *et al.*, 2019). Actual air seeding

pressures through pit membranes tend to be in the range of -0.4 to -2 MPa (Bartlett *et al.*, 2016),

thereby supporting a role for lipids in pit membranes in the air seeding process.

In an earlier paper, we proposed a hypothesis to reconcile the presence of apoplastic xylem lipids

with the CT theory (Schenk *et al.*, 2015) and later amended and modified that hypothesis in the

light of new data (Schenk *et al.*, 2017). To summarize that hypothesis briefly, gas penetrating

from embolized conduits into pit membranes of sap-filled conduits may snap off nanobubbles

inside confined, lipid-coated pore spaces, resulting in the creation of lipid-coated nanobubbles

(Fig. 6), which have in fact been found and visualized in xylem sap (Schenk *et al.*, 2017). Key to

this hypothesis is that both the low dynamic surface tension of lipids and snapping off of bubbles

at pore constrictions inside fibrous pit membranes would limit the bubbles' sizes (Park *et al.*,

2019) and keep them below a critical threshold that prevents bubble expansion into embolism

(Schenk *et al.*, 2015, Schenk *et al.*, 2017). This process could happen under normal xylem water

potentials and would allow xylem to operate without spreading embolisms via air seeding from

gas-filled conduits. The population of lipid-coated nanobubbles could also function as a gas

reservoir that serves as a buffer when dissolved gas in sap exceeds saturation under increasing

temperatures (Schenk *et al.*, 2016).

Below a more negative air seeding pressure threshold, gas could potentially still penetrate

through the entire length of pit membrane pores, causing a growing bubble to emerge from the

pit membrane, where nothing would restrict the bubble's growth and resulting in embolism. This

is the widely accepted air seeding process (Sperry and Tyree, 1988, Tyree and Zimmermann,

2002, Stroock *et al.*, 2014, Brodribb *et al.*, 2016), the underlying mechanism of which is still

poorly understood (Jansen *et al.*, 2018), but which is supported by a large body of research,

including measurements of pit membrane pore constrictions and air seeding pressures (Kaack *et*

al., 2019). What the new lipid-nanobubble hypothesis could explain is how plants can operate

under negative pressure at all without forming embolism even under mildly negative pressure, a

feat that engineers have found almost impossible to replicate (Smith, 1994). If this hypothesis,

which is not contradictory but complementary to the cohesion-tension theory, is correct, then

lipids are required for water transport under negative pressure. In fact, Dixon (1914) already

suggested the colloids increase the tensile strength of xylem sap over that of pure water.

Conclusions

This study provides conclusive evidence for substantial amounts of phospho- and galactolipids in

xylem conduits across the angiosperm phylogeny, and these findings force major questions about

the functions of these lipids for plant water transport and the cohesion tension theory, as well as

about the origins of lipids. Research on the functions of lipids in water that is under negative

pressure and in the presence of gas-water interfaces in pit membranes and bubbles will benefit

from an interdisciplinary approach to involve surface scientists and physicists with expertise on

the behavior of lipids at surfaces such as pulmonary surfactant layers (Zuo *et al.*, 2008, Zhang *et*

al., 2011, Kanduč *et al.*, 2013, Kanduč *et al.*, 2016). Moreover, possible interactions between

lipids and proteins, including enzymes, in xylem conduits need to be investigated. Even in the

absence of gas in conduits, lipids accumulated in pit membranes are likely to affect sap flow and

could be involved in the regulation of hydraulic conductance, the so-called ionic effect (Nardini

et al., 2011, Nardini *et al.*, 2012). Finally, if lipids are crucial for plant water transport, then they

should be found as well in gymnosperms, ferns, and perhaps even moss hydroids. The discovery,

or rather rediscovery (Scott *et al.*, 1960, Esau, 1965, Esau *et al.*, 1966, Wagner *et al.*, 2000), of

lipids in xylem conduits opens a vast field of new research questions.

FUNDING

The research was funded by research grants from the National Science Foundation (EAGER

IOS-1558108 and IOS- 1754850). SJ and LK acknowledge funding from the German Research

Foundation (DFG, project No. 383393940).

ACKNOWLEDGEMENTS

The authors thank Anne Basilio, Zoe Cuevas, Jessica Garcia, Ryan Cochoit, Alec Hunt, and Tilly

Duong for assistance with xylem sap extractions and vessel volume analyses, Jim Henrich at the

Los Angeles County Arboretum and Greg Pongetti at the Fullerton Arboretum for allowing

sampling of plants in their living collections, the Electron Microscopy Section of Ulm University

for preparing TEM samples, and the Core Facility for Confocal and Multiphoton Microscopy of

Ulm University for assistance with confocal imaging.

Table 1. Composition of xylem sap lipids (in nmol / mL) for the seven studied species determined

by mass spectrometry, arranged by head groups, and expressed as mean ± standard deviation (n =

3). LysoPG was excluded from these measurements, because trace amounts of a contaminant that

was also found in solvent interfered with the detection of LPG in stage 1 measurements.

	Laurus		*Liriodendron*		*Bambusa*		*Triadica*		*Geijera*		*Distictis*		*Encelia*	
	Mean	STD	Mean	STD	Mean	STD	mean	STD	mean	STD	mean	STD	Mean	STD
							Xylem sap							
DGDG	0.038	± 0.013	0.015	± 0.010	0.203	± 0.046	0.051	± 0.009	0.015	± 0.006	0.041	± 0.01	0.015	± 0.008
MGDG	0.099	± 0.052	0.050	± 0.041	0.158	± 0.022	0.042	± 0.007	0.032	± 0.012	0.050	± 0.016	0.021	± 0.010
PG	0.001	± 0.000	0.002	± 0.002	0.003	± 0.001	0.007	± 0.002	0.003	± 0.003	0.003	± 0.001	0.004	± 0.004
LysoPC	0.000	± 0.000	0.003	± 0.003	0.006	± 0.003	0.004	± 0.002	0.001	± 0.001	0.001	± 0.001	0.007	± 0.004
LysoPE	0.002	± 0.002	0.001	± 0.001	0.003	± 0.002	0.002	± 0.002	0.001	± 0.001	0.001	± 0.001	0.005	± 0.005
PC	0.022	± 0.003	0.040	± 0.017	0.088	± 0.046	0.148	± 0.042	0.070	± 0.031	0.087	± 0.032	0.071	± 0.053
PE	0.007	± 0.001	0.016	± 0.014	0.026	± 0.020	0.042	± 0.042	0.051	± 0.016	0.046	± 0.018	0.046	± 0.055
PI	0.007	± 0.003	0.009	± 0.009	0.027	± 0.009	0.068	± 0.018	0.035	± 0.027	0.051	± 0.030	0.071	± 0.081
PS	0.001	± 0.000	0.004	± 0.003	0.007	± 0.003	0.011	± 0.006	0.016	± 0.007	0.013	± 0.008	0.015	± 0.015
PA	0.004	± 0.002	0.041	± 0.03	0.112	± 0.049	0.120	± 0.047	0.223	± 0.125	0.153	± 0.090	0.262	± 0.144
Total	0.181	± 0.071	0.179	± 0.124	0.633	± 0.166	0.494	± 0.177	0.447	± 0.213	0.447	± 0.203	0.517	± 0.371
SQDG	n.d.		0.001	± 0.001	n.d.		0.003	± 0.000	0.003	± 0.001	0.003	± 0.001	0.009	± 0.003
							Contamination controls							
DGDG	0.003	± 0.001	0.005	± 0.007	0.005	± 0.004	0.001	± 0.001	0.001	± 0.002	0.002	± 0.001	0.000	± 0.000
MGDG	0.004	± 0.002	0.004	± 0.004	0.010	± 0.009	0.001	± 0.000	0.002	± 0.001	0.002	± 0.001	0.001	± 0.000
PG	0.000	± 0.000	0.000	± 0.000	0.000	± 0.001	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000
LysoPC	0.000	± 0.000	0.000	± 0.000	0.001	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000
LysoPE	0.003	± 0.002	0.000	± 0.000	0.001	± 0.001	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000
PC	0.002	± 0.002	0.003	± 0.001	0.008	± 0.013	0.008	± 0.005	0.007	± 0.001	0.003	± 0.000	0.012	± 0.004
PE	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.002	± 0.002	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000
PI	0.001	± 0.001	0.000	± 0.000	0.001	± 0.001	0.002	± 0.003	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000
PS	0.001	± 0.001	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.000	0.000	± 0.001	0.000	± 0.000
PA	0.000	± 0.000	0.007	± 0.004	0.002	± 0.002	0.011	± 0.006	0.009	± 0.006	0.002	± 0.000	0.006	± 0.001
Total	0.014	± 0.008	0.020	± 0.015	0.029	± 0.029	0.025	± 0.016	0.020	± 0.008	0.009	± 0.003	0.021	± 0.003
SQDG	n.d.		0.001	± 0.001	n.d.		0.003	±0.000	0.003	± 0.001	0.003	± 0.001	0.009	± 0.003

Galactolipids: DGDG = digalactosyldiacylglycerol, MGDG = and monogalactosyldiacylglycerol. Phospholipids:
LysoPG = lysophosphatidylglycerol, LysoPC = lysophosphatidylcholine, LysoPE = lysophosphatidylethanolamine, PA
= phosphatidic acid, PC = phosphatidylcholine, PE = phosphatidylethanolamine, PI = phosphatidylinositol, PS =
phosphatidylserine, PG = phosphatidylglycerol. SQDG = Sulfolipids.

Table 2. Comparison of lipid composition categorized by headgroups in polar lipids in xylem sap

of *Geijera* (n = 8) sampled in July 2019 and March 2020. Mean concentrations (nmol / mL)

were compared between dates using FDR-adjusted t-tests in Metaboanalyst.ca.

Lipid type	July 2019 Mean ± STD	March 2020 Mean ± STD	p-value (FDR adjusted)
DGDG	0.0143 ± 0.0046	0.0589 ± 0.0284	0.00027
MGDG	0.0230 ± 0.0096	0.0472 ± 0.0193	0.03036
PG	0.0137 ± 0.0059	0.0174 ± 0.0097	n.s.
LysoPG	0.0074 ± 0.0025	0.0072 ± 0.0045	n.s.
LysoPC	0.0033 ± 0.0024	0.0018 ± 0.0004	n.s.
LysoPE	0.0018 ± 0.0011	0.0009 ± 0.0002	n.s.
PC	0.0851 ± 0.0407	0.0860 ± 0.0348	n.s.
PE	0.0550 ± 0.0368	0.0512 ± 0.0181	n.s.
PI	0.0185 ± 0.0114	0.0224 ± 0.0119	n.s.
PS	0.0148 ± 0.0068	0.0114 ± 0.0031	n.s.
PA	0.5329 ± 0.3504	0.3050 ± 0.1206	n.s.
Total Polar	0.7698 ± 0.4546	0.6094 ± 0.2025	n.s.

713 Table 3. Comparison of lipid composition (in %) in paired samples (n = 4) of xylem sap and
714 wood from the same stems collected in March 2020. All concentrations were converted to
715 percentages of total polar lipids. Wood and sap samples are not independent from each other, so
716 statistical comparisons were not possible.

	Geijera		Distictis	
	Sap	Wood	Sap	Wood
	mean STD	mean STD	Mean STD	mean STD
DGDG	9.52 ± 0.84	7.23 ± 1.08	4.53 ± 1.93	6.05 ± 1.00
MGDG	7.74 ± 0.81	8.00 ± 1.54	4.33 ± 1.43	5.66 ± 0.78
PG	2.45 ± 0.55	5.78 ± 1.38	1.26 ± 0.30	2.73 ± 1.00
LysoPG	1.44 ± 1.66	0.16 ± 0.06	1.23 ± 0.75	0.50 ± 0.41
LysoPC	0.31 ± 0.15	0.18 ± 0.07	0.21 ± 0.07	0.50 ± 0.10
LysoPE	0.17 ± 0.08	0.15 ± 0.03	0.18 ± 0.08	0.30 ± 0.03
PC	11.64 ± 1.94	14.62 ± 5.38	23.55 ± 3.06	14.57 ± 3.23
PE	7.39 ± 2.40	10.66 ± 2.39	8.47 ± 1.37	9.01 ± 1.76
PI	3.73 ± 1.51	6.16 ± 0.83	7.35 ± 0.71	9.00 ± 0.87
PS	1.88 ± 0.62	1.57 ± 0.52	2.00 ± 0.77	1.68 ± 0.39
PA	53.74 ± 16.80	45.51 ± 4.85	46.88 ± 2.59	50.02 ± 3.85
Total	100	100	100	100

LITERATURE CITED

Aki, T., Shigyo, M., Nakano, R., Yoneyama, T. and Yanagisawa, S. (2008) Nano scale proteomics

revealed the presence of regulatory proteins including three FT-like proteins in phloem and xylem

saps from rice. *Plant & Cell Physiol.*, **49**, 767-790. https://doi.org/10.1093/pcp/pcn049

Alvarez, S., Goodger, J.Q.D., Marsh, E.L., Chen, S., Asirvatham, V.S. and Schachtman, D.P. (2006)

Characterization of the maize xylem sap proteome. *J. Proteome Res.*, **5**, 963-972.

https://doi.org/10.1021/pr050471q

Angeles, G., Bond, B., Boyer, J.S., Brodribb, T., Brooks, J.R., Burns, M.J., Cavender-Bares, J.,

Clearwater, M., Cochard, H., Comstock, J., Davis, S.D., Domec, J.C., Donovan, L., Ewers,

F., Gartner, B., Hacke, U., Hinckley, T., Holbrook, N.M., Jones, H.G., Kavanagh, K., Law,

B., Lopez-Portillo, J., Lovisolo, C., Martin, T., Martinez-Vilalta, J., Mayr, S., Meinzer, F.C.,

Melcher, P., Mencuccini, M., Mulkey, S., Nardini, A., Neufeld, H.S., Passioura, J.,

Pockman, W.T., Pratt, R.B., Rambal, S., Richter, H., Sack, L., Salleo, S., Schubert, A.,

Schulte, P., Sparks, J.P., Sperry, J., Teskey, R. and Tyree, M. (2004) The cohesion-tension

theory. *New Phytol.*, **163**, 451-452. https://doi.org/10.1111/j.1469-8137.2004.01142.x

Askenasy, E. (1895) Ueber das Saftsteigen. *Verh. Nat.-med. Ver. Heidelb. N. F.*, **5**, 325-345

Bargmann, B.O.R., Laxalt, A.M., Riet, B.T., Testerink, C., Merquiol, E., Mosblech, A., Leon-Reyes,

A., Pieterse, C.M.J., Haring, M.A., Heilmann, I., Bartels, D. and Munnik, T. (2009)

Reassessing the role of phospholipase D in the *Arabidopsis* wounding response. *Plant Cell*

Environ., **32**, 837-850. https://doi.org/10.1111/j.1365-3040.2009.01962.x

Bartlett, M.K., Klein, T., Jansen, S., Choat, B. and Sack, L. (2016) The correlations and sequence of

plant stomatal, hydraulic, and wilting responses to drought. *Proc. Natl. Acad. Sci.*, **113**, 13098-

13103. https://doi.org/10.1073/pnas.1604088113

Botté, C.Y. and Maréchal, E. (2014) Plastids with or without galactoglycerolipids. *Trends Plant Sci.*,

19, 71-78. https://doi.org/10.1016/j.tplants.2013.10.004

Bottier, C., Géan, J., Artzner, F., Desbat, B., Pézolet, M., Renault, A., Marion, D. and Vié, V. (2007)

Galactosyl headgroup interactions control the molecular packing of wheat lipids in Langmuir

films and in hydrated liquid-crystalline mesophases. *Biochim. Biophys. Acta (BBA) -*

Biomembranes, **1768**, 1526-1540. https://doi.org/10.1016/j.bbamem.2007.02.021

Brodribb, T.J., Bienaimé, D. and Marmottant, P. (2016) Revealing catastrophic failure of leaf

networks under stress. *Proc. Natl. Acad. Sci.*, **113**, 4865-4869.

https://doi.org/10.1073/pnas.1522569113

Buhtz, A., Kolasa, A., Arlt, K., Walz, C. and Kehr, J. (2004) Xylem sap protein composition is

conserved among different plant species. *Planta*, **219**, 610-618. https://doi.org/10.1007/s00425-

004-1259-9

Chafe, S.C. and Chauret, G. (1974) Cell wall structure in the xylem parenchyma of trembling aspen.

Protopl., **80**, 129-147. https://doi.org/10.1007/bf01666355

Charrier, G., Delzon, S., Domec, J.-C., Zhang, L., Delmas, C.E., Merlin, I., Corso, D., King, A.,

Ojeda, H. and Ollat, N. (2018) Drought will not leave your glass empty: Low risk of hydraulic

failure revealed by long-term drought observations in world's top wine regions. *Sci. Adv.*, **4**,

eaao6969. https://doi.org/10.1126/sciadv.aao6969

Choat, B., Ball, M., Luly, J. and Holtum, J. (2003) Pit membrane porosity and water stress-induced

cavitation in four co-existing dry rainforest tree species. *Plant Physiol.*, **131**, 41–48.

https://doi.org/10.1104/pp.014100

Christie, W.W. (1993) Preparation of lipid extracts from tissues. In *Advances in Lipid Methodology –*

Two (Christie, W.W. ed. Dundee: Oily Press, pp. 195-213.

Czaninski, Y. (1977) Vessel-associated cells. *IAWA Bull.*, **1977**, 51-55

Devaiah, S.P., Roth, M.R., Baughman, E., Li, M., Tamura, P., Jeannotte, R., Welti, R. and Wang,

X. (2006) Quantitative profiling of polar glycerolipid species from organs of wild-type

Arabidopsis and a PHOSPHOLIPASE Dα1 knockout mutant. *Phytochem.*, **67**, 1907-1924.

https://doi.org/10.1016/j.phytochem.2006.06.005

Dixon, H.H. (1914) On the tensile strength of sap. *Sci. Proc. Royal Dubl. Soc. N. S.*, **14**, 229-234

Dixon, H.H. and Joly, J. (1895) On the ascent of sap. *Phil. Trans. R. Soc. Lond. B*, **186**, 563-576

Djordjevic, M.A., Oakes, M., Li, D.X., Hwang, C.H., Hocart, C.H. and Gresshoff, P.M. (2007) The

Glycine max xylem sap and apoplast proteome. *J. Proteome Res.*, **6**, 3771-3779.

https://doi.org/10.1021/pr0606833

Domínguez, E., Heredia-Guerrero, J.A. and Heredia, A. (2015) Plant cutin genesis: unanswered

questions. *Trends Plant Sci.*, **20**, 551-558. https://doi.org/10.1016/j.tplants.2015.05.009

Dörmann, P. and Benning, C. (2002) Galactolipids rule in seed plants. *Trends Plant Sci.*, **7**, 112-118.

https://doi.org/10.1016/S1360-1385(01)02216-6

Dugé de Bernonville, T., Albenne, C., Arlat, M., Hoffmann, L., Lauber, E. and Jamet, E. (2014)

Xylem sap proteomics. In *Plant Proteomics* (Jorrin-Novo, J.V., Komatsu, S., Weckwerth, W. and

Wienkoop, S. eds): Humana Press, pp. 391-405.

Ellis, E.A. (2014) Staining sectioned biological specimens for transmission electron microscopy:

conventional and en bloc stains. In *Electron Microscopy: Methods and Protocols* (Kuo, J. ed.

New York: Humana Press, pp. 57-72.

Ellis, S.R., Brown, S.H., in het Panhuis, M., Blanksby, S.J. and Mitchell, T.W. (2013) Surface

analysis of lipids by mass spectrometry: More than just imaging. *Progr. Lid. Res.*, **52**, 329-353.

https://doi.org/10.1016/j.plipres.2013.04.005

Esau, K. (1965) *Plant anatomy* 2nd edn. New York: John Wiley & Sons.

Esau, K., Cheadle, V.I. and Gill, R.H. (1966) Cytology of differentiating tracheary elements. II.

Structures associated with cell surfaces. *Amer. J. Bot.*, **53**, 765-771.

https://doi.org/10.1002/j.1537-2197.1966.tb06832.x

Espino, S. and Schenk, H.J. (2011) Mind the bubbles: Achieving stable measurements of maximum

hydraulic conductivity through woody plant samples. *J. Exp. Bot.*, **62**, 1119-1132.

https://doi.org/10.1093/jxb/erq338

Fich, E.A., Segerson, N.A. and Rose, J.K.C. (2016) The plant polyester cutin: Biosynthesis, structure,

and biological roles. *Annu. Rev. Plant Biol.*, **67**, 207-233. https://doi.org/10.1146/annurev-arplant-

043015-111929

Fineran, B.A. (1997) Cyto- and histochemical demonstration of lignins in plant cell walls: an evaluation

of the chlorine water ethanolamine silver nitrate method of Coppick and Fowler. *Protopl.*, **198**,

186-201. https://doi.org/10.1007/BF01287568

Fujii, T., Harada, H. and Saiki, H. (1981) Ultrastructure of "amorphous layer" in xylem parenchyma

cell wall of angiosperm species. *Mokuzai Gakkaishi*, **27**, 149-156

Furse, S., Liddell, S., Ortori, C.A., Williams, H., Neylon, D.C., Scott, D.J., Barrett, D.A. and Gray,

D.A. (2013) The lipidome and proteome of oil bodies from *Helianthus annuus* (common

sunflower). *J. Chem. Biol.*, **6**, 63-76. https://doi.org/10.1007/s12154-012-0090-1

Gascó, A., Nardini, A., Gortan, E. and Salleo, S. (2006) Ion-mediated increase in the hydraulic

conductivity of Laurel stems: role of pits and consequences for the impact of cavitation on water

transport. *Plant Cell Environ.*, **29**, 1946-1955. https://doi.org/10.1111/j.1365-3040.2006.01570.x

Gonorazky, G., Laxalt, A.M., Dekker, H.L., Rep, M., Munnik, T., Testerink, C. and de la Canal, L.

(2012) Phosphatidylinositol 4-phosphate is associated to extracellular lipoproteic fractions and is

detected in tomato apoplastic fluids. *Plant Biol.*, **14**, 41-49. https://doi.org/10.1111/j.1438-

8677.2011.00488.x

Greenidge, K.N.H. (1952) An approach to the study of vessel length in hardwood species. *Amer. J. Bot.*,

39, 570-574. https://doi.org/10.1002/j.1537-2197.1952.tb13070.x

Hacke, U.G. and Sperry, J.S. (2003) Limits to xylem refilling under negative pressure in *Laurus nobilis*

and *Acer negundo*. *Plant Cell Environ.*, **26**, 303-311. https://doi.org/10.1046/j.1365-

3040.2003.00962.x

Hammer, Ø., Harper, D.A.T. and Ryan, P.D. (2001) PAST: Paleontological statistics software package

for education and data analysis. *Palaeont. Etectron.*, **4**, 9

Huynh, V.-B., Repellin, A., Zuily-Fodil, Y. and Pham-Thi, A.-T. (2012) Aluminum stress response in

rice: effects on membrane lipid composition and expression of lipid biosynthesis genes. *Physiol.*

Plant., **146**, 272-284. https://doi.org/10.1111/j.1399-3054.2012.01622.x

Jansen, S., Klepsch, M., Li, S., Kotowska, M.M., Schiele, S., Zhang, Y. and Schenk, H.J. (2018)

Challenges in understanding air-seeding in angiosperm xylem. *Acta Hort.*, **1222**, 13-20.

https://doi.org/10.17660/ActaHortic.2018.1222.3

Jansen, S. and Schenk, H.J. (2015) On the ascent of sap in the presence of bubbles. *Amer. J. Bot.*, **102**,

1561-1563. https://doi.org/10.3732/ajb.1500305

Jelínková, A., Malínská, K., Simon, S., Kleine-Vehn, J., Pařezová, M., Pejchar, P., Kubeš, M.,

Martinec, J., Friml, J., Zažímalová, E. and Petrášek, J. (2010) Probing plant membranes with

FM dyes: tracking, dragging or blocking? *Plant J.*, **61**, 883-892. https://doi.org/10.1111/j.1365-

313X.2009.04102.x

Kaack, L., Altaner, C.M., Carmesin, C., Diaz, A., Holler, M., Kranz, C., Neusser, G., Odstrcil, M.,

Schenk, H.J., Schmidt, V., Weber, M., Zhang, Y. and Jansen, S. (2019) Function and three-

dimensional structure of intervessel pit membranes in angiosperms: a review. *IAWA J.*, **40**, 673-

702. https://doi.org/10.1163/22941932-40190259

Kanduč, M., Schlaich, A., de Vries, A.H., Jouhet, J., Maréchal, E., Demé, B., Netz, R.R. and

Schneck, E. (2017) Tight cohesion between glycolipid membranes results from balanced water–

headgroup interactions. *Nat. Comm.*, **8**, 14899. https://doi.org/10.1038/ncomms14899

Kanduč, M., Schlaich, A., Schneck, E. and Netz, R.R. (2016) Water-mediated interactions between

hydrophilic and hydrophobic surfaces. *Langmuir*, **32**, 8767-8782.

https://doi.org/10.1021/acs.langmuir.6b01727

Kanduč, M., Schneck, E., Loche, P., Jansen, S., Schenk, H.J. and Netz, R.R. (2020) Cavitation in

lipid bilayers poses strict negative pressure stability limit in biological liquids. *Proc. Natl. Acad.*

Sci. USA, **117**, 10733-10739. https://doi.org/10.1073/pnas.1917195117

Kanduč, M., Schneck, E. and Netz, R.R. (2013) Hydration interaction between phospholipid

membranes: Insight into different measurement ensembles from atomistic molecular dynamics

simulations. *Langmuir*, **29**, 9126-9137. https://doi.org/10.1021/la401147b

Kates, M. (1957) Effects of solvents and surface-active agents on plastid phosphatidase C activity. *Can.*

J. Biochem. Physiol., **35**, 127-142. https://doi.org/10.1139/o57-016

Kolesnikov, Y.S., Nokhrina, K.P., Kretynin, S.V., Volotovski, I.D., Martinec, J., Romanov, G.A.

and Kravets, V.S. (2012) Molecular structure of phospholipase D and regulatory mechanisms of

its activity in plant and animal cells. *Biochem. (Moscow)*, **77**, 1-14.

https://doi.org/10.1134/s0006297912010014

Krasikov, V., Dekker, H.L., Rep, M. and Takken, F.L.W. (2011) The tomato xylem sap protein

XSP10 is required for full susceptibility to *Fusarium* wilt disease. *J. Exp. Bot.*, **62**, 963-973.

https://doi.org/10.1093/jxb/erq327

Krishnan, H.B., Natarajan, S.S., Bennett, J.O. and Sicher, R.C. (2011) Protein and metabolite

composition of xylem sap from field-grown soybeans (*Glycine max*). *Planta*, **233**, 921-931.

https://doi.org/10.1007/s00425-011-1352-9

Kwan, J.J. and Borden, M.A. (2010) Microbubble dissolution in a multigas environment. *Langmuir*, **26**,

6542-6548. https://doi.org/10.1021/la904088p

Li, N., Xu, C., Li-Beisson, Y. and Philippar, K. (2016) Fatty acid and lipid transport in plant cells.

Trends Plant Sci., **21**, 145-158. https://doi.org/10.1016/j.tplants.2015.10.011

Ligat, L., Lauber, E., Albenne, C., Clemente, H.S., Valot, B., Zivy, M., Pont-Lezica, R., Arlat, M.

and Jamet, E. (2011) Analysis of the xylem sap proteome of *Brassica oleracea* reveals a high

content in secreted proteins. *Proteomics*, **11**, 1798-1813. https://doi.org/10.1002/pmic.201000781

Lima, M.R.M., Machado, A.F. and Gubler, W.D. (2017) Metabolomic study of Chardonnay

grapevines double stressed with esca-associated fungi and drought. *Phytopathol.*, **107**, 669-680.

https://doi.org/10.1094/phyto-11-16-0410-r

Martínez-Cabrera, H.I., Jones, C.S., Espino, S. and Schenk, H.J. (2009) Wood anatomy and wood

density in shrubs: Responses to varying aridity along transcontinental transects. *Amer. J. Bot.*, **96**,

1388-1398. https://doi.org/10.3732/ajb.0800237

McLoughlin, F. and Testerink, C. (2013) Phosphatidic acid, a versatile water-stress signal in roots.

Front. Plant Sci., **4**, 525. https://doi.org/10.3389/fpls.2013.00525

Misra, B.B. (2016) The black-box of plant apoplast lipidomes. *Front. Plant Sci.*, **7**, 323.

https://doi.org/10.3389/fpls.2016.00323

Mongrand, S., Bessoule, J.-J., Cabantous, F. and Cassagne, C. (1998) The C16:3\C18:3 fatty acid

balance in photosynthetic tissues from 468 plant species. *Phytochem.*, **49**, 1049-1064.

https://doi.org/10.1016/S0031-9422(98)00243-X

Morris, H., Gillingham, M.A.F., Plavcová, L., Gleason, S.M., Olson, M.E., Coomes, D.A., Fichtler,

E., Klepsch, M.M., Martínez-Cabrera, H.I., McGlinn, D.J., Wheeler, E.A., Zheng, J.,

Ziemińska, K. and Jansen, S. (2018a) Vessel diameter is related to amount and spatial

arrangement of axial parenchyma in woody angiosperms. *Plant Cell Environ.*, **41**, 245-260.

https://doi.org/10.1111/pce.13091

Morris, H. and Jansen, S. (2016) Secondary xylem parenchyma - from classical terminology to

functional traits. *IAWA J.*, **37**, 1-15. https://doi.org/10.1163/22941932-20160117

Morris, H., Plavcová, L., Cvecko, P., Fichtler, E., Gillingham, M.A.F., Martínez-Cabrera, H.I.,

McGlinn, D.J., Wheeler, E., Zheng, J., Ziemińska, K. and Jansen, S. (2016) A global analysis

of parenchyma tissue fractions in secondary xylem of seed plants. *New Phytol.*, **209**, 1553-1565.

https://doi.org/10.1111/nph.13737

Morris, H., Plavcová, L., Gorai, M., Klepsch, M., Kotowska, M., Schenk, H.J. and Jansen, S.

(2018b) Vessel-associated cells in angiosperm xylem: Highly specialized living cells at the

symplast-apoplast boundary. *Amer. J. Bot.*, **105**, 153-162. https://doi.org/doi:10.1002/ajb2.1030

Nardini, A., Dimasi, F., Klepsch, M. and Jansen, S. (2012) Ion-mediated enhancement of xylem

hydraulic conductivity in four *Acer* species: relationships with ecological and anatomical features.

Tree Physiol., **32**, 1434-1441. https://doi.org/10.1093/treephys/tps107

Nardini, A., Salleo, S. and Jansen, S. (2011) More than just a vulnerable pipeline: xylem physiology in

the light of ion-mediated regulation of plant water transport. *J. Exp. Bot.*, **62**, 4701-4718.

https://doi.org/10.1093/jxb/err208

Nardini, A., Savi, T., Losso, A., Petit, G., Pacilè, S., Tromba, G., Mayr, S., Trifilò, P., Lo Gullo,

M.A. and Salleo, S. (2017) X-ray microtomography observations of xylem embolism in stems of

Laurus nobilis are consistent with hydraulic measurements of percentage loss of conductance.

New Phytol., **213**, 1068-1075. https://doi.org/10.1111/nph.14245

Oertli, J.J. (1971) The stability of water under tension in the xylem. *Z. Pflanzenphysiol.*, **65**, 195-209

Ohki, S. and Ohshima, H. (1985) Divalent cation-induced phosphatidic acid membrane fusion. Effect of

ion binding and membrane surface tension. *Biochim. Biophys. Acta*, **812**, 147-154.

https://doi.org/10.1016/0005-2736(85)90532-2

Park, J., Go, T., Ryu, J. and Lee, S.J. (2019) Air spreading through wetted cellulose membranes:

Implications for the safety function of hydraulic valves in plants. *Phys. Rev. E*, **100**, 032409.

https://doi.org/10.1103/PhysRevE.100.032409

Schenk, H.J., Espino, S., Rich-Cavazos, S.M. and Jansen, S. (2018) From the sap's perspective: the

nature of vessel surfaces in angiosperm xylem. *Amer. J. Bot.*, **105**, 174-187.

https://doi.org/10.1002/ajb2.1034

Schenk, H.J., Espino, S., Romo, D.M., Nima, N., Do, A.Y.T., Michaud, J.M., Papahadjopoulos-

Sternberg, B., Yang, J., Zuo, Y., Steppe, K. and Jansen, S. (2017) Xylem surfactants introduce

a new element to the cohesion-tension theory. *Plant Physiol.*, **173**, 1177-1196.

https://doi.org/10.1104/pp.16.01039

Schenk, H.J., Espino, S., Visser, A. and Esser, B.K. (2016) Dissolved atmospheric gas in xylem sap

measured with membrane inlet mass spectrometry. *Plant Cell Environ.*, **39**, 944-950.

https://doi.org/10.1111/pce.12678

Schenk, H.J., Steppe, K. and Jansen, S. (2015) Nanobubbles: a new paradigm for air-seeding in xylem.

Trends Plant Sci., **20**, 199-205. https://doi.org/10.1016/j.tplants.2015.01.008

Schmid, R. and Machado, R.D. (1968) Pit membranes in hardwoods—Fine structure and development.

Protopl., **66**, 185-204. https://doi.org/10.1007/BF01252532

Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of image

analysis. *Nat. Meth.*, **9**, 671. https://doi.org/10.1038/nmeth.2089

Schneider, H., Manz, B., Westhoff, M., Mimietz, S., Szimtenings, M., Neuberger, T., Faber, C.,

Krohne, G., Haase, A., Volke, F. and Zimmermann, U. (2003) The impact of lipid distribution,

composition and mobility on xylem water refilling of the resurrection plant Myrothamnus

flabellifolia. *New Phytol.*, **159**, 487-505. https://doi.org/10.1046/j.1469-8137.2003.00814.x

Schurr, U. (1998) Xylem sap sampling – new approaches to an old topic. *Trends Plant Sci.*, **3**, 293-298.

https://doi.org/10.1016/S1360-1385(98)01275-8

Scott, F.M., Sjaholm, V. and Bowler, E. (1960) Light and electron microscope studies of the primary

xylem of *Ricinus communis*. *Amer. J. Bot.*, **47**, 162-173. https://doi.org/10.2307/2439245

Shimojima, M. (2011) Biosynthesis and functions of the plant sulfolipid. *Progr. Lid. Res.*, **50**, 234-239.

https://doi.org/10.1016/j.plipres.2011.02.003

Shiva, S., Enninful, R., Roth, M.R., Tamura, P., Jagadish, K. and Welti, R. (2018) An efficient

modified method for plant leaf lipid extraction results in improved recovery of phosphatidic acid.

Plant Methods, **14**, 14. https://doi.org/10.1186/s13007-018-0282-y

Shiva, S., Vu, H.S., Roth, M.R., Zhou, Z., Marepally, S.R., Nune, D.S., Lushington, G.H.,

Visvanathan, M. and Welti, R. (2013) Lipidomic analysis of plant membrane lipids by direct

infusion tandem mass spectrometry. *Methods Mol. Biol.*, **1009**, 79-91.

https://doi.org/10.1007/978-1-62703-401-2_9

Smith, A.M. (1994) Xylem transport and the negative pressures sustainable by water. *Ann. Bot.*, **74**, 647-

651. https://doi.org/10.1006/anbo.1994.1166

Sperry, J.S., Hacke, U.G. and Wheeler, J.K. (2005) Comparative analysis of end wall resistivity in

xylem conduits. *Plant Cell Environ.*, **28**, 456-465. https://doi.org/10.1111/j.1365-

3040.2005.01287.x

Sperry, J.S. and Tyree, M.T. (1988) Mechanism of water stress-induced xylem embolism. *Plant*

Physiol., **88**, 581-587. https://doi.org/10.1104/pp.88.3.581

Stroock, A.D., Pagay, V.V., Zwieniecki, M.A. and Holbrook, N.M. (2014) The physicochemical

hydrodynamics of vascular plants. *Annu. Rev. Fluid Mechan.*, **46**, 615-642.

https://doi.org/10.1146/annurev-fluid-010313-141411

Su, W., Yeku, O., Olepu, S., Genna, A., Park, J.-S., Ren, H., Du, G., Gelb, M.H., Morris, A.J. and

Frohman, M.A. (2009) 5-Fluoro-2-indolyl des-chlorohalopemide (FIPI), a Phospholipase D

pharmacological inhibitor that alters cell spreading and inhibits chemotaxis. *Molec. Pharmacol.*,

75, 437-446. https://doi.org/10.1124/mol.108.053298

Testerink, C. and Munnik, T. (2011) Molecular, cellular, and physiological responses to phosphatidic

acid formation in plants. *J. Exp. Bot.*, **62**, 2349-2361. https://doi.org/10.1093/jxb/err079

Tyree, M.T., Salleo, S., Nardini, A., Lo Gullo, M.A. and Mosca, R. (1999) Refilling of embolized

vessels in young stems of laurel. Do we need a new paradigm? *Plant Physiol.*, **120**, 11-21.

https://doi.org/10.1104/pp.120.1.11

Tyree, M.T. and Zimmermann, M.H. (2002) *Xylem structure and the ascent of sap* 2nd edn. Berlin:

Springer-Verlag.

Venturas, M.D., Sperry, J.S. and Hacke, U.G. (2017) Plant xylem hydraulics: What we understand,

current research, and future challenges. *J. Integr. Plant Biol.*, **59**, 356-389.

https://doi.org/doi:10.1111/jipb.12534

Wagner, H.J., Schneider, H., Mimietz, S., Wistuba, N., Rokitta, M., Krohne, G., Haase, A. and

Zimmermann, U. (2000) Xylem conduits of a resurrection plant contain a unique lipid lining and

refill following a distinct pattern after desiccation. *New Phytol.*, **148**, 239-255.

https://doi.org/10.1046/j.1469-8137.2000.00755.x

Welti, R., Li, W., Li, M., Sang, Y., Biesiada, H., Zhou, H.-E., Rajashekar, C.B., Williams, T.D. and

Wang, X. (2002) Profiling membrane lipids in plant stress responses: Role of phospholipase Da

in freezing-induced lipid changes in *Arabidopsis*. *J. Biol. Chem.*, **277**, 31994-32002.

https://doi.org/10.1074/jbc.M205375200

Weschayanwiwat, P., Scamehorn, J. and Reilly, P. (2005) Surfactant properties of low molecular

weight phospholipids. *J. Surfact. Deterg.*, **8**, 65-72. https://doi.org/10.1007/s11743-005-0332-8

Westhoff, M., Schneider, H., Zimmermann, D., Mimietz, S., Stinzing, A., Wegner, L.H., Kaiser, W.,

Krohne, G., Shirley, S., Jakob, P., Bamberg, E., Bentrup, F.W. and Zimmermann, U. (2008)

The mechanisms of refilling of xylem conduits and bleeding of tall birch during spring. *Plant*

Biol., **10**, 604-623. https://doi.org/10.1111/j.1438-8677.2008.00062.x

Wheeler, J.K., Sperry, J.S., Hacke, U.G. and Huang, N. (2005) Inter-vessel pitting and cavitation in

woody Rosaceae and other vesseled plants: a basis for a safety vs. efficiency trade-off in xylem

transport. *Plant Cell Environ.*, **28**, 800-812. https://doi.org/10.1111/j.1365-3040.2005.01330.x

Wheeler, T.D. and Stroock, A.D. (2008) The transpiration of water at negative pressures in a synthetic

tree. *Nature*, **455**, 208-212. https://doi.org/10.1038/nature07226

Wisniewski, M. and Davis, G. (1995) Immunogold localization of pectins and glycoproteins in tissues of

peach with reference to deep supercooling. *Trees - Struct. Funct.*, **9**, 253-260.

https://doi.org/10.1007/BF00202015

Wisniewski, M., Davis, G. and Arora, R. (1991a) Effect of macerase, oxalic acid, and EGTA on deep

supercooling and pit membrane structure of xylem parenchyma of peach. *Plant Physiol.*, **96**,

1354-1359. https://doi.org/10.1104/pp.96.4.1354

Wisniewski, M., Davis, G. and Schaffer, K. (1991b) Mediation of deep supercooling of peach and

dogwood by enzymatic modifications in cell-wall structure. *Planta*, **184**, 254-260.

https://doi.org/10.1007/BF00197955

Yang, J., Michaud, J.M., Jansen, S., Schenk, H.J. and Zuo, Y.Y. (2020) Dynamic surface tension of

xylem sap lipids. *Tree Physiol.*, **40**, 433-444. https://doi.org/10.1093/treephys/tpaa006

Zhang, H., Fan, Q., Wang, Y.E., Neal, C.R. and Zuo, Y.Y. (2011) Comparative study of clinical

pulmonary surfactants using atomic force microscopy. *Biochim. Biophys. Acta (BBA) -*

Biomembranes, **1808**, 1832-1842. https://doi.org/10.1016/j.bbamem.2011.03.006

Zhang, Y., Carmesin, C., Kaack, L., Klepsch, M.M., Kotowska, M., Matei, T., Schenk, H.J., Weber,

M., Walther, P., Schmidt, V. and Jansen, S. (2020) High porosity with tiny pore constrictions

and unbending pathways characterise the 3D structure of intervessel pit membranes in

angiosperm xylem. *Plant Cell Environ.*, **43**, 116-130. https://doi.org/10.1111/pce.13654

Zhang, Y., Klepsch, M. and Jansen, S. (2017) Bordered pits in xylem of vesselless angiosperms and

their possible misinterpretation as perforation plates. *Plant Cell Environ.*, **40**, 2133-2146.

https://doi.org/10.1111/pce.13014

Zhou, Z., Marepally, S.R., Nune, D.S., Pallakollu, P., Ragan, G., Roth, M.R., Wang, L., Lushington,

G.H., Visvanathan, M. and Welti, R. (2011) LipidomeDB data calculation environment: online

processing of direct-infusion mass spectral data for lipid profiles. *Lipids*, **46**, 879-884.

https://doi.org/10.1007/s11745-011-3575-8

Zimmermann, U., Schneider, H., Wegner, L.H. and Haase, A. (2004) Water ascent in tall trees: does

evolution of land plants rely on a highly metastable state? *New Phytol.*, **162**, 575-615.

https://doi.org/10.1111/j.1469-8137.2004.01083.x

Zuo, Y.Y., Ding, M., Li, D. and Neumann, A.W. (2004) Further development of Axisymmetric Drop

Shape Analysis-Captive Bubble for pulmonary surfactant related studies. *Biochimica et*

Biophysica Acta (BBA) - General Subjects, **1675**, 12-20.

https://doi.org/10.1016/j.bbagen.2004.08.003

Zuo, Y.Y., Keating, E., Zhao, L., Tadayyon, S.M., Veldhuizen, R.A.W., Petersen, N.O. and

Possmayer, F. (2008) Atomic force microscopy studies of functional and dysfunctional

pulmonary surfactant films. I. Micro- and nanostructures of functional pulmonary surfactant films

and the effect of SP-A. *Biophys. J.*, **94**, 3549-3564. https://doi.org/10.1529/biophysj.107.122648

Zwieniecki, M.A., Melcher, P.J. and Holbrook, N.M. (2001) Hydrogel control of xylem hydraulic

resistance in plants. *Science*, **291**, 1059-1062. https://doi.org/10.1126/science.1057175

1025

FIGURE LEGENDS

FIG. 1. Experimental setup for xylem sap extraction via the vacuum method designed to ensure

that lipids in sap only come into contact with glass surfaces and are cooled immediately after

extraction to inhibit enzyme activity. Also shown are the proximal xylem surface at the

collection end of the stem from which cell contamination controls were collected after high-

pressure cleaning of the surface and the successive distal cuts that open vessels for vacuum

extraction. In lipid tracer (di17:0 PE) experiments, the tracer was added to the freshly cut distal

surfaces of the last three cuts (50, 45, and 40 cm stem length). See Methods section for detailed

explanations.

FIG. 2. Polar lipid composition of xylem sap from seven angiosperm species (n = 3). A. Chemical

composition of xylem sap lipids as determined by direct-infusion electrospray ionization triple-

quadrupole mass spectrometry. See Table S1 for the standards used and Figures S1-9 and Tables

S2-3 [**Supplementary Information**] for the complete data sets, including chains lengths and

degrees of saturation. Sulfolipids were only found in trace amounts (Tables S2 and S3) and are

not included in this figure. B. Most common numbers of acyl chain carbon atoms (mean ± SE) in

xylem sap lipids (PL = phospholipids, GL = galactolipids) and their degree of unsaturation as

determined by direct-infusion electrospray ionization triple-quadrupole mass spectrometry. See

Table S2 [**Supplementary Information**] and Figures S1-9 for the complete data set. C. Total

concentration of phospho- and galactolipids in xylem sap (mean ± SE) as a function of total

vessel xylem perimeter at the cut xylem surface (mean ± SE). Numbers under the abbreviated

generic names are the median vessel lengths in cm. Abbreviations: Bam = *Bambusa*, Dis =

Distictis, Enc = *Encelia*, Gei = *Geijera*, Lau = *Laurus*, Lir = *Liriodendron*, Tri = *Triadica*. D.

Polar lipid composition of xylem sap compared via principal component analysis (PCA), with

the first two components shown that explain 63% of the variation among species. Shaded areas

are 95% confidence regions.

FIG. 3. Polar lipid composition compared between xylem sap (n = 8) of *Geijera* sampled in July

2019 and March 2020. A. Results from a principal component analysis (PCA), with the first two

components shown. Shaded areas are 95% confidence regions. B. Heatmaps for polar lipids in,

based on Euclidean distance measures, a Ward clustering algorithm, and normalized data to

visualize differences between sampling times. The lipids shown are the 17 most different

between sampling times based on unpaired t-tests, adjusted for false discovery rates (FDR). All

lipids in this heatmap were significantly different ($p < 0.05$) between dates after FDR adjustment.

FIG. 4. Polar lipid composition compared between paired samples of xylem sap and wood of the

same stems in *Distictis* and *Geijera* (n = 4). A. and C. show results from principal component

analyses (PCA) for *Distictis* and *Geijera*, respectively, with the first two components shown.

Shaded areas are 95% confidence regions. B. and D. are heatmaps for polar lipids in *Distictis* and

Geijera, respectively, based on normalized data, Euclidean distance measures, and a Ward

clustering algorithm to visualize differences between xylem sap and wood. The lipids shown are

the 20 most different between sap and wood based on paired t-tests, adjusted for false discovery

rates (FDR). Lipids indicated in bold were flagged as different at $p < 0.05$ between xylem sap

and wood after paired t-tests with FDR adjustment. Note that lipids in xylem sap and wood

samples are not independent of each other, so t-tests were used only as qualitative indicators.

FIG. 5. Lipids in xylem and inter-vessel pits of *Laurus nobilis*. A-C: Super-resolution confocal

laser scanning microscopy images (false color) showing lignified cell walls in blue based on

lignin autofluorescence and FM1-43 dye bound to lipids in yellow. White arrows = intervessel

pit membranes. A-B: With FM1-43, C: Control without FM1-43. D-G: Transmission electron

micrographs of intervessel pits. D-E: without OsO_4; showing unusually dark, electron-dense

homogeneous pit membranes and linings on secondary walls, but no granular appearance. F-G:

with OsO_4, lipids visible as black aggregates lining the wall surface, pit chamber and inside the

pit membrane. Abbreviations: PA = pit aperture, PM = pit membrane, SW = secondary wall.

FIG. 6. Conceptual model of the formation of lipid-coated nanobubbles via snap-off inside pit

membranes that are coated and infiltrated with surface-active lipids. The different sizes of

micelles shown in this image reflect the wide range of nanoparticle sizes detected in xylem sap

and presumed to be bilayer to multilayer micelles, vesicles, and lipid-coated nanobubbles

(Schenk *et al*. 2017).

1082

1083

1084 SUPPLEMENTARY MATERIALS

Supporting figures:

1085

Figure S1. . Xylem sap concentrations of digalactosyldiacylglycerol (DGDG) in seven
angiosperms species.

Figure S2. Xylem sap concentrations of monogalactosyldiacylglycerol (MGDG) in seven
angiosperms species.

Figure S3. Xylem sap concentration of phosphatidylglycerol (PG) in seven angiosperms species.

Figure S4. Xylem sap concentration of phosphatidylcholine (PC) in seven angiosperms species.

Figure S5. Xylem sap concentration of phosphatidylethanolamine (PE) in seven angiosperms
species.

Figure S6. Xylem sap concentrations of phosphatidylinositol (PI) in seven angiosperms species.

Figures S7. Xylem sap concentrations of phosphatidylserine (PS) in seven angiosperms species.

Figures S8. Xylem sap concentrations of phosphatidic acid (PA) in seven angiosperms species.

Figure S9. Xylem sap concentrations of lysophosphatidylethanolamine (LPE) and
lysophosphatidylcholine (LPC) in seven angiosperms species.

Figure S10. Stems infiltrated with blue-colored silicone, cut at 0.7 cm from the injected surface.
The four spoke-like structures in *Distictis buccinatoria* xylem are areas of
interxylary phloem.

Figure S11. Total vessel area filled with silicone as a function of distance from the infiltrated
surface, shown for a sample of *Triadica sebifera*. The silicone-filled vessel volume
was calculated from these filled areas by linear interpolation.

Fig. S12. Principal component analyses (PCA) comparing polar lipid composition in xylem sap
for six angiosperm species to paired cell contamination controls from the same
stems.

Supporting tables:

1108

1109 Table S1. Standards used for mass spectrometry of xylem sap lipids.

1110 Table S2. Concentrations of polar lipids and sulfolipids in xylem sap.

1111 Table S3. Concentrations of polar lipids and sulfolipids in cell contamination controls for xylem
1112 sap.

Supporting experimental procedures:

1113

1114 Methods S1. Methods comparison of lipid extraction by partial freeze drying vs. SpeedVac.

1115