Reproducibility across single-cell RNA-seq protocols for spatial ordering analysis

Morten Seirup[1,2*], Li-Fang Chu[2], Srikumar Sengupta[2], Ning Leng[2,3,#a], Hadley Browder[4], Kevin Kapadia[4], Christina M. Shafer[2], Bret Duffin[2], Angela L. Elwell[2,#b], Jennifer M. Bolin[2], Scott Swanson[2], Ron Stewart[2], Christina Kendziorski[3], James A. Thomson[2,5,6*], Rhonda Bacher[7,*]

[1]Molecular and Environmental Toxicology Program, University of Wisconsin Madison, Madison, Wisconsin, United States of America
[2]Morgridge Institute for Research, Madison, Wisconsin, United States of America
[3]Department of Biostatistics and Medical Informatics, University of Wisconsin Madison, Madison, Wisconsin, United States of America
[4]Department of Statistics, University of Florida, Gainesville, Florida, United States of America
[5]Department of Cell & Regenerative Biology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, United States of America
[6]Department of Molecular, Cellular, & Developmental Biology, University of California Santa Barbara, Santa Barbara, California, United States of America
[7]Department of Biostatistics, University of Florida, Gainesville, Florida, United States of America
[#a]Current Address: Genentech, San Francisco, California, United States of America
[#b]Current Address: Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, United States of America

*Corresponding Authors:
Morten Seirup
E-mail: seirup@wisc.edu
James A. Thomson, V.M.D., Ph.D., Diplomate A.C.V.P.
E-mail: jthomson@morgridgeinstitute.org
Rhonda Bacher, Ph.D.
E-mail: rbacher@ufl.edu

Abstract

As newer single-cell protocols generate increasingly more cells at reduced sequencing depths, the value of a higher read depth may be overlooked. Using data from three different single-cell RNA-seq protocols that lend themselves to having either higher read depth (Smart-seq) or many cells (MARS-seq and 10X), we evaluate their ability to recapitulate biological signals in the context of pseudo-spatial reconstruction. Overall, we find gene expression profiles after spatial-reconstruction analysis are highly reproducible between datasets despite being generated by different protocols and using different computational algorithms. While UMI based protocols such as 10X and MARS-seq allow for capturing more cells, Smart-seq's higher sensitivity and read-depth allows for analysis of lower expressed genes and isoforms. Additionally, we evaluate trade-offs for each protocol by performing subsampling analyses, and find that optimizing the balance between sequencing depth and number of cells within a protocol is important for efficient use of resources. Our analysis emphasizes the importance of selecting a protocol based on the biological questions and features of interest.

Introduction

Single-cell RNA sequencing (scRNA-seq)[1–5] is a powerful tool for studying transcriptional differences between individual cells. The innovation of droplet-based techniques[6,7] and unique molecular identifiers (UMI)[8] has lowered the cost per cell and pushed the field towards obtaining data from tens of thousands of cells per experiment albeit at a reduced sequencing depth. Recent publications have compared the sensitivity, accuracy, and precision between several scRNA-seq techniques and report that the major trade-off between protocols is sensitivity, which is dependent on read depth[9,10]. With the push for sequencing an ever-increasing number of cells at the expense of read depth per cell, the value of a higher read depth might be overlooked. Here we investigate the reproducibility of biological signals across protocols that naturally lend themselves to generating data on more cells versus higher read depth.

Studies comparing protocols have mainly done so with respect to performance

on spike-ins or on technical variability alone[9,10]. Recently, Guo et al.[11] showed

agreement of cell types and signature genes between two platforms used for single-cell

RNA-seq – Fluidigm C1 and Drop-seq. However, few studies have examined

comparative agreement among protocols for biological inferences beyond clustering

and identifying differential gene expression, yet a key question of interest with single-

cell data is its ability to reflect temporal or spatial heterogeneity. For cells collected at a

given time, the underlying dynamic biological process is reflected in genome-wide

differences in gene expression. Computational algorithms that attempt to order cells in

pseudo-time or pseudo-space based on variability in gene expression have been

developed[4,12,13], and more than 45 existing algorithms were recently compared[14]. Yet,

as far as we know, no comparison of single-cell protocols exists for the question of cell

ordering.

Here, our evaluation is in the context of pseudo-spatial reconstruction in which

we compared three independently produced scRNA-seq datasets on the mouse liver

lobule. We chose to compare protocols on their ability to reflect the spatial patterning of

the liver lobule in which the parenchymal cells of the liver, hepatocytes, are organized

spatially in a polygonal shape around a central vein (Figure 1A). From the central vein,

a gradient of metabolic functions is performed extending to a portal triad at each

vertex[15–19]. The gradient of differences in gene expression patterns is referred to as the

zonation axis (from periportal (PP) to pericentral (PC))[20]. This coordinated spatial

organization provides a particularly interesting application of single-cell techniques. For

this study we obtained one dataset using Smart-seq—a full-length protocol, a second

dataset using MARS-seq[21]—a UMI and plate based protocol and the third dataset

generated using 10X[22]—a UMI and droplet protocol. Although the cell number and read

depth differ greatly across datasets, we find high reproducibility of gene expression

profiles after spatial-reconstruction analysis. Given the reproducibility and that each

protocol naturally lends itself to either producing more cells at a lower sequencing depth

or fewer cells at a higher depth, our results demonstrate the importance of carefully

evaluating the biological question and features of interest when selecting the

appropriate sequencing protocol. In applications focused on lower expressed genes or

on genes with high sequence similarity, increased read depth is preferable, whereas a

focus on identifying cell types based on more highly expressed genes will benefit from

collecting more cells. In an ideal situation a single cell assay would result in thousands

of cells that are all sequenced at a high read depth, but technical and financial

restrictions make this rarely possible.

Results

Differences in detection rates

By using the Fluidigm C1 coupled with the Smart-seq protocol, we were able to

identify on average around 38% (about 7100 genes) (Figure 1B) of all genes in the

genome expressed per cell, whereas the MARS-seq dataset finds on average 12%

(about 2200 genes) and the 10X dataset finds on average 6% (about 1100 genes)

(Figure 1B). This is in accordance with findings by Ziegenhain et al. 2017 when they

examined single-cell transcriptomic methods[9] and by Phipson et al[23] when comparing

biases in full-length versus UMI protocols. The increased sensitivity of the full-length

protocol is further illustrated in Figure 1C, which on a per gene level shows the

difference in detection fraction compared to the log fold change in mean expression

between the protocols. A difference in detection fraction of zero means that the gene is

detected in the same fraction of cells in both datasets. The difference across protocols

in log2 fold-change has a linear relationship with the difference in detection fractions,

which indicates a fairly constant increase in log2 expression as cells are sequenced

with greater sensitivity. At the intercept, a difference in detection equal to zero, the log2

fold change is 3.1 between Smart-seq and MARS-seq, indicating an experiment wide

increase in sensitivity in the Smart-seq protocol of approximately 9-fold. Between

Smart-seq and 10X, the increase in sensitivity is approximately 12-fold and there is a

similar level of sensitivity between MARS-seq and 10X. Not surprisingly, the vast

majority of genes are detected in a larger fraction of cells and have a higher expression

level in the more deeply sequenced dataset using the Smart-seq protocol. Although, it is

worth pointing out that around 6% of genes have higher detection using the MARS-seq

protocol (negative values on x-axis) and a few of these genes also have higher

expression levels (negative values on y-axis) than in the Smart-seq protocol. This

subset of genes better detected in the MARS-seq dataset have higher GC content and

are slightly longer (S1 Figure), which is consistent with previous reports of protocol

comparisons[23,24].

Figure 1. Illustration of the liver anatomy, and general comparison of the datasets.

A) Top. Illustration of the liver lobule identifying the portal triad along the outer edges

and the central vein in the middle. The color gradient represents metabolic zonation.

Bottom. Highlights the main differences between the datasets compared. B)

Comparison of gene detection fraction between the datasets. The detection fraction per

cell (y-axis) is shown for the two datasets (x-axis). C) Left. The log2 fold-change of

genes detected above an average expression level of zero in the Smart-seq dataset

compared to the MARS-seq dataset (y-axis), versus the difference in gene-level

detection fractions across datasets (x-axis). A linear regression line is overlaid and a

histogram of the x- and y-axis are shown opposite of each axis. Middle. Similar plot

shown for Smart-seq versus 10X. Right. Similar plot shown for 10X versus MARS-seq.

Reconstructing spatial profiles of liver zonation profiles

Next, to represent the spatial patterns across the liver lobule, the cells in the

three datasets were computationally ordered according to their expression profiles. The

MARS-seq dataset was spatially ordered by Halpern et al. 2017 by first performing

smFISH for six marker genes at various locations across the zonation axis, then single-

cell RNA-seq data obtained by MARS-seq assigned cells to one of nine zonation

locations based on each cell's expression profile of the six marker genes[21]. We ordered

the cells in the 10X dataset using the Monocle2 algorithm, which builds a trajectory

through cells based on the expression similarity among the most highly variable

genes[12]. For the Smart-seq protocol we used the computational algorithm Wave-Crest

to spatially order cells based on fifteen marker genes known in the literature to be

differentially expressed along the zonation axis (Figure 2A)[5]. The ordering procedure

uses the nearest insertion algorithm implemented in the Wave-Crest package, which

searches among the space of all possible orderings via a 2-opt algorithm by considering

insertion events and choosing orders which minimize the mean square error of a

polynomial regression on the marker genes expression. Of the 15 genes used, we

selected eight periportal expressed genes and seven pericentral expressed genes[20]. All

orderings assume the zonation profile and spatial organization can be represented in a

single dimension. A similar reconstructed order was obtained for the Smart-seq dataset

when applying Monocle2 (S2 Figure).

Figure 2. Pseudo-space reordering of hepatocytes, and prediction and validation of dynamically expressed genes. A) Top. Illustration of the pseudo-spatial reordering process. Bottom. Heatmap showing the pseudo-spatial reordering (x-axis) and the expression levels of the marker genes (y-axis) for the Smart-seq dataset. Pericentral cells are found on the left-hand side and Periportal cells are found on the right-hand side. B) Scaled expression profile (y-axis) of 8 dynamic genes based on the predicted pseudo-space reordering (x-axis) of the Smart-seq dataset (orange), the MARS-seq dataset (blue), and the 10X dataset (green). C) Immunohistochemistry staining of the genes highlighted in B). Above the staining is the log2 expression counts (y-axis) across the predicted pseudo-spatial order (x-axis) of the Smart-seq dataset. The left picture

shows the staining and the right picture is an enlarged section (black square). PP =

Periportal, PC = Pericentral.

Using the recreated order of the hepatocytes, we explored the dynamics of gene

expression across the periportal to pericentral axis. Figure 2B shows a subset of genes

that are predicted to be highly regulated across the axis, four of which were not in our

list of marker genes. Since the MARS-seq dataset placed cells into nine discrete zones

along the axis, we divided cells from the Smart-seq and 10X datasets into nine equally

sized groups in order to compare the reconstructed orderings. The zonation profiles in

Figure 2B have high agreement, with a median correlation of 0.95 between the three

datasets. Before proceeding, we also performed an additional experiment to validate

that our cell ordering and expression profiles reflect those of the liver lobule *in vivo*.

Remarkably, immunohistochemistry studies showed that selected marker gene protein

expression profiles also agreed with our spatial reconstructed scRNA-seq datasets: six

markers display a PC-high/PP-low profile and two markers display a PC-low/PP-high

profile in mouse liver lobule *in vivo* (Figure 2C). This confirmation in protein gradient

patterns corresponding to our reconstructed mRNA profiles provides us with confidence

for further analysis on the biological inference in comparing the three protocols in this

context.

Comparing marker gene expression across liver zonation profiles

An exciting prospect of single cell analysis is the identification of genes that have

non-monotonic or dynamic expression across pseudo-time or space. Several genes in

the bile acid synthesis pathway were shown by Halpern et al., 2017 to be non-

monotonically expressed in a pattern where the highest expression levels along the

lobule correspond to the functional placement of the genes in the bile acid synthesis

pathway (Cyp7a1, Hsd3b7, Cyp8b1, Cyp27a1 and Baat)[21]. We find that the expression

profiles for these genes are corroborated across the three datasets (Figure 3).

Figure 3. Comparison of zonation profiles across three datasets. Scaled

expression profile (y-axis) of 8 genes non-monotonically expressed from Halpern et al.

along the predicted pseudo-space reordering (x-axis) of the Smart-seq dataset

(orange), the MARS-seq dataset (blue), and the 10X dataset (green).

However, in the Smart-seq dataset, Cyp8b1 is found to have largely flat

expression levels along most of the lobule and lower expression toward the periportal

zone and Baat appears to have an opposite trend in the 10X dataset. Other genes

shown to be non-monotonically expressed such as Hamp, Igfbp2 and Mup3 in Halpern

et al., 2017 display similar non-monotonic expression profiles in the Smart-seq and 10X

datasets (Figure 3). The ability to identify gene expression profiles that are either high at

the PP end, high at the PC end, or high in the middle of the liver lobule confirms that the

sampling depth is sufficient to spatially reconstruct the liver lobule. We also investigated

the expression pattern of Glul in more detail as it is known to be expressed highly in a

one hepatocyte-wide band around the central vein[25]. Accordingly, the predicted

expression pattern found using all datasets demonstrated sufficient sampling of this

region (S3 Figure).

We further compared the zonation profiles between datasets by identifying genes

having significant differential expression along the reconstructed spatial order across

the periportal to pericentral axis. For genes displaying differential zonation in all

datasets (having adjusted p-value < .1), the Smart-seq versus MARS-seq dataset had

the highest median correlation (0.86), while the Smart-seq versus 10X had the lowest

median correlation (0.69). In Figure 4A we looked at significantly zonated genes within

the metabolic pathways in KEGG and found the median correlation between all datasets

ranged from 0.75 to 0.89. When all genes were considered the median correlation

ranged from 0 – 0.04.

Traditionally the liver lobule is divided into three zones, a periportal zone 1, a

pericentral zone 3, and transitioning zone 2[26,27]. The transitional nature of the liver axis

is reflected in the heatmap of metabolic genes that were significantly zonated in all

datasets (Figure 4B). Using k-means clustering, we found the Smart-seq data tended to

cluster into two distinct gene groups representing either the periportal or pericentral

zone. Examination of the two clusters by enrichment analysis of KEGG metabolic

pathways (Figure 4C) revealed that the predicted location along our reconstructed axis

of metabolic processes with known periportal or pericentral bias such as amino acid

metabolism (periportal), lipogenesis (pericentral), and CYP450 metabolism (pericentral)

corresponds to their known *in vivo* locations[27]. Despite using different reordering

algorithms and protocols, the three datasets show high agreement of expression along

the recovered pericentral to periportal axis among genes that are significantly zonated

in all datasets, and reliably mirror the *in vivo* patterning of the liver lobule (additional

KEGG categories are shown in S4 Figure).

Figure 4. Correlation and Gene Ontology analysis of genes between datasets.

A) Correlation analysis of significantly zonated genes annotated to the metabolic

pathways in KEGG between the datasets. The pairwise correlation is shown for each

dataset comparison. B) Heatmap of the expression level of genes that are significantly

differentially zonated in all datasets and enriched in the metabolic KEGG pathway. C)

Breakdown of KEGG enrichment analysis of the two k-mean clusters based on the

genes shown in B. Dot size represents the fraction of enriched genes in each ontology,

and the color represents the adjusted p-value for the enrichment.

Differences in gene profiles among lowly expressed genes and gene isoforms

When we look at genes with moderate and low expression levels, we find that the

datasets differ to a greater degree. We identified twenty-one genes that were classified

as significantly zonated along the periportal to pericentral axis in the Smart-seq dataset

that were not detected at all in the MARS-seq dataset and thirty-five such genes not

detected in the 10X dataset. Compared to the Smart-seq dataset, ten genes were

exclusively detected in the MARS-seq dataset and no genes were exclusive to the 10X

dataset. Figure 5A shows the six most highly expressed genes that we were able to

exclusively identify in the Smart-seq dataset having significant zonation (adjusted p-

value < .1).

Figure 5. Genes and isoforms found in the full-length dataset and not in the UMI

datasets. A) Six genes found to be zonally expressed in the Smart-seq dataset that

were not detected in either the MARS-seq or 10X datasets. The log2 of expression

values are represented on the y-axis and the pseudo-space ordered cells are found on

the x-axis. B) Examples of genes with two transcript variants expressed differently

across reordered cells from the Smart-seq dataset.

Further, an exciting field of study that benefits from an enhanced resolution of

scRNA-seq is isoform analysis[28–30]. Many genes in the genome have two or more

isoforms that are distinctly expressed and can change properties such as structure,

function, and localization of the resulting protein[31]. Due to the increased sensitivity of

full-length cDNA libraries generated by Smart-seq protocol, we were able to examine

genes with known isoforms and identify cases where the transcript variants for each

isoform has distinct expression from each other across the periportal to pericentral axis,

which is not possible with less sensitive protocols. In Figure 5B the transcript variants of

Romo1 are seen to display opposite trends in expression across the zonation axis,

where the Romo1 variant 3 is increasing in expression from the pericentral end towards

the periportal end and the Romo1 variant 1 is decreasing in expression along the same

axis. We also highlight genes Acox1 and Eif4a2 whose variants both show constant

expression across the zonation axis but at different levels. Both of these genes are

known to have isoform-specific expression in the liver lobule[32,33]. (For Ensembl and

ENTEREZ IDs for transcript variants see S6 Table).

Due to UMI based protocols capturing only one end of the transcript compared to

full-length cDNA procedures, there is an inability to resolve not just isoforms but also

many genes that are closely related. For instance, there were 242 concatenated genes

in the MARS-seq set that correspond to 539 unique genes. An example of this is seen

in S5 Figure where we highlight a concatenate of Ugt1a enzymes. Eight genes are

concatenated (annotated together) and when combined, the average expression level is

shown to be high at the pericentral end of the lobule and low at the periportal end.

Again, it is clear that not all the members of this concatenated group follow this trend as

Ugt1a6a can be seen to have consistent expression levels across the pericentral to

periportal axis.

Evaluating the trade-off within each protocol *in silico*

To further study the trade-offs between higher depth versus more cells, we

performed a subsampling experiment. For each dataset, we held either the number of

cells or the sequencing depth constant while varying the other. For the Smart-seq and

10X datasets, we evaluated the effect on the cell ordering as well as the gene-specific

zonation profiles. For the MARS-seq dataset, the assignment of each cell to a zonation

group depended on external data and was independent of the other cells profiled, and

thus we only evaluated the effect on zonation profiles. We estimate the MSE (mean

squared error) as the difference in zonation profiles in the subsampled dataset versus

the original dataset. In Figure 6A, the MARS-seq dataset displayed an approximately

linear tradeoff in zonation profile error for fewer cells at the original read depth.

However, at reduced read depths using the original 1,415 cells, the error increased

exponentially (Fig.6B). Within a dataset, we can compare the MSE between the two

trade-off scenarios and we find that for the MARS-seq dataset resequencing at the

same depth results in error that is equivalent to the reduction observed in MSE by going

from 600 to 1400 total cells. For the 10X dataset, we also find an approximately linear

tradeoff in zonation profile error for fewer cells at the original read depth (Fig. 6C).

However, at reduced read depths using the original 606 cells, we observe a gradual

increase in error as total depth decreases (Fig.6D). Similarly, by comparing the MSE

trade-off, it appears that resequencing at the same depth results in error that is

equivalent to reducing the total cells from 600 to around 400. Thus, in scenarios with

very low sequencing depth (average of 3-12k total UMIs per cell), sequencing deeper

may be more beneficial than adding more cells. For the Smart-seq dataset, we found

the spatial ordering to be quite robust to reduced sequencing depth, even as low as

50% fewer reads only marginal increased the average MSE as shown in Figure 6F. The

average sequencing depth for the Smart-seq cells was 3.5 million counts per cell, well

beyond the suggested sequencing saturation for single-cell data that occurs close to

one million total reads[34]. We do see more dramatic increases in error related to zonation

profiles when profiling fewer cells (Figure 6E). For Smart-seq data, sequencing to even

half of the current depth and increasing the number of cells would be beneficial.

Figure 6. Subsampling total numbers of cells and sequencing depth. A) For 25

subsamplings at various total numbers of cells in the MARS-seq dataset, the mean

squared error (MSE) of the zonation profile over 500 randomly selected genes is shown.

B) Similar to A, but for 25 subsamplings at various total read depths. C-D) Similar to A-

B, but for the Smart-seq dataset. E-F) Similar to A-B, but for the 10X dataset.

Discussion

In summary, we compared three scRNA-seq datasets of mouse hepatocytes

where two, MARS-seq and 10X, are wide but shallow and the other, Smart-seq, is

narrow but deeply sequenced. We find that the three different protocols present highly

reproducible liver zonation profiles in single cells, and for the vast majority of genes that

are highly expressed we observe highly comparable results. Our results were not

dependent on any one computational method or pre-processing pipeline. We do

however find that when we look at medium to low expressed genes, the increased

sensitivity of the C1/Smart-seq protocol is able to identify several genes exclusive to this

dataset. This increased sensitivity also allowed us to identify several genes with

isoforms that behaved differently across the periportal to pericentral axis. Though in

general, there are still limitations of short reads in regard to isoform analysis and if more

accuracy is needed, the newly developed technique ScISOr-seq[35] might be better

suited. We do however believe that this full-length data allows for more reliable

preliminary isoform analysis compared to either UMI method. However, the main

weakness of using fewer cells is that it is less likely that rare cell types will be sampled.

In cases where such rare cells are of high interest, protocols that produce a large

number of cells are preferable. In an ideal case, one would sample many cells and

sequence all of them deeply; unfortunately, this is not always possible in practice and

the decision of whether to sample many cells shallowly or fewer cells deeply comes

down to whether rare cell types are of interest or if higher resolution of the individual

cells is preferred.

Given the distinct advantages of the protocols, we emphasize that the biological

question should be the driving factor when deciding on protocol. Within a chosen

protocol, achieving balance between the sequencing depth and the number of cells is

still an important consideration for optimal use of resources. Based on our simulations

of datasets at opposite ends of the sequencing depth versus number of cells trade-off,

there is eventually a detriment to sacrificing reads for additional cells or sequencing

beyond the attainable sensitivity level on too few cells. We expect that the extent of the

cells versus depth trade-off will vary for other cell types or tissues and it will largely

depend on the heterogeneity of the biological system under study.

Methods

Animals and handling.

All animals were kept under standard husbandry conditions. A wildtype 8-week-old male

C57BL/6 (Jackson laboratories) was used in this experiment. Using isoflurane, the

mouse was anesthetized before euthanizing by cervical dislocation. Animal experiments

and procedures were approved by the University of Wisconsin Medical School's Animal

Care and Use Committee and conducted in accordance with the Animal Welfare Act

and Health Research Extension Act.

Cell isolation.

The euthanized mouse was pinned to a Styrofoam plate using 20 ga needles to aid in

dissection. The abdominal cavity was opened, and the portal vein exposed. A piece of

4-0 suture thread (Ethicon vicryl coated) was threaded under the portal vein and used to

secure a 26 ga catheter inserted into the portal vein (Butler Schein animal health 26 G

IV Catheter, Fisher Scientific). Hepatocytes were isolated using a 2-step perfusion

protocol. First, Liver Perfusion Medium (Gibco) warmed to 37°C was pumped through

the catheter for 10 minutes using a peristaltic pump at 7 ml/min flowrate. Then, Liver

Digest Medium (Gibco) warmed to 37°C was pumped through the liver at the same

settings for 10 minutes. After perfusion, the liver was excised and transferred to a 10 cm

dish containing 20 ml liver digest medium. The liver was dissected, allowing the cells to

spill into the media. The cells were then filtered through a 40 μm cell strainer into a 50

ml tube and 30 ml media (Williams E media + 2 μg/ml human insulin + 1x glutamax +

10% FBS) were added and placed on ice. The hepatocytes were purified by

centrifugation at 50 x G, 4 times for 3 minutes each, each time discarding the

supernatant and adding media.

Single cell RNA sequencing- Full-length dataset

Single-cell RNA sequencing was performed as previously described[4,5] with the following

modifications. In this study, we used small (5-10 μm), medium (10-17 μm), and large

(17-25 µm) plate sizes. ERCC RNA Spike-In (ThermoFisher Cat. No. 4456740) was

diluted in the lysis mix following the manufacturer's user guide and previous studies[36].

Single end reads of 51 bp were sequenced on an Illumina HiSeq 2500 system.

Sequencer outputs were processed using Illumina's CASAVA-1.8.2. The demultiplexed

reads were trimmed and filtered to eliminate adapter sequence and low-quality

basecalls. The reads were mapped to an mm10 mRNA transcript reference (extended

with ERCC transcripts) using bowtie-0.12.9[37]; expression estimates were generated

using RSEM v.1.2.3[38]. Using the Fluidigm C1 system to capture and synthesize cDNA

from single cells in the liver, we generated transcriptomes for 149 cells. To exclude low

quality transcriptomes, we removed cells in which the fraction of ERCC spike-in made

up 20% or more of the total assigned reads. This left 66 high quality cells that were

used in the downstream analysis. Finally, the data was normalized using SCnorm (R

package v 1.5.7)[39].

Pseudo-spatial reordering- Full-length dataset

For the full-length data, the cells were computationally ordered using the Wave-Crest

method as described in Chu et al. 2016[5]. For the reordering step, gene expression

values were rescaled to mean 0 and variance 1 to ensure the values across different

genes are comparable. The Wave-Crest algorithm implements an extended nearest

insertion algorithm that iteratively adds cells to the order and selects the insertion

location as the location producing the smallest mean squared error in a linear

regression of the proposed order versus gene expression. A 2-opt algorithm is then

used to find an optimal cell order by considering adjacent cell exchanges. The cell

ordering step uses the expression profiles of pre-selected known marker genes of liver

zonation. Thus, the resulting linear profile of ordered cells represents the periportal to

pericentral axis. The known marker genes used to construct the periportal to pericentral

axis in Wave-Crest include the following pericentral markers: cytochrome P450 7a1

(Cyp7a1), cytochrome P450 2e1 (Cyp2e1), ornithine aminotransferase (Oat),

cytochrome P450 1a2 (Cyp1a2), rh family, B glycoprotein (Rhbg), leucine-rich repeat-

containing G-protein coupled receptor 5 (Lgr5), glutamate-ammonia ligase (Glul); and

the following periportal markers: phosphoenolpyruvate carboxykinase 1 (Pck1), catenin

beta interacting protein 1 (Ctnnbip1), aldehyde dehydrogenase 1 family member B1

(Aldh1b1), sulfotransferase family 5A, member 1 (Sult5a1), cytochrome P450 2f2

(Cyp2f2), cathepsin C (Ctsc), serine dehydratase (Sds), and E-cadherin (Cdh1). All

markers were selected based on their expression ratio as reported by Braeuning et al.

2006[20].

A detection step was done to identify additional genes that follow the one-dimensional

periportal to pericentral axis by fitting a linear regression to the relationship between

each gene's expression and the Wave-Crest cell order. To determine if a gene is

significantly dynamic (differentially zonated) along the recovered axis, we tested

whether the regression slope is different from zero. We reported the Benjamini-

Hochberg adjusted p-values to control the false discovery rate. For genes having an

adjusted p-value < .01, the direction of the expression profile was assigned based on

the sign of the regression slope (periportal: positive slope, pericentral: negative slope).

We also calculated the linear fitting mean squared error (MSE) for each gene. Genes

with a smoother trend over the recovered cell order are expected to have a smaller

MSE. We report the full list of genes, sorted by their MSE, in S7 Table; scatter plots for

genes having adjusted p-value < .01 are shown in S8 File.

Pseudo-spatial reordering- 10X dataset

The 10X dataset was downloaded from the Tabula Muris compendium public resource

via Figshare[22] . The 10X data was originally processed using the CellRanger version

2.0.1. Within the liver cells, the authors originally identified 975 hepatocytes. For our

analysis, we performed a second quality control step to identify cells with low RNA

content, possible doublets, or dead/damaged cells, where we filtered cells based on the

total number of genes expressed per cell. Using the Seurat R package v3.1.5,

hepatocytes were further filtered to those having between 200 and 3000 genes detected

per cell (only one cell had more than 5000 genes detected per cell). Next, we clustered

the cells using Seurat, where a k-nearest neighbors (KNN) graph used was constructed

based on the first 20 principle components to create a shared nearest neighbors graph

based on the Jaccard index between each cell and its 20 nearest neighbors, as

implemented in the FindNeighbors function. Clusters were then identified by partitioning

this graph using the Louvain community detection algorithm with a resolution of .5, as

implemented in the FindClusters function. The cells clustered into three distinct larger

groups and we retained only the largest grouping of cells that clustered together,

resulting in 606 total cells. The data was then normalized using scran v1.12.1. Next, we

used Monocle v2.12.0 to order the cells, basing the ordering on the top 200 highly

variable genes estimated using the mean variance relationship via the

FindVariableFeatures function in Seurat. To determine if a gene is significantly dynamic

(differentially zonated) along the recovered axis, the Monocle2 function

differentialGeneTest was used to fit a spline on gene expression versus the estimated

pseudo-time.

Comparative Analysis

Smoothed densities (bean plots) with overlaid raw data, the mean, and a box

representing the interquartile range of the cellular detection fractions were created using

the pirateplot function in the yarrr R package (v0.1.5). The cellular detection fraction

was calculated per cell as the proportion of genes having expression greater than zero.

The fold-change for each gene between the two datasets (A versus B) was calculated

as the log2 fold-change of the dataset A over dataset B, where each gene mean was

calculated as the average expression among non-zero counts across all cells in the

datasets. The heatmap in Figure 2 of marker gene expression on the normalized Smart-

seq data was generated by setting values above the 95th percentile or below the 5nd

percentile to the 95th percentile or 5nd percentile value, respectively.

Due to the datasets having different dynamic ranges, we used scaled expression plots

to compare expression profiles, where the ordered cells in the full-length dataset and

10X were each divided into nine equally sized groups to correspond to the nine layers in

the UMI dataset. For the full-length and 10X dataset, for a given gene, the median (full-

length) or mean (10X) expression in each group was calculated, then the nine values

were scaled between zero and one. Smoothed fits were overlaid using the

smooth.spline function in R with the degrees of freedom parameter df=4. Expression

correlations along the zonation axis between datasets were calculated using Pearson

correlation. Enrichment of genes in KEGG pathways or GO was done using the R

package clusterProfiler (v. 3.10.1)[40]. For the enrichment analysis, since different

statistical methods were used to assess zonation profiles, genes were considered

significantly zonated if they had an adjusted p-value < .1 in all datasets. The heatmap in

Figure 3 is a smoothed heatmap, where a smoothing spline was first fit to the log

expression (pseudo-count of one added) of each gene using the smooth.spline function

in R with the smoothing parameter df=4 which provided profiles that were not over- or

underfit in either dataset. Then the smoothed expression was scaled and outliers above

the 98th percentile or below the 2nd percentile were set to the 98th percentile or 2nd

percentile value, respectively. Additional KEGG categories from this analysis can be

interactively viewed on Github

https://github.com/rhondabacher/scSpatialReconstructCompare-Paper.

Subsampling Analysis

In all subsamplings described below, each scenario was repeated a total of 25 times

and the zonation group means were scaled to be between zero and one.

For the MARS-seq dataset, zonation group means were recalculated on a subsampled

set of cells using the posterior probability matrix and original UMI counts from Halpern

et al. 2017. In each sampling, the mean squared error (MSE) was calculated based on a

random sample of 500 genes as $\sum_{i=1}^{500} \sum_{j=1}^{9} (Z_{i,j} - \hat{Z}_{i,j})^2 / 500$, where $Z_{i,j}$ represents the

mean expression of gene i in zonation group j in the original dataset and $\hat{Z}_{i,j}$ is the

corresponding value for the subsampled dataset. For subsampling at lower read depths,

we fixed the number of cells at the original total of 1415 cells and simulated each cell's

gene counts individually using a multinomial distribution. For each cell, the subsampled

total counts were set to X% of the original total read counts for that cell (for X =

(10,20,30,40,50,60,70,80,90,100)) and each gene's cell-specific probability was

calculated as its original count divided by the original total counts for that cell. The MSE

was calculated for each subsampled set as described above.

For the Smart-seq dataset, we reran Wave-Crest when subsampling the total number of

cells using the original parameter settings and marker genes. Then, as before, the

ordered cells were assigned zonation groups by dividing cells into nine equally sized

groups. The zonation profile error was estimated using MSE and calculated as

described above with the exception that since Wave-Crest orders can be flipped, we

calculated the MSE on the returned order and its reverse, and kept the minimum MSE

of the two. To evaluate the zonation profile error with lower read depths, we used a

similar approach as described above for the MARS-seq dataset, fixing the number of

cells to be the same as the original total of 66 and, since the order correlation was

shown to be consistently high, we used the original Wave-Crest order for every scenario

when evaluating zonation profile error. For the 10X dataset, the subsampling was

performed similarly as for the Smart-seq dataset, however Monocle2's ordering was

more variable as it was not based on marker genes and thus we did not fix the order

when evaluating the zonation profile error. Trade-offs in MSE are directly comparable

within a dataset but due to intrinsic differences in the original processing and in

subsampling, the MSE should not be compared across the datasets.

Immunohistochemistry

An 8-week-old male C57BL/6 mouse was anesthetized using isoflurane before

euthanizing by cervical dislocation. The liver was excised, sliced as thinly as possible

with a razor blade, and fixed in formaldehyde overnight. The liver slices were paraffin

embedded and sectioned. Sections were stained following the protocol published by

Abcam (http://www.abcam.com/ps/pdf/protocols/ihc_p.pdf). In short, the slices are

deparaffinized by dipping into sequential solutions of 100% xylene, 50-50% xylene-

ethanol, 100% ethanol, 95% ethanol, 70% ethanol, 50% ethanol, and tap water. The

antigens were then retrieved by placing the slides in Tris-EDTA buffer (10 mM Tris

Base, 1 mM EDTA Solution, 0.05% Tween 20, pH 9.0) and incubating them in a

decloaking chamber (Biocare Medical Decloaking Chamber #DC2008US) with the

following settings: delayed start 30 sec.; preheat 80°C, 2 min.; heat 101°C, 3 min. 30

sec.; and fan on. The slides were washed 2 x 5 min in TBS + 0.025% Triton X-100

before they were blocked for two hours at room temperature in 10% normal serum in

1% BSA. The appropriate primary antibody was then diluted in the same 10% normal

serum in 1% BSA, added to the slides, and incubated at 4ºC overnight in an incubation

chamber. The next day the slides were washed 2 x 5 min in TBS + 0.025% Triton X-100

followed by 15 min incubation in 0.3% H_2O_2 at room temperature. Next, the appropriate

secondary antibody was diluted into 10% normal serum in 1% BSA before it was added

to the slides and incubated for 1 hour at room temperature. The slides were then

washed 3 x 5 min in TBS before DAB (#ab103723) staining mixed according to

manufacturer instruction was applied and incubated under a microscope to stop the

reaction after sufficient staining. The slides were rinsed in tap water for 5 min before

being counterstained with Mayer's hematoxylin (#MHS1-100ML) for 30 sec. The stain

was developed in running tap water for 5 min. The slides were then dehydrated by

sequentially dipping in 50% ethanol, 70% ethanol, 95% ethanol, 100% ethanol, 50-50%

xylene-ethanol, and 100% xylene before Poly-Mount (#08381-120) was added and a

coverslip placed on top. The following primary antibodies were added: Aldh3a4 1:250

(AB184171), Cyp2e1 1:50 (AB28146), Cyp1a2 1:50 (R31007), Rgn 1:100 (NBP1-

80849), Oat 1:50 (AB137679), Cyp2f2 1:100 (SC-67283), Hal 1:50 (AV45694), and

Tbx3 1:50 (SC-31657). The following secondary antibodies were used: goat-anti-rabbit

HRP conjugated (ab97051) and donkey-anti-goat HRP conjugated (ab97110) at a

concentration of 1:500.

Acknowledgements

References
580

1. Tang, F. *et al.* mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods* **6**,

377–382 (2009).

2. Ramsköld, D. *et al.* Full-length mRNA-Seq from single-cell levels of RNA and individual

circulating tumor cells. *Nat Biotechnol* **30**, 777–782 (2012).

3. Islam, S. *et al.* Characterization of the single-cell transcriptional landscape by highly

multiplex RNA-seq. *Genome Research* **21**, 1160–1167 (2011).

4. Leng, N. *et al.* Oscope identifies oscillatory genes in unsynchronized single-cell RNA-seq

experiments. *Nat Methods* **12**, 947–950 (2015).

5. Chu, L.-F. *et al.* Single-cell RNA-seq reveals novel regulators of human embryonic stem cell

differentiation to definitive endoderm. *Genome Biol* **17**, 173 (2016).

6. Klein, A. M. *et al.* Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic

Stem Cells. *Cell* **161**, 1187–1201 (2015).

7. Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells

Using Nanoliter Droplets. *Cell* **161**, 1202–1214 (2015).

8. Islam, S. *et al.* Quantitative single-cell RNA-seq with unique molecular identifiers. *Nat*

Methods **11**, 163–166 (2014).

9. Ziegenhain, C. *et al.* Comparative Analysis of Single-Cell RNA Sequencing Methods.

Molecular Cell **65**, 631-643.e4 (2017).

10. Svensson, V. *et al.* Power analysis of single-cell RNA-sequencing experiments. *Nat Methods*

14, 381–387 (2017).

11. Guo, M. *et al.* Single cell RNA analysis identifies cellular heterogeneity and adaptive

responses of the lung at birth. *Nat Commun* **10**, 37 (2019).

12. Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381–386 (2014).

13. Ji, Z. & Ji, H. TSCAN: Pseudo-time reconstruction and evaluation in single-cell RNA-seq

analysis. *Nucleic Acids Res* **44**, e117–e117 (2016).

14. Saelens, W., Cannoodt, R., Todorov, H. & Saeys, Y. A comparison of single-cell trajectory

inference methods. *Nat Biotechnol* **37**, 547–554 (2019).

15. Burger, H.-J., Gebhardt, R., Mayer, C. & Mecke, D. Different capacities for amino acid

transport in periportal and perivenous hepatocytes isolated by digitonin/collagenase

perfusion. *Hepatology* **9**, 22–28 (1989).

16. Pösö, A. R., Penttilä, K. E., Suolinna, E. M. & Lindros, K. O. Urea synthesis in freshly

isolated and in cultured periportal and perivenous hepatocytes. *Biochem. J.* **239**, 263–267

(1986).

17. Tosh, D., Alberti, G. M. M. & Agius, L. Glucagon regulation of gluconeogenesis and

ketogenesis in periportal and perivenous rat hepatocytes. Heterogeneity of hormone action

and of the mitochondrial redox state. *Biochem. J.* **256**, 197–204 (1988).

18. Guzmán, M. & Castro, J. Zonation of fatty acid metabolism in rat liver. *Biochem. J.* **264**,

107–113 (1989).

19. Anundi, I., Lähteenmäki, T., Rundgren, M., Moldeus, P. & Lindros, K. O. Zonation of

acetaminophen metabolism and cytochrome P450 2E1-mediated toxicity studied in isolated

periportal and perivenous hepatocytes. *Biochemical Pharmacology* **45**, 1251–1259 (1993).

20. Braeuning, A. *et al.* Differential gene expression in periportal and perivenous mouse

hepatocytes. *FEBS Journal* **273**, 5051–5061 (2006).

21. Halpern, K. B. *et al.* Single-cell spatial reconstruction reveals global division of labour in the

mammalian liver. *Nature* **542**, 352–356 (2017).

22. The Tabula Muris Consortium *et al.* Single-cell transcriptomics of 20 mouse organs creates a

Tabula Muris. *Nature* **562**, 367–372 (2018).

23. Phipson, B., Zappia, L. & Oshlack, A. Gene length and detection bias in single cell RNA

sequencing protocols. *F1000Res* **6**, 595 (2017).

24. Parekh, S., Ziegenhain, C., Vieth, B., Enard, W. & Hellmann, I. The impact of amplification

on differential expression analyses by RNA-seq. *Sci Rep* **6**, 25533 (2016).

25. Gebhardt, R. & Mecke, D. Heterogeneous distribution of glutamine synthetase among rat

liver parenchymal cells in situ and in primary culture. *The EMBO Journal* **2**, 567–570

(1983).

26. Bhatia, S. N. *et al.* Zonal liver cell heterogeneity: effects of oxygen on metabolic functions of

hepatocytes. *Cell Eng* **1**, 125–135 (1996).

27. Kietzmann, T. Metabolic zonation of the liver: The oxygen gradient revisited. *Redox Biology*

11, 622–630 (2017).

28. Song, Y. *et al.* Single-Cell Alternative Splicing Analysis with Expedition Reveals Splicing

Dynamics during Neuron Differentiation. *Molecular Cell* **67**, 148-161.e5 (2017).

29. Karlsson, K., Lönnerberg, P. & Linnarsson, S. Alternative TSSs are co-regulated in single

cells in the mouse brain. *Mol Syst Biol* **13**, 930 (2017).

30. Arzalluz-Luque, Á. & Conesa, A. Single-cell RNAseq for the study of isoforms—how is that

possible? *Genome Biol* **19**, 110 (2018).

31. Wang, E. T. *et al.* Alternative isoform regulation in human tissue transcriptomes. *Nature*

456, 470–476 (2008).

32. Gruppuso, P. A., Tsai, S.-W., Boylan, J. M. & Sanders, J. A. Hepatic translation control in

the late-gestation fetal rat. *American Journal of Physiology-Regulatory, Integrative and*

Comparative Physiology **295**, R558–R567 (2008).

33. Oaxaca-Castillo, D. *et al.* Biochemical characterization of two functional human liver acyl-

CoA oxidase isoforms 1a and 1b encoded by a single gene. *Biochemical and Biophysical*

Research Communications **360**, 314–319 (2007).

34. Haque, A., Engel, J., Teichmann, S. A. & Lönnberg, T. A practical guide to single-cell RNA-

sequencing for biomedical research and clinical applications. *Genome Med* **9**, 75 (2017).

35. Gupta, I. *et al.* Single-cell isoform RNA sequencing characterizes isoforms in thousands of

cerebellar cells. *Nat Biotechnol* **36**, 1197–1202 (2018).

36. Brennecke, P. *et al.* Accounting for technical noise in single-cell RNA-seq experiments. *Nat*

Methods **10**, 1093–1095 (2013).

37. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

38. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

39. Bacher, R. *et al.* SCnorm: robust normalization of single-cell RNA-seq data. *Nat Methods*

14, 584–586 (2017).

40. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology* **16**,

284–287 (2012).

Supporting information captions

S1 Figure – Examining GC content and gene length in genes with a higher detection

fraction in either dataset. Top) The GC content (left) and gene length (right) are shown

for genes having a higher detection fraction in either the Smart-seq dataset (gray) or the

MARS-seq dataset (blue). A dotted line is shown for genes having a larger mean in

either dataset. The two lines closely correspond since the genes having a high detection

fraction typically have a higher mean. Bottom) Similar to the top for comparing the

Smart-seq and 10X datasets.

S2 Figure – Correlation between WaveCrest and Monocle methods for ordering cells in

the Smart-seq dataset.

S3 Figure – Expression of Glul. Scaled expression plots of Glul showing high correlation

among all three datasets.

S4 Figure – Correlation analysis of more KEGG pathways. A) Top left: Correlation

analysis for genes in the KEGG pathway "Complement and coagulation cascade". The

pairwise correlation is shown for each dataset comparison. Following are plots for the

eight highest correlated genes between the any two datasets in that pathway. On the

right is a smoothed heatmap of the Smart-seq expression data for the gene expression

of all significantly zonated genes enriched in that KEGG pathway. B) Similar to (A) but

for the "Drug metabolism – cytochrome P450" pathway. C) Similar to (A) but for the

"Biosynthesis of amino acids" pathway.

S5 Figure – Additional genes in Smart-seq dataset but not in the MARS-seq dataset.

Eight Ugt1a genes that were concatenated in the MARS-seq dataset (blue on all

graphs), but can be resolved in the Smart-seq dataset (orange line).

S6 Table – Ensembl and RefSeq ID's for genes with transcript variants.

S7 Table – Summary of genes with dynamic expression across the zonation axis

identified using Wave-Crest.

700 S8 File – Scatter plots of dynamic genes listed in S6 Table.

701 S9 Dataset – Normalized Smart-Seq single-cell data with cells in the Wave-Crest order.