Genome-wide changes in genetic diversity in a population of

Myotis lucifugus affected by white-nose syndrome

Lilley TM [1,2], Wilson IW [1], Field KA [3], Reeder DA [3], Vodzak ME[3], Turner G [4], Kurta A [5],

Blomberg AS [6], Hoff S [7], Herzog C [7], Paterson S [1].

[1]Institute of Integrative Biology, University of Liverpool, United Kingdom. [2]Finnish Museum

of Natural History, University of Helsinki, Finland. [3]Biology Department, Bucknell

University, 1 Dent Drive, Lewisburg, PA 12837, USA. [4]Pennsylvania Game Commission,

Harrisburg, PA, USA. [5]Department of Biology, Eastern Michigan University, Ypsilanti, MI,

USA. [6]Department of Biology, University of Turku, Finland. [7]Wildlife Diversity Unit, State

Department of Environmental Conservation, Albany, NY, USA.

Corresponding author: tmlill@liverpool.ac.uk

Abstract

Novel pathogens can cause massive declines in populations, but seldom lead to extirpation

of hosts. Rather, disease can act as a selective pressure on survivors, driving the evolution of

resistance or tolerance. Bat white-nose syndrome (WNS) is a rapidly spreading wildlife

disease in North America. The fungus causing the disease invades skin tissues of hibernating

bats, resulting in disruption of hibernation behaviour, premature energy depletion, and

subsequent death. We use whole-genome sequencing to investigate changes in allele

frequencies within a population of *Myotis lucifugus* in eastern North America to scan for

genetic resistance to WNS. Our results show low F_{ST} values within the population across

time, i.e. prior to WNS (Pre-WNS) compared to the population that has survived WNS (Post-

WNS). However, when dividing the population with a geographical, distance-based cut-off

between the states of Pennsylvania and New York, a sharp increase in values on scaffold

GL429776 is evident in the Post-WNS samples. Genes present in the diverged area are

associated with thermoregulation and promotion of brown fat production. Thus, although

WNS may not have subjected the entire *M. lucifugus* population to selective pressure, it could

allowed the rise of a selection for alleles in Pennsylvania through decreased gene flow within

the population, which may contribute to survival. However, the existence of remnant sub-

populations in the aftermath of WNS is likely due to multiple factors in bat life history.

Keywords: Opportunistic pathogen, white-nose syndrome, Selective pressure, Genetic

resistance.

1. Introduction

The emergence and spread of multiple infectious wildlife diseases during recent decades has had devastating consequences for biodiversity (Daszak et al., 2000; Smith et al., 2006). Unfortunately, for many taxa, anthropogenic activities and climate change appear to be the main causes of exposure to sources of infection (Gallana et al., 2013; Tompkins et al., 2015). With shifting pathogen ranges, disease-related declines in naïve wildlife often threaten the persistence of populations. Examples include chytridiomycosis, which has wiped out populations of amphibians globally (Daszak et al., 1999; Lips et al., 2006), and avian malaria, which caused the sharp decline of island populations of birds (van Riper et al., 1986). More recently, white-nose syndrome (WNS) has been described as one of the most rapidly spreading wildlife diseases ever recorded (Blehert et al., 2009; Frick et al., 2010). Since the discovery of WNS in North America in 2006, ten species of bats have been diagnosed with the disease in 33 U.S. states and 7 Canadian provinces (www.whitenosesyndrome.org, 2019). Genetic evidence suggests that *Pseudogymnoascus destructans*, the causative agent of WNS, was introduced to North America from Europe (Campana et al., 2017; Lorch et al., 2013; Minnis and Lindner, 2013; Ren et al., 2012; Wibbelt et al., 2010), where affected species do not experience associated mortality (Harazim et al., 2018; Puechmaille et al., 2011; Warnecke et al., 2012; Wibbelt et al., 2013; Zukal et al., 2016).

When a disease enters a naïve host population, the initial wave of infection often causes epizootics that result in mass mortality, which may extirpate local host populations or even cause species extinction (Daszak et al., 1999; De Castro and Bolker, 2005; Frick et al., 2010). However, extinctions due to a disease appear to occur seldomly, and a disease can act as a

selective pressure on survivors, driving the evolution of tolerance or resistance, which may

transform a disease from being epizootic to being enzootic (Boots et al., 2009; Karlsson et al.,

2014; Robinson et al., 2012). This may occur through rapid changes in the distribution of

genetic variants associated with disease susceptibility over short timescales (Gallana et al.,

2013). These powerful selective pressures pathogens exert on naïve host populations (Sironi

et al., 2015) and genetic signatures of infection-related selective sweeps may be detectable

for generations (Deschamps et al., 2016; Di Gaspero et al., 2012; Groot et al., 2002).

Detection of selective sweeps on particular genes or gene families can confirm or exclude

potential mechanisms of host susceptibility or pathogen virulence (Campbell and Tishkoff,

2008). Rapid, extreme bottlenecks (such as those caused by panzootic events), meanwhile,

are associated with a more stochastic loss of alleles, which does not necessarily indicate

selection (Luikart et al., 1998). Much of the research on disease-induced selection has

focused on the major histocompatibility complex (MHC), and indeed, diseases can drive the

evolution and maintenance of MHC diversity in natural populations (Davy et al., 2017a;

Jeffery and Bangham, 2000; Paterson et al., 1998; Spurgin and Richardson, 2010; Teacher

Amber G. F. et al., 2009; Zeisset and Beebee, 2014). However, factors not directly associated

with interactions between host and pathogen, such as environmental conditions and

competition with other species, can have a considerable influence on the manifestation of a

disease (Scholthof, 2007).

The psychrophilic (cold-growing) fungus *P. destructans*, (Lorch et al., 2011; Minnis and

Lindner, 2013; Warnecke et al., 2008) acts as an opportunistic pathogen of bats, invading the

skin tissues of hibernating hosts (Cryan et al., 2010; Meteyer et al., 2012). Susceptible

species, such as *Myotis lucifugus* and *M. septentrionalis*, have shown population declines

greater than 90% in affected hibernacula (Frick et al., 2015). The infection disrupts the

hibernation behaviour of susceptible species and leads to more frequent arousals from torpor,

evaporative water loss, premature energy depletion, and death of susceptible individuals due

to emaciation (McGuire et al., 2017; Reeder et al., 2012; Verant et al., 2014; Warnecke et al.,

2013; Willis et al., 2011). Naïve infected *M. lucifugus* upregulate genes involved in immune

pathways during the hibernation period (Davy et al., 2017b; Field et al., 2015, 2018; Lilley et

al., 2017). These responses are weak during torpor but are robust during the intermittent

arousals (Field et al., 2018; Luis and Hudson, 2006). Therefore, increased arousals may be

attempts by the host to counter the pathogen, in addition to quenching thirst, grooming,

expelling waste and possibly foraging (Bernard and McCracken, 2017; Willis et al., 2011),

and supplementing electrolytes (Cryan et al., 2013).

White-nose syndrome is a prime example of a disease which is manifested when factors

associated with the environment (i.e. temperature and humidity of the hibernaculum), the host

(and the host's response to infection) and the pathogen (optimum growth range, suitable host)

overlap, i.e. intersect within the "disease triangle" (Scholthof, 2007). Therefore bat species,

and populations within species, are affected differently according to hibernation behaviour

and prevailing micro-climate conditions in available hibernacula (Grieneisen et al., 2015;

Johnson et al., 2014; Langwig et al., 2015). As such, with an environmentally persistent

opportunistic pathogen such as *P. destuctans* capable of subsistence in the absence of the

host, loci not associated with MHC diversity or other immune response-associated factors

(Donaldson et al., 2017), such as the amount of body fat prior to hibernation (Cheng et al.,

2019), may also affect survival of species and populations. This highlights the importance of

examining the genome as a whole (Sparks et al., 2019).

The initial impacts of WNS on six species of hibernating bats in the northeastern and

midwestern USA vary from almost complete extirpation to arrested population growth

(Turner et al., 2011). After the initial decline caused by WNS in *M. lucifugus*, reports have

surfaced describing stabilisation at smaller sizes or even increases in numbers of individuals

(Dobony and Johnson, 2018; Frick et al., 2017; Langwig et al., 2015; Maslo et al., 2015).

Models parameterised with long-term data on fungal loads, infection intensity and counts of

M. lucifugus at hibernacula have suggested development of either tolerance or resistance in

these persisting populations (Frick et al., 2017), which is supported by reports on infected

individuals not arousing from torpor as frequently as during the acute phase of the zoonosis

(Lilley et al., 2016). Since WNS has inflicted massive population declines in some *M.*

lucifugus, there is a possibility that selection could occur for alleles conferring resistance or

tolerance from the standing genetic variation, which would confer to resistance or tolerance

could occur. Indeed, Maslo and Feffermann (2015) suggested the occurrence of evolutionary

rescue occurring and Donaldson et al. (2017) described changes in an immunome across *M.*

lucifugus populations obtained through a sequence capture method, which may be attributed

to selection by WNS. A study suggests the initially affected *M. lucifugus* population is

beginning to show signs of resistance to the pathogen (Langwig et al., 2017). However, the

possible selective pressures WNS has exerted on the population, which may not even be

related to immune responses (e.g. Cheng et al., 2019), have only recently been described at

the whole-genome level (Gignoux-Wolfsohn et al., 2018), however high-sequencing depth

studies are lacking.

Here, we utilise high-throughput whole-genome sequencing on the WNS-affected species *M.*

lucifugus, allowing us to look at entire genomes via single nucleotide polymorphisms (SNPs).

We compare genetic patterns (F_{ST} and heterozygosity) before and after the spread of WNS in

the eastern North American population to gain insight into whether the disease is causing

selection of major loci in surviving bats. We hypothesise that, due to the massive population

declines caused by WNS and the fact that the affected population is now beginning to

stabilise or slightly increase in size (Langwig et al., 2017), a soft selective sweep based on

standing genetic variation has resulted in differentiation in one or many regions of the

genome (Messer and Petrov, 2013). Adaptation could therefore be in the process of driving

the evolutionary rescue of a population, in which individuals resistant or tolerant to the

disease survived a bottleneck and passed these traits to their offspring (Carlson et al., 2014).

Alternatively, survival may be due to factors not related to selection for WNS-resistant

alleles, such as microbiome adaptation (Ange-Stark et al., 2019; Lemieux-Labonté et al.,

2017) or variation across hibernation site microhabitats. This study provides a foundation for

future work on genome-wide effects of WNS in *M. lucifugus* and other affected species of

bats.

2. Materials and methods

2.1. Ethics Statement

This study was carried out on bats from non-endangered species in strict accordance with

recommendations in the Guide for the Care and Use of Laboratory Animals of the National

Institutes of Health. All methods were approved by the Institutional Animal Care and Use

Committee at Bucknell University (protocol DMR-016). The bats were collected under

Pennsylvania Game Commission Special Use Permit #33085, and New York State

Department of Environmental Conservation Permit #427.

2.2. Sample collection and DNA extraction

We conducted whole-genome sequencing on a population of *M. lucifugus* prior to, and up to 10 years after, the onset of WNS. White-nose syndrome in bats was first observed in upstate New York, at Howe's Cave during the winter of 2006-2007 (Blehert et al., 2009). Our historic samples were obtained from individuals consisting of the eastern population of *M. lucifugus* (Miller-Butterworth et al., 2014; Vonhof et al., 2015) from New York (NY) (from the first batch of dead bats found at Howe's Cave in 2006) and from bats caught in central Pennsylvania (PA) in 2005-2006 (Fig. 1., Table 1.,Supp. Table 1.). The bats sampled from Howe's Cave, New York were affected by WNS, but because they were the first bats to be documented with the disease in North America, and moreover, because of the massive declines caused by the disease, we believe this batch of samples is representative of the population genetic structure of *M. lucifugus* prior to WNS. Bats in PA became affected by the disease in roughly 2008–2009 and had not been affected by WNS at the time of sampling. We sampled wing tissue from frozen specimens of known origin (Supp. Table 1.) that were collected and stored by the New York State Department of Environmental Conservation (Delmar, NY) and D. Reeder to obtain Pre-WNS samples. The remnants of the eastern North American population in NY and PA were sampled again in 2016, and genetic diversity was compared before and after WNS (Fig.1., Table 1., Supp. Table 1.). Post-WNS bats were captured using mist nets and harp traps during 2015 -2016 from a number of maternal colonies, with the exception of individuals from Williams Preserve Mine, NY and Woodward, Center County, PA, which were sampled from hibernacula. In addition to samples from PA and NY, we collected gDNA from two individuals from Upper-Peninsula Michigan (UPMI) for use in polymorphism detection. We did this in order to take advantage

of the greater number of polymorphic sites one would expect to arise from a more diverse

collection of individuals (see Section 2.3.). Tissue samples were collected using either 3.0

mm or 5.0 mm biopsy punches (Integra Milltex, Plainsboro, NJ) and stored in 95% ethanol

until extraction. We extracted DNA using QIAmp DNA Mini Kits (Qiagen, Hilden Germany)

and stored DNA at -80°C until sequencing.

2.3. Sequencing methods

Sequencing libraries were made for two separate sets of data: **(1)** Six individuals from PA (N

= 2), NY (N = 2) and UPMI (N= 2) were sequenced separately in order to initially identify a

set of polymorphic sites within the *M. lucifugus* genome within the focal population (See

Supp. Table 1. for information on individuals). Individual sequencing increases the

confidence of SNP discovery, relative to pooled sequencing, since individuals exhibit discrete

variation in the number of alleles they carry at each potential SNP site; **(2)** Equimolar pools

of DNA from multiple individuals from within our north-eastern US population (NY and PA)

at each of the two timepoints (pre- and post-WNS) to give four pooled sequencing libraries

(See Table 1. and Supp. Table 1. for information on individuals). TruSeq Nano libraries with

a 350 bp insert size were prepared from all samples and run on a HiSeq4000 platform to

generate 2x 150 bp reads providing ~15 Gbp of sequence data (~7.5x coverage) for each of

the six individuals **(1)** and ~90 Gbp of sequence data (~45x coverage) for each of the four

pools **(2)**. Adapter sequences were removed with Cutadapt v1.2.1 and trimmed with Sickle

1.200 with a minimum quality score of 20.

 2.4. Bioinformatics methods

First, reads from the six individually sequenced individuals (see 2.3.) were mapped onto the

M. lucifugus reference genome (https://www.broadinstitute.org/brown-bat/little-brown-bat-

genome-project) using Bowtie2 v2.2.5 (Langmead and Salzberg, 2012), with the '—

sensitive-local' preset. Reads were sorted using SAMtools v1.1 (Li et al., 2009) and

combined by read group (with duplicates removed using a 'lenient' validation stringency

setting) using Picard-tools v2.5.0 (http://broadinstitute.github.io/picard). SNPs were called

across all six individuals with the Genome Analysis Toolkit (GATK) v3.4 HaplotypeCaller

(McKenna et al., 2010) using default parameters. Within this SNP set, high-quality bivalent

SNPs were selected that met selection criteria across the six individuals (Quality Score > 100;

AF<1; 30>=DP>=100) and were input to the SAMtools mpileup function as a list of sites at

which pileup was to be generated. After SNP calling, these six individuals were discarded

from further analyses.

Reads from the four pooled samples were mapped to the *M. lucifugus* reference genome as

above and SAMtools mpileup was used to count alleles at high-quality SNP sites identified

above from the six sequenced individuals (i.e. reads matching either the reference or alternate

allele for bivalent loci). Sites with coverage in the bottom 10% or top 10% quantiles summed

across the pooled samples were excluded (since these could reflect inaccuracies in the

reference genome), leaving about 13.5 million sites.

Average heterozygosity within the pooled samples was calculated using npstats (Ferretti et

al., 2013), which controls for number of individuals within each sequencing pool and

sequencing depth, thus allowing robust comparisons between pooled samples. One million

SNPs were randomly picked and permuted, with 1000 sets of 1000 SNPs used to calculate

the observed heterozygosity within each pooled set of individuals and, from these, we

produced the median and 95% confidence intervals on heterozygosity.

Using allele counts at each locus to estimate allele frequencies, F_{ST} was calculated without

weighting for sampling effort ($F_{ST} = \left(\bar{p}(1-\bar{p}) - \overline{p(1-p)}\right)/\bar{p}(1-\bar{p})$, where p is the allele

frequency and either averaged between populations ($\bar{p}$) or the value $p(1-p)$ averaged

across populations). F_{ST} was separately calculated between the Pre-WNS and Post-WNS

samples and between subsections thereof, specifically PA pre- vs post-WNS, NY pre- vs

post-WNS, PA pre- vs NY pre-WNS, PA post- vs NY post-WNS, and all PA vs all NY).

Selection is expected to elevate F_{ST} at specific genomic regions relative to background levels

due to sampling noise or genetic drift, which are expected to act equally across the genome.

To identify such regions across the genome, a moving window was applied to calculate

median pairwise F_{ST} values for groups of 500 consecutive SNPs at a time to even out

sampling effects. Moving windows shifted 100 SNPs at a time, thus providing the best

balance between fine detail and practicality. Allele frequency estimates were weighted by the

number of individuals in each pool. This was done using a custom R script (see Data

availability section), run using R v3.4.1 (R Development Core Team, 2011). Results were

confirmed using Popoolation2 with a 10 kb sliding window and with defined pool sizes. This

gave the same qualitative results, but higher levels of background noise (Kofler et al., 2011).

Additionally, Popoolation2 was used to perform Fishers Exact Tests on individual SNPs in

order to test whether any very highly significant SNPs were missed by the moving window

approach.

From the comparison of the NY and PA pooled samples (i.e. the comparison that produced

the highest F_{ST} values), we visually identified across the whole genome windows with

median F_{ST} values greater than 5 standard deviations from the mean, as well as those with F_{ST}

values greater than 0.05, indicating at least 'moderate' genetic differentiation (Wright, 1978).

The numbers of such complete windows within gene models were counted for each gene to

identify genes potentially under selective pressure.

3. Results

3.1. Pre-WNS vs Post-WNS comparisons

To detect regions of the genome that may be under selection, we plotted F_{ST} across the *M.*

lucifugus genome, calculated from allele counts within moving windows covering 500 SNPs

and shifting 100 SNPs at a time. Comparing Pre- and Post-WNS samples revealed an overall

low level of population differentiation due to genetic structure (F_{ST} median of 0.0059; Table

2). Whilst a small number of windows have F_{ST} values slightly beyond our arbitrary cut-off

of 5 standard deviations above the mean, none have F_{ST} values exceeding 0.05, suggesting

that there is not even moderate selection in any region of the genome in Post-WNS samples

compared to Pre-WNS samples (Fig. 2, Table 2)(Wright, 1978). Similarly, using Fisher exact

tests from Popoolation2 at a threshold of $p < 10^{-6}$, no individual SNPs were found to be

consistently associated with differences Pre- versus Post-WNS among the four sets of

samples.

3.2. Comparisons within geographical locations

Due to our sampling and sequencing setup, we were also able to separately look at individuals of the same population from two discrete geographical locations within their range. In our case a divide was made between bats sampled in New York and Pennsylvania based on geographical distance between the sampling sites. Although population genetic analysis based on a considerably lower number of genetic markers assigns *M. lucifugus* on the eastern coast of North America to the same population (Vonhof et al., 2015), the species shows high degrees of philopatry (Davis and Hitchcock, 1965; Norquay et al., 2013), warranting the examination of these arbitrarily assigned pools of individuals outside the mean dispersal range of the species using more accurate, whole-genome sequencing based population genetic analyses.

A number of minor differences ($F_{ST} >$ 5 stdevs above mean, but $F_{ST} <$ 0.05) between these sets of samples are evident prior to arrival of WNS, as well as a peak in scaffold GL429835 with an F_{ST} value exceeding 0.05 (Fig. 3a). The high peak consists of 4 windows the midpoints of which span an 81,887 bp region within a coding sequence consisting of 47 exons, which encodes the NIPBL cohesion loading factor (NCBI Gene ID: 102430106). The highest peak in a comparison of PA populations Pre- and Post-WNS (Fig. 3b) lies in a similar position within this coding region, as does a peak in a similar comparison of NY samples (Fig. 3c), though neither exceed the F_{ST} value cut-off of 0.05.

3.3. Post-WNS comparisons between geographically isolated sample sets

When comparing the Post-WNS individuals sampled from PA with the Post-WNS individuals from NY, the peak on scaffold GL429835 is no longer visible, but a far more

pronounced spike in F_{ST} values can be clearly seen in scaffold GL429776 (Fig. 4a),

indicating relatively large allele frequency differences between the sample sets in this region

of the genome. This is also seen, to a lesser extent, when comparing all samples from NY

with all samples from PA (Fig. 4b). However, this significant peak in this comparison of all

samples collected does not affect average fixation indices across the genome relative to the

other comparison of all samples, Pre-WNS vs Post-WNS (Table 2). This peak could be

attributed to mapping error, but our coverage depths show an average depth of 66.33x across

the whole genome, whereas the focal region was sequenced at an average depth of 66.49x.

This means that there has been no collapsing of duplicated regions, nor poor coverage of the

region. Also, viewing the data on the NCBI genome viewer tool, with RepeatMasker

histogram enabled, suggested no immediately obvious increase in repetitive sequence in the

region of interest.

We examined scaffold GL429776 more closely to compare the F_{ST} values between

individuals from the arbitrarily selected sample sets from NY Post-WNS and PA Post-WNS

(Fig. 4a). Genes within scaffold GL429776 containing windows with fixation indices > 0.05

between the NY Post-WNS and PA Post-WNS samples, are presented in Table 4 and Fig. 4a.

To clarify why we saw no fixation at this site when comparing Pre-WNS and Post-WNS

populations (Fig. 3a), nor when we compared Pre-WNS samples from both geographical

origins (Figs. 3b and 3c), we calculated for the relevant subpopulations the proportion of

reference alleles called at each site included in the F_{ST} window analyses (Fig. 5). This

revealed a small decrease over time in the proportion of the reference alleles seen at this

locus in the PA subpopulation (Fig. 5a), commensurate with an increase in proportion of

reference alleles in the NY sample set (Fig. 5b). As such, existing differences between the

NY and PA sample sets were insufficient to cause a relative rise in fixation indices prior to

WNS spreading through the population (Fig 5c). Conversely, Post-WNS, allele proportions at

this site in both sample sets had not changed sufficiently to suggest selection *within* the

sample set but demonstrated clear differences *between* the NY and PA sample sets (Fig. 5d).

In the Post-WNS comparison, peaks which were above our arbitrary threshold of 5 standard

deviations above the mean across the whole genome, but below $F_{ST} = 0.05$, can also be

observed in small scaffolds on the right-hand side of the F_{ST} plots. The 63 windows

comprising these peaks lie in 47 scaffolds. Of these, high F_{ST} windows lie in only 6 protein-

coding regions across 6 scaffolds (Table 3). Two of these encode partial gene models. Of the

4 remaining models containing high F_{ST} regions, all but 1 is described on NCBI as a "Low

Quality Protein". The exception is a gene encoding the ankyrin 2 protein, within which 7 high

F_{ST} windows lie.

3.4. Measures of heterozygosity within subpopulations

We examined heterozygosity across the genome in the subpopulations divided by time and

location to measure whether the epizootic reduces genetic diversity. We found the following

values: PA pre-WNS: Median: 0.263 (0.252 – 0.274 95% CI); PA post-WNS: 0.247 (0.239 –

0.255); NY pre-WNS: 0.255 (0.247 – 0.263); and NY post-WNS: 0.256 (0.247 -0.265).

4. Discussion

Examining our data as a whole, we found low F_{ST} values between the Pre- and Post-WNS

individuals of our eastern North American *M. lucifugus* population, the absence of fixation

suggesting that WNS has not decreased the population in size sufficiently for the disease to

have an effect on genetic diversity. Also, we found no significant changes in within-

population heterozygosity Post-WNS, despite the massive population declines caused by

WNS over the past decade, and found overall high heterozygosity across the genome. These

results are in accordance with a similar, low-coverage study conducted by Gignoux-

Wolfsohn et al. (2018). Panmixia of populations in evolutionary time in bats to the east of the

Rocky Mountains has been previously documented (Vonhof et al., 2016; Wilder et al., 2015).

Gignoux-Wolfsohn et al. (2018) present evidence for a soft selective sweep, where selection

acts on standing genetic variation, between Pre- and Post-WNS populations. The analysis is

based on a change in minor allele frequencies due to selection, distinguished from

background demography using a simulation approach; and it shows a complex picture of

potentially up to 85 unlinked SNPs, each exhibiting small effects and with little or no

selective signal at adjacent SNPs. In contrast, a recent case study of selection on standing

genetic variation, by myxoma virus in rabbits, showed that soft selective sweeps caused

concerted changes in allele frequencies at sets of linked SNPs around coding loci (Alves et

al., 2019). The robust pooled sequencing approach allows us to examine for similar, larger

scale changes in the genome. We acknowledge that both methods have their advantages and

disadvantages, such as the increased possibility for false outliers in our pooled approach (Li

et al., 2008). Hovewer we took these into account with parameters used for aligning the reads

(Kofler et al., 2011), sequencing depth (Ferretti et al., 2013; Kofler and Schlötterer, 2014)

pool size (Futschik and Schlötterer, 2010; Gautier, 2015) and library preparation protocol

(Kofler et al., 2016).

The species originally described as *M. lucifugus* in North America (Fenton and Barclay,

1980) has since been divided into five non-sister species (Morales and Carstens, 2018). All of

the individuals sampled in our study belong to *M. lucifugus senso stricto*, with a range

extending from the east coast of North America to Alaska, and furthermore, to the same

previously assigned population (Miller-Butterworth et al., 2014). Previous nuclear genetic

studies show that differentiation is low, and there is no evidence for any major barriers to

nuclear gene flow across the range of *M. lucifugus*.

Due to our sampling design, we were able to separate bats geographically within our

sampling range. We made a cut-off point for a more in-depth investigation of bats sampled in

the states of Pennsylvania (PA) compared to New York (NY). Although the cut-off may seem

rather arbitrary, the reasoning behind this is geographic distance between sampling sites, and

diving the data between the two states is appropriate for our purposes as evidenced by the

results. Here, we see low F_{ST} values between the two groups of bats prior to the onset of

WNS, indicating an absence of selection across the genome, with the exception of a small

peak in scaffold GL429835, which lies within a gene model encoding the NIPBL cohesion

loading factor. Relatively high F_{ST} values are also visible within this region when comparing

the PA population before and after the spread of WNS, as well as the NY population pre- and

post-WNS. However, this peak is not evident when comparing PA and NY post-WNS (Fig.

4a). This suggests that this coding sequence was under selective pressures, manifesting as

differentiating allelic frequencies, in both subpopulations prior to the arrival of WNS, but that

any selection was nullified by the population-level impact of WNS (or another event

occurring during the same timeframe). In addition, this result confirms that although the pre-

WNS bats from NY were from the initial batch of bats found dead of WNS at Howe's cave in

2006, they were genetically similar to pre-WNS bats from PA, which were sampled before

the arrival of WNS. This is most likely because of the high frequency of initial mortality in

the population, which indiscriminately wiped out most of the population, eroding any genetic

differentiation. As for the protein NIPBL, it plays an important role in the function of the

cohesin complex, which is vital for chromosome segregation, DNA damage repair, and gene

expression regulation (Peters et al., 2008). As these results suggest the loss of low-level

functional diversity at this locus in our subpopulations, it would be interesting to further

investigate what impact this will have on the population as a whole.

The F_{ST} comparison of Post-WNS individuals sampled from PA and NY depicts considerable

genetic differentiation within scaffold GL429776; differentiation which did not exist in the

Pre-WNS comparison of individuals sampled from these geographical areas. This is a

difference which would have gone unnoticed had we not made the comparison between bats

sampled in NY and PA, or used fewer genetic markers instead of a whole-genome approach

(Miller-Butterworth et al., 2014; Vonhof et al., 2015). The region's appearance post-WNS

when considering all geographical locations together is not, however, reflected when the data

are broken down into representatives of each geographic location. A comparison of allele

frequencies of individual SNPs within this region shows that individuals sampled from PA

showed an increase in deviation from the reference genome over time, whilst a greater

proportion of reference alleles were seen in NY samples Post-WNS compared with Pre-

WNS. The differences in proportions, growing as they were over time, were insufficient to

exhibit a high fixation index Pre-WNS, but were notably different after the disease had

spread through the population. One could argue, that this is due to sampling at maternal

colonies and swarming sites, which has known effects on allele frequency differences

(Johnson et al., 2015). However, we sampled at a number of maternal colonies, which would

dilute this effect, and furthermore, recent studies suggest philopatry within maternal colonies

may not be as high as previously expected (Olivera-Hyde et al., 2019, although on *M.*

septentrionalis). Therefore we assume our sampling of multiple maternal colonies and

hibernation sites has removed any effects of relatedness within our sample sets.

This region within the scaffold containing the peak contains six annotated genes with very

different functions. Two, POC1B and CEP290, are involved in cilia development and

maintenance. The ATP2B1 gene encodes a protein which belongs to the family of P-type

primary ion transport ATPases, whereas TMTC3 is involved in the control of endoplasmic

reticulum stress response. MGAT4C is a glycosyltransferase that participates in the transfer

of N-acetylglucosamine (GlcNAc) to the core mannose residues of N-linked glycans. Finally,

KITLG produces a protein involved in mast cell development, migration and function, and

melanogenesis. KITLG has been documented as a target of evolution in recent studies and in

humans it has been linked with thermoregulation (Yang et al., 2018), which is a critical

component of bat life history (Studier and O'Farrell, 1972). In fact, upregulation of KITLG at

low temperature helps promote the production of brown fat for heat generation (Huang et al.,

2014). Bats utilise brown adipose tissue in arousals to normothermia from different degrees

of torpor (Thomas et al., 1990). There is also evidence of parallel evolution of pigmentation

in sticklebacks and humans linked to KITLG (Miller et al., 2007), as well as its role in

determining resistance/susceptibility to swine respiratory diseases in Erhulian pigs (Huang et

al., 2017). KITLG, by influencing mast cell development, could also play a role in either

protective or pathological immune responses to *P. destructans*, which may involve IgE-

mediated recognition of secreted fungal proteins. As such, temporal changes we found in

allele frequencies at this locus could, for instance, be linked to differences in hibernation site

temperatures, and/or thermoregulatory needs in bats sampled at different locations

(Humphries et al., 2002; Thomas et al., 1990). Spatial patterns, as seen here between the

arbitrarily pooled samples from PA and NY, could be due to local adaptation, in which strong

selective sweeps may be linked to gene variants favoured in local interactions (Hansen et al.,

2012; Kyle et al., 2014; Rico et al., 2015; Schoville et al., 2012). Whilst we lack data

concerning migration patterns of the bats studied here, as well as the environmental

conditions within their microfugia, future work would benefit from these data in a bid to

understand whether bats repeatedly frequent habitats with similar conditions and whether,

indeed, there is a significant difference in conditions across these Eastern states that explains

the rapid rise in allelic differentiation, and thus F_{ST} values, at this locus.

The increase in F_{ST} values in this scaffold GL429776 could, alternatively, be due to the

massive population decline caused by WNS (Langwig et al., 2017). Although not supported

by existing data, the decline could potentially have resulted in a decrease in connectivity and

gene flow between remnant local populations allowing for a local adaptation to arise

(Bohonak, 1999; Lilley, Anttila, & Ruokolainen, 2018). Genetically resolved examples of

local adaptations are rare (Hoekstra et al., 2006) and, in our focal system, pinpointing the

causality leading to local adaptation is especially difficult due to the cryptic life history of

bats. Determining how these variants are spread or localised among populations is essential

to understanding and managing the emergence of new selective pressures (Eizaguirre et al.,

2012; Kyle et al., 2014). We cannot fully exclude the possibility of WNS selecting for alleles

in our scaffold of interest though. It is likely that there is concurrent selection occurring

across multiple populations, and that selection is not acting to the same degree or even on the

432 same loci in different geographic areas affected by WNS (Messer and Petrov, 2013). This

433 also applies to interpreting the results on a smaller scale: the samples had not been collected

434 from exactly the same locations before and after WNS, which may contribute to our results.

435 Individuals sampled during the winter, at their hibernacula, may not inhabit the vicinity of

436 these locations during the summer, although dispersal distances are often short (Davis &

437 Hitchcock, 1965; but see Norquay et al., 2013). Understanding how local separate sub-

438 populations cope with pathogen pressure would benefit from an RNAseq-approach

439 identifying responses at transcript-level (Lilley et al., 2019).

440 The ankyrin 2 protein encoding gene, lying within the observed small scaffolds on the right-

441 hand side of the F_{ST} plots has a wide range of attributed Gene Ontology functions (The

442 UniProt Consortium, 2019), which reinforce the ubiquitous nature and far-reaching roles of

443 ankyrin-repeat-containing genes across organisms (Voronin and Kiseleva, 2007).

444 Alternatively, the peaks in this area may have arisen because only relatively few SNPs were

445 called there given the small size of the scaffolds, thus limiting our ability to generate high-

446 quality windows, hence the greater sampling noise.

447 Despite not seeing an overarching genetic signal associated with resistance in *M. lucifugus*,

448 remnant populations appear to be surviving year after year and even increasing in size

449 notwithstanding infection loads during the hibernation period (Langwig et al., 2017; Lilley et

450 al., 2016). The fact that the intensity of infection with *P. destructans* appears to be

451 significantly lower for remnant populations compared to colonies in the midst of the

452 epidemic phase and massive declines may suggest the development of resistance in these

453 populations (Bernard et al., 2017; Langwig et al., 2017; but see Lilley et al., 2019). However,

the role of environmental interactions during the hibernation period, when the bat hosts are

most vulnerable to the pathogen, have been largely neglected, although the effect of humidity

and temperature on the growth of the fungal pathogen and survival of the host have been

separately studied (Grieneisen et al., 2015; Marroquin et al., 2017; Verant et al., 2012).

Modelling also suggests that environmental factors, including the cave microbiome, have an

impact on the proliferation and infectivity of *P. destructans* (Hayman, Pulliam, Marshall,

Cryan, & Webb, 2016; Lilley et al., 2018). This is supported by the discovery of microbes in

hibernacula environments and bats that are able to retard the growth of the fungus (Micalizzi

et al., 2017; Zhang et al., 2014, 2015). In light of our results and considering bat life history

as a whole, adaptation and evolutionary rescue may not be a fast track for recovery of bat

populations affected by WNS (Maslo and Fefferman, 2015), although ultimately these will

contribute to long term survival of populations (Lilley et al., 2019). At the present, it is more

likely the variability in hibernation site environmental conditions and strong selection for

microbial taxa that inhibit *P. destructans* could explain why populations are recovering from

the zoonosis (Cheng et al., 2016; Lemieux-Labonté, Simard, Willis, & Lapointe, 2017; Lilley

et al., 2018).

It is still unclear how frequently genetic adaptation occurs in natural populations and under

what circumstances it is promoted (Schoville et al., 2012). Many studies have recorded

phenotypic changes in natural populations and attributed them to host-pathogen interactions

(Frank et al., 2014; Kilpatrick, 2006; Langwig et al., 2017; Råberg et al., 2009). However, it

is difficult to determine conclusively whether changes in phenotype are the result of selection

in the genome or a result of phenotypic plasticity (Paterson et al., 2010; Routtu and Ebert,

2015). Also, the assumed benefit of change (i.e. adaptation) is often not tested

experimentally, which would allow for inference of causality with the pathogen through

exclusion of other potential drivers.

To conclude, our results suggest WNS has not yet decidedly subjected populations of *M.*

lucifugus to selective pressures leading to genetic adaptation in remnant populations. No

population -wide signs of selection in comparisons of genomes of Pre-WNS and ten years

Post-WNS populations were observed (but see Donaldson et al., 2017; Gignoux-Wolfsohn et

al., 2018). This is something that is not unexpected with a species in which the effective

population size most likely numbered in tens of millions of individuals prior to the onset of

WNS. The disease has not caused a true population bottle-neck, as exemplified by our

measures of genetic diversity not varying between pre- and post-WNS samples. Our results

indicate that the existence and recent growth of remnant populations of *M. lucifugus* are more

likely attributable to other factors, such as microbiome adaptation and hibernation site

selection among others, rather than genetic adaption. However, we found increased

variability in a specific area of the genome in a comparison of Post-WNS samples from our

two different study states, relative to Pre-WNS samples. We suggest this is due to weakened

connectivity between bats at different locations allowing for local adaptations to enrich in the

absence of gene flow. Although genes in the high F_{ST} region of the genome were identified,

and genes such as KITLG provide interesting avenues of research, even with regards to

WNS, further investigation into the processes in bat life history to which these genes are

related are required to expound upon the existing gene annotations.

Acknowledgements

We thank Svenska Kulturfonden, H2020 Marie Skłodowska☐Curie Actions (TML, 706196) and the Natural Environment Research Council, UK for funding the study; staff at the Centre for Genomic Research, University of Liverpool, UK for DNA sequencing expertise; Jenni Prokkola and Riley F. Bernard for proofreading the manuscript; and Alice Lilley, Cali Wilson, Beth Rogers and Mike Scafini for assisting in sample collection.

Author contributions

TML and SP designed the research. TML, DMR, KAF, AK, GGR, MEV and CH collected the samples. TML prepared samples for sequencing. TML, IWW and SP analysed the data. TML, IWW and SP wrote the first draft of the manuscript, and all authors contributed to the final version.

Data availability statement

Raw data are available at ENA accession XXXXX and bioinformatics code at https://github.com/scottishwormboy/myoluc.

References

Alves, J.M., Carneiro, M., Cheng, J.Y., Matos, A.L. de, Rahman, M.M., Loog, L., Campos, P.F., Wales, N., Eriksson, A., Manica, A., et al. (2019). Parallel adaptation of rabbit populations to myxoma virus. Science *363*, 1319–1326.

Ange-Stark, M., Cheng, T.L., Hoyt, J.R., Langwig, K.E., Parise, K.L., Frick, W.F.,

Kilpatrick, A.M., MacManes, M.D., and Foster, J.T. (2019). White-nose syndrome

restructures bat skin microbiomes. BioRxiv 614842.

Bernard, R.F., and McCracken, G.F. (2017). Winter behavior of bats and the progression of

white□nose syndrome in the southeastern United States. Ecol. Evol. *7*, 1487–1496.

Bernard, R.F., Willcox, E.V., Parise, K.L., Foster, J.T., and McCracken, G.F. (2017). White-

nose syndrome fungus, Pseudogymnoascus destructans, on bats captured emerging from

caves during winter in the southeastern United States. BMC Zool. *2*, 12.

Blehert, D.S., Hicks, A.C., Behr, M., Meteyer, C.U., Berlowski-Zier, B.M., Buckles, E.L.,

Coleman, J.T.H., Darling, S.R., Gargas, A., Niver, R., et al. (2009). Bat white-nose

syndrome: an emerging fungal pathogen? Science *323*, 227–227.

Bohonak, A.J. (1999). Dispersal, gene flow, and population structure. Q. Rev. Biol. *74*, 21–

45.

Boots, M., Best, A., Miller, M.R., and White, A. (2009). The role of ecological feedbacks in

the evolution of host defence: what does theory tell us? Philos. Trans. R. Soc. Lond. B Biol.

Sci. *364*, 27–36.

Campana, M., Kurata, N., Foster, J.T., Helgen, L., Reeder, D., Fleischer, R., and Helgen,

K.M. (2017). White-nose syndrome fungus in a 1918 bat specimen from France. Emerg.

Infect. Dis. J. *23*, 1–1.

Campbell, M.C., and Tishkoff, S.A. (2008). African Genetic Diversity: Implications for

Human Demographic History, Modern Human Origins, and Complex Disease Mapping.

Annu. Rev. Genomics Hum. Genet. *9*, 403–433.

Carlson, S.M., Cunningham, C.J., and Westley, P.A.H. (2014). Evolutionary rescue in a

changing world. Trends Ecol. Evol. *29*, 521–530.

Cheng, T.L., Mayberry, H., McGuire, L.P., Hoyt, J.R., Langwig, K.E., Nguyen, H., Parise,

K.L., Foster, J.T., Willis, C.K.R., Kilpatrick, A.M., et al. (2016). Efficacy of a probiotic

bacterium to treat bats affected by the disease white-nose syndrome. J. Appl. Ecol. *54*, 701–

708.

Cheng, T.L., Gerson, A., Moore, M.S., Reichard, J.D., DeSimone, J., Willis, C.K.R., Frick,

W.F., and Kilpatrick, A.M. (2019). Higher fat stores contribute to persistence of little brown

bat populations with white-nose syndrome. J. Anim. Ecol. *0*.

Cryan, P.M., Meteyer, C.U., Boyles, J.G., and Blehert, D.S. (2010). Wing pathology of

white-nose syndrome in bats suggests life-threatening disruption of physiology. Bmc Biol. *8*,

135.

Cryan, P.M., Meteyer, C.U., Blehert, D.S., Lorch, J.M., Reeder, D.M., Turner, G.G., Webb,

J., Behr, M., Verant, M., Russe, R.E., et al. (2013). Electrolyte depletion in white-nose

syndrome bats. J. Wildl. Dis. *49*, 398–402.

Daszak, P., Berger, L., Cunningham, A.A., Hyatt, A.D., Green, D.E., and Speare, R. (1999).

Emerging infectious diseases and amphibian population declines. Emerg. Infect. Dis. *5*, 735–

748.

Daszak, P., Cunningham, A.A., and Hyatt, A.D. (2000). Emerging infectious diseases of

wildlife-threats to biodiversity and human health. Science *287*, 443–449.

Davis, W.H., and Hitchcock, H.B. (1965). Biology and migration of the bat, *Myotis lucifugus*,

in New England. J. Mammal. *46*, 296–313.

Davy, C.M., Donaldson, M.E., Rico, Y., Lausen, C.L., Dogantzis, K., Ritchie, K., Willis,

C.K.R., Burles, D.W., Jung, T.S., McBurney, S., et al. (2017a). Prelude to a panzootic: Gene

flow and immunogenetic variation in northern little brown myotis vulnerable to bat white-

nose syndrome. FACETS.

Davy, C.M., Donaldson, M.E., Willis, C.K.R., Saville, B.J., McGuire, L.P., Mayberry, H.,

Wilcox, A., Wibbelt, G., Misra, V., Bollinger, T., et al. (2017b). The other white-nose

syndrome transcriptome: Tolerant and susceptible hosts respond differently to the pathogen

Pseudogymnoascus destructans. Ecol. Evol. 7161–7170.

De Castro, F., and Bolker, B. (2005). Mechanisms of disease-induced extinction. Ecol. Lett.

8, 117–126.

Deschamps, M., Laval, G., Fagny, M., Itan, Y., Abel, L., Casanova, J.-L., Patin, E., and

Quintana-Murci, L. (2016). Genomic Signatures of Selective Pressures and Introgression

from Archaic Hominins at Human Innate Immunity Genes. Am. J. Hum. Genet. *98*, 5–21.

Di Gaspero, G., Copetti, D., Coleman, C., Castellarin, S.D., Eibach, R., Kozma, P., Lacombe,

T., Gambetta, G., Zvyagin, A., Cindrić, P., et al. (2012). Selective sweep at the Rpv3 locus

during grapevine breeding for downy mildew resistance. TAG Theor. Appl. Genet. Theor.

Angew. Genet. *124*, 277–286.

Dobony, C.A., and Johnson, J.B. (2018). Observed resiliency of little brown myotis to long-

term white-nose syndrome exposure. J. Fish Wildl. Manag. *9*, 168–179.

Donaldson, M.E., Davy, C.M., Willis, C.K.R., McBurney, S., Park, A., and Kyle, C.J. (2017).

Profiling the immunome of little brown myotis provides a yardstick for measuring the genetic

response to white-nose syndrome. Evol. Appl. *10*, 1076–1090.

Eizaguirre, C., Lenz, T.L., Kalbe, M., and Milinski, M. (2012). Divergent selection on locally

adapted major histocompatibility complex immune genes experimentally proven in the field.

Ecol. Lett. *15*, 723–731.

Fenton, M.B., and Barclay, R.M.R. (1980). *Myotis lucifugus*. Mamm. Species 1–8.

Ferretti, L., Ramos-Onsins, S.E., and Pérez-Enciso, M. (2013). Population genomics from

pool sequencing. Mol. Ecol. *22*, 5561–5576.

Field, K.A., Johnson, J., Lilley, T., Reeder, S., Rogers, E., Behr, M., and Reeder, D. (2015).

The white-nose syndrome transcriptome: activation of anti-fungal host responses in wing

tissue of hibernating bats. Plos Pathog. *11*, e1005168.

Field, K.A., Sewall, B.J., Prokkola, J.M., Turner, G.G., Gagnon, M., Lilley, T.M., White,

J.P., Johnson, J.S., Hauer, C.L., and Reeder, D.M. (2018). Effect of torpor on host

transcriptomic responses to a fungal pathogen in hibernating bats. Mol. Ecol.

Frank, C.L., Michalski, A., McDonough, A.A., Rahimian, M., Rudd, R.J., and Herzog, C.

(2014). The resistance of a North American bat species (Eptesicus fuscus) to White-Nose

Syndrome (WNS). PLoS ONE *9*, e113958.

Frick, W.F., Pollock, J.F., Hicks, A.C., Langwig, K.E., Reynolds, D.S., Turner, G.G.,

Butchkoski, C.M., and Kunz, T.H. (2010). An Emerging disease causes regional population

collapse of a common North American bat species. Science *329*, 679–682.

Frick, W.F., Puechmaille, S.J., Hoyt, J.R., Nickel, B.A., Langwig, K.E., Foster, J.T., Barlow,

K.E., Bartonička, T., Feller, D., Haarsma, A.-J., et al. (2015). Disease alters macroecological

patterns of North American bats. Glob. Ecol. Biogeogr. *24*, 741–749.

Frick, W.F., Cheng, T.L., Langwig, K.E., Hoyt, J.R., Janicki, A.F., Parise, K.L., Foster, J.T.,

and Kilpatrick, A.M. (2017). Pathogen dynamics during invasion and establishment of white-

nose syndrome explain mechanisms of host persistence. Ecology *98*, 624–631.

Futschik, A., and Schlötterer, C. (2010). The next generation of molecular markers from

massively parallel sequencing of pooled DNA samples. Genetics *186*, 207–218.

Gallana, M., Ryser-Degiorgis, M.-P., Wahli, T., and Segner, H. (2013). Climate change and

infectious diseases of wildlife: Altered interactions between pathogens, vectors and hosts.

Curr. Zool. *59*, 427–437.

Gautier, M. (2015). Genome-Wide Scan for Adaptive Divergence and Association with

Population-Specific Covariates. Genetics *201*, 1555-+.

Gignoux-Wolfsohn, S.A., Pinsky, M.L., Kerwin, K., Herzog, C., Hall, M., Bennett, A.B.,

Fefferman, N.H., and Maslo, B. (2018). Genomic signatures of evolutionary rescue in bats

surviving white-nose syndrome. BioRxiv 470294.

Grieneisen, L.E., Brownlee-Bouboulis, S.A., Johnson, J.S., and Reeder, D.M. (2015). Sex

and hibernaculum temperature predict survivorship in white-nose syndrome affected little

brown myotis (Myotis lucifugus). R. Soc. Open Sci. *2*, 140470.

Groot, N.G. de, Otting, N., Doxiadis, G.G.M., Balla-Jhagjhoorsingh, S.S., Heeney, J.L.,

Rood, J.J. van, Gagneux, P., and Bontrop, R.E. (2002). Evidence for an ancient selective

sweep in the MHC class I gene repertoire of chimpanzees. Proc. Natl. Acad. Sci. *99*, 11748–

11753.

Hansen, M.M., Olivieri, I., Waller, D.M., and Nielsen, E.E. (2012). Monitoring adaptive

genetic responses to environmental change. Mol. Ecol. *21*, 1311–1329.

Harazim, M., Horáček, I., Jakešová, L., Luermann, K., Moravec, J.C., Morgan, S., Pikula, J.,

Sosík, P., Vavrušová, Z., Zahradníková, A., et al. (2018). Natural selection in bats with

historical exposure to white-nose syndrome. BMC Zool. *3*, 8.

Hayman, D.T.S., Pulliam, J.R.C., Marshall, J.C., Cryan, P.M., and Webb, C.T. (2016).

Environment, host, and fungal traits predict continental-scale white-nose syndrome in bats.

Sci. Adv. *2*, e1500831.

Hoekstra, H.E., Hirschmann, R.J., Bundey, R.A., Insel, P.A., and Crossland, J.P. (2006). A

single amino acid mutation contributes to adaptive beach mouse color pattern. Science *313*,

101–104.

Huang, X., Huang, T., Deng, W., Yan, G., Qiu, H., Huang, Y., Ke, S., Hou, Y., Zhang, Y.,

Zhang, Z., et al. (2017). Genome-wide association studies identify susceptibility loci

affecting respiratory disease in Chinese Erhualian pigs under natural conditions. Anim.

Genet. *48*, 30–37.

Huang, Z., Ruan, H.-B., Xian, L., Chen, W., Jiang, S., Song, A., Wang, Q., Shi, P., Gu, X.,

and Gao, X. (2014). The stem cell factor/Kit signalling pathway regulates mitochondrial

function and energy expenditure. Nat. Commun. *5*, 4282.

Humphries, M.M., Thomas, D.W., and Speakman, J.R. (2002). Climate-mediated energetic

constraints on the distribution of hibernating mammals. Nature *418*, 313–316.

Jeffery, K.J., and Bangham, C.R. (2000). Do infectious diseases drive MHC diversity?

Microbes Infect. *2*, 1335–1341.

Johnson, J.S., Reeder, D.M., McMichael, J.W., Meierhofer, M.B., Stern, D.W.F., Lumadue,

S.S., Sigler, L.E., Winters, H.D., Vodzak, M.E., Kurta, A., et al. (2014). Host, pathogen, and

environmental characteristics predict white-nose syndrome mortality in captive little brown

myotis (*Myotis lucifugus*). Plos One *9*, e112502.

Johnson, L.N.L., McLeod, B.A., Burns, L.E., Arseneault, K., Frasier, T.R., and Broders, H.G.

(2015). Population Genetic Structure Within and among Seasonal Site Types in the Little

Brown Bat (Myotis lucifugus) and the Northern Long-Eared Bat (M. septentrionalis). PLOS

ONE *10*, e0126309.

Karlsson, E.K., Kwiatkowski, D.P., and Sabeti, P.C. (2014). Natural selection and infectious

disease in human populations. Nat. Rev. Genet. *15*, 379–393.

Kilpatrick, A.M. (2006). Facilitating the evolution of resistance to avian malaria in Hawaiian

birds. Biol. Conserv. *128*, 475–485.

Kofler, R., and Schlötterer, C. (2014). A guide for the design of evolve and resequencing

studies. Mol. Biol. Evol. *31*, 474–483.

Kofler, R., Pandey, R.V., and Schlötterer, C. (2011). PoPoolation2: identifying differentiation

between populations using sequencing of pooled DNA samples (Pool-Seq). Bioinformatics

27, 3435–3436.

Kofler, R., Nolte, V., and Schlötterer, C. (2016). The impact of library preparation protocols

on the consistency of allele frequency estimates in Pool☐Seq data. Mol. Ecol. Resour. *16*,

118–122.

Kyle, C.J., Rico, Y., Castillo, S., Srithayakumar, V., Cullingham, C.I., White, B.N., and

Pond, B.A. (2014). Spatial patterns of neutral and functional genetic variations reveal

patterns of local adaptation in raccoon (Procyon lotor) populations exposed to raccoon rabies.

Mol. Ecol. *23*, 2287–2298.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.

Methods *9*, 357–359.

Langwig, K.E., Frick, W.F., Reynolds, R., Parise, K.L., Drees, K.P., Hoyt, J.R., Cheng, T.L.,

Kunz, T.H., Foster, J.T., and Kilpatrick, A.M. (2015). Host and pathogen ecology drive the

seasonal dynamics of a fungal disease, white-nose syndrome. Proc. R. Soc. B-Biol. Sci. *282*,

20142335.

Langwig, K.E., Hoyt, J.R., Parise, K.L., Frick, W.F., Foster, J.T., and Kilpatrick, A.M.

(2017). Resistance in persisting bat populations after white-nose syndrome invasion. Phil

Trans R Soc B *372*, 20160044.

Lemieux-Labonté, V., Simard, A., Willis, C.K.R., and Lapointe, F.-J. (2017). Enrichment of

beneficial bacteria in the skin microbiota of bats persisting with white-nose syndrome.

Microbiome *5*, 115.

Li, H., Ruan, J., and Durbin, R. (2008). Mapping short DNA sequencing reads and calling

variants using mapping quality scores. Genome Res. *18*, 1851–1858.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence

Alignment/Map format and SAMtools. Bioinforma. Oxf. Engl. *25*, 2078–2079.

Lilley, T., Anttila, J., and Ruokolainen, L. (2018). Landscape structure and ecology influence

the spread of a bat fungal disease. Funct. Ecol. *32*, 2483–2496.

Lilley, T.M., Johnson, J.S., Ruokolainen, L., Rogers, E.J., Wilson, C., Schell, S.M., Field,

K.A., and Reeder, D.M. (2016). White-nose syndrome survivors do not exhibit frequent

arousals associated with *Pseudogymnoascus destructans* infection. Front. Zool. *13*, 12.

Lilley, T.M., Prokkola, J.M., Johnson, J.S., Rogers, E.J., Gronsky, S., Kurta, A., Reeder,

D.M., and Field, K.A. (2017). Immune responses in hibernating little brown myotis (*Myotis*

lucifugus) with white-nose syndrome. Proc R Soc B *284*, 20162232.

Lilley, T.M., Prokkola, J.M., Blomberg, A.S., Paterson, S., Johnson, J.S., Turner, G.G.,

Bartonička, T., Bachorec, E., Reeder, D.M., and Field, K.A. (2019). Resistance is futile:

RNA-sequencing reveals differing responses to bat fungal pathogen in Nearctic *Myotis*

lucifugus and Palearctic *Myotis myotis*. Oecologia.

Lips, K.R., Brem, F., Brenes, R., Reeve, J.D., Alford, R.A., Voyles, J., Carey, C., Livo, L.,

Pessier, A.P., and Collins, J.P. (2006). Emerging infectious disease and the loss of

biodiversity in a Neotropical amphibian community. Proc. Natl. Acad. Sci. U. S. A. *103*,

3165–3170.

Lorch, J.M., Meteyer, C.U., Behr, M.J., Boyles, J.G., Cryan, P.M., Hicks, A.C., Ballmann,

A.E., Coleman, J.T.H., Redell, D.N., Reeder, D.M., et al. (2011). Experimental infection of

bats with *Geomyces destructans* causes white-nose syndrome. Nature *480*, 376-U129.

Lorch, J.M., Muller, L.K., Russell, R.E., O'Connor, M., Lindner, D.L., and Blehert, D.S.

(2013). Distribution and environmental persistence of the causative agent of white-nose

syndrome, *Geomyces destructans*, in bat hibernacula of the Eastern United States. Appl.

Environ. Microbiol. *79*, 1293–1301.

Luikart, G., Allendorf, F.W., Cornuet, J.-M., and Sherwin, W.B. (1998). Distortion of allele

frequency distributions provides a test for recent population bottlenecks. J. Hered. *89*, 238–

247.

Luis, A.D., and Hudson, P.J. (2006). Hibernation patterns in mammals: a role for bacterial

growth? Funct. Ecol. *20*, 471–477.

Marroquin, C.M., Lavine, J.O., and Windstam, S.T. (2017). Effect of Humidity on
Development of Pseudogymnoascus destructans, the Causal Agent of Bat White-Nose
Syndrome. Northeast. Nat. *24*, 54–64.

Maslo, B., and Fefferman, N.H. (2015). A case study of bats and white-nose syndrome
demonstrating how to model population viability with evolutionary effects. Conserv. Biol.
29, 1176–1185.

Maslo, B., Valent, M., Gumbs, J.F., and Frick, W.F. (2015). Conservation implications of
ameliorating survival of little brown bats with white-nose syndrome. Ecol. Appl. *25*, 1832–
1840.

McGuire, L.P., Mayberry, H.W., and Willis, C.K.R. (2017). White-nose syndrome increases
torpid metabolic rate and evaporative water loss in hibernating bats. Am. J. Physiol.-Regul.
Integr. Comp. Physiol. *313*, R680–R686.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res. *20*, 1297–1303.

Messer, P.W., and Petrov, D.A. (2013). Population genomics of rapid adaptation by soft
selective sweeps. Trends Ecol. Evol. *28*, 659–669.

Meteyer, C.U., Barber, D., and Mandl, J.N. (2012). Pathology in euthermic bats with white
nose syndrome suggests a natural manifestation of immune reconstitution inflammatory
syndrome. Virulence *3*, 583–588.

Micalizzi, E.W., Mack, J.N., White, G.P., Avis, T.J., and Smith, M.L. (2017). Microbial

inhibitors of the fungus *Pseudogymnoascus destructans*, the causal agent of white-nose

syndrome in bats. PLOS ONE *12*, e0179770.

Miller, C.T., Beleza, S., Pollen, A.A., Schluter, D., Kittles, R.A., Shriver, M.D., and

Kingsley, D.M. (2007). Cis-regulatory changes in Kit ligand expression and parallel

evolution of pigmentation in sticklebacks and humans. Cell *131*, 1179–1189.

Miller-Butterworth, C.M., Vonhof, M.J., Rosenstern, J., Turner, G.G., and Russell, A.L.

(2014). Genetic structure of little brown bats (*Myotis lucifugus*) corresponds with spread of

white-nose syndrome among hibernacula. J. Hered. esu012.

Minnis, A.M., and Lindner, D.L. (2013). Phylogenetic evaluation of Geomyces and allies

reveals no close relatives of Pseudogymnoascus destructans, comb. nov., in bat hibernacula

of eastern North America. Fungal Biol. *117*, 638–649.

Morales, A.E., and Carstens, B.C. (2018). Evidence that Myotis lucifugus "Subspecies" are

Five Nonsister Species, Despite Gene Flow. Syst. Biol. *67*, 756–769.

Norquay, K.J.O., Martinez-Nuñez, F., Dubois, J.E., Monson, K.M., and Willis, C.K.R.

(2013). Long-distance movements of little brown bats (Myotis lucifugus). J. Mammal. *94*,

506–515.

Olivera-Hyde, M., Silvis, A., Hallerman, E.M., Ford, W.M., and Britzke, E.R. (2019).

Relatedness within and among northern long-eared bat (Myotis septentrionalis) colonies at a

local scale. Can. J. Zool. *97*, 724–735.

Paterson, S., Wilson, K., and Pemberton, J.M. (1998). Major histocompatibility complex

variation associated with juvenile survival and parasite resistance in a large unmanaged

ungulate population (Ovis aries L.). Proc. Natl. Acad. Sci. *95*, 3714–3719.

Paterson, S., Vogwill, T., Buckling, A., Benmayor, R., Spiers, A.J., Thomson, N.R., Quail,

M., Smith, F., Walker, D., Libberton, B., et al. (2010). Antagonistic coevolution accelerates

molecular evolution. Nature *464*, 275–278.

Peters, J.-M., Tedeschi, A., and Schmitz, J. (2008). The cohesin complex and its roles in

chromosome biology. Genes Dev. *22*, 3089–3114.

Puechmaille, S.J., Wibbelt, G., Korn, V., Fuller, H., Forget, F., Mühldorfer, K., Kurth, A.,

Bogdanowicz, W., Borel, C., Bosch, T., et al. (2011). Pan-European distribution of white-

nose syndrome fungus (*Geomyces destructans*) not associated with mass mortality. PLoS

ONE *6*, e19167.

R Development Core Team (2011). R: A language and environment for statistical computing.

Råberg, L., Graham, A.L., and Read, A.F. (2009). Decomposing health: tolerance and

resistance to parasites in animals. Philos. Trans. R. Soc. Lond. B. Biol. Sci. *364*, 37–49.

Reeder, D.M., Frank, C.L., Turner, G.G., Meteyer, C.U., Kurta, A., Britzke, E.R., Vodzak,

M.E., Darling, S.R., Stihler, C.W., Hicks, A.C., et al. (2012). Frequent arousal from

hibernation linked to severity of infection and mortality in bats with white-nose syndrome.

Plos One *7*, e38920.

Ren, P., Haman, K.H., Last, L.A., Rajkumar, S.S., Keel, M.K., and Chaturvedi, V. (2012).

Clonal Spread of Geomyces destructans among Bats, Midwestern and Southern United

States. Emerg. Infect. Dis. *18*, 883–885.

Rico, Y., Morris-Pocock, J., Zigouris, J., Nocera, J.J., and Kyle, C.J. (2015). Lack of Spatial

Immunogenetic Structure among Wolverine (Gulo gulo) Populations Suggestive of Broad

Scale Balancing Selection. PLOS ONE *10*, e0140170.

van Riper, C., van Riper, S.G., Goff, M.L., and Laird, M. (1986). The Epizootiology and

Ecological Significance of Malaria in Hawaiian Land Birds. Ecol. Monogr. *56*, 327–344.

Robinson, S.J., Samuel, M.D., Johnson, C.J., Adams, M., and McKenzie, D.I. (2012).

Emerging prion disease drives host selection in a wildlife population. Ecol. Appl. Publ. Ecol.

Soc. Am. *22*, 1050–1059.

Routtu, J., and Ebert, D. (2015). Genetic architecture of resistance in *Daphnia* hosts against

two species of host-specific parasites. Heredity *114*, 241–248.

Scholthof, K.-B.G. (2007). The disease triangle: pathogens, the environment and society. Nat.

Rev. Microbiol. *5*, 152–156.

Schoville, S.D., Bonin, A., François, O., Lobreaux, S., Melodelima, C., and Manel, S. (2012).

Adaptive Genetic Variation on the Landscape: Methods and Cases. Annu. Rev. Ecol. Evol.

Syst. *43*, 23–43.

Sironi, M., Cagliani, R., Forni, D., and Clerici, M. (2015). Evolutionary insights into host–

pathogen interactions from mammalian sequence data. Nat. Rev. Genet. *16*, 224–236.

Smith, K.F., Sax, D.F., and Lafferty, K.D. (2006). Evidence for the Role of Infectious

Disease in Species Extinction and Endangerment. Conserv. Biol. *20*, 1349–1357.

Sparks, A.M., Watt, K., Sinclair, R., Pilkington, J.G., Pemberton, J.M., McNeilly, T.N.,

Nussey, D.H., and Johnston, S.E. (2019). The genetic architecture of helminth-specific

immune responses in a wild population of Soay sheep (Ovis aries). BioRxiv 628271.

Spurgin, L.G., and Richardson, D.S. (2010). How pathogens drive genetic diversity: MHC,

mechanisms and misunderstandings. Proc. Biol. Sci. *277*, 979–988.

Studier, E.H., and O'Farrell, M.J. (1972). Biology of Myotis thysanodes and M. lucifugus

(Chiroptera: Vespertilionidae). I. Thermoregulation. Comp. Biochem. Physiol. A *41*, 567–

595.

Teacher Amber G. F., Garner Trenton W. J., and Nichols Richard A. (2009). Population

genetic patterns suggest a behavioural change in wild common frogs (Rana temporaria)

following disease outbreaks (Ranavirus). Mol. Ecol. *18*, 3163–3172.

The UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. Nucleic

Acids Res. *47*, D506–D515.

Thomas, D.W., Dorais, M., and Bergeron, J.-M. (1990). Winter Energy Budgets and Cost of

Arousals for Hibernating Little Brown Bats, Myotis lucifugus. J. Mammal. *71*, 475–479.

Tompkins, D.M., Carver, S., Jones, M.E., Krkošek, M., and Skerratt, L.F. (2015). Emerging

infectious diseases of wildlife: a critical perspective. Trends Parasitol. *31*, 149–159.

Turner, G.G., Reeder, D.M., and Coleman, J.T.H. (2011). A five-year assessment of mortality

and geographic spread of white-nose syndrome in North American bats and a look to the

future. Bat Res. News *52*, 13–27.

Verant, M.L., Boyles, J.G., Waldrep, W., Wibbelt, G., and Blehert, D.S. (2012).

Temperature-dependent growth of *Geomyces destructans*, the fungus that causes bat white-

nose syndrome. Plos One *7*, e46280.

Verant, M.L., Meteyer, C.U., Speakman, J.R., Cryan, P.M., Lorch, J.M., and Blehert, D.S.

(2014). White-nose syndrome initiates a cascade of physiologic disturbances in the

hibernating bat host. BMC Physiol. *14*, 10.

Vonhof, M.J., Russell, A.L., and Miller-Butterworth, C.M. (2015). Range-wide genetic

analysis of little brown bat (*Myotis lucifugus*) populations: estimating the risk of spread of

White-nose syndrome. PLoS ONE *10*, e0128713.

Vonhof, M.J., Amelon, S.K., Currie, R.R., and McCracken, G.F. (2016). Genetic structure of

winter populations of the endangered Indiana bat (Myotis sodalis) prior to the white nose

syndrome epidemic: implications for the risk of disease spread. Conserv. Genet. *17*, 1025–

1040.

Voronin, D.A., and Kiseleva, E.V. (2007). [Functional role of proteins containing ankyrin

repeats]. Tsitologiia *49*, 989–999.

Warnecke, L., Turner, J.M., and Geiser, F. (2008). Torpor and basking in a small arid zone

marsupial. Naturwissenschaften *95*, 73–78.

Warnecke, L., Turner, J.M., Bollinger, T.K., Lorch, J.M., Misra, V., Cryan, P.M., Wibbelt,

G., Blehert, D.S., and Willis, C.K.R. (2012). Inoculation of bats with European *Geomyces*

destructans supports the novel pathogen hypothesis for the origin of white-nose syndrome.

Proc. Natl. Acad. Sci. U. S. A. *109*, 6999–7003.

Warnecke, L., Turner, J.M., Bollinger, T.K., Misra, V., Cryan, P.M., Blehert, D.S., Wibbelt,

G., and Willis, C.K.R. (2013). Pathophysiology of white-nose syndrome in bats: a

mechanistic model linking wing damage to mortality. Biol. Lett. *9*, 20130177.

Wibbelt, G., Kurth, A., Hellmann, D., Weishaar, M., Barlow, A., Veith, M., Prueger, J.,

Goerfoel, T., Grosche, L., Bontadina, F., et al. (2010). White-nose syndrome fungus

(*Geomyces destructans*) in Bats, Europe. Emerg. Infect. Dis. *16*, 1237–1243.

Wibbelt, G., Puechmaille, S.J., Ohlendorf, B., Muehldorfer, K., Bosch, T., Goerfoel, T.,

Passior, K., Kurth, A., Lacremans, D., and Forget, F. (2013). Skin lesions in European

hibernating bats associated with *Geomyces destructans*, the etiologic agent of white-nose

syndrome. Plos One *8*, e74105.

Wilder, A.P., Kunz, T.H., and Sorenson, M.D. (2015). Population genetic structure of a

common host predicts the spread of white-nose syndrome, an emerging infectious disease in

bats. Mol. Ecol. *24*, 5495–5506.

Willis, C.K.R., Menzies, A.K., Boyles, J.G., and Wojciechowski, M.S. (2011). Evaporative

water loss is a plausible explanation for mortality of bats from white-nose syndrome. Integr.

Comp. Biol. *51*, 364–373.

Wright, S. (1978). Evolution and the Genetics of Populations, Volume 4. Variability within

and among Natural Populations. (Chicago: University of Chicago Press).

www.whitenosesyndrome.org (2019). White-nose syndrome occurance map - by year (2019).

Data   last   updated: 3/15/2019.   Available   at:

https://www.whitenosesyndrome.org/resources/map.

Yang, Z., Shi, H., Ma, P., Zhao, S., Kong, Q., Bian, T., Gong, C., Zhao, Q., Liu, Y., Qi, X., et

al. (2018). Darwinian Positive Selection on the Pleiotropic Effects of KITLG Explain Skin

Pigmentation and Winter Temperature Adaptation in Eurasians. Mol. Biol. Evol. 1–12.

Zeisset, I., and Beebee, T.J.C. (2014). Drift Rather than Selection Dominates MHC Class II

Allelic Diversity Patterns at the Biogeographical Range Scale in Natterjack Toads Bufo

calamita. PLOS ONE *9*, e100176.

Zhang, T., Victor, T.R., Rajkumar, S.S., Li, X., Okoniewski, J.C., Hicks, A.C., Davis, A.D.,

Broussard, K., LaDeau, S.L., Chaturvedi, S., et al. (2014). Mycobiome of the bat white nose

syndrome affected caves and mines reveals diversity of fungi and local adaptation by the

fungal pathogen *Pseudogymnoascus (Geomyces) destructans*. PLOS ONE *9*, e108714.

Zhang, T., Chaturvedi, V., and Chaturvedi, S. (2015). Novel *Trichoderma polysporum* strain

for the biocontrol of *Pseudogymnoascus destructans*, the fungal etiologic agent of bat white

nose syndrome. PLOS ONE *10*, e0141316.

Zukal, J., Bandouchova, H., Brichta, J., Cmokova, A., Jaron, K.S., Kolarik, M., Kovacova,

V., Kubátová, A., Nováková, A., Orlov, O., et al. (2016). White-nose syndrome without

borders: *Pseudogymnoascus destructans* infection tolerated in Europe and Palearctic Asia but

not in North America. Sci. Rep. *6*, 19829.

Table captions

Table 1. Samples used in study.

Table 2. Summary of individual SNP F_{ST} values across the *M. lucifugus* genome when

comparing Pre-WNS with Post-WNS; and samples from PA and NY (both Pre- and Post-

WNS).

Table 3. Genes within the *M. lucifugus* genome – excluding scaffold GL429776 – with

windows possessing F_{ST} values >= 0.05 when comparing PA and NY samples Post-WNS

Table 4. Genes within scaffold GL429776 with windows possessing F_{ST} values >= 0.05 when

comparing PA and NY samples Post-WNS.

Figure captions

Fig. 1. *M. lucifugus* sampling locations in Pennsylvania (PA) and New York (NY). Pre-

WNS-sites in grey circles and Post-WNS-sites in white circles. 1) Aitkin Cave, Mifflin

County, PA (n=3); 2) Snyder County, Penn, PA (n=19); 3) Lewisburg, Union Co, PA (n=3);

4) Northumberland County, PA (n=2); 5) Howes Cave, County Schoharie, NY (n=24); 6)

Allentown, Lehigh County, PA (n=32); 7) Woodward, Centre County, PA (n=14); 8)

Alexandria, Huntingdon County, PA (n=20); 9) Lackawaxen, Pike County, PA (n=4); 10)

Oley, County Berks, PA (n=12); 11) Frankstown, County Blair, PA (n=12); 12) Williams

Preserve Mine, NY (n=16); 13) Hoopers Mine, NY (n=24); 14) Delmar, Albany County, NY

(n=36); and 15) Willsboro, Essex County, NY (n=21).

Fig. 2. Fixation indices (F_{ST}) across the *M. lucifugus* genome, comparing the population

before and after the arrival of white-nose syndrome. Scaffold lengths presented in green and

blue. Solid red line indicates cut-off for F_{ST} values of 5 standard deviations from the mean

across the whole genome; dotted red line indicates cut-off of F_{ST}=0.05.

Fig 3. Fixation indices (F_{ST}) across the *M. lucifugus* genome between geographically

separated individuals of the study population a) comparing individuals from PA to

individuals from NY before the arrival of WNS; b) comparing individuals from PA before

and after the arrival of WNS; and c) comparing individuals from NY before and after the

arrival of WNS. Solid red line indicates cut-off for F_{ST} values of 5 standard deviations from

the mean across the whole genome; dotted red line indicates cut-off of F_{ST}=0.05.

Fig 4. Fixation indices (F_{ST}) across the *M. lucifugus* genome between geographically

separated individuals of the study population a) comparing individuals from PA to

individuals from NY after the arrival of WNS, with a zoomed plot of scaffold GL429776 and

gene models therein containing windows demonstrating F_{ST} values < 0.05; and b) comparing

individuals from PA to individuals from NY with Pre- and Post-WNS data combined. Solid

red line indicates cut-off for F_{ST} values of 5 standard deviations from the mean across the

whole genome; dotted red line indicates cut-off of F_{ST}=0.05.

Fig 5. Proportions of reference alleles called at individual SNP sites for the region of scaffold

GL429776 demonstrating high F_{ST} values when comparing a) individuals from Pennsylvania

before and after the arrival of WNS; b) individuals from NY before and after the arrival of

WNS; c) individuals from PA to individuals from NY before the arrival of WNS; and d)

individuals from PA to individuals from NY after the arrival of WNS.

Table 1.

Sample age	Classification	Total number of samples	Pennsylvania*	New York*
2006	Pre-WNS	49	25	24
2007	Pre-WNS	3	3	
2015	Post-WNS	28	28	
2016	Post-WNS	139	82	57

* WNS first detected in New York in 2007 and PA in 2009

Table 2.

	Pre- vs post-WNS	Between site (combining pre- and post-WNS)
Median (Interquartile range)	0.0059 (0.0013 – 0.0171)	0.0057 (0.0013 – 0.0165)
Mean (s.d.)	0.0127 (0.0173)	0.0123 (0.0170)

Table 3.

Scaffold	Location range (bp)	Gene ID	Protein Function	Number of overlapping high F_{ST} windows
NW_005871715.1	288334 - 351644	102435296	bifunctional heparan sulfate N-deacetylase/N-sulfotransferase 4-like (partial)	4
NW_005871771.1	26662 - 48938	102441697	E3 ubiquitin-protein ligase TRIM7	1
NW_005871823.1	81353 - 213965	102428352	Ankyrin 2	7
NW_005871905.1	175 - 139344	102428043	calcium/calmodulin-dependent protein kinase type II subunit delta-like	10

| NW_005872026.1 | 72042 - 101269 | 102443179 | N-deacetylase and N-sulfotransferase 4 (partial) | 1 |
| NW_005872062.1 | 33266 - 86607 | 102428048 | zinc finger GRF-type containing 1 | 1 |

Table 4.

Gene ID	F_{ST}	No of windows	No of windows	Start pos.	End pos.	Details

		(per 10kb)				
MGAT4C	0.0612	78	1.08	8999210	9724150	Glycosyltransferase that participates in the transfer of N-acetylglucosamine (GlcNAc) to the core mannose residues of N-linked glycans
CEP290	0.0679	12	1.01	10709418	10827912	Involved in early and late steps in cilia formation
ATP2B1	0.0826	12	0.89	12228178	12362389	The protein encoded by this gene belongs to the family of P-type primary ion transport ATPases characterised by the formation of an aspartyl phosphate intermediate during the reaction cycle. These enzymes remove bivalent calcium ions from eukaryotic cells against very large concentration gradients and play a critical role in intracellular calcium homeostasis.
KITLG	0.0706	10	1.13	11127130	11215731	Plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell

development, migration and function, and in melanogenesis.

POC1B	0.0686	7	0.74	12044678	12139811	Plays an important role in centriole assembly and/or stability and ciliogenesis
TMTC3	0.0872	6	1.00	10827719	10887502	Involved in the positive regulation of proteasomal protein degradation in the endoplasmic reticulum (ER), and the control of ER stress response.
GeneID:111824954	0.0693	5	1.14	11756928	11800690	uncharacterised LOC111824954
CUNH12orf50	0.0588	2	0.67	10663328	10693373	chromosome unknown C12orf50 homolog
GeneID:111824948	0.0816	2	1.13	11375011	11392720	uncharacterised LOC111824948
GeneID:111824949	0.0635	2	1.14	11802704	11820273	uncharacterised LOC111824949

CUNH12orf29	0.0488	1	0.71	10696290	10710421	chromosome unknown C12orf29 homolog
DUSP6	0.0603	1	2.26	11976039	11980455	Inactivates MAP kinases.
GeneID:111824955	0.0459	1	1.57	12101754	12108133	uncharacterised LOC111824955