Short Neuropeptide F regulates the starvation mediated enhanced locomotor Activity in *Drosophila*

Anna Geo[1, a], Himani Pathak[2, a], Anamika Elizabeth Kujur[1] , Sreesha R Sudhakar[2] and Nisha N Kannan [1,b]

[1]Chronobiology Laboratory and [2] Drosophila Energy and Metabolism Laboratory, School of Biology, Indian Institute of Science Education and Research (IISER), Thiruvananthapuram, Kerala 695551, India

Running title: sNPF governs starvation-mediated hyperactivity in fruit flies.

[a]AG and HP contributed equally to this work.
[b]Corresponding author. E-mail address: nishankannan@iisertvm.ac.in

Tel.: +0471-2778045; Fax: + 0471 2597427

Abstract:

The circadian clock regulates various behavioral, metabolic and physiological processes to occur at the most suitable time of the day. Internal energy stores and nutrient availability modulates the most apparent circadian clock mediated locmotor activity rhythm in *Drosophila*. Although previous studies unraveled the role of circadian clock in metabolism and activity rest rhythm, the precise pathway through which the circadian neuropeptidergic signaling regulates internal energy storage and the starvation-mediated increase in activity resembling foraging remains largely unclear. This study was aimed to elucidate the role of circadian neuropeptide, short neuropeptide F (sNPF) in triglyceride metabolism, starvation resistance and starvation-mediated increased locomotor activity in *Drosophila*. The results showed that *snpf* transcripts exhibits significant rhythmicity in wild type flies under 12:12 hour light-dark cycles (LD) and constant darkness (DD) whereas *snpf* transcript level in *period* null flies did not exhibit any significant rhythmicity under LD. Knockdown of sNPF in circadian clock neurons reduced the triglyceride level, starvation resistance and increased the starvation-mediated hyperactivity response after 24 hour of starvation. Further studies showed that knock down of sNPF receptors (sNPFR) expressed in insulin producing cells (IPC) increased the starvation resistance and reduced starvation-induced hyperactivity response after 24 hour of starvation. Collectively, our results suggest that transcriptional oscillation of *snpf* mRNA is endogenously controlled by the circadian clock and elucidate the role of sNPF in modulating locomotor activity in accordance with the nutrient availability in *Drosophila*.

Keywords: circadian, short neuropeptide F, hyperactivity, oscillation, starvation, triglyceride metabolism

Introduction

The circadian clock drives daily rhythms in a wide array of physiological and behavioral processes by scheduling it at the appropriate time of the day. This inherent timekeeping system coordinates the phase of two fundamental behavioral rhythms such as feeding and sleep/wake cycles and it is believed to aid the organism to adapt with the cyclic external environmental changes. The growing body of evidence indicates that the circadian clock not only mediates optimal phasing of behavioral rhythmicity during 24 hours of a day but also intimately ties with metabolism (Eckel-Mahan and Sassone-Corsi, 2013). While the clock orchestrates numerous metabolic pathways, the nutrient availability and metabolic status can in turn feed back to impinge on the functioning of the circadian clock (Damiola et al., 2000; Nakahata et al., 2009; Vollmers et al., 2009; Xu et al., 2011). A large body of evidence accumulated over the past two decades remarkably improved our understanding of the essential role of the clock in metabolism, feeding behavior, and the sleep wake cycle (Maury et al., 2010). However, little is known about the precise underlying pathways by which the circadian clock aids the organism to attain temporal harmonics between the sleep wake cycle and feeding behavior.

The association between the circadian clock, sleep wake cycle and metabolism was further substantiated by the findings that the circadian timing system drives rhythmicity of various peptides involved in metabolism that also play key role in the regulation of sleep wake cycles (Reviewed in Kalsbeek and Fliers, 2013). For instance, peptides involved in metabolism such as leptin, ghrelin and cholecystokinin display circadian rhythmicity in expression (Ahima et al., 1998; Kalra et al., 2003; Kalsbeek et al., 2001; Schade et al., 1993). Another appetite stimulating hormone orexin promotes wakefulness while stimulated by ghrelin and enhances sleep when inhibited by leptin (Sakurai, 2007) and orexigenic neuron activation is believed to be

under circadian control (Marston et al., 2008). Short neuropeptide F (sNPF) signaling plays a

role in feeding and sleep (Lee et al., 2004; Shang et al., 2013). sNPF receptor (sNPFR) signaling

in Insulin Producing Cells (IPCs) regulate the expression of Drosophila Insulin like peptides

(DILPS) and metabolism in Drosophila (Lee et al., 2008; Fadda et al., 2019). sNPF is expressed

in circadian clock small ventral lateral nurons (s-LNvs) and dorsal lateral neurons (LNds)

regulates sleep in *Drosophila* (Shang et al., 2013).

Feeding and sleep wake cycles are interconnected behaviors and the optimum balance

between them is disrupted under nutrient deprivation. Starvation suppresses sleep by promoting

hyperactivity in *Drosophila* and this behavioral change in response to nutrient unavailability is

governed by diverse neuropeptides such as allatostatin, octopamine, adipokinetic hormone,

glucagon and insulin (Chen et al., 2016; G. Lee and Park, 2004; Yang et al., 2015; Yu et al.,

2016). Sustained increase in locomotor activity under starvation condition partly resembles

foraging behavior (Yang et al., 2015). Feeding behavior appears to be under multiple controls

including the gustatory, olfactory and circadian neural circuits (Xu et al., 2008; Lin et al., 2019).

Significant progress has been made in understanding the circadian and neuronal basis of feeding

rhythm and starvation-mediated sleep changes in *Drosophila* (Xu et al., 2008; Keene et al.,

2010). However it remains unclear whether the circadian clock conveys the time of the day

information to coherently coordinate this starvation-mediated behavioral change with

metabolism through interlinked neuronal pathways or via independent neural circuitry.

In the present study, we aimed to elucidate the role of Short neuropeptide F in

triglyceride metabolism, starvation-mediated enhanced activity and starvation resistance in

Drosophila. The results showed that the circadian clock regulates the transcription of *snpf* gene.

sNPF expressed in circadian clock neurons delays the starvation-mediated hyperactivity

response, whereas sNPF receptor expressed in IPCs advances the starvation-mediated

hyperactivity response in *Drosophila.* sNPF expressed in clock neurons enhance the triglyceride

level and starvation resistance whereas SNP receptors expressed in the IPCs reduce the

starvation resistance suggesting the differential role of neuropeptide sNPF in clock neurons and

IPCs to regulate the starvation mediated locomotor activity in accordance with the nutrient

availability in *Drosophila*.

Materials and Methods

Fly stocks and maintenance

99   Following fly lines were used in this experiment: w^{1118} (BDSC #5905)*, tim-Gal4 and per^o*

100   (SheebaVasu Laboratory)*, dilp2-Gal4, UAS-snpf* ^{RNAi} and UAS-*snpfR* ^{RNAi} (Kweon Yu

101   Laboratory). All fly stocks were maintained in the *Drosophila* growth chamber (MIR-154,

102   Panasonic, Japan) at 25º C temperature, 75 ± 5% humidity and 12:12 hour (h) light-dark (LD)

103   cycle where lights came on at Zeitgeber Time 00 (ZT 00) and went off at ZT12.

Measurement of mRNA levels

To assess the mRNA transcription profiles of *snpf*, three replicates were used for each

time point. Each replicate contained 30 heads from 2-3 day old w^{1118} male flies. Total RNA was

extracted using QIAGEN RNAeasy Plus Mini kit; DNAase digestion was performed using

QIAGEN RNase–free DNase. cDNA was synthesized using SuperScript™ III First-Strand

Synthesis System (Invitrogen, Cat. No. 18080051) and real-time PCR was performed using Bio-

Rad CFX96TM with the cDNA template, Power SYBR® Green PCR Master Mix (Cat. #

4368702) from ThermoFisher Scientific to check temporal transcription profile of *snpf*. mRNA

levels were measured under LD cycle and constant darkness (DD). The molecular oscillation of

mRNA levels was determined by normalizing the mRNA level of the gene of interest with the

mRNA level of *rp49* at each time point. Knockdown efficiency of *tim-Gal4 > snpf* [RNAi] was

validated by qPCR. We used the following primers for qPCR. *snpf* forward primer 5'-

CCCGAAAACTTTTAGACTCA-3' and reverse primer 5'- TTTTCAAACATTTCCATCG -3´,

rp49 forward primer 5'- GCTAAGCTGTCGCACAAA-3' and reverse primer 5'-

TCCGGTGGGCAGCATGTG-3'.

Locomotor Activity assay

For activity/rest assay, freshly emerged male flies were pre-entrained for two days in the

LD cycle at 25 ^{o}C and a humidity of $75 \pm 5\%$ inside the incubator (MIR-154, Panasonic, Japan).

Two-day-old virgin males were individually introduced into activity tubes. Flies were provided

with standard cornmeal medium and their activity/rest behavior was recorded using Drosophila

Activity Monitors (DAM, Trikinetics, USA). The infrared beam that passes through the middle

of the activity tubes count the movement of the flies. Following acclimatization in activity tubes

for 12 hours, flies were transferred to 1% agar medium or to standard cornmeal medium in glass

tubes at ZT12 (during the onset of dark phase) to provide starved or fed conditions, respectively,

and locomotor activity was recorded for 36 hours at 25 ^{o}C. Each experiment was performed with

27-32 flies for each genotype and DAM readings were collected in every 1-minute interval and

the activity counts/ hour is plotted in the locomotor activity waveform. Flies died within 24 hour

of starvation were excluded and flies survived for 24-36 hours of starvation were used for the

analysis. Change in activity under starvation was calculated as [(Activity under starvation-

Activity under fed) /Activity under fed condition] x 100. Separate sets of flies were used for

recording the locomotor activity under fed and starved flies condition. Hence the change in

activity under starvation condition was calculated as [(Activity of individual fly under

starvation-Average activity of all the flies under fed) / Average activity of all the flies under fed]

x 100.

Triglyceride assay

This assay comprised of three replicates and each replicate containing five male flies

under fed condition was homogenized in homogenization buffer (0.05% Tween 20). The

homogenate was incubated at $70^{o}C$ for 5 minutes to inactivate the enzymes and centrifuged at

14000 rpm for 3 minutes and the supernatant was collected into a sample tube. Sigma serum

triglyceride determination kit TR0100 was used to estimate the triglyceride level in the sample

by measuring absorbance at 540 nm using TECAN Infinite M200 pro-multimode plate reader.

The triglyceride content was normalized to the total protein content of the flies estimated by

Quick Start™ Bradford 1X Dye Reagent (BioRad, Cat. # 500-0205). Freshly emerged flies feed

less and possess larval triglyceride energy storage. Larval fat cells persist only up to about two

days and are replaced by adult fat cells. Hence assays were carried out at ZT 12 in fed two-day-

old male flies to estimate the triglyceride levels in adult flies.

Starvation resistance Assay

151   Starvation resistance assay comprised of six replicates with each replicate vial containing

152   10-15 flies for each genotype. Five-day-old male flies were kept in the 1% agar medium vials.

153   The number of dead flies was recorded at every two-hour interval until all the flies died.

Statistical analysis

Cosinor analysis was implemented in MATLAB-R2016a to test for rhythmicity, to estimate the

peak phase and the amplitude (peak/trough ratio) of relative mRNA expression of *snpf* (Nelson et

al., 1979). The waveform of activity/rest rhythm, day time activity, nighttime activity and

change in locomotor activity were analyzed by using ANOVA followed by Tukey's post hoc

multiple comparisons. Log-rank test was used to compare and analyze the survivorship curves

under starvation. TGA levels was analyzed by Student's t-test. The statistical analyses were

implemented on Statistica for Windows Release 5.0 B (Statsoft, 1995). Error bars in all the

graphs represent standard error of the mean (SEM).

Results

snpf **transcript exhibit circadian oscillation**

To assess whether the circadian clock regulates *snpf* transcript oscillation, the

transcription profile of *snpf* was estimated in w^{1118} flies under LD and under constant darkness

(DD). The *snpf* mRNA level increased during light phase and decreased during the dark phase

resulting in a significant daily oscillation with a peak at ZT 8.1h under LD (Cosinor analysis,

p<0.001) (Fig. 1A). The *snpf* transcriptional oscillation persisted under DD with a peak at CT

8.5h (Cosinor analysis, *p*<0.002). However, the amplitude (peak/trough ratio) of the oscillation

decreased under DD compared to that obtained under LD (Fig. 1B). To confirm the role of

circadian clock in *snpf* transcript oscillation, further studies were carried out in *per^o* flies under

LD. *snpf* transcript level in *per^o* flies did not exhibit any significant rhythmicity under LD

(cosinor analysis, *p*> 0.05) (Fig. 1C). These results suggest that the transcriptional oscillation of

snpf mRNA is endogenously controlled by the clock.

sNPF expressed in circadian clock neurons suppress starvation-mediated hyperactivity

To assess the role of sNPF expressing circadian clock neurons in starvation-mediated

hyperactivity, we expressed *UAS-snpf ^{RNAi}* under the *tim-Gal4* driver (*tim-Gal4 > snpf ^{RNAi}*) and

recorded the locomotor activity under fed and starved LD conditions for 36 h. *tim-Gal4 > w^{1118},*

w ^{1118} > UAS-snpf ^{RNAi} control flies and the *tim-Gal4 > snpf ^{RNAi}* flies exhibited increased activity

under starved condition compared to what was observed under fed condition (*tim-Gal4 > w^{1118}*

flies; ANOVA, $p<0.0001$; $w^{1118} >$ *UAS-snpf* ^{RNAi}; ANOVA, $p<0.0001$; *tim-Gal4* $>$ *snpf* ^{RNAi} flies;

ANOVA, $p<0.0001$) (Fig. 2A- C). ANOVA followed by post-hoc multiple comparisons using

Tukey's test on activity data showed that starved *tim-Gal4* $>w^{1118}$ flies enhanced the activity at

25-27h of starvation (Fig. 2A) resulting in a significant increase in the total activity second dark

phase (25-36h) (ANOVA, $p<0.0001$) and during the light phase (13-24h) (ANOVA, $p<0.0001$)

compared to those observed under fed condition (Fig. E-F). $w^{1118} >$ *snpf* ^{RNAi} flies enhanced the

activity at 1-2h, 16-19h, 25-27h of and 32h starvation. This resulted in a significant increase in

the total activity of $w^{1118} >$ *snpf* ^{RNAi} flies during the first dark phase (1-12h) (ANOVA,

$p<0.0001$), light phase (13-24h) (ANOVA, $p<0.0001$) and second dark phase (25-36h)

(ANOVA, $p<0.0001$) under starvation compared to those observed under fed condition (Fig. 2B,

D-F). *tim-Gal4* $>$ *snpf* ^{RNAi} flies exhibited increased activity at 15h, 25-31h and 33h of starvation

resulting in a significant increase in the total activity during the light phase (13-24h) (ANOVA,

$p<0.0001$) and second dark phase (25-36h) (ANOVA, $p<0.0001$) under starvation compared to

those observed under fed condition (Fig. 2C, E-F). Comparison of change in activity between

controls and *tim-Gal4* $>$ *snpf* ^{RNAi} flies under starved vs fed condition revealed that *tim-Gal4* $>$

snpf ^{RNAi} flies exhibited significant increase in activity after 24 hours of starvation (25-36h) than

tim-Gal4 $> w^{1118}$ (ANOVA, $p<0.0001$) and $w^{1118} >$ *snpf* ^{RNAi} (ANOVA, $p<0.0001$) controls (Fig.

2G). These results suggest that sNPF in circadian clock neurons delays the starvation-mediated

increased locomotor activity response after 24h of starvation in *Drosophila*.

We further assessed the role of sNPF in regulating the triglyceride levels and starvation

resistance in flies expressing *UAS- snpf* ^{RNAi} under the *tim-Gal4* driver. *tim-Gal4* $>$ *snpf* ^{RNAi} flies

exhibited a significant decrease in the triglyceride level compared to both *tim-Gal4* $> w^{1118}$ (t-

test, $p<0.0001$) and $w^{1118} >$ *snpf* ^{RNAi} controls (t-test, $p<0.0001$) (Fig. 2H). In agreement with the

enhanced triglyceride level, *tim-Gal4 > snpf* ^{RNAi} flies exhibited a significant decrease in the

starvation resistance compared to both *tim-Gal4 > w*^{1118} (log-rank test , $p<0.0001$) and *w*^{1118} >

snpf ^{RNAi} controls (log-rank test , $p<0.0001$) (Fig. S1A). These results suggest that knock down

of *snpf* in circadian clock neurons reduces the triglyceride level and starvation resistance in

Drosophila.

sNPFR expressed in IPC promotes starvation-mediated hyperactivity

211   To test the role of sNPFR expressed in IPCs on starvation-mediated hyperactivity,

we expressed the *UAS-snpfR* ^{RNAi} under the *dilp2-Gal4* driver (*dilp2-Gal4 > snpfR* ^{RNAi}) and

recorded the locomotor activity of these flies under fed and starved condition for 36 h. *dilp2-*

Gal4 > w^{1118}, *w*^{1118} > *snpfR* ^{RNAi} control flies and the *dilp2-Gal4 > snpfR* ^{RNAi} flies exhibited

increased activity under starved condition compared to what was observed under fed condition

(*dilp2-Gal4 > w*^{1118} flies; ANOVA, $p<0.0001$; *w*^{1118} > *snpfR* ^{RNAi}; ANOVA, $p<0.0001$; *dilp2-*

Gal4 > snpfR ^{RNAi} flies; ANOVA, $p<0.0001$) (Fig. 3A- C). ANOVA followed by post-hoc

multiple comparisons using Tukey's test on activity data showed that starved *dlip2-Gal4 > w*^{1118}

flies enhanced the activity at 1h, 13-17h, 23-27h and 30-32h of starvation resulting in a

significant increase in the total activity during the first dark phase (1-12h) (ANOVA, $p<0.005$),

light phase (13-24h) (ANOVA, $p<0.0001$) and second dark phase (25-36h) (ANOVA, $p<0.0001$)

compared to those observed under fed condition (Fig. 3A, D-F). *w*^{1118} > *snpfR* ^{RNAi} flies enhanced

the activity at 2h, 16-17h, 25-27h and 29-31h of starvation. This resulted in a significant

increase in the total activity of *w*^{1118} > *snpfR* ^{RNAi} flies during the light phase (13-24h) (ANOVA,

$p<0.0001$) and second dark phase (25-36h) (ANOVA, $p<0.0001$) compared to those observed

under fed condition (Fig. 3B, E-F). *dilp2-Gal4 > snpfR* ^{RNAi} flies exhibited increased activity at

14-16h and 23- 28h of starvation resulting in a significant increase in the total activity only

during second dark phase (25-36h) (ANOVA, $p<0.001$) and under starvation compared to those

observed under fed condition (Fig. 3C, F). Starvation did not enhance the activity in

experimental flies ($dilp2\text{-}Gal4 > snpfR\ ^{RNAi}$) during the first 24 hours of starvation compared to

fed flies (Fig. 3C-E). Comparison of change in activity between controls and $dilp2\text{-}Gal4 >snpfR$

^{RNAi} flies under starved vs fed condition revealed that $dilp2\text{-}Gal4 > snpfR\ ^{RNAi}$ flies exhibited

decrease in activity than controls after 24 hours of starvation (25-36h) than $dilp2\text{-}Gal4 > w^{1118}$

(ANOVA, $p<0.0001$) and $w^{1118} > snpfR\ ^{RNAi}$ (ANOVA, $p<0.03$) control flies (Fig. 3G). These

results suggest knock down of the $snpfR$ expressed in IPCs delays the starvation-mediated

hyperactivity in Drosophila.

We also assessed whether $snpf$R expressed in the IPCs regulate the triglyceride levels

and starvation resistance in flies by expressing $UAS\text{-}snpfR\ ^{RNAi}$ under the $dilp2\text{-}Gal4$ driver.

$dilp2\text{-}Gal4 > snpfR\ ^{RNAi}$ flies did not exhibit any significant difference in the triglyceride level

compared to both $dilp2\text{-}Gal4 > w^{1118}$ (t-test, $p>0.05$) and $w^{1118} > snpfR\ ^{RNAi}$ controls (t-test,

$p>0.05$) (Fig. 3H). However, $dilp2\text{-}Gal4 > snpfR\ ^{RNAi}$ flies exhibited a significant increase in the

starvation resistance compared to both $dilp2\text{-}Gal4 > w^{1118}$ (log-rank test , $p<0.0001$) and $w^{1118} >$

$snpfR\ ^{RNAi}$ controls (log-rank test , $p<0.0001$) (Fig. S1B). These results suggest that knock down

of $snpfR$ in IPCs increases the starvation resistance in $Drosophila$.

Discussion

Behavior and metabolism are functionally intertwined processes that modulate each

other. However, it remains elusive how the circadian clock governs the major clock mediated

behavioral output locomotor activity rhythm, in synchrony with the metabolic status of an

organism. In the present study, we aimed to examine the specific role of circadian neuropeptide,

i.e., sNPF in lipid metabolism and starvation-mediated behavioral changes in *Drosophila*. The

results of the present study showed that *snpf* transcript exhibited a circadian oscillation in the fly

head. sNPF protein is expressed in a diverse subset of neurons in the adult fly brain including

olfactory receptor neurons, various small interneurons, kenyon cells of the mushroom bodies and

in the circadian clock neurons (Nässel., 2008). Among the ~150 clock neurons clustered in the

fly brain, two dorsal lateral neurons (LNds) and PDF expressing small ventral lateral neurons (s-

LNvs) express sNPF (Johard et al., 2009). Based on the previous findings, mRNA cycling is

essential for the circadian clock neuron function including the neuronal firing or electrical

excitability (Kula-Eversole et al., 2010; Shang et al., 2013). Hence it is quite likely that the *snpf*

transcript oscillation is more apparent in circadian clock neurons than other sNPF expressing

diverse neuronal subsets. Our results showed that *snpf* transcript level did not exhibit any

significant rhythmicity *period* null mutants confirming the endogenous regulation of *snpf*

transcript oscillation. Most of the previously identified cycling head transcripts are tissue or cell-

specific in the mammalian system (Storch et al., 2002). Ubiquitous mRNA cycling is relatively

rare. Further studies are required to confirm whether sNPF oscillation is indeed restricted to

clock neurons or if it also oscillates in non-clock neurons.

In agreement with the previous studies, we showed that starvation enhances locomotor

activity in *Drosophila* (Lee and Park 2015; Yang et al., 2015). Given that starvation-induced

hyperactivity and food intake are mutually inhibitory (Chen et al., 2015; Yang et al., 2015), it is

of great interest to understand how these two behaviors are dynamically regulated by the central

and peripheral neural circuits. sNPF expression in a large population of diverse neurons

indicates the multiple discrete functions of this neuropeptide (Nässel., 2008). So far studies

demonstrated that sNPF regulates feeding and its expression in the fan shaped body might be involved in fine-tuning the locomotor activity level (Lee et al., 2004; Kahsai et al., 2010). Expression of sNPF and sNPF receptor in the mushroom body and clock neurons regulate sleep in *Drosophila* (Chen et al., 2013; Shang et al., 2013). sNPF released from s-LNvs and LNds governs the Dorsal neuron 1 (DN1) late night Ca^{2+} activation and the Ca^{2+} dynamics in DN1 neurons are intricately linked to behavioural rhythm (Liang et al., 2017). Downregulation of sNPF in all pacemaker neurons increased nighttime locomotor activity under fed condition (Liang et al., 2017). The results of our present studies showed that knockdown of *snpf* in circadian clock neurons advances the starvation-mediated hyperactivity during the second dark phase, suggesting the role of sNPF expressing clock neurons in delaying hyperactivity under starvation. In the present study starvation condition was provided at the onset of dark phase (ZT 12) and flies expressing *snpf* RNAi in clock neurons exhibited enhanced starvation mediated hyperactivity after 24 hours of starvation during the second dark phase. sNPF is a sleep promoting molecule (Shang et al., 2013) and it regulates the Ca^{2+} activation during the nighttime in DN1 neurons (Liang et al., 2017) that possibly modulates the starvation-induced hyperactivity during the darkness. In the present study starvation condition was provided at the onset of dark phase (ZT 12), further studies are required on *tim-Gal4 > snpf* ^{RNAi} flies with the starvation starting at the onset of light phase (ZT00) to elucidate whether increased starvation-mediated hyperactivity observed after 24hr of starvation is restricted to the second dark phase or it would persist even under light phase after 24hr of starvation. A population of dorsally located *clk-*expressing neurons has been implicated in promoting sleep during starvation (Keene et al., 2010). Among the clock neurons, PDF positive morning cell s-LNvs and PDF negative evening cells LNds express sNPF (Johard et al., 2009). Morning cell specific downregulation of sNPF

impaired morning anticipatory activity under fed condition. sNPF from both M cells and E cells

are required for the Ca2+ rhythmicity, its phase and behavioral output (Liang et al., 2017).

Further studies are required to confirm the specific role of sNPF expressing PDF positive s-LNvs

and PDF negative LNds in starvation-mediated hyperactivity.

Although sNPF predominantly acts as an inhibitory modulator in neuronal networks

(Vecsey et al., 2014) sNPFR expressed in olfactory receptor neurons has excitatory effect and

promotes starvation induced food search behavior (Root et al., 2011). Knockdown of *snpfR* in

IPCs in an adult stage-specific manner does not affect the starvation-mediated sleep loss in

Drosophila (Shang et al., 2013), whereas the results of our present study showed that constitutive

knockdown of *snpfR* in IPCs suppress starvation-mediated hyperactivity after 24 hours of

starvation. While sNPFR expressed in IPCs promote starvation-mediated hyperactivity, sNPF

expressed in circadian clock neurons had an inhibitory effect on this starvation mediated

behavioral phenotype.

Knock down of sNPF in sNPF producing dorso lateral peptidergic (DLP) neurons

increased the starvation resistance and triglyceride levels in *Drosophila* (Kapan et al., 2012). In

contrast, we observed that $tim\text{-}Gal4 > snpf^{RNAi}$ flies exhibit decrease in the triglyceride level and

starvation resistance whereas knock down of *snpfR* expressed in IPCs enhanced only the

starvation resistance in *Drosophila* indicating the differential function of *snpf* expressing clock

neurons and *snpR* expressing IPCs in regulating the starvation resistance.

Taken together, our present study showed that transcriptional oscillation of *snpf*

mRNA is endogenously controlled by the circadian clock and sNPF expressed in circadian clock

neuron delays the starvation-mediated hyperactivity response. sNPF might promote starvation

induced locomotor activity partially through sNPF receptors expressed in the IPCs. The sNPF

appears to be a central neuropeptidergic pathway in regulating starvation mediated hyperactivity

, triglyceride metabolism and starvation resistance in *Drosophila*.

Acknowledgements

We thank Sabira A for the experimental assistance. We thank Jishy Varghese for the valuable

suggestions and for providing the infrastructure facility to carry out some of the experiments,

Kenji Tomioka for the valuable suggestions, SheebaVasu for providing the fly lines and Nikhil K

L for helping with the statistical analysis.

Declaration of competing interest

This work was supported by the Wellcome Trust/DBT India Alliance Fellowship

[IA/E/15/1/502329] awarded to NNK and the intramural fund from Indian Institute of Science

Education and Research, Thiruvananthapuram. The authors report no conflicts of interest. The

authors alone are responsible for the content and writing of the paper.

Author Contributions

Conceived and designed the experiments: NNK AG AEK HP. Performed the experiments: AG

AEK HP SRS. Analyzed the data: AG AEK HP. Contributed reagents/materials: NNK. Wrote

the paper: AG NNK.

Reference

Ahima, R.S., Prabakaran, D. and Flier, J.S. (1998). Postnatal leptin surge and regulation of

circadian rhythm of leptin by feeding. Implications for energy homeostasis and

neuroendocrine function. *J. Clin. Investig.* **101(5)**, 1020-1027. doi:10.1172/JCI1176.

Chen, J., Reiher. W., Hermann-Luibl, C., Sellami, A., Cognigni, P. and Kondo, S, et al.

(2016). Allatostatin A Signalling in *Drosophila* Regulates Feeding and Sleep and Is

Modulated   by   PDF.   *PLOS   Genet*.   **12(9)**,   e1006346.

doi:10.1371/journal.pgen.1006346.

Chen, W., Shi, W., Li, L., Zheng, Z., Li, T. and Bai, W. et al. (2013). Regulation of sleep by

the short neuropeptide F (sNPF) in *Drosophila melanogaster*. *Insect. Biochem. Molec*.

43(9), 809-819. doi:10.1016/j.ibmb.2013.06.003.

Chen, Y., Lin, Y.C., Kuo, T.W. and Knight, Z. A. (2015). Sensory detection of food rapidly

modulates arcuate feeding circuits. *Cell*. **160 (5)**, 829-841. doi:

10.1016/j.cell.2015.01.033.

Damiola, F., Le Minh, N., Preitner, N., Kornmann, B., Fleury-Olela, F. and Schibler, U.

(2000). Restricted feeding uncouples circadian oscillators in peripheral tissues from

the central pacemaker in the suprachiasmatic nucleus. *Genes and Dev*. **14(23)**, 2950-

2961. doi: 10.1101/gad.183500

Eckel-Mahan K.L. and Sassone-Corsi, P. (2013). Metabolism and the circadian clock

converge. *Physiol. Rev*. **93(1)**, 107-135. doi:10.1152/physrev.00016.2012.

Fadda, M., Hasakiogullari, I., Temmerman. L., Beets, I., Zels, S. and Schoofs, L. (2019).

Regulation of Feeding and Metabolism by Neuropeptide F and Short Neuropeptide F

in Invertebrates. *Front. Endocrinol*. **19**, 10:64. doi:10.3389/fendo.2019.00064

Johard, H.A., Yoishii. T., Dircksen. H., Cusumano, P., Rouyer, F., Helfrich-Forster, C. and

Nässel, D.R. (2009). Peptidergic clock neurons in *Drosophila*: ion transport peptide

and short neuropeptide F in subsets of dorsal and ventral lateral neurons. *J. Comp.*

Neurol. **516(1**), 59-73. doi:10.1002/cne.22099.

Kahsai, L., Martin, J.R. and Winther, A.M. (2010). Neuropeptides in the *Drosophila* central

complex in modulation of locomotor behavior. *J. Exp. Biol*. **213**, 2256-2265.

doi:10.1242/jeb.043190.

Kalra, S.P., Bagnasco, M., Otukonyong, E.E., Dube, M.G. and Kalra, P. S. (2003).

Rhythmic, reciprocal ghrelin and leptin signaling: new insight in the development of

obesity. *Regul. Pept*, **111(1-3)**, 1-11. doi.10.1016/S0167-0115(02)00305-1.

Kalsbeek, A., Fliers, E., Romijn, J.A., La Fleur, S.E., Wortel, J., Bakker., et al. (2001). The

suprachiasmatic nucleus generates the diurnal changes in plasma leptin levels.

Endocrinology. **142(6)**, 2677-2685. doi:10.1210/endo.142.6.8197.

Kalsbeek, A. and Fliers, E. (2013). Daily regulation of hormone profiles. *Handb. Exp.*

Pharmacol, **217**, 185-226. doi:10.1007/978-3-642-25950-0_8.

Kapan, N., Lushchak, O. V., Luo, J. and Nässel, D.R. (2012). Identified peptidergic neurons

in the Drosophila brain regulate insulin-producing cells, stress responses and

metabolism by coexpressed short neuropeptide F and corazonin. *Cell. Mol. life sci*. **69**

(**23)**, 4051-66. doi: 10.1007/s00018-012-1097-z.

Keene, A.C., Duboue, E.R., McDonald, D.M., Dus, M., Suh, G.S., Waddell, S., et al. (2010).

Clock and cycle limit starvation-induced sleep loss in *Drosophila*. *Curr. Biol*. **20(13)**,

1209-1215. doi:10.1016/j.cub.2010.05.029.

Kula-Eversole, E., Nagoshi, E., Shang, Y., Rodriguez, J., Allada, R. and Rosbash, M.
(2010). Surprising gene expression patterns within and between PDF-containing
circadian neurons in *Drosophila*. *PNAS*, **107(30)**, 13497-13502.
doi:10.1073/pnas.1002081107.

Lee, G. and Park, J.H. (2004). Hemolymph sugar homeostasis and starvation-induced
hyperactivity affected by genetic manipulations of the adipokinetic hormone-encoding
gene in *Drosophila melanogaster*. *Genetics*, **167(1)**, 311-323.
doi:10.1534/genetics.167.1.311.

Lee, K.S., You, K.H., Choo, J.K., Han, Y.M. and Yu, K. (2004). *Drosophila* short
neuropeptide F regulates food intake and body size. *J. Biol. Chemistry.* **279(49)**,
50781-50789. doi:10.1074/jbc.M407842200.

Lee, K.S., Kwon, O.Y., Lee, J.H., Kwon, K., Min, K. J., Jung, S. A. et al. (2008). *Drosophila*
short neuropeptide F signalling regulates growth by ERK-mediated insulin
signalling. *Nat. Cell Biol.* **10**, 468–475. DOI:10.1038/ncb1710.

Liang, X., Holy, T.E. and Taghert, P.H. (2017). A Series of Suppressive Signals within the
Drosophila Circadian Neural Circuit Generates Sequential Daily Outputs. *Neuron.* **21,**
94(6), 1173-1189.e4. doi: 10.1016/j.neuron.2017.05.007.

Lin, S., Senapati, B. and Tsao, C.H. (2019). Neural basis of hunger-driven behaviour in
Drosophila. *Open Biol.* **9(3)**, 180259. doi: 10.1098/rsob.180259.

Marston, O.J., Williams, R.H., Canal, M.M., Samuels, R.E., Upton. N. and Piggins, H.D.
(2008). Circadian and dark-pulse activation of orexin/hypocretin neurons. *Mol. Brain*.
1,19. doi:10.1186/1756-6606-1-19.

Maury, E., Ramsey, K.M. and Bass, J. (2010). Sleep, Circadian Rhythms and Metabolism. In:

Ahima RS, Editor. Metabolic Basis of Obesity. New York: Springer, p. 229-255.

Nakahata, Y., Sahar, S., Astarita, G., Kaluzova, M. and Sassone-Corsi, P. (2009). Circadian

control of the NAD+ salvage pathway by CLOCK-SIRT1. *Science*. **324(5927)**, 654-

657. doi:10.1126/science.1170803.

Nässel, D.R., Enell, L.E., Santos, J.G., Wegener, C. and Johard, H.A. (2008). A large

population of diverse neurons in the *Drosophila* central nervous system expresses

short neuropeptide F, suggesting multiple distributed peptide functions. *BMC*

Neurosci. **9**, 90. doi:10.1186/1471-2202-9-90.

Nelson, W., Tong, Y.L., Lee, J. K. and Halberg, F. (1979). Methods for cosinor-

rhythmometry. *Chronobiologia*. **6**, 305-323.

Root, C. M., Ko, K.I., Jafari, A. and Wang, J.W. (2011). Presynaptic facilitation by

neuropeptide signaling mediates odor-driven food search. *Cell*, **145**, 133-144. doi:

10.1016/j.cell.2011.02.008.

Sakurai, T. (2007). The neural circuit of orexin (hypocretin): maintaining sleep and

wakefulness. *Nat. Rev. Neurosci*. **8(3)**, 171-181. doi:10.1038/nrn2092.

Schade, R., Vick, K., Sohr, R., Ott, T., Pfister, C., Bellach, J. et al. (1993). Correlative

circadian rhythms of cholecystokinin and dopamine content in nucleus accumbens and

striatum of rat brain. *Behav. Brain Res*. **59(1-2)**, 211-214. doi: 10.1016/0166-

4328(93)90168-p

Shang, Y., Donelson, N.C., Vecsey, C. G., Guo, F., Rosbash. M. and Griffith, L. C. (2013).

Short neuropeptide F is a sleep-promoting inhibitory modulator. *Neuron*. **80(1)**, 171-

183. doi:10.1016/j.neuron.2013.07.029.

StatSoft. Statistica Vol. 1: General Conventions and Statistics 1. 1995, StatSoft Inc,Tulsa, OK.

Storch, K.F., Lipan, O., Leykin, I., Viswanathan, N., Davis, F.C., Wong, W.H. et al. (2002).

Extensive and divergent circadian gene expression in liver and heart. *Nature*.

417(6884), 78-83. doi:10.1038/nature744.

Vecsey, C.G., Pirez, N. and Griffith, L.C. (2014). The *Drosophila* neuropeptides PDF and

sNPF have opposing electrophysiological and molecular effects on central neurons. *J.*

Neurophysiol. **111(5)**: 1033-1045. doi:10.1152/jn.00712.2013.

Vollmers, C., Gill, S., DiTacchio, L., Pulivarthy, S.R., Le, H.D. and Panda, S. (2009). Time

of feeding and the intrinsic circadian clock drive rhythms in hepatic gene expression.

PNAS. **106(50)**, 21453-21458. doi:10.1073/pnas.0909591106.

Xu, K., Zheng, X. and Sehgal, A. (2008). Regulation of feeding and metabolism by neuronal

and peripheral clocks in *Drosophila*. *Cell Metab*. **8(4)**, 289-300.

doi:10.1016/j.cmet.2008.09.006.

Xu. K., DiAngelo, J.R., Hughes, M.E., Hogenesch, J.B. and Sehgal, A. (2011). The circadian

clock interacts with metabolic physiology to influence reproductive fitness. *Cell*

Metab. **13(6)**, 639-654. doi:10.1016/j.cmet.2011.05.001.

Yang, Z., Yu, Y., Zhang, V., Tian, Y., Qi, W. and Wang, L. (2015). Octopamine mediates

starvation-induced hyperactivity in adult *Drosophila*. *PNAS*, **112(16)**: 5219-5224.

doi:10.1073/pnas.1417838112.

Yu, Y., Huang, R., Ye, J., Zhang, V., Wu, C., Cheng, G. et al. (2016). Regulation of

starvation-induced hyperactivity by insulin and glucagon signaling in adult

Drosophila. *Elife*, **5**. doi:10.7554/eLife.15693.

Figure 1. Transcription of sNPF mRNA is clock controlled

(A) Transcription profile of *snpf* mRNA in w^{1118} flies under 12:12 h light- dark cycle (LD). White and grey shaded bars represent light and dark phase, respectively, under LD. (Cosinor analysis, $p<0.001$) (B) Transcription profile of *snpf* mRNA in w^{1118} flies constant darkness (DD). *snpf* transcript exhibited a significant daily oscillation with a peak at ZT 8.1h under LD and the oscillation persisted under DD with a peak at CT 8.5h. (Cosinor analysis, $p<0.002$)   (C) Transcription profile of *snpf* mRNA in *per*^{0} flies under 12:12 h light-dark cycle (LD). Three replicates were used for each time point. Each replicate contained 30 heads from 2-3 day old w^{1118} male flies. Error bars represent mean ± SEM.

Figure 2. Role of sNPF expressing circadian clock neurons in starvation-mediated hyperactivity. (A-C) Average locomotor activity counts/hour of *tim-Gal4 > w^{1118}*, *w^{1118} >snpf ^{RNAi}* and *tim-Gal4 > UAS-snpf ^{RNAi}* (*tim-Gal4 > snpf ^{RNAi}*) flies assayed with ad libitum food and under starvation for 36 h. Flies were transferred to fresh food or to 1% agar medium at the onset of dark phase to provide the fed and starved condition respectively. Activity counts per hour is plotted along the *y*-axis and hours of starvation on *x*-axis. Gray shaded bars indicate dark phase under LD for 36 h time window. Control flies *tim-Gal4 > w^{1118}* ($n= 27$) (ANOVA, $p<0.0001$), *w^{1118} > snpf ^{RNAi}* ($n= 30$) (ANOVA, $p<0.0001$) and *tim-Gal4 > snpf ^{RNAi}* ($n= 27$) (ANOVA, $p<0.0001$) exhibited enhanced locomotor activity under starvation condition compared to fed controls. (D-F) Total activity of *tim-Gal4 > w^{1118}*, *w^{1118} > snpf ^{RNAi}* and *tim-Gal4 >snpf ^{RNAi}* flies during the first dark phase (1-12h), light phase (13-24h) and second dark phase (25-36h) under fed and starved conditions. (G) Change in activity under starved condition compared to the fed

condition. Starved experimental flies (*tim-Gal4 > snpf* ^{RNAi}) exhibited a significant increase in

the activity after 24 h of starvation compared to both *tim-Gal4 > w* ^{1118} (ANOVA, p<0.0001) and

w ^{1118} *> snpf* ^{RNAi} (ANOVA, p<0.0001) controls. (H) Flies expressing *snpf* ^{RNAi} in clock neurons

exhibit reduced triglyceride levels on day 2 compared to both *tim-Gal4 > w* ^{1118} (t-test, *p*<0.0001)

and *w* ^{1118} *> snpf* ^{RNAi} controls (t-test, *p*<0.0001) (*N*=3; *n*= 5). The asterisks on locomotor activity

waveform and the vertical bar graphs indicate that the values are significantly different. Error

bars represent mean ± SEM. For further explanations see text.

Figure 3. Role of SNPF receptors expressed in IPC in regulating starvation-mediated

hyperactivity. (A-C) Average activity counts/hour of *dilp2-Gal4 > w* ^{1118}, *w* ^{1118} *> snpfR* ^{RNAi} and

dilp2-Gal4 > UAS-snpfR ^{RNAi} (*dilp2-Gal4 > snpfR* ^{RNAi}) flies assayed under the fed and starved

condition. . Control flies *dilp2-Gal4 > w* ^{1118} (*n*= 29) (ANOVA, *p*<0.0001), *w* ^{1118} *> snpfR* ^{RNAi} (*n*=

31) (ANOVA, *p*<0.0001) and *dilp2-Gal4 > snpfR* ^{RNAi} (*n*= 30) (ANOVA, *p*<0.0001) exhibited

enhanced locomotor activity under starvation condition compared to fed controls. (D-F) Total

activity of *dilp2-Gal4 > w* ^{1118}, *w* $^{111\ 8}$ *> snpfR* ^{RNAi} and *dilp2-Gal4 > snpfR* ^{RNAi} flies during first

dark phase (1-12h), light phase (13-24h) and second dark phase (25-36h) under fed and starved

conditions. (G) Change in activity under starved condition compared to the fed condition.

Experimental flies (*dilp2-Gal4 > snpfR* ^{RNAi}) exhibited reduction in the activity after 24 hours of

starvation compared to both *dilp2-Gal4 > w* ^{1118} (ANOVA, *p*<0.0001) and *w* ^{1118} *> snpfR* ^{RNAi}

(ANOVA, *p*<0.03) controls.   (H). Triglyceride levels of two day old *dilp2-Gal4 > w* ^{1118}, w ^{1118} >

snpfR ^{RNAi} and *dilp2-Gal4 > snpfR* ^{RNAi} flies (*N*=3; *n*= 5). Other details are the same as in Fig 2.

Supporting information

Figure S1. (A) Percentage survivorship of 5 day old flies expressing *snpf* ^{RNAi} in clock neurons

under starvation condition.   Five-day-old flies expressing *snpf* ^{RNAi} in clock neurons exhibit

reduced starvation resistance compared to the control flies (log-rank test , $p<0.0001$).   (B)

Percentage survivorship of 5 day old flies expressing $snpfR^{RNAi}$ in IPCs under starvation

condition. Flies expressing $snpfR^{RNAi}$ in IPCs exhibit reduced starvation resistance compared to

the control flies (log-rank test , $p<0.0001$).

A

B

C