Long transposon-rich centromeres in an oomycete reveal divergence of

centromere features in Stramenopila-Alveolata-Rhizaria lineages

Yufeng "Francis" Fang[1], Marco A. Coelho[1], Haidong Shu[2], Klaas Schotanus[1], Bhagya C.

Thimmappa[3†], Vikas Yadav[1], Han Chen[2], Ewa P. Malc[4], Jeremy Wang[4], Piotr A. Mieczkowski[4],

Brent Kronmiller[5], Brett M. Tyler[5], Kaustuv Sanyal[3], Suomeng Dong[2], Minou Nowrousian[6], and

Joseph Heitman[1*]

[1]Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham,

North Carolina 27710; [2]College of Plant Protection, Nanjing Agricultural University, Nanjing,

China; [3]Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for Advanced Scientific

Research, Bangalore, India; [4]Department of Genetics, University of North Carolina, Chapel Hill,

North Carolina 27599; [5]Center for Genome Research and Biocomputing and Department of

Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331; [6]Lehrstuhl fuer

Molekulare und Zellulaere Botanik, Ruhr-Universitaet Bochum, Bochum, Germany

*For correspondence: heitm001@duke.edu.

[†]Present address: Department of Biochemistry, Robert-Cedergren Centre for Bioinformatics and

Genomics, University of Montreal, 2900 Edouard-Montpetit, Montreal, H3T1J4, QC, Canada.

Abstract

Centromeres are chromosomal regions that serve as platforms for kinetochore assembly and spindle attachments, ensuring accurate chromosome segregation during cell division. Despite functional conservation, centromere DNA sequences are diverse and often repetitive, making them challenging to assemble and identify. Here, we describe centromeres in an oomycete *Phytophthora sojae* by combining long-read sequencing-based genome assembly and chromatin immunoprecipitation for the centromeric histone CENP-A followed by high-throughput sequencing (ChIP-seq). *P. sojae* centromeres cluster at a single focus at different life stages and during nuclear division. We report an improved genome assembly of the *P. sojae* reference strain, which enabled identification of 15 enriched CENP-A binding regions as putative centromeres. By focusing on a subset of these regions, we demonstrate that centromeres in *P. sojae* are regional, spanning 211 to 356 kb. Most of these regions are transposon-rich, poorly transcribed, and lack the histone modification H3K4me2 but are embedded within regions with the heterochromatin marks H3K9me3 and H3K27me3. Strikingly, we discovered a *Copia*-like transposon (CoLT) that is highly enriched in the CENP-A chromatin. Similar clustered elements are also found in oomycete relatives of *P. sojae*, and may be applied as a criterion for prediction of oomycete centromeres. This work reveals a divergence of centromere features in oomycetes as compared to other organisms in the Stramenopila-Alveolata-Rhizaria (SAR) supergroup including diatoms and *Plasmodium falciparum* that have relatively short and simple regional centromeres. Identification of *P. sojae* centromeres in turn also advances the genome assembly.

Author summary

Oomycetes are fungal-like microorganisms that belong to the stramenopiles within the Stramenopila-Alveolata-Rhizaria (SAR) supergroup. The *Phytophthora* oomycetes are infamous as plant killers, threatening crop production worldwide. Because of the highly repetitive nature of their genomes, assembly of oomycete genomes presents challenges that impede identification of centromeres, which are chromosomal sites mediating faithful chromosome segregation. We report long-read sequencing-based genome assembly of the *Phytophthora sojae* reference strain, which facilitated the discovery of centromeres. *P. sojae* harbors large regional centromeres fully embedded in heterochromatin, and enriched for a *Copia*-like transposon that is also found in discrete clusters in other oomycetes. This study provides insight into the oomycete genome organization, broadens our knowledge of the centromere structure, function and evolution in eukaryotes, and may help elucidate the high frequency of aneuploidy during oomycete reproduction.

Introduction

Accurate segregation of chromosomes during mitosis and meiosis is critical for the development and reproduction of all eukaryotic organisms. Centromeres are specialized regions of chromosomes that mediate kinetochore formation, spindle attachment, and sister chromatid segregation during cell division [1, 2]. The DNA coincident with functional centromeres typically consists of unusual sequence composition (e.g. AT-rich) and structure (e.g. repeats, transposable elements), low gene density, and transcription of non-coding RNA (ncRNA) as well as heterochromatic nature [3]. However, an active centromere is defined not by DNA sequences but by the deposition of a centromere-associated protein called centromere protein A (CENP-A, also known as CenH3) [1, 4]. CENP-A is a histone H3 variant, which replaces the canonical H3 in the nucleosomes at centromeres and provides the foundation for kinetochore assembly [1, 5, 6].

Despite the fact that centromere function is broadly conserved, centromeric sequences vary greatly in size and composition, ranging from "point" centromeres of 125 bp in length to "regional" centromeres consisting of up to megabases of repeated sequences to holocentromeres that extend along the entire length of the chromosome [1, 3, 7]. To date, point centromeres have been only reported in the budding yeast *Saccharomyces cerevisiae* and its close relatives, holocentromeres have been identified in some insects, plants and nematodes, represented by *Caenorhabditis elegans*, while regional centromeres are the most common type and found in nearly all eukaryotic phyla [1, 3]. Most animals and plants have large regional centromeres composed of satellite sequences that are organized into a variety of different higher order repeats [4, 8, 9]. Some plant centromeres also possess a different type of repeat called centromere-specific retroelements (CR) [10]. In comparison, all fungal centromeres identified to date do not contain satellite repeats and have diverse organizations. The size of fungal regional centromeres ranges from several kilobases, such as in *Candida albicans*, to hundreds of kilobases in *Neurospora crassa* [11, 12]. The centromeric sequences of fungal regional

centromeres can be composed of active or inactive clusters of transposable elements and thus

very repetitive, such as in *Cryptococcus* spp. and *N. crassa* [13, 14], or can be nonrepetitive and

very short, such as in the wheat pathogen *Zymoseptoria tritici* [15] and *C. albicans* [16].

Information on centromeres is limited in other eukaryotic lineages. The malaria pathogen

Plasmodium falciparum and the diatom *Phaeodactylum tricornutum* CENP-A binding regions are

characterized by short simple AT-rich sequences [17, 18], while the parasite *Toxoplasma gondii*

has a simple centromere without nucleotide bias [19].

Due to their highly repetitive nature, assembly of large regional centromeres presents a

significant challenge. Emerging long-read sequencing technologies, such as Pacific Bioscience

(PacBio) and Oxford Nanopore Technology (ONT), have led to substantial advances in resolution

of chromosomal structures including highly repetitive sequences such as centromeres. Using

these technologies, centromeres that were difficult to resolve using short-read sequencing, were

defined in various organisms, from fungi [13, 20, 21] to insects [22], plants [23] and humans [24].

Oomycetes are fungal-like organisms but belong to the stramenopila kingdom within the

Stramenopila-Alveolata-Rhizaria (SAR) supergroup [25, 26]. The SAR supergroup contains a

high diversity of lineages that include many important photosynthetic lineages (e.g. diatoms and

kelp), and important parasites of animals (e.g., *Plasmodium*, the causative agent of malaria) and

plants (e.g. oomycetes, or water molds) [27]. *Phytophthora* is a large oomycete genus (>160

species found to date) and contains some of the most devastating plant pathogens that destroy

a wide range of plants important in agriculture, forestry, ornamental and recreational plantings,

and natural ecosystems [28]. One notorious example is *Phytophthora infestans*, which caused

the Great Irish Potato Famine of the mid-1840s [29]. Today, *Phytophthora* species remain

significant threats to major food crops, causing multi-billion US dollars losses annually throughout

the world [28, 30]. *Phytophthora sojae* is a widespread soil-borne pathogen of soybean. Because

of its economic impact, and tractable genetic manipulation [31-33], *P. sojae* has become a model

species to study oomycete genetics, biology, and interactions with plants.

To date, the genomes of more than 20 *Phytophthora* species have been sequenced [34].

Their genomes are generally large and display complex features: they are diploid, highly

heterozygous for heterothallic species, and very repetitive, which makes genome assembly

challenging. The most contiguous oomycete genome assembly published to date is of the *P. sojae*

reference genome, which was generated based on Sanger random shotgun sequencing and

subsequent improvements involving gap closure and BAC sequencing [26, 35]. *P. sojae* genome

assembly v3.0 (www.jgi.doe.gov) spans ~82 Mb and contains 82 scaffolds; however, there are

~3 Mb of unresolved gaps (N's) persisting in the assembly. Recently, significant progress has

been made in genome assemblies of oomycetes based on long-read sequencing [36, 37];

however, the identity or the nature of the DNA sequences that form essential chromosomal

elements such as centromeres, remain unknown. In this study, using the evolutionarily conserved

kinetochore protein CENP-A as a tool, we investigated cellular dynamics of the kinetochore

complex in *P. sojae*, and uncovered the nature of the oomycete centromeres with the aid of long-

read genome sequencing and ChIP-seq (chromatin immunoprecipitation followed by high-

throughput sequencing) technologies. Our findings suggest that the centromeres of *P. sojae* are

divergent from those reported in other SAR lineages, and their features may be used to predict

centromeres in other oomycetes.

Results

GFP-tagging of CENP-A in *P. sojae* reveals clustered centromeres in different life stages

and throughout hyphal growth

Kinetochore protein homologs have been predicted in diverse eukaryotic lineages

including oomycete species [38]. To identify kinetochore proteins in *P. sojae*, we conducted

BLAST searches against the existing *P. sojae* genome database using the predicted oomycete

orthologs as query. Gene models of *P. sojae* kinetochore proteins were examined and corrected

based on RNA-seq data when necessary. Protein sequences were verified based on the presence

of corresponding motifs (S1 Fig and S1 File).

To examine centromere/kinetochore organization and localization in *P. sojae*, we selected

CENP-A, the hallmark of centromere identity in most organisms. The RNA-seq data did not

support the gene models of *CENP-A* that was instead verified by 3'-RACE and RT-PCR, followed

by Sanger sequencing (S2 A-B Fig). *P. sojae* CENP-A (PsCENP-A) has a conserved C-terminus

including the "CENP-A targeting domain" (CATD) (S2C Fig). *GFP* was fused to *CENP-A* at the N-

terminus and transiently expressed in *P. sojae* transformants with a constitutive promoter derived

from the *Bremia lactucae HAM34* gene (S2D Fig). Overexpressed GFP-CENP-A exhibited

nuclear localization with a single fluorescent focus in the nucleus (S2D Fig), suggesting that *P.*

sojae has a clustered centromere organization.

We also generated GFP labeled CENP-A expressed from the endogenous locus utilizing

CRISPR/Cas9-mediated gene replacement (Fig 1A and S3 Fig). Homokaryotic GFP-CENP-A

strains exhibited single GFP foci within nuclei from different *P. sojae* life stages (Fig 1B)*,*

confirming that the clustered centromere organization is a feature in *P. sojae.* In addition, we

tracked the centromere dynamics during hyphal growth, and found that the clustered centromere

pattern was maintained throughout *P. sojae* nuclear division (Fig 1C and S1 Movie).

Identification of centromeres in a long-read Nanopore-based assembly

To identify *P. sojae* centromeres, we performed native chromatin immunoprecipitation (N-

ChIP) using an anti-GFP antibody against the GFP-CENP-A fusion, followed by high-throughput

Illumina DNA sequencing. ChIP-seq reads were mapped to the latest Sanger genome assembly

(*P. sojae* V3 from JGI), which identified 12 scaffolds that showed relatively concentrated

enrichment of CENP-A reads (S4A Fig). CENP-A peaks appeared scattered in Scaffold 1 and

Scaffold 11, while more clustered in the other 10 scaffolds. However, further examination of each

CENP-A binding region revealed that most of the regions were interrupted by many sequence

gaps, which hampered analysis of the sequence features of the candidate centromeres. Thus, we

processed to re-sequence and re-assemble the reference *P. sojae* genome.

To improve the genome assembly of *P. sojae* reference strain P6497, we applied

Nanopore long-read sequencing and generated a *de novo* genome assembly with SMARTdenovo

together with polishing from PacBio and Sanger reads (S5A Fig and S1 Text). The resulting

assembly of the nuclear genome (Psojae2019.1) has a size of 86 Mb contained in 70 contigs,

with a contig N50 of 2 Mb (S5C Fig). Comparison of Psojae2019.1 to the Sanger assembly

indicated that Psojae2019.1 has more repetitive sequences and most regions were collinear (S5

B and S5C Fig, also see S1 Text for details). We also checked telomere repeats using a motif

proposed for oomycetes [39], and found 13 contigs (versus 7 in Sanger) that harbor telomeric

sequences at single ends (S1 Text and S2 File).

ChIP-seq reads derived from PsCENP-A were mapped to the new genome assembly

Psojae2019.1 (S2 Table), which initially revealed 16 regions exhibiting CENP-A enrichment. On

closer analysis, we found that the unassembled centromere in contig 20 was an artifact caused

by inaccurate genome assembly, as this region was duplicated with a centromere-containing

region in contig34 (S6F Fig). Of the 15 remaining CENP-A binding regions, 11 regions were

assembled within contigs, whereas four regions were disrupted at the edge of contigs (Fig 2). We

confirmed the integrity of the 10 centromeres assembled within contigs as they were completely

covered by long reads (S7 Fig), while the CENP-A peaks in Contig 37 and three broken ones (in

Contigs 9, 10, 57) lacked sufficient long-read coverage. We focused on the 10 verified CENP-A

regions for the further studies (Table 1). RNA-seq analysis indicated that all of the 10 CENP-A

regions exhibited low transcription, except the region in Contig 11. Contig 11 contained two

adjacent CENP-A peaks, one was 18 kb and the other was 114 kb, which were interrupted by a

21 kb transcriptionally active region (Fig 4C). Here, we define the entire region (two CENP-A

peaks, and the region between them) as one centromere (*CEN4*). Among the 10 CENP-A regions,

five have a length of ~190 kb, and three are ~160 kb, while *CEN3* and *CEN6* are significant larger

(>270 kb) (Table 1). All of these centromeres have a GC content comparable to the whole genome

(52.16 - 58.13% vs. 54.6%) (Table 1). Taken together, our CENP-A ChIP-seq analysis utilizing

the newly assembled genome indicates that *P. sojae* CENP-A prefers to bind large poorly

transcribed genomic regions with no specific DNA sequence bias.

To examine the correlation between the centromere regions identified in the new genome

assembly and in the Sanger assembly, we conducted synteny analysis using the genomic regions

flanking the centromeres. The locations of CENP-A found in the Psojae2019.1 assembly were

highly correlated with those in the Sanger assembly, except *CEN10* (Table 1, Fig 3 and S6 Fig).

Contig 51 was collinear with the Sanger scaffold 23; however, no enriched CENP-A signal was

detected for this scaffold, probably because the region corresponding to *CEN10* is interrupted by

gaps. Notably, two CENP-A binding regions in Sanger Scaffold 1 were found to correspond to

CEN8 and *CEN9*, and the smaller one (coordinates: 9,667-9,688 kb) was expanded from 20 kb

to 188 kb corresponding to *CEN8* (Table 1, S4B and S6H Figs). In addition, four contigs of the

Psojae2019.1 assembly (contigs 4, 38, 23, and 58) are collinear with Sanger Scaffold 1, and

telomere repeats are found at the ends of Contigs 4 and Contig 58, further suggesting that

Scaffold 1 of the Sanger genome is assembled incorrectly and should be split into two scaffolds

(S6H Fig). Overall, comparison of centromeres identified in the Sanger and Psojae2019.1

assemblies further confirms their authenticity and reflects some misassemblies that are present

in the Sanger genome assembly.

Due to large genome scales and potentially similar chromosome sizes, the karyotypes of

Phytophthora species cannot be well resolved by pulsed-field gel electrophoresis [32, 40]. The

chromosome number of *P. sojae* is not yet accurately known, but has been estimated to be

between 10 and 15 based on an earlier cytological study [41]. By comparing the location of

centromeres in the Sanger and Psojae2019.1 assembly, we could validate and predict the

configuration of 11 centromeres, namely *CEN1-CEN10*, and *CEN_C9* + *CEN_C48* (Table 1 and

S4 Table). Three centromeres, namely *CEN_C37*, *CEN_C10* and *CEN_C57*, are not fully

assembled. Thus, *P. sojae* is estimated to harbor 12-14 chromosomes, which is in agreement

with the previous cytological study.

***P. sojae* CENP-A regions are embedded within heterochromatin**

To define the epigenetic state of *P. sojae* centromeric regions, we performed ChIP-seq

with antibodies against two heterochromatin marks (H3K9me3, trimethylation of lysine 9 of

histone H3, and H3K27me3, trimethylation of lysine 27 of histone H3) and one euchromatin mark

(H3K4me2, dimethylation of lysine 4 of histone H3). The distribution of H3K9me3 and H3K27me3

is generally coincident throughout the genome, and both were colocalized with the CENP-A

binding regions (Fig 4 and S7 Fig). Intriguingly, the heterochromatic region extended 8 kb to 64

kb beyond each CENP-A binding region (Table 1 and Fig 4A), similar to pericentromeric

heterochromatin regions described in other species [14, 22, 42]. In contrast, the euchromatic mark

H3K4me2 was excluded from the CENP-A region and its flanking heterochromatic regions, and

generally overlapped with the mRNA transcriptional profile (Figs. 4 B, 4C and S7 Fig). Thus,

distribution of histone modifications suggests that the CENP-A regions are embedded in

heterochromatin, and we define the heterochromatic regions adjacent to the CENP-A peaks as

pericentric regions.

A *Copia*-like transposon (CoLT) is highly enriched in the *P. sojae* centromeres

The Psojae2019.1 genome assembly contains 31% repetitive sequences, the majority of

which are transposable elements (TEs) (S5D Fig). Our analysis showed that centromeres are

structurally organized differently, but all composed of many repetitive elements, mostly LTR-

retrotransposons (Figs 3, 5A, S6 and S7 Figs). To identify whether the centromeres in *P. sojae*

possess any common sequences or repeat elements, all identified CENP-A regions were subject

to multiple sequence alignment. This analysis found an ~5 kb sequence that is highly similar

(>98%) and shared among 10 centromeres (S8 Fig and S3 File). BLAST analyses with the

consensus 5 kb sequence against the genome revealed that although this element is not

exclusive to centromeres, it is significantly enriched in centromeres: approximately 90% of all

genomic copies of this element localized to centromeres (Fig 5B). Moreover, this element is

present as clusters in centromeric regions, and only sparsely found in other regions of the genome

(Fig 5B), further strengthening its association with centromeres.

In parallel, examination of the 5 kb sequence indicated that it resembled a *Copia*-like

transposon that comprises a *GAG* gene, and a *POL* gene encoding PR (protease), INT (integrase),

RT (reverse transcriptase), and RH (RNase H) domains in order (Fig 5C and S3 File). To identify

the long terminal repeats (LTRs) of CoLT, we analyzed the best Blast hit of the 5 kb sequence.

Examination of its flanking nucleotides and the LTR marks (5' …TG-3' and 5' …CA-3') enabled

us to identify the LTRs that are nearly identical (Fig 5C and S3 File). Hence, we named this

retroelement CoLT for <u>*Copia*</u>-<u>L</u>ike <u>T</u>ransposon. Phylogenetic analysis based on the conserved

RT domains of various reported retroelements confirmed the classification of CoLT as a

Ty1/*Copia* retrotransposon (Fig 6A). Notably, among the selected retroelements shown in Fig 6A,

P. sojae CoLT is distinct from previously studied oomycete *Copia* retrotransposons [43, 44], but

clusters together with *P. infestans Copia*-24_PIT-I annotated in Repbase [45], and with predicted

CoLT sequences found in other two oomycetes as described below. In addition, phylogenetic tree

also suggests that CoLT is close to *Copia* sequences found in plants (Fig 6A).

Further BLAST analysis employing the full-length CoLT including LTRs as a query

indicated that the Psojae2019.1 assembly, in total, harbors 80 CoLT elements of similar length

including 11 full-length elements that possess one long ORF encoding all of the *Copia* domains,

and 69 sequences that have frameshift mutations within domains (S4B File). Phylogenetic

analysis based on the DNA sequences of these elements revealed that the centromeric CoLT

copies cannot be distinguished from those found elsewhere in the genome, indicating that

centromeric and non-centromeric CoLT copies do not evolve separately; however, it is compelling

that almost all full-length CoLT copies (10 out of 11) are found in centromere (Fig 6B). Collectively,

our analyses suggest that CoLT, a retrotransposon that harbors all predicted functional domains

characteristic for a *Copia*-like transposon, is highly enriched in *P. sojae* centromeres, and is also

the only feature shared by all of the centromeres.

CoLT clusters are conserved in two *P. sojae* oomycete relatives and may be a hallmark of

oomycete centromeres

To examine if clustered CoLT elements found in *P. sojae* centromeres are also present in

other oomycete genomes, we conducted BLAST searches using the 5 kb consensus sequence

derived from *P. sojae* centromeres against the genome assemblies of two *P. sojae* relatives,

Bremia lactucae (downy mildew, lettuce pathogen) and *Phytophthora citricola* (citrus pathogen)*,*

which have relatively contiguous genome assemblies. Interestingly, similar CoLTs clusters were

observed in these genomes, and usually appeared once per contig (Figs 7A and 8A). To assess

if these clustered CoLTs were syntenic with the *P. sojae* centromere-containing contigs, we

examined the CoLT clusters that were present within Mb-long scaffolds/contigs. Synteny analysis

demonstrated that five regions in the *B. lactucae* genome that had CoLT clusters were syntenic

with *P. sojae* centromeres (Figs 7A*).* Unexpectedly, Scaffold 2 (original name, SHOA01000004.1,

see S5 File for details) contained two CoLT clusters that were syntenic with *P. sojae CEN3* and

CEN5 (Fig 7B), indicating that Scaffold 2 may be incorrectly assembled. It should be noted that

the *B. lactucae* genome assembly still has a large percentage of unresolved gaps likely due to its

highly heterozygous nature [37]. In comparison, all three selected regions that had clustered CoLT

clusters within *P. citricola* contigs (PcContigs) were syntenic with *P. sojae* centromeres

(*CEN3*/PcContig2, *CEN9*/PcContig1, *CEN5*/PcContig26) (Fig 8). However, a large number of the

CoLT clusters localized at contig ends, or were distributed across the length of short contigs (Fig

8A). This suggests that many of the centromeric regions in *P. citricola* were not fully assembled.

Taken together, we propose that the clustered CoLT elements may be used as criteria to predict

centromere regions in other oomycete species.

Discussion

In this study, we identified centromeres in the oomycete plant pathogen *P. sojae* by

combining long-read sequencing and ChIP-seq with the GFP tagged kinetochore protein CENP-

A. Cellular dynamics analysis revealed that *P. sojae* centromeres were clustered within nuclei in

different life stages and during vegetative growth. 10 fully assembled and five incompletely

assembled CENP-A binding regions were identified. The common features shared by these

regions include: a) a low level of transcription; b) a GC content similar to that of the whole genome;

c) repetitive sequences; d) enrichment for a specific *Copia*-like transposon; e) overlapping and

surrounding heterochromatin; and f) lack of H3K4me2.

While CENP-A is conserved among different organisms, centromere sequences evolve

rapidly [1, 46]. Although the filamentous fungal-like oomycetes are classified in the stramenopiles

of the SAR supergroup, it is intriguing to observe that the centromeres that we identified in *P.*

sojae are much larger and more complex, comparing to those reported in its stramenopile relative,

the diatom *P. tricornutum*, and those found in the parasites (*P. falciparum* and *T. gondii*) of the

alveolates (Fig 9). In the latter three cases, all centromeres are composed of non-repetitive

sequences. Surprisingly, *P. sojae* centromeres show structural similarity to several, only distantly

related, fungal species, such as *N. crassa* [14] and *Cryptococcus neoformans* [13]. These

features include an enrichment of transposons (or their remnants), and overlap with the

constitutive heterochromatin mark H3K9me2/3. Remarkably, the euchromatin mark H3K4me2

has been shown to be associated with centromeres in humans, mouse, *Drosophila*, *S. pombe*,

and rice [42, 47-49], but is excluded from other fungal regional centromeres reported to date and

in *P. sojae*. In humans and *D. melanogaster*, the CENP-A and pericentromeric heterochromatin

domains are spatially distinct, and the CENP-A domain is flanked by but does not overlap with

heterochromatin [42, 49, 50]. In contrast, the entire centromere of *P. sojae* is embedded in

heterochromatin. It is unknown if the distribution of heterochromatin regions affects centromere

distribution in *P. sojae*, but heterochromatin has been shown to be important for centromere

function and kinetochore assembly in *N. crassa* and *S. pombe* [14, 51, 52]. In addition, it is of

interest that *P. sojae* H3K9me3 and H3K27me3 fully overlap with the centromeric regions, which

have not been observed in centromeres of other species thus far, but was shown in human and

mouse pericentromeres [9, 53], and in the centromeres of *Z. tritici* accessory chromosomes (as

their whole accessory chromosomes are enriched for H3K27) [15]. On the other hand, these two

epigenetic marks generally coexist throughout the entire genome, suggesting it might be just a

general profile of H3K27me3 and H3K9me3 in *P. sojae*.

Transposable elements (and their relics) have been known as residents of the

centromeres and pericentromeres of many animals, plants, and fungi [54]. While animal

centromeres are associated with both satellite DNA and retroelements, satellite DNA is usually

regarded as the main sequence components [55]. Centromeres of many plants, such as maize

and rice, are built on centromere-specific retrotransposons (CR), and a certain CR is usually

unique to a particular chromosome [8]. Centromeres of *N. crassa* [14] and *C. neoformans* [13] are

composed of retrotransposons, and the retroelements in *C. neoformans* are centromere-specific

[13]. In comparison, although *P. sojae* regional centromeres include various transposons, many

of these elements are not limited to this region and can also be found in other genomic areas.

Our study shows that a specific *Copia*-like transposon (CoLT) is highly enriched in the *P. sojae*

centromeric regions and confines the CENP-A binding regions (Figs. 5A and S7 Fig). We

identified 11 CoLT homologs possessing intact domains that are typical for an active *Ty1/Copia*

retrotransposon (Fig 5B), indicating that they may be still active. High similarity of LTRs of each

CoLT indicate the transposition may occur recently (S3 File). Interestingly, two independent

studies employed the central and the 3'- terminal part of CoLT as probes (S3 File) for DNA

fingerprinting of *P. sojae* isolates, and polymorphisms were observed by Southern blotting among

strains isolated from different geographic locations [56, 57]. This suggest that CoLT may be

mobile. In contrast, RNA-seq and histone modification derived from the mycelial stage

demonstrated that all of the 11 copies were poorly transcribed and were heterochromatic (Fig 4B

and S7 Fig), indicating that the element may be inactive at least during mycelial growth. Therefore,

it remains to be determined in which life stages or conditions CoLT becomes active. A similar

distribution pattern of centromere-associated retrotransposons was recently found in *Drosophila*

melanogaster [22]. In *D. melanogaster*, a non-LTR retroelement named *G2/Jockey-3* was found

to be enriched in CENP-A chromatin, and this element is also associated with centromeres in its

sister species *D. simulans* [22]. Strikingly, the CoLT elements were found to be clustered in the

genomes of *P. sojae* oomycete relatives, and some of those regions were syntenic with *P. sojae*

centromeric regions. As most of the oomycete genome assemblies were not based on long-read

sequencing technology, and thus are very fragmented, it remains to be seen if the CoLT elements

have evolved to be widely utilized by oomycetes as a platform for CENP-A loading.

N-ChIP was implemented for this study, because several attempts to perform ChIP

analysis based on traditional formaldehyde-cross-linking strategies were unsuccessful. Cross-

linking with 1% formaldehyde caused degradation of DNA and failure of ChIP. *P. sojae*

transformants expressing GFP tagged CENP-A and CENP-C were both used for N-ChIP-seq.

However, only the GFP-CENP-A transformant produced significant enrichment, indicating that the

binding of CENP-C to chromosomes may be too weak to recover target DNA under native

conditions without cross-linking. Despite most of the ChIP-seq reads of CENP-A are clustered

and define the centromeres, we found that additional reads distributed sparsely out of

centromeres. This could be caused by mapping, because elements present in the centromeres

may also be found in non-centromeric regions, for instance, CoLT. In fact, we observed that all of

the CoLT elements including the truncated copies that are present outside of centromeres have

peaks of their chromatin marks, and thus are indistinguishable from the centromere-associated

ones. Future studies independent of genome assembly, such as fluorescence *in situ* hybridization

(FISH) and Hi-C will probably overcome these technical challenges and confirm the identity of

confirm the centromere regions mapped by ChIP-seq.

Our analysis showed that having an improved reference genome assembly based on long-

read sequencing technologies was crucial to the identification and characterization of

centromeres in *P. sojae*. Our attempt to characterize centromere sequences using the classical

Sanger assembly was not successful because most of the non-coding repetitive regions were not

assembled. While the N50 of the new genome assembly Psojae2019.1 is lower than that of the

Sanger assembly, the contigs do not contain gaps and many of the gaps present in the Sanger

assembly have been closed (S6 Fig). We tried to scaffold the assembly with different scaffolding

programs such as npScarf [58], SSPACE [59], LINKS [60] and the optical BioNano mapping (S1

Text and S9 Fig). Although these scaffolders improved the contiguity (up to 35 scaffolds using

SSPACE), they also generated multiple conflicts with the Sanger assembly, and most of the joins

could not be supported by evidence such as long read coverage (S3 Table and S10 Fig). Thus,

we opted to retain the contig-level assembly in our study. However, identification of centromeres

helped to resolve several structural problems present in the "classical" *P. sojae* Sanger assembly,

and revealed potential structural problems in other oomycete genome assemblies. On basis of

the presence of centromeres and predicted telomeres together with synteny analyses, we found

that three Sanger scaffolds/Psojae2019.1 contigs may represent full-length chromosomes,

namely Scaffold 2/Contigs [26+1+35+6] (S6A Fig), Scaffold 5/Contigs [17+36+7+49+45] (S6G

Fig); and partial Scaffold 1/Contigs [58+38+4] (S6H Fig). Notably, telomeres appear on the both

ends of Sanger Scaffold 5 and its syntenic contigs in Psojae2019 (S6G Fig). There are five *P.*

sojae centromeres that are not fully assembled. With the development of sequencing and

assembly technologies, a finalized chromosome-level genome assembly could help to assemble

those broken centromeres, and refine the centromere sequences that we identified.

Centromeres and their associated kinetochore network serve critical functions in genome

stability and replication. Failures in kinetochore assembly and attachment increase the probability

of chromosome mis-segregation leading to aneuploidy [61]. While these drastic genome changes

can be detrimental to the organism, formation of aneuploidy and polyploidy is an important

strategy orchestrated by pathogens to adapt to the environment during periods of stress [62].

Polyploidy and aneuploidy are prevalent in *Phytophthora* natural isolates and in progeny from

sexual reproduction [36, 63-66]. Interestingly, plant hosts can induce aneuploidy of the sudden

oak death pathogen *P. ramorum*, which enhances its phenotypic diversity and increases its

adaption to the environment [65]. Recently, a phenomenon termed dynamic extreme aneuploidy

(DEA) was described in a vegetable oomycete pathogen, *P. capsici*, in which high variability

among progeny produced by asexual spores was caused by ploidy variation [67]. However, the

mechanisms resulting in oomycete aneuploidy and/or polyploidy are understudied. As

centromeres are the functional and structural foundation for kinetochore assembly and proper

chromosome segregation, identification of centromeres and kinetochore proteins in *P. sojae* may

help to illuminate the mechanisms underlying oomycete genetic, genomic, and phenotypic

diversification.

Materials and Methods

***P. sojae* culture and transformation**

All the strains used in this study are listed in S5 Table. The reference *P. sojae* isolate

P6497 (race 2) used in this study was routinely grown and maintained in cleared V8 media at

25 °C in the dark. Transient gene expression assays based on an optimized polyethylene glycol

(PEG) mediated protoplast transformation protocol [31] was applied to examine the nuclear

localization of CENP-A. Stable and homokaryotic transformants expressing GFP tagged CENP-

A (driven by the strong promoter derived from *HAM34* gene) were chosen for ChIP-seq, which

were generated by passaging on V8 supplemented with 50 μg/mL G418 (Geneticin, AG Scientific,

San Diego, California, USA) for at least 5 times followed by zoospore isolation. Co-transformation

was employed to generate strains expressing both H2B-mCherry and GFP-CENP-A.

Transformation was performed as previously described [31]. Sporangia and zoospores were

induced by water flooding according to a method described previously [68].

Construction of plasmids

All the primers used in this study are listed in S6 Table. All GFP fusion constructs were

generated based on the plasmid backbone pYF3-GFP [69], in which StuI was used for the N-

terminal fusions, and HpaI was used for the C-terminal fusions.

3'-RACE was conducted to validate the gene model of CENP-A, according to the

manufacturer instruction (Invitrogen, 18373-019). All PCR-amplifications were performed using

Phusion High-Fidelity DNA Polymerase (NEB, M0530S).

CRISPR-mediated gene replacement

A sgRNA guide sequence whose PAM sequence overlapped with the start codon of

CENP-A was selected as the CRISPR/Cas9 targets. An oligo annealing strategy was used for

assembly of the sgRNA expression cassettes according to previously described methods [31].

HDR templates for CENP-A was assembled using NEBuilder® HiFi DNA Assembly (NEB,

E2621S). 5'-junction, 3'-junction and spanning diagnostic PCR were performed to genotype

mutants, utilizing the primers listed in S6 Table.

Microscopy imaging of *P. sojae* transformants

A Zeiss 780 inverted confocal microscope was adopted to examine the subcellular

localization of GFP tagged CENP-A driven by strong promoters. Images were captured using a

63 X oil objective with excitation/emission settings (in nm) 488/504-550 for GFP, and 561/605-

650 for mCherry. DeltaVision elite deconvolution microscope (Olympus IX-71 base) equipped with

Coolsnap HQ2 high resolution CCD camera was employed to examine the subcellular localization

of GFP tagged CENP-A produced from the native loci. Images were captured using a 100 X oil

objective (100x/1.40 oil UPLSAPO100X0 1-U2B836 WD 120 micron DIC ∞/0.17/FN26.5, UIS2)

with an excitation filter, 475/28 and an emission filter, 525/50 for GFP. Time-lapse experiments

were performed with 40 X oil objective (40x/0.65-1.35 oil UAPO40XOI3/340 1-UB768R WD 100

micron DIC ∞/0.17/FN22, UIS2, BFP1), with the same filters. Confocal images were edited using

microscope's built-in Zen 2012 software (Blue and/or Black edition according to different

purposes). DeltaVision images were edited using Fiji-ImageJ and Photoshop.

High molecular weight genomic DNA extraction and ONT sequencing

High molecular weight (HMW) genomic DNA (gDNA) from *P. sojae* was isolated by the

CTAB DNA extraction method. 1 g 3-day old fresh *P. sojae* liquid cultures were collected by

filtration and washed twice with sterile water. The resulting damp mycelial pads were frozen

immediately in liquid nitrogen in a pre-cooled mortar, then ground by a pestle. Mycelial powder

was transferred to a 50 ml Falcon tube and mixed gently with 10 ml room temperature *P. sojae*

CTAB extraction buffer (200 mM Tris·HCl pH=8.5, 250 mM NaCl, 25 mM EDTA pH=8.0, 2% SDS,

1% CTAB). The suspension was incubated in 65℃ for 15 minutes with mixing every 5 minutes.

An equal volume of phenol/chloroform/isoamyl alcohol (25:24:1, saturated with 10 mM Tris,

pH=8.0 and 1 mM EDTA) was added to the suspension and mixed gently by inverting the tube,

then centrifuge at 4℃, 5000 g for 15min. The supernatant was transferred to a new 50 ml tube

and treated with RNase A (final concentration, 100 µg/ml) at 37℃ for about 1 hour, followed by

proteinase K treatment (final concentration 200 µg/ml) at 50℃ for 2 hours. An equal volume of

chloroform was added to the solution and mixed gently by inverting the tube, then centrifuge, 4℃,

5000 g for 15min. The supernatant was transferred to a new 50 ml Falcon tube and DNA

precipitated by addition of an equal volume of isopropanol. The tube was mixed gently and

incubated on ice for 6 hours. The resulting white clump of DNA was spooled by a pipette tip and

washed once with 70% ethanol. The gDNA was air-dried for 15 minutes at room temperature and

dissolved in 100 µl sterile water. The quantity of DNA was examined by Qubit and the quality was

checked by pulsed field gel electrophoresis (PFGE).

1D Genomic DNA by Ligation kits (SQK-LSK108, for MinION); SQK-LSK109, for GridION)

were used to prepare the Oxford Nanopore library. Oxford Nanopore sequencing runs was

performed on SpotON R9.4 flow cells with MinKNOW V1.11.5 using MinION or SpotON R9.4.1

flow cells with MinKNOW V3.1.20 using GridION. All of the GridION sequence were basecalled

(on GridION, in real time) using Guppy v2.0.5.

Native ChIP-seq

Native ChIP was performed according to the ChIP protocol accompanying Gent, Wang

(70) with modifications. Briefly, 1-3 mg mycelia were collected from 1-1.5 L of ~3-day culture by

filtration system, and ground into fine powder in liquid nitrogen with pre-chilled mortars and pestles.

Nuclei were isolated and digested by micrococcal nuclease (M0247S, NEB) at 37 for 6 min. An

antibody against GFP (Abcam, ab290) was used to immunoprecipitate single nucleosomes

containing the GFP-CENP-A fusion. Antibodies H3K9me3 (Abcam, ab8898), H3K27me3 (Active

Motif, 39157), and H3K4me2 (Millipore, 07-030) were used to immunoprecipitate nucleosomes

with relevant modifications. ChIP-seq of GFP-CENP-A and H3K27me3 were performed by

Genewiz using Illumina NextSeq500 that generated 150 nucleotide paired-end reads (80%-85%

mappability, S2 Table); ChIP-seq of H3K9me3 and H3K4me2 were conducted by BGI using

Illumina Hiseq 4000 that produced 50 nucleotide single-end reads (~98% mappability, S2 Table).

Numbers of reads for each sample are listed in S2 Table.

Analysis of ChIP-seq and RNA-seq

To map ChIP-seq reads to the genomes, the quality of raw ChIP-seq reads were first

assessed by FastQC (v0.11.6). For ChIP-seq of CENP-A, H3K27me3, the resulting reads were

trimmed with fastx-clipper and mapped with Bowtie2 with default parameters [71] and aligned to

the genome assemblies. For H3K9me3 and H3K4me2, the ChIP-seq reads were polished by BGI

prior to be released and thus mapped to the genomes directly using the same Bowtie2 setup. In

case of replicates, the ChIP-seq reads were mapped randomly among all replicates. The aligned

file (.bam) was sorted and indexed by samtools (version 1.9). Subsequently the ChIP-ed and input

samples were analyzed with DeepTools (v3.2.0) "bamCompare" to calculate normalized ChIP

signal (log2[$ChIP_{RPKM}$/$Input_{RPKM}$]) and bigwig files were generated. Then .bw files were visualized

using the Integrative Genome Viewer (IGV). (https://software.broadinstitute.org/software/igv/).

Core centromeres (CENP-A binding regions) were defined by one continuous stretch of ChIP-seq

peaks (> 5 kb) in scaffolds or contigs in the Sanger or Psojae2019.1 genome assembly. The

boundaries of the 10 fully assembled CENP-A regions were defined by peaking calling employing

MACS v2.2.5 (–broad -q 0.00001 --broad-cutoff 0.00001). Peaks with enrichment fold smaller

than 2 were filtered out (S7 File). The boundaries of pericentric heterochromatin were defined by

the start of both RNA-seq and H3K4m2 peaks. To get profile mRNA, the existing RNA-seq reads

(SRA: SRR10283202) derived from the mycelial stage were aligned to the genomes using

HISAT2 (version 2.1.0), and the resulting files (.bam) were sorted and indexed by samtools

(version 1.9). The .bam file was converted to .tdf for visualization using IGV.

Genome assembly, analysis of genomic features and synteny comparison

Details of the *de novo* genome assembly is described in SI Text. To predict gene models,

first, the assembly Psojae2019.1 was subjected to repeat masking utilizing RepeatMasker [72]

based on a library of *de novo*-identified repeat consensus sequences that was generated by

RepeatModeler (www.repeatmasker.org/RepeatModeler.html). Next, the repeat-masked

assembly was used to predict gene models *ab initio* based on MAKER (v2.31.18) [73] with

predicted proteins from available *P. sojae* and *P. infestans* genome annotations as input [26, 74].

GC content was calculated in non-overlapping 5 kb windows using a modified Perl script

(gcSkew.pl, https://github.com/Geo-omics/scripts/blob/master/AssemblyTools/gcSkew.pl) and

plotted as the deviation from the genome average for each contig. Genes encoding ribosomal

RNA (18S, 5.8S, 25S, and 5S) and tRNA were inferred and annotated based on RNAmmer (v1.2)

[75] and tRNAscan-SE (v2.0) [76], respectively. To find telomeres, a custom-made Perl script was

used to search for the sequence "TTTAGGG" that was proposed for oomycetes telomeric

sequences [39]. Pairwise synteny comparison between the two *P. sojae* genome assemblies (i.e.

P. sojae V3 and Psojae2019.1) or between different oomycete species was conducted using

BLASTn. BLASTn hits and other genomic features were plotted using Circos (v0.69-6) [77].

Whole-genome alignment was computed with MashMap (https://github.com/marbl/MashMap)

employing default settings, and was visualized as a dot plot [78].

Bionano mapping

P. sojae protoplasts were generated from 2.5-day old mycelial and were embedded into

agarose. Bionano Prep Cell Culture DNA Isolation Protocol was employed for extracting the high

molecular weight DNA. DNA labelling with DLE-1 was performed according to the standard

protocols provided by Bionano Genomics (Document number 30206, version F). Labelled DNA

samples were loaded into two flow cells and run on a Saphyr system (Bionano Genomics). The

de novo assembly was performed using Bionano Solve 3.3. Standard parameters for Saphyr data

were used without "extend and split" and without haplotype refinement in order to create a single

map for each allele ("optArguments_nonhaplotype_noES_DLE1_saphyr.xml"). In the process of

de novo assembly, data generated from two flow cells were merged. An assembly graph was

generated during a pairwise comparison of all of the molecules with a p value threshold of 1e-11,

and was refined based on molecules aligned to the assembled maps with a p value threshold of

1e-12. After five rounds of extension and refinement, a final refinement was conducted with a p

value threshold of 1e-16. Then, the *de novo* assembled map was used to scaffold the sequence

assembly. When using the hybrid scaffold module of Bionano Solve 3.3 pipeline, the option of

"resolve conflicts" for sequence contigs and Bionano maps was selected. The standard hybrid

scaffold settings with a modified parameter (-E 0) was applied to remove discrepancies between

sequence assembly and Bionano *de novo* assembly. Sequence contigs were *in silico* digested,

based on the recognition sequence (CTTAAG) of DLE-1. Conflicts detection was accomplished

by aligning contig maps to Bionano maps with p value threshold of 1e-10. When divergence was

identified, the conflicts were resolved by cutting either the contig or the map, depending on the

quality of the genome map at the divergent position.

Analysis of transposable elements and identification of CoLT

To identify transposable elements in *P. sojae*, the new genome assembly was subjected

to RepeatMasker (Repbase v23.09) analysis and hits were mapped to this genome assembly.

The *Copia*-like transposon (CoLT) element was identified in a stepwise way by multiple

sequence alignments followed by extraction of a consensus sequence and BLASTn analyses.

Specifically, an approximately 5 kb consensus sequence was identified in the alignment of

centromere sequences (including incompletely assembled ones) utilizing the multiple alignment

program MAFFT, a plug-in in the Geneious R9 software (http://www.geneious.com), with default

parameters. Then the consensus sequence was used as a query to perform a BLASTn search

against the Psojae2019.1 genome assembly. The resulting sequence hits were used to map

against the genome, and hits longer than 500 bp were used for representing in the figures. The

longest sequence hit with highest identity was retrieved, and was used as a query to execute a

second round of BLASTn search against the NCBI database to further characterize the

sequence. The results of BLASTn analysis indicated that that the sequence was highly similar to

a *Copia*-like transposable element. To define the domains of the CoLT, this sequence was

further analyzed by repeat identification (utilizing a bioinformatics software Unipro UGENE [79]),

and by searches utilizing the Repbase database (https://www.girinst.org/) and NCBI CD-search

(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi).

To validate the classification of CoLT as a *Copia* retrotransposon, phylogenetic analyses

were performed based on the alignment of the reverse transcriptase (RT) domain of the

identified *P. sojae* CoLT, and of LTR (*Gypsy* and *Copia*) and LINE retrotransposons previously

characterized in other organisms. NCBI accession number of each TE sequence is shown in

Fig. 6A. CoLT sequences and the deduced RT domains of *P. sojae*, *B. lactucae* and *P. citricola*

are shown in S3 File. A total of 39 protein sequences were aligned with MAFFT v7.310 [80]

employing the L-INS-i strategy (--localpair --maxiterate 1000) and poorly aligned regions were

removed with TrimAl (-gappyout) [81]. A maximum-likelihood phylogeny was inferred using the

LG+I+G4 model of amino acid substitution in IQ-TREE v1.6.5 [82]. Branch support values were

obtained from 10,000 replicates of both the ultrafast bootstrap approximation (UFboot) [83] and

the SH-like approximate likelihood ratio test (SH-aLRT) [84]. Additionally, to investigate if the

centromeric CoLT sequences have a common origin, BLASTn was conducted using one full-

length CoLT (See S3 File for the sequence). All of the full-length CoLT copies were extracted,

and hits with less than 95% sequence identity and shorter than 95% of the query length were

removed. The obtained genomic coordinates were sorted, merged using 'bedtools merge', and

were employed to retrieve the corresponding nucleotide sequences with the aid of 'bedtools

getfasta' from the bedtools package (https://bedtools.readthedocs.io/en/latest/) [71]. CoLT

nucleotide sequences were subsequently aligned using MAFFT, as described above, and the

alignments were manually inspected. A ML phylogenetic tree was constructed based on 80 *P.*

sojae CoLT sequences (66 located within the defined centromeric regions and 14 located

elsewhere) using IQ-TREE and the TN+F+R2 model of DNA substitution. A full-length CoLT

sequence retrieved from the *P. citricola* genome using a similar procedure was used as

outgroup (S4 File). Truncated CoLT elements (i.e. without a full-length coding sequences) were

inferred upon automatic translation of the predicted coding region (see S4 File). All trees were

plotted with iTOL v4.3.3 [85].

Prediction of centromeric regions in *P. sojae* closely related species

To predict centromeres of the two oomycete species, namely *Phytophthora citricola*

P0716, (Genbank: GCA_007655245.1, with permission of the author) and *Bremia lactucae* SF5,

(GenBank: GCA_004359215.1) [37], BLASTn searches were conducted utilizing the *P. sojae*

Copia-like transposon (CoLT) as a query. Significant hits (>90% identity and > 500 bp) were

retrieved, and were plotted to all scaffolds of the *B. lactucae* assembly and to contigs > 10 kb of

the *P. citricola* assembly. For CoLT clusters that were localized within scaffolds or contigs, their

collinearities with the Psojae2019.1 assembly were further examined with BLASTn, and

visualized by Circos.

Data availability

The Nanopore sequencing raw reads are available in NCBI under the BioProject

PRJNA563922; All raw ChIP-seq reads can be accessed under NCBI's SRA listed in S2 Table.

The Psojae2019.1 genome assembly has been deposited at DDBJ/ENA/GenBank under the

accession WWEI00000000.

Acknowledgments

We thank BioNano Genomics Support, in particular, Yuanyuan Chang for technical support of BioNano mapping, Beth Sullivan at Duke University for critical reading and comments on the manuscript, and members of the Heitman, Sanyal and Garre labs for helpful discussions. M.N. would like to thank Ulrich Kück and Christopher Grefen for support by the Department of General and Molecular Botany/Molecular and Cellular Botany of Ruhr University. B.M.T. and B.K. were supported by Oregon State University. These studies were supported by NIH/NIAID grants R01 AI050113-15 and R37 MERIT award AI039115-21 to J.H and by the German Research Foundation (DFG, grant NO407/7-1 to MN). J.H. is also a Co-Director and Fellow of the CIFAR program, Fungal Kingdom: Threats & Opportunities.

References

1.   Kursel LE, Malik HS. Centromeres. Curr Biol. 2016;26(12):R487-R90. Epub 2016/06/22. doi: 10.1016/j.cub.2016.05.031. PubMed PMID: 27326706.

2.   Stimpson KM, Sullivan BA. Epigenomics of centromere assembly and function. Curr Opin Cell Biol. 2010;22(6):772-80. Epub 2010/08/03. doi: 10.1016/j.ceb.2010.07.002. PubMed PMID: 20675111.

3.   Buscaino A, Allshire R, Pidoux A. Building centromeres: home sweet home or a nomadic existence? Curr Opin Genet Dev. 2010;20(2):118-26. Epub 2010/03/09. doi: 10.1016/j.gde.2010.01.006. PubMed PMID: 20206496.

4.   Wang N, Dawe RK. Centromere size and its relationship to haploid formation in plants. Mol Plant. 2018;11(3):398-406. Epub 2017/12/27. doi: 10.1016/j.molp.2017.12.009. PubMed PMID: 29277426.

5.   Black BE, Foltz DR, Chakravarthy S, Luger K, Woods VL, Jr., Cleveland DW. Structural determinants for generating centromeric chromatin. Nature. 2004;430(6999):578-82. Epub 2004/07/30. doi: 10.1038/nature02766. PubMed PMID: 15282608.

6.   Guse A, Carroll CW, Moree B, Fuller CJ, Straight AF. In vitro centromere and kinetochore assembly on defined chromatin templates. Nature. 2011;477(7364):354-8. Epub 2011/08/30. doi: 10.1038/nature10379. PubMed PMID: 21874020; PubMed Central PMCID: PMCPMC3175311.

7.   Steiner FA, Henikoff S. Holocentromeres are dispersed point centromeres localized at transcription factor hotspots. Elife. 2014;3:e02025. Epub 2014/04/10. doi: 10.7554/eLife.02025. PubMed PMID: 24714495; PubMed Central PMCID: PMCPMC3975580.

8.   Comai L, Maheshwari S, Marimuthu MPA. Plant centromeres. Curr Opin Plant Biol. 2017;36:158-67. Epub 2017/04/16. doi: 10.1016/j.pbi.2017.03.003. PubMed PMID: 28411416.

9.   McNulty SM, Sullivan BA. Alpha satellite DNA biology: finding function in the recesses of the genome. Chromosome Res. 2018;26(3):115-38. Epub 2018/07/06. doi: 10.1007/s10577-018-9582-3. PubMed PMID: 29974361; PubMed Central PMCID: PMCPMC6121732.

10.   Jin W, Melo JR, Nagaki K, Talbert PB, Henikoff S, Dawe RK, et al. Maize centromeres: organization and functional adaptation in the genetic background of oat. Plant Cell. 2004;16(3):571-81. Epub 2004/02/20. doi: 10.1105/tpc.018937. PubMed PMID: 14973167; PubMed Central PMCID: PMCPMC385273.

11.   Smith KM, Galazka JM, Phatale PA, Connolly LR, Freitag M. Centromeres of filamentous fungi. Chromosome Res. 2012;20(5):635-56. Epub 2012/07/04. doi: 10.1007/s10577-012-9290-3. PubMed PMID: 22752455; PubMed Central PMCID: PMCPMC3409310.

12.   Yadav V, Sreekumar L, Guin K, Sanyal K. Five pillars of centromeric chromatin in fungal pathogens. PLoS Pathog. 2018;14(8):e1007150. Epub 2018/08/24. doi: 10.1371/journal.ppat.1007150. PubMed PMID: 30138484; PubMed Central PMCID: PMCPMC6107279.

13.   Yadav V, Sun S, Billmyre RB, Thimmappa BC, Shea T, Lintner R, et al. RNAi is a critical determinant of centromere evolution in closely related fungi. Proc Natl Acad Sci U S A. 2018;115(12):3108-13. Epub 2018/03/07. doi: 10.1073/pnas.1713725115. PubMed PMID: 29507212; PubMed Central PMCID: PMCPMC5866544.

14.   Smith KM, Phatale PA, Sullivan CM, Pomraning KR, Freitag M. Heterochromatin is required for normal distribution of *Neurospora crassa* CenH3. Mol Cell Biol. 2011;31(12):2528-42. Epub 2011/04/21. doi: 10.1128/MCB.01285-10. PubMed PMID: 21505064; PubMed Central PMCID: PMCPMC3133421.

15.   Schotanus K, Soyer JL, Connolly LR, Grandaubert J, Happel P, Smith KM, et al. Histone modifications rather than the novel regional centromeres of *Zymoseptoria tritici* distinguish core

and accessory chromosomes. Epigenet Chromatin. 2015;8(1):41. Epub 2015/10/03. doi:
10.1186/s13072-015-0033-5. PubMed PMID: 26430472; PubMed Central PMCID:
PMCPMC4589918.
16.   Sanyal K, Baum M, Carbon J. Centromeric DNA sequences in the pathogenic yeast
Candida albicans are all different and unique. Proc Natl Acad Sci U S A. 2004;101(31):11374-9.
Epub 2004/07/24. doi: 10.1073/pnas.0404318101. PubMed PMID: 15272074; PubMed Central
PMCID: PMCPMC509209.
17.   Hoeijmakers WA, Flueck C, Francoijs KJ, Smits AH, Wetzel J, Volz JC, et al.
Plasmodium falciparum centromeres display a unique epigenetic makeup and cluster prior to
and during schizogony. Cell Microbiol. 2012;14(9):1391-401. Epub 2012/04/18. doi:
10.1111/j.1462-5822.2012.01803.x. PubMed PMID: 22507744.
18.   Diner RE, Noddings CM, Lian NC, Kang AK, McQuaid JB, Jablanovic J, et al. Diatom
centromeres suggest a mechanism for nuclear DNA acquisition. Proc Natl Acad Sci U S A.
2017;114(29):E6015-E24. Epub 2017/07/05. doi: 10.1073/pnas.1700764114. PubMed PMID:
28673987; PubMed Central PMCID: PMCPMC5530661.
19.   Brooks CF, Francia ME, Gissot M, Croken MM, Kim K, Striepen B. *Toxoplasma gondii*
sequesters centromeres to a specific nuclear region throughout the cell cycle. Proc Natl Acad
Sci U S A. 2011;108(9):3767-72. Epub 2011/02/16. doi: 10.1073/pnas.1006741108. PubMed
PMID: 21321216; PubMed Central PMCID: PMCPMC3048097.
20.   Yadav V, Yang F, Reza MH, Liu S, Valent B, Sanyal K, et al. Cellular dynamics and
genomic identity of centromeres in cereal blast fungus. mBio. 2019;10(4):e01581-19. Epub
2019/08/01. doi: 10.1128/mBio.01581-19. PubMed PMID: 31363034; PubMed Central PMCID:
PMCPMC6667624.
21.   Navarro-Mendoza MI, Perez-Arques C, Panchal S, Niolas FE, Mondo SJ, Ganguly P, et
al. Early diverging fungus *Mucor circinelloides* lacks centromeric histone CENP-A and displays
a mosaic of point and regional centromeres. bioRxiv. 2019:706580.
22.   Chang CH, Chavan A, Palladino J, Wei X, Martins NMC, Santinello B, et al. Islands of
retroelements are major components of *Drosophila* centromeres. PLoS Biol.
2019;17(5):e3000241. Epub 2019/05/16. doi: 10.1371/journal.pbio.3000241. PubMed PMID:
31086362; PubMed Central PMCID: PMCPMC6516634.
23.   Lan T, Renner T, Ibarra-Laclette E, Farr KM, Chang TH, Cervantes-Perez SA, et al.
Long-read sequencing uncovers the adaptive topography of a carnivorous plant genome. Proc
Natl Acad Sci U S A. 2017;114(22):E4435-E41. Epub 2017/05/17. doi:
10.1073/pnas.1702072114. PubMed PMID: 28507139; PubMed Central PMCID:
PMCPMC5465930.
24.   Jain M, Olsen HE, Turner DJ, Stoddart D, Bulazel KV, Paten B, et al. Linear assembly of
a human centromere on the Y chromosome. Nat Biotechnol. 2018;36(4):321-3. Epub
2018/03/20. doi: 10.1038/nbt.4109. PubMed PMID: 29553574; PubMed Central PMCID:
PMCPMC5886786.
25.   Keeling PJ, Burki F. Progress towards the Tree of Eukaryotes. Curr Biol.
2019;29(16):R808-R17. Epub 2019/08/21. doi: 10.1016/j.cub.2019.07.031. PubMed PMID:
31430481.
26.   Tyler BM, Tripathy S, Zhang X, Dehal P, Jiang RH, Aerts A, et al. *Phytophthora* genome
sequences uncover evolutionary origins and mechanisms of pathogenesis. Science.
2006;313(5791):1261-6. Epub 2006/09/02. doi: 10.1126/science.1128796. PubMed PMID:
16946064.
27.   Grattepanche JD, Walker LM, Ott BM, Paim Pinto DL, Delwiche CF, Lane CE, et al.
Microbial diversity in the eukaryotic SAR clade: Illuminating the darkness between morphology
and molecular data. Bioessays. 2018;40(4):e1700198. Epub 2018/03/08. doi:
10.1002/bies.201700198. PubMed PMID: 29512175.

28.   Erwin DC, Ribeiro OK. Phytophthora diseases worldwide: American Phytopathological
Society (APS Press); 1996.
29.   Jiang RH, Tyler BM. Mechanisms and evolution of virulence in oomycetes. Annu Rev
Phytopathol. 2012;50:295-318. Epub 2012/08/28. doi: 10.1146/annurev-phyto-081211-172912.
PubMed PMID: 22920560.
30.   Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global
burden of pathogens and pests on major food crops. Nat Ecol Evol. 2019;3(3):430-9. Epub
2019/02/06. doi: 10.1038/s41559-018-0793-y. PubMed PMID: 30718852.
31.   Fang Y, Cui L, Gu B, Arredondo F, Tyler BM. Efficient genome editing in the oomycete
Phytophthora sojae using CRISPR/Cas9. Curr Protoc Microbiol. 2017;44:21A 1 1-A 1 6. Epub
2017/02/07. doi: 10.1002/cpmc.25. PubMed PMID: 28166383.
32.   Judelson HS, Coffey MD, Arredondo FR, Tyler BM. Transformation of the oomycete
pathogen *Phytophthora-megasperma* f.sp. *glycinea* occurs by DNA integration into single or
multiple chromosomes. Curr Genet. 1993;23(3):211-8. Epub 1993/03/01. doi:
10.1007/bf00351498. PubMed PMID: 8382110.
33.   Fang Y, Tyler BM. Efficient disruption and replacement of an effector gene in the
oomycete *Phytophthora sojae* using CRISPR/Cas9. Mol Plant Pathol. 2016;17(1):127-39. Epub
2015/10/29. doi: 10.1111/mpp.12318. PubMed PMID: 26507366; PubMed Central PMCID:
PMCPMC6638440.
34.   McGowan J, Fitzpatrick DA. Genomic, network, and phylogenetic analysis of the
oomycete effector arsenal. mSphere. 2017;2(6):e00408-17. Epub 2017/12/05. doi:
10.1128/mSphere.00408-17. PubMed PMID: 29202039; PubMed Central PMCID:
PMCPMC5700374.
35.   Tyler BM, Gijzen M. The *Phytophthora sojae* genome sequence: foundation for a
revolution. Genomics of plant-associated fungi and oomycetes: dicot pathogens: Springer;
2014. p. 133-57.
36.   Malar CM, Yuzon JD, Das S, Das A, Panda A, Ghosh S, et al. Haplotype-phased
genome assembly of virulent *Phytophthora ramorum* isolate ND886 facilitated by long-read
sequencing reveals effector polymorphisms and copy number variation. Mol Plant Microbe
Interact. 2019;32(8):1047-60. Epub 2019/02/23. doi: 10.1094/MPMI-08-18-0222-R. PubMed
PMID: 30794480.
37.   Fletcher K, Gil J, Bertier LD, Kenefick A, Wood KJ, Zhang L, et al. Genomic signatures
of heterokaryosis in the oomycete pathogen *Bremia lactucae*. Nat Commun. 2019;10(1):2645.
Epub 2019/06/16. doi: 10.1038/s41467-019-10550-0. PubMed PMID: 31201315; PubMed
Central PMCID: PMCPMC6570648.
38.   van Hooff JJ, Tromer E, van Wijk LM, Snel B, Kops GJ. Evolutionary dynamics of the
kinetochore network in eukaryotes as revealed by comparative genomics. EMBO Rep.
2017;18(9):1559-71. Epub 2017/06/24. doi: 10.15252/embr.201744102. PubMed PMID:
28642229; PubMed Central PMCID: PMCPMC5579357.
39.   Fulneckova J, Sevcikova T, Fajkus J, Lukesova A, Lukes M, Vlcek C, et al. A broad
phylogenetic survey unveils the diversity and evolution of telomeres in eukaryotes. Genome Biol
Evol. 2013;5(3):468-83. Epub 2013/02/12. doi: 10.1093/gbe/evt019. PubMed PMID: 23395982;
PubMed Central PMCID: PMCPMC3622300.
40.   Tooley PW, Carras MM. Separation of chromosomes of *Phytophthora* species using
CHEF gel electrophoresis. Experimental Mycology. 1992;16(3):188-96. doi: Doi 10.1016/0147-
5975(92)90027-O. PubMed PMID: WOS:A1992JQ55100003.
41.   Sansome E, Brasier C. Polyploidy associated with varietal differentiation in the
megasperma complex of *Phytophthora*. Transactions of the British Mycological Society.
1974;63(3):461-IN11.

42.   Sullivan BA, Karpen GH. Centromeric chromatin exhibits a histone modification pattern
that is distinct from both euchromatin and heterochromatin. Nature Structural & Molecular
Biology. 2004;11(11):1076-83. doi: 10.1038/nsmb845. PubMed PMID: WOS:000224847600017.
43.   Jiang RHY, Dawe AL, Weide R, van Staveren M, Peters S, Nuss DL, et al. Elicitin genes
in *Phytophthora infestans* are clustered and interspersed with various transposon-like elements.
Molecular Genetics and Genomics. 2005;273(1):20-32. doi: 10.1007/s00438-005-1114-0.
PubMed PMID: WOS:000228187500003.
44.   Basnayake S, Maclean DJ, Whisson SC, Drenth A. Identification and occurrence of the
LTR-Copia-like retrotransposon, PSCR and other Copia-like elements in the genome of
Phytophthora sojae. Curr Genet. 2009;55(5):521-36. Epub 2009/07/31. doi: 10.1007/s00294-
009-0263-9. PubMed PMID: 19641921.
45.   Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in
eukaryotic genomes. Mob DNA. 2015;6(1):11. Epub 2015/06/06. doi: 10.1186/s13100-015-
0041-9. PubMed PMID: 26045719; PubMed Central PMCID: PMCPMC4455052.
46.   Henikoff S, Ahmad K, Malik HS. The centromere paradox: stable inheritance with rapidly
evolving DNA. Science. 2001;293(5532):1098-102. Epub 2001/08/11. doi:
10.1126/science.1062939. PubMed PMID: 11498581.
47.   Volpe TA, Kidner C, Hall IM, Teng G, Grewal SI, Martienssen RA. Regulation of
heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. Science.
2002;297(5588):1833-7. Epub 2002/08/24. doi: 10.1126/science.1074973. PubMed PMID:
12193640.
48.   Li XY, Wang XF, He K, Ma YQ, Su N, He H, et al. High-resolution mapping of epigenetic
modifications of the rice genome uncovers interplay between DNA methylation, histone
methylation, and gene expression. Plant Cell. 2008;20(2):259-76. doi: 10.1105/tpc.107.056879.
PubMed PMID: WOS:000256415400006.
49.   Allshire RC, Karpen GH. Epigenetic regulation of centromeric chromatin: old dogs, new
tricks? Nat Rev Genet. 2008;9(12):923-37. Epub 2008/11/13. doi: 10.1038/nrg2466. PubMed
PMID: 19002142; PubMed Central PMCID: PMCPMC2586333.
50.   Blower MD, Sullivan BA, Karpen GH. Conserved organization of centromeric chromatin
in flies and humans. Dev Cell. 2002;2(3):319-30. Epub 2002/03/07. PubMed PMID: 11879637;
PubMed Central PMCID: PMCPMC3192492.
51.   Scott KC, Merrett SL, Willard HF. A heterochromatin barrier partitions the fission yeast
centromere into discrete chromatin domains. Curr Biol. 2006;16(2):119-29. Epub 2006/01/25.
doi: 10.1016/j.cub.2005.11.065. PubMed PMID: 16431364.
52.   Allshire RC, Madhani HD. Ten principles of heterochromatin formation and function. Nat
Rev Mol Cell Biol. 2018;19(4):229-44. Epub 2017/12/14. doi: 10.1038/nrm.2017.119. PubMed
PMID: 29235574.
53.   Greaves IK, Rangasamy D, Ridgway P, Tremethick DJ. H2A.Z contributes to the unique
3D structure of the centromere. Proc Natl Acad Sci U S A. 2007;104(2):525-30. Epub
2006/12/30. doi: 10.1073/pnas.0607870104. PubMed PMID: 17194760; PubMed Central
PMCID: PMCPMC1766418.
54.   Friedman S, Freitag M. Evolving centromeres and kinetochores. Advances in genetics.
98: Elsevier; 2017. p. 1-41.
55.   Brown JD, O'Neill RJ. The evolution of centromeric DNA sequences. eLS. 2001.
56.   Yin L, Wang Q, Ning F, Zhu X, Zuo Y, Shan W. Identification of a repetitive sequence
element for DNA fingerprinting in *Phytophthora sojae*. Wei Sheng Wu Xue Bao. 2010;50(4):524-
9. Epub 2010/06/22. PubMed PMID: 20560357.
57.   Xiao J, Miao M, Gao K, Xiang G, Yang S, Dong S, et al. Genetic diversity of
Phytophthora sojae in China based on RFLP. Scientia Agricultura Sinica. 2011;44(20):4190-8.
58.   Cao MD, Nguyen SH, Ganesamoorthy D, Elliott AG, Cooper MA, Coin LJ. Scaffolding
and completing genome assemblies in real-time with nanopore sequencing. Nat Commun.

2017;8:14515. Epub 2017/02/22. doi: 10.1038/ncomms14515. PubMed PMID: 28218240;
PubMed Central PMCID: PMCPMC5321748.
59.   Boetzer M, Pirovano W. SSPACE-LongRead: scaffolding bacterial draft genomes using
long read sequence information. BMC Bioinformatics. 2014;15(1):211. doi: 10.1186/1471-2105-
15-211. PubMed PMID: WOS:000338264200003.
60.   Warren RL, Yang C, Vandervalk BP, Behsaz B, Lagman A, Jones SJ, et al. LINKS:
Scalable, alignment-free scaffolding of draft genomes with long reads. Gigascience.
2015;4(1):35. Epub 2015/08/06. doi: 10.1186/s13742-015-0076-3. PubMed PMID: 26244089;
PubMed Central PMCID: PMCPMC4524009.
61.   Compton DA. Mechanisms of aneuploidy. Curr Opin Cell Biol. 2011;23(1):109-13. Epub
2010/09/03. doi: 10.1016/j.ceb.2010.08.007. PubMed PMID: 20810265; PubMed Central
PMCID: PMCPMC2997176.
62.   Todd RT, Forche A, Selmecki A. Ploidy variation in fungi: polyploidy, aneuploidy, and
genome evolution. Microbiol Spectr. 2017;5(4):599-618. Epub 2017/07/29. doi:
10.1128/microbiolspec.FUNK-0051-2016. PubMed PMID: 28752816; PubMed Central PMCID:
PMCPMC5656283.
63.   Elliott M, Yuzon J, C MM, Tripathy S, Bui M, Chastagner GA, et al. Characterization of
phenotypic variation and genome aberrations observed among *Phytophthora ramorum* isolates
from diverse hosts. BMC Genomics. 2018;19(1):320. Epub 2018/05/04. doi: 10.1186/s12864-
018-4709-7. PubMed PMID: 29720102; PubMed Central PMCID: PMCPMC5932867.
64.   Dobrowolski MP, Tommerup IC, Blakeman HD, O'Brien PA. Non-Mendelian inheritance
revealed in a genetic analysis of sexual progeny of *Phytophthora cinnamomi* with microsatellite
markers. Fungal Genet Biol. 2002;35(3):197-212. Epub 2002/04/04. doi:
10.1006/fgbi.2001.1319. PubMed PMID: 11929210.
65.   Kasuga T, Bui M, Bernhardt E, Swiecki T, Aram K, Cano LM, et al. Host-induced
aneuploidy and phenotypic diversification in the sudden oak death pathogen *Phytophthora*
ramorum. BMC Genomics. 2016;17(1):385. Epub 2016/05/22. doi: 10.1186/s12864-016-2717-z.
PubMed PMID: 27206972; PubMed Central PMCID: PMCPMC4875591.
66.   van der Lee T, Testa A, Robold A, van 't Klooster J, Govers F. High-density genetic
linkage maps of *Phytophthora infestans* reveal trisomic progeny and chromosomal
rearrangements. Genetics. 2004;167(4):1643-61. Epub 2004/09/03. doi:
10.1534/genetics.104.029652. PubMed PMID: 15342505; PubMed Central PMCID:
PMCPMC1470988.
67.   Lamour K, Shresthra S, Zhou Y, Liu X, Hu J. Dynamic extreme aneuploidy (DEA) in the
vegetable pathogen *Phytophthora capsici* sheds light on instant evolution and intractability.
bioRxiv. 2018:297788.
68.   Lin L, Ye W, Wu J, Xuan M, Li Y, Gao J, et al. The MADS-box transcription factor
PsMAD1 is involved in zoosporogenesis and pathogenesis of *Phytophthora sojae*. Front
Microbiol. 2018;9(2259):2259. Epub 2018/10/16. doi: 10.3389/fmicb.2018.02259. PubMed
PMID: 30319576; PubMed Central PMCID: PMCPMC6165875.
69.   Fang Y, Jang HS, Watson GW, Wellappili DP, Tyler BM. Distinctive nuclear localization
signals in the oomycete *Phytophthora sojae*. Front Microbiol. 2017;8:10. Epub 2017/02/18. doi:
10.3389/fmicb.2017.00010. PubMed PMID: 28210240; PubMed Central PMCID:
PMCPMC5288373.
70.   Gent JI, Wang N, Dawe RK. Stable centromere positioning in diverse sequence contexts
of complex and satellite centromeres of maize and wild relatives. Genome Biol. 2017;18(1):121.
Epub 2017/06/24. doi: 10.1186/s13059-017-1249-4. PubMed PMID: 28637491; PubMed Central
PMCID: PMCPMC5480163.
71.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2. Epub 2010/01/30. doi:

10.1093/bioinformatics/btq033. PubMed PMID: 20110278; PubMed Central PMCID:
PMCPMC2832824.
72.   Smit A, Hubley R, Green P. RepeatMasker Open-4.0. 2013–2015. 2015.
73.   Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, et al. MAKER: An easy-to-
use annotation pipeline designed for emerging model organism genomes. Genome Res.
2008;18:188-96.
74.   Haas BJ, Kamoun S, Zody MC, Jiang RHY, Handsaker RE, Cano LM, et al. Genome
sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. Nature.
2009;461:393-8. doi: 10.1038/nature08358
75.   Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW. RNAmmer:
consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids Res. 2007;35(9):3100-
8. Epub 2007/04/25. doi: 10.1093/nar/gkm160. PubMed PMID: 17452365; PubMed Central
PMCID: PMCPMC1888812.
76.   Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis
of transfer RNA genes. Nucleic Acids Res. 2016;44(W1):W54-7. Epub 2016/05/14. doi:
10.1093/nar/gkw413. PubMed PMID: 27174935; PubMed Central PMCID: PMCPMC4987944.
77.   Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: an
information aesthetic for comparative genomics. Genome Res. 2009;19(9):1639-45. Epub
2009/06/23. doi: 10.1101/gr.092759.109. PubMed PMID: 19541911; PubMed Central PMCID:
PMCPMC2752132.
78.   Jain C, Koren S, Dilthey A, Phillippy AM, Aluru S. A fast adaptive algorithm for
computing whole-genome homology maps. Bioinformatics. 2018;34(17):i748-i56. Epub
2018/11/14. doi: 10.1093/bioinformatics/bty597. PubMed PMID: 30423094; PubMed Central
PMCID: PMCPMC6129286.
79.   Okonechnikov K, Golosova O, Fursov M, team U. Unipro UGENE: a unified
bioinformatics toolkit. Bioinformatics. 2012;28(8):1166-7. Epub 2012/03/01. doi:
10.1093/bioinformatics/bts091. PubMed PMID: 22368248.
80.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. Epub
2013/01/19. doi: 10.1093/molbev/mst010. PubMed PMID: 23329690; PubMed Central PMCID:
PMCPMC3603318.
81.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3.
Epub 2009/06/10. doi: 10.1093/bioinformatics/btp348. PubMed PMID: 19505945; PubMed
Central PMCID: PMCPMC2712344.
82.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015;32(1):268-74. Epub 2014/11/06. doi: 10.1093/molbev/msu300. PubMed PMID: 25371430;
PubMed Central PMCID: PMCPMC4271533.
83.   Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. UFBoot2: improving the
ultrafast bootstrap approximation. Molecular Biology and Evolution. 2017;35(2):518-22.
84.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst Biol. 2010;59(3):307-21. Epub 2010/06/09. doi: 10.1093/sysbio/syq010.
PubMed PMID: 20525638.
85.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Res. 2019;47(W1):W256-W9. Epub 2019/04/02. doi:
10.1093/nar/gkz239. PubMed PMID: 30931475; PubMed Central PMCID: PMCPMC6602468.
86.   Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified
classification system for eukaryotic transposable elements. Nat Rev Genet. 2007;8(12):973-82.
Epub 2007/11/07. doi: 10.1038/nrg2165. PubMed PMID: 17984973.

87.   Xiong Y, Eickbush TH. Origin and evolution of retroelements based upon their reverse-
transcriptase sequences. Embo J. 1990;9(10):3353-62. doi: DOI 10.1002/j.1460-
2075.1990.tb07536.x. PubMed PMID: WOS:A1990EA35300039.
88.   Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: A resource for timelines,
timetrees, and divergence times. Mol Biol Evol. 2017;34(7):1812-9. Epub 2017/04/08. doi:
10.1093/molbev/msx116. PubMed PMID: 28387841.
89.   Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome
assembly evaluation with QUAST-LG. Bioinformatics. 2018;34(13):i142-i50. Epub 2018/06/29.
doi: 10.1093/bioinformatics/bty266. PubMed PMID: 29949969; PubMed Central PMCID:
PMCPMC6022658.
90.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094-100. Epub 2018/05/12. doi: 10.1093/bioinformatics/bty191. PubMed PMID:
29750242; PubMed Central PMCID: PMCPMC6137996.
91.   Sović I, Šikić M, Wilm A, Fenlon SN, Chen S, Nagarajan N. Fast and sensitive mapping
of nanopore sequencing reads with GraphMap. Nat Commun. 2016;7:11307.
92.   Marcais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast
and versatile genome alignment system. PLoS Comput Biol. 2018;14(1):e1005944. Epub
2018/01/27. doi: 10.1371/journal.pcbi.1005944. PubMed PMID: 29373581; PubMed Central
PMCID: PMCPMC5802927.

Figures and Tables

Fig 1. Subcellular localization of CENP-A in *P. sojae* at different life stages and during vegetative growth. (A) A schematic showing the generation of *GFP*-fused *CENP-A* utilizing CRISPR/Cas9 mediated gene replacement. (B) Subcellular localization of GFP-tagged CENP-A (expressed from the endogenous locus) in *P. sojae* hyphae, sporangia, and encysted zoospores. (C) Time-lapse images illustrating localization of GFP tagged CENP-A during hyphal growth. Dashed squares denote occurrence of nuclear division. Representative images are shown. Scale bars in all images, 5 µm.

Fig 2. Contigs in the Psojae2019.1 assembly demonstrating CENP-A enrichment based on ChIP-seq. (A) 10 fully assembled CENP-A binding sites presented contigs. (B) 5 incompletely assembled CENP-A binding regions. All contigs are drawn to scale and the ruler indicates the length of the contigs. All CENP-A profiles shown were normalized to input DNA. mRNA profiles are shown as log-scales. Solid stars indicate the CENP-A enriched regions within contigs; hollow stars denote broken centromeres at the edge.

Fig 3. A representative Circos visualization comparing centromere-containing genomic regions between the Sanger V3 Scaffold 2 and the corresponding contigs in the Psojae2019.1 assembly. For the inner circle, track A illustrates assembled contigs (in Psojae2019.1) or scaffold (in P. sojae V3) that are color coded as shown at the top. The locations of centromeres (CENP-A binding regions) are highlighted in yellow. Tracks B-E show the location of other genomic features as given in the key on the bottom. Blue and orange lines in track F link regions with collinearity extending over 2 kb, with orange lines corresponding to inversions. Grey box-shaded centromere-containing regions are magnified for detailed visualization (the two outer arcs).

Fig 4. Epigenetic state of *P. sojae* centromeres. (A) Schematics showing the *P. sojae* core

centromeres (CENP-A binding regions) and the pericentric regions of various lengths. Dark and

light grey bars indicate core centromeric and pericentric regions. Digits at the center indicate the

size of core centromeres; digits on the left denote the full-length of the centromeres (a combination

of core centromere and pericentromeric region). The right pericentric region of *CEN5* and the left

pericentric region of *CEN10*, indicated by dashed bars, are not fully assembled, and their lengths

are labeled with question marks. (B-C) Two centromeres (*CEN1* and *CEN4*) are shown as

representatives to compare CENP-A localization to the distributions of modified histones. A 400

kb region harboring the centromeric region is shown for *CEN1* and *CEN4*. Cyan block, a

transcriptionally active region (21 kb) that interrupts *CEN4*. Profiles of CENP-A, H3K9me3,

H3K27me3 and H3K4me2 shown were normalized to input. mRNA profiles are shown as log-

scales.

Fig 5. *P. sojae* centromeres are enriched for a *Copia*-like transposon (CoLT). (A) Distribution of

transposable elements (TEs) in *CEN1*. TEs were annotated using a *Phytophthora* TE library

from Repbase [45]. Tracks of different repeat families are color coded. The track "Other TEs"

includes all types of TEs beyond *Gypsy* and *Copia*. CoLT was composed of three elements

annotated in Repbase [45], namely Copia-24_PIT-I, Copia-24_PIT-LTR and Gypsy-P17-PR-I.

The profile of CENP-A shown was normalized to input. The mRNA track is shown as log-scales

and used to define the boundary of pericentric heterochromatin. (B) Location of CoLT elements

across all Psojae2019.1 contigs. (C) Diagram showing the domain structure of a representative

full-length CoLT sequence (see S3 File). The coding domain is characteristic of *Copia*

superfamily of retrotransposons, which consists of capsid protein (Gag), Gag-pre-integrase

(PR), integrase (INT), reverse transcriptase (RT), and RNase H (RH) domains, and diverges

from the *Gypsy* superfamily in the order of the RT and INT domains in their *POL* genes [86].

Fig 6. Phylogenetic analyses of CoLT. (A) Maximum-likelihood phylogeny of different retroelements. Protein sequences of the reverse transcriptase (RT) domains were used to construct the tree [87] (see S4A File for the complete DNA and protein sequences). The tree was rooted in the midpoint and branch support values (> 50%) shown at the three nodes were determined by 10,000 replicates of both ultrafast bootstrap approximation (UFBoot) and the Shimodaira-Hasegawa approximate likelihood ratio test (SH-aLRT). (B) Maximum-likelihood phylogeny of full-length CoLT elements identified in the Psojae2010.1 genome assembly. CoLT copies locate inside and outside of centromeres are denoted by blue and red circles. CoLT elements with full-length coding sequences (CDS) are depicted as filled circles. A CoLT sequence identified in *P. citricola* served as an outgroup. See S4A File for the CoLT copies used for the phylogenetic analysis.

Fig 7. Genomic distribution of CoLT in the *Bremia lactucae* genome. (A) Location of CoLT elements across all *B. lactucae* scaffolds >100 kb. For ease of analysis, scaffolds in *B. lactucae* assembly were sorted and re-named based on sizes (large to small). See S5 File for the original scaffold names. Regions underlined by green lines indicate both sides of the CoLT clusters were syntenic with the regions surrounding *P. sojae* centromeres. Regions underlined by blue indicate only one sides of the CoLT clusters were found to be syntenic with *P. sojae* centromere flanking sequences. (B) A representative Circos plot comparing *B. lactucae* Scaffold 2 that has clustered CoLTs with the corresponding Psojae2019 contigs. The outer track illustrates the assembled scaffold (in the sorted *B. lactucae* assembly) or contigs (in Psojae2019.1), which is color coded as shown at the top. Names of contigs possessing *P. sojae* centromeres are enclosed in circles. Yellow regions on the outer tracks indicate the locations of centromeres (CENP-A binding regions). Blue and orange lines in track F link regions with synteny extending over 2 kb, with orange lines corresponding to inversions. Two CoLT clusters are identified in the *B. lactucae* Scaffold 2 (s2; original scaffold, SHOA01000004.1), which are indicated by arrowheads.

1025

Fig 8. Genomic distribution of CoLT in the *Phytophthora citricola* genome. (A) Location of CoLT

across all *P. citricola* contigs >100 kb. (B-D) Circos plots comparing three *P. citricola* contigs that

have CoLT clusters with the corresponding Psojae2019 contigs. In each panel, the outer track

(bars) illustrates the contigs in the *P. citricola* or the Psojae2019.1 genome assemblies, which is

color-coded as shown below the Circos plot in panel B. Yellow regions indicate the locations of

centromeres (CENP-A binding regions). Names of contigs harboring *P. sojae* centromeres are

enclosed in circles. Blue and orange lines in track F link regions with synteny extending over 2

kb, with orange lines corresponding to inversions. CoLT clusters present in the *P. citricola* contigs

are indicated by arrowheads. The flanking sequences of each CoLT cluster are syntenic with the

regions surrounding *P. sojae* centromere.

1036

Fig 9. Diversity of centromere features within the SAR supergroup. Simplified schematics (not to

scale) showing the structure, epigenetic modifications, size and composition of centromeres

across SAR lineages. Asterisk, epigenetic state was not examined in the diatom centromeric

regions; however, several AT-rich DNA sequence can be employed for episome maintenance,

suggesting diatom centromere might not be epigenetically dependent. Histone modification

H3K27me2 was only tested in *P. sojae*. The phylogenetic tree was constructed using TimeTree

[88]. *Homo sapiens*, *Arabidopsis thaliana* and *Neurospora crassa* were used as representatives

of animals, plants and fungi for the phylogenetic analysis, and are used as outgroups illustrating

the evolutionary status of the SAR supergroup.

Table 1. Centromeres identified in the Psojae2019.1 assembly and their counterparts in the
Sanger assembly

		Psojae2019.1				Sanger V3	
Name	Contig	Position of core centromere, kb (size, kb)*	GC% of *CEN*	Position of pericentric region, kb (size, kb)		Scaffold	Position of core centromere (kb)*
CEN1	1	415-578 (163)	56.91	361-415 (54) 579-650 (71)		2	7086-7267
CEN2	2	4102-4295 (193)	55.51	4094-4102 (8) 4296-4305 (9)		8	2374-2420
CEN3	3	3138-3410 (272)	54.81	3223-3138 (15) 3412-3560 (48)		9	3075-3286
CEN4†	11	991-1174 (183)	58.13	944-991 (47) 1175-1205 (30)		4	995-1246
CEN5	18	1556-1706 (150)	57.93	1492-1556 (64) 1709-? (>22)		3	4078-4138
CEN6	34	697-880 (183)	57.65	643-696 (53) 882-904 (22)		6	1521-1726
CEN7	36	432-706 (274)	52.16	376-433 (57) 708-732 (24)		5	2049-2310
CEN8	38	302-490 (188)	57.01	288-302 (14) 490-515 (25)		1	9667-9688
CEN9	41	154-342 (188)	57.40	101-153 (52) 342-358 (16)		1	2921-3079
CEN10	51	36-151 (115)	57.93	?-36 (>36)† 152-216 (64)		-	-

1048

* The coordinates of core centromeres in the Psojae2019.1 assembly are defined by peak calling (S7 File). The positions of
centromeres in the Sanger assembly are defined by visualization of peaks.
†Contig11 contains a minor (coordinate, 991465 - 1009035, 18 kb) and a major (coordinate, 1060806 - 1174302, 114 kb) peak that
are separated by a 21 kb transcriptionally active region. The core centromere is composed of the minor and major CENP-A regions,
and the region between the two peaks.
‡One side of pericentric heterochromatin region is not fully assembled.

 Supporting information

 S1 Fig. Summary of the presence and absence of putative core kinetochore proteins

 identified in *P. sojae.* Kinetochore orthologs were identified based on BLAST searches. *P. sojae*

 CENP-A (in bold) was selected to track subcellular localization of centromere/kinetochore and

 profile centromere sequences. Sequences of *P. sojae* putative core kinetochore proteins are listed

 in S1 File.

 S2 Fig. Identification and expression of CENP-A in *P. sojae.* (A) Image of *CENP-A* model

 and RNA-seq log scale coverage taken from FungiDB. Arrows and triangle denote an

 erroneous *P. sojae* CENP-A gene model caused by an intron that was missed in the gene model

 prediction. (B) Electrophoresis image showing 3'-RACE result of *CENP-A*. 5'-primer JOHE45057

 (not shown in scale) served as a gene-specific primer for 3'-RACE (See S6 Table). (C) Alignment

 of *P. sojae* CENP-A with orthologs from different organisms. Ps, *P. sojae*; Pt, diatom

 (*Phaeodactylum tricornutum)*; Sc, *Saccharomyces cerevisiae;* Cn, *Cryptococcus neoformans; Hs,*

 Homo sapiens; At, *Arabidopsis thaliana*; Dm, *Drosophila melanogaster.* (D) Transient expression

 of GFP tagged CENP-A in *P. sojae* transformants. Upper panel, a plasmid constructed for

 transient expression of *CENP-A*. Expression of *P. sojae* CENP-A is driven by a constitutive

 promoter derived from the *B. lactucae HAM34* gene. Lower panel, subcellular localization of GFP-

 tagged CENP-A in the *P. sojae* transformants based on the constructs in (A).

 S3 Fig. Generation of *P. sojae* strains expressing *GFP* tagged *CENP-A* utilizing

 CRIPSR/Cas9 mediated genome editing. (A) Schematic of gene replacement of the

 endogenous *CENP-A* with *GFP-CENP-A*. Lightning bolts, sgRNA guideline sequences were

 designed overlapping the start codon of *CENP-A*. Primer pairs, JOHE50062/JOHE45358,

 JOHE45420/JOHE50063, JOHE50062/JOHE50063 were used for 5'-junction, 3'-junction, and

1080   spanning diagnostic PCR screening *GFP-CENP-A* mutants. See S6 Table for the primer

1081   information. (B) Representative genotyping results of zoospore isolated (homokaryotic) *GFP-*

1082   *CENP-A* strains (YFP10a1 and YFP10b1, see S5 Table for their genetic background). Products

1083   observed in the 5'- and 3'-junction PCR of wild type (WT) are non-specific amplicons.

1084

S4 Fig. Scaffolds in the Sanger assembly that are suggested to harbor putative

centromeres. (A) CENP-A enrichment identified in 12 scaffolds. Solid and hollow stars denote

CENP-A enrichment regions that are sequence-gap free or contain gaps, respectively. All CENP-

A profiles shown have been normalized to input. mRNA profiles are shown as log-scales. (B) Two

putative centromeric regions in Scaffold 1 were identified by poor transcription and the syntenic

regions in the Psojae2019.1 assembly.

1091

S5 Fig. Pipeline used for *de novo* assembly and metrics of the *P. sojae* genome assembly

Psojae2019.1. (A) Pipeline used for *de novo* assembly of Psojae2019.1. Box in grey, different

scaffolding programs were employed to enhance the contiguity of the assembly (See more details

in S10 Fig and S3 Table). As some of them generated conflict and sequence gaps, we opted to

use the contig-level assembly for the centromere study. (B) Dotplot comparison of the *de novo*

assembly Psojae2019.1 against the Sanger assembly. (C) Genome assembly metrics. *Statistics

for both the Psojae2019.1 and Sanger V3 assemblies were calculated by QUAST [89];

†Annotation was based on the repeat-masked assembly (See method). ‡Annotation was obtained

from FungiDB release 33 (https://fungidb.org/fungidb/). §Measured by RepeatModeler (See

methods). (D) Pie chart summarizing retroelements (LINE, SINE and LTR), DNA transposons and

other repeat sequences predicted in the Psoaje2019.1 assembly.

S6 Fig. Comparison of centromere-containing genomic regions between the Sanger (P.

sojae V3) and the Psojae2019.1 assemblies. In all panels (A-H), the outer tracks illustrate

assembled contigs (in Psojae2019.1) or scaffolds (in P. sojae V3), and are color coded as given

in key on the bottom. Yellow regions on the outer tracks indicate the location of centromeres

(CENP-A binding regions). Blue and orange lines in track F link regions with synteny extending

over 2 kb, with orange lines corresponding to inversions. Names of contigs that contain *P. sojae*

centromeres are enclosed in circles. Arrowheads indicate the shrunk centromeres present in the

Sanger scaffolds. (A) Comparison of Sanger Scaffold 2 (sca2) and its syntenic contigs in the

Psojae2019 assembly. (B) Comparison of Sanger Scaffold 8 (sca8) and its syntenic contigs in the

Psojae2019 assembly. (C) Comparison of Sanger Scaffold 9 (sca9) and its syntenic contigs in

the Psojae2019 assembly. (D) Comparison of Sanger Scaffold 4 (sca4) and its syntenic contigs

in the Psojae2019 assembly. Dots under *CEN4* indicate the regions showing CENP-A peaks, as

a transcriptionally active region is found in *CEN4*. (E) Comparison of Sanger Scaffold 3 (sca3)

and its syntenic contigs in the Psojae2019 assembly. (F) Comparison of Sanger Scaffold 6 (sca6)

and its syntenic contigs in the Psojae2019 assembly. Dashed lines under Psojae2019.1 Contigs

20 and Contig 34 indicate duplicated regions. (G) Comparison of Sanger Scaffold 5 (sca5) and

its syntenic contigs in the Psojae2019 assembly. (H) Comparison of Sanger Scaffold 1 (sca1) and

its syntenic contigs in the Psojae2019 assembly. The two centromeric regions in Scaffold 1 of the

Sanger assembly (P. sojae V3) suggested by CENP-A ChIP-seq are indicated by arrows. Part of

the Contig4 is inverted in the Psojae2019.1 assembly compared to the Sanger assembly

(inversion breakpoint indicated by an asterisk). Intriguingly, both versions can be supported by

some of the mapped reads indicating that both ends of the inverted region are repeat-rich and

composed of the same types of repeats so that reads map to both versions. Alternatively, this

may represent a structural rearrangement between the two haplotypes of the diploid genome, or

it could also represent a real difference between the P6497 culture used for the Sanger sequence

and the one used for the ONT assembly.

S7 Fig. Summary of features of each intact centromere and read coverage analysis of

centromere. For each panel (A-J), upper, an overview of the contig exhibiting enrichments of

both CENP-A and histone modifications (H3K9me3, H3K27me3 and H3K4me2), and distribution

of transposable elements. TE, transposable elements. CoLT, *Copia* like transposon. Middle, a

magnified view of the region shaded in the contig shown above; a diagram displaying the

correlation of core centromere (CENP-A) and heterochromatic regions. A 400 kb region is shown

for each centromere that has assembled pericentromeric regions on both sides. A 270 kb region

is shown for *CEN5*, as it is missing one side of the intact pericentric region. The length of Contig51

is close to 400 kb, only the entire contig is shown (J). Bottom, an image showing long-read

coverage. Canu-corrected Nanopore reads were mapped to all centromeres using Minimap2 [90],

except *CEN7*, which was verified by all Nanopore and PacBio reads (without Canu-correction)

using GraphMap [91]. For better visualization, indels of reads were masked. Asterisk in (B),

regions (underlined) that have high CENP-A enrichment but no CoLT and very low TE density

contains unknown repetitive sequences unique to *CEN2*.

1143

1144   **S8 Fig. MAFFT-based alignment of CENP-A binding regions reveals a 5 kb sequence that**

1145   **are highly similar among *P. sojae* centromeres.** Upper panel, alignment of CENP-A binding

1146   regions. Lower panel, magnified view of the conserved 5 kb region. Images were adapted from

1147   the alignment result generated by Geneious.

1148

1149   **S9 Fig. Representative contigs that are anchored by BioNano mapping and contigs that**

1150   **are suggested to be joined.** (A) Contig 2 representing most of the cases that only parts of contigs

1151   were anchored by Bionano molecules. (B) One case that the two contigs were not anchored by

1152   BioNano but suggested to be combined. (C-E) Three contigs that were anchored by BioNano and

suggested to be combined. (E, F) Two cases showing that contigs can be fully covered by BioNano molecules. Map102 was a fragment trimmed from Contig 32. NGS, next generation sequencing, i.e. Psojae2019.1; BNG, BioNano *de novo* assembly; HYBRID, hybrid scaffold.

S10 Fig. Dot plot comparison of scaffolded assemblies against the original Psojae2019.1 assembly and the Sanger assembly. (A-D) Assemblies generated by the scaffolding programs npSarf, SSPACE and LINKS, and BioNano mapping were aligned trespectively to the original SMARTdenovo assembly (Psojae2019.1), as well as the Sanger genome, and plotted utilizing the MUMmer package [92].

S1 Table. Metrics of ONT sequencing

S2 Table. Statistics of ChIP-seq samples

S3 Table. Metrics of scaffolded assemblies and their comparison to the Sanger V3 and Psojae2019.1 assembly.

S4 Table. Five incompletely assembled centromeres in the Psojae2019 assembly and their corresponding CENP-A regions mapped in the Sanger assembly

S5 Table. *P. sojae* strains used in the study

S6 Table. Primers used in this study

S1 Text. Nanopore sequencing and *de novo* assembly of the reference *P. sojae* genome

S1 Movie. Time-lapse experiment showing cellular dynamics of CENP-A during *P. sojae* vegetative growth.

S1 File (Separate file). Sequences of kinetochore orthologs identified in *P. sojae.*

S2 File (Separate file). 13 telomeres predicted in the Psojae2019.1 assembly.

S3 File (Separate file). The 5 kb consensus sequence shared by *P. sojae* centromeres, and

representative full-length CoLT homologs in *P. sojae*, *B. lactucae* and *P. citricola*.

S4 File (Separate file). Sequences used for building phylogenetic trees in Fig 6. (A)

Sequences employed for the phylogenetic analysis in Figure 6A. (B) Genomic coordinates

of the sequences used for constructing phylogenetic tree in Figure 6B.

S5 File (Separate file). Original names of the sorted *B. lactucae* scaffolds.

S6 File (Separate file). Bionano mapping report.

S7 File (Separate file). CENP-A peaks called by MACS in the 10 contigs that harbor fully

assembled centromeres

Figure 1

Figure 2

Figure 3

key:

Ⓐ contigs: | Telomeric repeats ▮ Centromeres ∙∙∙ assembly gaps

Ⓑ ▮ Copia-like TE (CoLT) enriched in the centromeres

Ⓒ Transposable elements (from the outside inward):
▮ LTR retrotransposons ▮ DNA transposons ▮ Other transposons

Ⓓ ╱╲ GC content — red above / blue below genome average
(5 kb non-overlapping window)

Ⓔ ▮ tRNA genes

Ⓕ BLASTn links (> 2 kb)

A

B

C

A

Contig 1

mRNA [0-10955]

CENP-A [1-6]

Ty1/Copia

Ty3/Gypsy

Other TEs

CoLT

B Contigs

P. sojae2019.1

key:
- Complete *CEN*
- Broken *CEN*
- *Copia*-like transposon (CoLT)
- Duplicated *CEN* region

C LTR — GAG — PR — INT — RT — RH — LTR

POL

Figure 6

A

| 0.1 |

88.5/88
90.1/85
99.9/100
91/99
92.2/100
73.8/-
80/-
90.9/98
88.8/56
82.1/64
66/90
83/95
90.3/98
78.2/97
100/100
82.6/91
82.6/66
100/100
84.6/52
86.5/-
92.7/-
69/-
91.8/75
99.9/100
96/99
92.8/63
78.3/-
80.9/-
85.2/62
99/99

Phytophthora infestans CopiaPi-2 | AY830099
Arabidopsis thaliana Copia | BAB84015
Oryza sativa Copia | AAR88589
Phytophthora infestans Copia-24 PIT-I 1p | Repbase
Bremia lactucae CoLT | Sca7 2388-7076
Phytophthora sojae CoLT | ctg38 322956-327751
Phytophthora citricola | Sca60 26953-31785
Drosophila melanogaster | P04146
Nicotiana tabacum Tnt1-Copia | P10978
Cryptococcus deneoformans TCN6 | XP_777044
Phaeodactylum tricornutum CoDi6.5 | ACA60907
Thalassiosira pseudonana CoDi6.3 | ACA60905
Nannochloropsis gaditana Copia | EWM20840
Saccharomyces cerevisiae Ty1 | ADB23365
Phytophthora infestans CopiaPi-3 | AY830100
Phytophthora sojae PSCR Copia | DQ485162
Cryptococcus deneoformans CNL1 | XP_571046
Drosophila melanogaster Jockey | P21328
Homo sapiens LINE-1 | AAD39215
Phytophthora infestans LINEPi-1 | AY830108
Drosophila melanogaster Gypsy | AAB50148
Thalassiosira pseudonana Gypsy | ACA60916
Saccharomyces cerevisiae Ty3 | Q99315
Phytophthora infestans GypsyPi-2 | AY830106
Phytophthora infestans GypsyPi-4 | AY830107
Bombyx mori Mag | S08405
Drosophila melanogaster Mdg3 | Q94885
Cryptococcus deneoformans TCN2 | XP_567971
Phytophthora infestans GypsyPi-1 | AAV92918
Phytophthora infestans GypsyPi-220 | AF490339
Phytophthora infestans GypsyPi-3 | AY830104
Arabidopsis thaliana Gypsy | AAD17358
Solanum lycopersicum Gypsy | Q9ZS84
Tricholoma matsutake MarY1 | BAA78625.
Cryptococcus deneoformans TCN1 | XP_571377
Passalora fulva Ctf-1 | AAF21678
Fusarium oxysporum Skippy | S60179
Magnaporthe grisea Maggy | AAA33420
Alternaria alternata REAL | BAA89272

LTR
Ty1/*Copia*
Ty3/*Gypsy*

Non-LTR

B

P. sojae

0.006

○ *CEN* CoLT
○ *Non-CEN* CoLT
● Full length CDS

P. citricola

Figure 7

Figure 8