Single-trial dynamics of competing reach plans in the human motor periphery

Abbreviated title: Competing reach plans in the motor periphery

Luc P. J. Selen[1], Brian D. Corneil[2-5] and W. Pieter Medendorp[1]

[1]Radboud University, Donders Institute for Brain, Cognition and Behaviour; 6500 HB Nijmegen,

The Netherlands

Departments of [2]Psychology and [3]Physiology & Pharmacology; University of Western Ontario;

London, ON, N6A 5B7; Canada

The [4]Brain and Mind Institute and [5]Robarts Research Institute;

Corresponding Author:

Dr. Luc Selen

luc.selen@donders.ru.nl

Keywords: motor planning; decision making; visually-guided reaching; electromyography

Number of Pages: 38

Number of Figures: 7

Words in Abstract: 248

Words in Significance Statement: 109

Words in Introduction: 650

Words in Discussion: 1600

Acknowledgement

27 This work was supported by operating grants from the Natural Sciences and Engineering
28 Research Council of Canada (NSERC) to BDC [RGPIN-311680], the Canadian Institutes of
29 Health Research (CIHR) to BDC [MOP-93796], a Vici grant from the Netherlands Organization
30 for Scientific Research to WPM [453-11-001].

ABSTRACT

Contemporary motor control theories propose competition between multiple motor plans before the winning command is executed. While most competitions are completed prior to movement onset, movements are often initiated before the competition has been resolved. An example of this is saccadic averaging, wherein the eyes land at an intermediate location between two visual targets. Behavioral and neurophysiological signatures of competing motor commands have also been reported for reaching movements, but debate remains about whether such signatures attest to an unresolved competition, arise from averaging across many trials, or reflect a strategy to optimize behavior given task constraints. Here, we recorded electromyographic activity from an upper limb muscle (m. pectoralis) while twelve (8 female) participants performed an immediate response reach task, freely choosing between one of two identical and suddenly presented visual targets. On each trial, muscle recruitment showed two distinct phases of directionally-tuned activity. In the first wave, time-locked ~100 ms of target presentation, muscle activity was clearly influenced by the non-chosen target, reflecting a competition between reach commands that was biased in favor of the ultimately chosen target. This resulted in an initial movement intermediate between the two targets. In contrast, the second wave, time-locked to voluntary reach onset, was not biased toward the non-chosen target, showing that the competition between targets was resolved. Instead, this wave of activity compensated for the averaging induced by the first wave. Thus, single-trial analysis reveals an evolution in how the non-chosen target differentially influences the first and second wave of muscle activity.

SIGNIFICANCE STATEMENT

Contemporary theories of motor control suggest that multiple motor plans compete for selection before the winning command is executed. Evidence for this is found in intermediate reach movements towards two potential target locations, but recent findings have challenged this notion by arguing that intermediate reaching movements reflect an optimal response strategy. By examining upper limb muscle recruitment during a free-choice reach task, we show early recruitment of a sub-optimal averaged motor command to the two targets that subsequently transitions to a single motor command that compensates for the initially averaged motor command. Recording limb muscle activity permits single-trial resolution of the dynamic influence of the non-chosen target through time.

INTRODUCTION

The environment offers multiple action opportunities, but ultimately only one action can be selected. Classic decision-making theories assume a two-stage process, where the brain selects an appropriate action, and then plans and executes the desired motor commands (Donders, 1969; McClelland, 1979). However, neurophysiological studies have suggested that multiple potential motor plans can be concurrently encoded and compete for selection within brain regions involved in eye (Christopoulos et al., 2018; McPeek and Keller, 2004; Port and Wurtz, 2003) or reach movements (Cisek and Kalaska, 2005; Klaes et al., 2011). Competition may also influence behavioral output. For example, when free to look to either one of two suddenly appearing visual targets, participants sometimes look to an in-between position (Chou et al., 1999; Findlay, 1982). Because such saccadic averaging is most prominent for short-latency saccades (Ottes et al., 1984; Walker et al., 1997), it is thought that target representations initially compete for selection, before resolving into a final decision (Kim and Basso, 2008; McPeek and Keller, 2002).

There is currently debate on whether reach motor plans can also be represented concurrently. Recent neurophysiological results (Dekleva et al., 2018) suggest that apparent concurrent encoding of multiple reach plans (Cisek and Kalaska,2005) may arise from averaging neural activity across many trials; while the represented alternative can vary trial-to-trial, only one alternative is represented on any given trial. Reach trajectories intermediate between two alternatives have been observed in 'go-before-you-know' paradigms (Chapman et al., 2010), in which reach movements start before the 'correct' target is unveiled. Such intermediate reaches have been ascribed to averaging of competing reach plans (Gallivan et al., 2017; Stewart et al., 2014; Enachescu et al., 2021), or to strategic optimization of success given task constraints (Haith et al., 2015; Hudson et al., 2007; Wong and Haith, 2017; Alhussein and Smith, 2021).

For reaching, target competition studies often impose a delay between target presentation and movement initiation or target identification (as in the 'go-before-you-know' paradigm). This approach differs from the immediate and free response paradigms that elicit saccadic averaging. Here, we employ an immediate response paradigm to show competition between potential reach targets at the individual trial level, studying humans reaching in a *free-choice, double-target task* (**Fig. 1b,c**). Unlike the 'go-before-you-know' paradigm, there is no correct target, and nothing is gained by strategically aiming between the two targets. We recorded electromyographic (EMG) activity (m. pectoralis) and analyzed timing and magnitude of recruitment in response to target presentation.

For reaches to a single visual target, two waves of directionally-tuned EMG-activity have been observed (Glover and Baker, 2019; Pruszynski et al., 2010; Wood et al., 2015): an initial stimulus-locked response ~100 ms after visual target onset, which we refer to as an express visuomotor response (EVR, Contemori et al., 2021a), followed by a larger wave of EMG-activity predictive of the onset of the reach movement (MOV, ~200-300 ms after the EVR). Muscle recruitment during the EVR and MOV-interval are governed by different processes. For example, the EVR is directed towards the stimulus location even during anti-reaches (Gu et al., 2016), is only influenced by the implicit but not explicit component of motor learning (Gu et al., 2019), and depends on stimulus properties (Glover and Baker, 2019; Kozak et al., 2019; Wood et al., 2015; Kozak and Corneil, 2021) and cueing (Contemori et al., 2021b). In contrast, MOV-epoch activity reflects the actual reach kinematics, but is not influenced by stimulus properties.

Examining EMG-activity in these intervals during free choice, double-target reaching, suggests a dynamically evolving interaction between the chosen and non-chosen target. During the EVR-interval, the non-chosen target influences muscle recruitment, revealing averaging that

109 is biased in favor of the ultimately selected target. This averaging produces a subtle attraction of

110 early reach kinematics to the non-chosen target. Subsequently, this non-chosen target influence

111 yields into a goal-directed motor command during the MOV-interval, compensating for the

112 earlier subtle attraction by bowing the reach trajectory away from the non-chosen target.

MATERIALS AND METHODS

Participants and Procedures

The experiment was conducted with approval from the institutional ethics committee from the

Faculty of Social Sciences at Radboud University Nijmegen, The Netherlands. Twelve

participants (eight females and four males), between 18 and 33 years of age (mean $\pm$SD = 24 $\pm$

5), gave their written consent prior to participating in the experiment. Three participants (one

female and two males) were self-declared left-handed, while the remaining participants were

self-declared right-handed. All participants were compensated for their time with either course

credits or a monetary payment and they were free to withdraw from the experiment at any time.

All participants had normal or corrected-to-normal vision and had no known motor impairments.

Reach Apparatus and Kinematic Acquisition

Participants were seated in a chair in front of a robotic rig. The participant's right arm was

supported by an air-sled floating on top of a glass table. All participants performed right-handed

horizontal planar reaching movements while holding the handle of the planar robotic

manipulandum (vBOT, Howard et al., 2009). The vBOT measured both the x- and y-positions of

the handle at a 1 kHz sampling rate. Throughout the whole experiment a constant load of 5 N in

the rightward direction, relative to the participant, was applied to increase the baseline activity

for the right pectoralis muscle (see below). All visual stimuli were presented within the plane of

the horizontal reach movements via a mirror, which reflected the display of a downward facing

LCD monitor (Asus – model VG278H, Taipei, Taiwan). The start position and the peripheral

visual targets were presented as white circles (0.5 and 1.0 cm radii, respectively) onto a black

background. Real-time visual feedback of the participant's hand position was given throughout

the experiment and was represented by a yellow cursor (0.25 cm in radius). Vision of the

physical arm, hand and manipulandum was occluded by the mirror.

EMG Acquisition

EMG-activity was recorded from the clavicular head of the right pectoralis major (PEC) muscle

using wireless surface EMG electrodes (Trigno sensors, Delsys Inc., Natick, MA, USA). The

electrodes were placed ~1 cm inferior to the inflection point of the participant's right clavicle.

Concurrent with the EMG recordings, we also recorded a photodiode signal that indicated the

precise onset of the peripheral visual targets on the LCD screen. Both the EMG and photodiode

signals were digitized and sampled at 1.11 kHz.

Experimental Paradigm

Each trial began with the onset of the start position located at the center of the screen, which was

also aligned with the participant's midline. Participants had to move their cursor into the start

position and after a randomized delay period (1-1.5 s) either one (*Single Target*, 25% of all trials,

Fig. 1a) or two peripheral targets appeared (*Double Targets*, 75%, **Fig. 1b,c**). All peripheral

targets were presented 10 cm away from the start position and at one of 12 equally spaced

locations around the start position (dotted circles in **Fig. 1a**). The onset of the peripheral targets

occurred concurrently with the offset of the start position. Participants were explicitly instructed

to reach as fast as possible towards one of the peripheral target locations during *Double Target*

trials. To ensure that the participants reached as fast as possible, the peripheral targets turned red

if the cursor had not moved out of the start position within 500 ms after the onset of the

peripheral targets. The trial ended as soon as the cursor entered one of the peripheral targets.

Note that it is highly unlikely and suboptimal for participants to make anticipatory movements,

given the high degree of spatial uncertainty of where targets would appear across trials.

For every participant, the experiment consisted of eight blocks, each block contained 240

trials, with 60 *Single Target* and 180 *Double Target* trials, pseudo-randomly interleaved. For the

Double Target trials, the two targets appeared either 60°, 120°, or 180° apart in equal likelihood.

Each possible single and double target configuration was presented five times in every block,

resulting in 40 repeats over the whole experiment. This design is expected to average out any

trial history effects.

Data Analyses

All data were analyzed using custom-written scripts in Matlab (version R2014b, Mathworks Inc.,

Natick, MA, USA). For both the 60° and 120° *Double Target* trials, we sorted trials based on

whether the final reach was directed to either the CW (**Fig. 1b**, red arrow) or CCW target

location (blue arrow). Thus, for all CW and CCW reach trials the non-chosen target location was

in the CCW and CW direction, respectively. Trials from the 180° *Double Target* condition

cannot be sorted in this way, since the non-chosen target location was always 180° away.

Reach onset detection and initial reach error: Reach onset was identified as the first time-point

after the onset of the peripheral targets at which the hand speed exceeded 2 cm/s. Reach reaction

time (RT) was calculated as the time between the onset of the peripheral targets and the initiation

of the reach movement. Initial reach direction was quantified as the angular direction of the

vector between the start position and the location of the hand at the time of reach onset. From

this, initial reach error was defined as the angular difference between the initial reach direction

and the direction of the final chosen target.

EMG processing and trial inclusion criteria: All EMG data were first rectified and aligned to

both the onset of the peripheral targets (as measured by the onset of the photodiode) and reach

initiation. To account for the difference in EMG recordings across the participants, we first

normalized EMG-activity for each participant by dividing against their own mean baseline

activity (i.e. mean EMG-activity over the 40 ms window prior to the stimuli onset). We then log-

normalized each participant's EMG-activity to account for the non-linearity of EMG-activity.

This normalization transforms the distribution of EMG values from an exponential distribution,

with many values close to zero and few large values, into a normal distribution. We specifically

examined two distinct epochs of EMG-activity: (1) The initial EVR, that is evoked 85-125 ms

after the onset of the visual stimuli (Gu et al., 2016, 2018, 2019), and (2) the later movement-

related response (MOV, -20 to 20 ms around reach initiation) associated with reach onset. To

prevent any overlap between these two different epochs (Gu et al., 2016, 2019; Kozak et al.,

2019), we excluded all trials with RTs less than 185 ms (~7% all trials). We also excluded with

RTs greater than 500 ms (<0.1% of all trials).

Receiver-Operating Characteristic Analysis: As done previously (Corneil et al., 2004;

Pruszynski et al., 2010), we used a time-series receiver-operating characteristic (ROC) analysis

to quantitatively detect the presence of an EVR. To do this, we first separated leftward (target

locations between 120° and 240° from straight right) and rightward (-60° to 60°) *Single Target*

trials. For each timepoint from 100 ms before to 300 ms after target onset, we calculated the area

under the ROC curve between the EMG-activity for leftward compared to rightward trials. This

metric indicates the probability that an ideal observer could discriminate the target location based

solely on the distribution of EMG-activity at that given timepoint. A value of 0.5 indicates

chance discrimination, whereas a value of 1 or 0 indicates perfect correct or incorrect

discrimination, respectively. We set the threshold for discrimination at 0.6, as this criterion

exceeds the 95% confidence intervals for EMG data that has been randomly shuffled through a

bootstrapping procedure (Chapman and Corneil, 2011). The discrimination time was defined as

the first timepoint after target onset at which the ROC metric was above 0.6 and remained above

that threshold for at least five out of the next 10 timepoints. We defined any participant with a

discrimination time less than 125 ms as a participant exhibiting a EVR. Based on this criterion,

11 of the 12 participants had a detectable EVR. All subsequent analyses were done on the 11

participants with an EVR.

Directional tuning of EMG-activity: We assumed cosine tuning (**Eq. 1**) between the log-

normalized EMG-activity and the chosen target location for both the EVR and MOV-epochs:

$$EMG(x) = A \times cos(x - \theta) \tag{1}$$

in which x is the chosen target location in degrees, starting CCW from straight right; $EMG(x)$

is the log-normalized EMG-activity for the given target location; A is the amplitude of the cosine

tuning; and θ is the preferred direction (PD) of the EMG-activity. We used Matlab's curve

fitting toolbox *fit* function to estimate both the A and θ parameters. We constrained our search

parameters such that $A > 0$ and $0° \leq \theta < 360°$. The initial search parameters were $A = 1$

and $\theta = 180°$. PDs of $0°$ and $180°$ would represent straight rightward and leftward,

respectively.

Model predictions: Previous studies have proposed different models of how the brain converts

multiple visual targets into a single motor command. Here we assumed a constant non-linear

cosine tuning between target locations and motor commands in *Single Target* trials to generate

the predicted responses during *Double Target* trials. Each model used parameters derived from

each participant's own *Single Target* data (**Fig. 2a**) to predict both the PD and amplitude of the

cosine tuning curves for *Double Target* trials. Thus, no free parameters were fitted in any of

these four models.

Model 1: The *winner-take-all model* (**Fig. 4a**) assumes that only the target location that

the participant reaches towards is converted into a motor command. Therefore,

$EMG(x_1|x_1, x_2) = EMG(x_1)$, where x_1 and x_2 are the chosen and non-chosen target

locations, respectively.

Model 2: The *spatial averaging model* (**Fig. 4b**) assumes that the two potential target

locations are first spatially averaged into an intermediate target location. Then that target location

is converted into a motor command. Therefore, $EMG(x_1|x_1, x_2) = EMG\left(\frac{x_1+x_2}{2}\right)$.

Model 3: The *motor averaging model* (**Fig. 4c**) assumes that the two potential target

locations are first converted into their own distinct motor commands and then averaged into a

single motor command. Therefore, $EMG(x_1|x_1, x_2) = 0.5 \times EMG(x_1) + 0.5 \times$

$EMG(x_2)$.

Model 4: The *weighted motor averaging model* (**Fig. 4d**) is a variation of the motor

averaging model. It assumes that the two target locations are first converted into their associated

motor commands, which are then differentially weighted before being averaged into a single

motor command. A higher weight is assigned to the chosen target compared to the non-chosen

target location. To estimate these weights indirectly, we used each participant's own *Single*

Target data. Previous studies have shown that the EVR magnitude is negatively correlated with

the ensuing RT for single target visually-guided reaches (Pruszynski et al., 2010; Gu et al.,

2016). We assumed that the trial-by-trial magnitude of the EVR reflected the 'readiness' to move

towards the target location. Thus, we performed a median RT split of the *Single Target* data to

get cosine tuning for both *Fast* and *Slow RT* trials (**Fig. 2a**). This results in *Fast RT* and *Slow RT*

amplitude and PD estimates, which were used to compute the tuning curves for the *Double*

Target trials. The *Fast RT* and *Slow RT* parameters were used for the chosen and non-chosen

target location, respectively. Therefore, $EMG(x_1|x_1, x_2) = 0.5 \times EMG_{Fast}(x_1) +$

$0.5 \times EMG_{Slow}(x_2)$.

To quantify the goodness-of-fit for each model, due to the non-linear interaction between

PD and normalized amplitude, we evaluated the total fit error between the predicted and

observed tuning curves. To do this, we took the sum of mean squared error for each of the 12

different reach directions (i.e. $x_1 = 0°, 30°, 60°, … 330°$) between the predicted and observed

tuning curves.

Statistical Analyses

Statistical analyses were performed using either one or two-sample *t*-tests or a one-way

ANOVA. For all post-hoc comparisons, we used a Tukey's HSD correction. The statistical

significance was set as $P < 0.05$. For the model comparison, significance was set at $P < 0.0083$,

Bonferroni corrected for the six possible comparisons between the four different models.

RESULTS

Under continuous EMG recording of the right PEC muscle, participants performed a free-choice

goal-directed center-out right-handed reach movement in response to the onset of either one (**Fig.**

1a, *Single Target*) or two visual targets (**Fig. 1b** and **c**, *Double Target* trials) that appeared

concurrently. The visual targets pseudo-randomly appeared at 12 different possible directions

equally spaced around the start position. For *Double Target* trials, the two visual stimuli had an

angular separation of either 60°, 120°, or 180°. Choice probability for the clockwise or

counterclockwise target in *Double Target* did not differ from 0.5 for both the 60° and 120° target

separation ($P_{CW,60} = 0.52 \pm 0.04; p = 0.06$ and $P_{CW,120} = 0.51 \pm 0.02; p = 0.13$).

Prior to examining the *Double Target* trials, we will first describe PEC EMG-activity

during the *Single Target* trials. **Figure 1a** shows the individual (middle panel) and mean log-

normalized EMG-activity (right) during left-outward (orange trace), straight outward (gray), and

right-outward *Single Target* trials (green) from a representative participant. All trials are aligned

to visual target onset and the individual trials were sorted based on the reach RTs (color squares).

Note, the increase and decrease of activity for the right PEC muscle for left-outward and right-

outward reach movement, respectively. Consistent with previous studies (Pruszynski et al., 2010;

Wood et al., 2015; Glover and Baker, 2019), we observed a reliable difference in EMG-activity

for the three different reach directions at two epochs: an initial EVR-epoch that occurs ~100 ms

after stimulus onset and a later MOV-epoch associated with reach RT (stochastically occurring

~200 ms after stimulus onset). Across our participants, the mean ±SEM discrimination time (see

Materials and Methods) for the EVR was 88 ±3 ms and the corresponding reach RT was 232 ±

3 ms. We calculated the EVR magnitude for a given trial as the mean log-normalized EMG-

activity during the EVR-epoch, 85-125 ms after stimulus onset (Gu et al., 2018, 2019) indicated

by the white dashed boxes and shaded panels in **Figure 1**. For this participant, we found a

reliable increase and decrease in EVR magnitude for left-outward and right-outward trials,

respectively, when compared to straight outward trials (1-way ANOVA, $F_{(2,105)} = 37.4$, $P < 10^{-12}$,

post-hoc Tukey's HSD, both $P < 0.001$).

Having established the profile of EMG-activity during the EVR-epoch on *Single Target*

trials, we next examined if the presence of a second non-chosen target during the *Double Target*

trials changed the EVR. For a direct comparison with **Figure 1a**, we first examined trials with

the same reach direction (i.e. straight outward) but with a different non-chosen target location

(60° CW, blue, or 60° CCW, red, from the target, **Figure 1b**). If the non-chosen target location

has no influence (i.e. no averaging), we would predict that the EVR magnitude resembles that

observed during outward reach movement during *Single Target* trials, which we overlaid in gray

in **Figure 1b**. Despite the same reach direction, we observed both an increase and a decrease of

EMG-activity during EVR-epoch for *Double Target* trials relative to the *Single Target* trials (1-

way ANOVA, $F_{(2,83)}$ = 16.2, $P < 10^{-5}$, post-hoc Tukey's HSD, $P = 0.01$ and $P = 0.004$,

respectively) when the non-chosen target was in the left-outward and right-outward locations,

respectively. This result suggests that EMG-activity during the EVR is systematically altered by

the presence of a second non-chosen target.

A second way to examine the EVR during *Double Target* trials is to compare EVR

magnitude on trials with the same two visual targets, but different reach directions. **Figure 1c**

shows the EMG-activity when both the left-outward (blue) and right-outward (red) targets were

presented to the representative participant. If the EVR averaged the locations of the two visual

targets completely, then we would predict that the resulting EMG-activity would not differ

regardless of the final reach direction. However, we observed a reliable difference in the EVR,

with it being slightly larger when the participant chose the left-outward versus right-outward

target (1-way ANOVA, $F_{(2,72)}$ = 7.06, $P = 0.002$, post-hoc Tukey's HSD, $P = 0.01$). This result

suggests that EMG-activity during the EVR is modulated by the chosen reach direction, even

when the same two visual targets are presented.

Systematic shifts in tuning of the EVR during Double Target trials

The results from **Figure 1b** and **c** demonstrate that the magnitude of the EVR during *Double*

Target trials depended on both the target configuration and the eventual reach direction. To

quantify the extent of averaging that occurred, we sought to compare how the directional tuning

of the EVR changed between *Single* and *Double Target* trials. Previously, it has been shown that

the log transformed EVR magnitude can be described by a cosine tuning function (Gu et al.,

2019; **Eq. 1**). For each tuning function we can extract both the preferred direction (PD) and the

amplitude of the fit. **Figure 2a** shows both individual trial data (dots) and the cosine tuning fit

(line) for the *Single Target* trials from the representative participant in **Figure 1a**. The PD of this

fit was 173° CCW (arrow) from straight rightward, indicating that the largest EVR magnitude

could be evoked by a visual target presented straight leftward of the start position. Importantly,

this cosine tuning between EVR magnitude and target location was not simply due to movement-

related EMG-activity from trials with the shortest RTs, as this relation was still present when we

performed a median RT split and re-fitted the data on either *Fast RT* (**Figure 2b**, dark line) or

Slow RT trials (light), separately. Across our participants, we found no systematic difference in

the PDs between *Fast* and *Slow RT* trials (**Figure 2c**, group mean ±SEM: PD = 169°± 3° and

162° ± 5°, respectively, paired *t*-test, $t_{(10)} = 1.30$, $P = 0.22$). We did find larger amplitudes (i.e.

larger EVR magnitudes) for *Fast* compared to *Slow RT* trials (**Figure 2d**, paired *t*-test, $t_{(10)} =$

7.89, $P < 10^{-4}$), which is consistent with previous studies demonstrating a negative correlation

between EVR magnitudes and RTs on a trial-by-trial basis (Pruszynski et al., 2010; Gu et al.,

2016). Note, we will leverage this relationship later in the modeling portion of the **RESULTS**.

We next fitted the EVR cosine tuning for the *Double Target* trials. For this we chose to

align the trials based on the participant's reach direction (**Figure 1b**) rather than controlling for

the visual target locations (**Figure 1c**) to accentuate the effect of the non-chosen target location.

Figure 3a shows the fits for the three different angular separations for the representative

participant. For both the 60° and 120° conditions, we generated two separate fits for when the

non-chosen target location was either CW (red) or CCW (blue) relative to the reach direction. To

give more intuition of how this figure relates to individual trials, the highlighted data (shaded

box in the left panel of **Figure 3a**) corresponds to the same trials as **Figure 1b**. The right panel

of **Figure 3a** shows the fit of EVR magnitude to the 180° *Double Target* condition. Note, the

data cannot be split because the non-chosen target location is always 180° away from the reach

direction. Despite the two targets being in diametrically opposite directions, the EVR was still

reliably tuned for the 180° condition ($r^2 = 0.19$, for this participant). Across participants, the

directional tuning of the EVR during the 180° *Double Target* trials was not reliably different

from that observed in the *Single Target* trials (paired *t*-test, $t_{(10)} = 1.92$, $P = 0.08$), although we

did find a systematic decrease in the amplitude of the fits (see below).

For both the 60° and 120° conditions, since we aligned our data relative to the final reach

direction, the only difference between CW and CCW trials was the non-chosen target location. If

the EMG-activity was the result of a perfect averaging between the two target locations, then we

would predict the difference in PD between CW and CCW trials (ΔPD) to be equal to the angular

separation between the two targets (i.e. ΔPD = 60° and 120°, respectively). If the EMG-activity

was only influenced by the chosen target direction, then we would predict no difference between

CW and CCW conditions (ΔPD = 0°). Consistent with the individual trial data from **Figure 1b**,

we observed signs of averaging, albeit incomplete, for the representative participant for both the

60° and 120° conditions, with ΔPDs of 49.3° and 53.0°, respectively (**Figure 3a**).

We found similar results of partial averaging across our participants for both the 60°

(**Figure 3b**, left panel, mean ±SEM, ΔPD = 38.6° ± 3.5°, one sample *t*-test against zero, $t_{(10)} =$

10.9, $P < 10^{-6}$) and 120° *Double Target* conditions (ΔPD = 47.2° ± 5.4°, one sample *t*-test, $t_{(10)} =$

8.7, $P < 10^{-5}$). To fairly compare the extent of averaging between the conditions, we converted

the ΔPD into an averaging ratio (**Figure 3b**, right panel): a value of 1 indicates complete

averaging (ΔPD = 60° and 120°, dashed lines) and a value of 0 indicates no averaging (ΔPD =

0°). Overall, we found that the extent of averaging decreases as the angular separation increased

from 60° to 120° (averaging ratio = 0.6 ± 0.06 and 0.39 ± 0.05 a.u., respectively, paired *t*-test,

$t_{(10)} = 3.81$, $P = 0.003$).

In addition to the changes in PD of the EVR tuning, we also quantified the changes in the

amplitude during *Double Target* trials. **Figure 3c** shows the mean amplitude for the three

conditions, normalized to each participant's own *Single Target* amplitude as a baseline. We

observed a systematic decrease in amplitude as a function of angular separation: 1.13 ± 0.04,

0.88 ± 0.04, and 0.63 ± 0.05 a.u. for the 60°, 120°, and 180° conditions, respectively (repeated

measures 1-way ANOVA, $F_{(2,20)} = 41.1$, $P < 10^{-7}$, post-hoc paired *t*-test, all $t_{(10)} > 5.5$, $P < 10^{-3}$).

The systematic changes in PD and amplitude will be interpreted based on different possible

averaging models tested below.

Model predictions of EMG-activity during the EVR-epoch for Double Target trials

Previous studies examining averaging behavior for both eye and reach movements have

proposed different models for how the two visual targets may be integrated into a single motor

command. These models make distinct predictions for how the PD and amplitude of the tuning

curves should change between *Single* and *Double Target* trials (see **Materials and Methods** for

exact details). **Figure 4,** right column, shows the predicted tuning curves generated from the four

different proposed models for both the 120° CW and CCW conditions, using the *Single Target*

data (dashed gray line) from the representative participants. **Model 1** is the *winner-takes-all*

model (**Figure 4a**), which proposes that the two visual targets compete for selection in a winner-

takes-all process, resulting in a motor command that is generated towards the winning target

location (Donders, 1969; McClelland, 1979). Effectively, there is no integration between the two

target locations at any stage of the process. Note this model is agnostic about whether the

competition for selection occurs at either a spatial or motor representation. **Model 2** is the *spatial*

averaging model (**Figure 4b**), which proposes that the two targets are first averaged into a spatial

representation, resulting in a motor command towards the intermediate spatial direction (Findlay,

1982; Glimcher and Sparks, 1993; Walker et al., 1997; Chou et al., 1999). **Model 3** is the *motor*

averaging model (**Figure 4c**), which proposes that the two targets are first converted into two

independent motor commands (Edelman and Keller, 1998; Port and Wurtz, 2003; Cisek and

Kalaska, 2005) and then averaged into a single motor command (Katnani and Gandhi, 2011;

Stewart et al., 2014; Gallivan et al., 2017). Finally, **Model 4** is the *weighted motor averaging*

model (**Figure 4d**), which is a variation of the motor averaging model. Once again, the two

targets are first converted into two separate motor commands, but a stronger weighting is given

towards the chosen compared to the non-chosen target location (Kim and Basso, 2008, 2010;

Pastor-Bernier and Cisek, 2011). The final motor command is then an average of these two

differentially weighted motor commands. This model can be conceptualized as a race between

two accumulators (Schall, 2001, Enachescu et al., 2021), with the eventual chosen target location

accumulating at a faster rate compared to the non-chosen target location. Instead of fitting the

weights of the chosen and non-chosen target locations, we decided to indirectly estimate them by

using the *Fast* and *Slow RT* tuning curves from the *Single Target* trials, respectively (**Figure 2b**).

Previous studies (Pruszynski et al., 2010; Gu et al., 2016, 2018) have linked trial-by-trial EVR

magnitude to the 'readiness' of the motor system towards a specific target location. Here, we

assumed that during *Double Target* trials the motor system reaches towards the more 'ready'

target location.

Weighted motor averaging model best explains EVR-related EMG-activity

Figure 5a and **b** summarize the four different model predictions (color lines) for both the ΔPD averaging ratio and normalized amplitude fits across the three different *Double Target* angular separation conditions relative to *Single Target* trials. The *winner-takes-all model* predicted no change in either ΔPD (i.e. averaging ratio = 0 a.u.) or amplitude (i.e. normalized amplitude = 1 a.u.). Both the *spatial* and *motor averaging models* predicted complete averaging (averaging ratio = 1 a.u.) for both the 60° and 120° conditions. The key difference between the two models was in the predicted amplitude, where the *spatial averaging model* predicted no change (amplitude = 1 a.u.), while the *motor averaging model* predicted a systematic decrease (amplitude < 1 a.u.). Finally, the *weighted motor averaging model* predicted both a partial averaging (0 < averaging ratio < 1) and a decrease in amplitude. The extent of these changes depended on each participant's own *Fast* and *Slow RT* fits.

 Figure 5a and **b** also show our observed group data (open bars) plotted against the predictions from the four models during the EVR-epoch. Note only the *weighted motor averaging model* (green lines) captured both the systematic decrease in averaging ratio and amplitude that was in the observed data. Since the parameters of all four models were derived from each participant's own *Single Target* trials and contained no free parameters, we can directly compare the four different models. **Figure 5c** illustrates the mean ± SEM of the fit error between the observed and predicted fits across the participants. We found that the *weighted motor averaging model* best predicted the observed tuning curves compared to the other three models during the EVR-epoch (repeated measures 1-way ANOVA, $F_{(3,30)} = 7.7$, $P < 10^{-3}$, post-hoc paired *t*-test, $t_{(10)} = 3.6$, 4.1, and 4.8, $P = 0.005$, 0.002, and 0.0001, compared to the *winner-takes-all, spatial,* and *motor averaging models*, respectively).

Winner-takes-all model best explains MOV-related EMG-activity

Up to this point, we have only examined the initial wave of EMG-activity time-locked to the onset of the two visual targets (i.e. during the EVR-epoch). Are there also signatures of averaging in the tuning of EMG-activity associated with movement onset (MOV-epoch) in the *Double Target* trials? We therefore examined EMG activity during the MOV-epoch, i.e. mean EMG-activity -20 to 20 ms around reach onset.

Figure 6a shows the EMG-activity during the MOV-epoch for individual trials for our exemplar participant, centered at the chosen target direction and split by the direction of the non-chosen target and the three target separations. On top the cosine tuning curves are shown. For this subject the amplitudes do not differ between target separations or non-chosen target direction. However, small shifts in preferred direction, away from the non-chosen target, can be observed for the 60° and 120° target separation.

Figure 6b and **c** show both the averaging ratio and amplitude across participants, based on fits to the EMG-activity during the MOV-epoch. Unlike the EVR-epoch, the *winner-takes-all model* best predicted EMG-activity around reach onset (**Fig. 6d**, repeated measures 1-way ANOVA, $F_{(3,30)} = 348.8$, $P < 10^{-22}$, post-hoc paired *t*-test, $t_{(10)} = 28.6, 21.8, 29.0$, all $P < 10^{-9}$, compared to the *spatial, motor,* and *weighted motor averaging models*, respectively). Although the *winner-takes-all* model provides the best explanation for our MOV-epoch data, we still observed an influence of the non-chosen target location with an averaging ratio shifting in the opposite direction, suggesting a *repulsion* from the non-chosen target location (averaging ratio = –0.13 ± 0.03 and –0.11 ± 0.01 a.u., for 60° and 120° *Double Target* trials, respectively, one sample *t*-test against zero, $t_{(10)} = -4.2$ and –11.7, both $P < 0.05$, **Fig. 6b**). However, in the next

section we will argue that this is not a genuine repulsion from the non-chosen target, but rather

compensation for the earlier attraction by the non-chosen target in the EVR-epoch.

Early kinematics show attraction to the non-chosen target location

Having established an opposite influence of the non-chosen target on the tuning of EMG-activity

during the EVR and MOV-epoch, we next determined whether the brief burst of muscle

recruitment during the EVR-interval carried any behavioral consequences. **Figure 7a** shows the

representative participant's initial reach error (i.e. the difference between the chosen target

location and the initial reach direction at the time of reach onset) for both *Single* and *Double*

Target trials. For the *Single Target* trials the distribution of initial reach direction is closely

centered on the actual target direction. However, for the *Double Target* trials the distributions of

initial reach direction are clearly shifted toward the non-chosen target. **Figure 7b** shows the

median initial reach error direction, averaged across participants. This initial reach error differed

significantly from zero for both target separations (Initial Reach Error = 15.2° ± 4.3° and 13.9°±

5.5°, paired *t*-test, $t_{(10)}$ = -11.7 and -8.4, both $P < 10^{-5}$, respectively). These initial reach errors

indicate an early attraction toward the non-chosen target, which is consistent with the averaging

of EMG-activity during the EVR-interval. Following this averaging during the EVR-interval, we

subsequently observed an opposite effect in the tuning of the EMG-activity in the MOV-epoch

for the *Double Target* trials. This is highlighted by a significant negative averaging ratio (**Figure**

6b). We surmise that this opposing effect corresponds to compensatory muscular activity that

corrects for the initial attraction of the arm toward the non-chosen target, bowing the arm back

toward the chosen target location (**Figure 7c).**

DISCUSSION

Contemporary theories of decision-making posit that multiple potential motor plans compete for selection (Cisek, 2007; Schall, 2001). Behavioral and neurophysiological results have shown such competition within the oculomotor system (Chou et al., 1999; Coren and Hoenig, 1971; Findlay, 1982; Ottes et al., 1984; Walker et al., 1997; Bhutani et al., 2012), but it is unclear whether such results generalize to reaching. Here, by measuring upper limb EMG during a reaching task that demands an immediate response, we demonstrate that the non-chosen target influences the earliest wave of muscle recruitment following target onset. Such evidence is apparent on single trials, implicating biased competition between the chosen and non-chosen target within upstream premotor areas soon after target appearance. This initial biased motor averaging affected the initial direction, but subsequently gave way to a goal-directed motor command that bowed the arm back onto a trajectory directed toward the chosen target.

The EVR is a trial-by-trial weighted average of motor commands

We tested different models of how the brain could have integrated the two visual targets and found that a *weighted-motor-averaging* model best explained partial averaging during the EVR-epoch. Such weighted-motor-averaging is apparent on a single trial and inconsistent with a recent interpretation that the apparent encoding of multiple alternatives in premotor cortex is caused by averaging of different alternatives across multiple trials (Dekleva et al., 2018). Instead, our result is consistent with contemporary theories for a deliberation process between multiple motor plans (Cisek, 2007; Schall, 2001; Enachescu et al., 2021). For example, previous neurophysiological studies have shown that experimentally manipulating the decision variable, via either target uncertainty (Basso and Wurtz, 1997; Dorris and Munoz, 1998), target

expectation (Basso and Wurtz, 1998; Bichot et al., 1996), or reward expectation (Pastor-Bernier

and Cisek, 2011; Rezvani and Corneil, 2008), modulates the neural representation of the

competing motor plans. Similarly, signatures of an evolving decision variable during deliberation

have been shown in the long latency reflex, when participants must indicate the direction of a

random-dot motion stimulus (Selen et al., 2012). Here, we exploited the fact that participants

randomly choose one of two visual targets and demonstrated post-hoc that the averaged EVR

was biased towards the chosen target. This suggests that either fluctuations along the

sensorimotor pathway (Faisal et al., 2008; Siegel et al., 2015) or idiosyncratic preferences based

on previous choices (Urai et al., 2019) biased both the initial EVR and the ultimate choice.

Interestingly, the influence of idiosyncratic preferences on the representation of alternatives was

also reported in premotor cortex (Dekleva et al., 2018).

While the weighted-motor-averaging model best explained the EVR, the fits were

imperfect (**Figure 5**). This is likely due to the arbitrary weighting of the EVR strength for the

chosen vs. non-chosen target, exploiting the inverse relationship between EVR magnitude and

reach RT (Gu et al., 2016; Pruszynski et al., 2010; Wood et al., 2015). We assume an

independent race between the motor programs to each of the two targets, with the one proceeding

faster (drawn from the shorter-than-average subset, having a larger EVR) 'winning' over the

non-chosen alternative (Rowe et al., 2010). This process is admittedly coarse, as it remains

unknown what the RT and EVR of the non-chosen alternative would have been. Regardless, only

the weighted-motor-averaging model captured the influence of the non-chosen target on both the

EVR tuning and magnitude (**Figure 5**); hence this model best captures the essence, if not the

magnitude, of the interaction between competing motor plans.

A *weighted-spatial-averaging* model, an extended version of **Model 2,** was not explicitly

evaluated. If we would allow the averaged target location to be somewhere between the

presented targets, instead of in the middle, the pre-motor circuitry would receive a 'go here'

signal that could change the shift of the tuning curve, but would not influence the amplitude of

the tuning. In contrast, our data show a systematic decrease in amplitude for the *Dual Target*

conditions, which can only be captured by the *weighted-motor-averaging* model.

Influence of task design on the EVR

Our results illustrate that different stages of decision-making influence distinct EMG

epochs in the motor periphery and thus suggest an influence of task design or stimulus properties

on these epochs. Indeed, the EVR is muted (Wood et al., 2015) or abolished (Pruszynski et al.,

2010) when a delay is imposed between stimulus presentation and movement onset.

Furthermore, the EVR is augmented when targets are temporally predictable (Kozak et al., 2020;

Contemori et al., 2021a). Given this, the EVR may be negligible or absent in delayed response

tasks (Cisek and Kalaska, 2005; Dekleva et al., 2018; Thura and Cisek, 2014), or in *'go-before-*

you-know' tasks introducing a delay between presentation of alternatives and initiation of the

reach. When an immediate response is required in the *'go-before-you-know'* paradigm, the

intermediate reaching movements skew toward the more salient stimulus (Wood et al., 2011),

paralleling the observation of earlier and larger-magnitude EVRs evoked by high-contrast (Wood

et al., 2015; Kozak and Corneil, 2021) or low-spatial frequency stimuli (Kozak et al., 2019).

The instruction to move rapidly reduces the production of intermediate reaches, possibly

due to adopting a control policy that maximizes task success (Wong and Haith, 2017). While the

impact of velocity instructions on the EVR is unknown, previous results suggest that the

magnitude, not timing, of the EVR would be modulated by changing control policy (Gu et al.,

2018, 2016). Furthermore, the EVR's short-latency makes the establishment of a control policy

after target presentation unlikely, but suggests a task dependent, preset control policy

implementing task instructions affecting relevant motor circuitry (Scott, 2016; Contemori et al.,

2022). Recently, Enachescu et al. (2021) provided a dynamic neural field model connected with

stochastic optimal feedback controllers. This model executes a weighted average of a continuum

of control policies for all possible reach directions, where competition and weighing of control

policies continues as the reach unfolds, consistent with the present findings.

Kinetic consequences of the EVR

EMG recordings permit the resolution of a decision-making dynamic at a level that

would be difficult, if not impossible, to resolve based on kinematics alone. For example, while

EMG-activity during the EVR was biased *toward* the non-chosen target, EMG-activity during

the MOV-interval was biased *away* from the non-chosen target (**Fig. 6**). At first glance, opposite

directions of EMG recruitment in these intervals seems paradoxical. The forces consequent to the

brief and smaller-magnitude EVR are undoubtedly less than those developed closer to the time of

reach initiation. However, the EVR has behavioral consequences, generating small forces toward

a stimulus (Gu et al., 2016). We show that forces from the averaged EVR bias the initial reach

toward the non-chosen target, but subsequent EMG compensates for their trial-specific kinematic

consequences. This suggests that voluntary control mechanisms are rapidly informed about trial-

specific kinematic consequences of the averaged EVR, using this information for feedforward

adjustments of the voluntary EMG-activity. These adjustments occur within 100ms after the

onset of EVR and are unlikely to be driven by visual feedback of the cursor.   A similar fast

mechanism has been reported for stretch reflexes (Pruszynski et al., 2009). If the background

load to a muscle increases, the mono-synaptic short latency reflex increases, but adjustments in

later phases, as quick as 45 ms after perturbation onset, already compensate for the stronger

adjustment in the first phase.

A shared neural substrate with the saccadic system

Our task incorporated many task features used to elicit saccadic averaging (Chou et al.,

1999; He and Kowler, 1989), including the requirement for an immediate response. Most

experiments on saccadic averaging have not been designed to dissociate between averaging at

the spatial or motor level. However, by contrasting two task instructions ("look at the last

presented target" vs "look at the targets in order of presentation") in a double-step paradigm,

Bhutani et al., (2012) provided evidence that saccadic averaging also takes place at the level of

the motor plan. Saccade kinematics offer a straightforward readout of the temporal evolution of

decision-making, paralleling our observations for EVRs. For example, the transition from an

averaged to a goal-directed command between the EVR and MOV-epoch resembles the

observation that averaging is strongest for short-latency saccades (Chou et al., 1999; Walker et

al., 1997). Further, EVR averaging diminishes with increasing angular target separation,

resembling observations for saccadic averaging (Chou et al., 1999; Vokoun et al., 2014).

Saccadic averaging has been related to the initial representation and subsequent resolution of

competing saccade plans within superior colliculus (Edelman and Keller, 1998; Port and Wurtz,

2003; Vokoun et al., 2014). Superior colliculus is also a potential substrate for the EVR via the

tecto-reticulo-spinal pathway (Corneil and Munoz, 2014; Glover and Baker, 2019; Gu et al.,

2016; Pruszynski et al., 2010; Contemori et al., 2021a, Kozak and Corneil, 2021; Kozak et al.,

2019). Thus, circumstantial evidence suggests that saccadic averaging and EVR averaging on

upper limb muscles may have a common collicular substrate. This subcortical substrate for the

deliberation process would agree with findings that M1 and PMd are mainly involved in

commitment to a choice (Derosiere et al., 2019; Thura and Cisek, 2020), but not the competition

between alternatives. Future neurophysiological experiments should investigate the causal

structure between weighted averaging of the EVR and the commitment to a single goal-directed

reach.

In summary, we examined neuromuscular activity during a free-choice reaching task to

two targets. We found that, similar to saccadic averaging, the earliest motor command in the

reaching system attests to a still-unresolved competition between multiple distinct motor plans.

However, this competition is rapidly resolved and by the time of movement onset the motor

system generates a goal directed reach movement that compensates for the averaging observed in

the early trajectory.

AUTHOR CONTRIBUTIONS

Conceptualization – LPS, BDC, and WPM; Methodology – LPS; Investigation – LPS; Writing,

Original Draft – LPS; Writing, Review and Editing – LPS, BDC and WPM; Funding Acquisition

– WPM; Resources – LPS and WPM.

DECLARATION OF INTERESTS

The authors declare no competing interesting.

 FIGURE LEGENDS

Figure 1: EVR from a representative participant. A. Individual (middle panel) and mean ±

SEM (right) log-normalized EMG-activity from the right PEC muscle during left-outward

(yellow), straight outward (gray), and right-outward (green) *Single Target* reach trials (left

panel). All EMG- activity is aligned to the onset of the peripheral visual target (thick black

vertical line). For the middle panel, each row represents EMG-activity within a single trial and

trials were sorted based on reach RT (colored squares). Dashed white box and shaded area in the

individual and mean EMG plots represent the EVR-epoch (85-125 ms after stimulus onset). **B.**

EMG-activity for *Double Target* trials when matched for the same outward reach movement.

The non-chosen target was either 60° CW (blue) or CCW (red) of the reach target. Same layout

as **A**. **C.** EMG-activity for 120° *Double Target* trials for the same visual target layout, but

different chosen target directions. Same layout as **A**. * $P < 0.05$.

Figure 2: Directional tuning of the EVR during *Single Target* trials. A. Cosine tuning of log-

normalized EVR magnitude as a function of the target direction for *Single Target* trials from the

representative participant in **Figure 1**. Dots indicate each trial, the solid line indicates the fit, and

the arrow indicates the PD of the fit. **B.** The cosine tuning is maintained regardless of the ensuing

reach RT, same data as **a**. but re-fitted for *Fast* (black) and *Slow RT* (gray) *Single Target* trials

separately. For illustration purposes only, we have staggered the individual trial data to illustrate

the difference between the two conditions. We did not stagger the cosine tuning curves. **C. D.**

Group ($n = 11$) mean ± SEM for the PD (c) and amplitude (**d**) of the fits between the *Fast* and

Slow RT trials. Each gray line indicates an individual participant, and the darker line indicates the

representative participant. * $P < 0.05$.

Figure 3: Systematic changes in directional tuning of the EVR during *Double Target* trials.

A. Fits for 60°, 120° and 180° conditions of the *Double Target* trials with all data aligned to the chosen target direction from the representative participant. For both the 60° and 120° conditions, the trials were sorted based on whether the non-chosen target was either CW (blue) or CCW (red) of the chosen target direction. Data in the shaded panel indicates the trials from **Figure 1b**. **B.** Group mean ± SEM shifts in PD (ΔPD) between the CW and CCW trials (left panel) and the normalized averaging ratio (right) for both 60° and 120° conditions across our participants. Dashed box indicates the predicted ΔPD if the EVR would be a complete average of the two targets (averaging ratio = 1 a.u.). **C.** Mean ± SEM amplitude of the fits for the three different *Double Target* conditions across our participants. The amplitudes were normalized to each participant's own amplitude fit from the *Single Target* trials. Each grey line indicates a different participant. * $P < 0.05$.

Figure 4: Model predictions of the tuning curves during *Double Target* trials. A. The *winner-takes-all* model chooses one visual stimulus as the target and converts it into the final motor command. **B.** The *spatial averaging* model averages the two visual stimulus directions into an intermediate target direction and that target direction is converted into a motor command. **C.** The *motor averaging* model first converts the two visual stimuli into two separated motor commands. Then it averages the two motor commands into a single motor command. **D.** The *weighted motor averaging* model first converts the two visual stimuli into two separate motor commands, but the cosine tuning have different weights. Then it averages the two motor

commands into a single motor command. For the right column: red curves for CW chosen target,

blue curves for the CCW chosen target and dashed grey curve the single target tuning curve.

Figure 5: Comparisons of model predictions and observed group data for *Double Target*

trial fits. a, b. The model predictions (colored lines, see legend for color coding) overlaid over

the observed mean ± SEM group data (open black bars) for EMG-activity during the EVR-epoch

(85 to 125 ms after stimuli onset) for both the averaging ratio (**a**) and amplitude (**b**). **c.** The mean

± SEM group model fit errors for the four different models.

Figure 6: EMG-activity and tuning properties in the MOV-epoch (-20 to 20 ms around

reach RT). a. Fits for 60°, 120° and 180° conditions of the *Double Target* trials with all data

aligned to the chosen target direction from the representative participant. For both the 60° and

120° conditions, the trials were sorted based on whether the non-chosen target was either CW

(blue) or CCW (red) of the chosen target direction. **b, c.** The model predictions overlaid over the

observed mean ± SEM group data (open black bars) for EMG-activity during the MOV-epoch

for both the averaging ratio (**b**) and amplitude (**c**). **d.** The mean ± SEM group model fit errors for

the four different models. * $P < 0.0083$.

Figure 7: Systematic repulsion away from the non-chosen target direction at the time of

reach RT. a. Histogram of reach error direction, relative to the chosen target direction, at the

time of reach RT for the representative participant during the experiment. For *Double Target*

trials, the location of the non-chosen target direction is shown as colored circles along the *x*-axis.

Vertical lines represent the median reach errors. **b.** Mean ±SEM of difference in median reach

error between CW and CCW during *Double Target* trials. Dashed boxes represent full averaging,

i.e. predictions from model 2 and 3. **c.** Initial reach errors converge to the target direction while

the reach unfolds and the reach percentage (RP) increases. RP = 0% corresponds to hand speed

above 2cm/s and RP = 100% to covering the target distance. Across subjects and conditions this

corresponds to a time window of 270 ± 23ms.

REFERENCES

Alhussein, L., Smith, M.A., 2021. Motor planning under uncertainty. Elife 10, e67019. https://doi.org/10.7554/elife.67019

Basso, M.A., Wurtz, R.H., 1998. Modulation of Neuronal Activity in Superior Colliculus by Changes in Target Probability. J Neurosci 18, 7519–7534. https://doi.org/10.1523/jneurosci.18-18-07519.1998

Basso, M.A., Wurtz, R.H., 1997. Modulation of neuronal activity by target uncertainty. Nature 389, 66–69. https://doi.org/10.1038/37975

Bhutani, N., Ray, S. and Murthy, A., 2012. Is saccade averaging determined by visual processing or movement planning? J Neurophysiol 108, 3161–3171. https://doi.org/10.1152/jn.00344.2012

Bichot, N.P., Schall, J.D., Thompson, K.G., 1996. Visual feature selectivity in frontal eye fields induced by experience in mature macaques. Nature 381, 697–699. https://doi.org/10.1038/381697a0

Chapman, C.S., Gallivan, J.P., Wood, D.K., Milne, J.L., Culham, J.C., Goodale, M.A., 2010. Reaching for the unknown: Multiple target encoding and real-time decision-making in a rapid reach task. Cognition 116, 168–176. https://doi.org/10.1016/j.cognition.2010.04.008

Chapman, B.B., Corneil, B.D., 2010. Neuromuscular recruitment related to stimulus presentation and task instruction during the anti-saccade task: Neck muscle activity during an anti-saccade task. Eur J Neurosci 33, 349–360. https://doi.org/10.1111/j.1460-9568.2010.07496.x

Chou, I., Sommer, M.A., Schiller, P.H., 1999. Express averaging saccades in monkeys. Vision Res 39, 4200–4216. https://doi.org/10.1016/s0042-6989(99)00133-9

Christopoulos, V.N., Kagan, I., Andersen, R.A., 2018. Lateral intraparietal area (LIP) is largely effector-specific in free-choice decisions. Sci Rep-uk 8, 8611. https://doi.org/10.1038/s41598-018-26366-9

Cisek, P., 2007. Cortical mechanisms of action selection: the affordance competition hypothesis. Philosophical Transactions Royal Soc B Biological Sci 362, 1585–1599. https://doi.org/10.1098/rstb.2007.2054

Cisek, P., Kalaska, J.F., 2005. Neural Correlates of Reaching Decisions in Dorsal Premotor Cortex: Specification of Multiple Direction Choices and Final Selection of Action. Neuron 45, 801–814. https://doi.org/10.1016/j.neuron.2005.01.027

Contemori, S., Loeb, G.E., Corneil, B.D., Wallis, G., Carroll, T.J., 2021a. The influence of
temporal predictability on express visuomotor responses. J Neurophysiol 125, 731–747.
https://doi.org/10.1152/jn.00521.2020

Contemori, S., Loeb, G.E., Corneil, B.D., Wallis, G., Carroll, T.J., 2021. Trial-by-trial
modulation of express visuomotor responses induced by symbolic or barely detectable cues. J
Neurophysiol 126, 1507–1523. https://doi.org/10.1152/jn.00053.2021

Contemori, S., Loeb, G.E., Corneil, B.D., Wallis, G., Carroll, T.J., 2022. Symbolic cues enhance
express visuomotor responses in human arm muscles at the motor planning rather than the
visuospatial processing stage. J Neurophysiol 128, 494–510.
https://doi.org/10.1152/jn.00136.2022

Coren, S., Hoenig, P., 1971. Effect of Non-Target Stimuli upon Length of Voluntary Saccades.
Percept Motor Skill 34, 499–508. https://doi.org/10.2466/pms.1972.34.2.499

Corneil, B.D., Olivier, E., Munoz, D.P., 2004. Visual responses on neck muscles reveal selective
gating that prevents express saccades. Neuron 42, 831-41. https://doi.org/10.1016/S0896-
6273(04)00267-3

Corneil, B.D., Munoz, D.P., 2014. Overt Responses during Covert Orienting. Neuron 82, 1230–
1243. https://doi.org/10.1016/j.neuron.2014.05.040

Dekleva, B.M., Kording, K.P., Miller, L.E., 2018. Single reach plans in dorsal premotor cortex
during a two-target task. Nat Commun 9, 3556. https://doi.org/10.1038/s41467-018-05959-y

Derosiere, G., Thura, D., Cisek, P., Duque, J., 2019. Motor cortex disruption delays motor
processes but not deliberation about action choices. Journal of neurophysiology 122, 1566–
1577. https://doi.org/10.1152/jn.00163.2019

Donders, F.C., 1969. On the speed of mental processes. Acta Psychol 30, 412–431.
https://doi.org/10.1016/0001-6918(69)90065-1

Dorris, M.C., Munoz, D.P., 1998. Saccadic Probability Influences Motor Preparation Signals and
Time to Saccadic Initiation. J Neurosci 18, 7015–7026. https://doi.org/10.1523/jneurosci.18-
17-07015.1998

Edelman, J.A., Keller, E.L., 1998. Dependence on Target Configuration of Express Saccade-
Related Activity in the Primate Superior Colliculus. J Neurophysiol 80, 1407–1426.
https://doi.org/10.1152/jn.1998.80.3.1407

Enachescu, V., Schrater, P., Schaal, S., Christopoulos, V., 2021. Action planning and control
under uncertainty emerge through a desirability-driven competition between parallel encoding
motor plans. Plos Comput Biol 17, e1009429. https://doi.org/10.1371/journal.pcbi.1009429

Faisal, A.A., Selen, L.P.J., Wolpert, D.M., 2008. Noise in the nervous system. Nat Rev Neurosci
9, nrn2258. https://doi.org/10.1038/nrn2258

Findlay, J.M., 1982. Global visual processing for saccadic eye movements. Vision Res 22, 1033–
1045. https://doi.org/10.1016/0042-6989(82)90040-2

Gallivan, J.P., Stewart, B.M., Baugh, L.A., Wolpert, D.M., Flanagan, J.R., 2017. Rapid
Automatic Motor Encoding of Competing Reach Options. Cell Reports 18, 1619–1626.
https://doi.org/10.1016/j.celrep.2017.01.049

Glover, I.S., Baker, S.N., 2019. Multimodal stimuli modulate rapid visual responses during
reaching. J Neurophysiol 122, 1894–1908. https://doi.org/10.1152/jn.00158.2019

Gu, C., Pruszynski, J.A., Gribble, P.L., Corneil, B.D., 2019. A rapid visuomotor response on the
human upper limb is selectively influenced by implicit motor learning. J Neurophysiol 121,
85–95. https://doi.org/10.1152/jn.00720.2018

Gu, C., Pruszynski, J.A., Gribble, P.L., Corneil, B.D., 2018. Done in 100 ms: path-dependent
visuomotor transformation in the human upper limb. J Neurophysiol 119, 1319–1328.
https://doi.org/10.1152/jn.00839.2017

Gu, C., Wood, D.K., Gribble, P.L., Corneil, B.D., 2016. A Trial-by-Trial Window into
Sensorimotor Transformations in the Human Motor Periphery. J Neurosci 36, 8273–8282.
https://doi.org/10.1523/jneurosci.0899-16.2016

Haith, A.M., Huberdeau, D.M., Krakauer, J.W., 2015. Hedging Your Bets: Intermediate
Movements as Optimal Behavior in the Context of an Incomplete Decision. Plos Comput Biol
11, e1004171. https://doi.org/10.1371/journal.pcbi.1004171

He, P., Kowler, E., 1989. The role of location probability in the programming of saccades:
Implications for "center-of-gravity" tendencies. Vision Res 29, 1165–1181.
https://doi.org/10.1016/0042-6989(89)90063-1

Hudson, T.E., Maloney, L.T., Landy, M.S., 2007. Movement Planning With Probabilistic Target
Information. J Neurophysiol 98, 3034–3046. https://doi.org/10.1152/jn.00858.2007

Kim, B., Basso, M.A., 2008. Saccade Target Selection in the Superior Colliculus: A Signal
Detection Theory Approach. J Neurosci 28, 2991–3007.
https://doi.org/10.1523/jneurosci.5424-07.2008

Klaes, C., Westendorff, S., Chakrabarti, S., Gail, A., 2011. Choosing Goals, Not Rules: Deciding
among Rule-Based Action Plans. Neuron 70, 536–548.
https://doi.org/10.1016/j.neuron.2011.02.053

Kozak, R.A., Kreyenmeier, P., Gu, C., Johnston, K., Corneil, B.D., 2019. Stimulus-Locked
Responses on Human Upper Limb Muscles and Corrective Reaches are Preferentially Evoked

by   Low   Spatial   Frequencies.   Eneuro   ENEURO.0301-19.2019.
https://doi.org/10.1523/eneuro.0301-19.2019

Kozak, R.A., Cecala, A.L., Corneil, B.D., 2020. An Emerging Target Paradigm to Evoke Fast
Visuomotor   Responses   on   Human   Upper   Limb   Muscles.   J   Vis   Exp.
https://doi.org/10.3791/61428

Kozak, R.A., Corneil, B.D., 2021. High-contrast, moving targets in an emerging target paradigm
promote fast visuomotor responses during visually guided reaching. J Neurophysiol 126, 68–
81. https://doi.org/10.1152/jn.00057.2021

McClelland, J.L., 1979. On the time relations of mental processes: An examination of systems of
processes in cascade. Psychol Rev 86, 287–330. https://doi.org/10.1037/0033-295x.86.4.287

McPeek, R.M., Keller, E.L., 2004. Deficits in saccade target selection after inactivation of
superior colliculus. Nat Neurosci 7, 757–763. https://doi.org/10.1038/nn1269

McPeek, R.M., Keller, E.L., 2002. Saccade Target Selection in the Superior Colliculus During a
Visual   Search   Task.   J   Neurophysiol   88,   2019–2034.
https://doi.org/10.1152/jn.2002.88.4.2019

Ottes, F.P., Gisbergen, J.A.M.V., Eggermont, J.J., 1984. Metrics of saccade responses to visual
double   stimuli: Two   different   modes.   Vision   Res   24,   1169–1179.
https://doi.org/10.1016/0042-6989(84)90172-x

Pastor-Bernier, A., Cisek, P., 2011. Neural Correlates of Biased Competition in Premotor Cortex.
J Neurosci 31, 7083–7088. https://doi.org/10.1523/jneurosci.5681-10.2011

Port, N.L., Wurtz, R.H., 2003. Sequential Activity of Simultaneously Recorded Neurons in the
Superior   Colliculus   During   Curved   Saccades.   J   Neurophysiol   90,   1887–1903.
https://doi.org/10.1152/jn.01151.2002

Pruszynski, J.A., King, G.L., Boisse, L., Scott, S.H., Flanagan, J.R., Munoz, D.P., 2010.
Stimulus-locked responses on human arm muscles reveal a rapid neural pathway linking
visual   input   to   arm   motor   output.   Eur   J   Neurosci   32,   1049–1057.
https://doi.org/10.1111/j.1460-9568.2010.07380.x

Pruszynski, J.A., Kurtzer, I., Lillicrap, T.P., Scott, S.H., 2009. Temporal evolution of "automatic
gain-scaling". J Neurophysiol 102, 992–1003. https://doi.org/10.1152/jn.00085.2009

Rezvani, S., Corneil, B.D., 2008. Recruitment of a Head-Turning Synergy by Low-Frequency
Activity   in   the   Primate   Superior   Colliculus.   J   Neurophysiol   100,   397–411.
https://doi.org/10.1152/jn.90223.2008

Rowe, J.B., Hughes, L., Nimmo-Smith, I., 2010. Action selection: A race model for selected and non-selected actions distinguishes the contribution of premotor and prefrontal areas. Neuroimage 51, 888–896. https://doi.org/10.1016/j.neuroimage.2010.02.045

Schall, J.D., 2001. Neural basis of deciding, choosing and acting. Nat Rev Neurosci 2, 33–42. https://doi.org/10.1038/35049054

Scott, S.H., 2016. A Functional Taxonomy of Bottom-Up Sensory Feedback Processing for Motor Actions. Trends Neurosci 39, 512–526. https://doi.org/10.1016/j.tins.2016.06.001

Selen, L.P.J., Shadlen, M.N., Wolpert, D.M., 2012. Deliberation in the Motor System: Reflex Gains Track Evolving Evidence Leading to a Decision. J Neurosci 32, 2276–2286. https://doi.org/10.1523/jneurosci.5273-11.2012

Siegel, M., Buschman, T.J., Miller, E.K., 2015. Cortical information flow during flexible sensorimotor decisions. Science 348, 1352–1355. https://doi.org/10.1126/science.aab0551

Stewart, B.M., Gallivan, J.P., Baugh, L.A., Flanagan, J.R., 2014. Motor, not visual, encoding of potential reach targets. Curr Biol 24, R953–R954. https://doi.org/10.1016/j.cub.2014.08.046

Thura, D., Cisek, P., 2020. Microstimulation of dorsal premotor and primary motor cortex delays the volitional commitment to an action choice. Journal of neurophysiology 123, 927–935. https://doi.org/10.1152/jn.00682.2019

Thura, D., Cisek, P., 2014. Deliberation and Commitment in the Premotor and Primary Motor Cortex during Dynamic Decision Making. Neuron 81, 1401–1416. https://doi.org/10.1016/j.neuron.2014.01.031

Urai, A.E., Gee, J.W. de, Tsetsos, K., Donner, T.H., 2019. Choice history biases subsequent evidence accumulation. Elife 8, e46331. https://doi.org/10.7554/elife.46331

Vokoun, C.R., Huang, X., Jackson, M.B., Basso, M.A., 2014. Response Normalization in the Superficial Layers of the Superior Colliculus as a Possible Mechanism for Saccadic Averaging. J Neurosci 34, 7976–7987. https://doi.org/10.1523/jneurosci.3022-13.2014

Walker, R., Deubel, H., Schneider, W.X., Findlay, J.M., 1997. Effect of Remote Distractors on Saccade Programming: Evidence for an Extended Fixation Zone. J Neurophysiol 78, 1108–1119. https://doi.org/10.1152/jn.1997.78.2.1108

Wong, A.L., Haith, A.M., 2017. Motor planning flexibly optimizes performance under uncertainty about task goals. Nat Commun 8, 14624. https://doi.org/10.1038/ncomms14624

Wood, D.K., Gu, C., Corneil, B.D., Gribble, P.L., Goodale, M.A., 2015. Transient visual responses reset the phase of low-frequency oscillations in the skeletomotor periphery. Eur J Neurosci 42, 1919–1932. https://doi.org/10.1111/ejn.12976

867   Wood, D.K., Gallivan, J.P., Chapman, C.S., Milne, J.L., Culham, J.C., Goodale, M.A., 2011.
868   Visual salience dominates early visuomotor competition in reaching behavior. J Vision 11,
869   16–16. https://doi.org/10.1167/11.10.16

a *Single Target*

b *Double Target* — Same Reach Direction

c *Double Target* — Same stimuli

EVR

Log (EMG) (a.u.)

Sorted Trial (#)

Time Aligned to Target Onset (ms)

Figure 1

Figure 2

Figure 3

Figure 4

a Averaging Ratio (a.u.)

b EVR epoch (85 to 110 after stimulus onset)
Normalized Amplitude (a.u.)

c Fit Error (a.u.)

60° 120°

60° 120° 180°

Winner-Takes-All — Spatial Averaging
Motor Averaging — Weighted Motor Averaging

Winner-Take-All Spatial Motor Weighted Motor

Figure 5

Figure 6

Figure 7