Revealing the structure of information flows discriminates similar animal social behaviors

Gabriele Valentini[1,2], **Nobuaki Mizumoto**[2,3], **Stephen C. Pratt**[2,7], **Theodore P. Pavlic**[2,4,5,6,7*°], **Sara Imari Walker**[1,4,7*°]

[1] Arizona State University, School of Earth and Space Exploration, Tempe, AZ 85287
[2] Arizona State University, School of Life Sciences, Tempe, AZ 85287
[3] Okinawa Institute of Science & Technology Graduate University, Onna-son, Okinawa, Japan, 904-0495
[4] Arizona State University, Beyond Center for Fundamental Concepts in Science, Tempe, AZ 85287
[5] Arizona State University, School of Computing, Informatics, and Decision Systems Engineering, Tempe, AZ 85287
[6] Arizona State University, School of Sustainability, Tempe, AZ 85287
[7] Arizona State University, ASU–SFI Center for Biosocial Complex Systems, Tempe, AZ 85287
*sara.i.walker@asu.edu; tpavlic@asu.edu °co-senior authors

Behavioral correlations stretching over time are an essential but often neglected aspect of interactions among animals. These correlations pose a challenge to current behavioral-analysis methods that lack effective means to analyze complex series of interactions. Here we show that non-invasive information-theoretic tools can be used to reveal communication protocols that guide complex social interactions by measuring simultaneous flows of different types of information between subjects. We demonstrate this approach by showing that the tandem-running behavior of the ant *Temnothorax rugatulus* and that of the termites *Coptotermes formosanus* and *Reticulitermes speratus* are governed by different communication protocols. Our discovery reconciles the diverse ultimate causes of tandem running across these two taxa with their apparently similar signaling mechanisms. We show that bidirectional flow of information is present only in ants and is consistent with the use of acknowledgement signals to regulate the flow of directional information.

Introduction

Social interactions among individuals unfold across different scales of space and time (Flack 2012). At short timescales, causal relationships can often be captured by experiments that manipulate an immediate stimulus to reveal its causal connection(s) to a stereotyped response, or fixed action pattern. In herring gulls, for example, the feeding behavior of chicks is visually triggered by a red spot on the lower bill of adult gulls—a causal relationship revealed through experiments in which changes in the color of this spot were shown to affect the likelihood that chicks will engage in feeding behavior (Tinbergen 1953). However, interactions are often more complex than this because they follow a protocol where rules are conditionally

40　applied over time contingent upon the outcome of previous interactions. In these cases, where short-term
41　histories may affect longer-term outcomes, the ability to make testable predictions requires quantitative
42　tools that can capture the dynamics of the interaction protocol at an intermediate time scale. Considering

Figure 1. Tandem run recruitment by ants and termites. (A) Tandem running pairs of the ant species T. rugatulus and the termite species C. formosanus and R. speratus. (B) A sampled tandem run trajectory within an idealised environment. The encoding schemas used to discretise spatial trajectories of each ant and each termite on the basis of (C) the rotation pattern and (D) the pausing pattern.

only short-term interactions, like the stimulus-and-response of herring-gull parent and chick, might not
explain functional differences observed at long time scales in otherwise similar behaviors.

Consider the tandem-running behavior of many ants and termites, in which one individual leads a
follower through their environment, the follower walking closely behind the leader throughout the run.
(Figure 1A). At short time scales, tandem runs in the ant *Temnothorax rugatulus* appear identical to those
in the termites *Coptotermes formosanus* and *Reticulitermes speratus*. They use similar signaling
mechanisms, in which the leader releases a short-range pheromone that attracts the follower (Möglich,
Maschwitz, and Hölldobler 1974; Bordereau and Pasteels 2010), while the follower taps the leader's body
with its antennae to indicate its continued presence (Möglich, Maschwitz, and Hölldobler 1974; Franks and
Richardson 2006; Nutting 1969; Vargo and Husseneder 2009). Upon removal of the follower, the leader
stops and waits for the follower to resume contact both in ants (Möglich, Maschwitz, and Hölldobler 1974;
Franks and Richardson 2006) and in termites (Mizumoto and Dobata 2019). At long time scales, however,
there exist clear functional differences in these seemingly similar behaviors. Leaders in *T. rugatulus* use
tandem runs as a recruitment mechanism that allows followers to learn a route and acquire navigational
information necessary to later repeat the same journey independently of the leader (Figure 1B). In contrast,
termites usually use tandem runs during mating, except for one example of recruitment in a basal termite
(Sillam-Dussès et al. 2007). In a mated pair, the male follows the female leader only to maintain spatial

cohesion when searching for a new home; once a suitable location is found, the termites remain there to
start a new colony, and neither partner ever retraces the route of their tandem run.

Short-term signaling mechanisms (i.e., their stimulus and response dynamics) are similar between ants
and termites and cannot explain species differences in the function of tandem runs (i.e., route learning
versus spatial cohesion). These differences are likely encoded at intermediate time scales, where it
becomes possible in principle to detect the communication protocol (*i.e.*, set of interaction rules) that
describe how and when leader and follower use each short-term signaling mechanism. However,
experimental manipulations that operate at intermediate time scales also interfere with and constrain normal
patterns of behavior over time. As we show in this study, information-theoretic methods can reveal the
structure of information flow between subjects based only on observational data from many repeated
interactions. Moreover, these model-free methods do not rely on a priori assumptions and can be applied
over different ecological scenarios allowing for comparisons across a wide taxonomic range (McCowan,
Hanser, and Doyle 1999).

Information theory provides a model-free formalism to explicitly quantify the effects of the interaction
between individuals across space and time (Cover and Thomas 2005; Lizier, Prokopenko, and Zomaya
2008). Whereas the generic concept of *entropy* quantifies the uncertainty in a distribution of outcomes, the
derived construct of *transfer entropy* quantifies the reduction of uncertainty about the future state of a
putative receiver given knowledge of the present state of the corresponding sender (Schreiber 2000).
Transfer entropy is well suited for studying message passing; it naturally incorporates temporal ordering,
from the sender's present to the receiver's future, and quantifies the additional predictive power gained
from the sender beyond what is contained in the receiver's past. In this way it accounts for autocorrelations
that might otherwise affect behavioral data (Mitchell et al. 2019). Previous studies have used symbolic
transfer entropy (Staniek and Lehnertz 2008) to reveal whether one animal is influencing another on the
basis of a single symbolic representation of behavioral data (Orange and Abaid 2015; Butail, Mwaffo, and
Porfiri 2016; Kim et al. 2018; Ward et al. 2018; Porfiri et al. 2019; Ray et al. 2019). We extend these methods
by applying transfer entropy to different symbolic representations of the same data to capture parallel
information flows within the same behavior (*e.g.*, patterns embedded in symbols representing the direction
of motion or the speed of motion as shown in Figures 1C and 1D). Using different symbolic representations
of the same raw data allows us to uncover the complex, multi-layered structure of causal relationships
between subjects. Following this approach, we provide evidence that the communication protocol used by
leaders and followers over intermediate time scales explains the functional differences between the tandem
runs of ants and termites despite their using similar signaling mechanisms at short time scales.

Results

We first used transfer entropy to find whether the leader's or the follower's behavior better predicts the *direction of motion* of the other runner along the route. In ants, the leader is demonstrating a known route to the follower (Franks and Richardson 2006), and in termites the leader is directing a random search for a new home across the environment with the follower in tow (Nutting 1969; Vargo and Husseneder 2009). In both cases, the leader is expected to be the best predictor of the direction of the pair's motion.

Figure 2. Information flow during tandem run recruitment. *(A) The predominant direction of predictive information given by the proportion of uncertainty reduction explained by the interaction between leader and follower (mean normalized transfer entropy and standard error). (B) A schematic illustration and (C) a mechanistic illustration of the regulation of information flow in ants' tandem runs.*

Consequently, we expect the leader's behavior to be more informative about the direction of the follower than the other way around in both ants and termites. To test this hypothesis, we coarse grained the spatial trajectories of each runner into sequences of clockwise and counterclockwise turns (Figure 1C and Materials and Methods). We then measured the flow of information between the pair averaged over the entire duration of the tandem run (*i.e.*, over intermediate time scales). We found that, as expected, the leader better predicts the direction of motion of the follower than vice versa across all three species (Figure 2A, rotation bars, and Table 2).

Next, we focused on the frequent brief interruptions that give tandem runs a distinctive stop-and-go appearance. During these interruptions, the follower breaks tactile contact with the leader, who then pauses while the follower performs a local random search (Franks et al. 2010; Mizumoto and Dobata 2019). When the follower again touches the leader, the latter resumes motion, and the pair continues on their way.

In ants, these frequent interruptions are believed to regulate the speed of the run to better enable followers

to acquire navigational information (Franks and Richardson 2006; Franklin et al. 2011). As termites do not

use tandem runs to learn a route, interruptions may be more consistent with accidental chance separations

from the leader. Thus, we hypothesize that in ants, but not in termites, followers better predict the cessation

and resumption of motion than do leaders. Under this hypothesis, followers send acknowledgment signals

(Figures 2B and 2C) similar to the use of utterances (*e.g.*, "mm-hmm" described by Jefferson 1984) or

gestures (*e.g.*, the nodding of one's head) in human conversations, as well as "ACK" messages in Internet

Figure 3. Spatiotemporal dynamics of the ant T. rugatulus. *(A) Distance between the centroids of runners as a function of time. (B) Average predictive information (measured with local transfer entropy) as a function of the distance between centroids of runners (smoothed conditional means with LOESS smoothing and 0.3 span). (C) Average speed of leader and follower as a function of the distance between their centroids for increasing and decreasing distance. Purple represents the leader, green represents the follower.*

protocols that confirm receipt of other content-laden packets (Cerf and Kahn 1974). If our hypothesis is

correct, we would expect the information-theoretic signature of the tandem pair's pausing pattern in ants to

differ from that of termites. To test this, we analyzed the spatial trajectories using a different representation

obtained by coarse graining them into sequences of *pauses and movements* (Figure 1D). As hypothesized,

we found that the leader remains the best source of predictive information in termites, but in ants the follower

instead controls the flow of information and better predicts the future pausing behavior of the leader (Figure

2A, pausing bars, and Table 2).

Side-by-side comparison of tandem-run trajectories (Figures 3A and 4A) shows that ants, but not

termites, evince a tension between cohesion and information acquisition. Leader and follower ants

repeatedly switch in and out of proximity regulation under the control of the follower (Figures 3B and 3C).

The predictive power of the leader's rotation pattern dominates at close distances up to two body lengths,

Figure 4. Spatiotemporal dynamics of the termite C. formosanus. (A) Distance between the centroids of runners as a function of time. (B) Average predictive information (measured with local transfer entropy) as a function of the distance between centroids of runners (smoothed conditional means with LOESS smoothing and 0.3 span). (C) Average speed of leader and follower as a function of the distance between their centroids for increasing and decreasing distance. Purple represents the leader, green represents the follower. See also Figure 4-figure

when the pair is undergoing sustained motion and seeking cohesion (point 1, rotation regime). When their

distance increases further, the follower becomes more informative, predicting pauses in the motion of the

leader (point 2, pausing regime). Their separation then decreases as the follower approaches the stationary

leader (point 3) and predicts her resumption of motion. When leader and follower are again in close

proximity, the leader begins to move away (point 4) and this pattern repeats. Large separations are

evidently generated by the follower ant and are unrelated to rotational course corrections.

In contrast to ants, the termite leader dominates both regimes of predictive information (Figure 4).

Furthermore, these regimes are inverted with respect to ants: rotation is predicted at larger distances and

pausing of motion at shorter distances. The distance between a leader and a follower is characterized by

oscillations with higher frequency but lower amplitude than those of the ants (cf. Figure 3A and Figure 4A).

These oscillations are largely within the rotation regime due to sustained motion. In this regime, tandem

runners frequently alternate between a phase in which the leader is the faster of the two and their distance

increases (Figure 4C, point 1a) and a phase in which the follower moves faster than the leader, reducing

the gap (point 2a). Sporadically, leader and follower can be found very close to each other (less than 0.89

body lengths, Figure 4B) where they enter the pausing regime. When this happens, the leader's motion

initially predicts the decrease and then the increase in speed of the follower (points 1b and 2b). The pausing

regime is then quickly abandoned, and rotation information regains dominance. This behavior is consistent

with relatively close proximity facilitating momentary large course corrections (Figure 4A, right inset).

Leader-initiated pauses in termites might serve some unknown function, *e.g.*, motor planning (Card and

Dickinson 2008; Hunt et al. 2016); however, unlike the case of ants, we uncover no evidence that the
termite pauses facilitate follower control over any aspects of the trajectory.
152
153   **Discussion**
154
Although both ants and termites have similar mechanisms for mutual signaling at short time scales,
route learning by ants requires a communication protocol at intermediate time scales different from that
needed solely to maintain spatial cohesion. In principle, both ant and termite followers can transfer
information to their leaders by signaling their presence through physical contact. However, in contrast to
ants, termite followers in our experiments transfer information only when establishing contact at the
beginning of a run and sporadically after accidental breaks. Once contact is established and the run is
proceeding steadily, termite followers cease to transfer information to their leaders who instead control
both the direction and the speed of the run. Although manipulation experiments acting at short time scales
can show bidirectional flow in all three species, we found evidence that communication at intermediate
time scales is consistently bidirectional only in ants (from leader to follower for rotations and from follower
to leader for pauses) whereas, with the exception of accidental breaks, it is consistently unidirectional in
termites (from leader to follower for both rotations and pauses).
The communication protocol followed by termites can be likened to a person leading another by
the hand. The protocol of ants reveals instead a more complex coordination of social behavior as leader
and follower systematically alternate between close contact and separation. We suggest that the ants'
intermittent motion and bidirectional feedback is akin to the pausing for acknowledgment observed between
machines on a computer network. In this case, communication theory can aid in understanding the
frequency of acknowledgments in terms of the receiver's informational capacity and the complexity of the
information being received (Cover and Thomas 2005). The selective exposure of a follower to navigational
information from the environment is akin to the sending of a complex message over a simple channel
which, in machine-to-machine communication, requires flow control mechanisms to prevent overwhelming
the receiver.
Tandem running by the related and behaviorally similar ant *T. albipennis* has also been likened to
teaching—a social behavior often used to distinguish humans from other animals—because the leader
modifies her behavior in the presence of a naïve follower at some cost and as a result of bidirectional
feedback (Franks and Richardson 2006; Richardson et al. 2007). Our results reveal how the regulation of
information flow might be an underappreciated requirement of teaching. This assumption could be
investigated by applying the methodology we put forward here to other examples of teaching known in the
animal kingdom (Hoppitt et al. 2008). Moreover, although social insects use cue-based mechanisms for
flow and congestion control of physical quantities, such as food or nesting material (Seeley 1989;
Prabhakar, Dektar, and Gordon 2012), our study is the first to reveal a protocol for the application of
stereotyped signals to control the flow of other information (*i.e.*, a non-physical quantity) in a non-human
organism. Furthermore, tandem running has evolved multiple times in the ants but not all instances

necessarily require acknowledgment signals (Kaur et al. 2017), and so comparing across taxa may reveal

the ecological context that led to the evolution of signals that regulate other signals.

Conclusions

Temporal correlations manifesting over intermediate time scales represent an important but often

neglected aspect in behavioral ecology (Mitchell et al. 2019). Complex spatiotemporal interactions among

individuals (*i.e.*, those interactions evolving over intermediate time scales) are difficult to study by direct

manipulation in highly controlled laboratory settings. Instead, quantitative and non-interventional methods

applied over longer observational periods can be used to capture the dynamical aspect of social

interactions, but these methods are generally underdeveloped and sporadically used in behavioral ecology.

As we have shown in this study, information theory offers tools such as transfer entropy that can

disentangle the temporal structure of the interaction between individuals.

Whereas the construct of transfer entropy has seen extensive applications in the neurosciences,

particularly to study effective connectivity in the brain (Vicente et al. 2011), its application in the field of

animal behavior is less frequent and has focused primarily on revealing leader–follower relationships in

fish (Butail, Mwaffo, and Porfiri 2016; Kim et al. 2018; Ward et al. 2018) and bats (Orange and Abaid 2015),

with more recent applications in the study of decision-making in humans (Grabow et al. 2016; Porfiri et al.

2019) and slime molds (Ray et al. 2019). These previous applications set out to answer the generic

question of whether one subject is influencing another on the basis of a single symbolic representation of

raw data (e.g., a single encoding that carries information only about the direction of motion). Such uses of

transfer entropy and other information-theoretic measures (McCowan, Hanser, and Doyle 1999) cannot

disentangle the complex structure of information flow between subjects when simultaneous aspects of their

interaction carry different forms of information transmitted in different directions and at different time scales.

By quantifying multiple, concurrent symbolic patterns (i.e., variation over time in both direction and speed)

and subsequently relating the structure of information flows latent in these data to each other, we have

shown how to uncover more complex communication protocols as opposed to simply identifying

distinguished individuals within a social interaction.

The methodology we put forward, which applies advanced information-theoretic measures to

different symbolic representations of the same dataset, has allowed us to show differences in the

communication protocol used by tandem-running ants and termites, and to explain the disparity in their

function. Our approach is sufficiently generic to enable the discovery of cryptic signaling behaviors in other

taxa and to provide deeper insights into behaviors whose function is poorly or partially understood (*e.g.*,

turn-taking (Flack 2013; Pika et al. 2018) and complex coordinated dances in birds (Ota, Gahr, and Soma

2015)). Furthermore, we have shown how the generality of this approach can extend traditional information-

theoretic analysis from a mechanistic focus on one species toward a comparison across a wide taxonomic

range. Such a common language of information processing can enable the posing of new questions,

hypotheses, and predictions for the evolution of information processing itself.

Materials and Methods

Data and source code are available in (Valentini et al. 2020).

Ant experiments. We used 6 colonies of *T. rugatulus* (between 30–60 individuals each) collected in the Pinal Mountains near Globe, Arizona, during September 2017. Each colony was kept in a plastic box (110 mm by 110 mm) with a nest, a water tube, and an agar-based diet (Bhatkar and Whitcomb 1970). Nests (50 mm by 75 mm) were composed of a balsa-wood slat with a central rectangular cavity (30 mm by 50 mm) and sandwiched between two glass slides (Figure 5A). The top slide had a 2 mm hole over the center of the nest cavity to allow ants to enter and leave the nest. We conducted emigration experiments to induce ants to perform tandem runs. To obtain sufficiently long tandem runs, we used a large experimental arena (370 mm by 655 mm) delimited by walls (37 mm tall) and subdivided by 5 barriers (10 mm by 310 mm) placed to form a contiguous corridor with alternating left and right turns (Figure 5B). The design and dimensions of this arena were informed by a preliminary analysis of termite experiments (see Computation of statistics). Both walls and barriers were coated with Fluon to prevent ants from leaving the experimental arena. A new nest was placed at one extremity of the corridor and was covered with a transparent red filter to encourage the ants, which prefer dark cavities (Franks et al. 2003), to move in. The nest housing a colony was transferred from its plastic box and placed at the other extremity of the corridor. Colony emigration was induced by removing the top slide of the occupied nest. We performed 6 experiments, one for each colony, and recorded them at 30 frames per second using a video camera with 1K resolution. For each colony, we then selected between 1 and 6 pairs of ants performing tandem runs obtaining a total of 20 samples. Selected tandem runs last more than 15 minutes and have the same pair of ants travelling between the two nests with no or minimal interaction with other members of the colony.

Figure 5. Experimental setup for *experiments performed with the ant* T. rugatulus. (*A*) Nest architecture with the entrance in the center of the roof. (*B*) Experimental arena partitioned into a corridor with the old nest (bottom right) and the new nest (top left) positioned at the extremities.

Termite experiments. Experiments with *C. formosanus* and *R. speratus* were performed as part of a study on sexually dimorphic movements of termites during mate search (Mizumoto and Dobata 2019). Alates

from 2 colonies of *C. formosanus* were collected in Wakayama, Japan, in June 2017; alates from 5 colonies

of *R. speratus* were collected in Kyoto, Japan, in May 2017. After controlled nuptial flight experiments,

termites that shed their wings were selected and used for tandem run experiments. Experiments were

performed in a Petri dish (145 mm Ø) filled with moistened plaster whose surface was scraped before each

trial. A female and a male termite were introduced in the experimental arena with the opportunity to tandem

run for up to 1 hour. A total of 17 experiments were performed for *C. formosanus* and 20 experiments for

R. speratus using different individuals. Tandem runs were recorded at 30 frames per second using a video

camera with a resolution of 640 by 480 pixels.

Data extraction. We extracted motion trajectories from video recordings of tandem runs by automatically

tracking the position over time of leaders and followers. Motion tracking was accomplished using the

UMATracker software platform (Yamanaka and Takeuchi 2018). Because we tracked the centroids of each

runner's body, the distance between individuals was always greater than zero even when leader and

follower were in contact with each other. All trajectories were sampled at 30 frames per second and

shortened to a duration of 15 minutes. Trajectories were then converted from pixels to millimeters using a

scaling factor estimated by measuring known features of the experimental arena with ImageJ (Schneider,

Rasband, and Eliceiri 2012). The body size of each runner (average ± standard deviation) was measured

from video recordings of the experiments using ImageJ (*T. rugatulus*: 2.34 ± 0.3 mm, *C. formosanus*: 8.89

± 0.42 mm, *R. speratus*: 5.5 ± 0.3 mm).

Encoding behavioral patterns into symbolic time series. We considered three possible behavioral

patterns for each runner: *pausing pattern*, *rotation pattern*, and their combination *pausing & rotation

pattern*. We did so by coarse-graining the space-continuous trajectories of each leader and each follower

using three different symbolic representations. Each spatial trajectory consists of a sequence $(q_1, q_2, ...)$ of

2-dimensional points, $q_i = \left(q_i^x, q_i^y \right)$, representing spatial coordinates over time which are then encoded

into a symbolic time series $X = (x_1, x_2, ...)$. To capture the time interval where the sender best predicts the

behavior of the receiver, we subsampled spatial trajectories in time before encoding the behavioral patterns

of each runner. We considered different sampling periods, starting from a short period of one sample every

33.3667 ms (29.97 Hz) to a long period of one sample every 1.5015 s (0.666 Hz) with an interval between

each period of 33.3667 ms (*i.e.*, sampling period $\in \{0.0334s, 0.667s, ..., 1.5015s\}$). Depending on the

sampling period, the resulting time series have a number of time steps between 599 and 26973.

Figure 6. Step-size distribution as a function of sampling period. *Probability density function of the step size for*
the ant T. rugatulus as a function of the sampling period. Purple represents the 10% probability mass used to define
the pause state; Green represents the remaining 90% of the probability mass defining the motion state. See also Figure
6-figure supplement 1 for the termite C. formosanus and Figure 6-figure supplement 2 for R. speratus.

The pausing pattern is encoded using two states: the *motion state* (M) and the *pause* state (P). The
motivation for this coding scheme is to capture when a tandem runner pauses while waiting for the other
to re-join the tandem run or to react to physical contact. Pauses, small adjustments of the position of the
runner, or changes due to noise in the sampled trajectories may each accidentally be considered as
genuine acts of motion. To prevent these spurious classifications, we used a threshold based on travelled
distance to distinguish segments of the trajectory into those identifying motion and those identifying pauses.
We first computed the probability distribution of step sizes, *i.e.*, the distance travelled by a runner between
two consecutive sampled positions q_i and q_{i+1} for each species and sampling period (Figure 6 and
supplements). These distributions show two distinct modes: short steps (i.e., low speed) characteristic of
pauses and long steps (i.e., high speed) characteristic of sustained motion. After inspection, we chose the
10th percentile of each probability distribution as the distance threshold used to distinguish between motion
and pauses. We therefore encoded as pause states all time steps in a given spatial trajectory with a
corresponding travelled distance in the 10th size percentile and the remaining 90% of time steps as motion
states. This threshold was varied in the interval $\{5\%, 6\%, \dots, 15\%\}$ during a perturbation analysis of
predictive information (see Computation of statistics).
The rotation pattern is also encoded using two states: *clockwise* (CW) and *counterclockwise* (CCW).
The direction of rotation at time i is obtained by looking at three consecutive positions, q_{i-1}, q_i, q_{i+1}, in the
spatial trajectory of each runner. The rotation is clockwise when the cross product $\overrightarrow{q_{i-1}q_i} \times \overrightarrow{q_i q_{i+1}}$ is positive,
counterclockwise when it is negative, and collinear when it is exactly zero. As we aim to model only
clockwise and counter-clockwise rotations, we do not consider any tolerance threshold to explicitly capture
collinear motion. Instead, in the rare occurrences of collinear motion, the direction of rotation at the previous
time step, $i - 1$, is copied over in the time series.
As a control for our choices of possible behavioral outcomes, we also considered a compound pausing
& rotation pattern that simultaneously encodes both components of tandem running. A possible approach

to do so is to use the time series of each behavioral pattern separately but then rely on multivariate
measures of predictive information. However, in our scenario, pauses have a mutually exclusive relation
with rotations that cannot be preserved using multivariate measures. The pausing & rotation pattern is
defined therefore using a ternary coding scheme that encodes motion bouts in the states *pause* (P),
clockwise (CW) and *counterclockwise* (CCW). As for the pausing pattern, the shortest 10% of steps in the
spatial trajectories are encoded as pausing (see Computation of statistics for a perturbation analysis of this
parameter). The remaining 90% of steps are encoded using states clockwise and counterclockwise
following the same methodology used for the rotation pattern.

Measuring predictive information. Our analysis of communication in tandem running is grounded in the
theory of information (Cover and Thomas 2005) and its constructs of entropy, conditional entropy, and
transfer entropy. We aim to quantify how knowledge of the current behavior of the sender allows us to
predict the future behavior of the receiver, that is, to measure causal interactions in a Wiener-Granger
sense (Bossomaier et al. 2016). We consider the behavioral patterns of leaders and followers as the series
of realizations $(l_i, i \geq 1)$ and $(f_i, i \geq 1)$ of two random variables, L and F, corresponding to the leader and
follower respectively. For simplicity, the following presentation focuses on predicting the future of the
follower, $F^{i+1} = (f_{i+1}, i \geq 1)$, from the present of the leader, L, but in our analysis we also consider how
much of the future of the leader, L^{i+1}, is predicted by the present of the follower, F.

The overall uncertainty about the future F^{i+1} of the follower is quantified by the (marginal) entropy
(Shannon 1948) $H(F^{i+1}) = - \sum_{f_{i+1}} p(f_{i+1}) \log_2 p(f_{i+1})$. Entropy measures the average amount of
information necessary to uniquely identify an outcome of F^{i+1}. Knowing the history of the follower may
reduce the uncertainty in the distribution of possible outcomes for the future of the follower, and the
reduction in uncertainty can be quantified by the difference between the marginal entropy and the entropy
after the historical information is considered. Let $f_i^{(k)} = \{f_{i-k+1}, ..., f_{i-1}, f_i\}$ represent the finite history with
length k of F up to the current time i and $F^{(k)}$ a new random variable defined over a series $\left(f_i^{(k)}, i \geq 1\right)$ of
k-histories. The amount of uncertainty about F^{i+1} that is left after accounting for its past behavior $F^{(k)}$ is
given by the conditional entropy

345
$$H\left(F^{i+1} \middle| F^{(k)}\right) = - \sum_{f_i^{(k)}, f_{i+1}} p\left(f_i^{(k)}, f_{i+1}\right) \log_2 \frac{p\left(f_i^{(k)}, f_{i+1}\right)}{p\left(f_i^{(k)}\right)},$$

for history length $1 \leq k < \infty$. $H(F^{i+1}|F^{(k)})$ represents the average amount of information necessary to
uniquely identify the future behavior of the follower given what we know about its past behavior.
A second step to obtain additional information about the future of the follower is to consider the time-
delayed effects of its interaction with the leader. Transfer entropy was introduced for this purpose
(Schreiber 2000). It measures the amount of information about the future behavior of the receiver given by
knowledge of the current behavior of the sender that is not contained in the receiver's past. Due to its time
directionality (*i.e.*, from the present of the sender to the future of the receiver), it is considered a measure

of information transfer or predictive information (Lizier and Prokopenko 2010). Transfer entropy is defined
as

$$T_{L \to F} = \sum_{f_{i+1}, f_i^{(k)}, l_i} p(f_{i+1}, f_i^{(k)}, l_i) \log_2 \frac{p\big(f_{i+1} \mid f_i^{(k)}, l_i\big)}{p\big(f_{i+1} \mid f_i^{(k)}\big)}$$

and measures the reduction of uncertainty of F^{i+1} given from knowledge of L which is not already given by
$F^{(k)}$. The logarithm in the above equation is known as local transfer entropy (Lizier, Prokopenko, and
Zomaya 2008) and tells us whether, at time i, the interaction $l_i \mid f_i^{(k)} \to f_{i+1} \mid f_i^{(k)}$ between the two
processes is informative (> 0) or misinformative (< 0). In our analysis, we look at local transfer entropy
averaged over the distance between leader and follower to understand the spatiotemporal dynamics of
communication during tandem running.

Due to the asymmetry of transfer entropy, $T_{L \to F} \neq T_{F \to L}$, we can obtain the predominant direction and
the magnitude of predictive information by studying the difference

$$T_{L \to F} - T_{F \to L}.$$

365　This quantity is positive when information flows predominantly from leader to follower ($T_{L \to F} > T_{F \to L}$) and
366　negative when it flows from follower to leader ($T_{L \to F} < T_{F \to L}$). Its value is known as net transfer entropy
367　(Porfiri 2018). Finally, as transfer entropy can be rewritten as $T_{L \to F} = H\big(F^{i+1} \big| F^{(k)}\big) - H(F^{i+1} \big| F^{(k)}, L)$, we
368　can normalise this quantity in the interval $[0; 1]$ simply by dividing it by the conditional entropy as in

$$\frac{T_{L \to F}}{H\big(F^{i+1} \big| F^{(k)}\big)} = \frac{H\big(F^{i+1} \big| F^{(k)}\big) - H(F^{i+1} \big| F^{(k)}, L)}{H\big(F^{i+1} \big| F^{(k)}\big)}.$$

Normalized transfer entropy (Porfiri 2018) is a dimensionless quantity that captures the proportion of the
future behavior F^{i+1} of the follower that is explained by the interaction with the leader at time i. When F^{i+1}
is completely predicted by L, the conditional entropy $H(F^{i+1} \big| F^{(k)}, L)$ is zero and normalized transfer
entropy is maximal and equal to 1; instead, when F^{i+1} is independent of L, $H\big(F^{i+1} \big| F^{(k)}\big) = H(F^{i+1} \big| F^{(k)}, L)$
and normalised transfer entropy is minimal and equal to 0.

Computation of statistics. We computed information-theoretic measures for both leaders and followers.
In our computations, we assume that the pausing and rotation patterns of ants and termites are peculiar
features of the species rather than of specific pairs of tandem runners. As such, rather than treating each
trial separately and then aggregating the results, we estimated the necessary probabilities from all
experimental trials together and obtained a single estimate of transfer entropy for each considered species
and parameter configuration. Our measures of predictive information are therefore averaged over all trials
of the same species. Probability distributions are estimated from the frequencies of blocks of consecutive
symbols within the time series. For example, the probability for a follower to have a history of rotations
$f_i^{(3)} = \{f_{i-2} = CW, f_{i-1} = CW, f_i = CCW\}$ is estimated by counting the number of times the symbols
$\{CW, CW, CCW\}$ occur consecutively at any point in the time series of any follower; this count is then

normalized by the number of samples to obtain a measure of probability. All information-theoretic measures
were computed in R 3.4.3 using the `rinform-1.0.1` package (Moore et al. 2018).
To ensure that the measured interactions are valid and not the result of artefacts that may arise due to
finite sample sets, we compared transfer entropy measured from the experimental data with measurements
from surrogate datasets artificially created by pairing independent time series (Porfiri 2018). To create a
surrogate dataset, we randomly paired the behavioral patterns of leaders and followers belonging to
different tandem runs, obtaining a dataset with the same size as the original. We then computed transfer
entropy for this surrogate data. Although leaders and followers from different runs are still influenced by
the same environmental cues, this randomization process breaks possible causal interactions within the
surrogate pair. For each species and parameter configuration, we repeated this randomization process 50
times obtaining 50 surrogate datasets that were used to estimate mean and standard error of transfer
entropy. Finally, measurements of transfer entropy for the experimental data were discounted by a
correction factor given by the estimated means.

Figure 7. Landscape of net information transfer. *Net transfer entropy (bits) as a function of the sampling period and of the history length. (A), (B), and (C) show the results for the rotation pattern, (D), (E), and (F) show the results for the pausing pattern, and (G), (H), and (I) show the results for the compound pausing & rotation pattern. The first, second, and third columns show the results, respectively, for T. rugatulus, C. formosanus, and R. speratus. Colors indicate the intensity and predominant direction of information transfer (purple, from leader to follower; green, from follower to leader); the red diamond symbol indicates the configuration with maximum magnitude. See also Figure 7-figure supplements 1–3.*

The sampling period of continuous spatial trajectories and the history length of transfer entropy define the parameter space of our study. The optimal choice of these parameters likely varies for different species and between leaders and followers within a species as a result of behavioral, morphological, and cognitive differences manifesting at different time scales. To choose a suitable parameter configuration and control for its robustness, we computed net transfer entropy for 900 different parameter configurations for each species (history length $k \in \{1, \dots, 20\}$ and sampling period $\{0.0334s, \dots, 1.5015s\}$). From the resulting landscapes of information transfer, which show robustness to variation of parameters, we then selected

the parameter configurations that maximize the net transfer of information (see Figure 7 and Table 1). A

similar analysis was preliminarily performed for increasing duration of tandem runs on the basis of the

termite data with the aim to inform the design of the experimental arena used for ants (see Figure 7-figure

supplement 1). These parameters, whose values converge for increasing length of the time series, are

relatively similar across behavioral patterns for both species of termites. Ants instead are characterized by

more diverse time scales likely because leaders and followers cause different aspects of tandem running

and, possibly, because they do so by following cognitive processes with different time constraints (see

Table 2 for summary statistics).

Table 1. Selected parameter configurations. *For each species and behavioral pattern, the table provides the number*
of tandem runs used in the study, the sampling period and corresponding number of time steps in each time series,
and the history length.

Species	#Tandem runs	Behavioral pattern	Sampling period (s)	#Time steps	History length k
T. rugatulus	20	Rotation	1.5015	599	9
		Pausing	0.9676	930	13
		Pausing & Rotation	1.2346	728	8
C. formosanus	17	Rotation	0.3670	2452	2
		Pausing	0.1668	5395	1
		Pausing & Rotation	0.3670	2452	2
R. speratus	20	Rotation	0.5005	1798	1
		Pausing	0.4671	1926	1
		Pausing & Rotation	0.5005	1798	2

427
428

For the chosen parameter configurations, we tested both the significance of our estimate of transfer

entropy with respect to surrogate data and that of leader–follower relations observed in Figure 2A. One-

sided two-sample Wilcoxon rank-sum tests with continuity correction show values of transfer entropy for

the experimental data significantly greater than those for surrogate data (Table 3, columns 3 and 4) in all

but four of these tests. One-sided paired Wilcoxon signed-rank tests with continuity correction were used

instead to test differences in causal interactions between leaders and followers and to confirm the effects

shown in Figure 2A. All leader–follower interactions were correctly identified by this analysis and none of

the significant tests is among the four cases mentioned above (Table 3, columns 5 and 6). Next, we

performed a perturbation analysis of the probability threshold used to separate pauses from motion in the

pausing pattern and in the pausing & rotation pattern ($\{5\%, 6\%, ..., 15\%\}$). Although the magnitude is subject

to some variation, the direction of information transfer that represents our primary observable remains

unaltered (see Figure 7-figure supplement 2). Finally, we also controlled for our choices of symbolic

representation on possible outcomes in the behavioral patterns by considering a compound pausing and

rotation pattern. Figure 7-figure supplement 3 shows the results of this analysis which closely resemble

those shown in Figure 3 for *T. rugatulus,* Figure 4 for *C. formosanus*, and Figure 4-figure supplement 1 for

R. speratus.

Table 2. Statistics about transfer entropy. *Mean value and standard error of transfer entropy for the experimental*
($T_{X \to Y}$) and surrogate ($T_{X \to Y}^s$) datasets and of normalized transfer entropy ($\tilde{T}_{X \to Y}$) from the predominant source of

Species	Behavioral pattern	$T_{L \to F}$	$T^s_{L \to F}$	$T_{F \to L}$	$T^s_{F \to L}$	$\tilde{T}_{L \to F}$ ($\tilde{T}_{F \to L}$)
T. rugatulus	Rotation	$.1488 \pm .0504$	$.0366 \pm .0026$	$.0368 \pm .0125$	$.0371 \pm .0073$	$.1197 \pm .0543$
	Pausing	$.0058 \pm .0033$	$.0047 \pm .0015$	$.0423 \pm .021$	$.0081 \pm .0044$	$.0795 \pm .0401$
	Pausing & Rotation	$.2656 \pm .034$	$.1429 \pm .0092$	$.1836 \pm .0368$	$.1847 \pm .0269$	$.1246 \pm .035$
C. formosanus	Rotation	$.2063 \pm .0731$	$.0009 \pm .0019$	$.0091 \pm .0077$	$.0012 \pm .0017$	$.2884 \pm .0844$
	Pausing	$.0303 \pm .0209$	$.0001 \pm .0003$	$.0064 \pm .0056$	$.0001 \pm .0001$	$.1532 \pm .0865$
	Pausing & Rotation	$.2593 \pm .072$	$.0024 \pm .0028$	$.013 \pm .0135$	$.0026 \pm .0014$	$.2991 \pm .0583$
R. speratus	Rotation	$.1804 \pm .1097$	$.0001 \pm .0003$	$.0027 \pm .0137$	$.0001 \pm .0003$	$.1949 \pm .119$
	Pausing	$.0228 \pm .0149$	$.0001 \pm .0002$	$.0055 \pm .0087$	$.0001 \pm .0001$	$.0698 \pm .0445$
	Pausing & Rotation	$.2385 \pm .1446$	$.0011 \pm .0013$	$.0081 \pm .0121$	$.0011 \pm .0007$	$.2082 \pm .1359$

Table 3. Results of hypothesis testing of information transfer between leaders and followers. *Columns 3 and 4 report the results of one-sided two-sample Wilcoxon rank-sum tests with continuity correction (p-value and W statistic) testing if the experimental dataset has significantly higher transfer entropy than the surrogate one. Columns 5 and 6 report the results of one-sided paired Wilcoxon signed-rank tests with continuity correction (p-value and V statistic) testing, respectively, if the leader is significantly more informative than the follower and vice versa. Significant p-values are reported in bold.*

Species	Behavioral pattern	$H_1: T_{L \to F} > T^s_{L \to F}$	$H_1: T_{F \to L} > T^s_{F \to L}$	$H_1: T_{L \to F} > T_{F \to L}$	$H_1: T_{F \to L} > T_{L \to F}$
T. rugatulus	Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 380$)	$p = .51$ ($W = 200$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 209$)	$p = 1.0$ ($V = 1$)
	Pausing	$p = .21$ ($W = 230$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 388$)	$p = 1.0$ ($V = 0$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 210$)
	Pausing & Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 400$)	$p = .7$ ($W = 181$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 207$)	$p = 1.0$ ($V = 3$)
C. formosanus	Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 289$)	$\boldsymbol{p} = \boldsymbol{.002}$ ($W = 226$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 153$)	$p = 1.0$ ($V = 0$)
	Pausing	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 272$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 272$)	$\boldsymbol{p} = \boldsymbol{.001}$ ($V = 138$)	$p = 1.0$ ($V = 15$)
	Pausing & Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 289$)	$\boldsymbol{p} = \boldsymbol{.001}$ ($W = 229$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 153$)	$p = 1.0$ ($V = 0$)
R. speratus	Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 380$)	$p = .3$ ($W = 220$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 201$)	$p = 1.0$ ($V = 9$)
	Pausing	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 380$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 340$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 195$)	$p = 1.0$ ($V = 15$)
	Pausing & Rotation	$\boldsymbol{p} < \boldsymbol{.001}$ ($W = 380$)	$\boldsymbol{p} = \boldsymbol{.018}$ ($W = 278$)	$\boldsymbol{p} < \boldsymbol{.001}$ ($V = 193$)	$p = 1.0$ ($V = 17$)

References

Bhatkar, A, and W H Whitcomb. 1970. "Artificial Diet for Rearing Various Species of Ants." *The Florida Entomologist* 53 (4): 229–32. https://doi.org/10.2307/3493193.

Bordereau, Christian, and Jacques M Pasteels. 2010. "Pheromones and Chemical Ecology of Dispersal and Foraging in Termites." In *Biology of Termites: A Modern Synthesis*, 279–320. Springer. https://doi.org/10.1007/978-90-481-3977-4_11.

Bossomaier, Terry, Lionel Barnett, Michael Harré, and Joseph T Lizier. 2016. *An Introduction to Transfer Entropy*. Springer.

Butail, Sachit, Violet Mwaffo, and Maurizio Porfiri. 2016. "Model-Free Information-Theoretic Approach to Infer Leadership in Pairs of Zebrafish." *Phys. Rev. E* 93 (4): 42411.

https://doi.org/10.1103/PhysRevE.93.042411.

Card, Gwyneth, and Michael H. Dickinson. 2008. "Visually Mediated Motor Planning in the Escape

Response of Drosophila." *Current Biology* 18 (17): 1300–1307.

https://doi.org/10.1016/J.CUB.2008.07.094.

Cerf, V, and R Kahn. 1974. "A Protocol for Packet Network Intercommunication." *IEEE Transactions on*

Communications 22 (5): 637–48. https://doi.org/10.1109/TCOM.1974.1092259.

Cover, Thomas M, and Joy A Thomas. 2005. *Elements of Information Theory*. *Elements of Information*

Theory. Hoboken, NJ, USA: John Wiley & Sons, Inc. https://doi.org/10.1002/047174882X.

Flack, Jessica C. 2012. "Multiple Time-Scales and the Developmental Dynamics of Social Systems."

Philosophical Transactions of the Royal Society B: Biological Sciences 367 (1597): 1802–10.

https://doi.org/10.1098/rstb.2011.0214.

———. 2013. "Animal Communication: Hidden Complexity." *Current Biology* 23 (21): R967--R969.

https://doi.org/10.1016/j.cub.2013.10.001.

Franklin, Elizabeth L, Thomas O Richardson, Ana B Sendova-Franks, Elva J H Robinson, and Nigel R

Franks. 2011. "Blinkered Teaching: Tandem Running by Visually Impaired Ants." *Behavioral*

Ecology and Sociobiology 65 (4): 569–79. https://doi.org/10.1007/s00265-010-1057-2.

Franks, Nigel R, Eamonn B Mallon, Helen E Bray, Mathew J Hamilton, and Thomas C Mischler. 2003.

"Strategies for Choosing between Alternatives with Different Attributes: Exemplified by House-

Hunting Ants." *Animal Behaviour* 65 (1): 215–23. https://doi.org/10.1006/anbe.2002.2032.

Franks, Nigel R, and Thomas O Richardson. 2006. "Teaching in Tandem-Running Ants." *Nature* 439

(7073): 153. https://doi.org/10.1038/439153a.

Franks, Nigel R, Thomas O Richardson, Samantha Keir, Stephen J Inge, Frederic Bartumeus, and Ana B

Sendova-Franks. 2010. "Ant Search Strategies after Interrupted Tandem Runs." *The Journal of*

Experimental Biology 213 (10): 1697 LP – 1708. https://doi.org/10.1242/jeb.031880.

Grabow, Carsten, James Macinko, Diana Silver, and Maurizio Porfiri. 2016. "Detecting Causality in Policy

Diffusion Processes." *Chaos: An Interdisciplinary Journal of Nonlinear Science* 26 (8): 83113.

https://doi.org/10.1063/1.4961067.

Hoppitt, William J E, Gillian R Brown, Rachel Kendal, Luke Rendell, Alex Thornton, Mike M Webster, and

Kevin N Laland. 2008. "Lessons from Animal Teaching." *Trends in Ecology & Evolution* 23 (9): 486–

93. https://doi.org/https://doi.org/10.1016/j.tree.2008.05.008.

Hunt, Edmund R, Roland J Baddeley, Alan Worley, Ana B Sendova-Franks, and Nigel R Franks. 2016.

"Ants Determine Their next Move at Rest: Motor Planning and Causality in Complex Systems."

Royal Society Open Science 3 (1): 150534. https://doi.org/10.1098/rsos.150534.

Jefferson, Gail. 1984. "Notes on a Systematic Deployment of the Acknowledgement Tokens 'Yeah' and

'Mm Hm.'" *Papers in Linguistics* 17: 197–206.

Kaur, Rajbir, Joby Joseph, Karunakaran Anoop, and Annagiri Sumana. 2017. "Characterization of

Recruitment through Tandem Running in an Indian Queenless Ant *Diacamma Indicum*." *Royal*

Society Open Science 4 (1): 160476. https://doi.org/10.1098/rsos.160476.

Kim, Changsu, Tommaso Ruberto, Paul Phamduy, and Maurizio Porfiri. 2018. "Closed-Loop Control of Zebrafish Behaviour in Three Dimensions Using a Robotic Stimulus." *Scientific Reports* 8 (1): 657. https://doi.org/10.1038/s41598-017-19083-2.

Lizier, Joseph T, and Mikhail Prokopenko. 2010. "Differentiating Information Transfer and Causal Effect." *The European Physical Journal B* 73 (4): 605–15. https://doi.org/10.1140/epjb/e2010-00034-5.

Lizier, Joseph T, Mikhail Prokopenko, and Albert Y Zomaya. 2008. "Local Information Transfer as a Spatiotemporal Filter for Complex Systems." *Phys. Rev. E* 77 (2): 26110. https://doi.org/10.1103/PhysRevE.77.026110.

McCowan, Brenda, Sean F Hanser, and Laurance R Doyle. 1999. "Quantitative Tools for Comparing Animal Communication Systems: Information Theory Applied to Bottlenose Dolphin Whistle Repertoires." *Animal Behaviour* 57 (2): 409–19. https://doi.org/10.1006/anbe.1998.1000.

Mitchell, David J, Antoine M Dujon, Christa Beckmann, and Peter A Biro. 2019. "Temporal Autocorrelation: A Neglected Factor in the Study of Behavioral Repeatability and Plasticity." *Behavioral Ecology*. https://doi.org/10.1093/beheco/arz180.

Mizumoto, Nobuaki, and Shigeto Dobata. 2019. "Adaptive Switch to Sexually Dimorphic Movements by Partner-Seeking Termites." *Science Advances* 5 (6): eaau6108. https://doi.org/10.1126/sciadv.aau6108.

Möglich, Michael, Ulrich Maschwitz, and Bert Hölldobler. 1974. "Tandem Calling: A New Kind of Signal in Ant Communication." *Science* 186 (4168): 1046–47. https://doi.org/10.1126/science.186.4168.1046.

Moore, Douglas G, Gabriele Valentini, Sara Imari Walker, and Michael Levin. 2018. "Inform: Efficient Information-Theoretic Analysis of Collective Behaviors." *Frontiers in Robotics and AI* 5: 60. https://doi.org/10.3389/frobt.2018.00060.

Nutting, W L. 1969. "Flight and Colony Foundation." *Biology of Termites*, 233–82.

Orange, N, and N Abaid. 2015. "A Transfer Entropy Analysis of Leader-Follower Interactions in Flying Bats." *The European Physical Journal Special Topics* 224 (17): 3279–93. https://doi.org/10.1140/epjst/e2015-50235-9.

Ota, Nao, Manfred Gahr, and Masayo Soma. 2015. "Tap Dancing Birds: The Multimodal Mutual Courtship Display of Males and Females in a Socially Monogamous Songbird." *Scientific Reports* 5 (November): 16614. https://doi.org/10.1038/srep16614.

Pika, Simone, Ray Wilkinson, Kobin H Kendrick, and Sonja C Vernes. 2018. "Taking Turns: Bridging the Gap between Human and Animal Communication." *Proceedings of the Royal Society B: Biological Sciences* 285 (1880): 20180598. https://doi.org/10.1098/rspb.2018.0598.

Porfiri, Maurizio. 2018. "Inferring Causal Relationships in Zebrafish-Robot Interactions through Transfer Entropy: A Small Lure to Catch a Big Fish." *Animal Behavior and Cognition* 5 (4): 341–67. https://doi.org/10.26451/abc.05.04.03.2018.

Porfiri, Maurizio, Raghu Ram Sattanapalle, Shinnosuke Nakayama, James Macinko, and Rifat Sipahi.

545  2019. "Media Coverage and Firearm Acquisition in the Aftermath of a Mass Shooting." *Nature*

546  *Human Behaviour* 3 (9): 913–21. https://doi.org/10.1038/s41562-019-0636-0.

547 Prabhakar, Balaji, Katherine N Dektar, and Deborah M Gordon. 2012. "The Regulation of Ant Colony

548  Foraging Activity without Spatial Information." Edited by Iain D. Couzin. *PLoS Computational Biology*

549  8 (8): e1002670. https://doi.org/10.1371/journal.pcbi.1002670.

550 Ray, Subash K, Gabriele Valentini, Purva Shah, Abid Haque, Chris R Reid, Gregory F Weber, and Simon

551  Garnier. 2019. "Information Transfer During Food Choice in the Slime Mold Physarum

552  Polycephalum." *Frontiers in Ecology and Evolution* 7: 67. https://doi.org/10.3389/fevo.2019.00067.

553 Richardson, Thomas O, Philippa A Sleeman, John M McNamara, Alasdair I Houston, and Nigel R Franks.

554  2007. "Teaching with Evaluation in Ants." *Current Biology* 17 (17): 1520–26.

555  https://doi.org/10.1016/j.cub.2007.08.032.

556 Schneider, Caroline A, Wayne S Rasband, and Kevin W Eliceiri. 2012. "NIH Image to ImageJ: 25 Years

557  of Image Analysis." *Nature Methods* 9: 671. https://doi.org/10.1038/nmeth.2089.

558 Schreiber, Thomas. 2000. "Measuring Information Transfer." *Physical Review Letters* 85 (2): 461–64.

559  https://doi.org/10.1103/PhysRevLett.85.461.

560 Seeley, Thomas D. 1989. "Social Foraging in Honey Bees: How Nectar Foragers Assess Their Colony's

561  Nutritional Status." *Behavioral Ecology and Sociobiology* 24 (3): 181–99.

562  https://doi.org/10.1007/BF00292101.

563 Shannon, C E. 1948. "A Mathematical Theory of Communication." *The Bell System Technical Journal* 23

564  (3): 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x.

565 Staniek, Matthäus, and Klaus Lehnertz. 2008. "Symbolic Transfer Entropy." *Phys. Rev. Lett.* 100 (15):

566  158101. https://doi.org/10.1103/PhysRevLett.100.158101.

567 Tinbergen, Niko. 1953. *The Herring Gull's World: A Study of the Social Behaviour of Birds. The Herring*

568  *Gull's World: A Study of the Social Behaviour of Birds.* Oxford, England, England: Frederick A.

569  Praeger, Inc.

570 Valentini, Gabriele, Nobuaki Mizumoto, Stephen C Pratt, Theodore P Pavlic, and Sara Imari Walker.

571  2020. "Data and Code from: Revealing the Structure of Information Flows Discriminates Similar

572  Animal Social Behaviors." Figshare. 2020. https://doi.org/10.6084/m9.figshare.9786260.

573 Vargo, Edward L, and Claudia Husseneder. 2009. "Biology of Subterranean Termites: Insights from

574  Molecular Studies of Reticulitermes and Coptotermes." *Annual Review of Entomology* 54: 379–403.

575  https://doi.org/10.1146/annurev.ento.54.110807.090443.

576 Vicente, Raul, Michael Wibral, Michael Lindner, and Gordon Pipa. 2011. "Transfer Entropy---a Model-

577  Free Measure of Effective Connectivity for the Neurosciences." *Journal of Computational*

578  *Neuroscience* 30 (1): 45–67. https://doi.org/10.1007/s10827-010-0262-3.

579 Ward, Ashley J W, T M Schaerf, A L J Burns, J T Lizier, E Crosato, M Prokopenko, and M M Webster.

580  2018. "Cohesion, Order and Information Flow in the Collective Motion of Mixed-Species Shoals."

581  *Royal Society Open Science* 5 (12): 181132. https://doi.org/10.1098/rsos.181132.

582 Yamanaka, Osamu, and Rito Takeuchi. 2018. "UMATracker: An Intuitive Image-Based Tracking
583   Platform." *Journal of Experimental Biology* 221 (16). https://doi.org/10.1242/jeb.182469.
584

Acknowledgements

586 This work was supported by NSF grant No. PHY-1505048. N.M. was supported by a JSPS Overseas
587 Research Fellowship. The authors would like to thank Dr. Anna Dornhaus for constructive criticism of the
588 manuscript.
589

Author contributions statement

Gabriele Valentini: Conceptualization; Data curation; Formal analysis; Investigation; Visualization;
Methodology; Writing - original draft, review and editing Nobuaki Mizumoto: Investigation; Writing - review
and editing Stephen Pratt: Conceptualization; Supervision; Funding acquisition; Writing - review and editing
Theodore Pavlic: Conceptualization; Supervision; Funding acquisition; Writing - original draft, review and
editing Sara Walker: Conceptualization; Supervision; Funding acquisition; Project administration; Writing -
review and editing.
597

Additional information

Accession codes: Data and code that support the findings of this study is available in "figshare" with identifier "10.6084/m9.figshare.9786260".

Competing interests: The authors declare no competing interest.

625 **Supplemental Information**

626
627 ***Figure 4-figure supplement 1. Spatiotemporal dynamics of the termite R. speratus.*** *(A) Distance between the*
628 *centroids of runners as a function of time. (B) Average predictive information (measured with local transfer entropy) as*
629 *a function of the distance between centroids of runners (smoothed conditional means with LOESS smoothing and 0.3*
630 *span). (C) Average speed of leader and follower as a function of the distance between their centroids for increasing*
631 *and decreasing distance. Purple represents the leader, green represents the follower.*

634
Figure 6-figure supplement 1. Step-size distribution as a function of sampling period. *Probability density function*
of the step size for the termite C. formosanus *as a function of the sampling period. Purple represents the 10% probability*

mass used to define the paus state, green represents the remaining 90% of the probability mass defining the motion
state.

Figure 6-figure supplement 2. Step-size distribution as a function of sampling period. *Probability density function*
of the step size for the termite R. speratus as a function of the sampling period. Purple represents the 10% probability
mass used to define the paus state, green represents the remaining 90% of the probability mass defining the motion
state.

645
Figure 7-figure supplement 1. Impact of amount of data on the landscape information transfer. *Net transfer*
entropy (bits) as a function of the sampling period and of the history length for increasing length of time series (1, 5,
10, and 15 minutes). (A) and (B) show the results for the rotation pattern and for the pausing pattern for C. formosanus.
(C) and (D) show the results for the rotation pattern and for the pausing pattern for T. rugatulus. Colors indicate the
intensity and predominant direction of information transfer (purple, from leader to follower; green, from follower to
leader); the red diamond symbol indicates the configuration with maximum absolute magnitude and each panel reports
the result of a one-sided paired Wilcoxon signed-rank test with continuity correction (p-value and V statistic) testing the
significance of the observed effect for this parameter configuration.

654
655

656
657 **Figure 7-figure supplement 2. Perturbation analysis of pause-motion threshold.** *Net transfer entropy as a function*
658 *of the probability threshold used to encode step sizes into the motion state and pausing state. (A) Results for the*
659 *pausing pattern. (B) Results for the compound pausing & rotation pattern. Sampling period and history length*
660 *correspond to those in Table 1. Positive values represent information transfer from leader to follower, negative values*
661 *represent information transfer from follower to leader.*

662
663

664
665 **Figure 7-figure supplement 3. Predictive information over distance between runners.** *Average local transfer*
666 *entropy as a function of the distance between centroids of runners for the compound pausing and rotation pattern*
667 *(smoothed conditional means with LOESS smoothing and* 0.3 *span). (A) T. rugatulus, (B) C. formosanus, and (C) R.*
668 *speratus.*