An unprecedented combination of metabolic pathways in the cryptic plastid of a non-photosynthetic

euglenophyte

Zoltán Füssy[1,2,#], Kristína Záhonová[1,2,3,#], Aleš Tomčala[1,§], Juraj Krajčovič[4], Vyacheslav Yurchenko[3],

Miroslav Oborník[1,5], Marek Eliáš[3,*]

[1] Institute of Parasitology, Biology Centre ASCR, 370 05 České Budějovice, Czech Republic

[2] Faculty of Science, Charles University, BIOCEV, 252 50 Vestec, Czech Republic

[3] Life Science Research Centre, Department of Biology and Ecology and Institute of Environmental

Technologies, Faculty of Science, University of Ostrava, 701 00 Ostrava, Czech Republic

[4] Department of Biology, Faculty of Natural Sciences, University of ss. Cyril and Methodius in Trnava,

917 01 Trnava, Slovakia

[5] University of South Bohemia, Faculty of Science, 370 05 České Budějovice, Czech Republic

[#] These authors contributed equally to this work.

[§] Present address: University of South Bohemia, Faculty of Fisheries and Protection of Waters,

CENAKVA, 370 05 České Budějovice, Czech Republic

[*] Corresponding author:

E-mail: marek.elias@osu.cz (ME)

Abstract

The non-photosyntetic alga *Euglena longa* harbours a cryptic plastid of unknown function. By a combination of bioinformatic and biochemical approaches we found out that this organelle houses a surprising set of metabolic processes. Biosynthesis of isoprenoid precursors and fatty acids is absent and the tetrapyrrole pathway is incomplete, whereas phospholipids and glycolipids are still being produced in the *E. longa* plastid. Unprecedented among non-photosynthetic plastids is the ability of this organelle to make tocopherols and a phylloquinone derivative. The most striking attribute is the presence of a linearized Calvin-Benson (CB) pathway including RuBisCO, together with ferredoxin-NADP$^+$ reductase and the ferredoxin/thioredoxin system. We hypothesize that the linear CB pathway is regulated by the redox status of the *E. longa* cell, in effect functioning as a redox valve bypassing the glycolytic oxidation of glyceraldehyde-3-phosphate to 3-phosphoglycerate. Altogether, the *E. longa* plastid defines a new class of relic plastids.

Abbreviations: ACP, acyl carrier protein; CBC, Calvin-Benson cycle; FAS, fatty acid synthesis; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; IPP, isopentenyl pyrophosphate; MEP, 2-C-methyl-d-erythritol 4-phosphate; MGDG/DGDG mono-/digalactosyldiacylglycerol; MVA, mevalonate; OH-PhQ, 5'-monohydroxyphylloquinone; PhQ, phylloquinone; SQDG, sulfoquinovosyldiacylglycerol; SP, signal peptide; TMD, transmembrane domain

Introduction

Plastids are organelles that evolved from a cyanobacterial endosymbiont in the ancestor of Archaeplastida and later found their way into other eukaryotic lineages by secondary or even higher-order endosymbioses (Keeling 2013; McFadden 2014; Ponce-Toledo et al. 2017). Photosynthetic harvesting of solar energy is supposedly the primary metabolic function and evolutionary advantage of plastid acquisition. However, plastids also host a variety of other metabolic pathways, such as biosynthesis of amino and fatty acids, isopentenyl pyrophosphate (IPP) and its derivatives (isoprenoids), and tetrapyrroles (Neuhaus and Emes 2000; Oborník and Green 2005; Van Dingenen et al. 2016). Hence, reversion of photosynthetic lineages to heterotrophy typically does not imply plastid loss, and non-photosynthetic plastids are found in many eukaryotic lineages (Wilson et al. 1996; Sanchez-Puerta et al. 2007; Slamovits and Keeling 2008; Janouškovec et al. 2015; Kamikawa et al. 2017; Hadariová et al. 2018). In these cases, metabolic integration

55   of the plastid presumably resulted in the host biochemistry being dependent on compound(s) supplied by

56   the organelle (Oborník et al. 2009; Lim and McFadden 2010; Janouškovec et al. 2015; Hadariová et al.

57   2018).

58   The most extensively studied relic plastid is undoubtedly the apicoplast of apicomplexan parasites

(especially of *Plasmodium falciparum* and *Toxoplasma gondii*). The knowledge of the apicoplast has

expanded tremendously since its discovery two decades ago (McFadden and Yeh 2017). The apicoplast is

ultimately derived from a red alga (Williamson et al. 1994; Moore et al. 2008; Janouškovec et al. 2010),

with an ochrophyte alga being a possible direct donor of the plastids in apicomplexans and their

photosynthetic relatives, chromerids (Ševčíková et al. 2015; Füssy and Oborník 2017). Except for the

recently characterized corallicolid apicomplexans (Kwong et al. 2019), the apicoplast genome contains only

genes related to gene expression, protein turnover, and assembly of FeS clusters, not considering a few short

hypothetical open reading frames of unknown function (Wilson et al. 1996; Janouškovec et al. 2015). The

essentiality of the apicoplast for parasite survival has attracted much attention, partly because this organelle

is a promising target for parasite-specific inhibitors (e.g. Miller et al. 2013; McFadden and Yeh 2017). So

far, three plastid pathways (encoded by the nuclear genome) seem to condition the apicoplast retention: non-

mevalonate IPP synthesis, haem synthesis, and type II fatty acid synthesis (FASII). IPP biosynthesis in

particular is vital for the bloodstream form of *P. falciparum* (Yeh and DeRisi 2011). On the other hand, the

mosquito and liver stages of *P. falciparum* are dependent on haem synthesis (Nagaraj et al. 2013), while

FASII is indispensable for pellicle formation in *Toxoplasma gondii* (Ke et al. 2014; Martins-Duarte et al.

2016).

75   Less is known about the actual metabolic functions of plastids in other non-photosynthetic algal

lineages. Many of them have a similar metabolic capacity as the apicoplast (Sanchez-Puerta et al. 2007;

Slamovits and Keeling 2008; Fernández Robledo et al. 2011), and some house an even more complex

metabolism that includes amino acid biosynthesis and carbohydrate metabolism pathways (Borza et al.

2005; Pombert et al. 2014; Smith and Lee 2014). Until recently, IPP synthesis seemed to be a process

conserved even in the most reduced relic plastids, such as the genome-lacking plastids of certain alveolates

(Matsuzaki et al. 2008; Janouškovec et al. 2015), but non-photosynthetic plastids lacking the characteristic

plastidial (MEP or DOXP) pathway of IPP biosynthesis are now known (Kamikawa et al. 2017; Graupner

et al. 2018; Dorrell et al. 2019). Thus, there is generally a metabolic reason for a plastid retention, although

the cases of plastid dependency differ between lineages.

85   An interesting group to study non-photosynthetic plastids are the euglenophytes. Like their prime

representative *Euglena gracilis*, most euglenophytes are mixotrophs containing a complex three-

membranes-bound photosynthetic plastid derived from a green alga belonging to Pyramimonadales (Turmel

et al. 2009; Leander et al. 2017; Jackson et al. 2018). However, non-photosynthetic mutants of *E. gracilis*,

induced by an antibiotic or mutagenic treatment, are often capable of heterotrophic living even after
presumed plastid loss (reviewed in Krajčovič et al. 2002; Hadariová et al. 2018). This might be enabled by
metabolic independence of the *E. gracilis* cell on the plastid. For instance, *E. gracilis* was shown to possess
two parallel haem synthesis pathways, one located in the mitochondrion/cytosol and another in the plastid
(Weinstein and Beale 1983). In the long run, this redundancy might predestine one of the pathways for loss,
provided that the other can efficiently supply the end-product to all compartments requiring it (Kořený and
Oborník 2011; Cihlář et al. 2016). On the other hand, localization of an essential pathway or its part
exclusively into the plastid will impose essentiality of the organelle as such.

Several lineages of euglenophytes independently became secondarily heterotrophic, but evidence
for the presence of a plastid has been provided only for *Euglena longa* (originally called *Astasia longa*), a
close relative of *E. gracilis* (Marin et al. 2003; Nudelman et al. 2003). The documentation of the organelle
at the cytological level is spurious (Webster et al. 1967; Kivic and Vesk 1974; Hachtel 1996), but the
complete plastid genome sequence was reported nearly two decades ago (Gockel and Hachtel 2000). As
expected, it lacks all the photosynthesis-related genes, except for *rbcL* encoding the large subunit of
ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). We recently documented the existence of a
nucleus-encoded small subunit (RBCS) of the *E. longa* RuBisCO enzyme synthesized as a precursor
polyprotein, but its processing into monomers could not be demonstrated (Záhonová et al. 2016). The
physiological role of the *E. longa* RuBisCO and the whole plastid remains unknown, although indirect
evidence suggests that the plastid is essential for *E. longa* survival (Siemeister et al. 1990a, b; Gockel et al.
1994; Hadariová et al. 2017).

To provide a key resource for investing the biology of *E. longa* and its plastid, we recently generated
a transcriptome assembly for this species and demonstrated that it is complete and directly comparable to
the transcriptome assemblies reported for *E. gracilis* (Záhonová et al. 2018). Evaluation of a set of high-
confidence candidates for plastid-targeted proteins enabled us to conclude that nucleus-encoded plastidial
proteins in *E. longa* employ N-terminal targeting presequences of the same two characteristic classes, as
known from *E. gracilis*. The *E. longa* transcriptome revealed various unusual features of the plastid
biogenesis and maintenance machinery shared with photosynthetic euglenophytes, but also supported the
lack of the photosynthesis-related machinery and suggested specific reductions of several plastidial house-
keeping functions presumably reflecting the loss of photosynthesis (Záhonová et al. 2018). However, the
repertoire of anabolic and catabolic pathways localized to the *E. longa* colourless plastid has not been
investigated and is the subject of the present paper.

To chart the main paths of the metabolic map of the *E. longa* plastids, we searched for homologs of
enzymes underpinning pathways known from plastids of other species. The reconstruction was greatly
facilitated by the recent characterization of the *E. gracilis* plastid metabolic network based on a proteomic

analysis of the organelle (Novák Vanclová et al. 2019). N-terminal regions of the candidates were evaluated

for characteristics of presequences predicting a specific subcellular localization to distinguish those likely

representing plastid-targeted proteins from enzymes located to other parts of the cell. Some of the

bioinformatics predictions were further tested by biochemical analyses. Our study provides the first

comprehensive view of a non-photosynthetic secondary plastid of green-algal origin and shows that the

metabolic capacity of the *E. longa* plastid is strikingly different from those of the apicoplast and other relic

plastids characterized in sufficient detail.

Results and Discussion

***E. longa* plastid lacks the MEP pathway of IPP biosynthesis, yet has kept the production of tocopherol**

and a phylloquinone derivative

Two parallel pathways of IPP biosynthesis exist in *E. gracilis* (Kim et al. 2004): the mevalonate (MVA)

pathway localized to the mitochondrion (first three enzymes) and the cytosol (the rest), and the plastid-

localized 2-C-methyl-D-erythritol (MEP) pathway, but only enzymes of the MVA pathway were found in

E. longa (Table S1, Fig. 1A). *E. longa* thus joins the group of recently discovered plastid-bearing eukaryotes

lacking the MEP pathway, namely the colourless diatom *Nitzschia* sp. NIES-3581 (Kamikawa et al. 2017)

and various colourless chrysophytes (Graupner et al. 2018; Dorrell et al. 2019). In contrast, the plastid-

localized MEP pathway in apicomplexans and related alveolates (i.e. Myzozoa) is essential as a source of

precursors for the synthesis of all cellular isoprenoids, since the cytosolic MVA pathway was lost from this

group (Janouškovec et al. 2015; Waller et al. 2016). The retention of the MEP pathway in the colourless

plastids of diverse non-photosynthetic chlorophytes (Figueroa-Martinez et al. 2015) is similarly explained

by the loss of the MVA pathway in this group (Lohr et al. 2012).

The MEP pathway in *E. gracilis* provides precursors for the synthesis in the plastid of terpenoid

compounds connected to photosynthesis, namely carotenoids and plastoquinone (Kim et al. 2004; Novák

Vanclová et al. 2019). As expected, the respective enzymes are all missing from *E. longa*. However, we

were surprised to find out that the *E. longa* plastid appears still involved in terpenoid metabolism,

specifically in its phytol branch. Photosynthetic eukaryotes generally produce three types of phytol

derivatives, tocopherols (vitamin E), phylloquinone (PhQ; vitamin K_1) and chlorophyll (its phytyl chain),

starting with a common precursor phytyl-PP, which is (directly or indirectly via salvage of phytol liberated

by chlorophyll degradation) made by reduction of geranylgeranyl-PP synthesized in the plastid by the MEP

pathway (Gutbrod et al. 2019). *E. gracilis* proved to be unusual not only because of lacking the conventional

geranylgeranyl-PP reductase (Novák Vanclová et al. 2019), but also for making phytol from precursors

provided by the MVA pathway (Disch et al. 1998; Kim et al. 2004). The route of phytol synthesis from the

cytosolic isoprenoids is currently unknown, though it was proposed that phytyl-PP is synthesized in the *E.*
gracilis plastid exclusively by step-wise phosphorylation of phytol by phytol kinase (VTE5) and phytyl
phosphate kinase (VTE6), which are enzymes normally employed for recycling phytol from chlorophyll
degradation (Novák Vanclová et al. 2019). This scheme is supported by the fact that *E. longa* has retained
VTE5 as well as VTE6, both proteins being highly similar to their *E. gracilis* orthologs and exhibiting
putative plastid targeting presequences (Fig. S1; Table S1). Since *E. longa* lacks chlorophyll, these two
enzymes must have a function independent of phytol recycling.

In addition of its role in chlorophyll synthesis, phytyl-PP is used by *E. gracilis* to make tocopherols
and a PhQ derivative, 5'-monohydroxyphylloquinone (OH-PhQ; Ziegler et al. 1989; Watanabe et al. 2017;
Novák Vanclová et al. 2019). All four enzymes mediating synthesis of α-tocopherol from phytyl-PP and
homogentisate were identified in *E. gracilis* and are localized to its plastid (Novák Vanclová et al. 2019).
Interestingly, orthologs of all four enzymes are found in *E. longa* as well, all with a typical plastid-targeting
presequence or at least with the N-terminal region being highly similar to the *E. gracilis* counterpart (Table
S1), consistent with their presumed plastidial localization (Fig. 1A). Homogentisate itself is apparently made
outside the plastid, most likely in the mitochondrion, as the enzyme responsible for its synthesis (4-
hydroxyphenylpyruvate dioxygenase) is not found in the *E. gracilis* plastid proteome and the respective
proteins have a predicted mitochondrial transit peptide in both *E. gracilis* and *E. longa* (Table S1). Our
analysis thus predicts that like its photosynthetic cousin, *E. longa* produces α-tocopherol. To test this
directly, we analysed extracts from *E. longa* by means of HPLC-MS/MS. For comparison, we employed
samples from *E. gracilis* grown at two different conditions (in light and in darkness). Tocopherols were
detected in both species (Fig. 1B), with α-tocopherol being the dominant form present in equivalent amounts
in all three samples (Fig. 1C). The signals of β- and/or γ-tocopherol (indistinguishable by our method) and
of δ-tocopherol suggest that tocopherol cyclase, and possibly also tocopherol O-methyltransferase, of both
Euglena species can process substrates with or without the 3-methyl group on the benzene ring (Fig. S2).

The synthesis of OH-PhQ in *E. gracilis* is understood only partially, with only three enzymes of the
pathway previously identified at the molecular level: the large multifunctional protein PHYLLO, apparently
localized to the cytosol and catalysing the first four steps leading to o-succinylbenzoate; MenA catalysing
phytylation of dihydroxynaphthoate localized in the plastid; and MenG (demethylnaphthoquinone
methyltransferase), possessing a typical N-terminal plastid-targeting presequence but not directly confirmed
as plastidial by proteomics (Novák Vanclová et al. 2019). Strikingly, *E. longa* expresses homologs of these
three *E. gracilis* proteins, although at low levels resulting in a low RNA-seq read coverage and thus
incomplete or inaccurate assembly of two of the respective transcripts (the PHYLLO sequence split into
several contigs, the MenA sequence with a frameshift corrected by RT-PCR; see Materials and Methods).
Nevertheless, N-terminal parts of both sequences were intact and confirmed the same subcellular

localization as in *E. gracilis* (Fig. 1A, Table S1). In agreement with these insights, OH-PhQ could be
detected in an extract from the *E. longa* culture (Fig. 1D, Fig. S3), although its abundance was smaller
compared to that in *E. gracilis* by an order of magnitude (Fig. 1E).

These findings document another unexpected function of the *E. longa* plastid, OH-PhQ synthesis,
albeit the plastid-associated part of the pathway cannot be presently reconstructed in full detail. One
uncertainty concerns the middle steps of the pathway, since like *E. gracilis* (see Novák Vanclová et al.
2019), *E. longa* also lack homologs of the conventional enzymes that are responsible for converting *o*-
succinylbenzoate to dihydroxynaphthoate and localized (at least in eukaryotes studied in sufficient detail)
in the peroxisome (Cenci et al. 2018). In contrast, tentative evidence was presented for the association of
the respective enzyme activities (presumably corresponding to enzymes non-orthologous to the
conventional ones) with the plastid envelope in *E. gracilis* (Seeger and Bentley 1991), raising a possibility
of a similar arrangement in *E. longa*. Secondly, the molecular identity of the putative hydroxylase catalysing
the final step of OH-PhQ synthesis is unknown, so its plastidial localization in *E. gracilis* or *E. longa* cannot
be ascertained. It is, nevertheless, likely given the fact that OH-PhQ is primarily needed in the plastid, at
least in photosynthetic eukaryotes (Ziegler et al. 1989). Thirdly, a previously unknown step – reduction of
the naphthoquinone ring – was recently demonstrated as a prerequisite for the reaction catalysed by MenG
to proceed in plants and cyanobacteria (Fatihi et al. 2015). The respective reductase is well conserved among
diverse plant and algal groups as well as cyanobacteria (Cenci et al. 2018), but we did not identify its close
homologs in any of the euglenophytes transcriptome assemblies, suggesting that euglenophytes employ an
alternative enzyme that yet needs to be characterized.

E. longa seems to be the first eukaryote with a non-photosynthetic plastid documented to have
retained the pathways (or final parts thereof) for tocopherols and OH-PhQ synthesis localized in the
organelle. The presence of tocopherols in *E. longa* is, however, not that surprising, as their function is not
restricted to photosynthetic tissues in plants and were detected also in bleached (i.e. non-photosynthetic) *E.*
gracilis mutants (Maeda and DellaPenna 2007; Watanabe et al. 2017). As potent lipophilic antioxidants
involved especially in membrane protection from damage caused by lipid peroxidation, tocopherols might
be used by *E. longa* as part of its protective mechanisms against reactive oxygen species generated by the
action of mitochondria and peroxisomes. The retention of OH-PhQ synthesis in *E. longa* may seem more
puzzling, given the fact that the best-established role of (OH-)PhQ in plants and algae is its functioning as
an electron carrier within the photosystem I (Ziegler et al. 1989; Brettel 1997). PhQ was additionally
proposed to serve as an electron acceptor in the process of disulfide bond formation in thylakoid lumenal
proteins required for proper function of photosystem II (Furt et al. 2010; Karamoko et al. 2011). A homolog
of the respective oxidoreductase (LTO1) exists in *E. gracilis* (Table S1), but not in the transcriptome data
from *E. longa*, consistent with the lack of photosystem II, and perhaps thylakoids entirely in the latter

species. Interestingly, PhQ was detected in the plasma membrane in plant tissues and proposed to be

involved in photosynthesis-unrelated redox processes at the cell surface (Lochner et al. 2003; Schopfer et

al. 2008). PhQ may even be synthesised directly in the plasma membrane, as a recent report from plants has

documented the existence of alternative forms of the terminal enzymes of PhQ biosynthesis that result from

alternative splicing and are localized to the plasma membrane rather than the plastid (Gu et al. 2018).

However, the protein sequences of both MenA and MenG enzymes in *E. longa* carry typical plastid-targeting

presequences, so they are unlikely to operate in the plasma membrane. We thus propose that OH-PhQ in *E.*

longa is involved in a hitherto uncharacterized, photosynthesis-unrelated plastid-resident process.

Figure 1: IPP and terpenoid-quinone biosynthesis in *E. longa* and its phototrophic relative *E. gracilis*.

A: Schematic comparison of the localization and evolutionary origin of enzymes (see colour-coding graphical legend below the "cells"). Abbreviations, IPP synthesis: ACAT – acetyl-CoA acetyltransferase, CDP-ME – 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol, CDP-MEP – 2-phospho-CDP-ME, CMK – CDP-ME kinase, CMS – CDP-ME synthase, DMAPP – dimethylallyl diphosphate, DXP – 1-deoxy-D-xylulose 5-phosphate, DXR – DXP reductase, DXS – DXP synthase, FPP – farnesyl siphosphate synthase, GGPS – geranylgeranyl-diphosphate synthase, HDR – HMB-PP reductase, HDS – HMB-PP synthase, HMB-PP – 4-hydroxy-3-methylbut-2-en-1-yl diphosphate, HMG-CoA – 3-hydroxy-3-methylglutaryl-CoA, HMGCR – HMG-CoA reductase, HMGCS – HMG-CoA synthase, IDI – isopentenyl-diphosphate delta-isomerase, MCS – MEcPP synthase, MDD – mevalonate-diphosphate decarboxylase, MEcPP – 2-C-methyl-D-erythritol 2,4-cyclodiphosphate, MEP – 2-C-methyl-D-erythritol 4-phosphate, MVK – mevalonate kinase, PMVK – phosphomevalonate kinase, PPS – unspecified polyprenyl-diphosphate synthase, ? – unclear substrate; Terpenoid-quinone synthesis: 4HPP – 4-hydroxyphenylpyruvate, HPD – hydroxyphenylpyruvate dioxygenase, HPT – homogentisate phytyltransferase, MMT – MPBQ/MPSQ methyltransferase, TAT – tyrosine aminotransferase, TC – tocopherol cyclase, TMT – tocopherol-O-methyltransferase, VTE5 – phytyl kinase, VTE6 – phytyl-phosphate kinase. **B:** MS/MS spectrum record of *E. longa* α-tocopherol and the proposed fragmentation pattern in positive ionization mode (inset). Monoisotopic masses of particular fragments were obtained by simulation in Xcalibur software. **C:** Semiquantitative comparison of tocopherol species in *E. longa*, heterotrophically (dark) grown *E. gracilis* and autotrophic *E. gracilis*. **D and E:** MS/MS spectrum record of *E. longa* 5-hydroxyphylloquinone and the proposed fragmentation pattern in positive ionization mode (inset); semiquantitative comparison of 5-hydroxyphylloquinone.

E. longa plastid plays a limited role in the metabolism of nitrogen-containing compounds

Some of the apparent oddities of the *E. longa* plastid do not stem from the loss of photosynthesis in this species but reflect unusual features of the plastid in euglenophytes in general. These particularly concern plastid functions in the metabolism of nitrogen-containing compounds. Plastid is commonly involved in nitrogen assimilation due to housing nitrite reductase (Giordano and Raven 2014; Sanz-Luque et al. 2015), but it was established a long time ago that *E. gracilis* cannot assimilate nitrate or nitrite (Oda et al. 1979; Kitaoka et al. 1989) and, accordingly, no nitrite reductase can be identified in the transcriptome data from this species and *E. longa*. The plastids of both *Euglena* species apparently also lack the enzymes working immediately downstream, i.e. glutamine synthetase and glutamine oxoglutarate aminotransferase (the GS/GOGAT system common in plastids of other groups; Fernandez and Galvan 2008; Dagenais-Bellefeuille and Morse 2013), indicating that the plastids rely on the import of organic nitrogen from other parts of the cell, like recently proposed for the plastid in chromerids (Füssy et al. 2019).

One of the most surprising insights of the recent proteomics-aided analysis of the *E. gracilis* plastid
metabolism was the paucity of pathways concerning amino acids (Novák Vanclová et al. 2019). *E. longa* is
apparently even more extreme in this regard, because it lacks counterparts of the (predicted or proteomically
verified) plastid-targeted forms of serine biosynthesis enzymes found previously in *E. gracilis*, i.e.
phosphoglycerate dehydrogenase and phosphoserine phosphatase. Thus, we could localize only two
elements of amino acid biosynthesis pathways to the *E. longa* plastid (Fig. S4): serine/glycine
hydroxymethyltransferase, whose obvious role is to provide the one-carbon moiety for formylmethionyl-
tRNA synthesis required for the plastidial translation; and one of the multiple isoforms of cysteine synthase
A, which (like in *E. gracilis*) apparently relies on O-acetyl-L-serine synthesized outside the plastid, due to
the absence of a discernible plastid-localized serine O-acetyltransferase (see Novák Vanclová et al. 2019,
and Table S2). This is not due to a general reduction of amino acid metabolism in *E. longa* or the
incompleteness of the sequence data, as its transcriptome assembly includes homologs of enzymes required
for the synthesis of all 20 proteinogenic amino acids, but the respective proteins have predicted localization
in compartments other than the plastid (Table S2).

Amino acids in the plastid are not only substrates of protein synthesis, but also serve as precursors
or nitrogen donors for the synthesis of various other compounds (Moffatt and Ashihara 2002; Gerdes et al.
2012). One such pathway, described in detail in the subsequent section, leads to tetrapyrroles. In contrast,
the spectrum of reactions related to the metabolism of other nitrogen-containing cofactors or their precursors
(B vitamins) is very limited in the plastids of both *Euglena* species. We identified only one such candidate
in *E. longa* – the reaction of vitamin B6 salvage catalysed by pyridoxamine 5'-phosphate oxidase, whereas
E. gracilis additionally expresses two plastid-targeted isoforms of pyridoxine 4-dehydrogenase (Table S3).
De novo synthesis or salvage of purines and pyrimidines is also absent from the plastid of both *Euglena*
species, except for one step present in *E. gracilis* but apparently not *E. longa*: the former exhibits two forms
of CTP synthase, one presumably cytosolic and another with a plastid-targeting presequence and found in
the plastid proteome reported by Novák Vanclová et al. 2019, whereas *E. longa* expresses an ortholog of
only the cytosolic version (Table S3). The lack of a separate plastidial CTP source in *E. longa* may reflect
the presumably lower magnitude of RNA synthesis (and possibly also less extensive phospholipid synthesis
requiring CTP in the reaction catalysed by CDP-diacylglycerol pyrophosphatase) in its plastid. Finally, *E.*
longa possesses an ortholog of an enzyme involved in the synthesis of polyamines (spermidine synthase)
found in the plastid proteome of *E. gracilis* (Novák Vanclová et al. 2019), but (in contrast to the *E. gracilis*
protein) its N-terminus does not fit the characteristics of a plastid-targeting presequence, suggesting that its
subcellular localization may be outside of the plastid (Fig. S4). However, we found out that both *E. longa*
and *E. gracilis* have another version of this enzyme with an obvious plastid-targeting-like presequence in
both species (Table S3), so we cannot rule out the possibility that polyamines are produced in the *E. longa*

plastid after all.

A residual tetrapyrrole biosynthesis pathway of unclear function is retained in the *E. longa* plastid

Most plastid-bearing eukaryotes synthesize protoporphyrin IX, the common precursor of haem and

chlorophyll, via a single pathway wholly or mostly localized to the plastid (Oborník and Green 2005; Cihlář

et al. 2016; Füssy and Oborník 2017). *E. gracilis* is one of the known exceptions, because it possesses two

independent protoporphyrin synthesis pathways, a mitochondrial-cytosolic and a plastid one (Fig. 2A;

Weinstein and Beale 1983; Kořený and Oborník 2011; Lakey and Triemer 2016). Whereas all enzymes of

the plastid pathway were identified at the sequence level, previous studies left some gaps in the enzyme

assignment to the different steps of the mitochondrial-cytosolic pathway (Kořený and Oborník 2011; Lakey

and Triemer 2016). The new sequence data from *E. gracilis* has now enabled us to identify all three missing

enzymes of the mitochondrial-cytosolic pathway; specifically novel, apparently cytosolic, isoforms of

uroporphyrinogen-III synthase (UROS) and uroporphyrinogen decarboxylase (UROD), and two variants of

oxygen-independent coproporphyrinogen oxidase (CPOXi), one cytosolic and one perhaps localized to the

mitochondrion. In addition, we found a third UROS homolog that carries a plastid-targeting presequence

and constitutes a divergent sister group to the previously known euglenophyte plastidial UROS isoform

(Fig. S5; Table S4). Indeed, it was detected in the experimentally determined plastid proteome, together

with the previously known plastidial UROS isoform (protein IDs 16898 and 15143 in Novák Vanclová et

al. 2019), but the functional significance of the plastidial UROS duplication remains unclear.

Our transcriptome data from *E. longa* revealed orthologs of all *E. gracilis* enzymes of the

mitochondrial-cytosolic pathway with the same predicted subcellular localization, barring a single enzyme

with an incomplete N-terminal sequence precluding confident prediction (Fig. 2A; Table S4). In contrast,

orthologs of only six of the *E. gracilis* plastid-targeted enzymes could be identified in the transcriptome of

E. longa: aminolevulinic acid dehydratase (ALAD), the two UROS isoforms, one of the UROD isoforms,

one isoform of CPOXi, and protoporphyrinogen oxidase (PPOX). The first three proteins have putative

plastid-targeting presequences, much like their *E. gracilis* orthologs. However, the respective CPOXi

protein has a strong mitochondrial targeting presequence, whereas the UROD protein is devoid of any

presequence and the presequence of the PPOX protein is markedly shorter and lacks the characteristics of a

plastid-targeting signal (Fig. 2B; Table S4). This suggests that only ALAD and the two UROS isoforms are

targeted to the plastid in *E. longa*, whereas the other enzymes have been retargeted to the cytosol or

mitochondrion in the evolution of the *E. longa* lineage. Strikingly, a putative *E. longa* homolog of the *E.*

gracilis plastid-localized porphobilinogen deaminase (PBGD), an enzyme for the reaction between those

catalysed by ALAD and UROS, was previously detected at both the RNA (by a Northern blot) and protein

level (by an immunoblot; Shashidhara and Smith 1991), but we could not identify the respective transcript

in the *E. longa* transcriptome data even when the raw reads were searched.

Intact isolated *E. gracilis* plastids are capable of chlorophyll synthesis from externally supplied 5-

aminolevulinic acid (ALA; Gomez-Silva et al. 1985), so it is possible that the *E. longa* plastid can

analogously import ALA from the cytoplasm, thus providing a substrate for the plastid-localized ALAD

isoform (Fig. 2A). The surprising absence of the previously documented plastid-localized PBGD in our

transcriptome data may be potentially accounted for by the lack of expression of the respective enzyme at

culture conditions employed by us. Hence, the pathway may theoretically proceed up to uroporphyrinogen

III (the product of the UROS enzyme) in the *E. longa* plastid, but the absence of enzymes for further

processing of this compound towards haem and/or chlorophyll (consistent with the lack of photosystems

and the cytochrome b_6f complex) indicates that it is consumed by another process.

Uroporphyrinogen III is indeed at the beginning of a separate branch converting it in three steps to

sirohaem (Dailey et al. 2017), which serves as a cofactor of several enzymes, including nitrite reductase and

sulfite reductase (Tripathy et al. 2010). The former enzyme is missing from the *Euglena* spp. (see above),

but sulfite reductase is present in both *E. longa* and *E. gracilis* (Table S4) and was detected in the plastid

proteome of the latter species (Novák Vanclová et al. 2019). Hence, sirohaem synthesis in the plastid to

support the function of sulfite reductase is an attractive hypothesis that might explain the retention of some

of the tetrapyrrole synthesis enzymes in the *E. longa* plastid. However, analyses of the transcriptome data

alone do not provide any clear answer as to whether this hypothesis is valid. Of the various alternative

enzymes mediating sirohaem synthesis in various taxa (Tripathy et al. 2010), only two have confidently

identified homologs in both *Euglena* spp. (Table S4). One of them is the multifunctional protein CysG,

which catalyses all reactions from uroporphyrinogen III to sirohaem, but the respective *Euglena* homolog

is devoid of any presequence in either species and is thus most likely cytosolic (Table S4). The other is a

stand-alone version of the enzyme (uroporphyrinogen-III C-methyltransferase) catalysing the first step of

sirohaem synthesis, called CobA or SirA (Table S4). The *Euglena* homologs exhibit a predicted signal

peptide, which is however immediately followed by a region belonging to the mature enzyme, so it is unclear

whether this protein is routed to the plastid (indeed it is not among the proteomically confirmed plastid

protein in *E. gracilis*). Hence, the role of the enzymes of the tetrapyrrole biosynthesis pathway in the *E.*

longa plastid remains unexplained.

Figure 2: Tetrapyrrole biosynthesis in *E. longa* and *E. gracilis*. A: Schematic comparison of the localization and evolutionary origin of enzymes (see colour-coding graphical legend). Abbreviations: ALAD – delta-aminolevulinate dehydrogenase, ALAS – delta-aminolevulinate synthase, CobA –

uroporphyrinogen-III C-methyltransferase, CPOX – coproporphyrinogen III oxidase, CPOXi – oxygen-independent CPOX, CysG – trifunctional enzyme of sirohaem synthesis (see text), FeCH – ferrochelatase, GSA-AT – glutamate semialdehyde-aminomutase, GTR – glutamyl-tRNA reductase, PBGD – porphobilinogen deaminase, PPOX – protoporphyrinogen oxidase, UROD – uroporphyrinogen decarboxylase, UROS – uroporphyrinogen-III synthase. **B-C:** Sequence alignment of UROD (B) and PPOX (C) presequences, showing the loss of the plastid-targeting motifs in *E. longa*. These targeting motifs represented by transmembrane domains are marked with frames in other euglenophyte sequences.

E. longa plastid does not make fatty acids but maintains phospholipid and glycolipid synthesis

Eukaryotes synthesize even-chain fatty acids by a single large multi-modular fatty acid synthase I (FASI) in the cytosol or by a multi-enzyme type II fatty acid synthesis complex in the plastid. *E. gracilis* possesses both systems (Zimorski et al. 2017). In contrast, *E. longa* encodes only a homolog of the FASI enzyme (Table S5), whereas enzymes of the type II fatty acid synthesis are absent (Fig. 3A) except for the acyl carrier protein (ACP). The loss of plastid-localized fatty acid synthesis in *E. longa* is not without precedent, as it has been also reported for the apicoplast of *Theileria parva* (which is fully dependent on fatty acid supply from host), the plastid of *Perkinsus marinus* (Janouškovec et al. 2015) and the chrysophyte "*Spumella*" sp. NIES-1846 (Dorrell et al. 2019). Nevertheless, the *E. longa* plastid has kept plastid-targeted enzymatic steps downstream of fatty acid synthesis. These include ACP and 4'-phosphopantetheinyl transferases (or holo-ACP synthase) crucial for the synthesis of an active form of ACP, which serves as a carrier of acyl chains in phospholipid and glycolipid biosynthesis (Lambalot and Walsh 1995). Next, *E. longa* possesses predicted plastid-targeted homologs of acyl-ACP synthetases (presumably activating fatty acids imported into the plastid from outside) and all enzymes required for the synthesis of phosphatidic acid (PA) and its subsequent conversion to phosphatidylglycerol (PG) (Fig. 3A; Table S5). It is worth noting that *E. longa* also has a parallel, plastid-independent, route of phosphatidylglycerol synthesis (Table S6).

No other reactions of phospholipid synthesis or decomposition beyond PG synthesis seem to operate in the *E. longa* plastid, but interestingly, we found homologs of the enzymes of the synthesis of galactolipids monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG) in the *E. longa* transcriptome (Fig. 3A, Table S5). All the proteins have a predicted N-terminal plastid-targeting presequence, consistent with the plastidial localization of galactolipid synthesis in all other plastid-bearing eukaryotes studied so far (Yuzawa et al. 2012). In support of the bioinformatic predictions, both MGDG and DGDG could be detected in lipid extracts from *E. longa* and *E. gracilis*, although galactolipid levels were significantly lower in *E. longa* than in the control sample of *E. gracilis* (Fig. 3B). The presence of DGDG was further confirmed by immunofluorescence using an anti-DGDG antibody, which showed DGDG to be present in small foci in the *E. longa* cells (white arrowheads in Fig. 3C), presumably

representing individual small plastids. In comparison, most of the volume of the photosynthetic *E. gracilis*
cells was stained, whereas the negative control, the primary osmotrophic (i.e. plastid-lacking) euglenoid
Rhabdomonas costata, did not stain at all.

The presence of galactolipids in plastids is generally being explained by their essentiality for the
proper functioning of the photosynthetic apparatus, but this view has been challenged by demonstration that
cyanobacterial mutants lacking galactolipids can photosynthesize normally (Awai et al. 2001). A
photosynthesis-independent role of galactolipids in plastid biology was proposed, too. Transit peptides of
plastid-targeted proteins exhibit affinities for MGDG and DGDG in the plastid envelope (Pinnaduwage and
Bruce 1996), suggesting a direct role of these lipids in plastid protein import. The photosynthesis-
independent role of galactolipids is indicated not only by their presence in the *E. longa* plastid documented
here, but also by previous reports from other non-photosynthetic algae, including the diatom *Nitzschia alba*
(Anderson et al. 1978) and the chlorophyte *Prototheca wickerhamii* (Borza et al. 2005), and from non-
photosynthetic tissues of plants (Awai et al. 2001; Kobayashi 2016). On the other hand, the apicoplast (Botté
et al. 2008; Botté et al. 2013) and most likely also the relic plastid of *Helicosporidium* (based on our analysis
of the respective genome sequence data generated by Pombert et al. 2014) lack galactolipid synthesis
completely. The reason for the differential retention of galactolipids in different colourless plastids remains
unclear.

In addition to MGDG and DGDG, we identified in samples from both *Euglena* spp. another
common glycolipid characteristic for plastids, sulfoquinovosyldiacylglycerol (SQDG) (Fig. 3B) (Hori et al.
2016). The presence of SQDG in *E. longa* is also not unprecedented among non-photosynthetic plastid-
bearing eukaryotes; see, e.g., its documented occurrence in the diatom *N. alba* or the dinoflagellate *Oxyrrhis*
marina (Anderson et al. 1978; Goddard-Borger and Williams 2017; Yoon et al. 2017). Of the three enzymes
of SQDG biosynthesis normally localized to the plastid, we found in both *Euglena* species only that
catalysing the final step (sulfoquinovosyltransferase; Fig. 3A). Interestingly, the standard eukaryotic version
of the enzyme, SQD2, is present only in *E. gracilis*, but both species proved to share another isoform
phylogenetically affiliated to bacterial SqdX version of the enzyme (Fig. 4). To the best of our knowledge,
this is the first encounter of SqdX in any eukaryote. The presence of SQD2 only in *E. gracilis* probably
relates to specific needs of its photosynthetic plastid. Indeed, *E. gracilis* contains a much larger amount of
SQDG compared to *E. longa* (Fig. 3B) and the profile of esterified fatty acids differs between the two species
(*E. longa* lacks SQDG forms with unsaturated longer chains; Table S7).

Details of the synthesis of the saccharide moieties of glycolipids in *E. longa* are also worth
considering (Fig. 3A). *E. longa* exhibits an ortholog of the *E. gracilis* UDP-glucose epimerase previously
identified in the plastid proteome (Table S5), explaining the source of UDP-galactose for galactolipid
synthesis. This enzyme seems to have been acquired by euglenophytes from a bacterial source (Fig. S6),

but, interestingly, *E. gracilis* encodes another enzyme, which corresponds to the plastidial UDP-glucose

epimerase, also called PHD1, known from plants and various algae (Li et al. 2011). The *E. gracilis* PHD1

possesses a predicted plastid-targeting presequence (Table S5) and is thus also likely plastidial (although

this is not confirmed by proteomic data reported by Novák Vanclová et al. 2019). This putative redundancy

in UDP-galactose is apparently not shared by *E. longa*, possibly because of a presumably much lower need

for galactolipid synthesis (Fig. 3B). The origin of the SQDG precursor UDP-sulfoquinovose in *E. longa*

remains obscure, because as noticed before (Novák Vanclová et al. 2019), euglenophytes in general lack the

conventional UDP-sulfoquinovose synthase SQD1/SqdB and probably employ an alternative, unrelated

enzyme. UDP-glucose, i.e. the common precursor of both UDP-galactose and UDP-sulfoquinovose, is most

likely imported from the cytoplasm, owing to the absence in *E. longa* of a candidate plastid-targeted version

of any of the relevant enzymes (UDP-glucose/UDP-sugar pyrophosphorylase).

Figure 3: Fatty acid and lipid biosynthesis in *E. longa* and *E. gracilis*. A: Schematic comparison of the

localization and evolutionary origin of enzymes. Abbreviations, fatty acid synthesis: ACC – acetyl-CoA
carboxylase, ACS – acetyl-CoA synthetase, ENR – enoyl-CoA reductase, Fas1 – malonyl-CoA/acetyl-CoA:
ACP transacylase, FASI – type I fatty acid synthase, FAT – fatty acyl-ACP thioesterase, HD – hydroxyacyl-
ACP dehydratase, KAR – ketoacyl-ACP reductase, KAS – ketoacyl-ACP synthase, TE – fatty acid
thioesterase, TRX – thioredoxin-regulated enzyme; glycolipid synthesis: AAS – acyl-ACP synthase, ACPS
– holo-ACP synthase, AGP-AT – acylglycerophosphate acyltransferase, G3P-AT – glycerol-3-phosphate
acyltransferase, G3PDH – glycerol-3-phosphate dehydrogenase, MGDG/DGDG – mono-/digalactosyl
diacylglycerol, MGDGS/DGDGS – MGDG/DGDG synthase, PAP – phosphatidic acid phosphatase, SQD1
– UDP-sulfoquinovose synthase, SQD2/SQDX – sulfoquinovosyl diacylglycerol (SQDG) synthase,
UGE/PHD1 – UDP-glucose epimerase, UGP3 – UDP-glucose pyrophosphorylase 3; phospholipid
synthesis: CDS – CDP-diacylglycerol synthase, PGP1 – phosphatidylglycerophosphate synthase, PGP-P –
phosphateidylglycerophosphate phosphatase. **B:** Semiquantitative comparison of glycolipids present in *E.*
longa and autotrophic *E. gracilis*. Note the logarithmic scale of the quantification units (peak area). Peak
area is an arbitrary unit expressing the intensity of the signal of a particular lipid species, recalculated
according to their respective ionization promptitude. As each lipid species have different ionization
promptitude, note that direct comparison can be done only within lipid class (for details, see Tomčala et al.
2017). **C-E:** Immunofluorescence micrographs using anti-DGDG antibody (C), DAPI (D) and differential
interference contrast (E). Autotrophic *E. gracilis* represents a positive control, while the aplastidic
euglenozoan *R. costata* was used as negative control.

Figure 4: Euglenophytes have replaced the eukaryotic form of sulfoquinovosyltransferase (SQD2) with a bacterial version (SqdX). The maximum-likelihood tree was inferred with IQ-TREE using the LG+F+G4 substitution model and ultra-fast bootstrapping. The UFboot support values are indicated at branches when higher than 75%.

A linearized Calvin-Benson pathway in the *E. longa* plastid

The presence of genes encoding both subunits of the enzyme RuBisCO in *E. longa* (Záhonová et al. 2016) raises the question whether the Calvin-Benson cycle (CBC) as a whole has been preserved in this organism to provide ribulose-1,5-bisphosphate, the RuBisCO substrate. A putative *E. longa* plastid triose-phosphate isomerase was described previously (Sun et al. 2008). We confirmed not only this candidate, but additionally identified homologs with putative plastid-targeting presequences for almost all remaining CBC enzymes (Table S8). Our identification was confirmed by phylogenetic analyses (supplementary Dataset S1), which showed specific relationships of the *E. longa* proteins to the previously characterized CBC enzymes from other euglenophytes (Markunas and Triemer 2016). However, two key CBC enzymes are apparently missing from the *E. longa* transcriptome: phosphoglycerate kinase (ptPGK) and glyceraldehyde-phosphate dehydrogenase (ptGAPDH). The homologs of these enzymes present in *E. longa* are not orthologous to the plastid-targeted isoenzymes from other euglenophytes and all clearly lack a plastid-targeting presequence (Table S8). Hence, these are presumably cytosolic enzymes involved in glycolysis/gluconeogenesis. The lack of ptPGK and ptGAPDH in *E. longa* means that the product of the RuBisCO carboxylase activity, 3-phosphoglycerate (3PG), cannot be converted (via 2,3-bisphosphoglycerate; 2,3-BPG) to glyceraldehyde-3-phosphate (GA3P) in the plastid and the cycle becomes a linear pathway (Fig. 5).

Assuming that the reactions catalyzed by fructose bisphosphatase, phosphoribulokinase, and RuBisCO are not reversible (Raines and Lloyd 2001), the flux through this linearized CB pathway most likely goes from GA3P to 3PG, with a net production of six molecules of 3PG from five molecules of GA3P due to fixation of three CO_2 molecules catalysed by RuBisCO. We thus need to define the origin of GA3P entering the pathway. Euglenophytes do not store starch in the plastid (instead they have cytosolic paramylon as a storage polysaccharide; Kiss et al. 1987), and indeed we did not find any glucose metabolism-related enzymes in the predicted *E. longa* plastid proteome. Hence, GA3P cannot be produced by a glycolytic route in the *E. longa* plastid. The presence of the plastid-targeted glycerophosphate dehydrogenase (Table S5) in principle allows for generation of GA3P from glycerol-3-phosphate (via dihydroxyacetone phosphate; DHAP; Fig. 3), which could possibly come from the degradation of glycerolipids in the plastid. However, no phospholipid-degradation enzymes (phospholipases) appear to localize to the plastid in *E. longa*. Hence, the primary function of glycerophosphate dehydrogenase is perhaps to operate in the reverse direction, i.e. to provide glycerol-3-phosphate for the plastid phospholipid and glycolipid synthesis (see above). The *E. longa* plastid thus most likely imports GA3P or DHAP from the cytosol (Fig. 5). This assumption is supported by the presence of several members of the plastid phosphate translocator (pPT) family (Fig. S7; Facchinelli and Weber 2011), including one phylogenetically closest to a previously characterized cryptophyte transporter with a preference for DHAP (Haferkamp et al. 2006). Concerning the opposite end of the hypothesized linear CB pathway, we did not identify any

candidate *E. longa* plastid-targeted enzyme that would metabolize 3PG further (see the absence of 3PG dehydrogenase discussed above), so this intermediate is most likely exported from the plastid into the cytosol, probably also by one of the members of the pPT family of transporters (Fig. 5).

The operation of CBC is inherently linked with the oxygenase side-activity of RuBisCO, which converts ribulose-1,5-bisphosphate into 3PG and phosphoglycolate instead of two molecules of 3PG that are produced by the regular carboxylase activity (Tabita et al. 2007). Phosphoglycolate is metabolized in the photorespiration pathway, initiated by phosphoglycolate phosphatase yielding glycolate. We did find a candidate plastid-targeted phosphoglycolate phosphatase in *E. longa* (Table S8), orthologous to a protein detected in *E. gracilis* plastid proteome, suggesting that the *E. longa* RuBisCO has the oxygenase activity and phosphoglycolate is metabolized in this species. However, we did not find in our *E. longa* transcriptome assembly any discernible homolog of the recently characterized transporter PLGG1 mediating glycolate export from the plastid in the canonical photorespiratory pathway (Pick et al. 2013), although *E. gracilis* does have it (Table S8). Since there is no obvious candidate for a plastid-targeted glycolate-metabolizing enzyme (glycolate oxidase, glyoxylate reductase, glycolaldehyde dehydrogenase) in *E. longa*, it is unclear how glycolate is removed from the plastid of this species. It is possible that the amount of glycolate produced in the *E. longa* plastid is low and that it can be exported by an alternative (PLGG1-independent) route, whose existence has been proposed also for plant plastids (Walker et al. 2016) and which might be sufficient for glycolate recycling in the semi-parasitic plant *Cuscuta campestris* capable of low-efficiency photosynthesis (Vogel et al. 2018).

Figure 5: Carbon metabolism in the plastids of *E. longa* and *E. gracilis*. The Calvin-Benson cycle (CBC) resident to this organelle is central to the plastid carbon metabolism, regulated by the ferredoxin/thioredoxin (Fd/Trx) system. Reduction of disulfide bonds by the Fd/Trx system activates FBP and PRK. FTR and FD of the Fd/Trx system require for their function a post-translationally added Fe-S prosthetic group provided by the Fe-S assembly system. GapN apparently mediates shuttling of reducing equivalent (NADPH) through the exchange of DHAP/GA3P and 3PG, reflecting the cytosolic $NADPH/NADP^+$ ratio and thus an overall metabolic state of the cell. In contrast, *E. gracilis* plastid is an energy-converting organelle, harvesting light into chemical energy bound as NADPH and ATP and subsequently using this bound energy to fix CO_2 into organic carbohydrates via the CBC. Enzyme abbreviations are colour-coded according to their inferred evolutionary origin, see the graphical legend. Abbreviations, CBC: ALDO – aldolase, DHAP – dihydroxyacetone-phosphate, FBP – fructose-1,6-bisphosphatase, GAPDH – glyceraldehyde-3-phosphate dehydrogenase, PGK – 3-phosphoglygerate kinase, PGP – phosphoglycolate phosphatase, PLGG1 – plastid glycolate/glycerate transporter, PRK – phosphoribulokinase, RBCL/RBCS – RuBisCO large/small subunit, RCA – RuBisCO activase, RPE – ribulose-5-phosphate epimerase; RPIA – ribulose-phosphate isomerase A, SBP – sedoheptulose-1,7-bisphosphatase, TKTL – transketolase, TPI – triose-phosphate isomerase, TPT – triose-phosphate translocator; Fd/Trx system: FD – ferredoxin; FNR – FD/NADP+ oxidoreductase, FTR

550 – FD/TRX oxidoreductase, TRX – thioredoxin, ATPS – ATP synthase, ATPC – ADP/ATP translocase,
551 LHC – light-harvesting complex.

The system of redox regulation of the Calvin-Benson pathway is conserved in *E. longa*

Although the photosynthetic machinery, including all photosystem I subunits, is missing from *E. longa*
(Záhonová et al. 2018), we found homologs (with clear plastidial localization) of the typical
"photosynthetic" (PetF-related) ferredoxin (Fd) and ferredoxin-NADP$^+$ reductase (FNR) (Table S9). These
two proteins are primarily involved in passing electrons from an activated photosystem I to NADP$^+$, which
is thus reduced to NADPH, the main donor of reducing equivalents for anabolic reactions in the plastid
(above all, CO_2 fixation). Phylogenetic analysis of FNR homologs from euglenophytes revealed the
existence of two different, yet related, clades affiliated to FNR from green algae (Fig. 6). One clade
comprises the *E. longa* FNR plus its orthologs from photosynthetic euglenophytes (*E. gracilis* and two
Eutreptiella strains), whereas the second clade is restricted to the photosynthetic species. Two different FNR
forms also exist in plants, one functioning in photosynthesis (production of NADPH dependent on the
function of the photosystem I; marked with "P" in Fig. 6) and the other being a "non-photosynthetic"
homolog ("NP" in Fig. 6) that allows the electron flow in the reverse direction, from NADPH to Fd (Vollmer
et al. 2001). In analogy with plants, we suggest that the two euglenophyte FNR clades functionally differ,
with one (that lacking a representative in *E. longa*) serving in photosynthesis and the other (present in *E.*
longa) mediating light-independent production of reduced Fd.

Multiple plastid enzymes depend on reduced Fd as an electron donor, namely glutamate synthase,
lipid desaturases, nitrite reductase, and sulfite reductase (Neuhaus and Emes 2000). As discussed in previous
sections, glutamate synthase and nitrite reductase are missing from *E. longa*, whereas all identified lipid
desaturases are predicted as targeted to the mitochondrion or the ER (Table S5). A two-subunit sulfite
reductase seems to be present in the plastid of *E. gracilis* (Novák Vanclová et al. 2019) and both subunits
have highly similar homologs in *E. longa* (Table S4), but this form of the enzyme utilizes NADPH rather
than ferredoxin as electron donor (Patron et al. 2008).

Another crucial role of Fd in plastids is to provide electrons to ferredoxin:thioredoxin reductase
(FTR) mediating reduction of the protein thioredoxin (Trx). The Fd/Trx system regulates several plastid
CBC enzymes in response to the redox status in the stroma and, in extension, to the photosynthetic activity
of the plastid (Fig. 5). An excess of NADPH leads to electrons being relayed from reduced ferredoxin to the
Fd/Trx system, which eventually reduces certain disulfide bonds in the target enzymes, thus changing their
activity (Schürmann and Buchanan 2008). This ensures activation of the CBC only when the photosynthetic
machinery works properly. Notably, FTR and Trx homologs with evident plastid-targeting presequences are
both present in *E. longa* (Table S9). Specific motifs necessary for the function of the Fd/Trx system

(Schürmann and Buchanan 2008) are conserved in the respective *E. longa* proteins (Fig. S8), consistent with

the Fd/Trx system being functional in this species. In addition, two *E. longa* CB pathway enzymes, fructose

bisphosphatase (two of the three isoforms present) and phosphoribulokinase, exhibit the conserved Trx

regulatory cysteine motifs similar to their orthologs in *E. gracilis* (Fig. S8, Table S10). Thus, we suggest

that the CB pathway in *E. longa* is sensitive to the redox status in the plastid, specifically to the concentration

of NADPH (Fig. 5).

Figure 6: The inferred phylogeny of FNR. The maximum-likelihood tree was inferred with IQ-TREE

using the LG+F+G4 substitution model and ultra-fast bootstrapping. The UFboot support values are

indicated at branches when higher than 75%. Euglenophyte species are in bold, and their putative

photosynthetic and non-photosynthetic homologs are depicted. P, photosynthetic; NP, non-photosynthetic.

May the *E. longa* plastid be involved in keeping the redox balance of the cell?

How to interpret in functional terms the retention of the linearized CB pathway and its putative redox

regulation in the non-photosynthetic plastid of *E. longa*? The key role of the pathway is supported by the

fact that production of the large RuBisCO subunit seems to be the *raison d'être* for the preservation of the

plastid genome in *E. longa* (Záhonová et al. 2016). The presence of CB enzymes in a non-photosynthetic

plastid is not without precedent, as it has been reported from a set of unrelated colourless algae and plants.

Some of them, e.g. the dinoflagellate *Crypthecodinium cohnii*, the dictyochophytes *Pteridomonas danica*

and *Ciliophrys infusionum*, the cryptophyte *Cryptomonas paramecium*, and some parasitic or

mycoheterotrophic land plants, are known to encode RuBisCO (Sekiguchi et al. 2002; Sanchez-Puerta et al.

2007; Donaher et al. 2009; Wicke et al. 2013; Hadariová et al. 2018), but how complete is the complement

of other CBC enzymes in these species is unknown. In contrast, transcriptomic or genomic analyses of other

colourless plastid-bearing taxa, such as the dinoflagellate *Pfiesteria piscicida*, the green alga

Helicosporidium sp. ATCC50920, the diatom *Nitzschia* sp. NIES-3581, and the non-photosynthetic

chrysophytes, revealed the presence of a subset of CB enzymes, including ptPGK and ptGAPDH, but not

of RuBisCO (Kim et al. 2013; Pombert et al. 2014; Kamikawa et al. 2017; Graupner et al. 2018). Hence,

the constellation of the CB enzymes retained in the *E. longa* plastid seems to be unique.

The CBC enzymes retained in various non-photosynthetic eukaryotes obviously do not serve to

sustain autotrophic growth, as ATP and the reducing power generated by photosynthesis are unavailable.

The CB pathway in *Nitzschia* sp. NIES-3581 was proposed to serve as a source of erythrose-4-P for the

synthesis of aromatic amino acid via the shikimate pathway in the plastid (Kamikawa et al. 2017). Although

not discussed in the respective report (Pombert et al. 2014), the CB pathway in the *Helicosporidium* plastid

may likewise serve to feed the co-localized shikimate pathway with erythrose-4-P. However, such a

rationalization of the CB pathway in the *E. longa* plastid would not work, since enzymes for aromatic amino

acid biosynthesis in this species are apparently localized to the cytosol (Table S2) and thus have access to

erythrose-4-P produced by the pentose phosphate pathway. In addition, the need to produce erythrose-4-P

in the *E. longa* plastid would not explain the retention of RuBisCO (absent in both *Nitzschia* and

Helicosporidium). A photosynthesis- and CBC-independent role of RuBisCO was described in oil formation

in developing seeds of *Brassica napus*, where refixation of CO_2 released during carbohydrate-to-fatty acid

conversion increases carbon use efficiency (Schwender et al. 2004). A similar explanation is unlikely to

hold for RuBisCO retention in *E. longa*, given the lack of fatty acid synthesis in its plastid and the apparently

much smaller significance of oil as a reserve substance in *E. longa* compared to *B. napus*.

We believe that the identification of the putatively functional Fd/Trx system, despite the absence of
the photosynthetic electron transport chain in this species, provides one of the key hints to understanding
the physiological role of the linear CB pathway in the *E. longa* plastid. The second potentially important
piece of the puzzle is provided by the proteomic data from *E. gracilis* (Novák Vanclová et al. 2019), which
indicated the presence of a unique form of GAPDH, the so-called non-phosphorylating GAPDH also
referred to as GapN, in the plastid (Table S8). This enzyme uses $NADP^+$ to directly oxidize GA3P to 3PG,
skipping the intermediate 2,3-BPG (and hence not leading to ATP generation) and producing NADPH rather
than NADH (Iddar et al. 2003). In plants, this enzyme is localized to the cytosol and is involved in shuttling
of reducing equivalents from the plastid by the exchange of GA3P and 3PG between the two compartments
(Rius et al. 2006). *E. longa* possesses a GapN homolog highly similar to the *E. gracilis* protein, including
an N-terminal presequence (Table S8), consistent with its presumed plastidial localization. It thus appears
that in *Euglena* spp. GapN mediates shuttling of reducing equivalents in the opposite direction than in plants,
i.e. from the cytosol to the plastid (Fig. 5). In case of *E. longa* this may be the main (if not the only)
mechanism of providing NADPH for the use in the plastid, whereas *E. gracilis* would utilize it when
photosynthetic production of NADPH is shut down. At the same time, the shuttle provides a mechanism of
linking the level of NADPH in the plastid with the cytosolic concentration of GA3P.

Taken together, we propose that in *E. longa* (at specific circumstances possibly also in *E. gracilis*)
the plastidial $NADPH/NADP^+$ ratio is directly influenced by the redox status of the cell, i.e. that it rises in
an excess of reducing power that slows down the glycolytic oxidation of GA3P in the cytosol. This
stimulates the linear CB pathway via the Fd/Trx system, effectively decreasing the level of GA3 by
converting it to 3PG without further increasing the reducing power in the cell. This conclusion is apparent
from considering the overall stoichiometries of the two alternative pathways from GA3 to 3PG:

glycolysis: 5 GA3P + 5 NAD^+ + 5 ADP + 5 PO_3^{2-} → 5 3PG + 5 NADH + 5 H^+ + 5 ATP

CB pathway: 5 GA3P + 3 CO_2 + 3 H_2O + 3 ATP → 6 3PG + 3 ADP + 2 PO_3^{2-}

The key difference is that the CB pathway does not produce NADH that needs to be reoxidized to keep the
glycolytic pathway running, since the fixed CO_2 effectively serves as an electron acceptor. Hence, turning
the CB bypass on may help the cell keep the redox balance when reoxidation of NADH is not efficient, e.g.
at hypoxic (or anoxic) conditions (although this happens at the expense of ATP). Indeed, euglenophytes in
their natural settings are probably often exposed to the shortage of oxygen, and anaerobiosis in *E. gracilis*
has been studied to some extent (Tucci et al. 2010; Zimorski et al. 2017). The anaerobic heterotrophic
metabolism of *E. gracilis* relies on fermentative degradation of the reserve polysaccharide paramylon
leading to production of wax esters (Yoshida et al. 2016). It is likely that *E. longa* exhibits a similar
metabolic adaptation to low levels of ambient oxygen as *E. gracilis*. However, details of the euglenophyte
anaerobic metabolism need to be worked out yet, and we propose that the plastid may be involved in it as a

kind of a redox valve. Work is ongoing to test this hypothesis and to illuminate further details of

physiological role of the linear CB pathway in the *E. longa* plastid.

Conclusions

Endosymbiotic organelles have proven to be extremely evolutionarily versatile. One of the manifestations

is the recurrent loss of key metabolic functions of the canonical forms of both mitochondria and plastids,

resulting in anaerobic mitochondria-related organelles (such as hydrogenosomes and mitosomes) and non-

photosynthetic plastids distributed across diverse eukaryotic branches. A lot of attention has been paid to

various mitochondrial derivatives and it is now well documented that they vary substantially in the

complement of functional pathways they have retained (Roger et al. 2017). The variation in functional

profiles of non-photosynthetic plastids is less well known, as the only example studied in detail is the

apicoplast of apicomplexan parasites. Nevertheless, a picture is emerging that independently evolved

colourless plastids may also exhibit a surprising degree of diversity in terms of their metabolic capacity.

Our analyses of the *E. longa* plastid stretch the breadth of variation among non-photosynthetic

plastids even further. The combination of pathways present (tocopherol and phylloquinone synthesis,

glycolipid synthesis and a linearized CB pathway including RuBisCO), absent (fatty acid, amino acid, and

isoprenoid precursor synthesis), and potentially residual (tetrapyrrole synthesis) makes the *E. longa* plastid

unlike any of the previously investigated non-photosynthetic plastids, including the apicoplast. However,

further work, combining additional *in silico* analyses (aimed, e.g., at potential plastid membrane transporters

mediating metabolite exchange with the cytosol) with biochemical and cytological investigations is needed

to achieve a more precise idea about the protein composition of the *E. longa* plastid and a better

understanding of its physiological roles.

Materials and Methods

Identification and annotation of plastid-targeted proteins

The analyses reported in this study were done using the *E. longa* transcriptome assembly reported previously

(Záhonová et al. 2018). Protein models for annotation were generated by a custom Geneious 8.1.6 (Kearse

et al. 2012) script that extracted all open reading frames longer than 297 bp, translated the sequences and

then filtered the protein models by a local BLAST+ ver.2.2.30 (Altschul et al. 1997) search against the

Swiss-Prot database (version 10/5/15, max E-value=10). Transcript models containing at least a partial

spliced leader sequence (TTTTTCG) at their 5'-end or 3'-end (within the first or last 35 nt) were translated

in the forward or reverse direction only, respectively.

Candidates for plastid-targeted proteins were identified using criteria for prediction of plastid-
targeting presequences described in detail by Záhonová et al. (2018). In the first step protein sequences were
gathered that fulfilled at least one of the following requirements: (i) the signal peptide was predicted by the
PrediSi v.2004 (Hiller et al. 2004) or PredSL v.2005 (Petsalaki et al. 2006) standalone programs; (ii) one or
two transmembrane domains at the N-terminus of the protein were predicted by standalone TMHMM 2.0c
(online version where graphical output was considered) (Krogh et al. 2001). The resulting set was then
filtered by checking for the presence of a plastid transit peptide, which was predicted by standalone
MultiLoc2.5 (Blum et al. 2009) after an *in silico* removal of the signal peptide or the first transmembrane
domain. Finally, protein models with a putative plastid-targeting presequence were automatically annotated
using InterProScan 5.21 (Jones et al. 2014) and the annotations were manually scrutinized to identify
proteins with an assignable specific metabolism-related function (enzymes of the metabolism of nucleic
acids and proteins were ignored in this study).

In parallel we searched with BLAST v.2.2.30 (including tBLASTn against the transcriptome
assembly and, in special cases, even against the whole set of raw RNAseq reads) for putative plastid proteins
by direct identification and evaluation of homologs of enzymes of specific biochemical pathways potentially
localized to the plastid; as queries we used respective protein sequences from *E. gracilis* (as identified by
Novák Vanclová et al. 2019), reference sequences from the KEGG PATHWAY Database
(https://www.genome.jp/kegg/pathway.html), or sequences identified by literature searches. In some cases,
a more sensitive homology detection algorithm HMMER 3.0 (Mistry et al. 2013) to identify homologs of
poorly conserved enzymes (e.g., UROS). For comparative purposes we used the same approach to identify
plastid-targeted proteins encoded by the transcriptome assemblies from *E. gracilis* reported by (Yoshida et
al. 2016) (accession GDJR00000000.1) and (Ebenezer et al. 2017) (accession GEFR00000000.1). Where
accessions are given for *E. gracilis* sequences, those starting with GDJR and GEFR belong to the former
and latter dataset, respectively.

For MenA cDNA resequencing, RNA was isolated using TRI Reagent (Thermo Fisher Scientific,
San Jose, USA) and mRNA was then extracted using the Dynabeads mRNA Purification kit (Thermo Fisher
Scientific). Reverse-transcription was performed with random hexamers and StrataScript III Reverse
Transcriptase (Thermo Fisher Scientific). For cDNA amplification, forward 5'-
GGTGCTGTTCTGCTCTCACT-3' and reverse 5'-CAGTGGGGATCAGAGATGCG-3' primers, and the
Q5 High-Fidelity DNA polymerase in a standard buffer (New England Biolabs) were used. Amplicons were
purified on MinElute PCR Purification columns (Qiagen, Hilden, Germany) and sequenced at the GATC
sequencing facility (Konstanz, Germany). The MenA cDNA sequence is deposited in GenBank with the
accession number MK484704.

Phylogenetic analyses

Phylogenetic analyses were employed to establish orthologous relationships among *E. longa* and *E. gracilis* genes or to illuminate the origin of the euglenophyte proteins of special interest. Homologs of target proteins were identified by BLAST v.2.2.30 searches in the non-redundant protein sequence database at NCBI (www.ncbi.nlm.nih.gov) and among protein models of selected organisms from JGI (Joint Genome Institute, jgi.doe.gov) and MMETSP (Marine Microbial Eukaryote Transcriptome Sequencing Project, marinemicroeukaryotes.org; Keeling et al. 2014). Sequences were aligned using the MAFFT v7.407 (Multiple Alignment using Fast Fourier Transform) tool with L-INS-I setting (Katoh and Standley 2013) and poorly aligned positions were eliminated by the trimAL v1.4.rev22 tool with "-automated1" trimming (Capella-Gutierrez et al. 2009). For presentation purposes, alignments were processed using the CHROMA software (Goodstadt and Ponting 2001). Maximum likelihood (ML) trees were inferred from the alignments using the LG+F+G4 model of IQ-TREE v1.6.9 (Nguyen et al. 2015), and employing the strategy of rapid bootstrapping followed by a "thorough" ML search with 1,000 bootstrap replicates (-m TEST -bb 1000). The list of species, and the number of sequences and amino acid positions are present in Tables S11-S22 for each phylogenetic tree.

Culture conditions

Euglena gracilis strain *Z* ("autotrophic" conditions) were cultivated statically under constant illumination at 26 °C in Cramer-Myers medium supplemented with ethanol (0.8% v/v) as a carbon source (Cramer and Myers 1952). *E. longa* strain CCAP 1204-17a (a gift from W. Hachtel, Bonn, Germany) and heterotrophic *E. gracilis* strain *Z* were cultivated in identical medium without illumination. The cultures of *E. longa* were not completely axenic, but the contaminating bacteria were kept at as low level as possible. *Rhabdomonas costata* strain PANT2 was provided by Vladimír Hampl (Department of Parasitology, Faculty of Science, Charles University in Prague, Czech Republic). It was isolated from a freshwater body in Pantanal (Brasil) and grown with an uncharacterised mixture of bacteria in Sonneborn's *Paramecium* medium (pH 7.4; Sonneborn 1950) at room temperature (inoculated every three to four weeks).

Mass spectrometry of structural lipids and terpenoids

For analysis of structural lipids, extracts from *E. longa* and autotrophic *E. gracilis* cellular pellets (four biological samples of different culture age) were obtained with chloroform and methanol solution (ratio – 2:1) following the method of (Folch et al. 1957) as modified by (Košťál and Šimek 1998). Samples were homogenized in extraction solution with glass beads using TissueLyser LT mill (Qiagen). Homogenates were dried, weighted, and resolved in 500 µl of chloroform and methanol (1:2) with internal standard PC 17:0/17:0 (Sigma Aldrich). Aliquots from each sample extract were used for lipid determination by HPLC

using a liquid chromatograph and autosampler Accela (Thermo Fisher Scientific). The samples (5 µL) were

injected and separated on the Gemini column 250 × 2 mm; i.d. 3 µm (Phenomenex, Torrance, USA). A

linear ion trap LTQ-XL mass spectrometer (Thermo Fisher Scientific) was used in both positive and

negative ion ESI mode. The settings of the system followed the methodology published earlier (Tomčala et

al. 2017). Data were acquired and processed using Xcalibur software version 2.1 (Thermo Fisher Scientific).

Particular compounds were determined based on m/z value, retention time, behaviour in positive and

negative ionization mode, and characteristic fragmentation pattern of target molecules (for details see

Tomčala et al. 2017).

Terpenoids were extracted from an autotrophic culture of *E. gracilis*, a heterotrophic culture of *E.*

gracilis, and a culture of *E. longa* of the same age in three repetitions. The same extraction procedure as for

lipid analysis was used. Sample aliquots were injected into the high-resolution mass spectrometry system

powered by Orbitrap Q-Exactive Plus with Dionex Ultimate 3000 XRS pump and Dionex Ultimate 3000

XRS Open autosampler (all by Thermo Fisher Scientific) and followed the settings described in (Tomčala

et al. 2017). Data were acquired and processed using Xcalibur software version 2.1. Identification of OH-

PhQ was achieved by considering the m/z value, fragmentation pattern, and high-resolution data.

Tocopherols (α, β/γ, and δ) were determined by the same characteristic as OH-PhQ and results were then

compared with commercially purchased standards (α-tocopherol from Sigma Aldrich, the other three

variants from SUPELCO).

Immunofluorescence assay

Immunofluorescence was performed as described previously (Botté et al. 2011). Briefly, cells were fixed in

4% paraformaldehyde for 30 minutes, cellular membranes were permeabilized for 10 minutes on ice with

0.1% non-ionic detergent Igepal CA-630 (Sigma-Aldrich) in PHEM buffer pH 6.9 (60 mM PIPES, 25 mM

HEPES, 10 mM EGTA, 2 mM $MgCl_2$), and background was masked with 3% BSA in PHEM buffer. DGDG

was detected using a polyclonal rabbit anti-DGDG antibody (1:25) that was a kind gift from Cyrille Y. Botté

(ApicoLipid Team, Laboratoire Adaptation et Pathogenie des Microorganismes, University of Grenoble I,

France), followed by incubation with a secondary Cy3-labeled polyclonal goat anti-rabbit antibody

(AP132C, 1:800, Merck Millipore, Billerica, USA). To improve the fluorescence lifetime, Fluoroshield™

with DAPI mounting medium (Sigma-Aldrich) was used. Cells were mounted on slides and observed with

a fluorescent microscope Olympus BX53 (Olympus, Tokyo, Japan). Photosynthetic *E. gracilis* served as a

positive control, and the primary osmotroph (i.e. aplastidic) *R. costata* as a negative control.

Acknowledgements

We thank Vladimír Hampl (Charles University) for providing the culture of *Rhabdomonas costata* and

Cyrille Y. Botté (University of Grenoble I) for providing the anti-DGDG antibody. We acknowledge

computation resources provided by CERIT-SC and MetaCentrum, Brno, Czech Republic, and the

infrastructure grant "Přístroje IET" (CZ.1.05/2.1.00/19.0388). We thank Laboratory of Analytical

Biochemistry and Metabolomics (Biology Centre, Czech Academy of Sciences) for free access to LC–MS

instruments. This study was supported by the Czech Science Foundation grants 17-21409S (to ME) and 16-

24027S (to MO), the Czech Academy of Sciences fellowship (to ZF), and the Scientific Grant Agency of

the Slovak Ministry of Education (grant VEGA 1/0535/17 to JK). This work was also part of the project

"Centre for research of pathogenicity and virulence of parasites", supported by the European Regional

Development Fund, within the Operational programme for Research, Development and Education

(CZ.02.1.01/0.0/0.0/16_019/0000759).

Competing interests

The authors declare that they have no financial or non-financial competing interests.

References

Altschul, SF, Madden, TL, Schaffer, AA, Zhang, J, Zhang, Z, Miller, W, Lipman, DJ. 1997. Gapped BLAST
and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res
25:3389-3402.

Anderson, R, Livermore, BP, Kates, M, Volcani, BE. 1978. The lipid composition of the non-photosynthetic
diatom *Nitzschia alba*. Biochim Biophys Acta 528:77-88.

Awai, K, Maréchal, E, Block, MA, Brun, D, Masuda, T, Shimada, H, Takamiya, K, Ohta, H, Joyard, J.
2001. Two types of MGDG synthase genes, found widely in both 16:3 and 18:3 plants, differentially
mediate galactolipid syntheses in photosynthetic and nonphotosynthetic tissues in *Arabidopsis*
thaliana. Proc Natl Acad Sci U S A 98:10960-10965.

Blum, T, Briesemeister, S, Kohlbacher, O. 2009. MultiLoc2: integrating phylogeny and Gene Ontology
terms improves subcellular protein localization prediction. BMC Bioinformatics 10:274.

Borza, T, Popescu, CE, Lee, RW. 2005. Multiple metabolic roles for the nonphotosynthetic plastid of the
green alga *Prototheca wickerhamii*. Eukaryot Cell 4:253-261.

Botté, C, Saidani, N, Mondragon, R, Mondragon, M, Isaac, G, Mui, E, McLeod, R, Dubremetz, JF, Vial, H,
Welti, R, et al. 2008. Subcellular localization and dynamics of a digalactolipid-like epitope in
Toxoplasma gondii. J Lipid Res 49:746-762.

Botté, CY, Yamaryo-Botté, Y, Janouškovec, J, Rupasinghe, T, Keeling, PJ, Crellin, P, Coppel, RL,
Maréchal, E, McConville, MJ, McFadden, GI. 2011. Identification of plant-like galactolipids in
Chromera velia, a photosynthetic relative of malaria parasites. J Biol Chem 286:29893-29903.

Botté, CY, Yamaryo-Botté, Y, Rupasinghe, TW, Mullin, KA, MacRae, JI, Spurck, TP, Kalanon, M, Shears,
MJ, Coppel, RL, Crellin, PK, et al. 2013. Atypical lipid composition in the purified relict plastid
(apicoplast) of malaria parasites. Proc Natl Acad Sci U S A 110:7506-7511.

Brettel, K. 1997. Electron transfer and arrangement of the redox cofactors in photosystem I. Biochimica et
Biophysica Acta (BBA) - Bioenergetics 1318:322-373.

Capella-Gutierrez, S, Silla-Martinez, JM, Gabaldon, T. 2009. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972-1973.

Cenci, U, Qiu, H, Pillonel, T, Cardol, P, Remacle, C, Colleoni, C, Kadouche, D, Chabi, M, Greub, G,
Bhattacharya, D, et al. 2018. Host-pathogen biotic interactions shaped vitamin K metabolism in
Archaeplastida. Sci Rep 8:15243.

Cihlář, J, Füssy, Z, Horák, A, Oborník, M. 2016. Evolution of the tetrapyrrole biosynthetic pathway in
secondary algae: conservation, redundancy and replacement. PLOS ONE 11:e0166338.

Cramer, M, Myers, J. 1952. Growth and photosynthetic characteristics of *Euglena gracilis*. Archiv
Mikrobiol 17:384-402.

Dagenais-Bellefeuille, S, Morse, D. 2013. Putting the N in dinoflagellates. Front Microbiol 4:369.

Dailey, HA, Dailey, TA, Gerdes, S, Jahn, D, Jahn, M, O'Brian, MR, Warren, MJ. 2017. Prokaryotic heme
biosynthesis: Multiple pathways to a common essential product. Microbiol Mol Biol Rev 81.

Disch, A, Schwender, J, Muller, C, Lichtenthaler, HK, Rohmer, M. 1998. Distribution of the mevalonate
and glyceraldehyde phosphate/pyruvate pathways for isoprenoid biosynthesis in unicellular algae
and the cyanobacterium *Synechocystis* PCC 6714. Biochem J 333 (Pt 2):381-388.

Donaher, N, Tanifuji, G, Onodera, NT, Malfatti, SA, Chain, PS, Hara, Y, Archibald, JM. 2009. The
complete plastid genome sequence of the secondarily nonphotosynthetic alga *Cryptomonas*
paramecium: reduction, compaction, and accelerated evolutionary rate. Genome Biol Evol 1:439-
448.

Dorrell, RG, Azuma, T, Nomura, M, Audren de Kerdrel, G, Paoli, L, Yang, S, Bowler, C, Ishii, KI,
Miyashita, H, Gile, GH, et al. 2019. Principles of plastid reductive evolution illuminated by
nonphotosynthetic chrysophytes. Proc Natl Acad Sci U S A 116:6914-6923.

Ebenezer, TE, Carrington, M, Lebert, M, Kelly, S, Field, MC. 2017. *Euglena gracilis* genome and
transcriptome: Organelles, nuclear genome assembly strategies and initial features. Adv Exp Med
Biol 979:125-140.

Fatihi, A, Latimer, S, Schmollinger, S, Block, A, Dussault, PH, Vermaas, WF, Merchant, SS, Basset, GJ.

2015. A dedicated type II NADPH dehydrogenase performs the penultimate step in the biosynthesis
of vitamin K_1 in *Synechocystis* and *Arabidopsis*. Plant Cell 27:1730-1741.

Fernandez, E, Galvan, A. 2008. Nitrate assimilation in *Chlamydomonas*. Eukaryot Cell 7:555-559.

Fernández Robledo, JA, Caler, E, Matsuzaki, M, Keeling, PJ, Shanmugam, D, Roos, DS, Vasta, GR. 2011.
The search for the missing link: a relic plastid in *Perkinsus*? Int J Parasitol 41:1217-1229.

Figueroa-Martinez, F, Nedelcu, AM, Smith, DR, Reyes-Prieto, A. 2015. When the lights go out: the
evolutionary fate of free-living colorless green algae. New Phytologist 206:972-982.

Folch, J, Lees, M, Sloane Stanley, GH. 1957. A simple method for the isolation and purification of total
lipides from animal tissues. J Biol Chem 226:497-509.

Furt, F, Oostende, Cv, Widhalm, JR, Dale, MA, Wertz, J, Basset, GJC. 2010. A bimodular oxidoreductase
mediates the specific reduction of phylloquinone (vitamin K1) in chloroplasts. Plant J 64:38-46.

Füssy, Z, Oborník, M. 2017. Chapter Six - Chromerids and Their Plastids. In: Y Hirakawa, editor. Advances
in Botanical Research: Academic Press. p. 187-218.

Füssy Z, Faitová T, Oborník M. 2019. Subcellular compartments interplay for carbon and nitrogen
allocation in *Chromera velia* and *Vitrella brassicaformis*. Genom Biol Evol, in press.
doi:10.1093/gbe/evz123.

Gerdes, S, Lerma-Ortiz, C, Frelin, O, Seaver, SM, Henry, CS, de Crecy-Lagard, V, Hanson, AD. 2012.
Plant B vitamin pathways and their compartmentation: a guide for the perplexed. J Exp Bot
63:5379-5395.

Giordano, M, Raven, JA. 2014. Nitrogen and sulfur assimilation in plants and algae. Aquatic Botany 118:45-
61.

Gockel, G, Hachtel, W. 2000. Complete gene map of the plastid genome of the nonphotosynthetic euglenoid
flagellate *Astasia longa*. Protist 151:347-351.

Gockel, G, Hachtel, W, Baier, S, Fliss, C, Henke, M. 1994. Genes for components of the chloroplast
translational apparatus are conserved in the reduced 73-kb plastid DNA of the nonphotosynthetic
euglenoid flagellate *Astasia longa*. Curr Genet 26:256-262.

Goddard-Borger, ED, Williams, SJ. 2017. Sulfoquinovose in the biosphere: occurrence, metabolism and
functions. Biochem J 474:827-849.

Gomez-Silva, B, Timko, MP, Schiff, JA. 1985. Chlorophyll biosynthesis from glutamate or 5-
aminolevulinate in intact *Euglena* chloroplasts. Planta 165:12-22.

Goodstadt, L, Ponting, CP. 2001. CHROMA: consensus-based colouring of multiple alignments for
publication. Bioinformatics 17:845-846.

Graupner, N, Jensen, M, Bock, C, Marks, S, Rahmann, S, Beisser, D, Boenigk, J. 2018. Evolution of
heterotrophy in chrysophytes as reflected by comparative transcriptomics. FEMS Microbiol Ecol

94.

Gu, X, Harding, S, Nyamdari, B, Aulakh, K, Clermont, K, Westwood, J, Tsai, C-J. 2018. A role for
phylloquinone biosynthesis in the plasma membrane as revealed in a non-photosynthetic parasitic
plant. bioRxiv:257519.

Gutbrod, K, Romer, J, Dormann, P. 2019. Phytol metabolism in plants. Prog Lipid Res 74:1-17.

Hadariová, L, Vesteg, M, Birčák, E, Schwartzbach, SD, Krajčovič, J. 2017. An intact plastid genome is
essential for the survival of colorless *Euglena longa* but not *Euglena gracilis*. Curr Genet 63:331-
341.

Hadariová, L, Vesteg, M, Hampl, V, Krajčovič, J. 2018. Reductive evolution of chloroplasts in non-
photosynthetic plants, algae and protists. Curr Genet 64:365-387.

Haferkamp, I, Deschamps, P, Ast, M, Jeblick, W, Maier, U, Ball, S, Neuhaus, HE. 2006. Molecular and
biochemical analysis of periplastidial starch metabolism in the cryptophyte *Guillardia theta*.
Eukaryot Cell 5:964-971.

Hachtel, W. 1996. DNA and gene expression in nonphotosynthetic plastids. In: M Pessarakli, editor.
Handbook of Photosynthesis. New York: Marcel Dekker. p. 349-355.

Hiller, K, Grote, A, Scheer, M, Munch, R, Jahn, D. 2004. PrediSi: prediction of signal peptides and their
cleavage positions. Nucleic Acids Res 32:W375-379.

Hori, K, Nobusawa, T, Watanabe, T, Madoka, Y, Suzuki, H, Shibata, D, Shimojima, M, Ohta, H. 2016.
Tangled evolutionary processes with commonality and diversity in plastidial glycolipid synthesis
in photosynthetic organisms. Biochim Biophys Acta 1861:1294-1308.

Iddar, A, Valverde, F, Serrano, A, Soukri, A. 2003. Purification of recombinant non-phosphorylating
NADP-dependent glyceraldehyde-3-phosphate dehydrogenase from *Streptococcus pyogenes*
expressed in *E. coli*. Mol Cell Biochem 247:195-203.

Jackson, C, Knoll, AH, Chan, CX, Verbruggen, H. 2018. Plastid phylogenomics with broad taxon sampling
further elucidates the distinct evolutionary origins and timing of secondary green plastids. Sci Rep
8:1523.

Janouškovec, J, Horák, A, Oborník, M, Lukeš, J, Keeling, PJ. 2010. A common red algal origin of the
apicomplexan, dinoflagellate, and heterokont plastids. Proc Natl Acad Sci U S A 107:10949-10954.

Janouškovec, J, Tikhonenkov, DV, Burki, F, Howe, AT, Kolísko, M, Mylnikov, AP, Keeling, PJ. 2015.
Factors mediating plastid dependency and the origins of parasitism in apicomplexans and their close
relatives. Proc Natl Acad Sci U S A 112:10200-10207.

Jones, P, Binns, D, Chang, HY, Fraser, M, Li, W, McAnulla, C, McWilliam, H, Maslen, J, Mitchell, A,
Nuka, G, et al. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics
30:1236-1240.

Kamikawa, R, Moog, D, Zauner, S, Tanifuji, G, Ishida, KI, Miyashita, H, Mayama, S, Hashimoto, T, Maier, UG, Archibald, JM, et al. 2017. A non-photosynthetic diatom reveals early steps of reductive evolution in plastids. Mol Biol Evol 34:2355-2366.

Karamoko, M, Cline, S, Redding, K, Ruiz, N, Hamel, PP. 2011. Lumen thiol oxidoreductase1, a disulfide bond-forming catalyst, is required for the assembly of photosystem II in *Arabidopsis*. The Plant Cell 23:4462.

Katoh, K, Standley, DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol 30:772-780.

Ke, H, Sigala, PA, Miura, K, Morrisey, JM, Mather, MW, Crowley, JR, Henderson, JP, Goldberg, DE, Long, CA, Vaidya, AB. 2014. The heme biosynthesis pathway is essential for *Plasmodium falciparum* development in mosquito stage but not in blood stages. J Biol Chem 289:34827-34837.

Kearse, M, Moir, R, Wilson, A, Stones-Havas, S, Cheung, M, Sturrock, S, Buxton, S, Cooper, A, Markowitz, S, Duran, C, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647-1649.

Keeling, PJ. 2013. The number, speed, and impact of plastid endosymbioses in eukaryotic evolution. Annu Rev Plant Biol 64:583-607.

Keeling, PJ, Burki, F, Wilcox, HM, Allam, B, Allen, EE, Amaral-Zettler, LA, Armbrust, EV, Archibald, JM, Bharti, AK, Bell, CJ, et al. 2014. The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans through transcriptome sequencing. PLOS Biol 12:e1001889.

Kim, D, Filtz, MR, Proteau, PJ. 2004. The methylerythritol phosphate pathway contributes to carotenoid but not phytol biosynthesis in *Euglena gracilis*. J Nat Prod 67:1067-1069.

Kim, GH, Jeong, HJ, Yoo, YD, Kim, S, Han, JH, Han, JW, Zuccarello, GC. 2013. Still acting green: continued expression of photosynthetic genes in the heterotrophic Dinoflagellate *Pfiesteria piscicida* (Peridiniales, Alveolata). PLOS ONE 8:e68232.

Kiss, JZ, Vasconcelos, AC, Triemer, RE. 1987. Structure of the euglenoid storage carbohydrate, paramylon. Am J Bot 74:877-882.

Kitaoka, S, Nakano, Y, Miyatake, K, Yokota, A. 1989. Enzymes and their functional location. In: DE Buetow, editor. Subcellular Biochemistry and Molecular Biology: Academic Press. p. 1-135.

Kivic, PA, Vesk, M. 1974. An electron microscope search for plastids in bleached *Euglena gracilis* and in *Astasia longa*. Can J Bot 52:695-699.

Kobayashi, K. 2016. Role of membrane glycerolipids in photosynthesis, thylakoid biogenesis and chloroplast development. J Plant Res 129:565-580.

Kořený, L, Oborník, M. 2011. Sequence evidence for the presence of two tetrapyrrole pathways in *Euglena*

gracilis. Genome Biol Evol 3:359-364.

Košťál, V, Šimek, P. 1998. Changes in fatty acid composition of phospholipids and triacylglycerols after
cold-acclimation of an aestivating insect prepupa. Journal of Comparative Physiology B 168:453-
460.

Krajčovič, J, Ebringer, L, Schwartzbach, SD. 2002. Reversion of endosymbiosis? In: J Seckbach, editor.
Symbiosis: Mechanisms and model systems. Dordrecht: Springer Netherlands. p. 185-206.

Krogh, A, Larsson, B, von Heijne, G, Sonnhammer, EL. 2001. Predicting transmembrane protein topology
with a hidden Markov model: application to complete genomes. J Mol Biol 305:567-580.

Kwong, WK, Del Campo, J, Mathur, V, Vermeij, MJA, Keeling, PJ. 2019. A widespread coral-infecting
apicomplexan with chlorophyll biosynthesis genes. Nature 568:103-107.

Lakey, B, Triemer, R. 2016. The tetrapyrrole synthesis pathway as a model of horizontal gene transfer in
euglenoids. J Phycol 53:198-217.

Lambalot, RH, Walsh, CT. 1995. Cloning, overproduction, and characterization of the *Escherichia coli*
holo-acyl carrier protein synthase. J Biol Chem 270:24658-24661.

Leander, BS, Lax, G, Karnkowska, A, Simpson, AGB. 2017. Euglenida. In: JM Archibald, AGB Simpson,
CH Slamovits, L Margulis, M Melkonian, DJ Chapman, JO Corliss, editors. Handbook of the
Protists. Cham: Springer International Publishing. p. 1-42.

Li, C, Wang, Y, Liu, L, Hu, Y, Zhang, F, Mergen, S, Wang, G, Schlappi, MR, Chu, C. 2011. A rice plastidial
nucleotide sugar epimerase is involved in galactolipid biosynthesis and improves photosynthetic
efficiency. PLoS Genet 7:e1002196.

Lim, L, McFadden, GI. 2010. The evolution, metabolism and functions of the apicoplast. Philos Trans R
Soc Lond B Biol Sci 365:749-763.

Lohr, M, Schwender, J, Polle, JE. 2012. Isoprenoid biosynthesis in eukaryotic phototrophs: a spotlight on
algae. Plant Sci 185-186:9-22.

Lochner, K, Doring, O, Bottger, M. 2003. Phylloquinone, what can we learn from plants? Biofactors 18:73-
78.

Maeda, H, DellaPenna, D. 2007. Tocopherol functions in photosynthetic organisms. Curr Opin Plant Biol
10:260-265.

Marin, B, Palm, A, Klingberg, M, Melkonian, M. 2003. Phylogeny and taxonomic revision of plastid-
containing euglenophytes based on SSU rDNA sequence comparisons and synapomorphic
signatures in the SSU rRNA secondary structure. Protist 154:99-145.

Markunas, CM, Triemer, RE. 2016. Evolutionary history of the enzymes involved in the Calvin-Benson
cycle in euglenids. J Eukaryot Microbiol 63:326-339.

Martins-Duarte, ES, Carias, M, Vommaro, R, Surolia, N, de Souza, W. 2016. Apicoplast fatty acid synthesis

is essential for pellicle formation at the end of cytokinesis in *Toxoplasma gondii*. J Cell Sci

129:3320-3331.

Matsuzaki, M, Kuroiwa, H, Kuroiwa, T, Kita, K, Nozaki, H. 2008. A cryptic algal group unveiled: a plastid

biosynthesis pathway in the oyster parasite *Perkinsus marinus*. Mol Biol Evol 25:1167-1179.

McFadden, GI. 2014. Origin and evolution of plastids and photosynthesis in eukaryotes. Cold Spring Harb

Perspect Biol 6:a016105.

McFadden, GI, Yeh, E. 2017. The apicoplast: now you see it, now you don't. Int J Parasitol 47:137-144.

Miller, LH, Ackerman, HC, Su, XZ, Wellems, TE. 2013. Malaria biology and disease pathogenesis: insights

for new treatments. Nat Med 19:156-167.

Mistry, J, Finn, RD, Eddy, SR, Bateman, A, Punta, M. 2013. Challenges in homology search: HMMER3

and convergent evolution of coiled-coil regions. Nucleic Acids Res 41:e121.

Moffatt, BA, Ashihara, H. 2002. Purine and pyrimidine nucleotide synthesis and metabolism. Arabidopsis

Book 1:e0018.

Moore, RB, Oborník, M, Janouškovec, J, Chrudimský, T, Vancová, M, Green, DH, Wright, SW, Davies,

NW, Bolch, CJ, Heimann, K, et al. 2008. A photosynthetic alveolate closely related to apicomplexan

parasites. Nature 451:959-963.

Nagaraj, VA, Sundaram, B, Varadarajan, NM, Subramani, PA, Kalappa, DM, Ghosh, SK, Padmanaban, G.

2013. Malaria parasite-synthesized heme is essential in the mosquito and liver stages and

complements host heme in the blood stages of infection. PLOS Pathog 9:e1003522.

Neuhaus, HE, Emes, MJ. 2000. Nonphotosynthetic metabolism in plastids. Annu Rev Plant Physiol Plant

Mol Biol 51:111-140.

Nguyen, LT, Schmidt, HA, von Haeseler, A, Minh, BQ. 2015. IQ-TREE: a fast and effective stochastic

algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268-274.

Novák Vanclová, AMG, Zoltner, M, Kelly, S, Soukal, P, Záhonová, K, Füssy, Z, Ebenezer, TE, Lacová

Dobáková, E, Eliáš, M, Lukeš, J, et al. 2019. Proteome of the secondary plastid of *Euglena gracilis*

reveals several metabolic quirks and traces of a colourful history. bioRxiv:573709.

Nudelman, MA, Rossi, MS, Conforti, V, Triemer, RE. 2003. Phylogeny of Euglenophyceae based on small

subunit rDNA sequences: Taxonomic implications. J Phycol 39:226-235.

Oborník, M, Green, BR. 2005. Mosaic origin of the heme biosynthesis pathway in photosynthetic

eukaryotes. Mol Biol Evol 22:2343-2353.

Oborník, M, Janouškovec, J, Chrudimský, T, Lukeš, J. 2009. Evolution of the apicoplast and its hosts: from

heterotrophy to autotrophy and back again. Int J Parasitol 39:1-12.

Oda, Y, Miyatake, K, Kitaoka, S. 1979. Inability of *Euglena gracilis* Z to utilize nitrate, nitrite and urea as

the nitrogen sources. Bulletin of the University of Osaka Prefecture. Series B. Agriculture and

Biology 31:43-48.

Patron, NJ, Durnford, DG, Kopriva, S. 2008. Sulfate assimilation in eukaryotes: fusions, relocations and
lateral transfers. BMC Evol Biol 8:39.

Petsalaki, EI, Bagos, PG, Litou, ZI, Hamodrakas, SJ. 2006. PredSL: a tool for the N-terminal sequence-
based prediction of protein subcellular localization. Genomics Proteomics Bioinformatics 4:48-55.

Pick, TR, Brautigam, A, Schulz, MA, Obata, T, Fernie, AR, Weber, AP. 2013. PLGG1, a plastidic glycolate
glycerate transporter, is required for photorespiration and defines a unique class of metabolite
transporters. Proc Natl Acad Sci U S A 110:3185-3190.

Pinnaduwage, P, Bruce, BD. 1996. In vitro interaction between a chloroplast transit peptide and chloroplast
outer envelope lipids is sequence-specific and lipid class-dependent. J Biol Chem 271:32907-
32915.

Pombert, JF, Blouin, NA, Lane, C, Boucias, D, Keeling, PJ. 2014. A lack of parasitic reduction in the
obligate parasitic green alga *Helicosporidium*. PLoS Genet 10:e1004355.

Ponce-Toledo, RI, Deschamps, P, Lopez-Garcia, P, Zivanovic, Y, Benzerara, K, Moreira, D. 2017. An
Early-Branching Freshwater Cyanobacterium at the Origin of Plastids. Curr Biol 27:386-391.

Raines, CA, Lloyd, JC. 2001. C3 carbon reduction cycle. eLS: John Wiley & Sons, Ltd.

Rius, SP, Casati, P, Iglesias, AA, Gomez-Casati, DF. 2006. Characterization of an *Arabidopsis thaliana*
mutant lacking a cytosolic non-phosphorylating glyceraldehyde-3-phosphate dehydrogenase. Plant
Mol Biol 61:945-957.

Roger, AJ, Muñoz-Gómez, SA, Kamikawa, R. 2017. The origin and diversification of mitochondria. Curr
Biol 27:R1177-R1192.

Sanchez-Puerta, MV, Lippmeier, JC, Apt, KE, Delwiche, CF. 2007. Plastid genes in a non-photosynthetic
dinoflagellate. Protist 158:105-117.

Sanz-Luque, E, Chamizo-Ampudia, A, Llamas, A, Galvan, A, Fernandez, E. 2015. Understanding nitrate
assimilation and its regulation in microalgae. Front Plant Sci 6:899.

Seeger, JW, Bentley, R. 1991. Phylloquinone (vitamin K_1) biosynthesis in *Euglena gracilis* strain *Z*.
Phytochemistry 30:3585-3589.

Sekiguchi, H, Moriya, M, Nakayama, T, Inouye, I. 2002. Vestigial chloroplasts in heterotrophic
stramenopiles *Pteridomonas danica* and *Ciliophrys infusionum* (Dictyochophyceae). Protist
153:157-167.

Shashidhara, LS, Smith, AG. 1991. Expression and subcellular location of the tetrapyrrole synthesis enzyme
porphobilinogen deaminase in light-grown *Euglena gracilis* and three nonchlorophyllous cell lines.
Proc Natl Acad Sci U S A 88:63-67.

Schopfer, P, Heyno, E, Drepper, F, Krieger-Liszkay, A. 2008. Naphthoquinone-dependent generation of

superoxide radicals by quinone reductase isolated from the plasma membrane of soybean. Plant
1070   Physiol 147:864.

1071   Schürmann, P, Buchanan, BB. 2008. The ferredoxin/thioredoxin system of oxygenic photosynthesis.
1072   Antioxid Redox Signal 10:1235-1274.

1073   Schwender, J, Goffman, F, Ohlrogge, JB, Shachar-Hill, Y. 2004. Rubisco without the Calvin cycle improves
1074   the carbon efficiency of developing green seeds. Nature 432:779-782.

1075   Siemeister, G, Buchholz, C, Hachtel, W. 1990a. Genes for ribosomal proteins are retained on the 73 kb
1076   DNA from *Astasia longa* that resembles *Euglena* chloroplast DNA. Curr Genet 18:457-464.

1077   Siemeister, G, Buchholz, C, Hachtel, W. 1990b. Genes for the plastid elongation factor Tu and ribosomal
1078   protein S7 and six tRNA genes on the 73 kb DNA from *Astasia longa* that resembles the chloroplast
1079   DNA of *Euglena*. Mol Gen Genet 220:425-432.

1080   Slamovits, CH, Keeling, PJ. 2008. Plastid-derived genes in the nonphotosynthetic alveolate *Oxyrrhis*
1081   *marina*. Mol Biol Evol 25:1297-1306.

1082   Smith, DR, Lee, RW. 2014. A plastid without a genome: evidence from the nonphotosynthetic green algal
1083   genus *Polytomella*. Plant Physiol 164:1812-1819.

1084   Sonneborn, TM. 1950. Methods in the general biology and genetics of *Paramecium aurelia*. Journal of
1085   Experimental Zoology 113:87-147.

1086   Sun, GL, Shen, W, Wen, JF. 2008. Triosephosphate isomerase genes in two trophic modes of euglenoids
1087   (euglenophyceae) and their phylogenetic analysis. J Eukaryot Microbiol 55:170-177.

1088   Ševčíková, T, Horák, A, Klimeš, V, Zbránková, V, Demir-Hilton, E, Sudek, S, Jenkins, J, Schmutz, J, Přibyl,
1089   P, Fousek, J, et al. 2015. Updating algal evolutionary relationships through plastid genome
1090   sequencing: did alveolate plastids emerge through endosymbiosis of an ochrophyte? Sci Rep
1091   5:10134.

1092   Tabita, FR, Hanson, TE, Li, H, Satagopan, S, Singh, J, Chan, S. 2007. Function, structure, and evolution of
1093   the RubisCO-like proteins and their RubisCO homologs. Microbiol Mol Biol Rev 71:576-599.

1094   Tomčala, A, Kyselová, V, Schneedorferová, I, Opekarová, I, Moos, M, Urajová, P, Kručinská, J, Oborník,
1095   M. 2017. Separation and identification of lipids in the photosynthetic cousins of Apicomplexa
1096   *Chromera velia* and *Vitrella brassicaformis*. J Sep Sci 40:3402-3413.

1097   Tripathy, BC, Sherameti, I, Oelmuller, R. 2010. Siroheme: an essential component for life on earth. Plant
1098   Signal Behav 5:14-20.

1099   Tucci, S, Vacula, R, Krajčovič, J, Proksch, P, Martin, W. 2010. Variability of wax ester fermentation in
1100   natural and bleached *Euglena gracilis* strains in response to oxygen and the elongase inhibitor
1101   flufenacet. J Eukaryot Microbiol 57:63-69.

Turmel, M, Gagnon, MC, O'Kelly, CJ, Otis, C, Lemieux, C. 2009. The chloroplast genomes of the green

algae *Pyramimonas*, *Monomastix*, and *Pycnococcus* shed new light on the evolutionary history of
prasinophytes and the origin of the secondary chloroplasts of euglenids. Mol Biol Evol 26:631-648.

Van Dingenen, J, Blomme, J, Gonzalez, N, Inzé, D. 2016. Plants grow with a little help from their organelle
friends. J Exp Bot 67:6267-6281.

Vogel, A, Schwacke, R, Denton, AK, Usadel, B, Hollmann, J, Fischer, K, Bolger, A, Schmidt, MH, Bolger,
ME, Gundlach, H, et al. 2018. Footprints of parasitism in the genome of the parasitic flowering
plant *Cuscuta campestris*. Nat Commun 9:2515.

Vollmer, M, Thomsen, N, Wiek, S, Seeber, F. 2001. Apicomplexan parasites possess distinct nuclear-
encoded, but apicoplast-localized, plant-type ferredoxin-NADP+ reductase and ferredoxin. J Biol
Chem 276:5483-5490.

Walker, BJ, South, PF, Ort, DR. 2016. Physiological evidence for plasticity in glycolate/glycerate transport
during photorespiration. Photosynth Res 129:93-103.

Waller, RF, Gornik, SG, Kořený, L, Pain, A. 2016. Metabolic pathway redundancy within the
apicomplexan-dinoflagellate radiation argues against an ancient chromalveolate plastid. Commun
Integr Biol 9:e1116653.

Watanabe, F, Yoshimura, K, Shigeoka, S. 2017. Biochemistry and physiology of vitamins in *Euglena*. In:
SD Schwartzbach, S Shigeoka, editors. *Euglena*: Biochemistry, Cell and Molecular Biology. Cham:
Springer International Publishing. p. 65-90.

Webster, DA, Hackett, DP, Park, RB. 1967. The respiratory chain of colorless algae: III. Electron
microscopy. J Ultrastruct Res 21:514-523.

Wedel, N, Soll, J, Paap, BK. 1997. CP12 provides a new mode of light regulation of Calvin cycle activity
in higher plants. Proc Natl Acad Sci U S A 94:10479-10484.

Weinstein, JD, Beale, SI. 1983. Separate physiological roles and subcellular compartments for two
tetrapyrrole biosynthetic pathways in *Euglena gracilis*. J Biol Chem 258:6799-6807.

Wicke, S, Müller, KF, de Pamphilis, CW, Quandt, D, Wickett, NJ, Zhang, Y, Renner, SS, Schneeweiss,
GM. 2013. Mechanisms of functional and physical genome reduction in photosynthetic and
nonphotosynthetic parasitic plants of the broomrape family. The Plant Cell 25:3711.

Williamson, DH, Gardner, MJ, Preiser, P, Moore, DJ, Rangachari, K, Wilson, RJM. 1994. The evolutionary
origin of the 35 kb circular DNA of *Plasmodium falciparum*: new evidence supports a possible
rhodophyte ancestry. Molecular and General Genetics MGG 243:249-252.

Wilson, RJ, Denny, PW, Preiser, PR, Rangachari, K, Roberts, K, Roy, A, Whyte, A, Strath, M, Moore, DJ,
Moore, PW, et al. 1996. Complete gene map of the plastid-like DNA of the malaria parasite
Plasmodium falciparum. J Mol Biol 261:155-172.

Yeh, E, DeRisi, JL. 2011. Chemical rescue of malaria parasites lacking an apicoplast defines organelle

function in blood-stage *Plasmodium falciparum*. PLOS Biol 9:e1001138.

Yoon, EY, Yang, AR, Park, J, Moon, SJ, Jeong, EJ, Rho, JR. 2017. Characterization of a new trioxilin and

a sulfoquinovosyl diacylglycerol with anti-inflammatory properties from the dinoflagellate

Oxyrrhis marina. Mar Drugs 15.

Yoshida, Y, Tomiyama, T, Maruta, T, Tomita, M, Ishikawa, T, Arakawa, K. 2016. De novo assembly and

comparative transcriptome analysis of *Euglena gracilis* in response to anaerobic conditions. BMC

Genomics 17:182.

Yuzawa, Y, Nishihara, H, Haraguchi, T, Masuda, S, Shimojima, M, Shimoyama, A, Yuasa, H, Okada, N,

Ohta, H. 2012. Phylogeny of galactolipid synthase homologs together with their enzymatic analyses

revealed a possible origin and divergence time for photosynthetic membrane biogenesis. DNA Res

19:91-102.

Záhonová, K, Füssy, Z, Birčák, E, Novák Vanclová, AMG, Klimeš, V, Vesteg, M, Krajčovič, J, Oborník,

M, Eliáš, M. 2018. Peculiar features of the plastids of the colourless alga *Euglena longa* and

photosynthetic euglenophytes unveiled by transcriptome analyses. Sci Rep 8:17012.

Záhonová, K, Füssy, Z, Oborník, M, Eliáš, M, Yurchenko, V. 2016. RuBisCO in non-photosynthetic alga

Euglena longa: divergent features, transcriptomic analysis and regulation of complex formation.

PLOS ONE 11:e0158790.

Ziegler, K, Maldener, I, Lockau, W. 1989. 5'-monohydroxyphylloquinone as a component of photosystem

I. Zeitschrift für Naturforschung C 44:468-472.

Zimorski, V, Rauch, C, van Hellemond, JJ, Tielens, AGM, Martin, WF. 2017. The mitochondrion of

Euglena gracilis. In: SD Schwartzbach, S Shigeoka, editors. *Euglena*: Biochemistry, cell and

molecular biology. Cham: Springer International Publishing. p. 19-37.

1159