Spatial distribution of lamin A determines nuclear stiffness and stress-mediated deformation

Luv Kishore Srivastava[1], Zhaoping Ju[2], Ajinkya Ghagre[1], Allen J. Ehrlicher[1,2]*

1. **Department of Bioengineering, McGill University, Montreal H3A 0E9**
2. **Department of Anatomy and Cell Biology, McGill University, Montreal H3A 0C7**

Abstract: The nucleus is the largest organelle and information center of the cell; while diverse cellular components have been identified as mechanotransduction elements, the deformation of the nucleus itself is emerging as a critical mechanosensory mechanism, suggesting that the nuclear stiffness is essential in determining responses to intracellular and extracellular stresses. The nuclear membrane protein, lamin A, is known to be a dominant component in nuclear stiffening; however, the quantitative relationship between lamin A expression and nuclear deformation is still unclear. Here we measure the nuclear moduli as a function of lamin A expression and applied stress, revealing a linear dependence of bulk modulus on lamin A expression. We also find that the bulk modulus is anisotropic, with the vertical axis of the nucleus being more compliant than the minor and major axis. To examine how lamin A influences nuclear mechanics at the sub-micron scale we correlated the spatial distribution of lamin A with 3D nuclear envelope deformation, revealing that local areas of the nuclear envelope with higher expression levels of lamin A have correspondingly lower local deformations, and that increased lamin A expression levels result in a narrower distribution of smaller deformations. These findings describe the complex dispersion of nuclear deformations as a function of lamin A expression and distribution and implicate a role in mechanotransduction.

Introduction: The nucleus is one of the most crucial organelles and the storehouse of DNA in the cell, integrating diverse biochemical cues for proper cell function[1]. In addition to biochemical cues, the nucleus has an increasingly clear role as a mechanosensory structure in the cell[2]. Nuclear

mechanotransduction is a process by which the nucleus detects mechanical cues and converts them

to biochemical signals, which alters cell physiology. As a mechanosensor, the nucleus' stiffness

determines specific deformations in response to applied stresses. The nucleus is notably stiffer

than the cytoplasm (0.5-3kPa)[3], with an effective Young's modulus ranging from 1–10kPa[4,5];

while these values appear relatively consistent in a given cell line[6] , multiple factors may cause

nuclear stiffness to change significantly. For example, the stiffness of nuclei varies 5-fold during

cell division[7], in stem cells the stiffness can increase 6-fold over the differentiation process[8], and

in many types of cancers the structure of the nucleus is altered and the stiffness is vastly reduced[9].

Many of these morphological and mechanical changes are the result of the changes in nuclear

envelope architecture[10].

The nuclear envelope is composed of double-membraned bilipid layers with a 30–50nm

perinuclear space between the membranes. The outer nuclear membrane is connected to the

endoplasmic reticulum, whereas the inner membrane is held in place by a consortium of more than

50 membrane proteins. Some of the key proteins present in the inner nuclear membrane include

emerin, SUN1, SUN2, lamins, and lamin associated proteins[11]. Lamins are type V intermediate

filaments localized under the nuclear membrane. The inner membrane is held in place by the

underlying nuclear lamina, which is mainly composed of A type and B type lamins. Lamin A has

a profound effect on the stiffness of the nucleus, but lamin B shows very little or no effect on the

nuclear stiffness, thus making lamin A a dominant component in nuclear mechanics[12].

Numerous studies have illustrated the significance of lamin A in determining the stiffness of the

nucleus. Suppression or knockdown of lamin A increases nuclear compliance, and nuclei in lamin

A knockout cells display 30% to 50% more deformation than WT when probed with substrate

stretch or magnetic microrheology, respectively[13]. Previous studies have also shown that nuclei in

lamin A knockout cells are more fragile, with a greater susceptibility to rupture under increased intranuclear pressure[14]. Nuclei isolated from cells transfected with shRNA for lamin A showed 1.5–1.7 times more bead displacement compared to wild type nuclei in magnetic bead microrheology, indicating the role of lamin A in determining the stiffness of the nucleus[15]. Conversely, over-expression of lamin A stiffens nuclei: when 3T3 fibroblasts were transfected with lamin A and seeded on vertical nanopillars, showed 40–50% less nanopillar induced deformation in both intact and isolated nuclei[14]. Lamin A expression and nuclear stiffness also directly affect the behavior of the cell; as an example, lamin A overexpression hinders 3D cell migration through micropores[16]; however, it seems to facilitate 2D migration as wild type cells are marginally faster than lamin A knockout cells[17]. While lamin A expression clearly influences nuclear mechanics and cell behavior, the vast majority of these studies have looked at qualitative treatments of lamin A, i.e., parsing cells into groups of overexpression, wild type, or knock-down. However, the quantitative scaling of nuclear stiffness and corresponding nuclear deformation as a function of lamin A expression remains unknown.

The expression of lamin A is highly variable, and changes in different stages of cell cycle[18], in stem cell differentiation[19], and as a function of microenvironment stiffness[20]. In addition to overall expression levels, the distribution of lamin A appears heterogeneous with the formation of foci and honeycomb patterns observed in the nuclear envelope[21]. It remains unknown how the spatial heterogeneity of lamin A affects nuclear deformation. Studies to date have focused on comparing gross nuclear stiffness changes in ensemble population lamin A expression levels via over-expression or suppression[22], but have not examined the quantitative dependence of nuclear stiffness on lamin A expression, or the impact of the spatial distribution of lamin A of deformation.

To understand the quantitative impact of lamin A structure on nuclear mechanics, here we measure

the deformability and bulk moduli of nuclei as a function of lamin A expression. We identify a

mechanical anisotropy of bulk compression, leading to increasing relative compliance in the major,

minor, and z axes, respectively. At the sub-micron level, we quantify the spatial distribution of

lamin A and show that dense lamin A structures attenuate local deformation, providing

heterogeneous nuclear deformations even under uniform nuclear stresses.

Methodology

Quantification of lamin A expression using antibodies and lamin A chromobody: To quantify

the total lamin A expression in the nucleus, the NIH 3T3 fibroblast cells were stably transfected

with GFP tagged lamin A chromobody[23] (Chromtek). To testify if the chromobody is an accurate

quantitative metric of lamin A expression and distribution, we compared its fluorescence with

antibody staining (Atto-647N Sigma) as a gold standard (see supplementary information Fig. S1).

For this, lamin A chromobody transfected cells were seeded onto gridded coverslips, fixed,

permeabilized and stained with Atto-647 lamin A antibodies and then imaged with a 63X/1.4 NA

oil immersion objective on a confocal microscope (Leica SP8). Fluorescence colocalization was

determined using Pearson's correlation coefficient of 0.88 ± 0.04 for 10 cells (see supplementary

information Fig. S2).

Cell culture & modulating nuclear lamin A expression: NIH 3T3 fibroblast cells were cultured

following standard protocols in DMEM media supplemented with 10% FBS and 1% Penicillin-

Streptomycin antibiotic. All cells were stably transfected with GFP tagged lamin A chromobody,

which labels the total lamin A present on the nuclear membrane without affecting its expression.

To over-express lamin A in intact cells, we transfected cells with m-Cherry tagged plasmid DNA

for lamin A which was a gift from Michael Davison (Addgene plasmid # 55068), whereas to

suppress lamin A expression we transfected the cells with RFP tagged inducible shRNA construct

for lamin A (Dharmacon).

Quantification of nuclear bulk moduli: The cells were synchronized in 0.1% FBS containing

media for 18 hours to eliminate the contribution of cell cycle on lamin A expression and

distribution. They were then attached on coverslips and exposed to media with 1%, 2.5%, 5%,

7.5% and 10% (wt/wt) 400 Da polyethylene glycol (PEG 400, Sigma) thus, exerting 847.7kPa,

936.5kPa, 1099.2kPa, 1290.3kPa, and 1514kPa osmotic stresses respectively for 25 minutes to

reach an equilibrium of compression[24] (see supplementary information Fig. S3), followed by

acquiring confocal microscopy XYZ-stacks (63x 1.4NA, Leica SP8). The cells compress due to

the osmotic pressure exerted by the hypertonic solution causing an efflux of water[25], and this

change in volume is measured and used to determine the bulk modulus by the relation $B = -\Delta P/$

$(\Delta V/V)$ where, B= bulk modulus, ΔP = osmotic pressure, ΔV= change in volume and V = original

volume[6].

3-D volume measurement of nuclei: XYZ stacks of GFP tagged lamin A chromobody cells were

imaged using a 63X/1.4 NA oil immersion objective on a Leica SP8 confocal microscope with a

z-step size of 0.3µm. 3-D visualization and measurement were carried out using ImageJ software.

Before 3-D measurement, the nuclear images were deconvolved using the 'Iterative deconvolve

3D' plugin in ImageJ. We measured nuclear volumes by counting the number of voxels of the

thresholded nuclei and then multiplied it by the size of each voxel to get the volume of the

thresholded region. To measure the major, minor and z-axis deformations, we acquired XYZ

stacks for different time intervals post-PEG addition. The stacks were summed in the Z dimension

using 'z-project' plugin in image J, followed by fitting the z-projections into an ellipse and

calculating the major and minor axis for different time intervals.

Nuclear strain mapping: To measure the local strain on the nuclear membrane, we captured

images of the GFP tagged lamin A chromobody nucleus pre and post-exposure to 2.5% PEG 400

(936.5kPa). To quantify the local deformation, a custom-made MATLAB code[26] was slightly

modified and used. Briefly, the code utilizes a fast Fourier transform (FFT) based cross-correlation

formulation in conjunction with the iterative deformation method (IDM). FIDVC calculates the

displacement between the images of GFP tagged nucleus by tracking the pixel intensities of

chromobody fluorescence. We mapped these local displacements with the lamin A expressions in

3-D space to determine the relationship between lamin A spatial distribution and nuclear

deformation.

Results:

Lamin A stiffens the nucleus: To measure the bulk compressibility of the nuclei, we exposed cell

nuclei of varying lamin A expression with different osmotic pressures using 400 Da PEG. Since

PEG is impermeable across the cell membrane, it creates a hyperosmotic environment for the cell

and nucleus resulting in their compression (Fig. 1a). This compression is due to the expulsion of

water as the cell and nuclei regain their original volume when returned to isotonic media[6,25].

Previous studies have shown that under external osmotic pressure nuclear volume changes along

with the concentration of intracellular macromolecules, thus exerting compressive forces on it[6,25].

The pressure-volume curve for nuclei of wild type NIH 3T3 fibroblast cells at different PEG 400

concentrations exerting different osmotic pressures shows that the nuclear compression was higher

in low lamin A expressing cells and vice-versa (Fig. 1b)

This is also reflected in the bulk moduli calculated from the change in the nuclear volume where

the cells expressing more lamin A are relatively stiffer. The bulk modulus for the highest lamin A

expressing cell at the highest exerted osmotic pressure is 3.48Mpa (Fig. 1c). A linear relationship

could be seen when we plotted bulk moduli as a function of lamin A fluorescence clearly a direct

relationship between lamin A expression and nuclear rigidity (Fig. 1d). We also observed the

nuclear volume as a function of time for different lamin A expression cells after PEG addition and

show that below lamin A fluorescence of 16.9 the rate of nuclear compression was more uniform,

whereas in nuclei with lamin A expression more than 16.9 showed a sudden fall in nuclear volume

followed by a very slow compression indicating different mechanical behavior of the nuclei at

below a certain level of lamin A expression (Fig. 1e & 1f).

Figure 1 | Hyperosmotic compression of the nucleus in 3T3 Fibroblast cells. a, 3D reconstruction of an example

nucleus within a cell at osmotic pressures of 743.2 and 1514.1kPa (scalebar=3µm). **b,** Normalized nuclear volumes

of 3T3 cells transfected with lamin A chromobody for wild type cells showing that stiffness increases as the protein

expression increases (fluorescence values in arbitrary units). **c,** Bulk moduli of 3T3 cells transfected with lamin A

chromobody for wild type showing that stiffness increases as the protein expression increases (n=8). **d,** Nuclear

stiffnes in terms of bulk moduli increasing as a function of lamin A fluorescence. **e,** Example 3T3 fibroblast nuclei

showing change in nuclear volume with time after PEG addition (scalebar = 2µm). **f,** Change in nuclear volume as a

function of time under 1514.1kPa osmotic pressure as a function of lamin A fluorescence.

Nuclear stiffness is anisotropic. To measure the anisotropic nuclear deformation, we exposed the

cells transfected with lamin A chromobody to PEG and imaged with confocal microscopy (Leica

SP8) after 25 minutes of PEG addition. Our results reveal strong anisotropic nuclear deformation

where the deformation in the z-axis was highest, followed by minor and major axis (Fig 2a). To

examine the dynamics of nuclear deformation under stress, we imaged the nuclei every 60 seconds

after PEG addition. Here, the nucleus quickly flattens in the z-axis, resulting in a transient

expansion along the major and minor axes, which starts to contract after 5 minutes (Fig. 2b). This

shows a time dependency to nuclear compression after PEG addition. Also, the deformation in z-

axis was inversely related with lamin A expression but the deformation along major and minor

axis did not show any relationship with lamin A expression (Fig. 2c).

Figure 2 | Anisotropy in nuclear compression under PEG. a, Example of a wild type nuclei in Y-Z and X-Z plane

before PEG addition (top) and after PEG addition (bottom) showing large deformation in the z-axis (scalebar = 3µm).

b, Example of a 3T3 fibroblast nucleus showing that the z-axis gets compressed instantly after PEG addition whereas

major and minor axes expand in the beginning followed by reduction in the length showing anisotropy in nuclear

compression with time. **c,** Nuclear deformation percentage along the major, minor and z-axis in wild type nuclei for

different lamin A expression (n=7) showing higher deformation along z-axis compared to the major and minor axis.

Lamin A density is spatially heterogeneous and leads to a variation in local compliance. While

overall lamin A expression levels are typically considered for overall nuclear compliance, we find

that the spatial distribution of lamin A is not uniform in the nuclear envelope (Fig. 3a and 3b). This

inhomogeneity also increases with overall lamin A expression, as revealed by plotting the variance

of local lamin A fluorescence intensity as a function of overall expression (Fig. 3c, supplementary

information Fig. S5). The histogram of local lamin A fluorescence shows narrow and high peak at

lower lamin A fluorescence levels but the peak broadens and flattens with increase in local lamin

A fluorescence levels indicating that lamin A distribution gets wider or heterogenous with increase

in lamin A expression (Fig. 3d). We also observed the bin span from Fig. 3d as a function of total

lamin A expression to show that the range of lamin A fluorescence covered increased with lamin

A expression showing a greater heterogeneity and vice-versa (Fig. 3e).

Figure 3│ Lamin A is spatially heterogeneous in the nucleus. a, Example 3T3 fibroblast cells transfected with lamin A chromobody showing variability in the spatial distribution along a cross-section nuclear membrane (scalebar = 3μm). **b,** Quantification of lamin A heterogeneity along the nuclear membrane circumference in panel **a**. **c,** Normalized variance of lamin A chromobody fluorescence as a function of total lamin A expression in different nuclei shows that higher expression of lamin A increases the heterogeneity of distribution (n=12). **d,** The histogram gets wider with increasing total lamin A fluorescence showing a wider range of local lamin A fluorescence or heterogeneity when overall lamin A expression increases. **e**. The bin span correlates linearly with lamin A expression showing a more heterogeneous and wider distribution at high lamin A fluorescence and vice-versa.

We then characterized the relationship between local lamin A expression and local nuclear deformation by quantifying the lamin A distribution via fluorescence and the nuclear deformation map in the X-Y and Z plane along the nuclear membrane using FIDVC (Fig. 4a, 4b, and 4c). These images suggest an inverse relationship between spatial variations of lamin A and deformation. To quantify the spatial variations, we plotted lamin A fluorescence with respective deformations along the nuclear circumference, showing the deformation as a function of the fraction of lamin A present at the nuclear membrane (Fig. 4d and Fig. 4e). We observed the least nuclear deformation in regions of highest lamin A expression and maximum deformation at regions of lowest lamin A expression. The ratio of applied osmotic stress to local nuclear deformation as a function of the lamin A expression yields the effective contribution of lamin A to nuclear stiffness; this reveals that the stiffness increases as lamin A expression increases locally (Fig. 4f). While this finding is consistent in all cells measured, some cells also presented low deformations in regions of low lamin A expression (see supplementary information Fig. S6), which may be attributed to the mechanical anisotropy of the underlying chromatin[27]. This may also contribute to the observed mechanical anisotropy of the nucleus along the major and minor axes of the nucleus, where the minor axis is deformed slightly more than the major axis (Fig. 4g). The correlation coefficient

between local deformation and local lamin A fluorescence for 10 cells was found to be negative showing a moderate anti-correlation between them but we did not find any relation of the correlation coefficient values with the total lamin A expression (Fig. 4h). This variation in anticorrelation could be due to the effect on the underlying chromatin.

Figure 4 | Local nuclear deformation as a function of lamin A distribution in 3T3 fibroblasts. a, Lamin A distribution along the nuclear membrane in X-Y plane. **b,** Example strain X-Y plane map of a 3T3 nucleus under 936.5kPa stress (scalebar = 5µm). **c,** Lamin A distribution (top) and Y-Z strain map (bottom) along the nuclear membrane. **d,** Lamin A fluorescence and nuclear deformation along the nuclear membrane. **e,** Deformation plotted against the ratio of local lamin A fluorescence at the membrane and total nuclear fluorescence. **f,** Ratio of stress to deformation (effective stiffness) plotted against the ratio of local lamin A fluorescence at the membrane and total nuclear fluorescence. **g,** Statistical distribution of overlaid major and minor axis. **h,** The correlation coefficient between local deformation and local lamin A fluorescence was negative showing anticorrelation but with no relationship with total lamin A expression.

Discussion: Here, we have quantitatively examined the role of lamin A in nuclear mechanics. Using osmotic pressure, we applied precisely controlled stresses and measured global volumetric changes. We find that the overall bulk modulus of the nucleus increasing with expression levels

of Lamin A. Decomposing nuclear compression, we identify anisotropic deformation of the

nucleus showing maximum deformation in z-axis followed by minor axis and then major axis,

respectively. Inspecting the distribution of lamin A more closely, we found distinct spatial

variations in concentration within the nuclear envelope; interestingly, the heterogeneity of lamin

A density increased with overall expression level. We suspected that these spatial variations in

concentration would lead to similar spatial variations in nuclear stiffness. By quantifying the

spatial variations of lamin A density and comparing this with a sub-micron resolution deformation

map of the nucleus, we indeed found a strong inverse correlation between local sub-micron lamin

A concentration and nuclear deformation, demonstrating that the spatial variation of lamin leads

to heterogeneous nuclear membrane deformation. Also, the local deformations along the minor

axes of the nuclei were found to be larger than the corresponding major axes, which was consistent

with our previous observation, as seen in the bulk deformation of the nuclei.

The expression and distribution of lamin A are critical in nuclear mechanics, with variations or

abnormalities in lamin A affecting nuclear stiffness leading to altered nuclear deformation.

Pathological conditions such as HGPS[28] and EDMD[29] are related to a mutation in the *lmna* gene

suggesting a potential role of nuclear deformation in nuclear mechanotransduction in the cell, but

the relationship between nuclear deformation and how it changes the gene expression is not clear.

The strongest evidence to this hypothesis comes from a recent study, where it is shown that when

the nucleus is physically compressed, there is translocation of an enzyme Histone deacetylase 3

(HDAC3) inside the nucleus. The role of this enzyme is to remove the acetyl group present on

histone residues which neutralize the positive charges present on the histones and inhibits its

interaction with the negatively charged DNA resulting in an open conformation of chromatin

known as euchromatin (non-compact expressing chromatin conformation). Thus, when the nucleus

is mechanically compressed, HDAC3 translocates in the nucleus to remove the acetyl group on

the histone leading to a tighter histone-DNA interaction promoting the non-expressing

heterochromatin (highly compact non-expressing chromatin) conformation of the chromatin[30].

Another factor contributing to nuclear stiffness is the tethering of chromatin itself with the nuclear

lamina[31]. Since the chromatin can be present as heterochromatin and euchromatin, it contributes

to the mechanical anisotropy of the nuclear envelope[32]. While lamin A dominates nuclear stiffness

under high mechanical stress, under small stresses, chromatin has a crucial role as well[27]. More

condensed heterochromatin regions are stiffer and experience less deformation compared to

euchromatin[33], making chromatin dynamics additionally important in nuclear

mechanotransduction. Nevertheless, the physical association between the nuclear lamina and the

chromatin, as well as the relationship between nuclear deformation and its direct effect on

epigenetic changes are still unclear and could prove to be essential in relating nuclear mechanics

with various pathological conditions.

Previous mechanotransduction studies have shown that mechanical forces can lead to translocation

of mechanosensitive proteins such as YAP from the cytoplasm to the nucleus under nuclear

tension[34], which affects properties like cell proliferation and differentiation in stem cells. Lamin

A could prove to be a critical factor in elucidating this relationship as it directly affects the nuclear

stiffness and its deformation. By manipulating lamin A expression, relating the change in nuclear

mechanics, and measuring downstream effects, we may better understand how nuclear stiffness

determines cellular fates, thus providing novel mechanics-centered strategies to correct defects in

diseases.

Acknowledgments: The authors thank members of Christian Frank's lab, University of

Wisconsin, for assistance with MATLAB FIDVC. AJE acknowledges support from NSERC

(RGPIN/05843-2014, EQPEQ/472339-2015, RTI/00348-2018), CIHR (Grant # 143327), and the

Canadian Foundation for Innovation (Project #32749).

References

1. Lamond, A. I. & Earnshaw, W. C. Structure and function in the nucleus. *Science* **280**, 547–553 (1998).

2. Kirby, T. J. & Lammerding, J. Emerging views of the nucleus as a cellular mechanosensor. *Nat. Cell Biol.*
20, 373–381 (2018).

3. Chen, J. *et al.* Cell mechanics, structure, and function are regulated by the stiffness of the three-dimensional
microenvironment. *Biophys. J.* **103**, 1188–1197 (2012).

4. Vaziri, A. & Mofrad, M. R. K. Mechanics and deformation of the nucleus in micropipette aspiration
experiment. *J. Biomech.* **40**, 2053–2062 (2007).

5. Caille, N., Thoumine, O., Tardy, Y. & Meister, J. J. Contribution of the nucleus to the mechanical properties
of endothelial cells. *J. Biomech.* **35**, 177–187 (2002).

6. Adele Khavari, A. J. E. Nuclei deformation reveals pressure distributions in 3D cell clusters. *PLoS One*
(2019).

7. Chu, F.-Y., Haley, S. C. & Zidovska, A. On the origin of shape fluctuations of the cell nucleus. *Proc. Natl.*
Acad. Sci. **114**, 10338–10343 (2017).

8. Pajerowski, J. D., Dahl, K. N., Zhong, F. L., Sammak, P. J. & Discher, D. E. Physical plasticity of the
nucleus in stem cell differentiation. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 15619–15624 (2007).

9. Lin, H.-H. *et al.* Mechanical phenotype of cancer cells: cell softening and loss of stiffness sensing.
Oncotarget **6**, (2015).

10. Bell, E. S. & Lammerding, J. Tumour Progression. **95**, 449–464 (2017).

11. Hetzer, M. W. The nuclear envelope. *Cold Spring Harb. Perspect. Biol.* **2**, a000539 (2010).

12. Dechat, T., Adam, S. A., Taimen, P., Shimi, T. & Goldman, R. D. Nuclear Lamins. 1–22 (2010).

13. Lammerding, J. *et al.* Lamin A / C deficiency causes Tema Grupo defective nuclear mechanics and
mechanotransduction. *J. Clin. Invest.* **113**, 370–378 (2004).

14. Hanson, L. *et al.* Vertical nanopillars for in situ probing of nuclear mechanics in adherent cells. *Nat.*
Nanotechnol. **10**, 554–562 (2015).

15. Guilluy, C. *et al.* Isolated nuclei adapt to force and reveal a mechanotransduction pathway in the nucleus.
Nat.Cell Biol. **16**, 376–381 (2014).

16. Harada, T. *et al.* Nuclear lamin stiffness is a barrier to 3D migration, but softness can limit survival. *J. Cell*
Biol. **204**, 669–682 (2014).

17. Wang, Y., Jiang, J., He, L., Gong, G. & Wu, X. Effect of lamin-A expression on migration and nuclear
stability of ovarian cancer cells. *Gynecol. Oncol.* **152**, 166–176 (2019).

18. Moir, R. D. *et al.* Review: The dynamics of the nuclear lamins during the cell cycle - Relationship between
structure and function. *J. Struct. Biol.* **129**, 324–334 (2000).

19. Schirmer, E. C. & Gerace, L. The stability of the nuclear lamina polymer changes with the composition of
lamin subtypes according to their individual binding strengths. *J. Biol. Chem.* **279**, 42811–42817 (2004).

20. Crisp, M. *et al.* Coupling of the nucleus and cytoplasm. *J. Cell Biol.* **172**, 41–53 (2006).

21.   Barateau, A. *et al.* A novel lamin a mutant responsible for congenital muscular dystrophy causes distinct
abnormalities of the cell nucleus. *PLoS One* **12**, 1–18 (2017).

22.   Lammerding, J. *et al.* Lamins a and C but not lamin B1 regulate nuclear mechanics. *J. Biol. Chem.* **281**,
25768–25780 (2006).

23.   Mayer, C. R., Arsenovic, P. T., Bathula, K., Denis, K. B. & Conway, D. E. Characterization of 3D Printed
Stretching Devices for Imaging Force Transmission in Live-Cells. *Cell. Mol. Bioeng.* **12**, 289–300 (2019).

24.   Zhou, E. H. *et al.* Universal behavior of the osmotically compressed cell and its analogy to the colloidal
glass transition. *Proc. Natl. Acad. Sci.* **106**, 10632–10637 (2009).

25.   Guo, M. *et al.* Cell volume change through water efflux impacts cell stiffness and stem cell fate. *Proc. Natl.*
Acad. Sci. (2017). doi:10.1073/pnas.1705179114

26.   Bar-Kochba, E., Toyjanova, J., Andrews, E., Kim, K. S. & Franck, C. A Fast Iterative Digital Volume
Correlation Algorithm for Large Deformations. *Exp. Mech.* **55**, 261–274 (2015).

27.   Stephens, A. D., Banigan, E. J., Adam, S. A., Goldman, R. D. & Marko, J. F. Chromatin and lamin A
determine two different mechanical response regimes of the cell nucleus. *Mol. Biol. Cell* **28**, 1984–1996
(2017).

28.   Goldman, R. D. *et al.* Accumulation of mutant lamin A progressive changes in nuclear architecture in
Hutchinson-Gilford progeria syndrome. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 8963–8968 (2004).

29.   Håkelien, A. M., Delbarre, E., Gaustad, K. G., Buendia, B. & Collas, P. Expression of the myodystrophic
R453W mutation of lamin A in C2C12 myoblasts causes promoter-specific and global epigenetic defects.
Exp. Cell Res. **314**, 1869–1880 (2008).

30.   Damodaran, K. *et al.* Compressive force induces reversible chromatin condensation and cell geometry–
dependent transcriptional response. *Mol. Biol. Cell* **29**, 3039–3051 (2018).

31.   Schreiner, S. M., Koo, P. K., Zhao, Y., Mochrie, S. G. J. & King, M. C. The tethering of chromatin to the
nuclear envelope supports nuclear mechanics. *Nat. Commun.* **6**, 1–13 (2015).

32.   Berendes, H. D. & Keyl, H. G. Distribution of DNA in heterochromatin and euchromatin of polytene nuclei
of Drosophila hydei. *Genetics* **57**, 1–13 (1967).

33.   Hampoelz, B. & Lecuit, T. Nuclear mechanics in differentiation and development. *Curr. Opin. Cell Biol.* **23**,
668–675 (2011).

34.   Driscoll, T. P., Cosgrove, B. D., Heo, S. J., Shurden, Z. E. & Mauck, R. L. Cytoskeletal to Nuclear Strain
Transfer Regulates YAP Signaling in Mesenchymal Stem Cells. *Biophys. J.* (2015).
doi:10.1016/j.bpj.2015.05.010