1
2
3 **Spatial and phylogenetic structure of Alpine stonefly community assemblages across**
4 **seven habitats using DNA-species**
5

6 Maribet Gamboa[1], Joeselle Serrana[1], Yasuhiro Takemon[2], Michael T. Monaghan[3,4], Kozo
7 Watanabe[1,5]
8
9
10 [1]Ehime University, Department of Civil and Environmental Engineering, Matsuyama, Japan
11
12 [2]Water Resources Research Center, Disaster Prevention Research Institute, Kyoto University,
13 6110011 Gokasho, Uji, Japan
14
15 [3]Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Müggelseedamm 301,
16 12587 Berlin, Germany
17
18 [4]Institut für Biologie, Freie Universität Berlin, Königin-Luise-Str. 1-3, 14195 Berlin, Germany
19
20 [5]Ehime University, Center Marine Environmental Studies (CMES), Matsuyama, Japan

Abstract

1. Stream ecosystems are spatially heterogeneous, with many different habitat patches distributed within a small area. The influence of this heterogeneity on the biodiversity of benthic insect communities is well documented; however, studies of the role of habitat heterogeneity in species coexistence and community assembly remain limited. Here, we asked how habitat heterogeneity influences spatial structure (beta biodiversity) and phylogenetic structure (evolutionary processes) of benthic stonefly (Plecoptera, Insecta) communities.

2. We sampled 20 sites along two Alpine rivers, including seven habitats in four different reaches (headwaters, meandering, bar-braided floodplain, and lowland spring-fed). We identified 21 morphological species and delineated 52 DNA-species based on sequences from mitochondrial *cox1* and nuclear ITS markers. Using DNA-species, we first analysed the patterns of variation in richness, diversity, and composition by quantifing the contribution of each of the four reaches and seven habitats to the overall DNA-species diversity using an additive partition of species diversity analysis and distance-based redundancy analysis. Using gene-tree phylogenies, we assessed whether environmental filtering led to the co-occurrence of DNA-species using a two-step analysis to find a phylogenetic signal.

3. The four reaches significantly contributed to DNA-species richness; with the meandering section displaying the highest contribution. However, we found that habitats had an effect on DNA-species diversity, where glide, riffle, and pool influenced the spatial structure of stonefly communities possibly due to a high species turnover.

4. Among the habitats, the pool showed significant phylogenetic clustering, suggesting high levels of evolutionary adaptation and strong habitat filtering. This community structure may be caused by long-term stability of the habitat and the similar requirements for co-ocurring species.

5. Our study shows the importance of different habitats on the spatial and phylogenetic structure of stonefly community assemblies and sheds light on the habitat-specific diversity that may help improve conservation practices.

KEYWORDS
DNA-species, habitats, spatial structure, phylogenetic structure, Plecoptera

1. INTRODUCTION

Understanding species diversity patterns and processes governing species coexistence in a community is one of the fundamental research objectives in ecology and biodiversity. The development of effective and suitable biodiversity conservation strategies requires knowledge of the processes driving community variation (Socolar, Gilroy, Kunin, & Edwards, 2016) along temporal and spatial gradients (Anderson et al., 2011). Habitat specialisation is known to influence changes in community composition along temporal and spatial gradients (e.g., Wiens et al., 2010); however, this change depends on the degree of specialisation of the species within a community (Rádková et al. 2014).

Stream ecosystems provide one of the most heterogeneous landscapes, given the dynamic interaction between spatial elements (e.g., topography) and ecological process (e.g., hydrology) (Benda et al., 2004; Tockner & Stanford, 2002). This interaction creates a variety of different habitats throughout the longitudinal (upstream-downstream) gradient of the river, classified as lotic (running water) and lentic (standing water) (Dobson & Frid, 1998; Calow & Petts, 1996; Hauer & Lamberti, 1996). At the biogeographic scale, habitat heterogeneity in a river channel defines environmental patches that impact aquatic community composition and species distribution (Brasil, Da Silva, Batista, Olivera, & Ramos, 2017; Dias-Silva, Cabetter, Juen, & De Marco, 2013); while at the local scale, creates environmental filtering to sort species with similar requirements (Saito, Cianciaruso, Siqueira, Fonseca-Gessner, & Povoine, 2016; Mykra, Heino, &, Muotka, 2007; Webb, Ackerly, McPeek, & Donoghue, 2002). Aquatic insects are abundant, diverse, and broadly distributed (Lancaster & Downes, 2013). Many studies have reported a strong positive relationship between aquatic insects species diversity and spatial habitat heterogeneity (e.g. Astorga et al., 2014; Karaus, Larsen, Guillong, & Tockner, 2013; Arscott, Tockner, & Ward, 2005; Benda et al., 2004; Batista, Buss, Dorville, & Nessimian, 2001), however, these were only based on one spatial scale, where the relationship between assembly process on both biogeographical and local scales remains limited to a few studies. Thus, understanding local trends in biodiversity is crucial to describe the interaction between habitat and species diversity at different scales.

One of the challenges in biodiversity research is that of species delimitation. Due to limitations in taxonomic expertise and incomplete descriptions of taxonomic groups, many species may remain unidentified or unclassified (Bickford et al., 2007), making it difficult to describe local biodiversity. Genetic methods developed over the last decade have helped with species identifications and the clarification of species boundaries of a wide range of taxa, including aquatic insects (e.g., Gamboa & Arrivillaga-Henriquez, 2019; Serrana, Miyake, Gamboa, & Watanabe, 2019; Gattoliatt, Rutschmann, Monaghan, & Sartori, 2018). The Generalized Mixed Yule Coalescent model (GMYC; Pons et al., 2006) is a method commonly used to inditify putative DNA-species, by quantifying the transition between inter- and intra-species branching events on an ultrametric tree (i.e., where the distance from the root of every branch is equal), using molecular markers, such as mitochondrial DNA (Fujisawa &

Barraclough, 2013; Mynott, Webb, & Suter, 2011), or based on the congruency between
mitochondrial and nuclear DNA (Rutschmann et al., 2016).

Stream ecologists have reported that local environmental heterogeneity may be the
strongest driver of community assembly patterns using putative DNA-species diversity of
aquatic insects along the river (Astorga et al., 2014; Baselga et al., 2013; Finn, Bonada, Múrria,
& Hughes, 2011). However, the majority of these observations is based on DNA-species
diversity changes along the river's longitudinal gradient (Finn, Zamora-Muñoz, Múrria, Sáinz-
Bariáin, & Alba-Tercedor, 2013; Gill, Harrington, Kondratieff, Zamudio, Poff, & Funk, 2013;
Jackson, Battle, White, Pilgrim, Stein, Miller, & Sweeney, 2013; Finn et al., 2011; Finn & LeRoy
Poff, 2011; Hughes, Schmidt, & Finn, 2009), where the influence of other river dimensions (e.g.,
habitats), has only been described in few species, such as beetles (Ribera & Vogler, 2008) or
caddisflies (Marten, Brandle, & Brandl, 2006). Therefore, investigating variations in spatial
patterns of aquatic insects community composition using DNA-species in the river channel and
their relationship with habitat heterogeneity will help to improve the understanding of the
process of community assembly in aquatic ecosystems.

The understanding of the community assembly not only has focused on spatial patterns
and composition changes analysis. The phylogenetic signal has been proposed as an
additional insight to understand the drivers of community assembly (Violle et al., 2011; Webb
et al., 2002). The basis of this approach is to compare ecological traits and phylogenetic
structures in communities with those expected under null models (i.e., phylogenetic signal).
Environmental filtering is thought to cause related species to co-occur (i.e., phylogenetic
clustering) more than what is expected by chance, as closely-related species are assumed to
share similar physiological requirements. By contrast, competitive exclusion causes the
opposite pattern (phylogenetic overdispersion) as species compete for the same limiting
resources (Cavender-Bares & Wilczek, 2003). Although phylogenetic clustering and
overdispersion have been observed in aquatic insect community assemblies at spatial scales
(Saito, Soininen, Fonseca-Gessner, & Siquiera, 2015a; Saito, Siquiera, & Fonseca-Gessner,
2015b; Saito et al., 2016) or based in specific trait-based approaches (e.g., respiration strategy;
Buchwalter, Jenkins, & Curtis, 2002, and anthropogenic pollutants; Martin, Cain, Luoma, &
Buchwalter, 2007), the assess of the phylogenetic structure in different habitats have not been
explored.

This study aims to investigate the impact of habitat heterogeneity on changes in
stonefly community composition in different riverine locations, by assessing the relative role of
seven aquatic habitats on the beta biodiversity and phylogenetic signal on stonefly community
composition using DNA-based species. Stoneflies (Plecoptera) are aquatic insects that show
more complex responses to habitat change than other insects (Bojková, Rádková, Soldán, &
Zahrádková, 2014), such as ecological (Lancaster & Downes, 2013) and evolutionary
responses (Gamboa & Watanabe, 2019). We hypothesised that species distribution and
community assembly are related to specific habitats and that habitat filtering acts on closely-
related species with similar physiological requirements. By contrast, high levels of species

diversity are expected to limit similarities in community structure and lead to phylogenetic overdispersion. Using two molecular markers (*cox1* and ITS) to delineate putative species, we aimed to: (1) quantify the contribution of four reaches and seven habitats to the overall DNA-species diversity, (2) determine the influence of habitats on community assembly, and (3) compare the phylogenetic structure of stoneflies communities in each of the seven habitats.

2. METHODS

2.1 Study sites and sampling collection

We selected seven habitat types (Fig. 1), waterfall (vertical descent of watercourse), riffle (high turbulence water flow), glide (low turbulence water flow), pool (flowing water > 1 m deep), wando (small-sized water in the head or tail on a sandy bar; as described by Ishida, Abekura, & Takemon, 2005), side-channel (small-sized water pass flowing on a sandy bar), and isolated pond (isolated still water habitat on the sandy bar) over 20 sampling sites in two Alpine rivers in northeastern Italy. Fifteen sampling sites were located on the Tagliamento River (T01 to T15) and five were located in its major tributary, the Fella River (F01 to F05) (Fig. 2). Sampling was performed across four geomorphologically distinct reaches (4 to 1556 m a.s.l.): constrained headwaters, meandering (zigzagging river movement), bar-braided floodplain (convergent river bankfull width), and lowland spring-fed streams in the floodplain (Doering, Uehlinger, & Tockner, 2013). All seven habitat types were sampled at each sampling site. Habitat diversity per site was measured using Simpsons Diversity index with at least one individual per habitat and sampling site.

We collected qualitative samples of stonefly nymphs using D-frame nets (250 µm mesh), spending one hour per site sampled (total sampling time: 20h). Between one and five replicates per habitat per site were collected (n = 188), where each replicate represented a different random location within each habitat sampled. All samples collected in the field were immediately immersed in >99 % ethanol and rinsed twice with fresh >99 % ethanol after being transported to the laboratory. Morphological species identification was conducted using a stereoscopic microscope (80 X) following the classifications in Consiglio (1980) and Fochetti and Tierno de Figueroa (2008). Field sampling (all sites/habitats) was carried out twice in summer 2009 (10 July to 10 September) and once in early spring 2010 (24 March to 15 April).

2.2 DNA extraction and sequencing analysis

A total of 375 stonefly nymphs were collected (see Results). Genomic DNA was extracted from each nymph using DNeasy Blood and Tissue kits, according to the manufacturer's instructions (Qiagen GmbH, Hilden, Germany). In all individuals, a 658 bp fragment of mitochondrial *cox1* was amplified using LCO-1490 and HCO-2198 primers (Folmer,

Black, Hoeh, Lutz, & Vrijenhoek, 1994) with an annealing temperature of 48 °C and 30 PCR
cycles. The internal transcribed spacer (ITS) nuclear marker (average 930 bp) was amplified in
a subset of 116 individuals chosen to represent all morphological taxonomic groups using ITS1
and ITS2 primers (McLain, Wesson, Oliver, & Collins, 1995) with an annealing temperature of
58 °C and 40 PCR cycles. PCR products were purified using the QIAquick PCR Purification Kit
(Qiagen) and sequenced in both directions using the aforementioned primers. *Cox1* sequences
were analysed on a 3500xL automated sequencer (Applied Biosystems) and ITS sequences
were analysed commercially by Eurofins – Operon, Tokyo. All sequencing data reported in this
study have been deposited into GenBank.
Forward and reverse reads were assembled and edited using CodonCode Aligner v
3.5 (Codon Code Corporation, Dedham, USA). Multiple sequence alignment was then
performed using ClustalW (align.genome.jp; Larkin et al., 2007). Highly variable regions of ITS
were removed using the Indel Module of SeqFIRE v 1.0.1 (https://omicstools.com/seqfire-tool;
Ajawatanawong, Atkinson, Watson-Haigh, MacKenzie, & Baldauf, 2012) with the default
settings. This reduced the ITS alignment length from 915-1016 bp to 790 bp. For both markers,
identical sequences were collapsed using CleanCollapse v. 1.0.5
(https://sray.med.som.jhmi.edu/SCRoftware/CleanCollapse/). Sequences of mtDNA and nDNA
markers were compared to the NCBI nucleotide database using blastn queries
(http://blast.ncbi.nlm.nih.gov) to verify species identification (>90 and 80 % similarity for mtDNA
and nDNA, respectively).
Genetic diversity for each marker was calculated as the number of polymorphic sites,
nucleotide diversity (Kimura 2-parameter model), and the number of haplotypes using DnaSp
v. 5.10 (Librado & Rozas, 2009).
204
205 **2.3 Data analysis**
206
207 **2.3.1 DNA-based species delimitation**
208
Putative DNA-species were delineated using the General Mixed Yule Coalescent
approach (GMYC; Fujisawa & Barraclough, 2013). First, ultrametric gene trees of *cox1* and ITS
were constructed independently using BEAST v. 1.8.3 (Drummond, Suchard, Xie, & Rambaut,
2012) with a relaxed lognormal clock and a coalescent prior run for 50 million generations. The
results were summarised with the Tree Annotator (BEAST package). A GMYC analysis was
run using the splits package (Ezard, Fujisawa, & Barraclough, 2014) in R v. 3.3 (R Core Team,
2014). We used the single-threshold GMYC model based on Fujisawa and Barraclough (2013).
The maximum likelihood of each GMYC model was tested using the likelihood ratio test against
a one-species null model (where the entire tree is considered as a single coalescent). Outputs
for *cox1* and ITS were compared and the results were combined to generate the list of DNA-
species.
220

2.3.2 Spatial structure of stonefly community assemblies

DNA-species diversity per reach and per habitat were calculated using the Shannon-Weaver and Simpsons diversity indices using the vegan package (Oksanen et al., 2012) in R. The contribution of each of the four reaches and seven habitat types to overall DNA-species diversity was quantified using an additive partition of species diversity analysis (Lande, 1996). This analysis allows alpha and beta values to be simultaneously calculated using an abundance matrix organized hierarchically among habitats (alpha diversity, α) and reaches (beta diversity, β) (Gering, Crist, & Veech, 2003). Simpsons and Shannon-Weaver diversity were partitioned into alpha and beta diversity to observe whether the overall value was greater than the mean values (Lande, 1996). The statistical significance of each component was tested with a randomisation procedure (10,000 randomisations) using the adipart function of the vegan package in R. We then partitioned beta diversity into turnover (i.e., the replacement of species by other species in different habitats) and nestedness (i.e., species loss or gain between habitats) components (Baselga et al., 2013) using a Sorenson dissimilarity matrix and the betadisper function of the vegan package in R. We used a null model with 10,000 randomisations to test if the results of each component were greater than expected by chance.

2.3.3 Habitat influence on community assembly

We performed a distance-based redundancy analysis (db-RDA; Legendre & Anderson, 1999) to quantify the influence of habitats on spatial community assemblies between sampling sites. We first conducted Principal Coordinates Analysis (PCoA) using a community dissimilarity matrix (the pairwise distance of DNA-species within each sampling site) calculated as the Bray-Curtis index. Then, we used the resulting eigenvalues as the "response" variable in the db-RDA models, while geographical distance and habitat influence were used as explanatory variables. The geographical distance between each pair of sampling site was calculated as the Euclidean distance extracted from the geographical coordinates as proposed by Gamboa and Watanabe (2019) using the Vicenty Ellipsoid package (Kamey, 2013) in R. To analyse the contribution of the different habitats to community assembly we calculated a contribution diversity approach matrix proposed by Lu, Wagner & Chen (2007). This approach calculated a differentiation coefficient to evaluate the distribution of DNA-species diversity in a given habitat within the sampling sites. We obtained the best model by first including all explanatory variables (seven habitats and geographical distance) then running nested models using the vegdist and capscale functions in the vegan package in R. After identifying the best model, we calculated the percentage of explained variance obtained by the db-RDA as suggested by Peres et al. (2006) using the ordistep function in vegan. We performed a permutation test (1000 permutations) and used ANOVA to test each explanatory variable of the db-RDA model for statistical significance using the varpart function in vegan.

We analysed multivariate dispersion as a measure of beta diversity based on a
community dissimilarity matrix using DNA-species to further investigate habitat influence on
beta diversity. We calculated beta diversity as the distance to group centroid (the homogeneity
of a community in a given habitat) based on a Bray-Curtis dissimilarity matrix. The community
matrix consisted of DNA-species richness in each sampling site at both rivers. We also
analysed an homogeneity of multivariate dispersions using a linear model in the betadisper
function in the R package vegan with a permutation test (1000 permutations).
267
268 **2.3.4 Phylogenetic structure of the stonefly community**
269
The phylogenetic relationship among species was estimated using Maximum-
Likelihood (ML) gene trees for *cox1* and ITS separately in PhyML 3.1 (Guindon & Gascuel,
2003), with the default settings under a GTR+I+G model as obtained in jModeltest 3.0 (Posada,
2008). Node support was determined by bootstrapping (10,000 times). We used sequences of
Orthoptera sp. as outgroups in the *cox1* (HM381647) and ITS (KT440350) trees based on the
reported phylogenetic relationship with stoneflies (Misof et al., 2014). Additional adult
specimens (Gamboa & Monaghan, 2015) were added on the *cox1* tree to corroborate nymph
species identification.
A congruency index (de Vienne, Giraud, & Martin, 2008) calculated using the web
version of MAST (http://max2.ese.u-psud.fr/icong/index.help.html), indicated that *cox1* and ITS
tree topologies were congruent (Icong = 1.9, p<0.05). Therefore, we concatenated the two
markers using the FASconCAT-G perl script (https://github.com/PatrickKueck/FASconCAT-G).
Using this matrix, we quantified multisite phylogenetic similarities (i.e., similarity measures
comparing more than two sampling sites) as proposed by Leprieur et al. (2012). We quantified
multisite phylogenetic turnover, nestedness and phylo-beta diversity index (known as Sorensen
similarity index) for every community by calculating the sum of the branch length in a phylogeny
connecting all species in a community. We then observed the proportion of shared branch
lengths between two sampling sites. We performed the analysis using the phylo.beta.multi
function in the R package betapart (Baselga & Orne, 2012). We also performed a two-step
analysis to check for a phylogenetic signal in stonefly communities within each habitat. First,
we estimated niche conservatism (i.e., closely related species are ecologically similar and thus
share similar habitats; Wiens et al., 2010) using Blomberg's K-statistic (Blomberg, Garland, &
Ives, 2003) and the phylogenetic independent contrast (PIC) test (Felsenstein, 1985) within
each habitat. The Blomberg's K test and PIC were implemented using the phytools package
(Revell, 2012) and picante package (Kembel et al., 2010) in R. Both tests were run for 10,000
iterations to obtain the null distribution. If the p-values of the observed versus random variance
were <0.05, we interpreted them as evidence of niche conservatism. Second, we calculated
the Net Relatedness Index (NRI) and the Nearest Taxon Index (NTI) to measure the degree of
phylogenetic clustering at the community level (Webb et al., 2002) within each habitat. NRI
measures the phylogenetic dispersion of a community assemblage (community associated with

the same habitat type) by comparing the observed mean pairwise phylogenetic distance between species in a community assemblage to the null model, while NTI measures the phylogenetic dispersion of a community assemblage by comparing the observed mean nearest phylogenetic neighbour distance between species in a community assemblage to the null model. Positive values of NRI and NTI indicate that community assemblages with similar habitat preferences are phylogenetically clustered (more closely related than expected), while negative values indicate phylogenetic overdispersion (more distantly related than expected). The NRI and NTI were calculated using the ses.mpd and ses.mntd functions, respectively, in R package picante, with 1000 replications.

3. RESULTS

We identified a total of 21 morphological species in 12 stonefly genera (Table S1) based on collections of 5-40 individuals (mean = 19) and 1-8 morphological species (mean = 4) from each of the 20 sampling sites (Table 1). There were 184 *cox1* haplotypes from the 375 sequenced individuals and 87 ITS genotypes from the subset of 116 sequenced individuals. Samples were dominated by *Leuctra major* (50 and 40 % of the total haplotypes (*cox1*) and genotypes (ITS), respectively; Table S1). For both markers, the log-likelihood of the GMYC model at the threshold (*cox1* = 2514.495; ITS = 2819.385) was significantly higher than the null model for a single coalescent (*cox1* = 2400.932; ITS = 2701.875), as shown by the likelihood ratio test (p < 0.001). Both genes delimited 52 individual putative species, 38-59 species (95% CI) for *cox1* and 40-64 species (95% CI) for ITS (Table S1). The *cox1* gene was composed of 42 clusters and ten singletons, while ITS was composed of 31 clusters and 21 singletons. Intraspecific nucleotide diversity for *cox1* ranged from 0-13 % among putative DNA-species, being the highest value for *L. major* (Table S1). Fourteen of the 21 putative morpho-species were split into multiple DNA-species. All DNA-species occurred in more than one habitat, sampling site, and reach.

3.1 Spatial structure of stonefly community assemblies

Among the four reaches, the meandering contained the highest number of DNA-species (n = 52) (Table 1) and the highest Shannon-Weaver (0.89) and Simpson (0.6) diversities (Table S2). The single most species-rich site was T08 in the meandering section (n = 15 DNA-species) (Table 1). The headwater section contained the highest number of cryptic species (25 of 31 total cryptic DNA-species). Among the seven habitats, riffles contained the highest number of DNA-species (n = 31) followed by glides (n = 25) (Table 2 and S2). Riffles was the most species-rich habitat in the meandering section.

The relative contribution of the four reaches and seven habitats to overall DNA-species diversity showed that reaches were the most important driver for richness (57 %, p < 0.001, Fig. 3), while habitats were the most important driver for species diversity (Shannon-Weaver

index; 88 %, p < 0.001). DNA-species turnover among habitats showed a significantly high species turnover (0.60) than species nestedness (0.16) to overall beta diversity (0.79) (p = 0.0016).

Spatial variation in DNA-species was only found to be significant in three habitats: riffle, glide, and pool (p<0.05, Fig. 4). Among the seven habitats, the riffle and glide had the highest diversity dissimilarity (F = 1.43, p = 0, Fig. 5), suggesting that the community in these two habitats were less similar among each other that in the remaining habitats.

3.2 Phylogenetic structure of stonefly community

The mtDNA phylogeny recovered all recognised morphospecies as clades; however, 11 morphospecies were composed of multiple lineages. These lineages likely reflect cryptic species, concurring with GMYC results. The nuclear ITS tree showed congruence with the *cox1* tree, except those of *Nemoura cinerea* and *N. mortini* which were located in the same clade as *Capnia nigra* and *C. vidua* (Fig. S1).

The stonefly community phylogenetic diversity among sampling sites showed that species turnover (0.927) contributed more than nestedness (0.055) to beta diversity (0.98) (p < 0.005). Pool habitats showed significant niche conservatism and phylogenetic clustering based on the two-step analysis of phylogenetic signal (Blomberg's K test, PIC, NRI, and NTI, all *p* < 0.05; Table 2), suggesting high similarity among closely-related species within this habitat. Additionally, significant negative values for NRI and NTI indices were reported (*p* < 0.05, Table 2) in the wando and side channel for NRI, and riffle for NTI, indicating that the community assemblage presents phylogenetic overdispersion.

4. DISCUSSION

Community assemblies within the stream are strongly related to habitat heterogeneity within a given area (Astorga et al., 2014). We observed that different habitat types influenced Alpine stoneflies DNA-species diversity, their spatial structure, and their evolution.

In this study, the habitats were the most important driver of DNA-species diversity (diversity partition analysis, Shannon index), concurring with previous studies of aquatic invertebrates (Astorga et al., 2014; Finn & LeRoy Poff, 2011; Finn, Theobald, Black, & Poff, 2006; Finn et al., 2011; Ribera & Vogler, 2008; Marten et al., 2006). Other than niche heterogeneity, one of the main determinants of beta diversity may be the dispersal ability of the organisms (Heino et al., 2015; Astorga et al., 2014; Finn & LeRoy Poff, 2011). A high species turnover is associated with poor dispersal ability (Thompson & Townsend, 2006). Our results showed a high species turnover among habitats (species turnover = 0.60), suggesting that non-random migration is likely to occur among habitats, probably caused by environmental filtering or spatial restrictions in their original habitat. Stoneflies have relatively poor dispersal abilities

within stream reaches (Consiglio, 1980; Fochetti & Tierno de Figueroa, 2008), and tend to avoid

migration among habitats due to predator pressure (Tiziano, Fenoglio, Lopez-Rodriguez,

Tierno de Figueroa, Grenna, & Cucco, 2010). The habitat heterogeneity along the Tagliamento

and Fella rivers could potentially reduce the intra-stream stonefly dispersal, and might explain

the large number of cryptic species found (GMYC and phylogenetic analyses). However, high

species turnover is also associated with sampling bias. Non-common species among habitats

could be a consequence of an un-equal sampling effort or habitat availability. In this study,

habitat availability was similar across most sampling sites (average = 0.75, range = 0.55-0.84,

Table 1) and despite a similar sampling effort at every site, some species were not detected.

Therefore, future studies will need to increase sampling effort to estimate if sampling bias could

impact our interpretations of species turnover.

Among the seven habitats, the riffle, glide, and pool had significant impacts on the

stonefly community structure (according to the db-RDA analysis) and two habitats (riffle and

glide) has significantly higher species diversity (according to the multivariate dispersion

analysis). The riffle and glide habitats characterised by a high and low turbulence flow,

respectively; both play an essential role in determining habitat suitability for many species of

aquatic insects (Lancaster & Downes, 2013; Benda et al., 2004), particulary stoneflies

(Lancaster & Downes, 2013; Batista et al., 2001). The dynamic nature of riffle and glide habitats

influence density-dependent local competition and high organisms migration with their turbulent

flow, leading to high levels of biodiversity (Hughes et al., 2009; Arscott et al., 2005; Batista et

al., 2001). On the contrary, the pool (a deep-high river section with a slow/inexistent flow) is

only weekly correlated with aquatic insect community diversity (Herrera-Vasquez, 2008) due to

the specialised taxa living in this stable habitat (Pastuchova, Lehotsky, & Greeskova, 2008).

Surprisingly, we also found that the pool habitat led to a phylogenetic clustering of the

stonefly community, according to the four indices used in this study (Blomberg's K, PIC, NRI,

NTI). Long-term adaptation (Lososova et al., 2015), dispersal limitations (Saito et al., 2015a,

2015b), colonisation history (Aizen, Gleiser, Sabatino, Gilarranz, Bascompte, & Verdu, 2015),

and low resource competition (Webb et al., 2002) are all associated with phylogenetic clustering,

particulary in habitats linked with long-term stability. Pools are rarely affected by physical

perturbations (Buffington, Lisle, Woodsmith, & Hilton, 2002), and they may persist despite high

river flow (Calow & Petts, 1996), discharge (Rolls, Leigh, & Sheldon, 2012) and drought (Lake,

2003). Therefore, habitats with low levels of perturbation enhanced niche conservatism,

clustering, and evolutionary adaptation of stonefly communities in Alpine rivers. However, a

greater competitive asymmetry (i.e., an unequal resource division) among distant phylogenetic

relatives and facilitation among close phylogenetic relatives can also cause phylogenetic

clustering (Mayfield & Levine, 2010; Sargent & Ackerly, 2008). Future studies should address

this concern by studying the distribution of resources (i.e., food) among habitats, to obtain a

clear picture of the evolutionary of co-occurring species.

In riffle, wando and side channel habitats, a phylogenetic overdispersion pattern was

detected, which is commonly observed in aquatic insects (Saito et al., 2016; Violle et al., 2011).

Efficient colonisation and high dispersal patterns are thought to be driving factors for the
phylogenetic overdispersion of several species groups, such as birds (Sobral & Cianciaruso,
2016), mammals (Cardillo, Gittleman, & Purvis, 2008) and insects (Violle et al., 2011). These
characteristics have been rarely documented in stoneflies (e.g., *Leutra ferruginea*, Macneale,
Peckarsky, & Likens, 2005), however, stonefly nymph dispersal is driven by intra-stream drift
(Stewart & Stark, 2008). Phylogenetic overdispersion is common in highly-disturbed habitats
which tend to preserve high phylogenetic diversity (Xiu et al., 2017). Habitats with high-turbulent
water flow (riffle) and high physical perturbation such as sandy bar habitats (wando and side-
channel) in the Tagliamento river show a 62 % habitat turnover (Tockner et al., 2003) mainly
driven by two annual flooding events (Ward et al., 1999). Therefore, the phylogenetic
overdispersion of stonefly communities at riffle, wando, and side-channel habitats is likely
driven by the high level of disturbance in these habitats.

Overall, the spatial and the phylogenetic structure of stonefly community assemblies in
the Tagliamento and Fella rivers were promoted by the availability and stability of habitats.
Among the four reaches, the meandering harboured the largest number of DNA-species (n =
52) and was highly diverse among habitats (Shannon-Weaver index: 0.89). Meanderings are
highly complex morphodynamic sections of the river supporting high benthic invertebrate
diversity, given the suitability and resilience of their habitats within fluvial ecosystems (Garcia,
Schnauder, & Push, 2012). However, the headwaters exhibited a high number of cryptic
species (78% of the total DNA-species richness), as previously shown by Jackson et al., 2013;
Murria, Bonada, Arnedo, Prat, & Vogler, 2013; Finn et al., 2011; Hughes et al., 2009. This
tendency is due to lower canalisation, loss of connectivity with the stream network, and low
levels of anthropogenic disturbance (Finn & LeRoy Poff, 2011; Finn et al., 2011). Cryptic
species are often observed in stoneflies (e.g., Viteck, Vincon, Graf, & Pauls, 2017), possibly
due to introgression (Boumans, Hogner, Brittain, & Johnsen, 2017) or hybridisation (Young,
Smith, Pilgrim, Fairchlid, & Schwartz, 2019) which remain unresolved species. Headwaters in
the Tagliamento and Fella rivers have narrow valleys and gorges with low sinuosity and
confinement (Tockner et al., 2003); therefore, this reach enhances intraspecific genetic
differences among stoneflies species. Both meandering and headwater reaches strongly
influenced the maintenance of stonefly biodiversity in alpine rivers; therefore, their
environmental protection should be a priority in achieving conservation practices.

In conclusion, we demonstrated that different habitats played an essential role in the
spatial and phylogenetic structure of stonefly community assemblies in alpine regions.
Community diversity was linked to turbulent water flow, while evolutionary adaptation was
linked with habitats with allow long-term stability. Our results highlight the importance of
molecular studies on quantitative measures of biodiversity in given area and the essential role
of habitats in a region, which is particulary important to conserve community assemblages with
threatened species or within threatened habitats/landscapes (Esteban & Finlay, 2010).

ACKNOWLEDGEMENTS
461
We thank Paul Schmidt-Yáñez for his assistance in the field and Micanaldo Francisco for his
assistance with the figures. MG was supported by the German Academic Exchange Service
(DAAD) fellowship (A/09/94531) and the Japan Society of the Promotion of Science
Postdoctoral Fellowship (PU17908). KW was supported by a European Union Marie-Curie
International Incoming Fellowship (PIIF-GA-2009-237026). MTM was partially supported by a
Japan Society for the Promotion of Science (JSPS) Fellowship (L-15543). This research was
supported by the JSPS (Grant Numbers: 24254003, 17H01666), the Sumitomo Electric
Industries Group Corporate Social Responsibility Foundation, the German Academic Exchange
Service (DAAD, Programm Projektbezogener Personenaustausch Japan, project 57402018)
and the Research Unit Program of Ehime University.
472
473
474   **CONFLICT OF INTEREST**
475
476   The authors declare no conflict of interest.
477
478
479   **DATA AVAILABILITY STATEMENT**
480
481   The data that support the findings of this study are available form the corresponding author
482   upon reasonable request.

483
484

485   **REFERENCES**
486
Aizen, M. A., Gleiser, G., Sabatino, M., Gilarranz, L. J., Bascompte, J., & Verdu, M. (2015). The
phylogenetic structure of plant-pollinator networks increases with habitat size and isolation.
Ecology Letters, *19*, 29–36.
Ajawatanawong, P., Atkinson, G. C., Watson-Haigh, N. S., MacKenzie, B., & Baldauf, S. L.
(2012). SeqFIRE: a web application for automated extraction of indel regions and
conserved blocks from protein multiple sequence alignments. *Nucleic Acids Research*, *40*,
340–347.
Anderson, M. J., Crist, T. O., Case, J. M., Vellend, M., Inouye, B. D., Freestone, A. L., ….
Swenson, N. G. (2011). Navigating the multiple meanings of ß diversity: A roadmap for the
practicing ecologist. *Ecology Letters*, *14*, 19–28.
Arscott, D. B., Tockner, K., & Ward, J. V. (2005). Lateral organization of aquatic invertebrates
along the corridor of a braided floodplain river. *Journal of North American Benthological
Society*, *24*, 934–954.

Astorga, A., Death, R., Death, F., Paavola, R., Chakraborty, M., & Muotka, T. (2014). Habitat
heterogeneity drives the geographical distribution of beta diversity: the case of New
Zealand stream invertebrates. *Ecology and Evolution*, *4*, 2693–2702.

Baselga, A., & Orme, C. D. L. (2012). Betapart: and R package for the study of beta diversity.
Methods in Ecology and Evolution, *3*, 808–812.

Baselga, A., Fujisawa, T., Crampton-Platt, A., Bergsten, J., Foster, P. G., Monaghan, M. T., &
Vogler, A. P. (2013). Whole-community DNA barcoding reveals a spatio-temporal
continuum of biodiversity at species and genetics levels. *Nature Communications*, *4*,
10.1038.

Batista, D. F., Buss, D. F., Dorville, L. F. M., & Nessimian, J. L. (2001). Diversity and habitat
preference of aquatic insects along the longitudinal gradient of the Mace river basin, Rio de
Janeiro, Brazil. *Revista Brasileira de Biologia*, *6*, 249–258.

Benda, L., Poff, N. L., Miller, D., Dunne, T., Reeves, G., Pess, G., & Pollock, M. (2004). The
network dynamics hypothesis: how channel networks structure riverine habitats.
BioScience, *54*, 413–427.

Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., Ingram, K. K., &
Das, I. (2007). Cryptic species as a window on diversity and conservation. *Trends in
Ecology and Evolution*, *22*, 148–155.

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in
comparative data: behavioral traits are more labile. *Evolution*, *57*, 717–745.

Bojková, J., Rádková, V., Soldán, T., & Zahrádková, S. (2014). Trends in species diversity of
lotic stoneflies (Plecoptera) in the Czech Republic over five decades. *Insect Conservation
and Diversity*, *7*, 252–262.

Boumans, L., Hogner, S., Brittain, J., & Johnsen, A. (2017). Ecological speciation by temporal
isolation in a population of the stonefly *Leuctra hippopus* (Plecoptera, Leuctridae). *Ecology
and Evolution*, 7, 1635–1649.

Brasil, L. S., Da Silva, N. F., Batista, J. D., Olivera, B., & Ramos, H. S. (2017). Aquatic insects
in organic and inorganic hábitats in the streams on the Central Brazilian savanna. *Revista
Colombiana de Entomologia*, *43*, 286–291.

Buchwalter, D. B., Jenkins, J. J., & Curtis, L. R. (2002) Respiratory strategy is a major
determinant of [^{3}H]water and [^{14}C]chlorpyrifos uptake in aquatic insects. *Canadian Journal
of Fisheries and Aquatic Science*, *59*, 1315–1322.

Buffington, J. M., Lisle, T. E., Woodsmith, R. D., & Hilton, S. (2002). Controls on the size and
occurrence of pools in coarse-grained forest rivers. *River Research and Applications*, *18*,
507–531.

Calow, P. P., & Petts, G. E. (1996). *The Rivers Handbook: Hydrological and Ecological
Principles*. Blackwell Science LTD, Victoria, Austria.

Cardillo, M., Gittleman, J. L., & Purvis, A. (2008). Global patterns in the phylogenetic structure
of island mammal assemblages. *Proceedings of the Royal Society B: Biological Science*,
275, 1549–1556.

Cavender-Bares, J., & Wilczek, A. (2003). Integrating micro and macroevolutionary processes
in community ecology. *Ecology*, *84*, 592–597.

Consiglio, C. (1980). *Plecoptteri*. Consiglio Nazionale delle Ricerche AQ/1/77, Verona, Italy.

Dobson, M. & Frid, C. (1998). *Ecology of Aquatic Systems*. Longman, Harlow, UK.

Doering, M., Uehlinger, U., & Tockner, K. (2013). Vertical hydrological exchange, and
ecosystem properties and processes at two spatial scales along a floodplain river
(Tagliamento, Italy). *Freshwater Science*, *32*, 12–25.

Dias-Silva, K., Cabetter, H. S. R., Juen, L., & De Marco JR, P. (2010). The influence of habitat
integrity and physical-chemical water variables on the structure of aquatic and semi-aquatic
Heteroptera. *Zoologia*, *27*, 918–930.

Drummond, A. J., Suchard, M. A., Xie, D., & Rambaut, A. (2012). Bayesian phylogenetics with
BEAUti and the BEAST 1.7. *Molecular Biology and Evolution*, *29*, 1969–1973.

Esteban, G. F., & Finlay, B. J., (2010). Conservation work is incomplete without cryptic
biodiversity. *Nature*, *463*, 293.

Ezard, T., Fujisawa, T., & Barraclough, T. (2014). *Splits: SPecies lImits*. Retrieved from http://R-
Forge.R-project.org/projects/splits/

Felsenstein, J. (1985). Phylogenies and the comparative method. *American Naturalist*, *125*, 1–
15.

Finn, D. S., & LeRoy Poff, N. (2011). Examining spatial concordance of genetic and species
diversity patterns to evaluate the role of dispersal limitation in structuring headwater
metacommunities. *Journal of North American Benthological Society*, *30*, 273–283.

Finn, D. S., Bonada, N., Múrria, C., & Hughes, J. M. (2011). Small but mighty: headwaters are
vital to stream network biodiversity at two levels of organization. *Journal of the North*
American Benthological Society, *30*, 963–980.

Finn, D. S., Theobald, D. M., Black, W. C., & Poff, N. L. (2006). Spatial population genetic
structure and limited dispersal in a Rocky Mountain alpine stream insect. *Molecular*
Ecology, *15*, 3553–3566.

Finn, D. S., Zamora-Muñoz, C., Múrria, C., Sáinz-Bariáin, M., & Alba-Tercedor, J. (2013).
Evidence from recently deglaciated mountain ranges that *Baetis alpinus* (Ephemeroptera)
could lose significant genetic diversity as alpine glaciers disappear. *Freshwater Science*,
33, 207–2016.

Fochetti, R., & Tierno de Figueroa, J. M. (2008). *Plecoptera*. Fauna D' Italia, Calderini, Italy.

Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for
amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan
invertebrates. *Molecular Marine Biology and Biotechnology*, *3*, 294–297.

Fujisawa, T., & Barraclough, T. G. (2013). Delimiting species using single-locus data and the
generalized mixed yule coalescent approach: a revised method and evaluation on
simulated data sets. *Systematic Biology*, *62*, 707–724.

Garcia, X.-F., Schnauder, I., & Push, M. T. (2012). Complex hydromorphology of meanders can
support benthic invertebrate diversity in rivers. *Hydrobiologia*, *685*, 49–68.

Gamboa, M. & Arrivillaga-Henriquez, J. (2019). Biochemical and molecular differentiation of
Anacroneuria species (Plecoptera, Insecta) in Andean National Park, Venezuela.
Systematics and Biodiversity, *17*, 669–678.

Gamboa, M. & Monaghan, M.T. (2015). Association of adult female and male stoneflies
(Plecoptera) from an Alpine river using wing morphometrics and mitochondrial DNA.
Aquatic insects, *36*, 1–8.

Gamboa, M., & Watanabe, K. (2019). Genome-wide signatures of local adaptation among
seven stoneflies species along nationwide latitudinal gradient in japan. *BMC Genomics*, *20*,
84.

Gattoliatt, J-L., Rutschmann, S., Monaghan, M. T., & Sartori, M. (2018). From molecular
hypothesis to valid species: description of three endemic species of Baetis
(Ephemeroptera: Baetidae) from the Canary Islands. *Arthropod Systematics and*
Phylogeny, *76*, 509–528.

Gering, J. C., Crist, T. O., & Veech, J. A. (2003). Additive partitioning of species diversity across
multiple spatial scales: implications for regional conservation of biodiversity. *Conservation*
Biology, *17*, 488–499.

Gill, B. A., Harrington, R. A., Kondratieff, B. C., Zamudio, K. R., Poff, N. L., & Funk, W. C.
(2013). Morphological taxonomy, DNA barcoding, and species diversity in southern Rocky
Mountain headwater streams. *Freshwater Science*, *33*, 288–301.

Guindon, S., & Gascuel, O. (2003). PhyML: A simple, fast and accurate algorithm to estimate
large phylogenies by maximum likelihood. *Systematic Biology*, *52*, 696–704.

Hauer, F. R., & Lamberti, G. A. (1996). *Methods in stream ecology*. Elsevier Science, California,
USA.

Herrera-Vasquez, J. (2008). Community structure of aquatic insects in the Esparza River, Costa
Rica. *Revista de Biologia Tropical*, *57*, 133–139.

Heino, J., Melo A. S., Siquiera, T., Soininen, J., Valanko, S., & Bini, L. M. (2015).
Metacommunity organization, spatial extent and dispersal in aquatic systems: patterns,
processes and prospects. *Freswater Biology*, *60*, 845–869.

Hughes, J. M., Schmidt, D. J., & Finn, D. S. (2009). Genes in streams: using DNA to understand
the movement of freshwater fauna and their riverine habitat. *BioScience*, *59*, 573–583.

Ishida, Y., Abekura, K., & Takemon, Y. (2005). Habitat characteristics of *Rhinogobius* sp. or
"Shimahiregata" in Shirokita wando. *Ecology and Civil Engineering*, *8*, 1–14.

Jackson, J. K., Battle, J. M., White, B. P., Pilgrim, E. M., Stein E. D., Miller, P. E., & Sweeney
B. W. (2013). Cryptic biodiversity in streams: a comparison of macroinvertebrate
communities based on morphological and DNA barcode identifications. *Freshwater*
Science, *33*, 312–324.

Karney CFF. (2013) Algorithms for geodesics. *Journal of Geodesy*. 87, 43–55.

Karaus, U., Larsen, S., Guillong, H., & Tockner, K. (2013). The contribution of lateral aquatic
habitats to insect diversity along river corridors in the Alps. *Landscape Ecology*, *28*, 1755–
1767.

Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., …
Webb, C. O. (2010). Picante: R tools for integrating phylogenies and ecology.
Bioinformatics, *26*, 1463–1464.

Lake, P.S. (2003). Ecological effects of perturbation by drought in flowing waters. *Freshwater*
Biology, *48*, 1161–1172.

Lancaster, J., & Downes, B. J. (2013). *Aquatic entomology*. Oxford University Press, United
Kingdom.

Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among multiple
communities. *Oikos*, *76*, 5–13.

Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H., …
Higgins, D. G. (2007). ClustalW and ClustalX version 2. *Bioinformatics*, *23*, 2947–2948.

Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: testing
multispecies responses in multifactorial ecological experiments. *Ecological Monographs*,
69, 1–24.

Leprieur, F, Albouy, C., De Bortoli, J, Cowman, P.F., Bellwood, D.R., & Mouillot, D. (2012).
Quantifying phylogenetic beta diversity: distinguishing between "true" turnover of lineages
and phylogenetic diversity gradients. *PlosOne*, *7*, e42760.

Librado, P., & Rozas, J. (2009). DnaSp v5: A software for comprehensive analysis of DNA
polymorphism data. *Bioinformatics*, *25*, 1451-1452.

Lososova, Z., Smarda, P., Chytry, M., Purschke, O., Pysek, P., Sadlo, L., … Winter, M. (2015).
Phylogenetic structure of plant species pools reflects habitats age on the geological time
scale. *Journal of Vegetation Science*, *24*, 820–833.

Lu, H.-P., Wagner, H.H., & Chen, X.-Y. (2007) A contribution diversity approach to evaluate
species diversity. *Basic and Applied Ecology*, *8*, 1–12.

Marten, A., Brandle, M., & Brandl, R. (2006). Habitat type predicts genetic population
differentiation in freshwater invertebrates. *Molecular Ecology*, *15*, 2643–2651.

Martin, C. A., Cain, D. J., Luoma, S. N., & Buchwalter, D. B. (2007). Cadmium ecophysiology
in seven stonefly (Plecoptera) species: Delineating sources and susceptibility.
Environmental Science & Technology, *41*, 7171–7177.

Mayfield, M. M., & Levine, J. M. (2010). Opposing effects of competitive exclusion on the
phylogenetic structure of communities. *Ecology Letters*, *13*, 1085–1093.

McLain, D. K., Wesson, D. M., Oliver, J. H., & Collins, F. H. (1995). Variation in ribosomal DNA
internal transcribed spacers 1 among eastern populations of *Ixodes scapularis* (Acari:
Ixodidae). *Journal of Medical Entomology*, *32*, 353–360.

Macneale, K. H., Peckarsky, B. L., & Likens, G. E., (2005). Stable isotopes identify dispersal
patterns of stonefly populations living along stream corridors. *Freshwater Biology*, *50*,
1117–1130.

Misof, B., Shanlin, L., Meusemann, K., Peters, R. S., Donath, A., Mayer C., … Zhou, X. (2014).
Phylogenomics resolves the timing and pattern of insect evolution. *Science*, *346*, 763–767.

Murria, C., Bonada, N., Arnedo, M. A., Prat, N., & Vogler, A. P. (2013). Higher β-and γ-diversity
at species and genetic levels in headwaters than in mid-order streams in *Hydropsyche*
(Trichoptera). *Freshwater Biology*, *58*, 2226–2236.

Mykra, H., Heino, J., & Muotka, T. (2007). Scale-related patterns in the spatial and
environmental components of stream macroinvertebrate assemblage variation. *Global*
Ecology and Biogeography, *16*, 149–159.

Mynott, J. H., Webb, J. M., & Suter, P. J. (2011). Adult and larval associations of the alpine
stonefly genus *Riekoperla* McLellan (Plecoptera: Gripopterygidae) using mitochondrial
DNA. *Invertebrate Systematics*, *25*, 11–12.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, R. B., O'Hara, R. B., … Wagner
H. (2012). Vegan. Community Ecology R Package. Retrieved from http://CRAN.R-
project/package-vegan

Pastuchova, Z., Lehotsky, M., & Greeskova, A. (2008). Influence of morphohydraulic habitat
structure on invertebrate communities (Ephemeroptera, Plecoptera and Trichoptera).
Biologia, *63*, 720–729.

Peres-Neto, P.R., Legendre, P., Dray, S. & Borcard, D. (2006). Variation partitioning of species
data matrices: estimation and comparison of fractions. *Ecology*, 87, 2614–2625.

Pons, J., Barraclough, T. G., Gomez-Zurita, J., Cardoso, A., Duran, D. P., Hazell S., … Vogler,
A. P. (2006). Sequence-based species delimitation for the DNA taxonomy of undescribed
insects. *Systematic Biology*, *55*, 595–609.

Posada, D. (2008). jModelTest: Phylogenetic Model Averaging. *Molecular Biology and*
Evolution, *25*, 1253–1256.

R Core Team (2014). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.

Revell, L. J. (2012). Phytools: An R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution*, *3*, 217–223.

Rádková, V., Bojková, J., Křoupalová, V., Schenková, J., Syrovátka V., & Horsák M. (2014).
The role of dispersal mode and habitat specialization in metacommunity structuring of
aquatic macroinvertebrates in isolated spring fens. *Freshwater Biology*, *59*, 2256–2267.

Ribera, I., & Vogler, A. P. (2008). Habitat type as a determinant of species range sizes: the
example of lotic-lentic differences in aquatic Coleoptera. *Biological Journal of the Linnean*
Society, *71*, 33–52.

Rolls, R. J., Leigh, C., & Sheldon, F. (2012). Mechanistic effects of low-flow hydrology on
riverine ecosystems: ecological principles and consequences of alteration. *Freshwater*
Science, *31*, 1163–1186.

Rutschmann S., Detering, H., Simon S., Funk, D. H., Gattolliat J.-L., Hughes, S. J., …
Monaghan, M. T. (2016). Colonization and diversification of aquatic insects on three
Macaronesian archipelagos using 59 nuclear loci derived from a draft genome. *Molecular*
Phylogenetics and Evolution, *107*, 27–38.

Sargent, R. D., & Ackerly, D. D. (2008). Plant-pollinator interactions and the assembly of plant
communities. *Trends in Ecology and Evolution*, *23*, 123–130.
Saito, V. S., Soininen, J., Fonseca-Gessner, A. A., & Siqiera, T. (2015a). Dispersal traits drive
the phylogenetic distance decay of similarity in neotropical stream metacommunities.
Journal of Biogeography, *42*, 2101–2111.
Saito, V. S., Siquiera, T., & Fonseca-Gessner, A. A. (2015b). Should phylogenetic and
functional diversity metrics compose macroinvertebrate multimetric indices for stream
biomonitoring? *Hydrobiologia*, *745*, 167–179.
Saito, V. S., Cianciaruso, M. V., Siqueira, T., Fonseca-Gessner, A. A., & Povoine, S. (2016).
Phylogenies and traits provide distinct insights about the historical and contemporary
assembly of aquatic insects communities. *Ecology and Evolution*, *6*, 2925–2937.
Serrana, J., Miyake, Y., Gamboa, M., & Watanabe, K. (2019). Comparison of DNA
metabarcoding and morphological identification for stream macroinvertebrate biodiversity
assessment and monitoring. *Ecological Indicators*, *101*, 963–972.
Socolar, J. B., Gilroy, J. J., Kunin, W. E., & Edwards, D. P. (2016). How should Beta-diversity
inform biodiversity conservation? *Trends in Ecology & Evolution*, *31*, 67–80.
Sobral, F. L., & Cianciaruso, M. V. (2016). Functional and phylogenetic structure of forest and
savanna bird assemblages across spatial scales. *Ecography*, *39*, 533–541.
Stewart, K. W., & Stark, B. P. (2008). Plecoptera. In *An introduction to the aquatic insects of
North America* (Eds. Merritt, R. W., Cummins, K. W., & Berg, M. B.), pp. 311–384. Iowa:
Kendall/Hunt Publishing Co.
Thompson, R., & Townsend, C. (2006). A truce with neutral theory: local deterministic factors,
species traits and dispersal limitation together determine patterns of diversity in stream
invertebrates. *Journal of Animal Ecology*, *75*, 476–484.
Tiziano, B., Fenoglio, S., Lopez-Rodriguez, M. J., Tierno de Figueroa, J. M., Grenna, M., &
Cucco, M. (2010). Do predators condition the distribution of prey within micro habitats? An
experiment with stoneflies (Plecoptera). *Journal Review of Hydrobiology*, *95*, 285–295.
Tockner, K., & Stanford, J. A. (2002). Riverine flood plains: present state and future trends.
Environmental Conservation, *29*, 308–330.
Tockner, K., Ward, J. V., Arscott, D. B., Edwards, P. J., Kollmann, J., Gurnell, A. M., … Maiolini,
B. (2013). The Tagliamento River: A model ecosystem of European importance. *Aquatic
Science*, *65*, 239–253.
de Vienne, D. M., Giraud, T., & Martin, O. C. (2007). A Congruence Index for Testing
Topological Similarity between Trees. *Bioinformatics*, *23*, 3119–3124.
Violle C., Nemergut, D. R., Pu, Z., & Jiang, L. (2011). Phylogenetic limiting similarity and
competitive exclusion. *Ecology Letters*, *14*, 782–787.
Viteck, S., Vincon, G., Graf, W., & Pauls, S. U. (2017). High cryptic diversity in aquatic insects:
an integrative approach to study the enigmatic *Leuctra inermis* species group (Plecoptera).
Arthropod Systematics & Phylogeny, *75*, 497–521.

737 Ward, J. V., Tockner, K., Edwards ,P. J., Kollmann, J., Bretschko, G., Gurnell, A. M., …

738   Rossaro, B. (1999). A reference system for the Alps: the 'Fiume Tagliamento'. *River*

739   *Research and Applications*, *15*, 63–75.

740 Webb, C. O., Ackerly, D. D., McPeek, M. A., & Donoghue, M. J. (2002). Phylogenies and

741   community ecology. *Annual Review of Ecology and Systematics*, *33*, 475–505.

742 Wiens, J. J., Ackerly, D. D., Allen, A. P., Anacker, B. L., Buckley, L. B., Cornell, H. V., …

743   Stephens, P. R. (2010). Niche conservatism as an emerging principle in ecology and

744   conservation biology. *Ecology Letters*, *13*, 1310–1324.

745 Xiu, J., Chen, Y., Zhang, L., Chai, Y., Wang, M., Guo Y., … Yue, M. (2017). Using phylogeny

746   and functional traits for assessing community assembly along environmental gradients: a

747   deterministic process driven by elevation. *Ecology and Evolution*, *7*, 5056–5069.

748 Young, M. K., Smith, R. J., Pilgrim, K. L., Fairchlid, M. P., & Schwartz, M. K. (2019). Integrative

749   taxonomy refutes a species hypothesis: The asymmetric hybrid origin of *Arsapnia arapahoe*

750   (Plecoptera, Capniidae). *Ecology and Evolution*, *9*, 1364–1377.

751 **TABLE 1** Sampling site characteristics, along with morphological and DNA-species richness in

752 the Tagliamento (sites T) and Fella (sites F) rivers.

Reaches	Sampling site	Altitude (m)	Habitat Simpsons diversity	Morpho-species	DNA species	Haplotypes
Headwater	T01	1174	0.77	2	3	3
	T02	1064	0.77	3	4	6
	T04	1556	0.76	3	5	5
	T06	1346	0.78	8	10	10
	F01	1042	0.80	4	9	9
	F03	978	0.80	3	4	15
Meandering	T05	486	0.76	6	10	12
	T08	625	0.81	5	15	12
	T13	183	0.80	5	10	13
	T15	5	0.81	2	5	2
	F02	769	0.80	4	10	13
	F04	488	0.76	2	2	2
Bar-braided (floodplain)	T03	661	0.81	6	10	13
	T07	524	0.78	3	6	10
	T09	393	0.76	3	11	11
	T10	287	0.78	5	8	13
	T12	131	0.84	5	9	16
	F05	298	0.82	1	1	1
Spring-fed	T11	170	0.55	4	11	15
	T14	4	0.55	1	1	3

755 **TABLE 2** Phylogenetic signal of stonefly communities in the seven habitats sampled in this
756 study. Bold values indicate statistical significance (p<0.05). Positive values indicate
757 phylogenetic clustering and negative values indicate overdispersion. N = total number of DNA-
758 species per habitat, K = Blomberg's K, PIC = phylogenetic independent contrasts, NRI = net
759 relatedness index, NTI = nearest taxon index

	N	K	PIC	NRI	NTI
Waterfall	21	2.30	2.41	-1.29	-0.79
Riffle	31	2.30	41.73	1.52	**-1.85**
Glide	25	2.71	26.47	0.11	-0.87
Pool	10	**4.03**	**76.80**	**2.53**	**1.79**
Wando	6	1.73	10.89	**-6.44**	-1.26
Side channel	4	1.25	64.83	**-2.06**	-1.06
Isolated Pond	17	2.09	16.59	0.15	-0.82

762   Figure caption

763   **FIGURE 1** Habitats types sampled at each sampling site (Fig. 2) along the Tagliamento and
764   Fella rivers. The arrows indicate the fow direction
765
766   **FIGURE 2** Sampling sites along the Tagliamento and Fella rivers. The rivers are located in the
767   northeast region of Italy and sampling sites were coded using T1-15 (Tagliamento) and F1-05
768   (Fella)
769
770   **FIGURE 3** Additive diversity partition of species richness and Shannon-Wiener diversity (based
771   on DNA-species) into observed and expected hierarchical components of reaches (light grey)
772   and habitats (dark grey). Both partition analyses were significant (<0.001) based on 10,000
773   randomizations. Note that the plots have different scales
774
775   **FIGURE 4** Distance based-redundancy analysis (dbRDA) of stonefly DNA-species among 20
776   sampling sites displaying the influence of habitats on community structure. T = Tagliamento
777   river, F = Fella river. Three habitats (pool, glide, and riffle) explained the highest proportion of
778   the total variation
779
780   **FIGURE 5** Beta diversity of stonefly communities in each of the seven habitats, based on the
781   distance of the values for each habitat to their centroid (community homogeneity in a given
782   habitat, see methods section 2.3.3). The bold line represents the median, while the bottom and
783   top of the box represent lower and upper quartiles, respectively

787
788

   Supplementary Material

   TABLE S1 DNA-species list for all sampling sites

Morphospecies	Total n	DNA-species	Haplotypes	Present sampling sites	Present habitats	Genetic Diversity (%)	Altitude range (m)
Systellognatha							
Perlodidae							
Isoperla cf. *andreinii*	2	1	1	1	1	0	1346
Isoperla grammatica	15	2	9	5	4	2	486-1346
Besdolus sp.	28	3	14	7	5	3.5	183-1556
Perlodes microcephalus	16	2	10	6	2	2	287-769
Perlodes intrincatus	4	1	3	3	2	0	287-625
Perlidae							
Dinocras cephalotes	13	2	8	4	2	2.6	4-183
Perla marginata	24	2	15	6	5	6.5	131-661
Perla grandis	4	2	3	2	1	3	183-625
Chloroperlidae							
Chloroperla susemicheli	13	5	9	2	5	9.5	287-661
Euholognatha							
Capniidae							
Capnia nigra	17	3	9	2	2	7	131-170
Capnia vidua	8	1	2	1	2	0	170
Nemouroidae							
Nemoura mortini	8	3	5	4	4	8.2	486-1346
Nemoura cinerea	10	1	2	1	2	0	1346
Amphinemoura sp.	2	1	1	1	1	0	524
Protonemoura cf. nimborum	3	2	3	3	4	5.2	393-1174
Protonemoura nimborella	7	1	1	1	1	0	1556
Protonemoura nitida	6	2	4	1	2	1.3	1346
Leuctricidae							
Leuctra fusca	25	7	12	6	3	3.7	131-287
Leuctra braueri	10	2	6	2	3	4.2	978-1346
Leuctra major	91	8	64	18	7	13.1	5-1556
Taenioterygidae							
Brachyptera risi	6	1	3	1	2	0	131

799 **TABLE S2** DNA-species richness among habitats per sampling site

Reaches	Sampling sites	Waterfall	Riffle	Glide	Pool	Wando	Side-channel	Isolated Pond	Shannon diversity	Simpson diversity
Headwater	T01	0	0	3	0	0	0	0	0.048602	7.68E-05
	T02	4	2	1	0	0	0	0	0.092222	0.000538
	T04	3	3	1	0	0	0	3	0.119007	0.001152
	T06	4	10	4	0	0	1	2	0.19427	0.005376
	F01	13	7	1	0	2	0	3	0.220693	0.008321
	F03	0	5	7	0	0	0	0	0.134995	0.00169
Meandering	T05	3	4	2	3	0	0	4	0.163554	0.003072
	T08	1	17	1	0	6	0	2	0.225542	0.008986
	T13	5	6	8	0	2	1	3	0.215707	0.00768
	T15	0	0	5	1	0	0	0	0.082351	0.000384
	F02	1	11	1	0	3	0	1	0.170096	0.003482
	F04	0	1	0	1	0	0	0	0.035297	2.56E-05
Bar-braided (floodplain)	T03	6	8	1	2	0	1	3	0.19427	0.005376
	T07	0	2	4	1	1	0	1	0.110493	0.000922
	T09	2	7	7	0	0	0	0	0.163554	0.003072
	T10	2	4	4	2	0	0	6	0.176427	0.003917
	T12	1	6	8	3	0	1	1	0.188504	0.004864
	F05	0	0	0	0	0	2	0	0.035297	2.56E-05
Spring-fed	T11	2	5	4	8	0	0	0	0.18256	0.004378
	T14	0	1	0	2	0	0	0	0.048602	7.68E-05
	Total	47	99	62	23	14	6	29	2.802043	0.936585

List of supplementary figures

FIGURE S1 Maximum likelihood phylogenetic trees of stonefly species located along

theTagliamento and Fella rivers. The *cox1* (mtDNA) tree is shown on the left and ITS (nDNA)

on the right. The numbers at each node indicate the likelihood bootstrap support values.

Asterisks show the adult specimen position. A subdivision in two morphological groups was

observed, Euholognatha and Systellogna

COI

Protonemoura nitida
Protonemoura nimborella
Protonemoura cf. *nimborum*
Amphinemoura sp.
Nemoura cinerea
Nemoura mortini
Brachyptera risi
Capnia vidua
Capnia nigra

Leuctra braueri
Leuctra fusca
Leuctra major

Euholognatha

Chloroperla susemicheli
Besdolus sp.
Perlodes intrincatus
Perlodes microcephalus
Isoperla cf. *andreinii*
Isoperla grammatica
Perla marginata
Perla grandis
Dinocras cephalotes

Systellognatha

Outgroup

ITS

Protonemoura nitida
Protonemoura nimborella
Protonemoura cf. *nimborum*
Amphinemoura sp.
Brachyptera risi   *Nemoura cinerea*
Nemoura mortini
Capnia vidua
Capnia nigra

Leuctra braueri
Leuctra fusca
Leuctra major

Chloroperla susemicheli
Besdolus sp.
Perlodes intrincatus
Perlodes microcephalus
Isoperla cf. *andreinii*
Isoperla grammatica
Perla marginata
Perla grandis
Dinocras cephalotes

Outgroup

809