# Systematic Pan-Cancer Analysis Reveals the Functional Roles of Ferroptosis Across Cancers

Zekun Liu[1, #], Qi Zhao[1, 2, #], Zhi-Xiang Zuo[1, #], Shu-Qiang Yuan[1, #], Kai Yu[1], Qingfeng Zhang[1],

Xiaolong Zhang[1], Hui Sheng[1], Han Cheng[3], Feng Wang[1], Rui-Hua Xu[1, 2, *], Ze-Xian Liu[1, 2, *]

[1] State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou 510060, China.

[2] Precision Diagnosis and Treatment for Gastrointestinal Cancer, Chinese Academy of Medical Sciences，Guangzhou 510060，China

[3] School of Life Sciences, Zhengzhou University, Zhengzhou 450001, China

[#]These authors contributed equally to this work.

[*]To whom correspondence should be addressed:

Dr. Ze-Xian Liu: liuzx@sysucc.org.cn

Dr. Rui-Hua Xu: xurh@sysucc.org.cn

## Abstract

Ferroptosis is a type of cell death that related to cancer, however, the characteristics of ferroptosis in cancers are still uncertain. Based on the data in The Cancer Genome Atlas, we found that most ferroptosis regulator genes (FRGs) were differentially expressed in tumors, copy number alterations (CNA) and DNA methylation contributed to their aberrant expression. We established the ferroptosis potential index (FPI) to reveal the functional roles of ferroptosis and noticed that the FPI was higher in tumors than in normal tissues in most cancers, and was associated with subtypes and clinical features. The FPI was negatively correlated with several metabolism pathways but positively associated with several important metastasis-related pathways and immune-related pathways. Higher FPI predicted worse prognosis in several tumors, while FPI and FRGs impacted drug sensitivity. Our study presents a systematical analysis of ferroptosis and its regulatory genes, and highlights the potential of ferroptosis-based cancer therapy.

## Keywords

ferroptosis; pan-cancer analysis; phenotype; survival; drug sensitivity

## Background

As a newly discovered type of programmed cell death, ferroptosis results from the accumulation of iron-dependent lipid hydroperoxides and leads to cytological changes, for which the features and mechanisms are different compared to typical cell death such as apoptosis [1]. Previous studies demonstrated that during the process of ferroptosis, sufficient and approachable cellular iron was required [1]. Inhibition of system $X_C^-$, which is a membrane $Na^+$-dependent cysteine-glutamate exchange transporter, and GPX4 can disrupt the oxidation-reduction balance and cause overwhelming lipid peroxidation that ultimately results in cell death [2-4]. The initiation and execution of ferroptosis are affected by multiple factors including amino acids, lipids, and iron metabolism, and regulated by various signaling pathways such as amino acid and glutathione metabolism, lipid metabolism, iron metabolism, and mevalonate pathway [2].

As the understanding of ferroptosis has increased, its complex biological function has been revealed [5, 6]. Furthermore, ferroptosis has been found to be closely related with various human diseases including periventricular leukomalacia, Huntington's disease and acute kidney injury [7-10]. Besides, the literatures have confirmed that ferroptosis could suppress tumor growth and kill tumor cells [3], and then plays an important role in cancers such as renal cell carcinomas and liver cancer [4, 11]. For example, erastin-induced ferroptosis decreases the growth of tumors formed from human CRC cells [12]. Ductal pancreatic cancer cells with a mutant *KRAS* gene were more susceptible to ferroptosis than wild type [3, 13]. Conversely, inducing tumor cell ferroptosis by small molecules has become an important strategy for the treatment of many tumors, such as hepatocellular carcinoma, kidney cancer, and pancreatic cancer [4, 13-15]. In addition, changes in the gene expression of tumor cells also affect ferroptosis, and a number of genes have been confirmed to regulate ferroptosis. For example, *ACSL4* participates in the biosynthesis and remodeling of polyunsaturated fatty acid-Pes, and the downregulation of *ACSL4* increases the resistance to ferroptosis [16, 17]. Furthermore, ferroptosis and its regulator genes (FRGs) were identified to be correlated with drug resistance [18]. For example, it was previously reported that liver cancer cells repressed ferroptosis by regulating the expression of *NRF2* or *MT-1G*, which promoted sorafenib resistance *in vitro* and in tumor xenograft models [11, 19]. Recently, Wang *et al.* reported that CD8+ T cells could promote tumor ferroptosis during cancer immunotherapy treatment [20]. Thus, ferroptosis might play important roles during cancer progression and treatment, and a systematic study for ferroptosis and its aberrances across

cancers will be helpful.

In the present study, for the first time, we performed a comprehensive analysis of genomic variations and expression profiles of the FRGs across 20 cancer types. Furthermore, we computationally modelled the ferroptosis level based on FRGs expression, and dissected the relations between ferroptosis and cancer clinical features. It was found that ferroptosis was associated with various cancer hallmarks, immune microenvironment, drug resistance and patient survival. These results highlighted the critical roles of ferroptosis in cancer and should be helpful for further investigations of ferroptosis-related molecular mechanisms and therapy development.

## Materials and Methods

### Datasets and Source

The mRNA expression data, copy number alteration thresholded data, masked copy number segmentation data, and DNA methylation 450K data of twenty cancers, including bladder urothelial carcinoma (BLCA), breast invasive carcinoma (BRCA), cholangiocarcinoma (CHOL), colon adenocarcinoma (COAD), endocervical cancers (CESC), esophageal carcinoma (ESCA), glioblastoma multiforme (GBM), head and neck squamous cell carcinoma (HNSC), kidney chromophobe (KICH), kidney renal clear cell carcinoma (KIRC), kidney renal papillary cell carcinoma (KIRP), liver hepatocellular carcinoma (LIHC), lung adenocarcinoma (LUAD), lung squamous cell carcinoma (LUSC), pancreatic adenocarcinoma (PAAD), prostate adenocarcinoma (PRAD), rectum adenocarcinoma (READ), stomach adenocarcinoma (STAD), thyroid carcinoma (THCA), uterine corpus endometrial carcinoma (UCEC), which had both tumors and normal samples were downloaded from Firehose (http://gdac.broadinstitute.org). Mutation data, miRNA-seq data, and clinical data were downloaded from the Xena Browser (https://xenabrowser.net/datapages/). Additional gene-centric RMA-normalized gene expression profiles and drug response data of over 1000 cancer cell lines were accessed from the Genomics of Drug Sensitivity in Cancer (GDSC) database (https://www.cancerrxgene.org/downloads) [21]. Immune associated data, including immune cell type fractions and immunophenoscore were obtained from TCIA (https://tcia.at/home) [22]. Integrated protein-protein interaction data was obtained from the Human Protein Reference Database (http://www.hprd.org/) and BioGRID (https://thebiogrid.org) [23, 24]. We obtained clinically actionable genes (CAGs) from a previous study [25] (https://software.broadinstitute.org/cancer/cga/target).

### Differential expression analysis of mRNA

To test genes differentially expressed between tumor and normal tissue, gene expression data for 20502 genes across 20 cancer types were downloaded from TCGA at FireBrowse (http://gdac.broadinstitute.org, 2016 January). Then, the fold change and adjusted $P$-value were calculated by the edgeR package [26]. We defined genes with an adjusted $P$-value less than 0.05 as the differential expression genes (DEGs).

**Establishing the Ferroptosis Potential Index Model**

The enrichment score (ES) of gene set that positively or negatively regulated ferroptosis was calculated using single sample gene set enrichment analysis (ssGSEA) in the R package 'GSVA' [27], and the normalized differences between the ES of the positive regulation gene set minus negative regulation gene set was defined as the ferroptosis potential index (FPI) to computationally dissect the ferroptosis levels/trends of the tissue samples.

**Somatic Copy-number Alteration (SCNA) and Mutation Analysis**

The heterozygosity and homozygosity of amplification and deletion were included to elevate the copy-number alteration of each gene, in which over five percent was regarded as high-frequency SCNA. Pearson's correlation between expression values and copy number segment values of each gene was calculated to evaluate the association between CNA and expression. The R package "DISCOVER" was employed to evaluate mutual exclusivity between FRGs and tumor suppressor genes or oncogenes across tumor samples in each cancer type [28]. The mutation and CNA events were integrated, while only homozygous amplification and deletion were included, and only protein-coding mutations were retained. For each cancer type, the genes were considered to be mutually exclusive if they had a q value of 0.05.

**DNA methylation analysis**

The R package "IlluminaHumanMethylation-450kanno.ilmn12.hg19" from Bioconductor was imported to annotate the methylation probe for the promoter of each gene. Differential methylation of each gene in tumor and normal samples was tested by the Wilcoxon signed rank test, and genes that were significantly hypomethylated or hypermethylated were identified using a *P*-value cutoff of 0.05. Pearson's correlation between the transcriptional expression of FRGs and the Beta value of the promoter DNA methylation were calculated and considered significant if the *P*-value < 0.05.

**miRNA expression analysis**

To investigate the mechanisms of dysregulation for ferroptosis regulator genes in cancer, we searched potential miRNAs which might regulate the FRGs based on miRNA-target intersections in starBase. The Spearman correlation between the expression of miRNA and FRGs was statistically evaluated

(adjusted *P*-value < 0.1, rho < -0.1). Cytoscape software was used to visualize the high-frequency interaction networks among FRGs and miRNA [29].

## Multivariate Regression Analysis of Gene Expression

To assess which factors had significant effects on FRG expression, the expression of each FRG was modeled by linear regression as a function of the median miRNA expression, the median Beta value of promoter methylation, and the copy number of the genes.

## Clinical Features Analysis

The R package "survival" was used to assess the prognosis potential of the FRGs and ferroptosis potential index among cancers. For survival analysis, the expression threshold was exhaustively tested and the one with most significant *P*-value was considered the best cut-off. To test the association between ferroptosis level and clinical features, the Pearson correlation was calculated between FPI and tumor stages, age, body mass index (BMI) and cigarette exposure per day, which were converted to numeric variables ("stage1" = 1, "stage2" = 2, etc.). Wilcoxon rank sum tests and Tukey's tests were used to determine the impact on the ferroptosis potential index for other clinical characteristics including race, remission status, and alcohol. The influence of different MSI statuses, histologic types, and molecular subtypes on FPI were also considered.

## Immune Features Analysis

To study the relationship of ferroptosis and immune microenvironments, we computed the Pearson correlation between FPI and immune parameters including immune cell fractions and 5 types of immunophenoscores.

## Identifying the FPI associated significant driver gene mutation

A total of 375 driver genes identified in previous pan-cancer research were included for analysis[30]. To test whether a driver gene's mutational status was significantly associated with ferroptosis among cancers, the rank-transformed FPI was modeled by linear regression as a function of the driver gene's mutational status, ignoring the synonymous variant. To diminish the confounding effects, the rank-transformed count of the total nonsynonymous mutations and the tumor type, which were encoded

as virtual variables, were included. To further characterize each cancer, the Benjamini-Hochberg method was used to correct the *P* values across 375 genes. Genes with an adjusted *P* value less than 0.05 for mutation status variable were significantly associated with FPI.

**Gene Set Enrichment Analysis**

To identify the pathways associated with ferroptosis, the samples of each tumor type were divided into two groups according to the FPI, consisting of the top 30% and bottom 30%. Then, the gene set enrichment analysis (GSEA) was performed [31].

**Correlation between Drug Sensitivity and FPI/FRG Expression**

To test the correlation between small molecular drugs and FPI and FRGs, the Pearson correlation coefficients for FPI, the expression value of FRGs and the area under the dose-response curve (AUCs) values were calculated, the results with |R| > 0.1 and *P*-value < 0.05 were considered as significantly correlated. To further investigate the influence of drugs on ferroptosis, the significant association between the expression of clinically actionable genes and FRGs were finished across cancer cell lines, and the associations were filtered with PPIs. Then the drugs which target CAGs were selected according to DrugBank [32] (https://www.drugbank.ca).

## Results

### Genetic alterations of ferroptosis regulator genes (FRGs) in cancers

In this study, we defined the 24 genes including *CDKN1A*, *HSPA5*, *TTC35*, *SLC7A11*, *NFE2L2*, *MT1G*, *HSPB1*, *GPX4*, *FANCD2*, *CISD1*, *FDFT1*, *SLC1A5*, *SAT1*, *TFRC*, *RPL8*, *NCOA4*, *LPCAT3*, *GLS2*, *DPP4*, *CS*, *CARS*, *ATP5G3*, *ALOX15*, and *ACSL4*, as ferroptosis regulator genes (FRGs), which were identified to play critical roles in regulating ferroptosis by previous studies [2]. To determine the patterns for dysregulation of FRGs in cancer, we examined the genomic data, including genetic variation, somatic copy number alternation (SCNA), mRNA expression and DNA methylation data of tumor and normal tissues from 20 cancer types. The analysis of nonsynonymous mutations in 20 cancer types showed that the mutation frequencies for FRGs were generally low (Figure S1A), while 13 FRGs in UCEC had mutation frequencies greater than 5%. Furthermore, *NFE2L2*, which encodes NRF2 and plays a master regulator role in anti-oxidant responses and has been shown to cause resistance to ferroptosis [11], showed relatively high mutation frequencies in multiple cancers compared to other FRGs.

To further dissect the genetic aberrance of FRGs in cancer, the percentage of SCNA was examined and the results showed that in general SCNA occurred at high frequencies (with over five percent of all samples) in most cancer types (Figure 1A), but all FRGs in THCA showed a low frequency of SCNA. It was observed that FRGs presented diverse SCNA profiles. For example, *TTC35*, *HSPB1*, *TFRC*, and *RPL8* were more prone to copy number gain than copy number loss in almost all tumors, but *SCL7A11* and *ALOX15* showed the opposite profile. Furthermore, we analyzed the co-occurrence of mutations and SCNA between FRGs and oncogenes/tumor suppressor genes (TSGs) previously identified by Bailey *et al.* [33]. The significance was calculated via the mutual exclusivity test by DISCOVER [28], with a false discovery rate of 1%. A total of 28 oncogenes (Figure S1B) and 43 TSGs (Figure S1C) were found to be altered with FRGs in the mutually exclusive manner in certain cancer types, and *FDFT1*, *RPL8*, *TFRC*, and *NFE2L2* were frequently exclusive to oncogenes or TSGs (Figure S1B-C).

### Aberrant expression of FRGs among cancer

Besides genetic alternations, differential expression analysis was performed to investigate alteration of gene expression patterns of FRGs, and the numbers of tumor and normal sample were

showed in Table 1. We found that all FRGs were differentially expressed in at least one cancer type. Several FRGs showed consistent expression patterns in cross-cancer analysis. *SLC7A11*, *FANCD2*, *CARS*, *SLC1A5* and *RPL8* were significantly upregulated in 15, 18, 12, 15, 14 types of cancers, respectively, while *NCOA4* was downregulated in 15 cancers (Figure 1B). Additionally, several FRGs showed miscellaneous cancer type specificities which has not been well characterized previously. For example, *HSPA5* was upregulated in most cancer types including breast invasive carcinoma (BRCA) (fold change [FC] = 1.87, adjust *P*-value = 1.42 ×10$^{-42}$) and LUAD (FC = 1.90, adjust *P*-value = 2.45 ×10$^{-29}$) but was significantly downregulated in THCA (FC = 0.68, adjust *P*-value = 2.67 ×10$^{-7}$). *DPP4*, which appears to play a suppressor role in the development of cancer [34-36], showed significant upregulation in KIRC, KIRP, ESCA, LUAD, GBM, and THCA, but was downregulated in CHOL, KICH, BRCA, LUSC, HNSC, and STAD. We also noticed that *DPP4* had opposed expression profiles in different subtypes of tumors in the lung and kidney. This demonstrated that FRGs might play different roles in different cancers.

Since SCNA in tumors plays a critical role in regulating gene expression, we evaluate the effects of SCNA on gene expression of FRGs. The Pearson correlation between gene expression and the copy number from masked copy number segment of TCGA was examined. The results showed that the expression of most FRGs was obviously correlated to SCNA in most tumors (Figure 1C). For example, the expression of citrate synthase (*CS*), which participates in oxidative metabolism, was significantly associated with SCNA in all cancer. This result indicates that the aberrance of copy number for FRGs is common in most cancers and can have an influence on gene expression.

Besides SCNA, the methylation of promoter can regulate gene expression, and aberrant DNA methylation of the promoter is associated with tumorigenesis [37]. We observed that FRGs showed complex methylation patterns in the 20 cancer types (Figure 1D), only *CDKN1A* showed hypomethylation in 11 tumors consistently. For example, we observed that *DPP4* showed hypermethylation in seven cancer types and hypomethylation in four cancer types. Although there were differences in the pattern of methylation for FRGs, a negative relationship was observed between gene expression and DNA methylation overall (Figure 1E). This result demonstrated that promoter DNA methylation may regulate the expression of FRGs in tumors.

Besides SCNA and DNA methylation, microRNAs (miRNAs) can regulate gene expression and become involved in cancer development [38]. To determine which miRNAs are involved in regulating

ferroptosis, we performed the analysis to reveal the network of miRNA-FRGs. The starBase database was used to infer miRNA that potentially targeting FRGs, and those miRNAs that were significantly negatively correlated with gene expression were thought to be involved in the regulation of FRGs [39]. As the network showed (Figure 1F), all of the interactions between miRNA and FRGs occurred in more than six cancers. It is obvious that FRGs could be targeted by miRNA with high frequency, including *ACSL4*, *NCOA4*, and *CDKN1A* (Figure 1F). To further understand the aberrances of the miRNA-FRG network in tumors, we conducted differential expression analysis of miRNA and counted the expression aberrances of miRNA across cancer types (Figure S1D). It was found that several miRNAs had consistent expression trends in tumors, for example, has-miR-93 targeted *CDKN1A* and was downregulated in 11 cancers. On the other hand, miRNA had different expression trends in different tumors. For example, has-miR-375 targeted *ACSL4* in five tumors, but it showed upregulation in two cancers and downregulation in three cancers. This suggests that miRNA play a regulatory role in FRGs expression, and the aberrant expression of miRNA in the tumors could impact ferroptosis.

Since SCNA, DNA methylation, and miRNA all can regulate FRGs expression, but the contribution of a single factor on gene expression is not clear. The linear regression approach was applied to analyze the contribution of each factor and diminish the confounding effects. As the results showed (Figure S1E), SCNA was positively correlated with gene expression, while methylation and miRNA showed a negative correlation. We further observed multiple regulated patterns in FRGs, and the expression of several FRGs was only related to a single factor in several tumors, while others were related to multiple factors. For example, *ACSL4* expression was only associated with miRNA in seven cancers, including BLCA, CESC, KIRC, KIRP, LUAD, PRAD, and UCEC. SCNA played the only significant regulator role for *CARS* in CHOL, ESCA and colorectal cancer. However, the expression of *NCOA4* and *NFE2L2* was significantly regulated by three factors in seven and ten tumors, respectively. Thus, all the FRGs show diverse regulation patterns in different cancers. This suggests that the expression regulation patterns of all FRGs were tumor-specific.

**Computational modeling the ferroptosis level among cancers**

To further understand the role of ferroptosis in tumorigenesis and investigate the factors or biological processes associated with ferroptosis, the ferroptosis potential index (FPI) was modeled

based on the enrichment score (ES) of positive regulatory components calculated by ssGSEA minus that of negative regulatory components. We performed multiple analyses to validate the performance of FPI and test the robustness of this index. First, we collected three independent gene expression datasets of tumor cell lines that were treated with erastin or withaferin A (WA), which were reported as an inducer of ferroptosis [1, 40], or ferrostatin that was identified as an inhibitor of ferroptosis [41]. The FPI was calculated and analyzed for the gene expression datasets in neuroblastoma cells (GSE112384), clear cell carcinoma cells (GSE121689) and liver cancer cells (GSE104462) (Figure S2). The results showed that erastin and WA increased the FPI remarkably in all of this three cell lines, while ferrostatin could slightly decreased the FPI compared to control group (Figure S2A, S2D, S2G). Because the increases of the *CHAC1* and *PTGS2* mRNA and ACSL4 protein were associated with cells undergoing ferroptosis, but these changes were not consistent in all experiments [2], we compared the mRNA expression of these three genes in this three cell lines. There were no significant changes in *PTGS2* and *ACSL4* but *CHAC1* was upregulated after treated with erastin or WA (Figure S2B, S2E, S2H). Then we examined Pearson's correlation between FPI and the mRNA expression value of *CHAC1*, as the results showed (Figure S2C, S2F, S2I) they were significantly highly associated and all of coefficient were over 0.7. Thus, the FPI could be used to represent the potential level of ferroptosis based on the transcriptome data.

Based on the computed marker of ferroptosis, we compared the differences of FPI between tumor and normal tissues with the cancer genome atlas (TCGA) data (Figure 2A). Of note, significant differences were found in most cancers such as lung cancer and gastrointestinal cancer, and higher FPI was observed in most tumors except for BRCA and UCEC. We also noticed that the FPI of normal tissues was higher than tumors in female cancers (BRCA, UCEC, and CESC). To further dissect the factors involved in these different FPI patterns, we examined the FPI for different subtypes of BRCA. The results presented in Figure 2B showed that all the estrogen receptor (ER)-positive, progesterone receptor (PR) positive and HER2 positive patients had lower FPI than the negative patients, respectively. Furthermore, triple negative breast cancer (TNBC) samples showed higher ferroptosis level than non-TNBC samples (Figure 2C), which was consistent with previous study [42]. Furthermore, among the different subtypes of kidney cancer, the FPI of tumor samples was higher than normal in KIRP and KIRC, while the FPI in KICH was lower (Figure 2A). In addition, to further dissect the variances of ferroptosis level in different histological types, the FPI in the esophagus

squamous cell carcinoma (ESCC) and esophagus adenocarcinoma (ESAD) were compared. The results showed that FPI in ESCC was significantly higher than ESAD and normal samples, while ESAD was higher than normal but not significant (Figure 2D).

Besides the cancer types and subtypes, we analyzed the correlation between the FPI and remarkable molecular features such as microsatellite instability (MSI) and driver gene mutations. The FPI decreased obviously in tumors with high-level of MSI (MSI-H) compared to non-MSI-H tumors in gastrointestinal tumors (Figure 2E). The FPI was remarkably lower in COAD and STAD patients with MSI-H, and slight decrease was also observed in ESCA and READ (Figure 2F). Furthermore, we analyzed the correlation between FPI and 375 driver genes that frequently mutated in tumors based on regression-based analysis. Through controlling for cancer type and tumor mutation burden (TMB), 81 genes were found to be correlated, and most of them were negatively associated with FPI (adj.*p* < 0.1, Figure S3A). Strikingly, somatic mutations, namely *NFE2L2*, *BRAF*, and *TP53*, were positively associated with FPI in pan-cancer associations, but a mutation in *TP53* was negatively correlated with ferroptosis in LUSC. This observation was verified by the comparison of FPI between the *TP53* mutant and wild type groups (Figure S3B), consistent with previous finding that showed that *TP53* played a dual role in regulating ferroptosis [43]. The influences of *KRAS* mutations on FPI were also investigated, and the results showed that *KRAS* mutations had a significantly negative effect on ferroptosis in gastric tumors (Figure S3C), and we observed that *KRAS* mutations positively impacted FPI in hepatocellular carcinoma (Figure S3A, S3C). Furthermore, the differential expression of FRGs between wild type and mutant tumors for *TP53* (Figure S3D) and *KRAS* (Figure S3E) were found to be ubiquitous in most cancers. Taken together, ferroptosis was negatively related with MSI-H in cancer.

**Association between ferroptosis and pathways in cancer**

To further elucidate the related cellular signaling of ferroptosis in cancer, gene set enrichment analysis (GSEA) was performed based on the transcriptome of two tumor groups with the top and bottom thirty percent of FPI. It was observed that metabolism related pathways in KEGG usually enriched in the tumor with lower FPI, and pathways frequently enriched (>6 cancers) were presented in Figure 3A. For example, terpenoid backbone biosynthesis and steroid biosynthesis were enriched in the low FPI group in 19 and 16 cancers, respectively (Figure 3A). Peroxisome, biosynthesis of unsaturated fatty acids, etc. were also significantly correlated with lower FPI in many cancer types

(Figure 3A). Furthermore, the relations between cancer hallmarks and FPI were also analyzed, and the results showed that 15 hallmarks were frequently significantly correlated with FPI (Figure 3B). For example, KRAS signaling, epithelial-mesenchymal transition, IL6 JAK STAT3 signaling, WNT Beta-catenin signaling were enriched in the high FPI group, which indicated that ferroptosis was positively related with these oncogenic pathways (Figure 3B). Also, metabolism related hallmarks were observed to be negatively related with ferroptosis, which was consistent with the pathway analysis (Figure 3B).

Since previous studies showed that ferroptosis was related to the immune response process [5], here we investigated the association between ferroptosis and immune microenvironment in tumor. The results showed that in several cancers FPI was weakly negatively correlated with immunophenoscore (IPS) (Figure 3C), which could predict the response of the immune checkpoint blockade in melanoma tissue [22]. Furthermore, FPI was positively correlated with MHC and effector cells (EC) but negatively correlated with immunomodulators (CP) and suppressor cells (SC) in most cancers (Figure 3C). The associations between ferroptosis and immune cell types were further evaluated in detail, and the results showed that FPI showed a positive correlation with macrophages in most cancers and a negative association with the activated dendritic cells, activated mast cells, plasma cells and T follicular helper cells (Figure S4A). Thus, there might be close connections between ferroptosis and immune microenvironment, and more investigations were needed to reveal the details.

**Clinical relevance of ferroptosis and its regulators**

Since ferroptosis was involved in cellular metabolism and immune microenvironment, it was proposed that ferroptosis and its regulators should be correlated with cancer survival and other clinical characteristics. Based on the clinical data from TCGA, the survival analysis was performed and the results were consistent since it is found that the lower FPI predicted a better survival in five cancers including GBM, KIRC, KIRP, LIHC, and LUAD (Figure 4). We further dissected the factors that might impact FPI including tumor stage, cigarette smoking, age, and gender. As the results shown (Figure S4B), tumor stages were positively correlated with FPI in LIHC, KIRC, KIRP, and COAD. Only the FPI in LUAD tumors showed a positive correlation with cigarettes smoked per day. Age was positively associated with FPI in GBM and PRAD but showed a negative correlation in seven cancers such as

BRCA and LIHC. It was observed that FPI was remarkably different between genders (Figure S4C), FPI for females was higher than males in KIRC, LIHC, LUAD, and LUSC, but lower in STAD. Significant difference in FPI were also observed among different races in BLCA, BRCA and ESCA. Furthermore, we found that the FPI was correlated cancer metastasis, recurrence, primary and follow outcome in several cancers. In addition, we evaluated the association between FPI and other clinical characteristics including alcohol abuse, fatty liver, hemochromatosis, and viral infection (HPV, HBC, HCV, and EBV), while no significant results were observed. These results indicate that ferroptosis might play critical roles in cancer survival, and clinical factors have tumor-specific effects on FPI.

To further dissect the clinical relevance of ferroptosis in cancer, the roles of FRGs in cancer survival were analyzed. The survival analysis showed that all FRGs were associated with overall survival in at least six cancer types (Figure S4D), but the correlations were miscellaneous since most FRGs could both serve as protective and risk factors in different cancers. For example, patients with higher expression of *FANCD2* showed better survival in kidney cancer, LIHC, and LUAD, but showed worse survival in READ, STAD, and BLCA (Figure S4D). However, high expression of *CARS* and *MT1G* showed consistent better survival among cancers including KICH, UCEC, PAAD, and HNSC (Figure S4D). Thus, the functional roles of FRGs in cancer survival should be further explored.

To study the potential clinical implications of ferroptosis, we examined the association between the gene expression of FRGs and 156 clinically actionable genes (CAGs), including 136 genes for targeted therapy and 20 genes for immunotherapy across cancer types [25, 44]. As a FRG, *CDKN1A* regulates iron uptake and *GPX4* abundance [2], was also a CAG targeted by anti-cancer drugs, such as the inorganic compound arsenic trioxide [45]. We found that all 136 CAGs and FRGs had significant co-expression relationships that were significantly different across cancers as shown in Figure S5A. The number of CAGs correlated with FRGs ranged from 45 in LUSC to 132 in THCA (Figure S5B). The FRG-CAG correlation pairs ranged from 69 in LUSC to 581 in THCA (Figure S5C). For example, *GPX4* co-expressed with *MAP2K2* in 20 cancer types, and showed a significantly positive correlation with *TNFRSF4*, which are targeted genes for immunotherapy. *ACSL4* is significantly correlated with 17 of 20 genes targeted for immunotherapy, suggesting that *ACSL4* has a potential effect on cancer immunotherapy (Figure S6A). Furthermore, besides co-expression, the protein-protein interactions among drug-targeted CAGs and FRGs were analyzed, and the results presented in Figure S6B showed that the interactions were obvious. For example, *EGFR*, the target

of lapatinib, gefitinib and afatinib, was positively associated with FRGs including *DPP4* (R = 0.15) and *HSPA5* (R = 0.31), and negatively correlated with *FDFT1* (R = -0.24), while there were protein-protein interactions between EGFR and DPP4, HSPA5, FDFT1 (Figure S6B). These results indicated that ferroptosis might be involved in drug effects by interacting with targeted clinically actionable genes. Taken together, our studies suggest that clinically actionable genes are closely related to FRGs, which highlights the significance of ferroptosis in cancer treatment, including both immunotherapy and targeted therapy.

To further understand the correlation between ferroptosis and drug sensitivity, the area under the percent viability curve (AUC) approach was employed to evaluate drug sensitivity [46] and calculate the correlation between FPI and drug sensitivity across cancer cell lines. Drugs associated with FPI were also tested for their correlation with FRG expression. We identified that FPI was significantly associated with 64 drugs, including 12 that were negatively correlated and 52 that were positively correlated (Figure S6C). To explore the effects of each FRG on drug sensitivity, we also analyzed the associations between the drug sensitivity of 64 cancer drugs and the expression of FRGs, and found 521 significantly correlated pairs (Figure S6C). Among them, the expression of *FANCD2* was correlated with AUC in 57 drugs, while no significant relation was observed for *CISD1* (Figure S6C). Furthermore, the associations classified the FRGs into two group, one group included genes such as *FANCD2* and *ATP5G3*, which were positively associated AUC of docetaxel, trametinib, etc. (Figure S6C). The other group included *SLC7A11*, *SAT1*, *HSPA5*, etc., which were negatively associated (Figure S6C). Taken together, these results suggest that ferroptosis was correlated to the sensitivity of multiple drugs.

## Discussion

Ferroptosis is driven by damage of cell membranes caused by the loss of activity of GPX4 [47]. Researchers identified that ferroptosis is closely related to tumorigenesis and plays an important role in cancer treatment [4, 48]. However, there is a lack of systematic studies on ferroptosis and its regulator genes across cancer types. In this study, we employed multi-omics data and clinical data across 20 cancer types from TCGA and revealed the global alterations of ferroptosis regulator genes at genetic, epigenetic and transcriptional levels. We also used ssGSEA to process expression data to establish FPI to characterize ferroptosis and addressed which genetic and nongenetic factors (including drug and patient phenotype) were related with FPI. Different molecular types affected the ferroptosis in breast cancer and gastrointestinal cancer, which meant that the responses of different molecular subtypes to treatment may be related to ferroptosis. On the other hand, molecular subtypes need to be considered when ferroptosis is applied as a therapeutic strategy.

The mechanism by which ferroptosis regulates tumor cell growth and proliferation is still unclear, but the relationship we observed between FPI and hallmarks of cancer could improve the understanding the role of ferroptosis. The results of GSEA demonstrate that the level of ferroptosis is closely related to tumor-associated hallmarks in most cancers. Ferroptosis genes can play both oncogene and tumor suppressor roles in cancer [43], and the FPI acts as a protective or risk factor across cancer types. We also found that several common clinical factors affected ferroptosis, such as cigarette smoking, BMI and tumor stage. As we know, cigarette smoking is a risk factor for esophageal cancer, lung cancer and kidney cancer [49-51]. In our results, the number of cigarettes per day was positively correlated with the ferroptosis potential index in LUAD, which may be because the cigarette-induced oxidation reaction promoted lipid peroxidation [52-54]. FPI also varied between the sexes in several cancers, including LIHC, LUAD, LUSC, and STAD and between different races in BRCA, BLCA, and ESCA, which implied that gender and race need to be considered when using ferroptosis as a treatment strategy. We also noticed that better clinical outcomes or status in several cancer types also have lower FPI, which further confirmed the dual role of ferroptosis. Thus, a different strategy of regulating the ferroptosis of tumor cells may benefit patients and improve prognosis.

Furthermore, we showed that FRGs were co-expressed with most clinically actionable genes and interacted with genes modulated by drugs, suggesting that studying ferroptosis may improve the strategy for cancer therapy. We observed that the co-expression of FRGs and CAGs could be roughly

divided into two groups, and drugs that target clinical genes could have complex regulatory effects on ferroptosis. Most strikingly, the expression of *GPX4* was associated with *MAP2K2* in 20 cancers, which indicated that *MAP2K2* may play an important role in ferroptosis. To further explore drug sensitivity and ferroptosis, we observed that the FPI could characterize the sensitivity of many drugs, the AUC of many drugs was inversely associated with the FPI in cancer cell lines, which implied that regulating the ferroptosis of tumor cells may improve the therapeutic effect of cancer treatments. However, opposite effects were observed regarding the effects of drugs on ferroptosis. Thus, further detailed studies should be carried out to determine the functions and mechanisms of ferroptosis in different cancers.

## Conclusions

In the present study, we presented a systematical analysis of ferroptosis and its regulator genes across cancers. Most FRGs were aberrantly expressed in tumors among various cancer types, while frequent CNA and differential DNA methylation contributed greatly. We established the FPI to evaluate the ferroptosis level and found that the FPI were higher in tumors than in the adjacent normal tissues in most cancers and were associated with clinical features, and cancer metastasis, recurrence, outcome and drug sensitivity. These findings highlighted the potential of ferroptosis-based cancer therapy.

**List of abbreviations**

FRGs: ferroptosis regulator genes; FPI: ferroptosis potential index; CAN: copy number alterations; BLCA: bladder urothelial carcinoma; BRCA: breast invasive carcinoma; CHOL: cholangiocarcinoma; COAD: colon adenocarcinoma; CESC: endocervical cancers; ESCA: esophageal carcinoma; GBM: glioblastoma multiforme; HNSC: head and neck squamous cell carcinoma; KICH: kidney chromophobe; KIRC: kidney renal clear cell carcinoma; KIRP: kidney renal papillary cell carcinoma; LIHC: liver hepatocellular carcinoma; LUAD: lung adenocarcinoma; LUSC: lung squamous cell carcinoma; PAAD: pancreatic adenocarcinoma; PRAD: prostate adenocarcinoma; READ: rectum adenocarcinoma; STAD: stomach adenocarcinoma; THCA: thyroid carcinoma; UCEC: uterine corpus endometrial carcinoma; CAGs: clinically actionable genes; DEG: differential expression gene; ES: enrichment score; ssGSEA: single sample gene set enrichment analysis; SCNA: Somatic Copy-number Alteration; GSEA: gene set enrichment analysis; MSI: microsatellite instability; TMB: tumor mutation burden; IPS: immunophenoscore; EC: effector cells; CP: immunomodulators; SC: suppressor cells

# Declarations

## Ethics approval and consent to participate

Not applicable

## Consent for publication

Not applicable

## Availability of data and materials

Not applicable

## Competing interests

The authors declare that they have no competing interests

## Funding

This work was supported by grants from National Key R&D Program of China (2018YFC1313300 to RH.X.), Program for Guangdong Introducing Innovative and Entrepreneurial Teams (2017ZT07S096 to ZX.L.), the Natural Science Foundation of China (81972239 to ZX.L.), Pearl River S&T Nova Program of Guangzhou (201906010088 to ZX.L.), CAMS Innovation Fund for Medical Sciences (2019-I2M-5-036 to RH.X.), Natural Science Foundation of Guangdong Province (2017A030313485 to RH.X.); and Science and Technology Program of Guangdong (2019B020227002 to RH.X.).

## Authors' contributions

ZX.L. and RH.X. designed and supervised the experiments. Z.L., Q.Z., ZX.Z., and SQ.Y. performed the data analysis with contributions from K.Y., QF.Z., X.Z., H.S., H.C., and F.W.. ZX.L. and Z.L. wrote the manuscript with contributions of all authors. All authors reviewed the manuscript.

## Acknowledgements

Not applicable

# References

1. Dixon SJ, Lemberg KM, Lamprecht MR, Skouta R, Zaitsev EM, Gleason CE, Patel DN, Bauer AJ, Cantley AM, Yang WS, et al: **Ferroptosis: an iron-dependent form of nonapoptotic cell death.** *Cell* 2012, **149:**1060-1072.

2. Stockwell BR, Friedmann Angeli JP, Bayir H, Bush AI, Conrad M, Dixon SJ, Fulda S, Gascon S, Hatzios SK, Kagan VE, et al: **Ferroptosis: A Regulated Cell Death Nexus Linking Metabolism, Redox Biology, and Disease.** *Cell* 2017, **171:**273-285.

3. Yu H, Guo P, Xie X, Wang Y, Chen G: **Ferroptosis, a new form of cell death, and its relationships with tumourous diseases.** *J Cell Mol Med* 2017, **21:**648-657.

4. Yang WS, SriRamaratnam R, Welsch ME, Shimada K, Skouta R, Viswanathan VS, Cheah JH, Clemons PA, Shamji AF, Clish CB, et al: **Regulation of ferroptotic cancer cell death by GPX4.** *Cell* 2014, **156:**317-331.

5. Matsushita M, Freigang S, Schneider C, Conrad M, Bornkamm GW, Kopf M: **T cell lipid peroxidation induces ferroptosis and prevents immunity to infection.** *J Exp Med* 2015, **212:**555-568.

6. Yang WS, Stockwell BR: **Ferroptosis: Death by Lipid Peroxidation.** *Trends Cell Biol* 2016, **26:**165-176.

7. Skouta R, Dixon SJ, Wang J, Dunn DE, Orman M, Shimada K, Rosenberg PA, Lo DC, Weinberg JM, Linkermann A, Stockwell BR: **Ferrostatins inhibit oxidative lipid damage and cell death in diverse disease models.** *J Am Chem Soc* 2014, **136:**4551-4556.

8. Inder T, Mocatta T, Darlow B, Spencer C, Volpe JJ, Winterbourn C: **Elevated free radical products in the cerebrospinal fluid of VLBW infants with cerebral white matter injury.** *Pediatr Res* 2002, **52:**213-218.

9. Linkermann A, Skouta R, Himmerkus N, Mulay SR, Dewitz C, De Zen F, Prokai A, Zuchtriegel G, Krombach F, Welz PS, et al: **Synchronized renal tubular cell death involves ferroptosis.** *Proc Natl Acad Sci U S A* 2014, **111:**16836-16841.

10. Friedmann Angeli JP, Schneider M, Proneth B, Tyurina YY, Tyurin VA, Hammond VJ, Herbach N, Aichler M, Walch A, Eggenhofer E, et al: **Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice.** *Nat Cell Biol* 2014, **16:**1180-1191.

11. Sun X, Ou Z, Chen R, Niu X, Chen D, Kang R, Tang D: **Activation of the p62-Keap1-NRF2 pathway protects against ferroptosis in hepatocellular carcinoma cells.** *Hepatology* 2016, **63:**173-184.

12. Xie Y, Zhu S, Song X, Sun X, Fan Y, Liu J, Zhong M, Yuan H, Zhang L, Billiar TR, et al: **The Tumor Suppressor p53 Limits Ferroptosis by Blocking DPP4 Activity.** *Cell Rep* 2017, **20:**1692-1704.

13. Eling N, Reuter L, Hazin J, Hamacher-Brady A, Brady NR: **Identification of artesunate as a specific activator of ferroptosis in pancreatic cancer cells.** *Oncoscience* 2015, **2:**517-532.

14. Louandre C, Marcq I, Bouhlal H, Lachaier E, Godin C, Saidak Z, Francois C, Chatelain D, Debuysscher V, Barbare JC, et al: **The retinoblastoma (Rb) protein regulates ferroptosis induced by sorafenib in human hepatocellular carcinoma cells.** *Cancer Lett* 2015, **356:**971-977.

15. Louandre C, Ezzoukhry Z, Godin C, Barbare JC, Maziere JC, Chauffert B, Galmiche A: **Iron-dependent cell death of hepatocellular carcinoma cells exposed to sorafenib.** *Int J*

*Cancer* 2013, **133:**1732-1742.

16. Dixon SJ, Winter GE, Musavi LS, Lee ED, Snijder B, Rebsamen M, Superti-Furga G, Stockwell BR: **Human Haploid Cell Genetics Reveals Roles for Lipid Metabolism Genes in Nonapoptotic Cell Death.** *ACS Chem Biol* 2015, **10:**1604-1609.

17. Doll S, Proneth B, Tyurina YY, Panzilius E, Kobayashi S, Ingold I, Irmler M, Beckers J, Aichler M, Walch A, et al: **ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid composition.** *Nat Chem Biol* 2017, **13:**91-98.

18. Lu B, Chen XB, Ying MD, He QJ, Cao J, Yang B: **The Role of Ferroptosis in Cancer Development and Treatment Response.** *Front Pharmacol* 2017, **8:**992.

19. Sun X, Niu X, Chen R, He W, Chen D, Kang R, Tang D: **Metallothionein-1G facilitates sorafenib resistance through inhibition of ferroptosis.** *Hepatology* 2016, **64:**488-500.

20. Wang W, Green M, Choi JE, Gijón M, Kennedy PD, Johnson JK, Liao P, Lang X, Kryczek I, Sell A, et al: **CD8+ T cells regulate tumour ferroptosis during cancer immunotherapy.** *Nature* 2019.

21. Yang W, Soares J, Greninger P, Edelman EJ, Lightfoot H, Forbes S, Bindal N, Beare D, Smith JA, Thompson IR, et al: **Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker discovery in cancer cells.** *Nucleic Acids Res* 2013, **41:**D955-961.

22. Charoentong P, Finotello F, Angelova M, Mayer C, Efremova M, Rieder D, Hackl H, Trajanoski Z: **Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade.** *Cell Rep* 2017, **18:**248-262.

23. Peri S, Navarro JD, Amanchy R, Kristiansen TZ, Jonnalagadda CK, Surendranath V, Niranjan V, Muthusamy B, Gandhi TK, Gronborg M, et al: **Development of human protein reference database as an initial platform for approaching systems biology in humans.** *Genome Res* 2003, **13:**2363-2371.

24. Keshava Prasad TS, Goel R, Kandasamy K, Keerthikumar S, Kumar S, Mathivanan S, Telikicherla D, Raju R, Shafreen B, Venugopal A, et al: **Human Protein Reference Database--2009 update.** *Nucleic Acids Res* 2009, **37:**D767-772.

25. Van Allen EM, Wagle N, Stojanov P, Perrin DL, Cibulskis K, Marlow S, Jane-Valbuena J, Friedrich DC, Kryukov G, Carter SL, et al: **Whole-exome sequencing and clinical interpretation of formalin-fixed, paraffin-embedded tumor samples to guide precision cancer medicine.** *Nat Med* 2014, **20:**682-688.

26. Robinson MD, McCarthy DJ, Smyth GK: **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.** *Bioinformatics* 2010, **26:**139-140.

27. Hanzelmann S, Castelo R, Guinney J: **GSVA: gene set variation analysis for microarray and RNA-seq data.** *BMC Bioinformatics* 2013, **14:**7.

28. Canisius S, Martens JW, Wessels LF: **A novel independence test for somatic alterations in cancer shows that biology drives mutual exclusivity but chance explains most co-occurrence.** *Genome Biol* 2016, **17:**261.

29. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T: **Cytoscape: a software environment for integrated models of biomolecular interaction networks.** *Genome Res* 2003, **13:**2498-2504.

30. Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, Carter SL, Stewart C, Mermel CH, Roberts SA, et al: **Mutational heterogeneity in cancer and the**

search for new cancer-associated genes. *Nature* 2013, **499:**214-218.

31.     Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP: **Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.** *Proc Natl Acad Sci U S A* 2005, **102:**15545-15550.

32.     Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, Sajed T, Johnson D, Li C, Sayeeda Z, et al: **DrugBank 5.0: a major update to the DrugBank database for 2018.** *Nucleic Acids Res* 2018, **46:**D1074-D1082.

33.     Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, Colaprico A, Wendl MC, Kim J, Reardon B, et al: **Comprehensive Characterization of Cancer Driver Genes and Mutations.** *Cell* 2018, **173:**371-385 e318.

34.     Wesley UV, McGroarty M, Homoyouni A: **Dipeptidyl peptidase inhibits malignant phenotype of prostate cancer cells by blocking basic fibroblast growth factor signaling pathway.** *Cancer Res* 2005, **65:**1325-1334.

35.     Masur K, Schwartz F, Entschladen F, Niggemann B, Zaenker KS: **DPPIV inhibitors extend GLP-2 mediated tumour promoting effects on intestinal cancer cells.** *Regul Pept* 2006, **137:**147-155.

36.     Pro B, Dang NH: **CD26/dipeptidyl peptidase IV and its role in cancer.** *Histol Histopathol* 2004, **19:**1345-1351.

37.     Shen H, Laird PW: **Interplay between the cancer genome and epigenome.** *Cell* 2013, **153:**38-55.

38.     Macfarlane LA, Murphy PR: **MicroRNA: Biogenesis, Function and Role in Cancer.** *Curr Genomics* 2010, **11:**537-561.

39.     Li JH, Liu S, Zhou H, Qu LH, Yang JH: **starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data.** *Nucleic Acids Res* 2014, **42:**D92-97.

40.     Hassannia B, Wiernicki B, Ingold I, Qu F, Van Herck S, Tyurina YY, Bayir H, Abhari BA, Angeli JPF, Choi SM, et al: **Nano-targeted induction of dual ferroptotic mechanisms eradicates high-risk neuroblastoma.** *J Clin Invest* 2018, **128:**3341-3355.

41.     Zhang X, Du L, Qiao Y, Zhang X, Zheng W, Wu Q, Chen Y, Zhu G, Liu Y, Bian Z, et al: **Ferroptosis is governed by differential regulation of transcription in liver cancer.** *Redox Biol* 2019, **24:**101211.

42.     Kettner NM, Voicu H, Finegold MJ, Coarfa C, Sreekumar A, Putluri N, Katchy CA, Lee C, Moore DD, Fu L: **Circadian Homeostasis of Liver Metabolism Suppresses Hepatocarcinogenesis.** *Cancer Cell* 2016, **30:**909-924.

43.     Kang R, Kroemer G, Tang D: **The tumor suppressor protein p53 and the ferroptosis network.** *Free Radic Biol Med* 2018.

44.     Mak MP, Tong P, Diao L, Cardnell RJ, Gibbons DL, William WN, Skoulidis F, Parra ER, Rodriguez-Canales J, Wistuba, II, et al: **A Patient-Derived, Pan-Cancer EMT Signature Identifies Global Molecular Alterations and Immune Target Enrichment Following Epithelial-to-Mesenchymal Transition.** *Clin Cancer Res* 2016, **22:**609-620.

45.     Hassani S, Khaleghian A, Ahmadian S, Alizadeh S, Alimoghaddam K, Ghavamzadeh A, Ghaffari SH: **Redistribution of cell cycle by arsenic trioxide is associated with demethylation and expression changes of cell cycle related genes in acute**

promyelocytic leukemia cell line (NB4). *Ann Hematol* 2018, **97:**83-93.

46.     Basu A, Bodycombe NE, Cheah JH, Price EV, Liu K, Schaefer GI, Ebright RY, Stewart ML, Ito D, Wang S, et al: **An interactive resource to identify cancer genetic and lineage dependencies targeted by small molecules.** *Cell* 2013, **154:**1151-1161.

47.     Feng H, Stockwell BR: **Unsolved mysteries: How does lipid peroxidation cause ferroptosis?** *PLoS Biol* 2018, **16:**e2006203.

48.     Shen Z, Song J, Yung BC, Zhou Z, Wu A, Chen X: **Emerging Strategies of Cancer Therapy Based on Ferroptosis.** *Adv Mater* 2018, **30:**e1704007.

49.     Mao WM, Zheng WH, Ling ZQ: **Epidemiologic risk factors for esophageal cancer development.** *Asian Pac J Cancer Prev* 2011, **12:**2461-2466.

50.     Pesch B, Kendzia B, Gustavsson P, Jockel KH, Johnen G, Pohlabeln H, Olsson A, Ahrens W, Gross IM, Bruske I, et al: **Cigarette smoking and lung cancer--relative risk estimates for the major histological types from a pooled analysis of case-control studies.** *Int J Cancer* 2012, **131:**1210-1219.

51.     Chow WH, Dong LM, Devesa SS: **Epidemiology and risk factors for kidney cancer.** *Nat Rev Urol* 2010, **7:**245-257.

52.     Barreiro E, Peinado VI, Galdiz JB, Ferrer E, Marin-Corral J, Sanchez F, Gea J, Barbera JA, Project EiC: **Cigarette smoke-induced oxidative stress: A role in chronic obstructive pulmonary disease skeletal muscle dysfunction.** *Am J Respir Crit Care Med* 2010, **182:**477-488.

53.     Guan SP, Tee W, Ng DS, Chan TK, Peh HY, Ho WE, Cheng C, Mak JC, Wong WS: **Andrographolide protects against cigarette smoke-induced oxidative lung injury via augmentation of Nrf2 activity.** *Br J Pharmacol* 2013, **168:**1707-1718.

54.     Louhelainen N, Myllarniemi M, Rahman I, Kinnula VL: **Airway biomarkers of the oxidant burden in asthma and chronic obstructive pulmonary disease: current and future perspectives.** *Int J Chron Obstruct Pulmon Dis* 2008, **3:**585-603.

## Table

**Table 1** – The abbreviations and numbers of samples for the 20 types of tumors investigated in this study.

| Cancer Type | Abbreviation | Number of Samples |
|---|---|---|
| Bladder urothelial carcinoma | BLCA | 412 |
| Breast invasive carcinoma | BRCA | 1098 |
| Cervical and endocervical cancers | CESC | 307 |
| Cholangiocarcinoma | CHOL | 51 |
| Colon adenocarcinoma | COAD | 460 |
| Esophageal carcinoma | ESCA | 185 |
| Glioblastoma multiforme | GBM | 613 |
| Head and Neck squamous cell carcino | HNSC | 528 |
| Kidney Chromophobe | KICH | 113 |
| Kidney renal clear cell carcinoma | KIRC | 537 |
| Kidney renal papillary cell carcinoma | KIRP | 323 |
| Liver hepatocellular carcinoma | LIHC | 377 |
| Lung adenocarcinoma | LUAD | 585 |
| Lung squamous cell carcinoma | LUSC | 504 |
| Pancreatic adenocarcinoma | PAAD | 185 |
| Prostate adenocarcinoma | PRAD | 499 |
| Rectum adenocarcinoma | READ | 171 |
| Stomach adenocarcinoma | STAD | 443 |
| Thyroid carcinoma | THCA | 503 |
| Uterine Corpus Endometrial Carcinon | UCEC | 560 |

## Figures

**Figure 1** – Differential expression of the ferroptosis regulator genes (FRGs). The somatic copy number alternations (A) and differential expression (B) of FRGs among cancers. (C) The correlation between and somatic copy number alternations and expression of FRGs. The differential methylation (D) and its correlation (E) with expression of FRGs. (F) The miRNA-mRNA network for FRGs.

**Figure 2** – The ferroptosis potential index (FPI) and its relations with histological types and molecular subtypes among cancers. (A) The different FPI between tumor and adjacent normal tissues among cancers. (B-C) The FPI for different molecular subtypes of breast cancer. (D) The different FPI among different histological types of esophageal carcinoma. The FSI for different MSI status of overall (E) and detailed (E) digestive system neoplasm.

**Figure 3** – Relations between the ferroptosis and signaling pathways and immunophenotypes. Enrichment analysis for metabolism pathway (A) and cancer signaling (B) between high- and low-FPI tumor tissues. NES is the normalized enrichment score in the GSEA algorithm. (C) Correlations between immunophenotypes and the FPI, and dots indicate statistically significant results (*P* value<0.05).

**Figure 4** – Kaplan–Meier analysis of overall survival according to the FPI among cancers.

**Additional Figure S1** – Molecular alternations of FRGs. The mutation frequencies (A), mutual exclusivity between FRGs and oncogenes (B) /tumor suppressor genes (C) for FRGs among cancers. (D) The differential expressed FRGs-related miRNAs. The correlation between FRGs expression and somatic copy number alternation, DNA methylation and miRNAs.

**Additional Figure S2** – Established the FPI in cell lines and examined the performance of the FPI. The boxes showed the FPI between withaferin A (WA), erastin or ferrostatin and the control (A, D, G). The boxes showed the comparison of three genes between stimulators and control in neuroblastoma cells (B), clear cell carcinoma cells (E) and liver cancer cells (H). The correlation between the expression of *CHAC1* and the FPI in neuroblastoma cells (C), clear cell carcinoma cells (F) and liver cancer cells (I).
correlation between FPI and the expression of well-known FRGs including PTGS2 (B), ACSL4 (C), CHAC1 (D) and driver genes (E).

**Additional Figure S3** – The correlation between the FPI and mutation status of driver genes (A). The FPI between mutant and wild type tumors for *TP53* (B) and *KRAS* (C). The differential expression of FRGs between mutant and wild type tumors for *TP53* (D) and *KRAS* (E).

**Additional Figure S4** – The correlation between ferroptosis and immune cells (A) and clinical characteristics (B-C). The overall prognosis abilities of FRGs (D).

**Additional Figure S5** – (A) The co-expression between FRGs and 136 genes for targeted therapy. (B) The number of CAGs correlated with FRGs. (C) The number of FRG-CAG correlation pairs.

**Additional Figure S6** – (A) The co-expression between FRGs and 20 genes for immunotherapy. (B) The interactions between drug, drug target and FRGs. (C) The correlation between expression of FRGs and the area under the dose-response curve (AUCs) for drugs.