Representations of complex contexts: A role for hippocampus

Halle R. Dimsdale-Zucker, Maria E. Montchal, Zachariah M. Reagh, Shao-Fang Wang, Laura A. Libby, Charan Ranganath

Abstract. The hippocampus plays a critical role in supporting episodic memory, in large part by binding together experiences and items with surrounding contextual information. At present, however, little is known about the roles of different hippocampal subfields in supporting this item-context binding. To address this question, we constructed a task in which items were affiliated with differing types of context – cognitive associations that vary at the local, item level and membership in temporally organized lists that linked items together at a global level. Participants made item recognition judgments while undergoing high-resolution fMRI imaging. We performed voxel pattern similarity analyses to answer the question of how human hippocampal subfields represent retrieved information about cognitive states and the time at which a past event took place. As participants recollected previously presented items, activity patterns in the CA23DG subregion carried information about prior cognitive states associated with these items. We found no evidence to suggest reinstatement of information about temporal context at the level of list membership, but exploratory analyses revealed representations of temporal context at a coarse level in conjunction with representations of cognitive contexts. Results are consistent with characterizations of CA23DG as a critical site for binding together items and contexts in the service of memory retrieval.

Introduction. Converging evidence suggests that the hippocampus (HC) plays a critical role in memory for events and their episodic details (Eichenbaum et al., 2007; Scoville & Milner, 1957; Vargha–Khadem et al., 1997). The HC is a circuit of interconnected subfields with different anatomical (Amaral & Witter, 1989; Burwell, 2000; Kesner & Rolls, 2015; Lavenex & Amaral, 2000; Witter et al., 2000) , and, hence, computational (Marr, 1971; Norman & O'Reilly, 2003; Schapiro et al., 2017; Treves & Rolls, 1994) properties. Evidence from animal models has suggested that different hippocampal subfields uniquely contribute to memory through different computational specializations (e.g., pattern separation and completion; for reviews, see (Liu et al., 2015; Yassa & Stark, 2011) that enable one to encode the spatial and task variables that create a context for episodic memories (Kesner & Rolls, 2015; Levy, 1996; Lisman, 1999; Mankin et al., 2015; Norman & O'Reilly, 2003; Wallenstein et al., 1998). This is particularly true when task demands highlight the need to bind together item-in-context representations, such as learning where to search for a reward given an animal's current location (McKenzie et al., 2014).

Evidence in humans generally corroborates the research in animal models, suggesting that, as a whole, the HC supports episodic memory by binding together information about items and the context in which they were encountered (Davachi, 2006; Eacott & Gaffan, 2005; Eichenbaum et al., 2007; O'Keefe & Nadel, 1978; Ranganath, 2010; Y. Zheng et al., 2021). At present, however, there is little known about the roles of different hippocampal subfields in item-context binding. Many studies have used high-resolution fMRI to examine item recognition (Bakker et al., 2008; Carr et al., 2010; Chen et al., 2011; Lacy et al., 2011; LaRocque et al., 2013; Reagh et al.,

2014; Suthana et al., 2015; Viskontas et al., 2009; Yassa et al., 2011; Zeineh et al.,

2003) or representations of spatial contexts (Brown et al., 2014, 2014; Chanales et al.,

2017; Copara et al., 2014, 2014; Dimsdale-Zucker et al., 2018; Stokes et al., 2015;

Suthana et al., 2011; L. Zheng et al., 2021).

Cognitive theories of episodic memory have conceptualized context in at least

two ways. One view emphasizes context as a variable that changes over time (Estes,

1955; Howard & Kahana, 2002) due to ongoing fluctuations in the participant's

environment and cognitive state (e.g., the frank passage of time, changes in stimuli,

shifting mental and physical states, etc.). Some high-resolution imaging studies have

revealed evidence suggesting that the hippocampus can represent information about

temporal context associated with particular items or associations between items based

on temporal contiguity (Copara et al., 2014; Deuker et al., 2016; Dimsdale-Zucker et al.,

2018; Nielson et al., 2015; Schapiro et al., 2013).

Another view is that episodic memories are associated with "cognitive contexts"

(Diana et al., 2012, 2013)—relatively discrete cognitive states that are tied to attentional

priorities, task demands, or currently relevant goals (Aly & Turk-Browne, 2016a, 2016b;

Antony et al., 2021; Diana et al., 2008, 2012). These views are not mutually exclusive,

and some models suggest that context in episodic memory may be represented through

a combination of temporal context and cognitive representations of task states (Lohnas

et al., 2015; Polyn et al., 2009).

In order to understand whether shifting cognitive contexts affiliated with encoding

items could drive differences in representations across hippocampal subfields, we

constructed a task in which we experimentally manipulated the cognitive context

affiliated with studied items by varying the encoding question associated with an item

(Davachi et al., 2003; Diana et al., 2008, 2012, 2013; Dzulkifli & Wilding, 2005; J. D.

Johnson et al., 2009; M. K. Johnson et al., 1997; McDuff et al., 2009; Nolde et al., 1998;

Polyn et al., 2009; Ranganath et al., 2004; Ritchey et al., 2015). To manipulate temporal

context, we organized items into repeated lists, such that items in the same list would

be expected to have stronger intra-list temporal associations than items in different lists.

This allowed us to test the extent to which hippocampal subfields represent information

about cognitive (i.e., encoding question) and temporal (i.e., list membership) contextual

associations. We then scanned participants while they recollected these objects and

used voxel pattern similarity (PS) analyses (Dimsdale-Zucker & Ranganath, 2018;

Kriegeskorte et al., 2008) to examine patterns of activity in hippocampal subfields

reflected spontaneous retrieval of information from the study phase about the temporal

and cognitive context associated with each item.

Method.

Participants. 32 participants were recruited from the community and were

compensated $50 for their time. This study was approved by the Institutional Review

Board at the University of California, Davis. Four participants were excluded due to

missing behavioral data, two participants were excluded for excessive motion that

prevented tracing of hippocampal subfields, one participant was excluded due to an

experimenter error at data collection that resulted in the incorrect stimuli being seen,

and one participant was excluded because they only had one run of usable data after

discarding motion-contaminated and data-collection contaminated runs. The results

below reflect data from 24 remaining participants (Mage = 22.85 years, SD = 3.06

years, Nfemale = 13). One of these 24 participants was excluded from behavioral

cognitive and temporal context analyses due to partially missing data; since the brain

imaging data for this participant were complete and did not depend on this behavior

being recorded, they were included in all other analyses.

Encoding. Participants viewed eight 36-item lists of still pictures of everyday

objects (e.g., contact lens case, french fries; http://cvcl.mit.edu/mm/uniqueObjects.html;

see Figure 1). Object assignment to list and presentation order of objects within a list

were uniquely randomized for each participant via the Matlab randperm function. To

encourage participants to learn temporal relationships amongst items in a list (Palombo

et al., 2019), each list was presented three times in a mini-block before subjects saw

items from the next list (e.g., 1, 1, 1, 2, 2, 2...8, 8, 8). A label with a list number (e.g.,

"List 1") appeared at the beginning of each list prior to seeing any objects. Mini-blocks

were separated with a self-paced break. Presentation order of objects within a list was

identical for all three list presentations.

Objects remained on the screen for 2.5 seconds (timing and presentation

parameters were controlled via Presentation [Neurobehavioral Systems, Inc., Berkeley,

CA, www.neurobs.com]) while the participant made a yes/no button response to an

orienting question (cognitive context). To manipulate cognitive context, each object was

associated with one of four questions: Would this item fit in a refrigerator?, Would this

item fit in a bathtub?, Would you find this item in a convenience store?, Would you find

this item in a supermarket?. Each of the four questions was presented equally often in

each block, and question/object pairs remained the same across all three list

presentations. Participants were instructed that this was a decision-making task and

that there would be some repetition but to concentrate on doing the task. Participants

were not aware that memory for these questions would be tested later, thus the learning

of question (cognitive) and temporal context information was incidental. Grouping items

into lists was meant to induce a shared context representation for items in the same list.

We reasoned that repetition of lists during the study phase would enhance contextual

relationships between items in the same list. A similar temporal grouping strategy has

been used previously to encourage connections between items in this fashion (Palombo

et al., 2019). In contrast to the inter-item links facilitated by list membership, previous

work has shown that interspersing questions throughout the encoding phase can result

in fast, shifting variations in context that are aligned to changes in cognitive state (Polyn

et al., 2009). Thus, to target cognitive context, each item was affiliated with one of four

orienting questions (for a similar manipulation of cognitive context see (Diana et al.,

2012). We note that because we did not collect fMRI data during the encoding phase,

we are limited to looking at reactivated representations of cognitive and temporal

contexts at the time of retrieval.

Scanned object recognition. While in the MRI scanner, participants saw each

of the 288 old objects from encoding as well as 72 new objects presented one at a time

for 2.5 seconds with a jittered ITI ranging from 2-15 seconds (mean ITI jitter = 6

seconds). Objects were divided into 6 runs (60 trials per run). Object order within a run

was pseudo-randomized such that objects with the same encoding question always had

at least one intervening object (e.g., fridge, convenience store, bathtub, fridge,

supermarket, fridge, etc.) to help minimize encoding context reinstatement biases on PS

results (see Multivariate Results below). Proximity of objects from encoding mini-blocks

(1-8) was not considered in the pseudo-randomization.

While in the scanner, participants were instructed to indicate via button press

whether or not they remembered the object on a 4-point scale: 1=new, 2=familiar (old

but no remembered details), 3=remembered non-temporal details (e.g., the encoding

question, something about the object itself, or an association they made with the

object), 4=remembered temporal detail (e.g., in what list or when they had seen the

object during encoding). Responses for remembered judgments were collapsed into a

single response bin for behavioral and fMRI analyses.

Source memory: Cognitive context. After completing MRI scanning,

participants returned to the lab where they completed a cognitive context source

memory task. In this phase, participants saw all 288 studied objects from encoding and

were asked to indicate which encoding question (fridge/bathtub/convenience

store/grocery store) had been associated with the object. Objects were presented

across four blocks of 72 trials each. Within each block, there were an equal number of

objects from each encoding mini-block (1-8). Presentation order of objects was uniquely

randomized by participant within each source memory block. Objects appeared on the

screen until the participant had made their source memory judgment. There was no

opportunity to guess or skip objects.

Source memory: Temporal context. After completing the cognitive context

source memory test, participants again saw the 288 old objects from encoding and this

time were asked to indicate in which mini-block (1-8) the object had appeared. Objects

were presented centrally on the screen and a number line displaying the numeric

options 1-8 appeared below. Participants were told to use the numbers at the top of the

keypad to make their response. Objects were again divided across four blocks of 72

trials with a different randomization order than was used in the task context source

memory test. Objects remained on the screen until the participant had made their

response. There was no opportunity to guess or skip temporal context judgments.

encoding (unscanned)

encoding question

list 1 repetition 1   list 1 repetition 2   list 1 repetition 3

fridge?

bathtub?

supermarket?

convenienece store?

list 2 repetition 1   list 2 repetition 2   list 2 repetition 3

convenienece store?

bathtub?

supermarket?

supermarket?

• • •

list 8 repetition 1   list 8 repetition 2   list 8 repetition 3

bathtub?

supermarket?

fridge?

convenienece store?

object recognition (fMRI)

same cognitive context   different temporal context

pattern similarity analysis

similar temporal context   same temporal context

Figure 1. Task structure. During the encoding phase, each participant studied lists of 36 objects that were each randomly paired with one of four encoding questions ("Would this item fit in a fridge?", "Would this item fit in a bathtub?", "Would you find this item in a supermarket?", "Would you find this item in a convenience store?"). Each list was repeated three times in a row to promote learning of the temporal relationships amongst the items. Objects appeared in the same order and with the same question (cognitive context) across all repetitions. High-resolution functional magnetic resonance brain imaging (fMRI) was used to examine hippocampal activity patterns during a recognition memory test for these objects, allowing us to examine activity pattern similarity as a function of whether pairs of items were encoded within the same or similar temporal contexts (i.e., studied in the same list or temporally proximal lists), and/or the same cognitive context (i.e., associated encoding question).

fMRI acquisition and pre-processing. Scans were acquired on a Siemens

Skyra 3T scanner with a 32 channel head coil. Two sets of structural images were

acquired to enable subfield segmentation: A T1-weighted magnetization prepared rapid

acquisition gradient echo (MP-RAGE) pulse sequence image (1 mm isotropic voxels),

and a high-resolution T2-weighted image (TR = 4200 ms; TE= 93 ms; field of view =

200 mm^2; flip angle = 139°; bandwidth = 199 Hz/pixel; voxel size = 0.4 x 0.4 x 1.9 mm;

58 coronal slices acquired perpendicular to the long-axis of the hippocampus). High-

resolution functional (T2*) images were acquired using a multiband gradient echo planar

(EPI) imaging sequence (TR = 2010 ms; TE = 25 ms; field of view = 216 mm; image

matrix = 144 x 152; flip angle = 79°; bandwidth = 1240 Hx/pixel; partial phase Fourier =

6/8; parallel imaging = GRAPPA acceleration factor 2 with 72 reference lines; multiband

factor = 2; 52 oblique axial slices acquired parallel to the long-axis of the hippocampus

slices; voxel size = 1.5 mm isotropic).

SPM8 (http://www.fil.ion.ucl.ac.uk/spm/) was used for image pre-processing.

Functional EPI images were realigned to the first image and resliced. No slice timing

correction was performed due to the acquisition of multiple simultaneous slices with the

multiband sequence (capabilities to handle multiband timing do not exist in SPM8). Co-

registration between the native-space ROIs defined in T2 space and the functional

images was done with SPM's Coregister: Estimate and Reslice procedure. This

procedure uses a linear normalized mutual information cost-function between a

reference (mean functional) image and source (T2) image to compute and apply a

voxel-by-voxel affine transformation matrix. This transformation matrix was then applied

to the subfield ROIs that had been defined in T2 space (see *ROI segmentation*) to bring

them into register with the functional images. The T1 image was co-registered to the

mean EPI. Then, nonlinear spatial normalization parameters were derived by

segmenting the coregistered T1 image. Quality assurance included identifying suspect

timepoints via custom code (https://github.com/memobc/memolab-fmri-qa) defined as

time-points in excess of 0.5 mm frame displacement (based on (Power et al., 2012) or

1.5% global mean signal change (based on ARTRepair recommendations, (Mazaika et

al., 2005)). Runs were excluded if the frame displacement exceeded the voxel size. As

reported earlier, three participants were excluded for motion in excess of these

thresholds; of the 24 subjects included in the analyses, 9 had runs excluded based on

these thresholds (mean number of removed runs = 0.92, SD = 1.38; ranging from 0-4

runs).

Pattern similarity analyses. PS analyses were conducted on beta maps

generated from unsmoothed data in native subject space. Following the least squares

separate procedure described by Mumford (2012), single trial models were generated to

estimate the unique beta map for every trial in a run (N=60). Within each single trial

model, the first regressor modeled the trial of interest with a stick function, the second

regressor modeled all other trials in that run, six regressors were used to capture

motion, and any additional spike regressors as identified by our QA scripts were used to

capture additional residual variance. Voxel-wise patterns of hemodynamic activity were

separately extracted for each ROI from the single trial beta images. To ensure robust

ability to detect differences in PS, we required temporal signal-to-noise ratios (TSNR) in

a region to be above 20 (approximately 2 standard deviations below the mean global

TSNR of 50.4). This required the removal of entorhinal cortex and its subregions (mean

TSNR ranged between 10-20), despite its compelling role in the representation of

temporal context (Bellmund et al., 2019; Montchal et al., 2019)

Within each ROI, correlations (Pearson's r) were computed between these trial-

wise betas to yield a trial-by-trial correlation matrix that related each voxel's signal on a

trial to all other trials across all runs. We restricted comparisons to those trials for which

participants made a correct "remember" response (during MRI scanning). Trials were

sorted on the basis of encoding context (cognitive, temporal), however source memory

for these contexts was not taken into consideration. Correlation values were z-

transformed prior to statistical analysis. Statistical analyses tested for differences in

correlations between trial pairs on the basis of encoding context (cognitive context:

same vs. different encoding question; temporal context: same vs. different encoding list,

or similar vs. different list half). To more accurately characterize both within- and across-

subject error variance (Baayen et al., 2008; Clark, 1973; Dixon, 2008; Jaeger, 2008;

Mumford & Poldrack, 2007; Singmann & Kellen, 2019), we implemented a mixed-

modelling approach to evaluate statistical significance with the *lme4* packing in R (Bates

et al., 2014); for a similar approach, see (Dimsdale-Zucker et al., 2018). We used an

alpha of $p < 0.05$ and performed permutation testing for all of the pattern similarity

results reported in the manuscript. Only between-run correlations were used to

maximize the number of possible trial pairs without mixing within- and between-run

correlations. Trial pairs of interest were extracted from these trial-by-trial correlation

matrices.

All relevant code (https://github.com/hallez/tempcon_pub), a reproducible

compute environment (https://doi.org/10.24433/CO.0129473.v1), and relevant data

(https://osf.io/qfcjg/) are available online.

ROI definition. Hippocampal subfields were defined following the procedure

reported in (Dimsdale-Zucker et al., 2018; see Figure 2). In short, the ASHS automated

segmentation procedure was used to delineate subfields in subject-native space

(Yushkevich et al., 2010). We restricted our analyses to hippocampal body where

discriminating subfields is most agreed upon. Medial temporal lobe cortical regions were

manually-traced (see the Libby and Ranganath protocol in (Yushkevich et al., 2015)). In

accordance with prior findings suggesting functional distinctions between anterior and

posterior parahippocampal gyrus (Aminoff et al., 2007; Baldassano et al., 2013, 2016),

we subdivided parahippocampal cortex one slice posterior to the wing of the ambient

cistern (Frankó et al., 2014).

Because ROI definition and selection for comparisons was a priori, we performed

independent analyses on each ROI. As Poldrack and Mumford (2009) suggest, taking

this approach does not necessitate multiple comparison correction. Furthermore, we

were primarily interested in characterizing the content of reactivated encoding

contextual information within *each* subfield rather than making direct comparisons

between the similarities or differences in what types of information each subfield

primarily represented.

anterior ⟶ posterior

Figure 2. Top: Segmentations of hippocampal subfields (Cornu Ammonis [CA] 1, a combined CA2, CA3, and dentate gyrus [CA23DG] region, and subiculum [SUB]) as well as cortical medial temporal lobe regions (perirhinal cortex [PRC], anterior parahippocampal cortex [aPHC], and posterior parahippocampal cortex [pPHC]). Segmentations are depicted for a representative subject in the coronal plane of a T2 image. Slices move from anterior to posterior from left-to-right. Bottom: Segmentations following reslicing to functional (EPI) space displayed on a mean functional image from the same subject.

Results.

  Behavioral results. During MRI scanning participants performed a recognition memory test requiring judgments as to whether each item was recognized on the basis of recollection of specific item and source information from the study phase (see *Method*). Pattern similarity analyses were restricted to correctly remembered items. On average, this yielded 200 trials (SD = 51.9; minimum trials = 71; maximum trials = 282) from which we constructed all possible trial pairs and then compared activity patterns (see fMRI Results).

Correct remember judgments were the most common response (mean hit rate =

0.69, SD = 0.19), and, for these items, participants showed high accuracy at

remembering the associated encoding task context (mean hit rate = 0.71, SD = 0.12).

Memory for the exact ("same") temporal context (list 1-8) was poor (mean hit rate =

0.18, SD = 0.03). Memory for task context and exact temporal context for items that

were correctly remembered as studied were not significantly correlated with one another

(r = -0.31, p = 0.16).

We reasoned that, even if participants were unable to recall the exact list identity,

they might have memory for the "similar" temporal context associated with each item.

Visual inspection of temporal source errors indicated that participants were more likely

to incorrectly identify an item as coming from a list near in time to when it was actually

studied as compared to lists further away (e.g., if an item was actually studied in List 3,

participants would have been more likely to incorrectly say it had been in List 2 or 4 than

lists further away in time; see Appendix A).

We therefore performed an exploratory analysis in which we re-scored each trial

according to whether the participant could accurately determine whether it was

presented in the "similar" (first half (lists 1-4) or the second half (lists 5-8)) versus

"different" (across halves, i.e., list 1/list5, list 1/list 6, etc.) temporal context of the

encoding phase. On this metric, memory for similar temporal context (mean accuracy

for list half = 0.59, SD = 0.04) was reliably greater than chance of 0.5, t(22) = 8.74, p <

0.001. Thus, although participants did not have access to the precise list in which an

item had been studied, they were able to retrieve information about the item's temporal

context at a coarse level.

Appendix A. Temporal (list) source memory errors were more likely to occur at nearer
than further lags. That is, items were more likely to be incorrectly attributed to a list that
was closer in time (+/-1 or +/-2 lists) to the list in which it was actually located at
encoding. We take this as evidence that although exact memory for which of the eight
encoding lists an item was studied in was relatively poor, participants had access to the
relative temporal context of an item at encoding.

fMRI results. We next tested whether activity patterns in the hippocampal

subfields (CA1, CA23DG, subiculum[1]) during memory retrieval carried information about

the context in which the item was previously encountered. Specifically, we examined

voxel PS during retrieval as a function of whether pairs of trials shared a temporal

---

[1] For medial temporal neocortical regions, see reported results in Appendix H.

(same list vs. different list [*same* temporal context] and/or cognitive (i.e., same encoding

task vs. different encoding task) context when the items were originally learned.

We first considered when items came from the same temporal encoding context

(same vs. different list). No hippocampal subfield showed significant PS differences

between retrieval of items from the same list as compared with items from different lists

(all $X^2 < 0.5$, all ps > 0.40; see Appendix C for statistics and Appendix B for visualization

of PS values between lists).

332
333   **Appendix B.** Pattern similarity split by list across subfields of interest. No subfield
334   showed a reliable relationship in PS levels across list. We take this as evidence that list
335   membership was not a salient feature that participants reactivated at the time of
336   retrieval. This aligns with participants' poor source memory for the list in which an item
337   was studied during encoding.
338
339   We next considered whether hippocampal voxel patterns during recollection of

340   studied items carried information about the cognitive context (encoding question)

341   associated with the item during encoding. In CA23DG, we found that PS was higher

342   during retrieval of items associated with the same cognitive context compared to items

343   associated with different cognitive contexts ($X^2(1) = 4.63$, $p^{perm1000} = 0.031$; Figure 3[2]).

---

[2] For ease of visualization of the effects, for pattern similarity analyses we have chosen to plot means with
individual subject points to illustrate the observed data. This does not exactly recapitulate the way the statistical
comparisons were computed since these were performed as mixed models. That is, these figures show the raw
pattern similarity values for these experimental conditions but do not account for the other effects present in the

344   No other subfield showed significant effects of cognitive context (all $X^2 < 0.3$, all ps >

345   0.50; see Appendix D) nor the combination of same versus different temporal and

346   cognitive contexts (all $X^2 < 0.80$, all ps > 0.30; see Appendix D) on PS[3].

348
349
Figure 3. Pattern similarity values in CA23DG during memory retrieval carry information
about cognitive encoding contexts. A. Mean pattern similarity scores, with scatter of
individual subject observations for the combination of different encoding contexts. Mean
PS values were greater for same as compared to different cognitive contexts. B.
Permuted chi-square values to determine significance of cognitive context main effect.
Observed chi-square value is depicted in blue and the significance threshold ($p = 0.05$)
is plotted in a dashed red line.
357

---

models that were used to determine significance. To better visualize the observed statistical effects, we have plotted the permutation distributions that were used to determine model significance.

[3] At the suggestion of an anonymous reviewer, we repeated all of our analyses for the key effects we saw in CA23DG using a Bayesian modelling framework. We fit all models using Hamiltonian Monte Carlo No-U-Turn sampling as implemented by the brms package in the R computing environment (Bürkner, 2017). For all models, we fit 4 chains of 10,000 sampling iterations (5,000 warmup) each for a total of 20,000 post-warmup samples. This sampling iteration size was used to avoid cases where the tail effective sample size was low (as indicated by Stan warning messages). Extraction and transformation of posterior draws after models were fit was done using the *tidybayes* package and the *tidyverse* collection of packages in R (Kay, 2022; Wickham et al., 2019). This approach was informed by best practices as advised by Paul Alexander Bloom (personal communication). Results from these Bayesian analyses replicated the results we report in the main analyses in the paper that were implemented via mixed models in *lmer*. However, for full transparency, we have included the Bayesian effect estimates for all contrasts as well as the Highest Density Interval (which can be interpreted similarly to a confidence interval, Turkkan & Pham-Gia, 1993) in Appendix D. Furthermore, we include graphical depictions of the model posterior distribution and conditional effect estimates for any contrast that was deemed significant (see Appendices E-G).

We next performed a series of exploratory analyses to consider whether PS

might depend on whether items shared a "similar" temporal context at encoding (same

vs. different half of the encoding phase). The choice of list half was informed by the fact

that participants were behaviorally able to discriminate coarse temporal context reliably

greater than chance, however we present these analyses as exploratory because

participants were not explicitly asked to make temporal source judgments at the

granularity of list half. This analysis revealed no significant effects of similar temporal

context alone (all $X^2 < 2.5$, all ps > 0.10; see Appendix D).

We next considered whether hippocampal voxel patterns during recollection of

studied items carried information about the cognitive context (encoding question)

associated with the item during encoding when we defined temporal context as "similar".

Consistent with our previous analysis, in CA23DG, PS for pairs of items that were

associated with the same cognitive context was higher than for pairs of items that were

associated with different cognitive contexts ($X^2(1) = 4.68$, $p^{perm1000} = 0.031$). This effect

was qualified by a significant similar temporal by cognitive context interaction ($X^2(1) =$

8.11, $p^{perm1000} = 0.004$; see Figure 4), such that the effect of cognitive context in

CA23DG was larger for items that were in similar temporal contexts (same half) than for

items that were in different temporal contexts (different half) particularly when these

items shared the same cognitive context. No other subfield showed significant effects of

cognitive context (all $X^2 < 0.30$, all ps > 0.80; see Appendix D) nor the combination of

similar versus different temporal and cognitive contexts (all $X^2 < 3$, all ps > 0.080; see

Appendix D) on PS.

381
382
Figure 4. Pattern similarity values in CA23DG during memory retrieval carry information
about similar temporal and cognitive encoding contexts. A. Mean pattern similarity
scores, with scatter of individual subject observations for the combination of different
encoding contexts. B. Permuted chi-square values to determine significance of
cognitive-by-(similar)temporal interaction. Observed chi-square value is depicted in blue
and the significance threshold (p = 0.05) is plotted in a dashed red line.
389
390

Appendix C. Pattern similarity findings in hippocampal subfields. Italics indicate significant modulation of pattern similarity values by cognitive, temporal, or interaction of cognitive and temporal encoding contexts.

	"Same" Temporal Context			"Similar" Temporal Context		
	Cognitive	**Temporal**	**Cognitive x Temporal**	**Cognitive**	**Temporal**	**Cognitive x Temporal**
CA1	PS levels did not vary by cognitive context $X^2(1)$ = 0.08, $p^{perm1000}$ = 0.773	PS levels did not vary by same/different temporal context $X^2(1)$ = 0.11, $p^{perm1000}$ = 0.744	PS levels did not vary by the interaction between same temporal and cognitive contexts $X^2(1)$ = 0.03, $p^{perm1000}$ = 0.871	PS levels did not vary by cognitive context ($X^2(1)$ = 0.08, $p^{perm1000}$ = 0.776)	PS levels did not vary by similar temporal context ($X^2(1)$ = 2.42, $p^{perm1000}$ = 0.120)	PS levels did not vary by the interaction between similar temporal and cognitive contexts ($X^2(1)$ = 0.84, $p^{perm1000}$ = 0.360)
CA23DG	*PS values were greater for same as compared to different cognitive contexts $X^2(1)$ = 4.63, $p^{perm1000}$ = 0.031*	PS levels did not vary by same/different temporal context $X^2(1)$ = 0.44, $p^{perm1000}$ = 0.505	PS levels did not vary by the interaction between same temporal and cognitive contexts $X^2(1)$ = 0.71, $p^{perm1000}$ = 0.400	*PS values were greater for same as compared to different cognitive contexts $X^2(1)$ = 4.68, $p^{perm1000}$ = 0.031*	PS levels did not vary by similar temporal context $X^2(1)$ = 0.08, $p^{perm1000}$ = 0.783	*PS was greater for items that shared a similar temporal context as well as a cognitive context but was lower when these items did not share a cognitive context $X^2(1)$ = 8.11, $p^{perm1000}$ = 0.004*
Subiculum	PS levels did not vary by cognitive context $X^2(1)$ = 0.27, $p^{perm1000}$ = 0.605	PS levels did not vary by same/different temporal context $X^2(1)$ = 0.13, $p^{perm1000}$ = 0.721	PS levels did not vary by the interaction between same temporal and cognitive contexts $X^2(1)$ = 0.03, $p^{perm1000}$ = 0.856	PS levels did not vary by cognitive context ($X^2(1)$ = 0.27, $p^{perm100}0$ = 0.602)	PS levels did not vary by similar temporal context ($X^2(1)$ = 0.11, $p^{perm1000}$ = 0.745)	PS was *marginally* higher for items that were studied relative to the same question but studied in different list halves as compared to all other conditions ($X^2(1)$ = 2.91, $p^{perm1000}$ = 0.088)

Appendix D. Bayesian model results for pattern similarity findings in CA23DG.
Here we report effect estimates for contrasts as estimated by the *brms* package in the R computing environment (Bürkner, 2017). Highest density intervals (abbreviated in the table as "posterior intervals" [PI]) were calculated for each contrast. Such intervals are roughly analogous to confidence intervals, and, thus, must not contain zero for the contrast to be considered significant (Turkkan & Pham-Gia, 1993). Italics are used to emphasize contrasts that were deemed significant based on this criteria.

	"Same" Temporal Context			"Similar" Temporal Context		
	Cognitive	**Temporal**	**Cognitive x Temporal**	**Cognitive**	**Temporal**	**Cognitive x Temporal**
CA23DG	*PS values were greater for same as compared to different cognitive contexts; same > different cognitive context estimate = 0.0004, PI = (0.00005, 0.0008)*	PS levels did not vary by same/different temporal context; different > same list estimate = 0.000182, PI = (-0.0003, 0.0007)	PS levels did not vary by the interaction between same temporal and cognitive contexts; different list & same cognitive context > same list & different cognitive context estimate = -0.0005, PI = (-0.002, 0.00007)	*PS values were greater for same as compared to different cognitive contexts; same > different cognitive context estimate = 0.0004, PI = (0.00005, 0.0008)*	PS levels did not vary by similar temporal context; different half > same half estimate = 0.00005; PI = (-0.00003, 0.00004)	*PS was greater for items that shared a similar temporal context as well as a cognitive context but was lower when these items did not share a cognitive context. For items in different list halves, PS values were relatively similar regardless of same vs. different cognitive context; different list half & same cognitive context > same list half & different cognitive context estimate = -0.004, PI = (-0.002, -0.00003)*

Appendix E. Bayesian model posterior distributions and conditional effects for the influence of cognitive context (for "same" temporal context items) on PS values in CA23DG. A. Posterior distributions for the same > different cognitive context contrast, representing differences in mean pattern similarity values between these conditions. Shaded distributions represent all posterior contrast samples, vertical red line represents zero, black point below shaded distribution represent effect estimates as determined by the model's posterior predictive distribution for the linear predictor, and error bars around this point represent 95% credible intervals. Since the 95% CI does not overlap with zero, we can determine that cognitive context had a significant influence on PS values in CA23DG. B. Conditional effect estimates for the influence of same and different cognitive context on pattern similarity values, accounting for other effects present in the model.

Appendix F. Bayesian model posterior distributions and conditional effects for
the influence of cognitive context (for "similar" temporal context items) on PS
values in CA23DG. A. Posterior distributions for the same > different cognitive context
contrast, representing differences in mean pattern similarity values between these
conditions. Shaded distributions represent all posterior contrast samples, vertical red
line represents zero, black point below shaded distribution represent effect estimates as
determined by the model's posterior predictive distribution for the linear predictor, and
error bars around this point represent 95% credible intervals. Since the 95% CI does not
overlap with zero, we can determine that cognitive context had a significant influence on
PS values in CA23DG. B. Conditional effect estimates for the influence of same and
different cognitive context on pattern similarity values, accounting for other effects
present in the model.
436

437
438
439

Appendix G. Bayesian model posterior distributions and conditional effects for the influence of the combined effect of "similar" temporal context and cognitive context on PS values in CA23DG. A. Posterior distributions for the [(different list half & same cognitive context) > (same list half & different cognitive context)] contrast, representing differences in mean pattern similarity values between these conditions. Shaded distributions represent all posterior contrast samples, vertical red line represents zero, black point below shaded distribution represent effect estimates as determined by the model's posterior predictive distribution for the linear predictor, and error bars around this point represent 95% credible intervals. Since the 95% CI does not overlap with zero, we can determine that the combined influence of similar temporal context and cognitive context had a significant influence on PS values in CA23DG. Specifically, we see that PS values were greater for items affiliated both with the same cognitive context and the same list half ("similar" temporal context) than those affiliated with different cognitive contexts (comparing teal vs. red points in the left column of the graph). PS values were relatively equivalent for items affiliated with different list halves regardless of cognitive context (same/different). B. Conditional effect estimates for the influence of similar temporal context and cognitive context on pattern similarity values, accounting for other effects present in the model.

Appendix H. Pattern similarity findings in medial temporal lobe neocortical areas.
461 Italics indicate significant modulation of pattern similarity values by cognitive, temporal,
462 or interaction of cognitive and temporal encoding contexts.

	"Same" Temporal Context			"Similar" Temporal Context		
	Cognitive	**Temporal**	**Cognitive x Temporal**	**Cognitive**	**Temporal**	**Cognitive x Temporal**
Anterior PHC	*PS was higher during retrieval of items associated with the same cognitive context compared to items associated with different cognitive contexts, $X^2(1) = 5.92$, $p^{perm1000} = 0.015$*	*PS was higher during retrieval of items associated with the same as compared to different list, $X^2(1) = 4.79$, $p^{perm1000} = 0.029$*	PS levels did not vary by the interaction between same temporal and cognitive contexts $X^2(1) = 0.22$, $p^{perm1000} = 0.639$	*PS was higher during retrieval of items associated with the same cognitive context compared to items associated with different cognitive contexts, $(X^2(1) = 5.83$, $p^{perm1000} = 0.016$*	*PS was higher during retrieval of items associated with a different temporal context as compared to items with similar temporal encoding contexts, $(X^2(1) = 7.39$, $p^{perm1000} = 0.007$*	PS levels did not vary by the interaction between similar temporal and cognitive contexts $(X^2(1) = 1.42$, $p^{perm1000} = 0.233$
Posterior PHC	PS levels did not vary by cognitive context $X^2(1) = 0.64$, $p^{perm1000} = 0.422$	PS levels did not vary by same/different temporal context $X^2(1) = 2.88$, $p^{perm1000} = 0.090$	*PS was greater for items that shared a temporal context as well as a cognitive context but was lower when these items did not share a cognitive context, $(X^2(1) = 9.44$, $p^{perm1000} = 0.002$*	PS in pPHC did not vary with respect to cognitive context alone, $(X^2(1) = 0.68$, $p^{perm1000} = 0.411)$	PS levels did not vary by similar temporal context $X^2(1) = 2.07$, $p^{perm1000} = 0.150$	*PS was greater for items that shared a similar temporal context as well as a cognitive context but was lower when these items did not share a cognitive context $X^2(1) = 4.67$, $p^{perm1000} = 0.031$*
PRC	PS was *marginally higher* during retrieval of items associated with the same cognitive context compared to items associated with different cognitive contexts $X^2(1) = 3.53$, $p^{perm1000} = 0.060$	PS levels did not vary by same/different temporal context $X^2(1) = 0.074$, $p^{perm1000} = 0.786$	PS levels did not vary by the interaction between same temporal and cognitive contexts $X^2(1) = 0.96$, $p^{perm1000} = 0.327$	PS was *marginally higher* during retrieval of items associated with the same cognitive context compared to items associated with different cognitive contexts PRC, $(X^2(1) = 3.55$, $p^{perm1000} = 0.051$	PS levels did not vary by similar temporal context $X^2(1) = 0.11$, $p^{perm1000} = 0.743$	PS levels did not vary by the interaction between similar temporal and cognitive contexts $X^2(1) = 0.15$, $p^{perm1000} = 0.703$

463
464

Discussion.

The goal of the present study was to test whether hippocampal subfields represent information about experimentally-manipulated cognitive and temporal contexts (i.e., encoding task and list identity) during recollection of studied items. Results showed that cognitive context significantly influenced activity patterns in CA23DG. We saw no evidence for a precise representation of temporal (list) context, although exploratory analyses revealed that when using a coarse definition of temporal context (first vs second half of study phase), that CA23DG carried information about the *conjunction* of cognitive and temporal context. This suggests that patterns of activity within CA23DG may support successful memory retrieval by enabling recollection of task-relevant cognitive states or goals.

Cognitive Context Representation in CA23DG

The hippocampus is a prime example of how form can influence function. Computational models (Marr, 1971; for a non-Hebbian instantiation, see Y. Zheng et al., 2021) propose that the combination of sparse coding in DG and the dense recurrent collaterals that make up the majority of the inputs to CA3 (Amaral & Witter, 1989) enable the hippocampus to retrieve or recover a memory trace given a noisy or incomplete retrieval cue (i.e., "pattern completion" (Marr, 1971; O'Reilly & McClelland, 1994; Yassa & Stark, 2011). A second factor that is key to understanding hippocampal function is that CA3 and DG are critical sites for binding of information about items and contexts (Knierim et al., 2006; Y. Zheng et al., 2021). Considerable evidence suggests that lateral and medial entorhinal cortex receive distinct cortical inputs carrying information about items and contexts, respectively, and this information is subsequently

integrated via convergent inputs in DG and CA3 (Amaral & Witter, 1989; Burwell, 2000;

Eichenbaum et al., 2007; Libby et al., 2012; Rolls & Kesner, 2006). Based on these

characteristics, we would expect that the DG-CA3 circuit should play a crucial role in

recollecting contextual information about previously encountered items.

Using a paradigm adapted from Diana and colleagues (2012, 2013), in which

different objects were encoded with unique cognitive context questions, we

demonstrated that activity patterns in CA23DG carried information about an item's

cognitive context during retrieval. Pattern similarity values were greater for items that

had been affiliated with the same, as compared to different, cognitive tasks at encoding.

This result is striking, given that participants were only shown items from the study

phase, and they were not instructed to explicitly recall information about the study tasks.

Our results therefore accord with the idea that CA23DG plays an important role in

linking item and context information, such that encountering a familiar item can trigger

the recovery of contextual information via pattern completion (see also Grande et al.,

2019).

Our findings are relevant to the idea that, during conscious recollection, the

hippocampus reinstates representations of discrete cognitive contexts that encompass

one's current attentional priorities, task set, and/or goals (Aly & Turk-Browne, 2016a,

2016b; Davachi, 2006; Davachi et al., 2003; Diana et al., 2012; Eichenbaum et al.,

2007; J. D. Johnson et al., 2009; Libby et al., 2018; Ranganath et al., 2004; Ritchey et

al., 2015). For instance, Aly and colleagues (Aly & Turk-Browne, 2016a, 2016b) found

that voxel patterns in hippocampal subfields carried information about attentional

orientation during a memory encoding task ("attend to artistic features" vs. "attend to

spatial layout"). Converging evidence in rodents also suggests that task-relevant

contextual features were the most salient factor in describing hippocampal ensemble

activity similarity (conceptually analogous to the pattern similarity metrics used in human

fMRI) (McKenzie et al., 2014). Our results build on this idea by highlighting CA23DG as

a site where binding of cognitive contextual information occurs in the service of episodic

memory retrieval.

518   An intriguing possibility that could be addressed in future work is whether

representational patterns would change if cognitive contexts were designed to

systematically relate to features of the items. In the present design, item/question

pairing was uniquely randomized for each participant and thus this question cannot be

explored here. However, prior work has suggested that the hippocampus may play a

particular role in representing an item's context (e.g., the type of associated judgment)

whereas when context is posed as a feature of the item (e.g., its color) that medial

temporal cortical regions such as perirhinal cortex are engaged (Staresina & Davachi,

2008; although see Diana, Yonelinas, & Ranganath, 2008 for some suggestions that

perirhinal involvement in item/context unitization may be limited to familiarity). Thus, if

cognitive context questions were selected to represent *features* of the items, it seems

that there would be competing hypotheses about whether hippocampus would still treat

this information as an item's context or whether medial temporal lobe cortical regions

would additionally be recruited.

Contributions of temporal context

According to Tulving's definition (Tulving, 1972, 1983, 1984), episodic memories

are organized by temporal context. Temporal context has been formally operationalized

in terms of random fluctuations in cognitive states over time (Estes, 1955), and a time-

weighted average of recently processed items and experiences (Howard & Kahana,

2002; Norman et al., 2008; Sederberg et al., 2008). These models would predict that

activity patterns during memory retrieval should reflect information about the temporal

context associated with each study item (e.g., Deuker et al., 2016; Jenkins &

Ranganath, 2016; Manning et al., 2011; Nielson et al., 2015).

In the present study, when asked to choose in which of eight lists an item had

been studied, participants were essentially at chance. This is not surprising because

models suggest that temporal context is relative, not absolute (Kahana, 1996; Manning

et al., 2011). In other words, people can use recovered context to gauge the relative

recency of two previously encountered items but this might not be sufficient for precise

memory for the temporal position of an item (DuBrow & Davachi, 2013, 2014; Ezzyat &

Davachi, 2014; Hintzman, 2001, 2004, 2005; Jenkins & Ranganath, 2010, 2016;

Lewandowsky & Murdock, 1989; Montchal et al., 2019) without the use of additional

heuristics and reconstructive strategies (Friedman, 1993).

If the hippocampus carries information about the relative temporal context of past

events, we might expect that hippocampal activity patterns would be more similar during

recollection of items that were learned in close temporal proximity than for items that

were studied far apart in time. Consistent with this prediction, fMRI studies have shown

that recollection of items encountered either in virtual reality (Copara et al., 2014;

Deuker et al., 2016; Dimsdale-Zucker et al., 2018) or in the real-world (Nielson et al.,

2015), yield activity patterns within the hippocampus that show graded similarity on the

basis of both temporal and spatial contextual features.

Here, we found no evidence to suggest that activity patterns in any subfield

carried information about list context. There are a number of potential reasons for this

null result, but we suspect that the most important factor is that, unlike previous studies

that reported temporal context representations in the hippocampus, we manipulated

cognitive contexts independently of temporal information.

Our task was designed such that participants changed cognitive contexts often

within a list. Prior work has shown that when ongoing temporal contexts are interrupted

by changes in item-level cognitive contexts that both behavioral (Polyn et al., 2009) and

neural (Polyn et al., 2012) responses are shaped by both these local and global

contextual features. Thus, by varying the encoding question within each study list,

participants might have segmented each list into micro-contexts (Clewett & Davachi,

2017; DuBrow et al., 2017; Zacks et al., 2001; Zacks & Swallow, 2007) thereby

disrupting associations based on temporal contiguity. A related possibility is that, during

encoding, participants might have prioritized cognitive context over temporal contiguity,

because the encoding questions were more salient and relevant.

Although we did not find evidence of precise temporal context representation in

the hippocampus, an exploratory analysis did reveal evidence to suggest conjunctive

representation of temporal context—"similar" or "coarse" temporal context—and

cognitive context in CA23DG. In this analysis, temporal context was operationalized on

the basis of which half of the study phase that an item had been previously encountered

(first half: lists 1-4; second half: lists 5-8). Results revealed that activity patterns in

CA23DG were shaped by the *conjunction* of coarse temporal context and cognitive

context, which aligns with the binding role CA3 is thought to play (Burwell, 2000;

Eichenbaum et al., 2007; Rolls & Kesner, 2006).

Shared versus distinct contextual representations in the hippocampus

Our results conflict with a prior study, in which, we found that CA23DG activity

patterns during memory retrieval were less similar across pairs of items that were

associated with the same episodic context (i.e., objects seen within the same movie)

than across pairs of items that were associated with different spatial or episodic

contexts (Dimsdale-Zucker et al., 2018). The contrast between the present results and

those of Dimsdale-Zucker et al. (2018) is relevant to many standard resolution fMRI

studies showing that that hippocampal activity patterns are sometimes more similar

across events that have a shared context (Horner et al., 2015; Libby et al., 2018;

Milivojevic et al., 2015; Pidgeon & Morcom, 2016; Schlichting et al., 2015), and

sometimes less similar between overlapping events (Chanales et al., 2017; Favila et al.,

2016; Kim et al., 2017; Schlichting et al., 2015; L. Zheng et al., 2021). Computational

work by Ritvo (2019) suggests that these different results may be due to competition

between items that are to be learned. If there is relatively low competition, then

CA23DG might assign similar representations to overlapping events, whereas, if

competition is high, the representations can be orthogonalized. We speculate that

dynamic might be at play in our studies. In Dimsdale-Zucker et al. (2018), multiple

sequences of items were encoded in the same two virtual reality contexts, potentially

creating a significant degree of interference across items, but, in the present study, the

use of multiple encoding tasks probably reduced contextual overlap across items,

thereby reducing competition. This explanation can be directly tested in a future high-

resolution imaging study.

General Conclusions

Our results show that activity patterns in the CA23DG region of the human

hippocampus carry information about cognitive contexts. This finding can help explain

how we represent continuously unfolding episodes in a changing world. Outside of the

laboratory, we are constantly multitasking between competing goals and

responsibilities. The hippocampus, and, specifically CA23DG, may allow us to

differentiate between experiences that are associated with different tasks.

Gender, Race, and Ethnicity Bias Statement

Recent work in several fields of science has identified a bias in citation practices

such that papers from women and other minority scholars are under-cited relative to the

number of such papers in the field (Bertolero et al., 2020; Caplar et al., 2017; Chatterjee

& Werner, 2021; Dion et al., 2018; Dworkin et al., 2020; Fulvio et al., 2021; Maliniak et

al., 2013; Mitchell et al., 2013; Wang et al., 2021). Here we sought to proactively

consider choosing references that reflect the diversity of the field in thought, form of

contribution, gender, race, ethnicity, and other factors. First, we obtained the predicted

gender of the first and last author of each reference by using databases that store the

probability of a first name being carried by a woman (Dworkin et al., 2020; Zhou,

2020/2022; Zhou et al., 2020). By this measure (and excluding self-citations to the first

and last authors of our current paper), our references contain 7.65%

woman(first)/woman(last), 14.53% man/woman, 26.29% woman/man, and 51.53%

man/man. This method is limited in that a) names, pronouns, and social media profiles

used to construct the databases may not, in every case, be indicative of gender identity

and b) it cannot account for intersex, non-binary, or transgender people. Second, we

obtained predicted racial/ethnic category of the first and last author of each reference by

databases that store the probability of a first and last name being carried by an author of

color (Ambekar et al., 2009; Sood & Laohaprapanon, 2018). By this measure (and

excluding self-citations), our references contain 3.36% author of color (first)/author of

color(last), 10.63% white author/author of color, 15.17% author of color/white author,

and 70.84% white author/white author. This method is limited in that a) names and

Florida Voter Data to make the predictions may not be indicative of racial/ethnic identity,

and b) it cannot account for Indigenous and mixed-race authors, or those who may face

differential biases due to the ambiguous racialization or ethnicization of their names.

We look forward to future work that could help us to better understand how to support

equitable practices in science.

Acknowledgements. The authors wish to thank Paul Alexander Bloom for his help in

fitting and interpreting the Bayesian models.

References

Aly, M., & Turk-Browne, N. B. (2016a). Attention promotes episodic encoding by stabilizing

hippocampal representations. *Proceedings of the National Academy of Sciences*,

201518931. https://doi.org/10.1073/pnas.1518931113

Aly, M., & Turk-Browne, N. B. (2016b). Attention Stabilizes Representations in the Human

Hippocampus. *Cerebral Cortex*, *26*(2), 783–796. https://doi.org/10.1093/cercor/bhv041

Amaral, D. G., & Witter, M. P. (1989). The three-dimensional organization of the hippocampal

formation: A review of anatomical data. *Neuroscience*, *31*(3), 571–591.

Ambekar, A., Ward, C., Mohammed, J., Male, S., & Skiena, S. (2009). Name-ethnicity

classification from open sources. *In Proceedings of the 15th ACM SIGKDD International*

Conference on Knowledge Discovery and Data Mining, 49–58.

Aminoff, E. M., Gronau, N., & Bar, M. (2007). The parahippocampal cortex mediates spatial and

nonspatial associations. *Cerebral Cortex (New York, N.Y.: 1991)*, *17*(7), 1493–1503.

https://doi.org/10.1093/cercor/bhl078

Antony, J. W., Hartshorne, T. H., Pomeroy, K., Gureckis, T. M., Hasson, U., McDougle, S. D., &

Norman, K. A. (2021). Behavioral, Physiological, and Neural Signatures of Surprise during

Naturalistic Sports Viewing. *Neuron*, *109*(2), 377-390.e7.

https://doi.org/10.1016/j.neuron.2020.10.029

Baayen, H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random

effects for subjects and items. *Journal of Memory and Language*, *59*(4), 390–412.

https://doi.org/10.1016/j.jml.2007.12.005

Bakker, A., Kirwan, C. B., Miller, M., & Stark, C. E. L. (2008). Pattern separation in the human

hippocampal CA3 and dentate gyrus. *Science (New York, N.Y.)*, *319*(5870), 1640–1642.

https://doi.org/10.1126/science.1152882

Baldassano, C., Beck, D. M., & Fei-Fei, L. (2013). Differential connectivity within the

Parahippocampal Place Area. *NeuroImage*, *75*, 228–237.

https://doi.org/10.1016/j.neuroimage.2013.02.073

Baldassano, C., Esteva, A., Fei-Fei, L., & Beck, D. M. (2016). Two Distinct Scene-Processing

Networks Connecting Vision and Memory. *ENeuro*, *3*(5), ENEURO.0178-16.2016.

https://doi.org/10.1523/ENEURO.0178-16.2016

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting Linear Mixed-Effects Models using

lme4. *10.18637/Jss.V067.I01*, 1–48.

Bellmund, J. L., Deuker, L., & Doeller, C. F. (2019). Mapping sequence structure in the human

lateral entorhinal cortex. *ELife*, *8*, e45333. https://doi.org/10.7554/eLife.45333

Bertolero, M., Dworkin, J., David, S., Lloreda, C. L., Srivastava, P., Stiso, J., Zhou, D., Dzirasa, K.,

Fair, D., Kaczkurkin, A., Marlin, B. J., Shohamy, D., Uddin, L., Zurn, P., & Bassett, D.

(2020). *Racial and ethnic imbalance in neuroscience reference lists and intersections with*

gende. https://doi.org/10.1101/2020.10.12.336230

Brown, T. I., Hasselmo, M. E., & Stern, C. E. (2014). A high-resolution study of hippocampal and

medial temporal lobe correlates of spatial context and prospective overlapping route

memory. *Hippocampus*, *24*(7), 819–839. https://doi.org/10.1002/hipo.22273

Bürkner, P.-C. (2017). **brms**: An *R* Package for Bayesian Multilevel Models Using *Stan*. *Journal of*

Statistical Software, *80*(1). https://doi.org/10.18637/jss.v080.i01

Burwell, R. D. (2000). The parahippocampal region: Corticocortical connectivity. *Annals of the*

New York Academy of Sciences, *911*, 25–42.

Caplar, N., Tacchella, S., & Birrer, S. (2017). Quantitative evaluation of gender bias in

astronomical publications from citation counts. *Nature Astronomy*, *1*(6), 0141.

https://doi.org/10.1038/s41550-017-0141

Carr, V. A., Rissman, J., & Wagner, A. D. (2010). Imaging the Human Medial Temporal Lobe with

High-Resolution fMRI. *Neuron*, *65*(3), 298–308.

https://doi.org/10.1016/j.neuron.2009.12.022

Chanales, A. J. H., Oza, A., Favila, S. E., & Kuhl, B. A. (2017). Overlap among Spatial Memories

Triggers Repulsion of Hippocampal Representations. *Current Biology: CB*, *27*(15), 2307-

2317.e5. https://doi.org/10.1016/j.cub.2017.06.057

Chatterjee, P., & Werner, R. M. (2021). Gender Disparity in Citations in High-Impact Journal

Articles. *JAMA Network Open*, *4*(7), e2114509.

https://doi.org/10.1001/jamanetworkopen.2021.14509

Chen, J., Olsen, R. K., Preston, A. R., Glover, G. H., & Wagner, A. D. (2011). Associative retrieval

processes in the human medial temporal lobe: Hippocampal retrieval success and CA1

mismatch detection. *Learning & Memory*, *18*(8), 523–528.

https://doi.org/10.1101/lm.2135211

Clark, H. H. (1973). The language-as-fixed-effect fallacy: A critique of language statistics in

psychological research. *Journal of Verbal Learning and Verbal Behavior*, *12*(4), 335–359.

https://doi.org/10.1016/S0022-5371(73)80014-3

Clewett, D., & Davachi, L. (2017). The ebb and flow of experience determines the temporal

structure of memory. *Current Opinion in Behavioral Sciences*, *17*, 186–193.

https://doi.org/10.1016/j.cobeha.2017.08.013

Copara, M. S., Hassan, A. S., Kyle, C. T., Libby, L. A., Ranganath, C., & Ekstrom, A. D. (2014).

Complementary Roles of Human Hippocampal Subregions during Retrieval of

Spatiotemporal Context. *The Journal of Neuroscience*, *34*(20), 6834–6842.

https://doi.org/10.1523/JNEUROSCI.5341-13.2014

Davachi, L. (2006). Item, context and relational episodic encoding in humans. *Current Opinion in*

Neurobiology, *16*(6), 693–700. https://doi.org/10.1016/j.conb.2006.10.012

Davachi, L., Mitchell, J. P., & Wagner, A. D. (2003). Multiple routes to memory: Distinct medial

temporal lobe processes build item and source memories. *Proceedings of the National*

Academy of Sciences of the United States of America, *100*(4), 2157–2162.

https://doi.org/10.1073/pnas.0337195100

Deuker, L., Bellmund, J. L., Schröder, T. N., & Doeller, C. F. (2016). An event map of memory

space in the hippocampus. *ELife*, *5*, e16534. https://doi.org/10.7554/eLife.16534

Diana, R. A., Yonelinas, A. P., & Ranganath, C. (2008). High-Resolution Multi-Voxel Pattern

Analysis of Category Selectivity in the Medial Temporal Lobes. *Hippocampus*, *18*(6), 536–

541. https://doi.org/10.1002/hipo.20433

Diana, R. A., Yonelinas, A. P., & Ranganath, C. (2012). Adaptation to cognitive context and item

information in the medial temporal lobes. *Neuropsychologia*, *50*(13), 3062–3069.

https://doi.org/10.1016/j.neuropsychologia.2012.07.035

Diana, R. A., Yonelinas, A. P., & Ranganath, C. (2013). Parahippocampal cortex activation during

context reinstatement predicts item recollection. *Journal of Experimental Psychology.*

General, *142*(4), 1287–1297. https://doi.org/10.1037/a0034029

Dimsdale-Zucker, H. R., & Ranganath, C. (2018). Representational Similarity Analyses: A

Practical Guide for Functional MRI Applications. In D. Manahan-Vaughan (Ed.),

Handbook of in Vivo Neural Plasticity Techniques. Elsevier.

Dimsdale-Zucker, H. R., Ritchey, M., Ekstrom, A. D., Yonelinas, A. P., & Ranganath, C. (2018).

CA1 and CA3 differentially support spontaneous retrieval of episodic contexts within

human hippocampal subfields. *Nature Communications*, *9*(1), 294.

https://doi.org/10.1038/s41467-017-02752-1

Dion, M. L., Sumner, J. L., & Mitchell, S. M. L. (2018). Gendered Citation Patterns across Political

Science and Social Science Methodology Fields. *Political Analysis*, *26*(3), 312–327.

https://doi.org/10.1017/pan.2018.12

Dixon, P. (2008). Models of accuracy in repeated-measures designs. *Journal of Memory and*

Language, *59*(4), 447–456. https://doi.org/10.1016/j.jml.2007.11.004

DuBrow, S., & Davachi, L. (2013). The influence of context boundaries on memory for the

sequential order of events. *Journal of Experimental Psychology. General*, *142*(4), 1277–

1286. https://doi.org/10.1037/a0034024

DuBrow, S., & Davachi, L. (2014). Temporal memory is shaped by encoding stability and

intervening item reactivation. *The Journal of Neuroscience: The Official Journal of the*

Society for Neuroscience, *34*(42), 13998–14005.

https://doi.org/10.1523/JNEUROSCI.2535-14.2014

756 DuBrow, S., Rouhani, N., Niv, Y., & Norman, K. A. (2017). Does mental context drift or shift?

757  *Current Opinion in Behavioral Sciences*, *17*(Supplement C), 141–146.

758  https://doi.org/10.1016/j.cobeha.2017.08.003

759 Dworkin, J. D., Linn, K. A., Teich, E. G., Zurn, P., Shinohara, R. T., & Bassett, D. S. (2020). The

760  extent and drivers of gender imbalance in neuroscience reference lists. *Nature*

761  *Neuroscience*, *23*(8), 918–926. https://doi.org/10.1038/s41593-020-0658-y

762 Dzulkifli, M. A., & Wilding, E. L. (2005). Electrophysiological indices of strategic episodic retrieval

763  processing. *Neuropsychologia*, *43*(8), 1152–1162.

764  https://doi.org/10.1016/j.neuropsychologia.2004.11.019

765 Eacott, M. J., & Gaffan, D. (2005). The roles of perirhinal cortex, postrhinal cortex, and the

766  fornix in memory for objects, contexts, and events in the rat. *The Quarterly Journal of*

767  *Experimental Psychology. B, Comparative and Physiological Psychology*, *58*(3–4), 202–

768  217. https://doi.org/10.1080/02724990444000203

769 Eichenbaum, H. B., Yonelinas, A. P., & Ranganath, C. (2007). The Medial Temporal Lobe and

770  Recognition Memory. *Annual Review of Neuroscience*, *30*(1), 123–152.

771  https://doi.org/10.1146/annurev.neuro.30.051606.094328

772 Estes, W. K. (1955). Statistical theory of distributional phenomena in learning. *Psychological*

773  *Review*, *62*(5), 369–377. https://doi.org/10.1037/h0046888

774 Ezzyat, Y., & Davachi, L. (2014). Similarity Breeds Proximity: Pattern Similarity within and across

775  Contexts Is Related to Later Mnemonic Judgments of Temporal Proximity. *Neuron*,

776  *81*(5), 1179–1189. https://doi.org/10.1016/j.neuron.2014.01.042

Favila, S. E., Chanales, A. J. H., & Kuhl, B. A. (2016). Experience-dependent hippocampal pattern

differentiation prevents interference during subsequent learning. *Nature*

Communications, *7*. Scopus. https://doi.org/10.1038/ncomms11066

Frankó, E., Insausti, A. M., Artacho-Pérula, E., Insausti, R., & Chavoix, C. (2014). Identification of

the human medial temporal lobe regions on magnetic resonance images: Human Medial

Temporal Lobe Landmarks. *Human Brain Mapping*, *35*(1), 248–256.

https://doi.org/10.1002/hbm.22170

Friedman, W. J. (1993). Memory for the time of past events. *Psychological Bulletin*, *113*(1), 44–

66. https://doi.org/10.1037/0033-2909.113.1.44

Fulvio, J. M., Akinnola, I., & Postle, B. R. (2021). Gender (Im)balance in Citation Practices in

Cognitive Neuroscience. *Journal of Cognitive Neuroscience*, *33*(1), 3–7.

https://doi.org/10.1162/jocn_a_01643

Grande, X., Berron, D., Horner, A. J., Bisby, J. A., Düzel, E., & Burgess, N. (2019). Holistic

Recollection via Pattern Completion Involves Hippocampal Subfield CA3. *Journal of*

Neuroscience, *39*(41), 8100–8111. https://doi.org/10.1523/JNEUROSCI.0722-19.2019

Hintzman, D. L. (2001). Judgments of frequency and recency: How they relate to reports of

subjective awareness. *Journal of Experimental Psychology: Learning, Memory, and*

Cognition, *27*(6), 1347–1358. https://doi.org/10.1037/0278-7393.27.6.1347

Hintzman, D. L. (2004). Time versus items in judgment of recency. *Memory & Cognition*, *32*(8),

1298–1304. https://doi.org/10.3758/BF03206320

Hintzman, D. L. (2005). Memory strength and recency judgments. *Psychonomic Bulletin &*

Review, *12*(5), 858–864. https://doi.org/10.3758/BF03196777

Horner, A. J., Bisby, J. A., Bush, D., Lin, W.-J., & Burgess, N. (2015). Evidence for holistic episodic

recollection via hippocampal pattern completion. *Nature Communications*, *6*, 7462.

https://doi.org/10.1038/ncomms8462

Howard, M. W., & Kahana, M. J. (2002). A Distributed Representation of Temporal Context.

Journal of Mathematical Psychology, *46*(3), 269–299.

https://doi.org/10.1006/jmps.2001.1388

Jaeger, T. F. (2008). Categorical Data Analysis: Away from ANOVAs (transformation or not) and

towards Logit Mixed Models. *Journal of Memory and Language*, *59*(4), 434–446.

https://doi.org/10.1016/j.jml.2007.11.007

Jenkins, L. J., & Ranganath, C. (2010). Prefrontal and medial temporal lobe activity at encoding

predicts temporal context memory. *The Journal of Neuroscience: The Official Journal of*

the Society for Neuroscience, *30*(46), 15558–15565.

https://doi.org/10.1523/JNEUROSCI.1337-10.2010

Jenkins, L. J., & Ranganath, C. (2016). Distinct neural mechanisms for remembering when an

event occurred. *Hippocampus*, *26*(5), 554–559. https://doi.org/10.1002/hipo.22571

Johnson, J. D., McDuff, S. G. R., Rugg, M. D., & Norman, K. A. (2009). Recollection, familiarity,

and cortical reinstatement: A multivoxel pattern analysis. *Neuron*, *63*(5), 697–708.

https://doi.org/10.1016/j.neuron.2009.08.011

Johnson, M. K., Kounios, J., & Nolde, S. F. (1997). Electrophysiological brain activity and memory

source monitoring. *NeuroReport*, *8*(5), 1317–1320.

Kahana, M. J. (1996). Associative retrieval processes in free recall. *Memory & Cognition*, *24*(1),

103–109. https://doi.org/10.3758/BF03197276

Kay, M. (2022). *tidybayes: Tidy Data and Geoms for Bayesian Models*.

https://zenodo.org/record/5823492

Kesner, R. P., & Rolls, E. T. (2015). A computational theory of hippocampal function, and tests of

the theory: New developments. *Neuroscience and Biobehavioral Reviews*, *48*, 92–147.

https://doi.org/10.1016/j.neubiorev.2014.11.009

Kim, G., Norman, K. A., & Turk-Browne, N. B. (2017). Neural Differentiation of Incorrectly

Predicted Memories. *Journal of Neuroscience*, *37*(8), 2022–2031.

https://doi.org/10.1523/JNEUROSCI.3272-16.2017

Knierim, J. J., Lee, I., & Hargreaves, E. L. (2006). Hippocampal place cells: Parallel input streams,

subregional processing, and implications for episodic memory. *Hippocampus*, *16*(9),

755–764. https://doi.org/10.1002/hipo.20203

Kriegeskorte, N., Mur, M., & Bandettini, P. (2008). Representational Similarity Analysis—

Connecting the Branches of Systems Neuroscience. *Frontiers in Systems Neuroscience*, *2*.

https://doi.org/10.3389/neuro.06.004.2008

Lacy, J. W., Yassa, M. A., Stark, S. M., Muftuler, L. T., & Stark, C. E. L. (2011). Distinct pattern

separation related transfer functions in human CA3/dentate and CA1 revealed using

high-resolution fMRI and variable mnemonic similarity. *Learning & Memory*, *18*(1), 15–

18. https://doi.org/10.1101/lm.1971111

LaRocque, K. F., Smith, M. E., Carr, V. A., Witthoft, N., Grill-Spector, K., & Wagner, A. D. (2013).

Global similarity and pattern separation in the human medial temporal lobe predict

subsequent memory. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *33*(13), 5466–5474. https://doi.org/10.1523/JNEUROSCI.4293-12.2013

Lavenex, P., & Amaral, D. G. (2000). Hippocampal-neocortical interaction: A hierarchy of

associativity. *Hippocampus*, *10*(4), 420–430. https://doi.org/10.1002/1098-

1063(2000)10:4<420::AID-HIPO8>3.0.CO;2-5

Levy, W. B. (1996). A sequence predicting CA3 is a flexible associator that learns and uses

context to solve hippocampal-like tasks. *Hippocampus*, *6*(6), 579–590.

https://doi.org/10.1002/(SICI)1098-1063(1996)6:6<579::AID-HIPO3>3.0.CO;2-C

Lewandowsky, S., & Murdock, B. B. (1989). Memory for serial order. - PsycNET. *Psychological*

Review, *96*(1), 25–57.

Libby, L. A., Ekstrom, A. D., Ragland, J. D., & Ranganath, C. (2012). Differential connectivity of

perirhinal and parahippocampal cortices within human hippocampal subregions

revealed by high-resolution functional imaging. *The Journal of Neuroscience: The Official*

Journal of the Society for Neuroscience, *32*(19), 6550–6560.

https://doi.org/10.1523/JNEUROSCI.3711-11.2012

Libby, L. A., Reagh, Z. M., Bouffard, N. R., Ragland, J. D., & Ranganath, C. (2018). The

Hippocampus Generalizes across Memories that Share Item and Context Information.

Journal of Cognitive Neuroscience, 1–12. https://doi.org/10.1162/jocn_a_01345

Lisman, J. E. (1999). Relating Hippocampal Circuitry to Function. *Neuron*, *22*(2), 233–242.

https://doi.org/10.1016/S0896-6273(00)81085-5

Liu, K. Y., Gould, R. L., Coulson, M. C., Ward, E. V., & Howard, R. J. (2015). Tests of pattern

separation and pattern completion in humans—A systematic review. *Hippocampus*.

https://doi.org/10.1002/hipo.22561

Lohnas, L. J., Polyn, S. M., & Kahana, M. J. (2015). Expanding the scope of memory search:

Modeling intralist and interlist effects in free recall. *Psychological Review*, *122*(2), 337–

363. https://doi.org/10.1037/a0039036

Maliniak, D., Powers, R., & Walter, B. F. (2013). The Gender Citation Gap in International

Relations. *International Organization*, *67*(4), 889–922.

https://doi.org/10.1017/S0020818313000209

Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S., & Leutgeb, J. K. (2015). Hippocampal CA2

activity patterns change over time to a larger extent than between spatial contexts.

Neuron, *85*(1), 190–201. https://doi.org/10.1016/j.neuron.2014.12.001

Manning, J. R., Polyn, S. M., Baltuch, G. H., Litt, B., & Kahana, M. J. (2011). Oscillatory patterns

in temporal lobe reveal context reinstatement during memory search. *Proceedings of*

the National Academy of Sciences, *108*(31), 12893–12897.

https://doi.org/10.1073/pnas.1015174108

Marr, D. (1971). Simple memory: A theory for archicortex. *Philosophical Transactions of the*

Royal Society of London. Series B, Biological Sciences, *262*(841), 23–81.

Mazaika, P. K., Whitfield-Gabrieli, S., & Cooper, J. C. (2005). *Detection and repair of transient*

artifacts in fMRI data [Poster]. Human Brain Mapping.

McDuff, S. G. R., Frankel, H. C., & Norman, K. A. (2009). Multivoxel Pattern Analysis Reveals

Increased Memory Targeting and Reduced Use of Retrieved Details during Single-

Agenda Source Monitoring. *Journal of Neuroscience*, *29*(2), 508–516.

https://doi.org/10.1523/JNEUROSCI.3587-08.2009

McKenzie, S., Frank, A. J., Kinsky, N. R., Porter, B., Rivière, P. D., & Eichenbaum, H. (2014).

Hippocampal representation of related and opposing memories develop within distinct,

hierarchically organized neural schemas. *Neuron*, *83*(1), 202–215.

https://doi.org/10.1016/j.neuron.2014.05.019

Milivojevic, B., Vicente-Grabovetsky, A., & Doeller, C. F. (2015). Insight Reconfigures

Hippocampal-Prefrontal Memories. *Current Biology*, *25*(7), 821–830.

https://doi.org/10.1016/j.cub.2015.01.033

Mitchell, S. M., Lange, S., & Brus, H. (2013). Gendered Citation Patterns in International

Relations Journals. *International Studies Perspectives*, *14*(4), 485–492.

https://doi.org/10.1111/insp.12026

Montchal, M. E., Reagh, Z. M., & Yassa, M. A. (2019). Precise temporal memories are supported

by the lateral entorhinal cortex in humans. *Nature Neuroscience*, *22*(2), 284.

https://doi.org/10.1038/s41593-018-0303-1

Mumford, J. A., & Poldrack, R. A. (2007). Modeling group fMRI data. *Social Cognitive and*

Affective Neuroscience, *2*(3), 251–257. https://doi.org/10.1093/scan/nsm019

Mumford, J. A., Turner, B. O., Ashby, F. G., & Poldrack, R. A. (2012). Deconvolving BOLD

activation in event-related designs for multivoxel pattern classification analyses.

NeuroImage, *59*(3), 2636–2643. https://doi.org/10.1016/j.neuroimage.2011.08.076

Nielson, D. M., Smith, T. A., Sreekumar, V., Dennis, S., & Sederberg, P. B. (2015). Human

hippocampus represents space and time during retrieval of real-world memories.

Proceedings of the National Academy of Sciences, *112*(35), 11078–11083.

https://doi.org/10.1073/pnas.1507104112

Nolde, S. F., Johnson, M. K., & D'Esposito, M. (1998). Left prefrontal activation during episodic

remembering: An event-related fMRI study. *NeuroReport*, *9*(15), 3509–3514.

Norman, K. A., Detre, G., & Polyn, S. M. (2008). Computational models of episodic memory. In

The Cambridge handbook of computational psychology (pp. 189–225). Cambridge

University Press. https://doi.org/10.1017/CBO9780511816772.011

Norman, K. A., & O'Reilly, R. C. (2003). Modeling hippocampal and neocortical contributions to

recognition memory: A complementary-learning-systems approach. *Psychological*

Review, *110*(4), 611–646. https://doi.org/10.1037/0033-295X.110.4.611

O'Keefe, J., & Nadel, L. (1978). *The hippocampus as a cognitive map*. Clarendon Press ; Oxford

University Press.

O'Reilly, R. C., & McClelland, J. L. (1994). Hippocampal conjunctive encoding, storage, and

recall: Avoiding a trade-off. *Hippocampus*, *4*(6), 661–682.

https://doi.org/10.1002/hipo.450040605

Palombo, D. J., Di Lascio, J. M., Howard, M. W., & Verfaellie, M. (2019). Medial Temporal Lobe

Amnesia Is Associated with a Deficit in Recovering Temporal Context. *Journal of*

Cognitive Neuroscience, *31*(2), 236–248. https://doi.org/10.1162/jocn_a_01344

Pidgeon, L. M., & Morcom, A. M. (2016). Cortical pattern separation and item-specific memory

encoding. *Neuropsychologia*, *85*, 256–271. Scopus.

https://doi.org/10.1016/j.neuropsychologia.2016.03.026

Poldrack, R. A., & Mumford, J. A. (2009). Independence in ROI analysis: Where is the voodoo?

Social Cognitive and Affective Neuroscience, *4*(2), 208–213.

https://doi.org/10.1093/scan/nsp011

Polyn, S. M., Kragel, J. E., Morton, N. W., McCluey, J. D., & Cohen, Z. D. (2012). The neural

dynamics of task context in free recall. *Neuropsychologia*, *50*(4), 447–457.

https://doi.org/10.1016/j.neuropsychologia.2011.08.025

Polyn, S. M., Norman, K. A., & Kahana, M. J. (2009). A context maintenance and retrieval model

of organizational processes in free recall. *Psychological Review*, *116*(1), 129–156.

https://doi.org/10.1037/a0014420

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but

systematic correlations in functional connectivity MRI networks arise from subject

motion. *NeuroImage*, *59*(3), 2142–2154.

https://doi.org/10.1016/j.neuroimage.2011.10.018

Ranganath, C. (2010). Binding Items and Contexts: The Cognitive Neuroscience of Episodic

Memory. *Current Directions in Psychological Science*, *19*(3), 131–137.

https://doi.org/10.1177/0963721410368805

Ranganath, C., Yonelinas, A. P., Cohen, M. X., Dy, C. J., Tom, S. M., & D'Esposito, M. (2004).

Dissociable correlates of recollection and familiarity within the medial temporal lobes.

Neuropsychologia, *42*(1), 2–13.

Reagh, Z. M., Watabe, J. M., Ly, M., Murray, E., & Yassa, M. A. (2014). Dissociated Signals in

Human Dentate Gyrus and CA3 Predict Different Facets of Recognition Memory. *Journal

of Neuroscience*, *34*(40), 13301–13313. https://doi.org/10.1523/JNEUROSCI.2779-

14.2014

Ritchey, M., Montchal, M. E., Yonelinas, A. P., & Ranganath, C. (2015). Delay-dependent

contributions of medial temporal lobe regions to episodic memory retrieval. *ELife*, *4*,

e05025. https://doi.org/10.7554/eLife.05025

Ritvo, V. J. H., Turk-Browne, N. B., & Norman, K. A. (2019). Nonmonotonic Plasticity: How

Memory Retrieval Drives Learning. *Trends in Cognitive Sciences*.

https://doi.org/10.1016/j.tics.2019.06.007

Rolls, E. T., & Kesner, R. P. (2006). A computational theory of hippocampal function, and

empirical tests of the theory. *Progress in Neurobiology*, *79*(1), 1–48.

https://doi.org/10.1016/j.pneurobio.2006.04.005

Schapiro, A. C., Rogers, T. T., Cordova, N. I., Turk-Browne, N. B., & Botvinick, M. M. (2013).

Neural representations of events arise from temporal community structure. *Nature

Neuroscience*, *16*(4), 486–492. https://doi.org/10.1038/nn.3331

Schapiro, A. C., Turk-Browne, N. B., Botvinick, M. M., & Norman, K. A. (2017). Complementary

learning systems within the hippocampus: A neural network modelling approach to

reconciling episodic memory with statistical learning. *Phil. Trans. R. Soc. B*, *372*(1711),

20160049. https://doi.org/10.1098/rstb.2016.0049

Schlichting, M. L., Mumford, J. A., & Preston, A. R. (2015). Learning-related representational

changes reveal dissociable integration and separation signatures in the hippocampus

and prefrontal cortex. *Nature Communications*, *6*, 8151.

https://doi.org/10.1038/ncomms9151

Scoville, W. B., & Milner, B. (1957). Loss of recent memory after bilateral hippocampal lesions.

Journal of Neurology, Neurosurgery, and Psychiatry, *20*(1), 11–21.

Sederberg, P. B., Howard, M. W., & Kahana, M. J. (2008). A context-based theory of recency and

contiguity in free recall. *Psychological Review*, *115*(4), 893–912.

https://doi.org/10.1037/a0013396

Singmann, H., & Kellen, D. (2019). An introduction to mixed models for experimental

psychology. In D. H. Spieler & E. Schumacher, *New Methods in Neuroscience and*

Cognitive Psychology. Psychology Press.

Sood, G., & Laohaprapanon, S. (2018). Predicting Race and Ethnicity From the Sequence of

Characters in a Name. *ArXiv:1805.02109 [Stat]*. http://arxiv.org/abs/1805.02109

Staresina, B. P., & Davachi, L. (2008). Selective and shared contributions of the hippocampus

and perirhinal cortex to episodic item and associative encoding. *Journal of Cognitive*

Neuroscience, *20*(8), 1478–1489. https://doi.org/10.1162/jocn.2008.20104

Stokes, J., Kyle, C., & Ekstrom, A. D. (2015). Complementary roles of human hippocampal

subfields in differentiation and integration of spatial context. *Journal of Cognitive*

Neuroscience, *27*(3), 546–559. https://doi.org/10.1162/jocn_a_00736

Suthana, N. A., Donix, M., Wozny, D. R., Bazih, A., Jones, M., Heidemann, R. M., Trampel, R.,

Ekstrom, A. D., Scharf, M., Knowlton, B., Turner, R., & Bookheimer, S. Y. (2015). High-

resolution 7T fMRI of Human Hippocampal Subfields during Associative Learning.

Journal of Cognitive Neuroscience, *27*(6), 1194–1206.

https://doi.org/10.1162/jocn_a_00772

Suthana, N. A., Ekstrom, A., Moshirvaziri, S., Knowlton, B., & Bookheimer, S. (2011).

Dissociations Within Human Hippocampal Subregions During Encoding and Retrieval of

Spatial Information. *Hippocampus*, *21*(7), 694–701. https://doi.org/10.1002/hipo.20833

Treves, A., & Rolls, E. T. (1994). Computational analysis of the role of the hippocampus in

994   memory. *Hippocampus*, *4*(3), 374–391. https://doi.org/10.1002/hipo.450040319

995   Tulving, E. (1972). Episodic and semantic memory. In E. Tulving & W. Donaldson (Eds.),

996   *Organization of memory* (pp. 381–403). Academic Press.

997   Tulving, E. (1983). *Elements of episodic memory*. Oxford University Press.

998   Tulving, E. (1984). Précis of Elements of episodic memory. *Behavioral and Brain Sciences*, *7*(02),

999   223–238. https://doi.org/10.1017/S0140525X0004440X

1000   Turkkan, N., & Pham-Gia, T. (1993). Computation of the highest posterior density interval in

1001   bayesian analysis. *Journal of Statistical Computation and Simulation*, *44*(3–4), 243–250.

1002   https://doi.org/10.1080/00949659308811461

1003   Vargha–Khadem, F., Gadian, D. G., Watkins, K. E., Connelly, A., Paesschen, W. V., & Mishkin, M.

1004   (1997). Differential Effects of Early Hippocampal Pathology on Episodic and Semantic

1005   Memory. *Science*, *277*(5324), 376–380. https://doi.org/10.1126/science.277.5324.376

1006   Viskontas, I. V., Carr, V. A., Engel, S. A., & Knowlton, B. J. (2009). The neural correlates of

1007   recollection: Hippocampal activation declines as episodic memory fades. *Hippocampus*,

1008   *19*(3), 265–272. https://doi.org/10.1002/hipo.20503

1009   Wallenstein, G. V., Eichenbaum, H., & Hasselmo, M. E. (1998). The hippocampus as an

1010   associator of discontiguous events. *Trends in Neurosciences*, *21*(8), 317–323.

1011   Wang, X., Dworkin, J. D., Zhou, D., Stiso, J., Falk, E. B., Bassett, D. S., Zurn, P., & Lydon-Staley, D.

1012   M. (2021). Gendered citation practices in the field of communication. *Annals of the*

1013   *International Communication Association*, *45*(2), 134–153.

1014   https://doi.org/10.1080/23808985.2021.1960180

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller,

K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the

Tidyverse. *Journal of Open Source Software*, *4*(43), 1686.

https://doi.org/10.21105/joss.01686

Witter, M. P., Wouterlood, F. G., Naber, P. A., & Van Haeften, T. (2000). Anatomical

Organization of the Parahippocampal-Hippocampal Network. *Annals of the New York*

Academy of Sciences, *911*(1), 1–24. https://doi.org/10.1111/j.1749-

6632.2000.tb06716.x

Yassa, M. A., Lacy, J. W., Stark, S. M., Albert, M. S., Gallagher, M., & Stark, C. E. L. (2011).

Pattern Separation Deficits Associated With Increased Hippocampal CA3 and Dentate

Gyrus Activity in Nondemented Older Adults. *Hippocampus*, *21*(9), 968–979.

https://doi.org/10.1002/hipo.20808

Yassa, M. A., & Stark, C. E. L. (2011). Pattern separation in the hippocampus. *Trends in*

Neurosciences, *34*(10), 515–525. https://doi.org/10.1016/j.tins.2011.06.006

Yushkevich, P. A., Amaral, R. S. C., Augustinack, J. C., Bender, A. R., Bernstein, J. D., Boccardi, M.,

Bocchetta, M., Burggren, A. C., Carr, V. A., Chakravarty, M. M., Chételat, G., Daugherty,

A. M., Davachi, L., Ding, S.-L., Ekstrom, A., Geerlings, M. I., Hassan, A., Huang, Y., Iglesias,

J. E., … Hippocampal Subfields Group (HSG). (2015). Quantitative comparison of 21

protocols for labeling hippocampal subfields and parahippocampal subregions in in vivo

MRI: Towards a harmonized segmentation protocol. *NeuroImage*, *111*, 526–541.

https://doi.org/10.1016/j.neuroimage.2015.01.004

Yushkevich, P. A., Wang, H., Pluta, J., Das, S. R., Craige, C., Avants, B. B., Weiner, M. W., &

Mueller, S. (2010). Nearly Automatic Segmentation of Hippocampal Subfields in In Vivo

Focal T2-Weighted MRI. *NeuroImage*, *53*(4), 1208–1224.

https://doi.org/10.1016/j.neuroimage.2010.06.040

Zacks, J. M., & Swallow, K. M. (2007). EVENT SEGMENTATION. *Current Directions in*

Psychological Science, *16*(2), 80–84. https://doi.org/10.1111/j.1467-8721.2007.00480.x

Zacks, J. M., Tversky, B., & Iyer, G. (2001). Perceiving, remembering, and communicating

structure in events. *Journal of Experimental Psychology. General*, *130*(1), 29–58.

https://doi.org/10.1037/0096-3445.130.1.29

Zeineh, M. M., Engel, S. A., Thompson, P. M., & Bookheimer, S. Y. (2003). Dynamics of the

Hippocampus During Encoding and Retrieval of Face-Name Pairs. *Science*, *299*(5606),

577–580. https://doi.org/10.1126/science.1077775

Zheng, L., Gao, Z., McAvan, A. S., Isham, E. A., & Ekstrom, A. D. (2021). Partially overlapping

spatial environments trigger reinstatement in hippocampus and schema representations

in prefrontal cortex. *Nature Communications*, *12*(1), 6231.

https://doi.org/10.1038/s41467-021-26560-w

Zheng, Y., Liu, X. L., Nishiyama, S., Ranganath, C., & O'Reilly, R. C. (2021). *Correcting the Hebbian*

Mistake: Toward a Fully Error-Driven Hippocampus (p. 2021.10.29.466546).

https://doi.org/10.1101/2021.10.29.466546

Zhou, D. (2022). *CleanBib* [Jupyter Notebook]. https://github.com/dalejn/cleanBib (Original

work published 2020)

Zhou, D., Cornblath, E. J., Stiso, J., Teich, E. G., Dworkin, J. D., Blevins, A. S., & Bassett, D. S.

(2020). *Gender Diversity Statement and Code Notebook v1.0*. Zenodo.

https://doi.org/10.5281/zenodo.3672110

1061