1   **Broadening the ecology of fear: non-lethal effects arise from diverse responses to**
2   **predation and parasitism**
3

4   D.R. Daversa[1,2]*, R.F. Hechinger[3], E. Madin[4], A. Fenton[1], A.I. Dell[2,5,6] E. Ritchie[7], J. Rohr[8], V.H.W.
5   Rudolf[9] **
6
[1] Institute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom
[2]National Great Rivers Research and Education Centre (NGRREC), East Alton, IL, 62024, USA
[3]Scripps Institution of Oceanography, University of California, San Diego, United States of
America
[4]Hawai'i Institute of Marine Biology, University of Hawai'i, Kane'ohe, HI 96744, United States of
America
[5]Department of Biology, Washington University of St. Louis, St. Louis, MO 63130
[6]Dept of Biology, Saint Louis University, Saint Louis, MO 63130
[7]School of Life and Environmental Sciences, Centre for Integrative Ecology (Burwood campus),
Deakin University, Geelong, Victoria 3220, Australia
[8]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA
[9]BioSciences, Rice Univ., Houston, TX, USA
19

20   **Key Words:** Natural enemies, Sub-lethal effects, Trait-mediated effects, Trophic interactions,
21   Animal behaviour, Community ecology, Food webs, Risk effects
22

23   **Running Title:** Non-lethal effects of predation and parasitism
24

25   **Author Contributions:** This review was broadly conceived during a 2016 Gordon Conference
26   group meeting on predator-prey interactions. DRD and EM organized initial planning and idea
27   development. DRD performed the systematic review and meta-analysis with input from the
28   other authors. DRD, KL, RH, and AF constructed the framework and drafted the manuscript. All
29   authors contributed to the revision process and ideas expressed in the final piece.
30

31   **Data accessibility statement:** The data and code supporting the review and data analyis are
32   archived in Dryad and will be made publicly available should the paper be accepted after peer
33   review.
34

***Corresponding author:** David Daversa; ddaversa@gmail.com
****Not all authors are listed.  The full author list will be listed once the paper is published after

peer review.

**Abstract**

The ecology of fear demonstrates how prey responses to avoid predation cause non-lethal

effects at all ecological scales. Parasites also elicit defensive responses in hosts with associated

non-lethal effects, which raises the longstanding, yet unresolved question of how non-lethal

effects of parasites compare with those of predators. We developed a framework for

systematically answering this question for all types of predator and parasite systems. Our

framework predicts that trait responses and their non-lethal effects should be strongest from

predators and parasites that do not kill individuals to feed on them, but which nevertheless

damage fitness. Analysing trait response data on amphibians, which have been well-studied for

this area of research, showed that individuals generally responded more directly to short-term

predation risks than to parasitism. Apart from studies using amphibians, there have been few

direct comparisons of responses to predation and parasitism, and none have incorporated

responses to micropredators, parasitoids, or parasitic castrators, or examined their long-term

consequences. Addressing these and other data gaps highlighted by our general framework can

advance the field toward understanding how non-lethal effects shape real food webs, which

contain multiple predator and parasite species.

**Introduction**

Few people watch the movie "Jaws" without experiencing a fear that can make them

think twice before swimming in the ocean. Wildlife are no different. Animals avoid foraging

where predators are or have been, reduce activity levels, seek refuges, and exhibit other

responses in 'fear' of being eaten by predators [1,2]. Fear responses exemplify non-lethal effects

of predators on prey traits, and are more generally called 'trait responses'. Trait responses can

trigger a broad range of additional non-lethal effects, including 'trait-mediated effects' that

range from reduced individual fitness to trophic cascades [3–5] that can destabilize

communities [6]. Wolves, for example, frighten elk away from exposed foraging grounds into

sheltered habitats with less nutritious vegetation, which then reduces elk birth rates [7] and

alters vegetation structure [8]. By eliciting trait responses, predators impact prey, and wider

communities, without direct killing.

Recent surges in disease outbreaks make clear that parasites also elicit trait responses

with non-lethal effects. To reduce infection risk, hosts may avoid infected conspecifics [9–11],

defend against infectious propagule attack [12], or avoid risky areas, such as faeces

representing a hot spot of undetectable nematode eggs [13–15]. Basic emotions like 'disgust'

[15–18] and the age-old cliché "avoid like the plague" suggest that parasite avoidance is

interwoven in our own history as much as is our fear of predators.

Trait responses to parasites are not confined to avoidance tactics like those made in fear

or disgust. Because parasitism is not immediately lethal, hosts can respond, and incur non-lethal

effects, after a successful attack by a parasite [19–21]. Immune responses are one of myriad

host responses made after infection that can have non-lethal effects [22]. Post-attack trait

responses elicited by parasites has led some to hypothesize that parasites cause stronger

cumulative non-lethal effects than predators [5,23]. A formal framework for comparing the full

range of trait responses to predators and parasites is lacking, however, leaving uncertainty over

the similarities and differences in how non-lethal effects accumulate across different predator-

prey and host-parasite interactions.

Here, we compare trait responses to predation and parasitism, considering how

variation in their frequency and strength may drive differences in how non-lethal effects accrue

in prey and hosts. Building on recent conceptual developments [21,24], our goal is to establish a

quantitative foundation for estimating non-lethal effects in real ecosystems, which contain

multiple predator and parasite species. We draw from consumer-resource theory to construct a

general framework for estimating trait responses that accounts for the variable consumer

strategies of predators and parasites. Unlike current conceptual frameworks for fear and

disgust, we deconstruct interactions into sequential phases to consider trait responses before,

during, and after an attack, thus allowing us to compare and contrast the complete diversity of

predatory and parasitic consumers. We use this framework to form specific predictions

regarding how trait responses, and their non-lethal effects, should manifest from interactions

with different types of predators and parasites. For brevity, we focus on adaptive trait

responses for defence against predation or parasitism, but our framework can also consider

maladaptive responses, such as those occurring from parasite manipulation [25]. We then

report on a systematic review of the comparative literature on trait responses. We also

analysed data on larval amphibians, the most common animal group used by the reviewed

studies, to assess the empirical evidence for our predictions. We conclude by highlighting

unresolved questions about trait responses and their non-lethal effects on communities and

ecosystems.

**A general trait-response framework for examining non-lethal effects**

*Existing trait response frameworks*

104            The ecology of fear in predator-prey systems provides a strong, yet incomplete

foundation for examining trait responses and their impact on population and community

dynamics. 'Fear responses' denote trait responses to the risk of predation, before a predator

attack. Prey movement away from foraging habitats when predators are nearby is a well-

studied 'fear response' [1,3,26]. As Lima and Dill (1990) pointed out in their seminal framework

[1],  prey may also exhibit defensive trait responses during predator attack and even after being

captured, phases not covered within the standard domain of the 'ecology of fear'. Systematic

examination of non-lethal effects must go beyond fear to consider multiple trait responses made

throughout interactions. This becomes especially obvious when incorporating responses to

parasites.

114            In stark contrast to predation, host responses after parasite attack, that is, after

infection, can continue as parasites feed on individuals. The same is true for prey of

micropredators, which act like parasites by feeding without killing. This distinguishing

characteristic of host-parasite and micropredator-prey interactions is not factored into the Lima

and Dill (1990) framework, nor it is factored into new conceptual frameworks for disgust

[15,24], a form of parasite avoidance. Surviving while being fed on by parasites or

micropredators opens up a broad range of responses that slow or stop feeding, or otherwise

minimize its impact. Immune responses are a clear example of host responses made during

parasite feeding that do not occur in predator-prey systems. Immune responses and other trait

responses during parasite and micropredator feeding also cause non-lethal effects to

individuals and broader ecosystems [5,21], making their inclusion in systematic examinations

of non-lethal effects essential.

*A general trait response framework*

Although trait responses take diverse forms, all responses may be included in our

proposed framework (Fig. 1). The framework is informed by a general consumer-resource

model [27]. The general consumer-resource model features a single mathematical logic for

characterizing the dynamics of all types of host-parasite and predator-prey systems (and other

consumer-resource interactions, e.g. decomposer-carcass; plant-pollinator) by breaking down

interactions temporally, during which individuals transition through discrete sequential states

(Fig. S1). Prey and hosts transition through four states – susceptible, exposed, ingested,

resistant (i.e. invulnerable; Fig. S1). For clarity, here we use the term 'consumed' in place of

'ingested'. At the same time, predators and parasites transition through three states - questing,

attacking, consuming (Fig. S1). Individuals transition between states following sequential

biological processes: contact, attack failure and success, and feeding (further details of the

model are provided in the supplementary material).

We distinguish three trait response classes in our framework that coincide with the

biological processes driving predator-prey and host-parasite interactions: **avoid contact,**

**counter attack,** and **combat consumption** (Fig. 1). All three responses can reduce the fitness

costs of being consumed, albeit through distinct mechanisms and to varying degrees depending

on their efficacy (i.e. the degree of change they cause to the interaction sequence). *Susceptible*

prey and hosts may **avoid contact** with *questing* consumers before an attack. Effective

avoidance reduces the rate that *questing* predators and parasites transition to *attacking*, and

prey and hosts benefit from transitioning more slowly from *susceptible* to *exposed* states (Table

S3). Prey and hosts that become *exposed* may **counter attack** to increase the probability that

attacks fail. Countering-attack includes "fight or flight" responses, like hares sprinting to

burrows when being chased by lynx, or tadpoles rapidly and erratically swimming when being

attacked by trematode cercariae [12]. Finally, prey and hosts that are *consumed* (i.e. eaten or

infected) and remain alive may **combat consumption**. Combating consumption shortens or

slows predator and parasite feeding rates [i.e. 'resistance' in parasitology [19]], or lessens

damage from the feeding [i.e. 'tolerance' in parasitology [20]]. Responses that reduce feeding

include behaviours like social grooming by primates [28] and adaptive immune responses to

parasitism [22]. Increasing tissue repair and protecting high-risk areas of the body from

parasite feeding, as tadpoles do in response to trematodes [12], exemplify responses that

combat consumption by reducing damage, without affecting feeding rates. Collectively, the

combination of three trait responses with two consumer types results in six general pathways

for non-lethal effects to arise.

*Factors shaping the strength, duration, and frequency of trait responses*

With the full range of trait responses classified and integrated into consumer-resource

dynamics, we can now consider the conditions that determine which responses predators and

parasites are likely to elicit, as well as how strong and frequent they are likely to be. The extent

to which prey and hosts avoid, counter, and combat consumers depends first and foremost on

their basic physical and sensory abilities [29]. For instance, tadpoles cannot physically leave

ponds when predators are present. They may, however, reduce activity to avoid contact [30].

Individuals must also be able to detect consumer threats to respond to them. Prey and hosts use

both visual and non-visual cues to detect predation and parasitism risk, making sensory

limitations (e.g. sight, hearing, and smell) potential constraints on trait responses [31,32].

Impediments to either risk detection or the ability to act on detected risk should reduce the

frequency of trait responses, or even preclude them.

Even when prey and hosts have the physical and sensory capacity to mount responses,

trade-offs may mediate the frequency and strength of trait responses [29]. The ecology of fear

provides ample evidence for the role of trade-offs in shaping trait responses to predators

[33,34]. Whether via reductions in foraging, reproduction, or energy levels, fitness-related costs

of mounting responses compete with the benefits of responses (i.e. the costs of not responding),

making the frequency and strength that individuals exhibit trait responses a matter of

economics [29]. An important point not recognized by the ecology of fear (or disgust) is that

trade-offs may change over different phases of any given interaction. The frequency, duration,

and strength of trait responses will therefore depend not just on their absolute costs but also on

the relative costs compared to other possible responses. Exemplified by certain host-parasite

interactions, avoiding contact may be more costly than becoming infected, potentially driving

stronger and more frequent combat responses after becoming infected. The extent to which

'fear' and 'disgust' influence prey and host populations will depend on the feasibility to counter

attacks and combat consumption. Phase-specific trade-off situations are accounted for in our

framework through functions that link responses to state-specific mortality and  fecundity rates,

which expresses responses costs.  Those costs are balanced by benefits of responses, expressed

through response impacts on state transition rates, as described in the above section.

*Predicting trait responses and non-lethal effects across different predators and parasites*

The timing, frequency, and strength of trait responses in prey and hosts will also depend

on traits of the predators and parasites, which leads to predictions for how trait responses and

non-lethal effects may vary across the different types of consumers. Predators and parasites

comprise distinct 'consumer strategies'; that is, basic ways by which individuals feed on and

impact other organisms [27,35]. One key difference involves the fitness consequences of

predation and parasitism on individual prey or hosts. Predators and most parasitoids eliminate

future reproductive success of their prey or hosts by killing them, and parasitic castrators do so

by blocking host reproduction. In contrast, feeding by most other types of parasites and

micropredators is not so deadly and does not completely eliminate future fitness gains. Given

their great risk of consumption, we predict that predators, parasitoids, and parasitic castrators

will elicit strong trait responses in their prey and hosts, resulting in overall strong non-lethal

effects.  Similar patterns may also emerge with other parasites with strong negative fitness

impacts, such as highly virulent pathogens, or the micropredators that vector them. However,

non-vectoring micropredators and less damaging parasites should elicit relatively weaker

responses and weaker associated non-lethal effects. Differences in the fitness consequences of

being consumed should then drive variation in non-lethal effects not just between predators

and parasites, but also between different types of parasites, with some parasites being more

similar to predators than to other types of parasites.

We can go further by considering the timing of prey and host death during interactions.

Predators usually kill prey before or shortly after commencing to feed. This is not the case for

most parasites and their hosts. Even parasitoids and castrators can have a substantial amount of

feeding time before hosts are killed or castrated. Recognizing this basic difference in timing of

prey and host mortality, or "reproductive death", leads to two clear predictions for how non-

lethal effects of predators and parasites differ. First, prey responses to predators will be

constrained to avoiding contact and countering attack (Fig. 1b). Second, parasites and

micropredators will elicit all three trait responses (Fig. 1c-d), resulting in more diverse non-

lethal effects on hosts than prey of predators.

Integrating the above hypotheses leads to the prediction that, all else being equal,

parasites and micropredators that entail high fitness costs should cause the strongest overall

non-lethal effects from trait responses. This includes parasitic castrators, parasitoids, highly

damaging pathogens, and the micropredators that vector them. The high fitness costs of feeding

by these consumers should drive strong avoidance and counter-responses, resulting in strong

non-lethal effects from those responses. Yet, by feeding on living organisms, these consumers

also permit combat responses as a source of non-lethal effects, as evidenced in many hosts of

parasitoids, parasitic castrators, and other parasites [5,36]. Collectively, we expect non-lethal

effects from this triad of trait responses to supersede those arising solely from 'fear' responses

(i.e. avoidance of contact and counter of attack) to predators. Exceptions may exist, such as

some parasitoids that paralyze hosts during the initial attack, which can make hosts incapable of

combatting consumption [36]. On average, however, our framework indicates that non-lethal

effects will be strongest not solely from consumers that impose the highest risk of death, but

rather from those that impose such strong costs while keeping the consumed host or prey alive.

**Studies to measure trait responses to predation and parasitism**

We reviewed the literature that compares trait responses to predation and parasitism to

assess support for our predictions. Specially, we compiled studies that directly compared trait

responses to both predation and parasitism risk in a single resource species (see supporting

material for details of literature search). The vast majority of studies we found measured trait

responses in larval frogs (i.e. 'tadpoles', N = 106 entries across 13 studies, Table S1).

Behavioural traits were most common, with activity level being the most reported trait (Table

S1, Table S2). We did not find studies that measured immunological trait responses, likely

because this is distinct to host-parasite systems. The studies spanned the following consumer

strategies: solitary predators, trophically transmitted parasites, typical parasites, and pathogens

(Table S1); in no cases were responses to parasitoids, parasitic castrators, micropredators, or

social predators considered. Predator-induced trait responses were only measured during

*questing* predator states, representing avoidance of contact, whereas measurements of parasite-

induced responses included all three states: *questing* (N=9), *attacking* (N=5), and *consuming*

(N=24). The limited data we found constrained our ability to make comparisons, although some

general patterns did emerge.

Analysis of the data on tadpoles, the main animal group used as prey and hosts revealed

considerable variation in the magnitude and direction of tadpole responses to predators (Fig.

2a), parasites (Fig. 2b), and their combination (Fig. 2c). However, on average and across all

consumer and resource states, predator-induced trait responses were stronger than parasite-

induced trait responses (Fig. 3a, Table S4). These patterns were evident after controlling for

consumer state (comparing questing predators to questing parasites) (Fig. 3b), though they

were weaker (Table S4), likely owing to lower power of the data subset. Distinguishing between

trait response types (avoid, counter, combat) revealed that tadpoles did respond to parasites,

but only by combatting parasites post-infection and at lower magnitudes than their responses

to avoid predator contact (Fig. 3b, Table S4). Tadpoles also responded to the simultaneous

presence of predators and parasites on average, and at similar magnitudes to their responses to

predators alone (Fig. 3a). The strong tadpole responses to predators primarily represented

reductions in activity levels (Fig. S7a, Table S4), and they were most evident when measuring

group-level responses as opposed to individual-level responses (Fig. S7b, Table S4). Across the

host-parasite interactions studied, responses did not differ between pathogens and trophically-

transmitted parasites, the two parasite strategies investigated (Table S3).

**Discussion**

Our framework reveals that non-lethal effects from predation and parasitism are a

function of diverse and potentially interacting responses by prey and hosts, exhibited at

different time points in consumer-resource interactions. The current literature on non-lethal

effects provides a very limited portrayal of trait responses, which restrained our ability to test

the predictions that emerged from the general framework. Current comparative data comprise

brief snapshots of interactions that do not fully capture the temporal dynamics of trait

responses (and their non-lethal effects). Longitudinal data that tracks individuals through all

interaction phases could provide insight into how non-lethal effects accrue from multiple trait

responses. Longitudinal data could also detect interactive effects between trait responses. For

example, hosts that invest heavily in immune defences may exhibit weaker avoidance of contact

with parasites, particularly if avoidance conflicts with feeding, reproducing, and other fitness-

related activities. The growing literature on 'fear' and 'disgust' of parasites, which focuses

exclusively on certain forms of parasite avoidance, will benefit by being contextualized within

the broader suite of trait responses available to hosts.

Trait response data on tadpoles, although limited, underscore the importance of

distinguishing nuances in the timing of trait responses. Pooling all trait response measurements

suggested that parasites generally did not elicit changes in tadpole activity levels as did

predators. However, accounting for response timing revealed significant changes in activity

levels that were confined to post-infection periods of interaction. Adult amphibians show

similar tendencies to combat infections rather than avoid them [37]. Combat responses appear

to be a primary trait response to parasitism in this group of animals, whose role in non-lethal

effects can most fully be estimated by treating the responses as separate from those made

earlier in interactions.

The post-infection combat responses apparent in tadpoles are particularly notable

considering that immune responses were not factored in. Immune responses are a very

common form of combatting parasitism that can be exhibited for prolonged periods. The

combined non-lethal effects of tadpole parasites were therefore likely much stronger than the

analysed data suggest. Additionally, non-lethal effects of parasitism could have occurred from

host phenotypic changes caused by parasite manipulation [25,38], and even directly from

parasite feeding independent of defensive responses. For instance, general energy drain or

direct tissue damage caused by parasite infection can substantially impact host performance

[39,40]. The diverse sources of non-lethal effects of parasites not relevant to prey collectively

could have rivalled in magnitude the stronger predator avoidance observed in tadpoles.

Accounting for distinct predator and parasite strategies led to trait response predictions

that did not align with the predator versus parasite dichotomy. Pinpointing the sources of

variation in non-lethal effects therefore demands a functional approach rather than a taxonomic

approach to distinguishing predators and parasites [35]. Parasitoids, and parasitic castrators in

particular, could offer rewarding insights because they share functional characteristics with

both predators and parasites. Avoidance, counter, and combat responses to parasitoids are

well-documented [36], but how their frequency and strength compare with responses to

predators and less debilitating parasites is unknown. The latter can also be said for parasitic

castrators. Given the high fitness consequences of infection from parasitoids and castrators,

together with infections that are not immediately lethal, their combined non-lethal effects may

very well be the strongest of all predator and parasite functional groups.

Regardless of how individuals respond to predators and parasites alone, risks of

predation and parasitism in real ecosystems rarely occur in isolation. Future research could

apply our framework to investigate the additive and interactive non-lethal effects of

simultaneous exposure to predators and parasites. Our analysis of the tadpole data suggests

that responses to simultaneous exposure are non-additive, perhaps owing to the prioritization

of responses to the more severe threat (Fig. S5). Although not a focus of this review, evidence

for increasing predation of parasitized prey [41–43], and increasing parasitism in predator-rich

environments [44], provide further indication that predators and parasites interact to impose

non-lethal effects in non-additive ways. Fewer studies have considered single responses that

defend against both predators and parasites. Nevertheless, there were several cases where

tadpole responses to predators and parasites responses were in the same direction (i.e. a

reduction in the trait expression). Trait responses that effectively deter both predators and

parasites may mitigate the non-lethal effects incurred from the essential task of defending

oneself against being eaten.

**Conclusion**

Whether through fear or through infection, consumers elicit costly trait responses in

their resources that give rise to non-lethal effects at the level of individuals, with consequences

for communities and ecosystems more generally. A general consumer-resource model helped us

to develop a framework for systematically comparing trait responses to various types of

predators and parasites. Different types of predators and parasites should elicit different trait

responses, and therefore, have different non-lethal effects, given differences in consumer

strategies that influence when and how strongly they impact prey and hosts. However, many

predator and parasite strategies have not yet been tested comparatively. Expanding research of

non-lethal effects to regularly consider different predator and parasite strategies sets the

foundation for exploring how non-lethal effects manifest in the multi-dimensional food webs

found in real ecosystems.

**Acknowledgements**

We thank Gordon Research Conferences and Andy Sih, chair of the 2016 meeting "Predator-

Prey Interactions", for providing the forum that inspired this work; the EGLIDE group (Amy

Pedersen, Amy Sweeny, Saudamini Venkatesan, Dishon Muloi, Alexandra Morris, Shaun Keegan

and Kayleigh Gallagher), the EEGID group at University of Liverpool (Mike Begon, Greg Hurst,

David Montagnes, Mark Viney, Steve Parratt), the Garner lab at the Zoological Society of London

(Trent Garner, Chris Owen, Lola Brooks, Stephen Price, Goncalo Rosa, Bryony Allan), and Chris

Carbone for their comments on earlier versions of the review; and Daniel Noble, Raj Whitlock,

and Wolfgang Viechtbauer for advice on the meta-analysis. Any use of trade, product, or firm

names in this publication is for descriptive purposes only and does not imply endorsement by

the U.S. Government. This manuscript benefited from NSF Ecology of Infectious Diseases grant

(OCE-1115965) and a grant from the Natural Environment Research Council UK

(NE/N009800/1).

**References**

1. Lima SL, Dill LM. 1990 Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology* **68**, 619–640.

2. Dröge E, Creel S, Becker MS, M'soka J. 2017 Risky times and risky places interact to affect prey behaviour. *Nature Ecology & Evolution* **1**, 1123–1128. (doi:10.1038/s41559-017-0220-9)

3. Brown JS. 1998 The ecology of fear: optimal foraging, game theory, and trophic interactions. *IFAC Proceedings Volumes* **31**, 31. (doi:10.1016/S1474-6670(17)38332-5)

4. Ritchie EG, Johnson CN. 2009 Predator interactions, mesopredator release and biodiversity conservation. *Ecology Letters* **12**, 982–998. (doi:10.1111/j.1461-0248.2009.01347.x)

5. Buck JC, Ripple WJ. 2017 Infectious agents trigger trophic cascades. *Trends in Ecology & Evolution* **32**, 681–694. (doi:10.1016/j.tree.2017.06.009)

6. Pringle RM *et al.* 2019 Predator-induced collapse of niche structure and species coexistence. *Nature* **570**, 58–64. (doi:10.1038/s41586-019-1264-6)

7. Creel S, Christianson D, Liley S, Winnie JA. 2007 Predation risk affects reproductive physiology and demography of Elk. *Science* **315**, 960–960. (doi:10.1126/science.1135918)

8. Fortin D, Beyer HL, Boyce MS, Smith DW, Duchesne T, Mao JS. 2005 Wolves influence elk

movements: behavior shapes a trophic cascade in Yellowstone National Park. *Ecology* **86**,

1320–1330.

9. Milinski M, Bakker TCM. 1990 Female sticklebacks use male coloration in mate choice and

hence avoid parasitized males. *Nature* **344**, 330.

10. Kavaliers M, Colwell DD, Braun WJ, Choleris E. 2003 Brief exposure to the odour of a

parasitized male alters the subsequent mate odour responses of female mice. *Animal*

*Behaviour* **65**, 59–68. (doi:10.1006/anbe.2002.2043)

11. Behringer DC, Butler MJ, Shields JD. 2006 Avoidance of disease by social lobsters. *Nature*

**441**, 421–421. (doi:10.1038/441421a)

12. Sears BF, Snyder PW, Rohr JR. 2013 Infection deflection: hosts control parasite location with

behaviour to improve tolerance. *Proceedings of the Royal Society B: Biological Sciences* **280**,

20130759–20130759. (doi:10.1098/rspb.2013.0759)

13. Hart BL. 1994 Behavioural defense against parasites: interaction with parasite invasiveness.

*Parasitology* **109**, S139–S151.

14. Curtis VA. 2014 Infection-avoidance behaviour in humans and other animals. *Trends in*

*Immunology* **35**, 457–464. (doi:10.1016/j.it.2014.08.006)

15. Weinstein SB, Buck JC, Young HS. 2018 A landscape of disgust. *Science* **359**, 1213–1214.

(doi:10.1126/science.aas8694)

16. Curtis V, de Barra M. 2018 The structure and function of pathogen disgust. *Philosophical*

*Transactions of the Royal Society B: Biological Sciences* **373**, 20170208.

(doi:10.1098/rstb.2017.0208)

17. Dougherty ER, Seidel DP, Carlson CJ, Spiegel O, Getz WM. 2018 Going through the motions:

incorporating movement analyses into disease research. *Ecology Letters* **21**, 588–604.

(doi:10.1111/ele.12917)

18. Tybur JM, Çınar Ç, Karinen AK, Perone P. 2018 Why do people vary in disgust? *Philosophical*

*Transactions of the Royal Society B: Biological Sciences* **373**, 20170204.

(doi:10.1098/rstb.2017.0204)

19. Rigby MC, Hechinger RF, Stevens L. 2002 Why should parasite resistance be costly? *Trends*

*in parasitology* **18**, 116–120.

20. Raberg L, Graham AL, Read AF. 2009 Decomposing health: tolerance and resistance to

parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*

**364**, 37–49. (doi:10.1098/rstb.2008.0184)

21. Buck JC. 2019 Indirect effects explain the role of parasites in ecosystems. *Trends in*

*Parasitology* (doi:10.1016/j.pt.2019.07.007)

22. Hawley DM, Altizer SM. 2011 Disease ecology meets ecological immunology: understanding

the links between organismal immunity and infection dynamics in natural populations.

*Functional Ecology* **25**, 48–60. (doi:10.1111/j.1365-2435.2010.01753.x)

23. Rohr JR, Swan A, Raffel TR, Hudson PJ. 2009 Parasites, info-disruption, and the ecology of

fear. *Oecologia* **159**, 447–454. (doi:10.1007/s00442-008-1208-6)

24. Doherty J-F, Ruehle B. 2020 An integrated landscape of fear and disgust: the evolution of

avoidance  behaviors amidst a myriad of natural enemies. *Front. Ecol. Evol.* **8**, 564343.

(doi:10.3389/fevo.2020.564343)

25. Lafferty KD, Shaw JC. 2013 Comparing mechanisms of host manipulation across host and

parasite taxa. *Journal of Experimental Biology* **216**, 56–66. (doi:10.1242/jeb.073668)

26. Creel S, Winnie J, Maxwell B, Hamlin K, Creel M. 2005 Elk alter habitat selectioin as an

antipredator response to wolves. *Ecology* **86**, 3387–3397. (doi:10.1890/05-0032)

27. Lafferty KD, DeLeo G, Briggs CJ, Dobson AP, Gross T, Kuris AM. 2015 A general consumer-

resource population model. *Science* **349**, 854–857. (doi:10.1126/science.aaa6224)

28. Hart BL, Hart LA. 2018 How mammals stay healthy in nature: the evolution of behaviours to

avoid parasites and pathogens. *Philosophical Transactions of the Royal Society B: Biological*

*Sciences* **373**, 20170205. (doi:10.1098/rstb.2017.0205)

29. Ydenberg RC, Dill LM. 1986 The economics of fleeing from predators. *Advances in the Study*

*of Behavior* **16**, 229–249.

30. Hossie T, Landolt K, Murray DL. 2017 Determinants and co-expression of anti-predator

responses in amphibian tadpoles: a meta-analysis. *Oikos* **126**. (doi:10.1111/oik.03305)

31. Kats LB, Dill LM. 1998 The scent of death: Chemosensory assessment of predation risk by

prey animals. *Écoscience* **5**, 361–394. (doi:10.1080/11956860.1998.11682468)

32. Behringer DC, Karvonen A, Bojko J. 2018 Parasite avoidance behaviours in aquatic

environments. *Phil. Trans. R. Soc. B* **373**, 20170202. (doi:10.1098/rstb.2017.0202)

33. Laundré JW. 2010 Behavioral response races, predator–prey shell games, ecology of fear,

and patch use of pumas and their ungulate prey. *Ecology* **91**, 2995–3007. (doi:10.1890/08-

2345.1)

34. Gaynor KM, Brown JS, Middleton AD, Power ME, Brashares JS. 2019 Landscapes of fear:

patial patterns of risk perception and response. *Trends in Ecology & Evolution* **34**, 355–368.

(doi:10.1016/j.tree.2019.01.004)

35. Lafferty KD, Kuris AM. 2002 Trophic strategies, animal diversity and body size. *Trends in*

*Ecology & Evolution* **17**, 507–513.

36. Abram PK, Brodeur J, Urbaneja A, Tena A. 2019 Nonreproductive effects of insect

parasitoids on their hosts. *Annu. Rev. Entomol.* **64**, 259–276. (doi:10.1146/annurev-ento-

011118-111753)

37. Daversa DR, Manica A, Bosch J, Jolles JW, Garner TWJ. 2018 Routine habitat switching alters

the likelihood and persistence of infection with a pathogenic parasite. *Functional Ecology*

**32**, 1262–1270. (doi:10.1111/1365-2435.13038)

38. Poulin R, Brodeur J, Moore J. 1994 Parasite manipulation of host behaviour: Should hosts

always lose? *Oikos* , 479–484.

39. Munger JC, Karasov WH. 1989 Sublethal parasites and host energy budgets: tapeworm

infection in white-footed mice. *Ecology* **70**, 904–921. (doi:10.2307/1941358)

40. Delahay RJ, Speakman JR, Moss R. 1995 The energetic consequences of parasitism: effects of

a developing infection of *Trichostrongylus tenuis* (Nematoda) on red grouse (*Lagopus*

*lagopus scoticus*) energy balance, body weight and condition. *Parasitology* **110**, 473.

(doi:10.1017/S0031182000064817)

41. Lafferty KD, Morris AK. 1996 Altered behavior of parasitized killifish Increases

susceptibility to predation by bird final hosts. *Ecology* **77**, 1390–1397.

(doi:10.2307/2265536)

42. Johnson PTJ, Stanton DE, Preu ER, Forshay KJ, Carpenter SR. 2006 Dining on disease: how

interactions between infection and environment affect predation risk. *Ecology* **87**, 1973–

1980. (doi:10.1890/0012-9658(2006)87[1973:DODHIB]2.0.CO;2)

43. Rae J, Murray D. 2019 Pathogen vs. predator: ranavirus exposure dampens tadpole

responses to perceived predation risk. *Oecologia* **191**, 325–334. (doi:10.1007/s00442-019-

04501-1)

44. Stephenson JF, Van Oosterhout C, Mohammed RS, Cable J. 2015 Parasites of Trinidadian

guppies: evidence for sex-and age-specific trait-mediated indirect effects of predators.

*Ecology* **96**, 489–498.

**Fig. 1. A general trait response framework and predictions. (a)** Resources can mount three sequential responses to consumers, each with distinct effects on the interaction: avoid contact, counter attack, combat consumption. Examples of each type of response are listed in italicized text. Responses may be constrained by physical and sensory limitations, as well as trade-offs against other fitness-related activities (e.g. feeding and reproducing). The general framework can be tailored to specific types of consumer-resource interactions, such as interactions between field mice and **(b)** owl predators, **(c)** biting flies and infective nematodes (red worms in faeces), or **(d)** between caterpillars and parasitoid wasps. The lack of a combat stage in **(b)** illustrates that resources rarely respond during consumption by predators.

**Fig. 2. Forest plots of effect sizes used in the meta-analysis.** The distribution of effect sizes for responses elicited by the presence of **(a)** predators, **(b)** parasites, and **(c)** their combined presence resulting from resource adjustments in activity (blue), space use (red), and morphological/physiological traits (grey). Error bars denote the 95% confidence intervals.

**Fig. 3. Relative magnitude of responses to predation vs. parasitism. (a)** The estimated mean magnitude of trait responses to predation cues (blue), parasitism cues (red), and both cues (grey). **(b)** Mean trait responses to predators (blue) and parasites (red) when distinguishing by the type of trait responses, as defined in our framework. Only avoidance responses to questing predators

486     were found in our literature review, likely owing the low probability of surviving attack or

487     consumption by predators. Responses to the combined presence of predators and parasites are

488     not shown in **(b)** because only one study with this treatment had predators and parasites in the

489     same state. Lines denote the mean response magnitudes, boxes denote the standard error of the

490     mean, and error bars denote the 95% confidence intervals.

**(a)  general trait response framework**

**(b) responses to predators**

**(c) responses to parasites and micropredators**

**(d) responses to parasitoids**

     Fig. 2

Predators

**(a)**
mean ± CI:
0.78 ± 0.35

**(c)**
mean ± CI:
0.89 ± 0.40

Parasites

**(b)**
mean ± CI:
0.21 ± 0.36

−7 −6 −5 −4 −3 −2 −1 0 1 2 3 4 5 6 7
Response magnitude (Hedge's d ± 95% CI)

● activity
● space use
● morphology/physiology

501    Fig. 3