1 **Beyond the ecology of fear: non-lethal effects of predators are strong whereas**
2 **those of parasites are diverse**
3

4 D.R. Daversa[1,2*], R.F. Hechinger[3], E. Madin[4], A. Fenton[1], *et al.*
5 [1]Institute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom
6 [2]Institute of Zoology, Zoological Society of London, London, United Kingdom
7 [3]Scripps Institution of Oceanography, University of California, San Diego, United States of America
8 [4]Hawai'i Institute of Marine Biology, University of Hawai'i, Kane'ohe, HI 96744, United States of America
9

10 **Key Words:** Natural enemies, Sub-lethal effects, Trait-mediated effects, Trophic
11 interactions, Animal behaviour, Community ecology, Food webs, Risk effects,
12

13 **Running Title:** Non-lethal responses to predation and parasitism
14
15

*Corresponding author: David Daversa, Institute of Integrative Biology, University of Liverpool,
Liverpool, United Kingdom; ddaversa@gmail.com

Abstract

Individuals can alter their behaviour and other traits to reduce threats from predators

and parasites. However, predators and parasites likely elicit different responses,

which subsequently lead to different non-lethal effects. We created a sequentially

structured framework to examine trait responses to distinct predatory and parasitic

consumers. We predicted that parasites with strong negative effects on host fitness

should act like predators and elicit strong responses before attack. We also predicted

that less damaging parasites and micropredators should elicit diverse responses

across multiple interaction stages, because their hosts and prey remain alive while

being eaten. A meta-analysis indicated that predators do tend to elicit stronger

responses than parasites before attack, whereas parasites generally elicit responses

after attack, albeit weaker than pre-attack responses to predators. Organisms exposed

simultaneously to predator and parasite cues responded similarly when exposed to

predator cues alone, suggesting that individuals prioritize anti-predator responses

over responses to less harmful parasites. Extending these findings requires

addressing knowledge gaps concerning responses to different consumer types, costs of

immune responses, and cumulative effects of repeated responses. Expanding research

beyond the predator vs. parasite dichotomy toward a broader consumer-resource

perspective will facilitate understanding of non-lethal effects in complex, multi-

trophic food webs.

Introduction

"Whenever I swim in the ocean… I feel increasingly panicky and … I must leave

the water" is a typical response to the 1975 film Jaws (Cantor 2004). As with

moviegoers, many species respond to predators by changing behaviours, physiology,

or even appearance to avoid being eaten. These non-lethal effects of predators, known

as 'trait responses', are pervasive and take many forms, such as seeking shelter (Creel

et al. 2005) or maturing faster to reach less vulnerable life stages (Raffel *et al.* 2010).

Such trait responses influence how individuals interact with the broader community,

driving 'trait-mediated effects' that range from reduced individual fitness to trophic

cascades (Werner & Peacor 2003; Ritchie & Johnson 2009; Buck & Ripple 2017) that

can destabilize communities (Pringle *et al.* 2019). Wolves, for example, frighten elk

away from exposed foraging grounds into sheltered habitats with less nutritious

vegetation, which then reduces elk birth rates (Creel *et al.* 2007) and alters vegetation

structure (Fortin *et al.* 2005). Predators can therefore impact species and communities

without directly killing prey, just as *Jaws* kept many people from swimming in the

summer of 1975.

Perhaps less well recognised is that parasites also elicit trait responses in hosts

with associated non-lethal effects. To reduce infection risk, hosts may avoid infected

conspecifics (Milinski & Bakker 1990; Kavaliers *et al.* 2003a; Behringer *et al.* 2006),

defend against infectious propagule attack (Sears *et al.* 2013), or avoid risky areas,

such as faeces representing a hot spot of undetectable nematode eggs (Hart 1994;

Curtis 2014; Weinstein *et al.* 2018). Furthermore, in stark contrast to predation,

parasitism is not immediately lethal, so hosts can also respond *after* successful

parasite attack through various physiological and behavioural responses (Rigby *et al.*

2002; Raberg *et al.* 2009; Buck 2019). For instance, a caterpillar can initiate an

immune response to prevent being killed by a parasitoid wasp (Abram *et al.* 2019).

Basic emotions like "disgust" (Curtis & de Barra 2018; Tybur *et al.* 2018; Weinstein *et*

al. 2018) and the age-old cliché "avoid like the plague" suggest that parasite avoidance

is interwoven in our own history as much as is our fear of predators. The diverse trait

responses elicited by parasites has led some to hypothesize that parasites actually

impose stronger cumulative non-lethal effects than predators (Rohr *et al.* 2009; Buck

& Ripple 2017).

In this review, we compare trait responses to predation and parasitism,

considering how they may overlap and differ. Although predators and parasites

threaten most species in natural ecosystems, trait responses to predators and

parasites have been largely studied in isolation. As a result, how trait responses to

parasites compare with trait responses to predators is still unclear. We used a general

consumer-resource model to develop hypotheses and predictions for how key life

history differences among predators and parasites, such as the number of attacks they

make in a lifetime and whether they kill organisms while eating them, should influence

the likelihood and magnitude of trait responses at different interaction stages. We then

conducted a systematic review and meta-analysis to: (a) assess the literature that

compares trait responses to different forms of predation and parasitism, (b) compare

average response magnitudes between predation and parasitism, and (c) test how

factors related to resources, consumers, and study designs influence trait responses.

We conclude by pointing to several unresolved questions concerning how non-lethal

species interactions affect community and ecosystem dynamics.

A general trait-response framework for examining non-lethal effects

Theoretical framework

Predators and parasites employ various 'consumer strategies', that is, how

individuals find, attack, and consume organisms (Lafferty & Kuris 2002; Lafferty *et al.*

2015). For example, predators have short feeding times (i.e., seconds to days) and eat

multiple organisms in a lifetime, whereas parasites feed on hosts for up to months or

even years, but die or transform after a single feeding interaction. Predators kill prey

before or while consuming them, but mosquitoes and other micropredators do not.

Although some parasites eliminate host fitness, exemplified by parasitoids and

parasitic castrators, others infect hosts without substantial fitness impacts. These and

other differences in consumer strategies likely affect how, and to what extent,

organisms respond to them (Buck 2019). To account for these differences when

predicting the non-lethal effects of predation and parasitism, we draw on consumer-

resource theory to develop a general trait-response framework that applies across

both host-parasite and predator-prey systems.

We use, as a scaffolding, the general model for consumer-resource population

dynamics developed by Lafferty *et al.* (2015), which collates the key elements of all

consumer-resource models into a single, temporally compartmentalized structure

(Fig. 1, Box 1). Briefly, predators and parasites are 'consumers', whereas their prey

and hosts are 'resources'. Interactions are broken into sequential transitions between

up to three discrete consumer states and up to four corresponding resource states

(circles in Box 1). State transitions — i.e., mortality, contact, attack failure and success,

and feeding (arrows in Box 1) — occur at various rates. Basic differences among

consumer – resource systems are incorporated with a set of binary parameters that

alter model structure (Box 1).

We derive three basic trait responses from the model: **avoid contact, counter**

attack, combat consumption (Fig. 1)[1]. This temporal sequence implies that trait

responses to minimize consumption can be driven by multiple biological mechanisms.

---

[1] These classifications are similar to "avoidance", "escape attack", and "escape capture" defined by Lima
& Dill (1990) for behavioural responses of prey to predator encounters, though we propose this
alternative terminology as a more comprehensive categorization of the diversity of trait responses
elicited by different predators and parasites.

In initial interactions with *questing* consumers, *susceptible* resources may **avoid**

contact in various ways. Avoidance serves to reduce the rate that *questing* consumers

transition to attacking, with the benefit that *susceptible* resources transition more

slowly to *exposed* states (Table 1). Avoidance responses may manifest as trait

adaptations [i.e. constitutive responses (Westra *et al.* 2015)], such as camouflaged

coloration shown by many species to reduce visibility to predators (e.g. Stevens &

Merilaita 2009), or as induced plastic traits, exemplified by herbivores moving away

from nematode-infected faeces (Hutchings *et al.* 2001; Weinstein *et al.* 2018) or wolf

scents (Creel *et al.* 2005). Resources that become *exposed* to *attacking* consumers may

counter attack to increase attack failure (Box 1). Countering attack includes "fight or

flight" responses, like hares sprinting to burrows when being chased by lynx, or

tadpoles jolting their bodies when being attacked by trematode cercariae (Sears *et al.*

2013). Finally, resources—particularly hosts of parasites—that become *ingested* may

combat consumption. Combating consumption shortens or slows consumer feeding

rates [i.e. 'resistance' in parasitology (Rigby *et al.* 2002)], or lessens the damage of

being eaten [i.e. 'tolerance' in parasitology (Raberg *et al.* 2009)]. Responses that

shorten or slow feeding include behaviours like social grooming by primates (Hart &

Hart 2018) and adaptive immune responses to parasitism (Hawley & Altizer 2011).

Increasing tissue repair and protecting high-risk areas of the body from feeding, as

tadpoles do for trematodes (Sears *et al.* 2013), are ways that resources combat

consumption by reducing damage without affecting consumer feeding rates. Whether,

and to what degree, resources combat, avoid, or counter, will depend on the different

pressures imposed by different consumers, which are modelled by altering rates of

contact, attack, and/or consumption. Distinguishing these three model-derived

resource trait responses makes it easier to compare, contrast, and make predictions

about the different ways that consumers exert non-lethal effects on resources.

Constraints on trait responses

Despite the clear benefits that trait responses provide to resources, various

constraints can limit a resource's ability to avoid, counter, and combat its consumers.

Hence, predicting the timing and magnitude of trait responses also requires

accounting for constraints on resources. Constraints can arise from basic limitations

on detecting consumers and mounting responses, or from trade-offs. First, resources

can mount responses only if they can detect consumer threats. Resources use visual

and non-visual cues to detect predation and parasitism risk, making sensory

limitations - e.g., sight, hearing, and smell – a potential constraint on trait responses.

Second, resources must possess the morphology, physiology, and energy level to

mount specific responses to consumers. For instance, tadpoles cannot physically leave

ponds when predators are present as can adult amphibians, so they may instead

reduce activity levels to avoid consumer contact (Hossie *et al.* 2017). Third, because

trait responses can compete with essential activities like feeding, reproducing, or

maintenance (Dröge *et al.* 2017; Hart & Hart 2018), trade-offs can also constrain

resource responses. For example, moose can afford to avoid wolves more in summer

than in winter when food is scarce (Oates *et al.* 2019). These three constraints likely

interact to jointly influence response timing and magnitudes. For example, food

scarcity may interact with moose body size or background nutritional levels to

determine their ability to avoid contact with wolves. Considering these constraints,

along with the potential benefits of response, leads to several hypotheses and

associated predictions for how resources should respond to different consumer

threats.

Hypotheses and predictions concerning trait responses against predators and parasites

Hypothesis 1: Severe fitness consequences of consumption will favour strong

trait responses—and strong non-lethal effects—at early interaction stages.

Severe fitness consequences of consumption make combat responses very risky,

placing a premium on mounting defensive responses at earlier interaction stages

when consumers are questing or attacking. This leads to the intuitive prediction that

predators should elicit strong avoidance and counter responses because successful

predation leads to death. Perhaps less intuitively, this hypothesis also leads to the

prediction that parasitoids, which regularly kill their hosts (Lafferty & Kuris 2002;

Abram *et al.* 2019), and parasitic castrators, which reproductively kill their hosts, will

be similar to predators in eliciting strong responses at the two early interaction stages.

Other types of parasites with strong negative fitness impacts, such as certain

pathogens, may place similar pressures on hosts by producing detrimental infections.

In contrast, micropredators and less harmful parasites should elicit the weakest

avoidance and counter responses of all consumer types.

Hypothesis 2: Resources that remain alive while being eaten can implement and

concentrate responses—and incur non-lethal effects—during consumption.

In stark contrast to most prey of predators, hosts are alive while parasites feed, and

prey are alive when micropredators feed. Those resources can therefore mount

combat responses while being ingested. This leads to the prediction that, compared to

predators, many parasites and micropredators will evoke more types of responses

with a more even distribution among the three interaction stages. Because hosts of

castrators and many parasitoids remain alive while being eaten, these predictions

apply for these types of parasites as well. A less obvious prediction is that the ability

to combat consumption may lead to resources concentrating responses in the third

interaction stage for parasites – including parasitoid and parasitic castrators - and

micropredators, particularly in cases where constraints or trade-offs limit avoidance

and combat responses.

Hypothesis 3: Detection ability determines trait response timing and magnitude.

Regardless of the fitness consequences of consumption, an inability to detect questing

predators and parasites will preclude mounting avoidance responses. Under this

hypothesis, the strength of responses should increase with the ease of detecting

consumers. For instance, to the extent that visual detection is important to elicit

defensive responses, we predict that resources avoid questing predators more than

questing micropredators and parasites, because the generally larger size of questing

predators relative to questing micropredators and parasites make them easier to see.

Exceptions will exist, however; some parasites have large searching stages (e.g., some

hymenopteran wasp parasitoids), which would permit easy detection and subsequent

avoidance by hosts. Further, certain predators, like some ambush predators and filter

feeders, are actually *not* detectable while questing, but only during attack, which

precludes avoiding contact but favours countering attack. Hence, this hypothesis does

not predict consistent differences among consumer strategies (e.g. predators,

parasites, micropredators, etc.), and could perhaps be most strongly tested using

predator and parasite species that span a range of detectability.

These three hypotheses are not mutually exclusive and can be integrated to

predict how trait responses vary among the different types of predation and

parasitism in realistic situations (Fig. 1b-d). For instance, foraging mice will avoid

questing owls by hiding in burrows (Fig. 1b). Burrow use may vary with flyover

frequency, which predicts mouse-owl contact rates. Burrow use is traded-off against

the cost of reducing mouse feeding, and consequently likely depends on mouse

nourishment. Mice may also detect infective nematode eggs in feeding fields and then

avoid contact by moving to other locations (Fig. 1c). Avoiding contact with nematodes

is likely constrained by mouse ability to detect eggs and should also depend on

movement costs and food availability in new locations. Unlike avoiding owls, the

strength with which mice avoid nematodes may also depend on their ability to combat

infection (consumption by the nematode), perhaps by mounting an immune response.

Effective immune function would favour mice avoiding substantial avoidance costs

(e.g., if mice avoid rich feeding areas with nematodes for poor feeding areas lacking

nematodes), which would concentrate non-lethal effects in consumption stages. All

three responses may also be expected from caterpillar hosts of wasp parasitoids, with

the difference that high risk of death via successful consumption may favour a shift to

earlier avoidance and counter responses (Fig. 1d). What remains unclear is whether

the broader range of responses, and potential concentration of combat responses, to

less virulent parasites like nematodes together might, when combined, rival the

magnitude of the avoidance and counter responses to consumers with severe fitness

consequences. To compare the overall magnitude of trait responses to parasites and

predators, we conducted a systematic review of and meta-analysis on available

literature.

Systematic review and meta-analysis of trait responses to predation and

parasitism

We systematically reviewed studies that measured the magnitude of trait

responses elicited by predators and parasites. We then performed a meta-analysis on

the compiled data to test the predictions established above. Our broad goal with the

meta-analysis was to assess how the type and magnitude of trait responses vary by

consumer strategy as defined by Lafferty and Kuris (2002): solitary predators,

trophically transmitted parasites, typical parasites, and pathogens, parasitoids,

parasitic castrators, micropredators, or social predators. Many studies have measured

trait responses to predators and parasites alone, but we limited our review to studies

that measured comparable responses to both a predator and a parasite for the same

resource species. To focus on trait responses conferring defence, we did not consider

trait changes originating from general parasite pathology, or from parasite adaptive

manipulation of host traits (Poulin *et al.* 1994; Moore 2002; Lafferty & Shaw 2013),

though our framework can accommodate such effects.

Detailed information about data collection, extraction and analyses are

available in the Supplementary Material. Briefly, we compiled 129 entries from 15

studies, which included 44 predator-prey interactions, 44 host-parasite interactions,

and 41 simultaneous interactions with predators and parasites (Table S1). Although

studies measured several morphological, behavioural, and physiological responses,

behavioural traits were most common, with activity level being the most reported trait

(Fig. S1). No studies fitting our criteria measured physiological or immunological trait

responses. The studies spanned the following consumer strategies: solitary predators,

trophically transmitted parasites, typical parasites, and pathogens (Table S1). We

therefore could not consider responses to parasitoids, parasitic castrators,

micropredators, or social predators. Hereafter, we broadly distinguish between

predators (i.e. solitary predators) and parasites (i.e. trophically-transmitted, typical, or

pathogens). Predator-induced trait responses were only measured during the *questing*

predator state (Fig. S1), whereas measurements of parasite-induced responses

included questing (10), attacking (9), and consuming (25) states (Fig. S1). There were

77 entries for individual-level responses and 49 entries for group-level responses.

We calculated the standardized mean difference (Hedge's *d*) (Koricheva *et al.*

2013) from included studies as the measure of trait response magnitudes, whereby

positive effects denoted reductions in trait values (e.g., reduced activity level or mass),

except in measures of space use that measured time in a refuge or distance from a

consumer cue (e.g. positive effect would mean an increase in refuge use). We reversed

the sign of these values so that positive effect sizes would denote reductions in use of

risky habitats, indicative of defence (see Table S1 for further details). Because studies

often included a treatment containing both a predator and parasite cue, we also

estimated the magnitude of trait responses to the combined presence of predators and

parasites. We did not have predictions for how these responses would compare to

those made to predators or parasites by themselves. We also considered the following

factors related to consumers, resources, and experimental designs that may have

influenced response magnitudes: a) the type of trait measured to quantify responses

(trait type), b) whether studies used a trait value of individuals or proportions of

individuals in group that exhibited the focal trait (analysis scale); c) whether

responses were elicited under the physical presence of the consumer, or solely by

indirect cues such as used media or ingested conspecifics (consumer presence), d) the

genus and species of the consumer and resource, and e) whether the consumer was in

a questing, attacking or consuming state (Box 1) when the trait response was

measured. We also assessed consumer state and consumer strategy effects using only

the parasite data because data on responses to predators were limited to one

consumer state (questing) and strategy (solitary predator).

Results

We found considerable variation in response magnitude and direction to

predators (Fig. 2a), parasites (Fig. 2b), and their combination (Fig. 2c). However,

although individual parasite-induced effects were sometimes just as strong as

predator-induced effects (Fig. 2), on average and across all stages, predator-based trait

responses were stronger than parasite-based trait responses (Table S2, Fig. 3a). These

patterns were also evident after controlling for consumer state (i.e., questing

predators vs. questing parasites) (Table S2, Fig. 3b). Nevertheless, distinguishing

between parasite states (questing, attacking, or consuming) revealed that parasites, on

average, did elicit responses, but only while they were consuming (i.e. infecting) their

hosts (Table S2. Fig. 3b). The simultaneous presence of predators and parasites also

elicited responses on average, and they were similar in magnitude to trait responses

elicited by predators alone ($z = 0.10$, $p = 0.476$; Fig. 3a). Predators and the

simultaneous presence of predators and parasites elicited reductions in activity but,

on average, did not influence space use or morphological/physiological traits (Table

S2, Fig. 3c). Whether traits were measured at the individual level or group level (e.g.

proportions) influenced response magnitudes, with group-level responses being

stronger (Table S2, Fig. d). Responses were not contingent on the consumers being

physically present; indirect cues of the consumers elicited similar responses (Table

S2). Across the host-parasite interactions studied, responses did not depend on the

specific strategy of parasites (pathogens, trophically transmitted parasites, or typical

parasites; Table S2), and there was insufficient replication to consider how consumer

or resource taxon influenced responses. For that reason, these results mostly pertain

to amphibians as resources (Table S1).

Discussion

We used a general consumer-resource model to construct a framework that can

be broadly applied across many predator-prey and host-parasite systems to predict

trait responses and non-lethal effects. The framework identified plausible

mechanisms driving the timing and magnitude of trait responses, including the fitness

consequences associated with being eaten, whether individuals are alive while being

eaten, and the ease of detecting consumers. From these hypothesized mechanisms, we

generated testable predictions regarding how trait responses should differ between

predator-prey and host-parasite interactions. We generally predict severe fitness

consequences of predator consumption to drive strong avoidance and counter

responses in prey before any contact is made. We also predict that, in general, host

responses to parasites are weaker than those of prey, but also more diverse; a range of

behavioural, morphological, and physiological responses can be made throughout all

interaction stages (Rigby *et al.* 2002; Raberg *et al.* 2009), and potentially concentrated

to combat consumption. These general predictions were supported by our meta-

analysis on the existing literature to directly compare trait responses to predatory

versus parasitic consumers. However, the meta-analysis data predominantly related

to a rather narrow taxonomic range of predators and parasites, and several limitations

of the included studies (Box 2) suggest ways forward to corroborate and expand the

results.

Several overlooked distinctions in how organisms respond to parasites likely led

to underestimation of overall response magnitudes to parasitism in our meta-analysis.

None of the included studies measured individual responses in more than one

interaction stage, even though parasites commonly evoke responses in all interaction

stages. Longitudinal data on individual responses to multiple consumer states will

more comprehensively quantify trait response magnitudes, and may even reveal

interactive effects between trait responses (Boxes 2,3). Given that immunological

responses may be the most common type of host response to parasitism, non-lethal

effects arising from host combat responses, in particular, are likely to be much

stronger than our meta-analysis suggests. Additionally, non-lethal effects of parasitism

can also arise from host phenotypic changes caused by parasite manipulation (Poulin

et al. 1994; Lafferty & Shaw 2013), and even directly from parasite feeding

independent of defensive responses. For instance, general energy drain (Munger &

Karasov 1989; Delahay *et al.* 1995) or direct tissue damage caused by parasite

infection can substantially impact host performance (Palstra *et al.* 2007). Such

pathological effects are not driven by a host response, but nevertheless represent non-

lethal effects. These distinct types of non-lethal effects of parasitism could collectively

rival in magnitude the stronger predator avoidance that we observed. To test this

hypothesis would require longitudinal data on predator responses as well, which our

review indicates is also lacking in the literature. Although we expect combat

responses to predation to be rare, there are exceptional cases of prey defending

themselves while being ingested, particularly for some slow predators that do not kill

their prey before consuming them (e.g., sea stars eating mussels). In such cases,

predators could also evoke combat responses.

Considering distinct predator and parasite consumer strategies led to more

comprehensive trait response predictions that did not completely align with the

predator versus parasite dichotomy. However, our systematic review revealed a

paucity of literature to develop this more holistic approach at understanding trait

responses, suggesting a fruitful area of future research on non-lethal effects. For

example, we predict parasitoids and parasitic castrators should act like predators to

elicit strong pre-contact responses, but share with other parasites the ability to elicit

combat responses during consumption. Micropredators, by not killing their prey when

feeding, should act like parasites to elicit responses across all interaction stages.

Studies that focus solely on micropredators (Kavaliers *et al.* 2003b, 2005) and

parasitoids (Abram *et al.* 2019), do find that resources avoid, counter, and combat

these consumers, though we still lack direct comparisons of the magnitudes of these

responses with similar responses to predators and typical parasites. Furthermore

there remains much to be known regarding the extent to which detectability

constrains or enables responses. Testing these more specific, yet meaningful, trait

response predictions requires distinguishing not just between predators and

parasites, but also different types of predators and parasites. A synthesis of trait

responses from multiple single-consumer studies may permit tests of these

predictions, but will come with the potential expense of error from inconsistent study

designs. To minimize such error, we encourage experiments that directly compare

trait responses to a broader range of consumer strategies and resource species.

Regardless of how individual resources respond to predators and parasites

alone, risks of predation and parasitism in the wild rarely occur in isolation. The non-

additive predator and parasite effects that we observed in the meta-analysis may be

indicative of priority effects, whereby the first type of exposure elicits the stronger

response, though we could not test for this. Additionally, predators may frequently

interfere with parasite responses by imposing stronger immediate threats to survival.

Trade-offs between defences against predation and parasitism, such as adjustments in

activity by tadpoles (Koprivnikar & Urichuk 2017) and shoaling behaviour by guppies

(Stephenson *et al.* 2015), could also explain why it might be difficult to respond

effectively to different simultaneous threats. However, although trade-offs between

predator and parasite defences have been considered previously (Orlofske *et al.* 2012;

Stephenson *et al.* 2015; Koprivnikar & Urichuk 2017), our meta-analysis highlights

that anti-predator and anti-parasite responses may also be complementary in that

predators and parasites can elicit similar responses, and most of the significant

predator and parasite responses reported by studies were in the same direction (i.e. a

reduction in the trait expression). A well-demonstrated example comes from tadpoles

(Box 3), which reduce activity levels (Marino *et al.* 2014; Preston *et al.* 2014; Gallagher

et al. 2019), increase refuge use (Han *et al.* 2011) and reduce time in foraging habitats

(Koprivnikar & Penalva 2015) when either exposed to predators or infected with

parasites. When responses deter both predators and parasites, avoiding predation

may also inadvertently aid in the avoidance of parasites.

Conclusion

Whether through fear or through infection, consumers elicit costly trait

responses in their resources that give rise to non-lethal effects at the level of

individuals, communities, and ecosystems. A general consumer-resource model

helped to develop a framework to predict trait responses to various consumer types.

Owing to differences in consumer strategies that influence when and how strongly

they impact resources, we expect different trait responses to different types of

predators and parasites, and therefore, different non-lethal effects. However, many

consumer strategies have not yet been tested in a comparative fashion. Expanding

research of non-lethal interactions beyond the predator vs. parasite dichotomy toward

a broader consumer-resource perspective sets the foundation for exploring how non-

lethal effects manifest in the complex, multi-trophic food webs found in real

ecosystems.

Acknowledgements

We thank Gordon Research Conferences and Andy Sih, chair of the 2016 meeting

"Predator-Prey Interactions", for providing the forum that inspired this work; the

EGLIDE group (Amy Pedersen, Amy Sweeny, Saudamini Venkatesan, Dishon Muloi,

Alexandra Morris, Shaun Keegan and Kayleigh Gallagher), the EEGID group at

University of Liverpool (Mike Begon, Greg Hurst, David Montagnes, Mark Viney, Steve

Parratt), the Garner lab at the Zoological Society of London (Trent Garner, Chris Owen,

Stephen Price, Goncalo Rosa, Bryony Allan), and Chris Carbone for their comments on

earlier versions of the review; and Daniel Noble, Raj Whitlock, and Wolfgang

Viechtbauer for advice on the meta-analysis. Any use of trade, product, or firm names

in this publication is for descriptive purposes only and does not imply endorsement by

the U.S. Government. This manuscript benefited from NSF Ecology of Infectious

Diseases grant (OCE-1115965) and a grant from the Natural Environment Research

Council UK (NE/N009800/1).

References

Abram, P.K., Brodeur, J., Urbaneja, A. & Tena, A. (2019). Nonreproductive effects of

insect parasitoids on their hosts. *Annu. Rev. Entomol.*, 64, 259–276.

Behringer, D.C., Butler, M.J. & Shields, J.D. (2006). Avoidance of disease by social

lobsters. *Nature*, 441, 421–421.

Buck, J.C. (2019). Indirect effects explain the role of parasites in ecosystems. *Trends in*

Parasitology.

Buck, J.C. & Ripple, W.J. (2017). Infectious agents trigger trophic cascades. *Trends in*

Ecology & Evolution, 32, 681–694.

Cantor, J. (2004). "I'll Never Have a Clown in My House" -- Why movie horror lives on.

Poetics Today, 25, 283–304.

Creel, S., Christianson, D., Liley, S. & Winnie, J.A. (2007). Predation risk affects

reproductive physiology and demography of Elk. *Science*, 315, 960–960.

Creel, S., Winnie, J., Maxwell, B., Hamlin, K. & Creel, M. (2005). Elk alter habitat

selectioin as an antipredator response to wolves. *Ecology*, 86, 3387–3397.

Curtis, V. & de Barra, M. (2018). The structure and function of pathogen disgust.

Philosophical Transactions of the Royal Society B: Biological Sciences, 373,

20170208.

Curtis, V.A. (2014). Infection-avoidance behaviour in humans and other animals.

Trends in Immunology, 35, 457–464.

Daversa, D.R., Manica, A., Bosch, J., Jolles, J.W. & Garner, T.W.J. (2018). Routine habitat

switching alters the likelihood and persistence of infection with a pathogenic

parasite. *Functional Ecology*, 32, 1262–1270.

Delahay, R.J., Speakman, J.R. & Moss, R. (1995). The energetic consequences of

parasitism: effects of a developing infection of *Trichostrongylus tenuis*

(Nematoda) on red grouse (*Lagopus lagopus scoticus*) energy balance, body

weight and condition. *Parasitology*, 110, 473.

Dröge, E., Creel, S., Becker, M.S. & M'soka, J. (2017). Risky times and risky places

interact to affect prey behaviour. *Nature Ecology & Evolution*, 1, 1123–1128.

Fortin, D., Beyer, H.L., Boyce, M.S., Smith, D.W., Duchesne, T. & Mao, J.S. (2005). Wolves

influence elk movements: behavior shapes a trophic cascade in Yellowstone

National Park. *Ecology*, 86, 1320–1330.

Gallagher, S.J., Tornabene, B.J., DeBlieux, T.S., Pochini, K.M., Chislock, M.F., Compton,

Z.A., *et al.* (2019). Healthy but smaller herds: Predators reduce pathogen

transmission in an amphibian assemblage. *J Anim Ecol*, 1365-2656.13042.

Han, B.A., Searle, C.L. & Blaustein, A.R. (2011). Effects of an infectious fungus,

Batrachochytrium dendrobatidis, on amphibian predator-prey interactions.

PLoS ONE, 6, e16675.

Hart, B.L. (1994). Behavioural defense against parasites: interaction with parasite

invasiveness. *Parasitology*, 109, S139–S151.

Hart, B.L. & Hart, L.A. (2018). How mammals stay healthy in nature: the evolution of

behaviours to avoid parasites and pathogens. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, 373, 20170205.

Hawley, D.M. & Altizer, S.M. (2011). Disease ecology meets ecological immunology:

understanding the links between organismal immunity and infection dynamics

in natural populations: Disease ecology meets ecological immunology.

Functional Ecology, 25, 48–60.

Hossie, T., Landolt, K. & Murray, D.L. (2017). Determinants and co-expression of anti-

predator responses in amphibian tadpoles: a meta-analysis. *Oikos*, 126.

Hutchings, M.R., Gordon, I.J., Kyriazakis, I. & Jackson, F. (2001). Sheep avoidance of

faeces-contaminated patches leads to a trade-off between intake rate of forage

and parasitism in subsequent foraging decisions. *Animal Behaviour*, 62, 955–

964.

Kavaliers, M., Colwell, D.D., Braun, W.J. & Choleris, E. (2003a). Brief exposure to the

odour of a parasitized male alters the subsequent mate odour responses of

female mice. *Animal Behaviour*, 65, 59–68.

Kavaliers, M., Colwell, D.D. & Choleris, E. (2003b). Learning to fear and cope with a

natural stressor: individually and socially acquired corticosterone and

avoidance responses to biting flies. *Hormones and Behavior*, 43, 99–107.

Kavaliers, M., Colwell, D.D. & Choleris, E. (2005). Kinship, familiarity and social status

modulate social learning about ?micropredators? (biting flies) in deer mice.

Behavioral Ecology and Sociobiology, 58, 60–71.

Koprivnikar, J. & Penalva, L. (2015). Lesser of two evils? foraging choices in response

to threats of predation and parasitism. *PLOS ONE*, 10, e0116569.

Koprivnikar, J. & Urichuk, T.M.Y. (2017). Time-lagged effect of predators on tadpole

behaviour and parasite infection. *Biology Letters*, 13, 20170440.

Koricheva, J., Gurevitch, J. & Mengersen, K. (2013). *Handbook of Meta-analysis in*

Ecology and Evolution. Princeton University Press.

Lafferty, K.D., DeLeo, G., Briggs, C.J., Dobson, A.P., Gross, T. & Kuris, A.M. (2015). A

general consumer-resource population model. *Science*, 349, 854–857.

Lafferty, K.D. & Kuris, A.M. (2002). Trophic strategies, animal diversity and body size.

Trends in Ecology & Evolution, 17, 507–513.

Lafferty, K.D. & Shaw, J.C. (2013). Comparing mechanisms of host manipulation across

host and parasite taxa. *Journal of Experimental Biology*, 216, 56–66.

Lima, S.L. & Dill, L.M. (1990). Behavioral decisions made under the risk of predation: a

review and prospectus. *Canadian Journal of Zoology*, 68, 619–640.

Marino, J.A., Holland, M.P. & Middlemis Maher, J. (2014). Predators and trematode

parasites jointly affect larval anuran functional traits and corticosterone levels.

Oikos, 123, 451–460.

Milinski, M. & Bakker, T.C.M. (1990). Female sticklebacks use male coloration in mate

choice and hence avoid parasitized males. *Nature*, 344, 330.

Moore, J. (2002). *Parasites and the Behavior of Animals*. Oxford University Press.

Munger, J.C. & Karasov, W.H. (1989). Sublethal parasites and host energy budgets:

tapeworm infection in white-footed mice. *Ecology*, 70, 904–921.

Oates, B.A., Merkle, J.A., Kauffman, M.J., Dewey, S.R., Jimenez, M.D., Vartanian, J.M., *et al.*

(2019). Antipredator response diminishes during periods of resource deficit for

a large herbivore. *Ecology*, 100, e02618.

Orlofske, S.A., Jadin, R.C., Preston, D.L. & Johnson, P.T. (2012). Parasite transmission in

complex communities: predators and alternative hosts alter pathogenic

infections in amphibians. *Ecology*, 93, 1247–1253.

Palstra, A.P., Heppener, D.F.M., van Ginneken, V.J.T., Székely, C. & van den Thillart,

G.E.E.J.M. (2007). Swimming performance of silver eels is severely impaired by

the swim-bladder parasite Anguillicola crassus. *Journal of Experimental Marine*

Biology and Ecology, 352, 244–256.

Poulin, R., Brodeur, J. & Moore, J. (1994). Parasite manipulation of host behaviour:

Should hosts always lose? *Oikos*, 479–484.

Preston, D.L., Boland, C.E., Hoverman, J.T. & Johnson, P.T.J. (2014). Natural enemy

ecology: comparing the effects of predation risk, infection risk and disease on

host behaviour. *Functional Ecology*, 28, 1472–1481.

Pringle, R.M., Kartzinel, T.R., Palmer, T.M., Thurman, T.J., Fox-Dobbs, K., Xu, C.C.Y., *et al.*

(2019). Predator-induced collapse of niche structure and species coexistence.

Nature, 570, 58–64.

Raberg, L., Graham, A.L. & Read, A.F. (2009). Decomposing health: tolerance and

resistance to parasites in animals. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 364, 37–49.

Raffel, T.R., Hoverman, J.T., Halstead, N.T., Michel, P.J. & Rohr, J.R. (2010). Parasitism in

a community context: trait-mediated interactions with competition and

predation. *Ecology*, 91, 1900–1907.

Relyea, R.A. & Werner, E.E. (1999). Quantifying the relation between predator-induced

behavior and growth performance in larval anurans. *Ecology*, 80, 2117.

Rigby, M.C., Hechinger, R.F. & Stevens, L. (2002). Why should parasite resistance be

costly? *Trends in parasitology*, 18, 116–120.

Ritchie, E.G. & Johnson, C.N. (2009). Predator interactions, mesopredator release and

biodiversity conservation. *Ecology Letters*, 12, 982–998.

Rohr, J.R., Swan, A., Raffel, T.R. & Hudson, P.J. (2009). Parasites, info-disruption, and

the ecology of fear. *Oecologia*, 159, 447–454.

Sears, B.F., Snyder, P.W. & Rohr, J.R. (2013). Infection deflection: hosts control parasite

location with behaviour to improve tolerance. *Proceedings of the Royal Society*

B: Biological Sciences, 280, 20130759–20130759.

Stephenson, J.F., Van Oosterhout, C., Mohammed, R.S. & Cable, J. (2015). Parasites of

Trinidadian guppies: evidence for sex-and age-specific trait-mediated indirect

effects of predators. *Ecology*, 96, 489–498.

Stevens, M. & Merilaita, S. (2009). Animal camouflage: current issues and new

perspectives. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, 364, 423–427.

Tybur, J.M., Çınar, Ç., Karinen, A.K. & Perone, P. (2018). Why do people vary in disgust?

Philosophical Transactions of the Royal Society B: Biological Sciences, 373,

20170204.

Van Buskirk, J. (2001). Specific induced responses to different predator species in

anuran larvae. *Journal of Evolutionary Biology*, 14, 482–489.

Weinstein, S.B., Buck, J.C. & Young, H.S. (2018). A landscape of disgust. *Science*, 359,

1213–1214.

Werner, E.E. & Peacor, S.D. (2003). A review of trait-mediated indirect interactions in

ecological communities. *Ecology*, 84, 1083–1100.

Westra, E.R., van Houte, S., Oyesiku-Blakemore, S., Makin, B., Broniewski, J.M., Best, A.,

et al. (2015). Parasite Exposure Drives Selective Evolution of Constitutive

versus Inducible Defense. *Current Biology*, 25, 1043–1049.

(a) general trait response framework for consumer-resource interactions

Consumer state	questing	attacking	consuming

Avoid Contact
e.g. refuge use, camouflage, inactivity

benefits: reduce contact with consumers

constraints: trade-offs with feeding and reproduction, detection

Counter attack
e.g. sprinting, erratic movement

benefits: increase attack failure

constraints: energy levels, morphology and physiology

Combat Consumption
e.g. immune activation, habitat switching, tissue repair

benefits: shorted or slowed feeding, reduced damage from feeding

constraints: trade-offs with physiological maintenance, energy levels

Resource State	susceptible/resistant	exposed	being ingested
	time available to avoid	time available to counter	time available to combat

(b) responses to predators

(c) responses to parasites and micropredators

(d) responses to parasitoids

Fig. 1. A general trait response framework and predictions (a) Resources can respond to consumers to reduce contact rates, attack rates, and feeding rates, but might be constrained by physical and sensory limitations, as well as trade-offs against other fitness-related activities (e.g. feeding and reproducing). The general framework can be tailored to specific types of consumer-resource interactions, such as interactions between field mice and **(b)** owl predators, **(c)** biting flies and infective

577   nematodes (red worms in faeces), or **(d)** between caterpillars and parasitoid wasps.

578   The opacity of the circles denotes the predicted strength of responses. The lack of a

579   combat stage in **(b)** illustrates that resources rarely repond during consumption by

580   predators.

Fig. 2. Forest plots of studies used in the meta-analysis. The distribution of effect

sizes for responses elicited by the presence of **(a)** predators, **(b)** parasites, and **(c)**

their combined presence resulting from resource adjustments in behavioural (grey)

587   and morphological/physiological traits (blue). Error bars denote the 95% confidence

588   intervals.

Fig. 3. Relative magnitude of responses to predation vs. parasitism. (a) The

estimated mean magnitude of trait responses to predation cues (blue), parasitism cues

(red), and both cues (grey). **(b)** Mean trait responses to predators (blue) and parasites

(red) during each consumer state comprising the interaction. Only pre-attack

responses to questing predators were found in our literature review, likely owing the

low probability of surviving attack or consumption by predators. Responses to the

combined presence of predators and parasites are not shown because only one study

with this treatment had predators and parasites in the same state. **(c, d)** Estimated

mean trait response magnitudes to predation cues (blue), parasitism cues (red), and

both cues (grey), broken down by **(c)** the type of trait measured and **(d)** the scale at

which trait responses measured. Lines denote the mean response magnitudes, boxes

denote the standard error of the mean, and error bars denote the 95% confidence

intervals.

Resource Response	Resource transition	Consumer transition	processes affected	Examples	Consumer Type	Potential costs to resource	Costs to consumer
Avoid Contact	↑S-R	↓Q-A	Contact	burrow use when predators are present; vaccination from parasitic infection	All	reduced food intake and reproduction	Hunger
	↓S-E	↓Q-A	Contact	camoflaged morphology of prey, leaving habitats with cues of parasite infestation	All	reduced food intake and reproduction	energy loss, hunger
Counter attack	↑E-S	↑A-Q	Attack failure / Resource Recovery S	Fleeing from approaching predators	Predator, micropredator	energy loss or allocation trade-offs	energy loss, hunger
	↑E-S	↑A death	Consumer death / Resource recovery to S	Deflecting parasitic propagules or micropredators; venemous biting of predators	All	energy loss or allocation trade-offs	Death
	↑E-R	↑A-Q	Attack failure / Resource recovery to S	fleeing approaching predators via dispersal to areas outside of consumer distribution	Predator	energy loss or allocation trade-offs	Energy loss, hunger
	↑E-R	↑A death	Consumer death / Resource recovery to S	Developing antibodies after deflecting parasitic propagules	All	energy loss or allocation trade-offs	Death, hunger
Combat consumption	↑I-S	↑C-Q	Consumption ended / Resource recovery to S	swatting, jolted body movement, posionous flesh	Micropredator	energy loss or allocation trade-offs	Failure,
	↑I-S	↑C death	Consumer Death / Resource recovery to S	Grooming, immune activation, habitat switching	Micropredator, parasite	energy loss or allocation trade-offs, reduced food intake and reproduction	Death
	↑I-R	↑C-Q	Consumption ended / Resource recovery to S	dispersal to areas outside of consumer distribution	Micropredator	energy loss or allocation trade-offs	Failure, hunger
	↑I-R	↑C death	consumer death / Resource recovery to S	Sterile immunity	Pathogens	energy loss or allocation trade-offs	Death, hunger

Table 1. Trait responses and their consequences for consumer-resource

dynamics. The three types of trait responses that predators and parasites may elicit

(Column 1) have distinct effects on interaction dynamics (columns 2-3), in part

because each response affects specific biological processes (Column 3). For columns 2

and 3: S = susceptible, E = exposed, I = ingested, R = resistant, Q = questing, A =

attacking, C = consuming, as per Lafferty et al. 2015. Arrows denote when responses

increase transition rates and down arrows denote when trait responses decrease

transition rates.

618
619

Box 1. Overview of the general consumer resource model.

The Lafferty *et al.* (2015) model helps identify and compare different consumer-

resource interactions using a singular and formalized mathematical logic. During

interactions the consumer can be in one of three states: "questing", "attacking", or

"consuming". The resource, here being a prey or host, moves through a corresponding

four states; they are "exposed" when under attack, being "ingested" while consumers

are consuming them, and can either be "susceptible" or "resistant" to new attack. State

transitions are driven by multiple biological processes, which can be represented by

simple per capita rates or complex functions. Consumers and resources transition

from questing and susceptible states, respectively, according to detection rates of each

organism and attack rates of the consumer, while they transition to consuming and

ingested states, respectively, according to attack success rates of the consumer (Fig.

1a). Handling rates determine transitions back to questing and susceptible states or

alternatively, death rates of consumers and resources following consumption. All

states have death rates that allow for specific costs to be specified. The expanded

model shown here can be formally simplified (through setting some rates to zero to

subsume states) to represent nearly all classic consumer-resource models.

The expanded model does not represent a specific consumer-resource interaction.

Instead, a set of binary parameters (switches) makes it easy to specify distinct

consumer strategies. For instance, the "fatal attack" parameter (f) defines whether

resources are dead (f = 1) or alive (f = 0) after consumption, which distinguishes

predators and parasitoids from other parasites and micropredators. The "joint death"

parameter defines whether the consumer dies if its resource dies (j = 1) or remains

living (j = 0), which distinguishes all parasites (which have an intimate relationship

with their host) from predators and micropredators. Finally, the "multiple attacks"

parameter defines whether the consumer attacks once in a lifetime (m = 0), as

parasites and parasitoids do, or more than once in a lifetime (m = 1), as

micropredators and predators do.

Box 2. Future directions for research on non-lethal effects

There remain several limitations in the comparative data on trait responses to, and

non-lethal effects of, predators and parasites. Addressing the following limitations will

lead to more comprehensive estimates of non-lethal effects in diverse, real

ecosystems:

1. Comparative experiments of trait responses that cover a broader range of

consumer-resource systems can determine how generalizable patterns are across

systems, and opens avenues to consider factors, like phylogenetics of consumer and

resource, on response magnitudes. Most comparative trait response studies identified

by our systematic review used larval amphibians and their aquatic consumers. The

experimental designs of those studies provide examples of how comparisons of trait

responses to different types of consumers can be performed with other species to test

the generality of our meta-analysis results.

2. Comparative studies that include responses to parasitic castrators,

parasitoids and micropredators are needed to provide novel tests of the

mechanisms, such as detection and fitness consequences, underlying trait responses to

consumer threats (see Discussion). Comparisons of trait responses to different

consumer types are currently limited to typical and trophically-transmitted parasites,

pathogens, and solitary predators – only three of the 10 consumer life histories found

in natural ecosystems (Lafferty & Kuris 2002; Lafferty *et al.* 2015).

3. Considering the full array of parasite non-lethal effects on hosts will be

important to truly compare predators and parasites. For instance, though immune

responses are one of the most common forms of anti-parasite defence, we were not

able to include immune responses into comparisons of trait responses to predators

and parasites. Because immune responses can be costly to initiate and maintain, such

host responses certainly cause non-lethal effects. Further, parasites may alter host

traits in ways that do not involve defensive responses, including by pathological

energy drain or tissue damage, or via adaptive host manipulation of host phenotypes.

All these effects occur during the consumption phase and will be uniquely pronounced

for parasites compared to predators.

4. Longitudinal data on prey and host responses are needed to consider how

multiple responses elicited by single consumers collectively shape the magnitude of

non-lethal effects. Longitudinal data may even reveal interactive effects between trait

responses, such that mounting one type of response affects the strength of other

responses. For example, parasites whose infections are not highly costly (e.g., many

gastrointestinal helminths) may cause hosts to prioritize combat responses to

minimize trade-offs of avoiding contact against fitness-related activities, like feeding

and reproducing (Moore 2002; Hart & Hart 2018). This occurrence may explain why

newts, for example, do not avoid infective questing parasites in breeding ponds, but

infections from those parasites drive the same individuals to leave those ponds

(Daversa *et al.* 2018). Prioritization of combat responses to parasitism provides one

explanation for our finding that average responses to parasites were strongest after

parasites commenced feeding.

5. Linking trait-response magnitudes to trait-mediated effects can provide

estimates of how the non-lethal impacts of consumers on individual traits correspond

with the their broader impacts on individual fitness, and community and ecosystem

dynamics. Future work can extend the length of prey and host monitoring to link trait

responses to individual fitness. Mesocosm and field experiments mirroring the

experimental designs of the studies in our meta-analysis can introduce primary

producers and other species in food webs into the picture, allowing associations

between response magnitudes to predators and parasites and trophic flows and

cascades to be quantified.

Box 3. Larval anurans as a case study for interactive non-lethal effects of

predation and parasitism.

The literature on trait responses to consumers is mostly about frogs. Larval anurans

(i.e. tadpoles) were the most studied resource species in our meta-analysis (Table S1),

and how tadpole behaviour responds to predation risk (Relyea & Werner 1999; Van

Buskirk 2001; Hossie *et al.* 2017) and parasitism (Han *et al.* 2011; Preston *et al.* 2014;

Gallagher *et al.* 2019) is well-documented for several species. Questing dragonfly

larvae cause tadpoles to reduce activity levels in order to avoid contact, a trait

response that can reduce tadpole feeding rates. Similarly, parasitic trematodes can

affect tadpole activity, but tadpole responses seem more variable and depend on the

parasite state; *questing* trematode cercariae either elicit no change in activity (Preston

et al. 2014) or increase activity (Rohr *et al.* 2009; Raffel *et al.* 2010), whereas *attacking*

cercariae cause strong activity spikes in tadpoles that increase attack failure rates

(Sears *et al.* 2013). However, *consuming* trematode states (larval metacercarial stages

in tissues) cause tadpoles to reduce activity (Preston *et al.* 2014).

The rich literature using this exemplar system offers guidelines for how studies

could be designed to quantify non-lethal effects of predators and parasites in other

systems. In general, simultaneous interactions with both predators and parasites elicit

reductions in activity similar to encounters with predators alone (Fig. 1a). Although

this finding suggests that tadpoles prioritize predator avoidance over parasite

avoidance, trait responses elicited during these multi-trophic interactions may be

more complex. For example, because trematode-induced changes in tadpole activity

vary by parasite state, trait responses and their non-lethal effects may vary depending

on the order of encounters, otherwise known as priority effects. Being exposed to

attacking trematode cercariae during or after encounters with predatory insects poses

a clear trade-off between increasing activity to counter cercariae attack and

decreasing activity to avoid contact with the predator. Under the hypothesis that

severe fitness consequences of consumption will favour strong trait responses

(Hypothesis 1 in main text), the tadpoles should sustain reduced activity levels, giving

rise to non-lethal effects in the form of increased infection rates and reduced feeding

rates (Fig B3a). By contrast, being exposed to attacking cercariae before encounters

with predatory insects should first elicit activity increases, followed by reduced

activity in individuals that become infected. If a questing predator is then encountered,

infections should facilitate contact avoidance of the predator (Fig. B3b). Nevertheless,

the possibility exists that encounters with questing predators while being eaten by

parasites elicits additive reductions in tadpole activity that give rise to strong non-

lethal effects from reduced feeding. These studies underscore how different consumer

states can have different non-lethal effects, and highlight how non-lethal effects of

predation and parasitism can interact. Studies that adopt similar experimental designs

in other systems would greatly help broaden out our understanding of resource

responses to consumers more generally.