1
2
3 **CaMKII oxidation is a performance/disease trade-off in vertebrate evolution**
4

5 Qinchuan Wang[1], Erick O. Hernández-Ochoa[2], Meera C. Viswanathan[1], Ian D. Blum[3], Jonathan
6 M. Granger[1], Kevin R. Murphy[1], An-Chi Wei[4], Susan Aja[5,6], Naili Liu[7], Corina M. Antonescu[8],
7 Liliana D. Florea[8], C. Conover Talbot Jr.[9], David Mohr[10], Kathryn R. Wagner[3,7], Sergi Regot[11],
8 Richard M. Lovering[12], Mark N. Wu[3], Anthony Cammarato[1], Martin F. Schneider[2], Gabriel S.
9 Bever[1,13], Mark E. Anderson*[1]
10

1 Department of Medicine, Johns Hopkins School of Medicine, Baltimore, MD, USA
2 Department of Biochemistry and Molecular Biology, University of Maryland School of
Medicine, Baltimore, MD, USA
3 Department of Neurology, Johns Hopkins School of Medicine, Baltimore, MD, USA
4 Department of Electrical Engineering, Graduate Institute of Biomedical Electronics and
Bioinformatics, National Taiwan University, Taipei, Taiwan
5 Department of Neuroscience, Johns Hopkins School of Medicine, Baltimore, MD, USA
6 Center for Metabolism and Obesity Research, Johns Hopkins School of Medicine,
Baltimore, MD, USA
7 Center for Genetic Muscle Disorders, Kennedy Krieger Institute, Baltimore, MD, USA
8 Johns Hopkins Computational Biology Consulting Core, Baltimore, MD, USA
9 Institute for Basic Biomedical Sciences, Johns Hopkins School of Medicine, Baltimore,
MD, USA
10 Johns Hopkins School of Medicine Genetic Resources Core Facility
11 Department of Molecular Biology & Genetics, Johns Hopkins School of Medicine,
Baltimore, MD, USA
12 Department of Orthopaedics, University of Maryland School of Medicine, Baltimore,
MD, USA
13 Center for Functional Anatomy & Evolution, Johns Hopkins School of Medicine,
Baltimore, MD, USA
31
32 * Correspondence: Mark E. Anderson, MD, PhD, mark.anderson@jhmi.edu

33
34

Reactive oxygen species (ROS) contribute to health and disease. CaMKII is a widely
expressed enzyme whose activation by oxidation of regulatory domain methionines (ox-
CaMKII) contributes to cardiovascular disease, asthma, and cancer. Here we integrate
comparative genomic and experimental data to show that CaMKII activation by ROS
arose more than half-a-billion years ago on the vertebrate stem lineage where it constituted
a bridge between ROS and increased intracellular Ca^{2+} release, exercise responsive gene
transcription, and improved performance in skeletal muscle. These enhancements to fight-
or-flight physiology were likely key in facilitating a well-evidenced shift in the behavioural
ecology of our immediate chordate ancestors, and, in turn, the evolutionary success of
vertebrates. Still, the ox-CaMKII innovation for augmenting performance must be
considered a critical evolutionary trade-off, as it rendered us more susceptible to common
and often fatal diseases linked to excessive ROS.

The Ca^{2+}-and calmodulin-dependent protein kinase II (CaMKII) is a multifunctional enzyme that
augments intracellular Ca^{2+} flux, and regulates gene transcription[1]. CaMKII is initially activated
by binding calmodulin, but post-translational modifications[2-5] of conserved regulatory domain
residues convert CaMKII into a Ca^{2+}- and calmodulin-independent conformation by preventing
the regulatory domain from occluding the active site (Fig. 1a). These post-translational
modifications include autophosphorylation at threonine 287 (T287, numbered in accordance with
CaMKIIγ isoform)[4,5], O-GlcNAclyation at serine 280 (S280)[3], and oxidation at cysteine
281/methionine 282 (CM; CaMKIIα) or methionine 281/282 (MM; CaMKIIγ, δ and β)[2].
CaMKII activity using the oxidative pathway (ox-CaMKII) is elevated in tissues of patients with
cardiovascular diseases[6,7], human cancer cell lines[8], and asthmatic human pulmonary
epithelium[9], suggesting ox-CaMKII contributes to common human diseases. Mutant knock-in
mice where the MM module in CaMKIIδ was replaced by ROS-resistant VV (valine 281/282)
mutations are protected against a range of diseases that involve elevated oxidative stress,
including stroke, cardiac arrhythmia, ischemia-reperfusion injury, sudden death, and asthma[6,7,10-
13]. Given its clinical relevance, we sought to better understand the biological role of ox-CaMKII
by exploring its evolutionary origins and functional implications.
64
MM residues confer ROS sensitivity to CaMKII in vertebrates and invertebrates
The phylogenetic distribution of ox-CaMKII supports the inference that an oxidation-sensitive
amino acid pair in the regulatory domain is a derived feature of vertebrates, one that emerged on
the vertebrate stem lineage sometime between approximately 570 and 480 million years ago (Fig
1b; Extended Data Fig. 1). This origin contrasts markedly with that of the T287 and S280
regulatory pathways. These appear to have evolved concurrent with CaMKII itself and are
features ancestral to all metazoans, antedating the CM module by an additional ~500 million
years (Fig 1b). The consistent expression among crown vertebrates of an oxidation-sensitive pair
at regulatory loci 281/282 indicates this vertebrate oxidative pathway has never been lost. It also
indicates that ox-CaMKII is likely integrated within functionally and evolutionarily beneficial
cascades, but at the cost of enhanced susceptibility to common, ROS-mediated diseases.
76
To test whether the MM module was sufficient to convert an invertebrate CaMKII into a ROS
sensor, we mutated the ROS-resistant VV residues in *Drosophila melanogaster* into MM
(Extended Data Fig. 2). *Drosophila melanogaster* is advantageous as a model, in part, because it
has only one *CaMKII* gene. We found that the *CaMKII^{MM}/CaMKII^{MM}* flies (referred to as MM
flies hereafter) showed a significantly higher mortality when fed sucrose solutions dosed with
paraquat, a ROS-inducing toxin[14] (Fig. 1c). These results establish a phylogenetically justified
inference that the evolutionary appearance of these residues did confer ROS sensitivity through
CaMKII activity in our stem-vertebrate ancestors.
85
The vertebrate stem lineage was witness to a major shift in behavioural ecology that set the stage
for the modern (crown) vertebrate radiation and eventually our own evolutionary origin[15]. Out
was the sessile, filter-feeding existence that served our deuterostome ancestors well and that still
characterizes the adults of our closest living chordate relatives — the lancelets and tunicates. In
was the metabolically costly strategy of being an active marine predator. A large series of
structural innovations supported this dramatic transition by fundamentally altering the way in
which these stem vertebrates received, processed, and acted on environmental stimuli. A few

examples include: a prechordal head with complex organs of special sensation and a fully
differentiated forebrain, a cartilaginous internal skeleton, muscular pharyngeal arches supporting
advanced respiration, a more powerful heart pumping blood containing haemoglobin-rich red-
blood cells, and a sympathetic nervous system and elaborate endocrine glands supporting a fight-
or-flight physiological response. Many of these novel functional complexes were developmental
products of a newly evolved population of highly migratory and multipotent neural crest cells,
and, ultimately, the entire suite of derived vertebrate features may well owe its existence to a
greatly expanded genetic tool kit made available by one, and possibly, two full rounds of genome
duplication[16]. At some level, all of these innovations promote an increasing level of activity that
must be enacted through the skeletal (striated) musculature; so, it was with the physiology of
skeletal muscle that we tested for potential beneficial effects of ox-CaMKII.
104
CaMKII activity contributes to skeletal muscle function[17,18], so we hypothesized that gaining the
MM motif allowed ROS to enhance skeletal muscle performance through ox-CaMKII. We set
out to test whether MM residues could dynamically respond to ROS in muscle fibres to increase
CaMKII activity. To measure the dynamic change of CaMKII activity in muscle fibres, we
developed a fluorescent reporter that translocates from the nucleus to the cytosol in response to
increased activity of CaMKII (kinase translocation reporter, or KTR, Extended Data Fig. 3a and
b and Methods)[19]. We validated the reporter in RPE-1 cells by showing that it translocated into
the cytosol when the cells were treated by histamine (Extended Data Fig. 3c and d), which
transiently increased cytosolic Ca^{2+} concentration[20] (data not shown). This translocation was
enhanced by co-expressed exogenous CaMKII, and blunted by co-expressed kinase-dead
CaMKII mutant[21] ($CaMKII^{K43M}$) and by the CaMKII-specific inhibitory protein CaMKIIN[22],
supporting that the KTR translocation was driven by CaMKII activity. The results show that the
cytosol/nuclear distribution of the KTR is a sensitive measurement of cellular CaMKII activity
(Extended Data Fig. 3e). To determine the role of MM residues for ROS-induced CaMKII
activity in muscle fibres, we developed a knock-in mouse where the MM residues of CaMKIIγ
were replaced with VV (Extended Data Fig. 4a and 4b, homozygous *CaMKIIγ^{VV}/CaMKIIγ^{VV}*
mice are referred to as VV mice hereafter). We selected *Camk2g* as the background because we
found that CaMKIIγ is the most abundant isoform in mouse skeletal muscle (Extended Data Fig.
5a). The knock-in mutation did not change the mRNA expression from the *Camk2g* gene
(Extended Data Fig. 5b). We introduced the CaMKII-KTR into the flexor digitorum brevis
(FDB) skeletal muscles of MM (referring to wild type or WT mice) and VV mice by
electroporating plasmids[23] encoding the reporter and isolated the FDB muscle fibres after KTR
was expressed (see Methods). We found that the increase of CaMKII activity in response to
electrical stimulation required the MM motif (Fig. 1d and e). Addition of the antioxidant N-
acetylcysteine (NAC) eliminated the rise in CaMKII activity in both VV and MM (WT) FDB
fibres in response to stimulation (Fig. 1d and e). The results showed that ROS contribute to the
activation of myofibre CaMKII, a process dependent on the MM module of CaMKII.
132
Ox-CaMKII promotes exercise performance
Based on our hypothesis that ox-CaMKII arose in vertebrates to enhance skeletal muscle
performance, we next asked whether loss of the MM residues affected exercise by testing the
mice with maximal coerced treadmill exercise (Fig. 2a). MM (WT) mice ran farther (Fig. 2b),
and attained higher maximal speeds (Fig. 2c) compared to VV littermates. Although oxidative
stress and the MM motif were necessary for normal CaMKII activity in stimulated muscle fibres

(Fig. 1d and e), the reduced exercise performance in VV mice could be due to muscle extrinsic
factors, such as motivation or metabolism. Blood lactate accumulation correlates with perceived
effort during progressively intensifying exercise[24]. However, we found no difference in blood
lactate concentration between MM (WT) and VV mice either before or after running (Extended
Data Fig. 6a), suggesting that perceived effort was similar. Furthermore, the VV and MM (WT)
mice engaged equally in voluntary wheel running (Extended Data Fig. 6b), suggesting that the
difference in forced running performance in VV mice was unlikely a consequence of reduced
motivation for running. Exercise demands uninterrupted energy supply, and the depletion of
blood glucose can be a limiting factor for endurance running in mice[25]. Furthermore, CaMKIIγ
promotes hepatic gluconeogenesis[26], an important source of blood glucose during exercise[27].
However, we found no difference in blood glucose between MM (WT) and VV mice before or
immediately after running (Extended Data Fig. 6c), suggesting that the MM module does not
contribute to blood glucose maintenance under these conditions. In addition, we found no
significant difference in the body weight, lean mass, and fat mass between MM (WT) and VV
mice; nor did we find a significant difference in their activity inside cages, food intake, oxygen
consumption rate (VO_2), CO_2 production rate (VCO_2), respiratory control ratio (RER), or energy
expenditure (data not shown) when the mice were monitored individually in the Comprehensive
Lab Animal Monitoring System (CLAMS), suggesting that the energy metabolism of MM (WT)
and VV mice are similar. Based on these negative findings, we next determined whether the
diminished exercise performance observed in VV mice might result from reduced muscle
function *in vivo*. We determined the reduction in quadriceps contractility of repeated maximal
isometric contractions elicited by direct repetitive electrical stimulation of femoral nerves, in
anesthetized mice, as a measure of muscle fatigue (Fig. 2d). This *in vivo* approach allows for the
direct examination of muscle function at body temperature, with intact blood flow and
neuromuscular communication, while minimizing potential confounding factors derived from
circulatory and nervous system feedback[28]. The VV mice exhibited earlier and enhanced fatigue
compared to MM (WT) littermate mice (Fig. 2e and 2f). The reduced performance of the VV
mice was unlikely due to developmental defects or gross pathological remodelling, as we found
that the VV and MM (WT) mice have similar muscle weight to body weight ratios and grip
strength (Extended Data Fig. 7a-g). In addition, the contents of mitochondrial complexes in
muscles (Extended Data Fig. 8a), and oxidative phosphorylation and glycolysis capacities of
isolated FDB muscle fibres were all similar between MM (WT) and VV mice (Extended Data
Fig. 8b and c). Furthermore, we found no significant change in the fatigue-resistant type I fibres,
and noted significant but subtle switches among type II fibres in VV quadriceps muscles, which
were unlikely to explain the reduced endurance capacity (Extended Data Fig. 9a and b). Taken
together, these data support a view that ox-CaMKII enhances dynamic responses to exercise and
skeletal muscle performance.
176
Intracellular Ca^{2+} grades myofilament interactions, thereby serving as an essential signal for
muscle performance, and CaMKII promotes intracellular Ca^{2+} release in excitable tissues,
including skeletal muscle[17,18]. We used a validated *in vitro* model of skeletal muscle fibre
fatigue[29], under conditions where we monitored the intracellular Ca^{2+} transients (see Methods,
Fig. 2g). The VV fibres had reduced Ca^{2+} transients under basal conditions compared to MM
(WT) counterparts (Fig. 2h). Fatigue is marked by reduced intracellular Ca^{2+} transients[30], and
VV fibres showed significantly greater reductions in these Ca^{2+} transients compared to MM
(WT) (Fig. 2i). In order to test whether the exaggerated fatigue Ca^{2+} phenotype in VV muscle

fibres was a consequence of ROS-signalling, we treated MM (WT) fibres with NAC. The MM
fibres exposed to NAC phenocopied the Ca^{2+} release profiles measured in VV fibres (Fig. 2h-j).
We next used a genetic approach to reduce ROS by isolating muscle fibres from $Ncf1^{-/-}$ mice that
lack p47[31], an essential protein cofactor for NADPH oxidases that are an important source of
ROS in skeletal muscles[32-34]. Similar to NAC treated MM (WT) fibres, the $Ncf1^{-/-}$ fibres shared a
phenotype of diminished Ca^{2+} transients resembling VV muscle fibres (Fig 2. h-j). Taken
together, we interpret these data as supporting a model where ox-CaMKII contributes to
enhanced skeletal muscle performance, at least in part, by connecting ROS to mobilization of
intracellular Ca^{2+}.
194
Ox-CaMKII regulates acute transcriptional responses to exercise
Exercise imposes metabolic, mechanical and redox stresses on skeletal muscle leading to
transcriptional adaptation that is partly orchestrated by CaMKII[35,36]. We next measured
transcriptional responses to submaximal exercise in skeletal muscles, comparing poly(A)$^+$
transcriptomes by RNA sequencing from MM (WT) and VV littermate mice under identical
conditions of speed, time, distance and feeding conditions (Fig. 3a, see Methods). Principal
components analysis showed that sedentary MM (WT) and VV muscles had very similar
transcriptional profiles (Fig. 3b). In contrast, transcriptional responses to exercise by VV
muscles were present, but diminished compared to MM (WT) (Fig. 3b). We found that 582
genes were significantly up or down regulated (multiple-test false discovery rate-adjusted q-
value < 0.05) in the MM (WT) samples, whereas only 216 genes reached the same threshold of q
< 0.05 in the VV muscles (Fig. 3c and Supplementary table 1). Among the significantly changed
genes in VV muscles, most (180, or 83%) were recapitulated by the MM (WT) muscles. To
further compare the transcriptional responses of MM (WT) and VV muscles at the level of
individual genes, we ranked the exercise-responsive genes identified in the MM (WT) muscles
based on their $\log_2$ (fold change) values, as diagrammed (Fig. 3d, left panel), and plotted these
genes in heat map palettes (Fig. 3d, middle panel). The changes of the same set of genes in the
VV muscles were shown in separate palettes (Fig. 3d, right panel), but in the same order as that
of the MM palettes. It is clear that the MM module had heterogeneous effects on the response of
individual genes to exercise: most exercise responsive genes preserved their qualitative
responses in VV muscles, while the up- or down-regulations of a small number of genes were
completely blunted or even reversed. The results suggest that ox-CaMKII plays important, but
specified, roles in the acute transcriptional response of skeletal muscles to exercise.
218
We next used QIAGEN Ingenuity Pathway Analysis[37] to extract biological pathway information
from genes that showed a large shift (more than ±2σ) in expression in response to exercise. We
found that in the MM (WT) muscles, eight out of the ten most significantly enriched biological
function terms were related to inflammation (Fig. 3e). Strikingly, exercise induced lesser
changes of the genes involved in the inflammatory response in the VV muscles (Fig. 3e). In
contrast, the top 10 most enriched biological functions in VV muscles included terms such as
"differentiation of muscle cells" and "growth of muscle tissue", which were expected for the
adaptive response to exercise[36]. Importantly, the MM (WT) muscles shared a similar pathway
enrichment for these biological functions (Fig. 3f). We then directly compared the
transcriptomes of exercised MM (WT) and VV muscles to identify a list of genes that showed
the most prominent differences (more than ±2σ) between genotypes under this post-exercise
condition. When these differentially regulated genes were analysed by Ingenuity Pathway

Analysis, the results (Fig. 3g) further supported the prominent difference in inflammatory
responses between exercised MM (WT) and VV muscles: exercised MM (WT) muscles showed
significant enrichments ($p < 0.05$) and activation (z score ≥ 2.0) of multiple biological functions
related to inflammation (Fig. 3g). Our results suggest that ox-CaMKII plays an important role in
coupling ROS to the activation of physiological inflammatory response pathways, a well-
established adaptive response to a single bout of unaccustomed exercise[38]. Under disease
conditions CaMKII has been shown to promote inflammation in the heart[39-41] and airway[11], and
to function in mast cells[11], macrophages[42-45] and T cells[46]. Our data unambiguously established
ox-CaMKII as a molecular connection between inflammatory responses and physiological ROS
signalling.
241
MM enacted the performance/disease trade-off in flies
For it to be fixed by natural selection, the MM module likely provided fitness benefits to the
ancestral vertebrates. Using *Drosophila melanogaster* as a model, we next tested whether the
MM module could exert beneficial effects on physiological performance in an invertebrate. We
reasoned that if a performance benefit is conferred by the MM module in flies it would suggest
that the cellular context of invertebrates was permissive for the physiological benefits of ox-
CaMKII. Since climbing involves insect leg muscles that are functionally and physiologically
analogous to skeletal muscles of vertebrates[47], we tested the climbing ability MM and VV (WT)
flies. We placed the flies into vertical race-tracks too narrow for flying, but wide enough for
climbing. When the flies were dislodged to the bottom of the race-tracks by vertical orientation
of the apparatus, they climbed upwards as an innate escape response (Fig. 4a). Strikingly, the
MM flies climbed at a significantly higher velocity than VV (WT) flies (Fig. 4b, control
condition). The superior climbing performance conferred by the MM module was dependent on
the physiological redox state, because ingesting food supplemented with the antioxidant NAC for
24 hours dose-dependently reduced the performance of MM but not VV (WT) flies (Fig. 4b,
NAC-treated conditions). The MM flies climbed at similar velocities to VV (WT) flies after
treatment by NAC (Fig. 4b). These results suggested that the MM module was capable of
enhancing physiological responses to ROS in an invertebrate and likely also in ancestral
vertebrates. To determine whether the MM module plays a direct role in *Drosophila*
melanogaster striated muscles, we evaluated the performance of denervated hearts in MM and
VV (WT) flies. In *Drosophila* the heart is a muscular tube that is experimentally accessible (Fig
4c). The hearts of MM flies had significantly better basal performance evidenced by significantly
higher shortening velocity and relaxation rate (Fig. 4d and e). Similar to the vertical climbing
assay, the performance benefits of the MM module in heart tubes were lost after exposure to
NAC (Fig 4d and e). We interpreted these studies to show that ox-CaMKII was capable of
producing ROS-driven physiological enhancements across evolutionarily distant species.
268
The striking benefits of MM modules on motor function and cardiac performance in flies (Fig. 4)
contrasted with the fact that MM module promoted death when the flies were exposed to lethal
doses of paraquat (Fig. 1c). Conceivably, when ROS increase above an optimal level, the MM
module transduces the toxic ROS signal into excessive CaMKII activity that is detrimental to the
flies. To test this possibility, we examined the effects of a sublethal dose of paraquat (4 mM for
24 hours) on climbing (Fig. 5a). After paraquat treatment, the MM flies exhibited significantly
reduced climbing velocity, whereas VV (WT) flies were unaffected (Fig. 5a). The notion that
excessive CaMKII activity is detrimental to motor function in invertebrates is supported by the

observation that a hyperactive mutation of CaMKII impaired motor function of *Caenorhabditis*
elegans[48]. We further examined the effects of very low dose paraquat feeding (1 mM for 3 to 6
days) on spontaneous ambulatory activity. We found that there was no difference in spontaneous
ambulation between MM and VV (WT) flies at baseline, whereas exposure to food containing
paraquat significantly reduced daily ambulatory activity counts only in MM flies (Fig 5b).
Similarly, the benefits of the MM module on performance of denervated hearts were completely
abrogated when the fly hearts were exposed to 10 mM paraquat for 90 minutes (Fig 5c and d);
and longer exposure (150 minutes) to paraquat disrupted the contraction of significantly higher
portions of MM than VV hearts (Fig 5e, and supplementary video 1). Taken together, effects of
the MM module in fly CaMKII are strikingly similar to those in mice, strongly supporting the
case for a performance/disease trade-off. The MM module promotes motor function in both
species in response to physiological ROS, but switches from an asset to a liability when the
organisms are challenged by pathological oxidative stress.
290
Finally, we tested whether insertion of the MM module in CaMKII could establish unique
connections between ROS and gene expression in flies, potentially mirroring the situation in
mice (Fig. 3). Many mammalian transcription regulators targeted by CaMKII have orthologous
counterparts in *Drosophila melanogaster* (http://flybase.org/). We fed our MM and VV (WT)
flies with 10 mM paraquat for 24 hours at 25 °C, a regimen that induced elevated mortality in
MM flies (Fig. 1c), and is known to induce a stereotyped transcriptional response in *Drosophila*
melanogaster[49]. We focused on a subset of the paraquat-induced fly genes[49] whose paralogues
in mice were altered by exercise. We randomly selected some of these genes and confirmed by
RT-qPCR that all were significantly regulated by paraquat in MM and VV flies (Fig. 5f).
Strikingly, none of these genes showed a difference in expression between MM and VV flies
consuming control food, while a subset exhibited significant differences between MM and VV
flies after paraquat feeding (Fig. 5f). The results suggest that introducing the MM module to fly
CaMKII bridges ROS to the expression of a specific set of genes, reminiscent of the
transcriptional effects of the MM module in exercising mouse skeletal muscles (Fig. 3).
305
Discussion
Our studies provide new *in vivo* and *in vitro* evidence that ox-CaMKII directly orchestrates
connections between ROS, intracellular Ca^{2+} and gene transcription that lead to physiological
advantages in mice, and, presumably, other vertebrates. Our pattern-based phylogenetic results
indicate this advantage evolved concurrent with the establishment of the modern vertebrate body
plan and its highly active behavioural ecology. It seems likely that ox-CaMKII was a key
innovation in facilitating the heightened physiological output required of these derived
anatomical systems and thus played a key role in the initial establishment and continued
evolutionary success of vertebrates. The conservation of the CM/MM module in all isoforms of
vertebrate CaMKII further suggests that ox-CaMKII plays diverse physiological roles, beyond
those uncovered by this study in the skeletal muscles. The formative evolutionary role of the
MM module is paired with considerable irony, given the well-recognized contributions of ox-
CaMKII to major chronic and life-threatening human diseases. The striking observation that the
MM module enacts the performance/disease trade-off in flies, an invertebrate diverged from our
common ancestors for more than 500 million years is particularly worth noting. It suggests that
the MM module is a concise but highly impactful ROS sensor, and once it was obtained by
CaMKII in the ancestral vertebrate, the MM (CM) module was sufficient to couple ROS to a

wide range of CaMKII targets important for enhanced performance, gene expression, disease and
death. The totality of this information strongly supports the conclusion that the CM/MM module
is an evolutionary trade-off in vertebrates that uses ROS to enhance physiological performance,
while simultaneously bestowing sensitivity to ROS for promoting chronic diseases, many of
which transpire in older organisms, beyond the reach of natural selection.
328

329 **Acknowledgement**
330 We thank Drs. Hal Dietz and Gregg Semenza for their insightful comments and suggestions,
331 Teresa Ruggle for assistance in graphic design, Benjamin Garlow for assistance in developing
332 KTR, Jinying Yang for managing mice, and Tran Nguyen for maintaining fly stocks.
333
334 This work was supported by the National Institutes of Health (R35-HL140034 to MEA, R37-
335 AR055099 to EOH and MFS, R01-AR059179 and R21-AR067872-01 to RML, R01-HL124091
336 to MCV and AC, R01-NS079584 to MNW) and a MOST grant (MOST-107-2636-B-002 -001 to
337 MEA).
338

Author contributions
Q.W., G.S.B. and M.E.A. contributed to the conception and design of the work; Q.W., E.O.H.,
M.C.V., I.D.B., J.M.G., K.R.M., A.W., S.A., N.L., D.M., K.R.W. and R.M.L. contributed to the
acquisition of the data; Q.W., E.O.H., M.C.V., I.D.B., K.R.M., A.W., S.A., C.M.A., L.D.F.,
C.C.T., D.M., S.R., M.N.W., A.C. and M.S.F. analysed and interpreted the data; Q.W., G.S.B.
and M.E.A drafted the manuscript; Q.W., G.S.B. and M.E.A. substantively revised the
manuscript.
346

347 **Data availability statement**
348 Raw sequencing data and the gene-expression matrix are available in the Gene Expression
349 Omnibus (GEO) under accession number GSE132520. All other data are available from the
350 corresponding author upon reasonable request.
351

352
353

Fig. 1 | The MM motif in CaMKII arose in vertebrates, allowing CaMKII activation by ROS. **a**, CaMKII activation is initiated by binding Ca^{2+}/CaM, but Ca^{2+}/CaM independent CaMKII activity is sustained by post-translational (PTM) modifications of the regulatory

domain. The CaMKII regulatory domain sequences of *Drosophila melanogaster* CaMKII and
Mus musculus CaMKIIγ are shown. **b**, Phylogenetic origin and conservation of key residues for
the activating PTMs of CaMKII (divergence time estimates from Kumar et al[50]). mya: million
years ago; bya: billion years ago. **c**, Replacing the VV residues of *Drosophila melanogaster*
CaMKII with MM increased mortality caused by exposure to 25 mM and 50 mM paraquat (PQ)
incorporated in 5% sucrose solution at room temperature (21 °C, left panel) after 24 hours (24-h)
and 48 hours (48-h), and by exposure to 10 mM paraquat at 25 °C after 24 hours (right panel).
Only one out of 1020 flies fed control food (5% sucrose solution) died (not shown). *P*-values
from Fisher's exact test are shown above the square brackets. N = 216 and 217 respectively for
MM and VV flies treated at the room temperature; n = 510 for both genotypes of flies treated by
either control or paraquat solutions under the 25 °C test condition. **d**, Representative confocal
micrographs of MM and VV mouse FDB muscle fibres expressing CaMKII-KTR before and
after field electrical stimulation in the absence or presence of 2 mM N-acetylcysteine (NAC).
Arrows indicate nuclei. **e**, Quantification of CaMKII activity (cytosolic to nuclear CaMKII-KTR
signal ratio) in MM and VV fibres before and after field stimulation in the absence or presence
of 2 mM NAC (Two-way ANOVA followed by Sidak's multiple comparisons test. n = 14 nuclei
from 6 fibres (MM) and n = 21 nuclei from 7 fibres (VV) in the left panel, and n = 11 nuclei
from 5 fibres (MM) and n = 16 nuclei from 6 fibres (VV) in the right panel. *P*-values are shown
above the brackets).
378

379
380
Fig. 2 | Ox-CaMKII supports exercise performance and enhances Ca²⁺ transients in mouse
skeletal muscle fibres. **a**, Protocol for treadmill exercise. **b**, Running distance, and **c**, maximal
speed attained prior to exhaustion. Unpaired Student's *t* test, two-tailed, *P*-value shown above
the square brackets, n = 20 MM (WT) mice and n = 27 VV mice. **d**, Experimental apparatus and
electrical stimulation protocol for assessing quadriceps muscle performance *in vivo*. **e**, Averaged

traces of quadriceps torque (normalized against maximum) from MM and VV mice during
optimized nerve stimulation, n = 10 VV mice, and n = 11 MM mice. **f**, Quantification of fatigue
defined by torque reduction divided by initial torque of each individual mouse, as shown in e.
Unpaired Student's *t* test, two-tailed, *P*-value shown above the square bracket. **g**, Protocol for
field electrical stimulation of isolated FDB muscle fibres loaded with the Ca^{2+} sensitive
fluorescent dye Rhod2. Rhod2 fluorescence during one cycle of electrical stimulation in MM and
VV fibres, an MM fibre treated by NAC, and a *Ncf1$^{-/-}$* fibre, before (h) and after (i) fatigue. **j**,
Quantification of peak Ca^{2+} transients as measured in h and i. Before fatigue: n = 42 MM, n = 27
VV, n = 8 MM + NAC, and n = 32 *Ncf1-/-* fibres; after fatigue: n = 40 MM, n = 27 VV, n = 8
MM + NAC, and n = 32 *Ncf1-/-* fibres; *P*-values are shown above the brackets; Dunnett's
multiple comparisons tests comparing all other groups to VV fibres. Horizontal lines in b, c, f,
and j indicate medians.
398

Figure 3

Fig. 3 | Ox-CaMKII is important for acute transcriptional responses to exercise. a, Protocol
for submaximal exercise and muscle collection. **b,** Principal components analysis (PCA) of
RNA sequencing results of sedentary (sed) and exercised (ex) muscle samples. The distance
between samples in PCA corresponds to similarity (near) or difference (far) in their
transcriptional profiles (n = 4 mice for each group). **c,** Numbers and overlap of significantly

changed (false discovery rate-adjusted q-value < 0.05) genes in response to exercise in MM and
VV muscles. Arrows indicates up- (↑) or down-regulation (↓) when comparing exercised
muscles to sedentary muscles. Left panel of **d**, diagram of the layouts for arranging genes in the
middle and right panels; middle panel of **d**, genes whose expression was significantly (q <0.05)
changed in response to exercise in the MM (WT) muscles are ordered according to the diagram
in the left panel, and their $\log_2$ (fold changes) are represented by colour; right panel, the $\log_2$
(fold change) of the same genes in response to exercise in the VV muscles are shown. **e**, Top-10
most significantly (smallest P-values) changed functions identified by Ingenuity Pathway
Analysis comparing transcriptomes of exercised MM muscles to their sedentary counterparts.
Corresponding enrichment P-values of the same functions in the exercised VV muscles are
plotted for comparison. **f**, Top-10 most significantly (smallest P-values) changed functions
identified by Ingenuity Pathway Analysis comparing transcriptomes of exercised VV muscles to
their sedentary counterparts. Corresponding enrichment P-values of the same functions in the
exercised MM muscles are plotted for comparison. **g**, Ingenuity pathway analysis directly
comparing transcriptomes of exercised MM and VV muscles. In (e-g), activation, depression or
bidirectional changes of the biological functions are determined by the z-score of Ingenuity
Pathway Analysis for each pathway ($z \geq 2.0$ for activation, $z \leq -2.0$ for depression, otherwise for
bidirectional changes).
424

426
427
Fig. 4 | MM module couples ROS to improved performance in *Drosophila melanogaster*. a,
Diagram for climbing test. **b**, Vertical climbing velocity of flies treated by control food (5%
sucrose) or food containing 10 mM or 50 mM NAC for 24 hours; n = 93 control-treated MM, n =
90 10 mM NAC-treated MM, n = 86 50 mM NAC-treated MM, n = 94 control-treated VV, n =
91 10 mM NAC-treated VV, n = 94 50 mM NAC-treated VV. Horizontal lines indicate
medians. *P*-values shown above the brackets, Tukey's multiple comparisons test. **c**, diagram of a
fly heart (in blue color). **d** and **e**, Cardiac performance indices of MM and VV hearts before and
after 60-minute treatment by 10 mM NAC. The shortening velocity (d) and relaxation rate were
assessed. n = 27 MM hearts per condition and n = 29 VV hearts per condition. Horizontal lines
indicate medians. *P*-values shown about the brackets in (d) and (e), Tukey's multiple
comparisons test.
439

Figure 5

Fig. 5 | MM module sensitizes flies to pathological oxidative stress. a, Vertical climbing velocity of flies treated by control food or food containing 4 mM paraquat for 24 hours at 25 °C; n = 68 control-treated MM, n = 74 paraquat-treated MM, n = 57 control-treated VV, n = 75 paraquat-treated VV flies. Horizontal lines indicate medians. *P*-value from Sidak's multiple comparisons test shown in the graph. **b**, Daily activity counts of MM and VV flies consuming a

control or paraquat (1mM) diet. Diets started at day 1 and behaviour monitoring occurred
between days 3 and 6 (Points and error bars are mean ± SEM. P-values calculated using Tukey's
multiple comparisons test for effect of paraquat, n = 29 control MM, n = 21 paraquat-treated
MM, n = 31 control VV, and n = 21 paraquat-treated VV flies). **c and d,** Cardiac performance of
hearts bathed first in control artificial haemolymph and then in haemolymph containing 10 mM
of paraquat for 90 minutes. n = 26 hearts per genotype and the P-values from Tukey's multiple
comparisons test are shown. Horizontal lines indicate medians. **e,** All MM and VV hearts
showed normal contraction before paraquat treatment, however, after exposure to 10 mM
paraquat for 150 minutes, significantly more MM hearts became hypokinetic, severely
hypokinetic or akinetic (examples of categorical cardiac performance are in the supplementary
video 1). n = 26 hearts per genotype, P-value from Chi-square test. **f,** expression heat map of a
subset of paraquat responsive genes that was quantified by RT-qPCR after RNA was extracted
from flies ingesting control food (Control, 5% sucrose solution) or paraquat (10 mM of paraquat
in 5% sucrose solution) for 24 hours at 25 °C (n = 8 biological replicates per group, each
containing 15 males and 15 females). All of these genes were significantly upregulated by
paraquat ($P < 0.05$, not shown, two-Way ANOVA). No genes showed significant differences
between MM and VV flies fed control food, whereas a subset of genes had significantly higher
expression in MM than in VV flies after exposure to paraquat (P-values shown in the table.
Sidak's multiple comparisons test).
468

Extended Data Fig. 1 | Sequence alignment of CaMKII regulatory domains reveals the emergence and conservation of ox-CaMKII in vertebrates. The paired oxidation sensitive amino acid residues (CM and MM at positions corresponding to residues 281 and 282 of mouse CaMKIIγ, highlighted yellow) are required for ox-CaMKII and are present only in the vertebrates. In contrast, none of the sampled invertebrate CaMKII possess the CM/MM pair (highlighted in light blue).

a

b

VV/VV

MM/MM

c

478
479
Extended Data Fig. 2 | Generation of *CaMKII^MM*/*CaMKII^MM* flies using CRISPR. a,
Schematics of the CRISPR guide designs and the single strand template (ssODN-1R) that
mediated the homology directed recombination, resulting in mutations of codons from encoding
VV to encoding MM, and introduction of the silent mutations for NsiI/NsiI-HF restriction site.
Primers-F1/R2 anneal outside of the range homologous to the ssODN-1R to amplify the genomic
region for genotyping. **b**, Chromatograms of sequencing results from the PCR products amplified
by primers F1 and R2 from homozygous VV/VV (top) and MM/MM (bottom) flies. The VV/VV
(wild type) and MM/MM flies are referred to as VV and MM flies in the text for brevity. **c**,
Agarose gel electrophoresis of PCR products from flies backcrossed into the w1118 or iso31
genetic background for 5 generations. The genotypes of flies were determined by digesting the
493 bp PCR products with the restriction enzyme NsiI. PCR products amplified from the VV
allele were resistant to NsiI, while those from the MM allele were cut into 345 bp and 148 bp
fragments.
493

495
496
Extended Data Fig. 3 | Design and validation of the CaMKII activity reporter, CaMKII-
KTR. a, Schematic of the CaMKII kinase activity translocation reporter CaMKII-KTR
(abbreviated as KTR). The N-terminus of the KTR is a well-characterized CaMKII-interacting
domain from HDAC4 (AA582-624[51]), followed by a nuclear localization signal (NLS) and a
nuclear exporting signal (NES). The high affinity CaMKII substrate consensus sequence
(LXRXX**S**V) was built into both the NLS and NES (see Method). The C-terminus of the

CaMKII-KTR is an enhanced green fluorescent protein (EGFP). **b**, The KTR shuttles between
the nucleus and cytosol. Phosphorylation by CaMKII decreases the strength of the NLS while
increases the strength of the NES, resulting in a net translocation of the KTR into the cytosol.
The ratio between the cytosolic and nuclear signals of the KTR corresponds to the overall
activity of CaMKII inside the cells. **c**, Fluorescent images of KTR transfected into RPE-1 cells
alone, co-transfected with CaMKII, with kinase-dead CaMKII^{K43M}, or with a CaMKII-specific
inhibitor CaMKIIN. The arrowheads indicate nuclei. Cells were imaged at time -60 seconds (s),
0, 60 s, and 120 s, and were treated at time 0 with 50 μM of histamine. Treatment with vehicle
(medium) did not elicit a response and is not shown. **d**, Quantification of cytosolic to nuclear
KTR signal ratios in RPE-1 cells as exemplified in **c** immediately before and 60 seconds after the
histamine treatment. *P*-values are shown in the graph, Sidak's multiple comparisons test for
repeated measurement comparing before and after histamine treatment. **e**, Changes in KTR
cytosolic to nuclear signal ratios 60 s after histamine treatment compared to time 0 in cells
shown in **d**. Horizontal lines indicate the means. *P*-values are shown in the graph, Dunnett's
multiple comparisons test. In (d) and (e), n = 36 KTR transfected cells, n = 19 KTR + CaMKII
cells, n = 30 KTR + CaMKII^{K43M} cells, and n = 30 KTR + CaMKIIN cells. Data in (d) and (e)
were from at least 2 independent experiments.
520
521

a

b

522
523
Extended Data Fig. 4 | Generation of knock-in *CaMKII^VV^/CaMKII^VV^* mice. a, Schematics of
the *CaMK2g* endogenous locus (top), targeting vector (middle), and targeted locus (bottom).
The MM residues are encoded by exon 10. The gene targeting was carried out on ES cells of
C57BL6/n background. **b**, chimeric mice bearing the targeted VV allele were crossed with Cre
mice to remove the Neo-pGK cassette between the two LoxP sites. The resulting
*CaMKII^MM^/CaMKII^VV^* heterozygous mice were generated backcrossed to C57BL/6J mice for >7
generations.
531

533
534
Extended Data Fig. 5 | Expression of CaMKII isoforms in representative striated muscles
quantified by RT-qPCR. a, Expression of CaMKII isoforms in the heart, soleus, TA (tibialis
anterior), and EDL (extensor digitorum longus) muscles relative to *Gapdh*. Each tissue was
represented by three individual wild type animals. **b**, RT-qPCR assay for the expression of
Camk2g in gastrocnemius muscles from MM and VV mice; n = 5 for each genotype. In a and b,
horizontal lines indicate means.

543
544
Extended Data Fig. 6 |Blood lactate, glucose, and voluntary running. a, Lactate
concentrations measured with a drop of blood from tail tips before and immediately after
treadmill exercise, n = 14 MM and n = 20 VV mice, no statistically significant differences were
present between genotypes either before or after exercise. **b**, Counts of wheel rotations during
24 hours on the 6[th] day of running wheel access by individual MM (n = 11) and VV (n = 11)
mice, no statistically significant difference was found between genotypes. **c**, Blood glucose
concentration measured from the same mice at the same time as in (a). In a-c, horizontal lines
indicate the medians.
553

555
556 **Extended Data Fig. 7 | MM (WT) and VV skeletal muscles show no difference in weight**
557 **and strength. a-f**, Muscle to body weight ratios of representative skeletal muscles, **a**, EDL
558 (extensor digitorum longus). **b**, Soleus. **c**, TA (tibialis anterior). **d**, Gastrocnemius. **e**, Triceps. **f**,
559 Quadriceps. **g**, Grip force of front paws. Horizontal bars indicate means, n = 5 MM and n = 5
560 VV mice from (a-g).
561

562
563

564
565
Extended Data Fig. 8 | Oxidative and glycolytic capacities of MM and VV muscles are similar. **a**, Western blot of gastrocnemius muscle extracts showed no difference in the expression of representative protein subunits from mitochondrial oxidative phosphorylation complexes (n = 5 mice for each genotype). Isolated flexor digitorum brevis (FDB) fibres were analysed for (**b**) oxygen consumption (OCR) and (**c**) extracellular acidification (ECAR) rates. Points and error bars are mean ± SEM. No statistically significant (multiple t-test adjusted for false-discovery) differences were found between MM and VV fibres, n = 5 MM (WT) mice and n = 4 VV mice in b and c.

576
Extended Data Fig. 9 | Analysis of skeletal muscle fibre types in MM and VV mice. a, an
example cross sectional image of quadriceps muscle with immunostaining for myosin heavy
chain isoforms I (blue), IIA (green), IIB (red), and IID (black, unstained). Note that this image
was chosen to show the staining of all four types of myosin heavy chain isoforms, and the
proportions of isoforms in this image do not reflect the entire cross section of the muscle because
fibre types are not uniformly distributed throughout the cross-section of the muscles. **b**,
quantification of the percentages of fibre types determined by myosin heavy chain isoform
expression in MM (n = 5 mice, 1 section per mouse, 20110 fibres counted in total) and VV (n =
5 mice, 1 section per mouse, 22922 fibres counted in total) mice (Fisher's exact test comparing
the same fibre types between MM and VV muscles).
587

Methods

Animal use
All animal handling procedures were in accordance with National Institutes of Health guidelines
and were approved by the Institutional Animal Care and Use Committees of Johns Hopkins
University School of Medicine.

Generation of *CaMKII^MM* point mutation in *Drosophila melanogaster* by CRISPR mediated
gene editing
The genomic sequence of the *Drosophila melanogaster CaMKII* gene was used in the CRISPR
guide design tool (http://crispr.mit.edu/) to create the CRISPR guides. The guide #1
(GTTACAGCAACGCGAACGTG) was chosen due to its close proximity to the codons
encoding V281 and V282 and its lack of high probability off-targets. The guide #1 was ordered
as complementary oligomeric DNA (Integrated DNA Technologies) and cloned into the pU6-
BbsI-ChIRNA plasmid[46] [the pU6-BbsI-chiRNA was a gift from Melissa Harrison & Kate
O'Connor-Giles & Jill Wildonger (Addgene plasmid # 45946 ; http://n2t.net/addgene:45946 ;
RRID:Addgene_45946)]. A single strand 176nt ultramer DNA oligo (ssODN-1R) was designed
as the template for HDR-mediated point mutations and was ordered from Integrated DNA
Technologies (Extended data Fig. 2a). The sequence of ssODN-1R is
"AACATTGTCGTAAGTATGGCTCCCTTTAGCTTGCGCCGCGCATTAAATTTCTTGAGA
CAGTCTACGGTTTCTTGGCGATGCATCATGGAAGCGACTCGTTCGCGTTGCTGTAAC
AATGTTTTTTCATTATCTTTATGTAAACCTAAGAGAAAAATTAGTCTGCACTTACACA
AATC". Injection of the ssODN-1R and the guide RNA encoding plasmids into fly embryos was
carried out by Rainbow Transgenic Flies, Inc (3251 Corte Malpaso Unit 506 Camarillo CA).
Genotyping was carried out by PCR amplification from genomic DNA, extracted from the wings
of the flies, with primer-F (GTCGGTTATCCACCCTTTTG), and primer-R
(GACGCCAAGTATATTGATGTGG) followed by Sanger sequencing and NsiI/NsiI-HF
digestion. The flies with the correct *CaMKII^MM* allele were backcrossed with iso31 flies[52] for five
generations to minimize the possibility of carrying off-targets from the CRISPR-mediated gene
editing.

Phylogenetic survey of CaMKII
Most CaMKII orthologues listed in Supplementary Fig. 1 were identified in the Interpro database
(http://www.ebi.ac.uk/interpro/entry/IPR013543/taxonomy), based on the criteria that the sequences
have the conserved CaMKII association domain, and a kinase domain. Additional sequences
were uncovered by BLAST in the NCBL nucleotide database and translated into proteins
(https://blast.ncbi.nlm.nih.gov/Blast.cgi). The CaMKII sequences were aligned using the
Molecular Evolutionary Genetics Analysis software MEGA-X[53]
(https://www.megasoftware.net/). The evolution of an oxidation-sensitive amino acid pair at loci
281/282 of the CaMKII regulatory domain is an unambiguous synapomorphy of crown-clade
vertebrates among Deuterostomata. The initial identity of this pair was CM, with the 281
cysteine likely being subsequently replaced by a methionine in one of the two paralogs that
resulted from a full round of genome duplication that occurred prior to the origin of the
vertebrate crown clade. Our phylogenetic survey did recover a small number of non-
deuterostome taxa that also exhibit oxidizable residues at those same regulatory loci. The large
phylogenetic separation between these taxa, both individually and collectively, from

Deuterostomata leaves it clear that they evolved independent of the vertebrate condition; they are
also MM rather than the CM of the earliest vertebrates. As a greater taxonomic diversity of
metazoan genomes become available, a meaningful probabilistic analysis of ox-CaMKII
evolution outside of Deuterostomata will be possible. But that analysis is highly unlikely to
question the evolutionarily unique nature of vertebrate ox-CaMKII.
639
640 **Generation of *CaMKII^{VV}* knock-in mutation in mice**
641 *CaMKII^{VV}* knock-in mice were generated by GenOway (https://www.genoway.com/) with mouse
642 embryonic stem cells of the C57BL/6 background as specified in Extended Data Fig. 4. The mice
643 used in experiments had been further backcrossed to C57BL/6J mice (The Jackson Laboratory,
644 000664) and all experiments were carried out with littermates.
645
Western blotting for mitochondrial complexes
Protein extracts were prepared from frozen tissue with T-PER Tissue Protein Extraction Reagent
(Thermo Scientific, #78510) in the presence of protease (Sigma-Aldrich, P8340) and
phosphatase (Sigma-Aldrich, P0044) inhibitors. The primary antibodies were the Total OXPHOS
Rodent WB Antibody Cocktail from Abcam (ab110413), and GAPDH (D16H11) XP® Rabbit
mAb (Cell Signalling, #5174). Data were collected by LI-COR Odyssey Fc (Lincoln, Nebraska
68504 USA).
653
654 **Skeletal muscle fibre typing**
655 Skeletal muscle fibre composition was determined by immunostaining following a standard
656 method[54]. The primary antibodies BA-F8-c (myosin heavy chain, slow), SC-71-c (Myosin Heavy
657 Chain Type IIA), BF-F3-c (Myosin Heavy Chain Type IIB), 6H1-s (myosin heavy chain, fast,
658 IIX) were obtained from Developmental Studies Hybridoma Bank (University of Iowa).
659
Paraquat treatment and behaviour study of flies
To determine the effects of paraquat on mortality, newly eclosed flies were sorted under CO_2
anesthetization and placed into individual vials, and each vial received 10 males and 10 females.
When the flies reached 5 to 7 days old, they were transferred into vials containing a filter paper
pad (cut from Bio-Rad #1704085) soaked with 600 μL of 5% sucrose solution (control) or 5%
sucrose solutions containing 10 mM, 25 mM or 50 mM paraquat (Sigma-Aldrich, # 856177).
Mortality of flies was recorded at 24 and 48 hours after initiation of the treatment.
667
To test the negative geotaxis (climbing), females were collected and aged, as above, and kept in
vials in groups of 10 flies. For low dose paraquat treatment, the flies were treated by 5% sucrose
or 5% sucrose + 4 mM paraquat for 24 hours at 25 °C. They were then transferred into vertical
test tracks made from 25 mL serological pipette tubes. During the climbing test, the flies were
dislodged to the bottom of the tubes by rapidly tapping the vials on the desktop for 10 times and
climbing was video recorded for subsequent analysis. Each group of flies was tested for 10
consecutive trials at 30 seconds intervals. The vertical distances the flies climbed in 6 seconds
since the last tap (time 0) were used to calculate the vertical velocity of climbing. Flies that
initiated flight or paused during the 6-second time window were excluded from the analysis. We
found that the flies performed reproducibly from the second to tenth trials and presented data
from trial 2 in Fig. 4b and Fig. 5a.
679

680   To determine the effects of very low dose paraquat (1 mM) on daily ambulatory activity,
681   individual 1-week old female flies were anesthetized by CO_2 and loaded into tubes containing
682   control or paraquat-containing food and monitored by *Drosophila* Activity Monitoring System
683   (Trikinetics). Fly behaviour was recorded from day 3 to day 6.
684
Drosophila **cardiac physiological analysis**
Dorsal cardiac tubes of ten-day-old female *CaMKII^WT* (denoted VV (WT)) and *CaMKII^MM*
(denoted MM) flies were dissected in oxygenated artificial hemolymph[55]. Myogenic contractions
of cardiac tissue were recorded using the Hamamatsu Orca Flash 2.8 CMOS camera on a Leica
DM5000B TL microscope with a 10x immersion lens at ~120 frames per second at baseline and
after 90 and 150 minutes following the addition of 10 mM paraquat, or 60 minutes after the
addition of 10 mM NAC. Cardiac physiological indices were determined using the semi-
automated optical heartbeat analysis program[56,57]. Significant differences between genotypes
before and after paraquat treatment for 90 minutes were determined by two-tailed Mann-Whitney
tests. After 150 minutes in paraquat, many of the MM hearts no longer contracted and the cardiac
indices could not be meaningfully derived. We, therefore, categorized the contractions as normal,
hypokinetic (part of the heart contracting), severe hypokinetic (only twitching could be observed
in part of the heart), and akinetic (no movement). Representative videos for each category are
shown in supplementary video 1. The categorical data were assessed using a Chi-square test.
699
Construction and validation of a CaMKII activity sensor CaMKII-KTR
The CaMKII-KTR was constructed based on the principles previously published[19] and described
in the Extended Data Fig. 3. Specifically, the sensor consists (from N-terminus to C-terminus) of
a CaMKII-binding region derived from HDAC4, a linker, a nuclear localization signal (NLS), a
nuclear exporting signal (NES) and a fluorescent protein. Optimized CaMKII phosphorylation
sites were built into the NLS and NES while keeping the NLS and NES functional. The protein
sequence of the CaMKII-KTR, excluding the enhanced green fluorescent protein, is
EQELLFRQQALLLEQQRIHQLRNYQASMEAAGIPVSFGSHRPLKRTASVNEDEAPSKKPL
ARTASVSSRLERLTLQSS. A cDNA encoding this sequence was ordered as a codon-optimized
gene block (gBlock_HDAC4-NLS-NES) from Integrated DNA Technologies
(gccaccatgGAACAGGAACTGCTCTTCCGGCAACAGGCACTTCTGTTGGAGCAGCAACG
AATCCATCAACTTAGAAACTACCAAGCATCAATGGAAGCAGCCGGGATTCCTGTCTC
CTTCGGATCTCACAGACCTCTCAAAAGGACAGCTAGTGTAAACGAGGACGAAGCAC
CTTCAAAGAAACCCTTGGCTAGGACCGCTAGTGTCAGTAGTCGACTGGAGCGGTTGA
CACTTCAAAGTTCC). The gBlock_HDAC4-NLS-NES was cloned into Cerulean-N1 vector[58]
(Cerulean-N1 was a gift from Michael Davidson & Dave Piston, Addgene plasmid # 54742 ;
http://n2t.net/addgene:54742 ; RRID:Addgene_54742) by In-fusion cloning technology (In-
Fusion® HD Cloning Plus CE, Takara, CA). The Cerulean encoding region was then replaced by
a stretch of cDNA encoding eGFP, derived from pEGFP-C1 (Takara, CA).
To validate the response of the CaMKII-KTR to intracellular activity of CaMKII, we transfected
(FuGENE® HD Transfection Reagent, Promega, WI) the CaMKII-KTR or co-transfected it with
CaMKII, CaMKII^{K43M}, and CaMKIIN constructs into RPE-1 cells and stimulated the cells with
50 μM histamine. Before cells were imaged, we replaced the culture medium with Live Cell
Imaging Solution supplemented with 4.5 g/L glucose (ThermoFisher Scientific, A14291DJ), and
stained their nuclei with Hoechst 33342 (ThermoFisher Scientific, #62249) for 20 minutes to
facilitate identification of the nuclei. Fluorescent images were collected using an Olympus IX83

epifluorescence microscope equipped with an ORCA Flash 4.0 sCMOS camera and
UPLSAPO20X NA0.75 objective lens. Cells were maintained at 37°C in an OkoLabs stage top
incubator. Image analyses were carried out in CellProfiler[59], which identified the nuclei and five-
pixel-wide cytosolic rings surrounding the nuclei. The cytosolic to nuclear KTR signal ratios
were calculated using the median intensities measured from the nuclei and cytosolic rings of
individual cells.
732
Mouse treadmill exercise
Exercise capacity tests were carried out with the Exer 3/6 Rodent treadmill (Columbus
Instrument, Columbus, OH). Prior to exercise capacity testing, the mice (12 to 15-week-old)
were acclimated to the treadmill for three sessions on three consecutive days. The treadmill was
set to 10° inclination and the speed was set to 0, 5, and 10 m/min for the first, second and third
acclimation sessions respectively. The electric shock grid at the rear end of the treadmill was
turned on and set at stimulation intensity of 9 and frequency of 3 Hz. During exercise capacity
testing, each mouse was placed into a lane of the treadmill. The genotype of the animals was
blinded to the operator. The exercise protocol consisted of the following steps: (1) 10 m/min for
2 minutes for warm up, (2) continuous acceleration from 15 m/min at a rate of 0.6 m/min^2 until
the mouse was exhausted. Exhaustion was determined when the mouse stayed on the shock grid
continuously for 5 seconds and was determined by the same observer for all experiments.
Glucose and lactate were measured from a drop of blood from the tail tip before and immediately
after exercise, using an OneTouch Ultra 2 glucometer (Lifescan, Inc) and a Lactate Plus lactate
meter (Nova Biomedical) respectively.
748
Mouse voluntary wheel running, and accompanying metabolic data
Voluntary wheel running data were collected from mice tested for 6 days in an open-circuit
indirect calorimeter outfitted with running wheels (Comprehensive Lab Animal Monitoring
System, Columbus Instruments) at the Center for Metabolism and Obesity Research service core.
Data were collected continuously (Oxymax software, v.5.9, Columbus Instruments). Days 1-5 of
acclimation to wheel running were monitored for expected daily increases in number of wheel
rotations; the analysis of day 6 is presented. The instrument also provided data for voluntary
physical activity in the main cage as indexed by counts of infrared beam breaks, intakes of
powdered diet (2018, Envigo), as well as rates of O_2 consumption (VO_2, ml/kg/hr) and CO_2
production (VCO_2, ml/kg/hr). Oxymax software calculated the respiratory exchange ratio (RER
= VCO_2/VO_2) to assess the oxidized fuel mixture being oxidized, and the rates of energy
expenditure (EE, kcal/kg/hr; EE= $VO_2 \times [3.815 + (1.232 \times RER)]$. The standard outputs of
indirect calorimetry data as per-kg/hr were also renormalized and analysed as per-kg-lean-
mass/hr. Body composition data for lean mass and fat mass were obtained using an EchoMRI-
100 at the Johns Hopkins University Phenotyping service core. None of the measures from these
experiments showed group differences; data from voluntary wheel running are presented.
765
766 **Assessments muscle function *in vivo***
767 Grip strength measurements were carried out as described previously using a grip strength meter
768 (Columbus Instruments, Columbus, OH, USA)[60]. Each mouse performed grip strength test until
769 6 successful attempts were accumulated, and the maximal force among the six attempts was
770 taken as the grip strength.
771

In vivo quadriceps torque measurement was described previously[28]. Briefly, the mice were
anesthetized under 4% isoflurane and then maintained at 1%. Then their pelvis, torso and femur
were stabilized on the apparatus. Afterwards, the distal leg was taped to a lever arm, which was
connected with a torque cell. The femoral nerve was stimulated subcutaneously to induce
maximal quadriceps muscle contractions and the torque produced was recorded by a connected
computer for subsequent analysis. The voltage of the stimulation was optimized prior to the
studies to produce the maximal torque.
779
In cellulo study of skeletal muscle fibres
Electroporation of DNA into flexor digitorum brevis (FDB) skeletal muscles of mice, muscle
fibre culture, measurements of cytosol/nucleus distribution of CaMKII-KTR, and action
potential-induced Ca^{2+} transient imaging followed our previous reports[23,29]. For N-acety-L-
cysteine (NAC) treatment, the fibres were incubated for 20 minutes with 2 mM NAC (Sigma-
Aldrich, St. Louis, MO; catalogue # A-7250). Fibres for study were randomly chosen, and where
noted, 2-3 nuclei from the same fibre were studied.
787
For Seahorse study, FDB skeletal muscle fibers were isolated one day before the experiments
and plated to a laminin-pretreated Seahorse XF96 Cell Culture Microplates overnight. Cell
metabolism and bioenergetic analyses of muscle fibers were performed using an Agilent XF96
Extracellular Flux Analyzer. XF Cell Mitochondrial Stress Test kit and Glycolysis Stress Test kit
were used to measure mitochondria respiration capacities and cellular glycolysis capacities
following the manufacturer's protocol. In the mitochondrial stress assay, muscle fibers were
incubated in the muscle fiber assay medium (120 mM NaCl, 3.5 mM KCl, 1.3 mM $CaCl_2$, 0.4
mM KH_2PO_4, 1 mM $MgCl_2$, 5 mM HEPES, and 10 mM glucose, pH 7.4) followed by port
injections of final concentration of 1 µM oligomycin, 0.5 µM FCCP, 10 mM pyruvate and 0.5
µM antimycin A/rotenone. In the glycolysis stress assay, glucose was not included in the initial
muscle fiber assay medium, and then 2 mM glutamine, 10mM glucose, 1µM oligomycin, and
50mM 2-DG were injected sequentially. The oxygen consumption rate (OCR) and extracellular
acidification rate (ECAR) were analyzed using Seahorse Wave software. OCR and ECAR were
normalized by the number of skeletal muscle fibers per well.
802
Mouse treatment and sample collection for RNA sequencing
Male mice between 13 and 15 weeks of age were used for the RNA sequencing experiments. The
mice were first acclimated to the treadmill (Exer 3/6, Columbus Instrument, Columbus, OH) for
10 minutes on three consecutive days. The treadmill inclined at 10° and the speed was 0, 5
m·min^{-1}, and 10 m·min^{-1} for day 1, 2 and 3 respectively. To minimize the effects of training on
the skeletal muscles, the mice rested for 7 days before sample collection. On the day of sample
collection, food was withdrawn from the mice at 9:00 AM to minimize the effects of food intake
on signalling and gene transcription in the muscles. At 12:00 PM, the mice ran on a treadmill set
to 10° of inclination. The treadmill speed was set at 10 m/min for 2 minutes to allow the mice to
warm up. The speed was then increased to 15 m/min and then continuously ramped up from 15
m/min to 23 m/min at a rate of acceleration of 0.6 m·min^{-2}. When the running protocol ended,
the mice had run 350 meters, which was lower than the average running capacity of VV mice
tested by the same treadmill protocol. Any mice that did not finish the protocol were excluded
from subsequent sample collection. After exercise, the mice were allowed to rest for 3 hours with
access to water but not food. Then they were euthanized by cervical dislocation after being

anesthetized by isoflurane. The quadriceps muscles were quickly excised, frozen, and stored in
liquid nitrogen.
820
RNA extraction, quantification and quality control
To extract high quality total RNA, the quadriceps muscles were processed first in the Trizol
reagent (ThermoFisher Scientific, Catalog # 15596018) and then purified by RNeasy mini
columns (Qiagen, catalog # 74104) as follows. To avoid sampling bias, the entire quadriceps
muscles were homogenized in Trizol reagent at the weight (mg) to volume (µL) ratio of 1:15.
One mL of homogenate was processed following the manufacture's protocol until the step of
phase separation. Then, 0.5 mL of the aqueous phase was mixed with an equal volume of 70%
ethanol for subsequent RNA purification by RNeasy mini kit (Qiagen, catalog # 74104) with on-
column DNAse (RNase-Free DNase Set, Qiagen catalog # 79254) treatment. The concentration
of the RNA was determined by Qubit fluorometric quantitation (Qubit RNA BR Assay Kit,
ThermoFisher Scientific, catalog # Q10210). The integrity of the total RNA was determined by a
Fragment Analyzer (Advanced Analytical Technologies, Inc). The average RNA Quality
Number (RQN) was 9.04 ± 0.08 (Mean ± SEM, n=24).
834
RNA sequencing library preparation
1 µg of total RNA from each sample was used for RNA sequencing library preparation with the
TruSeq® Stranded mRNA Library Prep kit (Illumina, catalog # 20020594). The libraries were
barcoded by TruSeq® RNA Unique Dual Indexes (Illumina, catalog # 20022371) and quantified
by qPCR on a Bio-Rad CFX Connect Real-Time PCR detection system with the NEBNext
Library Quant Kit for Illumina (New England Biolabs, catalog # E7630L). The libraries were
normalized to 10 nM, pooled and sequenced.
842
RNA sequencing and data analyses
RNA sequencing was carried out at Johns Hopkins School of Medicine Genetic Resources Core
Facility on a NovaSeq 6000 sequencing system (Illumina) with a S1 flow cell for 200 cycles,
generating 100 bp paired-end reads. Sequencing data processing was carried out by the Johns
Hopkins Computational Biology Consulting Core. The statistics for sequencing data analysis are
presented in supplementary tables 1-3. The RNA sequencing data were submitted to GEO
repository and were assigned record number GSE132520.
850
Biological interpretation of sequencing results was carried out with the Ingenuity Pathway
Analysis (IPA) platform (QIAGEN Ingenuity Systems, Redwood CA, USA) at the Johns
Hopkins Deep Sequencing & Microarray Core Facility. The differentially expressed genes (over
2σ) between exercised and sedentary groups within the same genotypes and between the MM
and VV samples after exercise were mapped by IPA to known pathways and biological functions
of a curated Knowledge Base (QIAGEN Bioinformatics IPA Winter Release 2018). IPA
evaluated the statistical significance of over-representation of the differentially expressed genes
in each pathway or biological function, and a *P*-value < 0.05 was defined as statistically
significant. In addition, IPA calculated the whether a pathway or biological function can be
considered activated ($z \geq 2.0$), inactivated ($z \leq -2.0$) or bidirectional altered ($-2.0 < z < 2.0$)
based the up- and down-regulation of genes involved the pathway or biological function.
862

RT-qPCR
For samples from mice, we converted 1 µg of total RNA from each sample into cDNA with the
iScript™ Reverse Transcription Supermix (Bio-Rad, Hercules, CA catalog # 1708840). 2 ng of
cDNA were used in each qPCR reaction on a CFX Connect Real-time PCR detection system
(Bio-Rad, Hercules, CA) with SsoAdvanced™ Universal SYBR® Green Supermix (Bio-Rad,
Hercules, CA catalog # 1725271). The primers for *CaMK2b* (qMmuCID0021273), *CaMK2d*
(qMmuCIP0030149), and *CaMK2g* (qMmuCIP0030022) were pre-validated primePCR primers
from Bio-Rad. The *Camk2a* primers (AGGTGTGTGAAGGTGCTGG, and
TGGAGTCGGACGATATTGGG) were designed with NCBI primer design tool
(https://www.ncbi.nlm.nih.gov/tools/primer-blast/) and validated in house to recognize all
splicing variants containing the kinase domain. qPCR data were analysed by the software Bio-
Rad CFX Manager 3.1, using *Gapdh* expression as the loading control.
875
For flies, the total RNA from adults was prepared with Direct-Zol RNA miniprep kit with on-
column DNase treatment (ZYMO Research, #2050). cDNA synthesis and qPCR were carried out
as described above. The primers for qPCR were obtained from the FlyPrimerBank[61] and were
further validated for efficiency and specificity. Expression of RP49 was used for normalization.
The primer IDs and sequences are as follows: RP49 (PD41810:
AGCATACAGGCCCAAGATCG, TGTTGTCGATACCCTTGGGC), Thor (PD43730:
CAGATGCCCGAGGTGTACTC, CATGAAAGCCCGCTCGTAGA), Sulfiredoxin/CG6762
(PD42226: GCATCGATGAGACCCACCTG, GATCCACAGCAGGTCGATGG), Gadd45
(PD42384: GGCCTTTTGCTACGAGAACG, CGCAGTAGTCGACTAGCTGG), Socs36E
(PP11279: ATGGGTCATCACCTTAGCAAGT, TCCAGGCTGATCGTCTCTACT), GstD2
(PP27238: AAACCGCGTTTGGATTTCTCG, GTGGAGACAGTGGACAGGAT), Ank2
(PD41602: TGTGGTCATGTTAGGGTGGC, TTCAAAGCCCTTGCATTGGC), Kay
(PA60087: ACTCCAACGCTTCGTACAACGATA, CACTTGAAGTATCGGTCGTGTC), Lk6
(PA60203: CAAACGCCCAGTAACATC, GCTGTAGGACCACACGCTTGAC), Atf3
(PP9314: AAGACGCCAGAGATCCTCAAC, GCAACTGGAATGACTGCTGTC), Drep3
(PP34108: GACGATGGTTTGGACGATGC, TGTTCCTCGTGATGTCCTTGA), Htk (PP1052:
TACCTGGTACATTACACAGGCT, GTGCGAGTTTTCTGCTTGGA), SNF4Aγ (PP10614:
ACCTCCGCCAAGTTGGTTG, CGCACACCGTTGTAGACGA), Sra (PP5475:
CCGATGCACCTGATCCGAC, TTGTTCTTGCTTCTGCCGTTG), Axud1 (PP35701:
GAGATAATCGTACTAGGCGATGC, GCGGAGTCAAGAATGTTGTCAA).
896

897 **Supplementary table 1 | Statistics of differential expression analysis by Cuffdiff2**
898

#Categories	MM (sed) vs. VV (sed)	MM (ex) vs. VV (ex)	MM (sed) vs. MM (ex)	VV (sed) vs. VV (ex)
(1) gene_exp.diff	30529	30599	30598	30618
(2) OK	14628	14689	14825	14628
(3) signif	93	96	743	324
(4) signif.ann	41	46	**582**	**216**
(5) signif.ann.fpkm2	40	41	490	194
(6) signif.ann.fpkm2.logFC1.5	0	3	34	24
(7) signif.ann.fpkm5	33	31	365	157
(8) signif.ann.fpkm5.logFC1.5	0	3	26	21
(9) signif.novel	52	50	161	108
(10) signif.novel.fpkm2	29	27	77	44
(11) signif.novel.fpkm2.logFC1.5	20	25	47	33

899 [#](1) Number of loci; (2) Testable loci; (3) Significant genes (P-val ≤ 0.05, q-val ≤ 0.05); (4)
900 Significant and annotated (known) genes; of which: (5) at least one of FPKM1 and FPKM2 ≥
901 2.0; (6) additionally, log2 fold change ≥ 1.5; (7) at least one of FPKM1 and FPKM2 ≥ 5.0; (8)
902 additionally, log2 fold change ≥1.5; (9) Significant and unannotated ('novel') loci; of which:
903 (10) at least one of FPKM1 and FPKM2 ≥ 2.0; (11) additionally, log2 fold change ≥1.5.
904

907 **Supplementary table 2 | Alignment statistics of RNA sequencing reads[#]**

Sample	InputReads	Aligned (R1+R2)	(R1+R2)%	Concordant%
VV1 (sed)	41,945,048	39,070,968	93.1	91.90%
VV2 (sed)	42,016,122	38,807,529	92.3	91.00%
VV3 (sed)	46,189,514	41,951,595	90.8	89.40%
VV4 (sed)	43,144,230	39,254,593	90.9	89.60%
VV5 (ex)	47,807,944	42,458,900	88.8	87.30%
VV6 (ex)	38,539,280	35,076,702	91.0	89.60%
VV7 (ex)	49,663,280	44,645,070	89.8	87.90%
VV8 (ex)	49,188,874	44,338,048	90.1	88.80%
MM1 (sed)	46,952,868	43,008,663	91.5	90.20%
MM2 (sed)	46,419,456	41,927,251	90.3	88.90%
MM3 (sed)	41,815,766	39,202,326	93.7	92.50%
MM4 (sed)	48,554,206	45,274,822	93.2	91.90%
MM5 (ex)	42,053,606	38,226,221	90.8	89.50%
MM6 (ex)	39,374,142	36,293,822	92.1	90.90%
MM7 (ex)	50,489,586	46,161,870	91.4	90.20%
MM8 (ex)	47,541,966	43,658,172	91.8	90.60%

908 [#] Reads were aligned with Tophat2.

912 **Supplementary table 3 | RNA sequencing read classification**

Sample	Aligned	Intergenic	Intron	Exon	Exon-intron	(Ex+Ex-in) %
VV1 (sed)	39,070,968	10,871,032	1,151,817	20,572,924	6,475,195	69.2
VV2 (sed)	38,807,529	10,173,751	1,056,398	20,822,770	6,754,610	71.1
VV3 (sed)	41,951,595	9,850,891	1,100,707	23,295,361	7,704,636	73.9
VV4 (sed)	39,254,593	9,358,135	1,067,585	21,523,146	7,305,727	73.4
VV5 (ex)	42,458,900	9,795,942	1,104,903	23,546,149	8,011,906	74.3
VV6 (ex)	35,076,702	8,350,481	1,020,203	19,380,828	6,325,190	73.3
VV7 (ex)	44,645,070	12,762,540	1,450,013	22,758,083	7,674,434	68.2
VV8 (ex)	44,338,048	12,196,685	1,273,665	23,333,460	7,534,238	69.6
MM1 (sed)	43,008,663	14,389,489	1,223,644	20,909,218	6,486,312	63.7
MM2 (sed)	41,927,251	11,612,157	1,202,400	22,166,915	6,945,779	69.4
MM3 (sed)	39,202,326	9,690,241	1,091,394	21,601,873	6,818,818	72.5
MM4 (sed)	45,274,822	11,798,947	1,183,309	24,310,729	7,981,837	71.3
MM5 (ex)	38,226,221	9,375,367	1,089,510	20,889,204	6,872,140	72.6
MM6 (ex)	36,293,822	9,136,213	1,069,954	19,752,989	6,334,666	71.9
MM7 (ex)	46,161,870	12,957,951	1,406,060	24,129,718	7,668,141	68.9
MM8 (ex)	43,658,172	12,897,911	1,309,884	22,353,997	7,096,380	67.5

913
914
915
916

917
918
919 **References**
920

1   Anderson, M. E., Brown, J. H. & Bers, D. M. CaMKII in myocardial hypertrophy and
heart failure. *J Mol Cell Cardiol* **51**, 468-473, doi:10.1016/j.yjmcc.2011.01.012 (2011).
2   Erickson, J. R. *et al.* A dynamic pathway for calcium-independent activation of CaMKII
by methionine oxidation. *Cell* **133**, 462-474, doi:10.1016/j.cell.2008.02.048 (2008).
3   Erickson, J. R. *et al.* Diabetic hyperglycaemia activates CaMKII and arrhythmias by O-
linked glycosylation. *Nature* **502**, 372-376, doi:10.1038/nature12537 C2 - PMC3801227
(2013).
4   Lou, L. L., Lloyd, S. J. & Schulman, H. Activation of the multifunctional
Ca2+/calmodulin-dependent protein kinase by autophosphorylation: ATP modulates
production of an autonomous enzyme. *Proc Natl Acad Sci U S A* **83**, 9497-9501 (1986).
5   Miller, S. G. & Kennedy, M. B. Regulation of brain type II Ca2+/calmodulin-dependent
protein kinase by autophosphorylation: a Ca2+-triggered molecular switch. *Cell* **44**, 861-
870 (1986).
6   Purohit, A. *et al.* Oxidized Ca(2+)/calmodulin-dependent protein kinase II triggers atrial
fibrillation. *Circulation* **128**, 1748-1757, doi:10.1161/CIRCULATIONAHA.113.003313
(2013).
7   Luo, M. *et al.* Diabetes increases mortality after myocardial infarction by oxidizing
CaMKII. *J Clin Invest* **123**, 1262-1274, doi:10.1172/JCI65268 (2013).
8   Hart, P. C. *et al.* MnSOD upregulation sustains the Warburg effect via mitochondrial
ROS and AMPK-dependent signalling in cancer. *Nat Commun* **6**, 6053,
doi:10.1038/ncomms7053 (2015).
9   Sanders, P. N. *et al.* CaMKII is essential for the proasthmatic effects of oxidation.
Science Translational Medicine **5**, 195ra197, doi:10.1126/scitranslmed.3006135 C2 -
PMC4331168 (2013).
10   Gu, S. X. *et al.* Protein methionine oxidation augments reperfusion injury in acute
ischemic stroke. *JCI Insight* **1**, doi:10.1172/jci.insight.86460 (2016).
11   Qu, J. *et al.* Oxidized CaMKII promotes asthma through the activation of mast cells. *JCI*
Insight **2**, e90139, doi:10.1172/jci.insight.90139 (2017).
12   Swaminathan, P. D. *et al.* Oxidized CaMKII causes cardiac sinus node dysfunction in
mice. *J Clin Invest* **121**, 3277-3288, doi:10.1172/JCI57833 (2011).
13   Wu, Y., Wang, Q., Feng, N., Granger, J. M. & Anderson, M. E. Myocardial death and
dysfunction after ischemia-reperfusion injury require CaMKIIdelta oxidation. *Sci Rep* **9**,
9291, doi:10.1038/s41598-019-45743-6 (2019).
14   Bus, J. S. & Gibson, J. E. Paraquat: model for oxidant-initiated toxicity. *Environ Health*
Perspect **55**, 37-46, doi:10.1289/ehp.845537 (1984).
15   Gee, H. *Across the bridge : understanding the origin of the vertebrates.* (The University
of Chicago Press, 2018).
16   Green, S. A., Simoes-Costa, M. & Bronner, M. E. Evolution of vertebrates as viewed
from the crest. *Nature* **520**, 474-482, doi:10.1038/nature14436 (2015).
17   Rose, A. J., Alsted, T. J., Kobbero, J. B. & Richter, E. A. Regulation and function of
Ca2+-calmodulin-dependent protein kinase II of fast-twitch rat skeletal muscle. *J Physiol*
580, 993-1005, doi:10.1113/jphysiol.2006.127464 (2007).

18   Tavi, P. *et al.* Calmodulin kinase modulates Ca2+ release in mouse skeletal muscle. *J*
Physiol **551**, 5-12, doi:10.1113/jphysiol.2003.042002 (2003).
19   Regot, S., Hughey, J. J., Bajar, B. T., Carrasco, S. & Covert, M. W. High-sensitivity
measurements of multiple kinase activities in live single cells. *Cell* **157**, 1724-1734,
doi:10.1016/j.cell.2014.04.039 C2 - PMC4097317 (2014).
20   Morioka, E., Kanda, Y., Koizumi, H., Miyamoto, T. & Ikeda, M. Histamine Regulates
Molecular Clock Oscillations in Human Retinal Pigment Epithelial Cells via H-1
Receptors. *Frontiers in Endocrinology* **9**, doi:ARTN 108
10.3389/fendo.2018.00108 (2018).
21   Hanson, P. I., Meyer, T., Stryer, L. & Schulman, H. Dual role of calmodulin in
autophosphorylation of multifunctional CaM kinase may underlie decoding of calcium
signals. *Neuron* **12**, 943-956 (1994).
22   Chang, B. H., Mukherji, S. & Soderling, T. R. Characterization of a calmodulin kinase II
inhibitor protein in brain. *Proc Natl Acad Sci U S A* **95**, 10890-10895,
doi:10.1073/pnas.95.18.10890 (1998).
23   Shen, T. *et al.* DNA binding sites target nuclear NFATc1 to heterochromatin regions in
adult skeletal muscle fibers. *Histochem Cell Biol* **134**, 387-402, doi:10.1007/s00418-010-
0744-4 (2010).
24   Noble, B. J., Borg, G. A., Jacobs, I., Ceci, R. & Kaiser, P. A category-ratio perceived
exertion scale: relationship to blood and muscle lactates and heart rate. *Med Sci Sports*
Exerc **15**, 523-528 (1983).
25   Fan, W. *et al.* PPARdelta Promotes Running Endurance by Preserving Glucose. *Cell*
Metab **25**, 1186-1193 e1184, doi:10.1016/j.cmet.2017.04.006 (2017).
26   Ozcan, L. *et al.* Calcium signaling through CaMKII regulates hepatic glucose production
in fasting and obesity. *Cell Metabolism* **15**, 739-751, doi:10.1016/j.cmet.2012.03.002 C2
- PMC3348356 (2012).
27   Holloszy, J. O. & Kohrt, W. M. Regulation of carbohydrate and fat metabolism during
and after exercise. *Annu Rev Nutr* **16**, 121-138,
doi:10.1146/annurev.nu.16.070196.001005 (1996).
28   Iyer, S. R., Valencia, A. P., Hernandez-Ochoa, E. O. & Lovering, R. M. In Vivo
Assessment of Muscle Contractility in Animal Studies. *Methods in molecular biology*
(Clifton, N.J.) **1460**, 293-307, doi:10.1007/978-1-4939-3810-0_20 (2016).
29   Liu, Y., Hernandez-Ochoa, E. O., Randall, W. R. & Schneider, M. F. NOX2-dependent
ROS is required for HDAC5 nuclear efflux and contributes to HDAC4 nuclear efflux
during intense repetitive activity of fast skeletal muscle fibers. *Am J Physiol Cell Physiol*
303, C334-347, doi:10.1152/ajpcell.00152.2012 (2012).
30   Allen, D. G., Lamb, G. D. & Westerblad, H. Skeletal muscle fatigue: cellular
mechanisms. *Physiological Reviews* **88**, 287-332, doi:10.1152/physrev.00015.2007
(2008).
31   Jackson, S. H., Gallin, J. I. & Holland, S. M. The p47phox mouse knock-out model of
chronic granulomatous disease. *J Exp Med* **182**, 751-758 (1995).
32   Khairallah, R. J. *et al.* Microtubules underlie dysfunction in duchenne muscular
dystrophy. *Sci Signal* **5**, ra56, doi:10.1126/scisignal.2002829 (2012).
33   Powers, S. K., Talbert, E. E. & Adhihetty, P. J. Reactive oxygen and nitrogen species as
intracellular signals in skeletal muscle. *J Physiol* **589**, 2129-2138,
doi:10.1113/jphysiol.2010.201327 (2011).

34   Prosser, B. L., Khairallah, R. J., Ziman, A. P., Ward, C. W. & Lederer, W. J. X-ROS
signaling in the heart and skeletal muscle: stretch-dependent local ROS regulates
[Ca(2)(+)]i. *J Mol Cell Cardiol* **58**, 172-181, doi:10.1016/j.yjmcc.2012.11.011 (2013).
35   Chin, E. R. The role of calcium and calcium/calmodulin-dependent kinases in skeletal
muscle plasticity and mitochondrial biogenesis. *The Proceedings of the Nutrition Society*
63, 279-286, doi:10.1079/PNS2004335 (2004).
36   Egan, B. & Zierath, J. R. Exercise metabolism and the molecular regulation of skeletal
muscle adaptation. *Cell Metabolism* **17**, 162-184, doi:10.1016/j.cmet.2012.12.012 (2013).
37   Kramer, A., Green, J., Pollard, J., Jr. & Tugendreich, S. Causal analysis approaches in
Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-530,
doi:10.1093/bioinformatics/btt703 (2014).
38   Peake, J., Nosaka, K. & Suzuki, K. Characterization of inflammatory responses to
eccentric exercise in humans. *Exerc Immunol Rev* **11**, 64-85 (2005).
39   Ling, H. *et al.* Ca2+/Calmodulin-dependent protein kinase II delta mediates myocardial
ischemia/reperfusion injury through nuclear factor-kappaB. *Circ Res* **112**, 935-944,
doi:10.1161/CIRCRESAHA.112.276915 (2013).
40   Singh, M. V. *et al.* Ca2+/calmodulin-dependent kinase II triggers cell membrane injury
by inducing complement factor B gene expression in the mouse heart. *J Clin Invest* **119**,
986-996, doi:10.1172/JCI35814 (2009).
41   Suetomi, T. *et al.* Inflammation and NLRP3 Inflammasome Activation Initiated in
Response to Pressure Overload by Ca(2+)/Calmodulin-Dependent Protein Kinase II delta
Signaling in Cardiomyocytes Are Essential for Adverse Cardiac Remodeling. *Circulation*
138, 2530-2544, doi:10.1161/CIRCULATIONAHA.118.034621 (2018).
42   Cai, B. *et al.* MerTK signaling in macrophages promotes the synthesis of inflammation
resolution mediators by suppressing CaMKII activity. *Sci Signal* **11**,
doi:10.1126/scisignal.aar3721 (2018).
43   Liu, X. *et al.* CaMKII promotes TLR-triggered proinflammatory cytokine and type I
interferon production by directly binding and activating TAK1 and IRF3 in macrophages.
Blood **112**, 4961-4970, doi:10.1182/blood-2008-03-144022 (2008).
44   Huang, W., Ghisletti, S., Perissi, V., Rosenfeld, M. G. & Glass, C. K. Transcriptional
integration of TLR2 and TLR4 signaling at the NCoR derepression checkpoint. *Mol Cell*
35, 48-57, doi:10.1016/j.molcel.2009.05.023 (2009).
45   Huang, W. *et al.* Coronin 2A mediates actin-dependent de-repression of inflammatory
response genes. *Nature* **470**, 414-418, doi:10.1038/nature09703 (2011).
46   Bui, J. D. *et al.* A Role for CaMKII in T Cell Memory. *Cell* **100**, 457-467,
doi:10.1016/s0092-8674(00)80681-9 (2000).
47   Peckham, M., Molloy, J. E., Sparrow, J. C. & White, D. C. Physiological properties of
the dorsal longitudinal flight muscle and the tergal depressor of the trochanter muscle of
Drosophila melanogaster. *J Muscle Res Cell Motil* **11**, 203-215 (1990).
48   Reiner, D. J., Newton, E. M., Tian, H. & Thomas, J. H. Diverse behavioural defects
caused by mutations in Caenorhabditis elegans unc-43 CaM kinase II. *Nature* **402**, 199-
203, doi:10.1038/46072 (1999).
49   Brown, J. B. *et al.* Diversity and dynamics of the Drosophila transcriptome. *Nature* **512**,
393-399, doi:10.1038/nature12962 (2014).

50   Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. *Mol Biol Evol* **34**, 1812-1819,
doi:10.1093/molbev/msx116 (2017).

51   Backs, J., Song, K., Bezprozvannaya, S., Chang, S. & Olson, E. N. CaM kinase II
selectively signals to histone deacetylase 4 during cardiomyocyte hypertrophy. *The*
Journal of Clinical Investigation **116**, 1853-1864, doi:10.1172/JCI27438 C2 -
PMC1474817 (2006).

52   Ryder, E. *et al.* The DrosDel collection: a set of P-element insertions for generating
custom chromosomal aberrations in Drosophila melanogaster. *Genetics* **167**, 797-813,
doi:10.1534/genetics.104.026658 (2004).

53   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-
1549, doi:10.1093/molbev/msy096 (2018).

54   Bloemberg, D. & Quadrilatero, J. Rapid determination of myosin heavy chain expression
in rat, mouse, and human skeletal muscle using multicolor immunofluorescence analysis.
PLoS One **7**, e35273, doi:10.1371/journal.pone.0035273 (2012).

55   Vogler, G. & Ocorr, K. Visualizing the beating heart in Drosophila. *J Vis Exp*,
doi:10.3791/1425 (2009).

56   Fink, M. *et al.* A new method for detection and quantification of heartbeat parameters in
Drosophila, zebrafish, and embryonic mouse hearts. *Biotechniques* **46**, 101-113,
doi:10.2144/000113078 (2009).

57   Cammarato, A., Ocorr, S. & Ocorr, K. Enhanced assessment of contractile dynamics in
Drosophila hearts. *Biotechniques* **58**, 77-80, doi:10.2144/000114255 (2015).

58   Rizzo, M. A., Springer, G. H., Granada, B. & Piston, D. W. An improved cyan
fluorescent protein variant useful for FRET. *Nat Biotechnol* **22**, 445-449,
doi:10.1038/nbt945 (2004).

59   McQuin, C. *et al.* CellProfiler 3.0: Next-generation image processing for biology. *PLoS*
Biol **16**, e2005970, doi:10.1371/journal.pbio.2005970 (2018).

60   Cohen, T. V., Kollias, H. D., Liu, N., Ward, C. W. & Wagner, K. R. Genetic disruption
of Smad7 impairs skeletal muscle growth and regeneration. *J Physiol* **593**, 2479-2497,
doi:10.1113/JP270201 (2015).

61   Hu, Y. *et al.* FlyPrimerBank: an online database for Drosophila melanogaster gene
expression analysis and knockdown evaluation of RNAi reagents. *G3 (Bethesda, Md.)* **3**,
1607-1616, doi:10.1534/g3.113.007021 (2013).

1087