High-resolution synchrotron imaging studies of intact fresh roots reveal soil bacteria promoted bioremediation and bio-fortification

KEYWORDS

Bioremediation, Selenium, Iron, Plant uptake, Micronutrients, Synchrotron Imaging

AUTHORS

Hanna Help [1] [*], Merja Lusa [2] [†], Ari-Pekka Honkanen [1] [†], Ana Diaz [3] [‡], Mirko Holler [3] [‡], Murielle Salomé [4], Peter Cloetens [4], Henrik Mäkinen [1], Simo Huotari [1] and Heikki Suhonen [1]

1 Department of Physics, P.O.Box 64, FI-00014 University of Helsinki, Finland
2 Department of Chemistry, P.O.Box 55, FI-00014 University of Helsinki, Finland
3 Paul Scherrer Institut, Forschungsstrasse 111, 5232 Villigen PSI, Switzerland
4 ESRF - The European Synchrotron, CS 40220, 38043 Grenoble Cedex 9, France
* Corresponding author
† These authors contributed equally
‡ These authors contributed equally

Author ORCiD IDs:
H.H. 0000-0003-2501-9851
M.L. 0000-0001-5709-7753
A.-P.H. 0000-0002-6822-3062
A.D. 0000-0003-0479-4752

M.H. 0000-0001-8141-0148
P.C. 0000-0002-4129-9091
H.M. 0000-0003-0101-2946
S.H. 0000-0003-4506-8722
H.S.0000-0002-0337-6308

ABSTRACT

Plant-microbe interactions can be utilized in bio-based processes such as bioremediation and biofortification, either to remove hazardous radionuclides and heavy metals from the soil, or to increase the accumulation of desired elements into crops to improve their quality. Optimizing such elegant plant-microbe interactions requires detailed understanding of the chemical element compositions of fresh plant tissues at cellular organelle resolution. However, such analyses remain challenging because conventional methods lack the required spatial resolution, contrast or sensitivity. Using a novel combination of nanoscale-resolution 3D cryogenic synchrotron-light ptychography, holotomography and fluorescence tomography, we show how soil bacteria interact with *Arabidopsis thaliana* and promote the uptake of various metals. Co-cultivation with *Pseudomonas* sp. strain T5-6-I alters root anatomy and increases levels of selenium (Se), iron (Fe) and other micronutrients in roots. Our approach highlights the interaction of plants and microbes in bioremediation and bio-fortification on the subcellular level.

MAIN TEXT

Soilborne bacteria can modify soil chemical landscapes altering the mobility and uptake rates of various chemical elements, including hazardous radionuclides and heavy metals, but also macro- and micro-nutrients into plant tissues (Ojuederie *et al.,* 2018, Desai *et al.,*

2019). For tailoring plant-microbe interactions to improve the efficiency of bio-based processes like bioremediation and biofortification, conclusive data on chemical element composition from intact fresh plant tissues at cellular organelle resolution is required. Currently, however, there is a lack of quantitative knowledge on the ultimate fate of accumulated elements in plants on a subcellular level. This is unfortunate, as understanding in which tissue and cellular organelle various chemical elements and species accumulate would help increase their uptake and translocation rates e.g. by modifying tissue and organelle specific transporter activity via genetic engineering.

Selenium (Se) is an essential micronutrient for mammals, but in high-valence states it is toxic at elevated concentrations. In alkaline agricultural soils, selenium mostly exists as selenate ($Se(VI)O_4^{2-}$) whereas in acidic forest soils it exists as selenite ($Se(IV)O_3^{2-}$). The environmental releases of Se in these forms is potentially hazardous (Gupta & Gupta, 2016). Selenate and selenite differ in their absorption and mobility within the plant (Li *et al.,* 2008). As the environmental mobility and biological effects of Se are mainly dictated by its chemical speciation, it is interesting that the chemical speciation can be affected by some soilborne bacteria, including *Pseudomonas* sp. strain T5-6-I, via reduction and oxidation (redox) reactions. T5-6-I forms intracellular aggregates of reduced Se^0 when incubated in $Se(IV)O_3^{2-}$ containing solutions, and co-cultivation with plants enhances Se(IV) uptake and accumulation *in planta* (Lusa *et al.,* 2015, 2017 and 2019). Indirect evidence suggests that in plants Se accumulates in cellular vacuoles (Gupta & Gupta, 2016, Santa Cruz *et al.,* 2017) as it does in human tissues (Bodnar *et al.,* 2017). Transmission electron microscopy (TEM) (Durán *et al.,* 2018, Lusa *et al.,* 2019) has revealed aggregate formation in plant cell compartments after Se treatments. However, it was not concluded if the observed aggregates were selenium. While direct chemical element localization has been done from fresh plant roots (Wang *et al.,* 2013, Hu *et al.,* 2014, Zhao *et al.,* 2014, Krajcarová *et al.,* 2017) and seeds (Rodrigues *et al.,* 2018) using synchrotron radiation x-ray fluorescence microscopy, the resolution has not been sufficient to determine subcellular element concentrations. Using sectioning techniques and 2D imaging methods, such as nano-secondary ion mass spectrometry (nano SIMS), laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS), or ColorEM (electron microscope) one could potentially reach desired resolution and sensitivity (Ondrasek *et al.,* 2019, Pirozzi *et al.,* 2018). However, these techniques can only show small fraction of the total volume and are subject to artefacts from the sectioning process. Therefore, the question whether plants form intracellular vacuole localized Se granules in their roots, similarly to *Pseudomonas* sp. strain or humans, or if Se is deposited elsewhere in plant tissues (e.g. cell walls) has not been answered conclusively.

We established the role of bacteria on the accumulation and distribution of Se in the elongation zone of roots, the key uptake sites for water and nutrients, by using multimodal synchrotron nanotomography on fresh *Arabidopsis thaliana* root material. We did this by imaging frozen native state specimens cultivated in Se(IV) and Se(IV) + *Pseudomonas* sp. strain T5-6-I supplemented growth media with three state-of-the-art synchrotron x-ray imaging techniques: X-ray ptychography (PXCT) (Dierolf *et al.,* 2010) and

nanoholotomography (XNH) (da Silva *et al.* 2018), both giving high resolution 3D images of
the electron density, as well as x-ray nanofluorescence microscopy (XRNF) giving
quantitative, high resolution 2D and 3D images of trace element distributions in the plants.
These techniques gave subcellular organelle resolution, allowing to address whether the
aggregates reported with EM studies (Durán *et al.,* 2018, Lusa *et al.,* 2019) are indeed
composed of Se.
98
Our findings show that, even though Se levels varied between different treatments, tissue
profiles of Se and Se+bacteria treated samples were similar. Most importantly; soil bacteria
appeared to promote Se accumulation in young *Arabidopsis thaliana* lateral root tips. Using
tissue segmentation analysis and subcellular measurements, we were able to determine
that Se concentrates predominantly to cytoplasmic domains, cell walls and vacuolic
membranes in roots. Based on our results we were able to conclude that Se does form
aggregates in *Arabidopsis thaliana* plants, however not in a similar manner as has been
reported with bacterial or mammalian cells, where Se aggregates predominantly in vacuolic
compartments. Interestingly, bacterial co-culturing also promoted accumulation of Fe and
other micronutrients into root tissues. Fe was predominantly localized in cell walls showing
different localization pattern than Se. The amount of Fe in bacteria co-cultured root tissues
exceeded Se levels by a magnitude and could partially explain the observed increase in root
cell wall density. From an agricultural point, soil bacteria promoted Fe biofortification would
be a cost-effective way to improve the nutritional value of crops growing in micronutrient
poor soils.
114

RESULTS

Root tissue densities and physical features differ between treatments

Overall, in all imaged lateral root samples (Figure 1 A), cell walls had the highest electron
density, followed by vacuole membranes, cytoplasm (including endoplasmic reticulum and
different cell organelles like mitochondria) and vacuolic fluid (least dense) (Figure 2,
Extended Figure 1). Interestingly, we found that bacteria increased cell wall densities by
>7% compared to control samples, while the cytoplasm and vacuolic fluid densities were
similar in all samples irrespective of treatment (Table 1). The vacuolic membrane densities
of the Se+bacteria treated root were ~16% higher than in Se-alone supplemented sample.
This difference is large enough to be directly seen in electron density visualizations (Figure
2, panels I and L).
126
The ground tissues (the two outermost cell layers, epidermis and underlying cortex cells) of
control roots (Figure 2 A and D) appeared denser than the underlying endodermal and stele
cells. Se treated roots (Figure 2 B and E) did not exhibit obvious differences in tissue density
profiles between the cell types. Se+bacteria supplemented roots exhibit higher density of
endodermis, pericycle and the stele cells (vasculature) than in the control or Se
supplemented plants. In addition, Se+bacteria supplemented roots exhibited pronounced
cytoplasmic and vacuolic membrane features and numerous high-density aggregates,
mainly in ground tissue layers (Figure 2, panels K and L). Apart from density features, 3D
visualizations of the datasets revealed differences in cellular differentiation rates between

control, Se supplemented and Se+bacteria supplemented plants (Figure 3). Compared to control root (Figure 3 A, D), Se treated roots exhibited enhanced vacuolization of cortical cells, but no other visible abnormalities (Figure 3 B, E). On the contrary, Se+bacteria supplemented roots exhibited both enhanced vacuolic expansion and abnormal cell division patterns (Figure 3, C and F). These aberrations in cellular division and differentiation rates appeared to correlate with the increased numbers of root hairs in *Arabidopsis* roots (Figure 3 panels G, H and I), similar to what was reported on kale (Lusa *et al.,* 2019).

Micronutrient accumulation differs between treatments

We found that not only the uptake of Se is modified by bacteria, but most notably that Fe uptake is promoted, and to smaller extent also that of Br and Cr (Figure 4, Extended Figure 2). The uptake or distribution of other detected elements as Al, Au, Ca, Cl, Cu, K, Mn, Ni, P, Pb and S was not modified by the presence of Se or bacteria. In Figure 4 the distinct cellular features of the XNH datasets are matched to XRNF-microscopy maps to visualize Se, Fe, Cr and Ca distribution (Figure 4). Top row of figure 4 illustrates the electron density slices that correspond to chemical element maps (1, control root; 2 and 3, Se supplemented roots; 4 and 5 Se+bacteria supplemented roots). As expected, control roots did not contain Se above the detection limit (Graph 1), while considerable levels of Se were detected in all root tissues of plants supplemented with Se. Overall, the tissue Se profiles of roots supplemented with Se-alone were comparable to Se+bacteria roots – however clear root-to-root variation in the overall levels of Se was seen between biological replicates of Se-alone and Se+bacteria treated samples (Graph 1). On tissue level, accumulation of Se appears quite even between intracellular and cell wall areas in Se and Se+bacteria supplemented roots (Graph 2). The Se+bacteria supplemented root #2 contained more Se than other measured samples. Interestingly, the Fe levels in control and Se supplemented roots were very low compared to Se+bacteria supplemented roots, which contained over 10-times more Fe than the other measured samples (Graphs 1 and 2). Unexpectedly, the levels of Fe of Se supplemented roots were even lower than in control root, with the fluorescent signal of Se supplemented roots localized mainly to the superficial cell facing growing medium (Figure 5, Graph 2).

To better understand the element accumulation at subcellular level, close ups of Se and Fe maps were overlaid with electron density images (Figure 5). Again, control root (A) showed no sign of Se and low levels of Fe. In Se supplemented roots (B and C) Se appeared to be unevenly distributed, and Fe levels were barely detectable. In Se+bacterial supplemented roots (D and E) both Se and Fe signals were clearly visible. On subcellular level Se appeared visually to be primarily located in cytoplasm and internal membranes in various root cells. In bacteria co-cultured roots, apart from cytoplams and vacuoles, Se was also visually detected in some cell walls (Figure 5 D), supporting data presented in Graph 2. In all samples from which Fe could be detected, Fe signal was localized almost exclusively to cell walls and hyperaccumulated into the junctions of cells, showing a very different localization pattern than Se (Figure 5, Graphs 2 and 3). Following visual inspections, Se and Fe levels were quantified at different subcellular compartments from the XRNF maps (masks of the selected areas and measurements are shown in Graph 3). Results indicated that in all samples, cell

walls & cytoplasmic areas contained on average equal levels of Se, which exceed Se levels
in vacuolic compartments (Graph 3). Within vacuoles, the membranes contained more Se
than fluid filled areas, at least in the case of Se+bacteria supplemented root #1 that exhibited
clearest subcellular compartmentalization required in such fine detail measurements
(Extended Figure 3). Segmentation analysis showed that in all measured samples Fe was
predominantly localized to cell walls (Graph 3), supporting visual observations. When the
Se and Fe levels of Se+bacteria supplemented roots were compared on across all tissues,
a ~3.5-fold difference was observed in averaged mean values between Fe and Se. In cell
walls specifically, the difference was even more pronounced as mean Fe levels were >12-
times higher than Se levels (online material: Se_Fe statistics.xlsx). Levels of Cr and Br
accumulation were also increased in cells of bacteria co-cultured roots – however the
increase was not as high as for Fe. Cr appeared to be mainly localized to cell walls, while
Br was more prominent within the cells (Extended figure 2). While the overall Ca signal
intensity in cell walls appeared similar for all treatments (Figure 4, Extended Figure 2), few
local higher Ca signal peaks were observed in cell walls of damaged epidermal cells of
Se+bacteria supplemented roots (Extended Figure 4). The XNH 3D surface reconstruction
of such damaged area is shown in Extended Figure 4. The concentration of Ca in these
locations was ~2.5 times higher than in neighboring epidermal cell walls. It is not clear
whether the damage was caused by the *Pseudomonas* bacteria visible on the surface of the
root – or if it was caused by sample preparation and mounting onto cryo-sample holder.
200
Finally, Se-Fe density scatter plots were generated from each sample (Extended Figure 5)
and used to segment root areas based on their densities. When plotted between Se and Fe
allocation domains, the plotted areas were in line with the XRNF maps (Figures 4 and 5). It
was initially assumed that the dense intracellular aggregates located at vacuolic membranes
and around nuclei of many samples, especially visible in Se+bacteria supplemented root
(Extended figure 6), were Se enriched protein aggregates (as was suggested by Gupta &
Gupta*,* 2016). However, the majority of these aggregates seen in both PXCT and XNH
datasets did not match with XRNF Se maps, indicating that the aggregates consist of other
chemical element(s). Interestingly, both Zn and Ca aggregates were found in the epidermal
cells of most imaged samples (shown in Extended Figure 2), localizing mainly to vacuoles.
In many cases these elements matched to the dense aggregates measured using XNH,
indicating that these micronutrients can make dense aggregates in plant cells (as was
suggested for Zn by Arrivault *et al.,* 2006)(data not shown).
214
DISCUSSION
Bacterial co-culturing affects root anatomy, tissue density and chemical composition
The electron density data was used to 1) visualize tissue and subcellular organelle anatomy
in fine detail and 2) measure tissue densities within intact 3D samples. Each co-centric root
tissue type (epidermis, cortex, endodermis, pericycle and stele, illustrated in Figure 2 and
Extended Figure 7) could be identified based on their position, shape and cell numbers.
Visualization of subcellular contents allowed comparisons between treatments in processes
like cellular clearing and increase in vacuolic size and water content (seen especially
endodermis, pericycle and stele in control roots), which occur during cellular differentiation,

as the volume of cytoplasmic domain reduces upon cellular differentiation (Perrot-Rechenmann, 2010). For example, Se+bacteria supplemented roots exhibited altered cellular division and differentiation rates and higher density of vascular tissues (Figure 2) compared to both control and Se supplemented roots. Their increased intracellular density could possibly be attributed to 1) higher levels of sugars, fats and proteins in cytoplasm, 2) proportionally larger cytoplasmic cell fractions in cells, and to 3) increased number and size of subcellular aggregates (especially in epidermis and cortex cells). In addition to intracellular cytoplasmic density increase, Se+bacteria treated roots showed increased density of extracellular cell walls (table 1). This finding could at least partially attribute to the fact that the bacteria promote Fe fortification of cell walls. While some Se *did* localize to cell walls in bacteria supplemented roots, the difference in magnitude indicates that Se is not a major contributor to the observed cell wall density differences between different treatments (Fe levels in roots supplemented with Se+bacteria were >12-times higher than Se levels). Oppositely, as the electron densities of Se+bacteria treated root vacuolic membranes were ~17 higher than in control root, *and* Fe did not accumulate into vacuoles, it is quite possible that the observed increase in vacuolic membrane density in bacteria co-cultured roots was attributed primarily to Se aggregate formation.

Se toxicity and bacteria elicitor promoted growth abnormalities

The observed treatment specific differences in root anatomy, including differences in root size (diameter), defects in maintaining proper cell divisions, and growth arrest, both in Se and Se+bacteria supplemented roots compared to control roots, could in part be attributed to biotic Se stress (toxicity) as the Se supplemented roots showed slight reduction in growth (Lusa *et al.,* 2019, Extended Figure 8). However, as the cell division defects, and meristem differentiation abnormalities were exaggerated in the Se+bacteria supplemented roots (Figure 3), it is unlikely that the phenotypes were caused by Se toxicity alone. While deviations from normal meristem patterning could at least partially explain the observed increased root hair phenotype seen in bacteria co-cultured *Arabidopsis thaliana* roots (Figure 3), there is an additional explanation to the observed pleiotropic root phenotype: the bacteria could possibly excrete plant phytohormone like elicitors to the growing media. Presence of bioactive auxin (or inactive precursors or auxin like compounds) could explain the observed phenotypes of our Se+bacteria supplemented roots: the Se+bacteria roots had larger meristems, exhibited extra root hair formation, showed root growth arrest (seen as white meristems, Extended Figure 8) and odd root branching architecture (Extended Figure 8). In fact, analyzed roots resembled closely several classical *Arabidopsis* auxin mutants, including abnormal root architecture of *superroot* (Boerjan et al., 1995), *yucca*-mutants (Zhao *et al.,* 2001, Cheng *et al.,* 2006), and pleiotropic morphological and cellular differentiation defected *auxin resistant 1*-mutants (Lincoln *et al.,*1990). Moreover, the Se+bacteria supplemented roots showed an exposure time dependent increase in lateral root emergence phenotype (data not shown), similar to what was reported by Singh *et al.* in 2015. The bacterium induced auxin accumulation scenario is supported by the fact that several soil bacteria (similar to used *Pseudomonas* strain) are known on elicit auxin overproduction like root phenotypes in plants across the plant kingdom, most famous example being soil dwelling microbes belonging to the *Agrobacterium* genus (Esitken 1970,

Cardarelli *et al.,* 1987, Amselem & Tepfer, 1992). The exact mechanism behind the putative
auxin related phenotype remains to be determined, but could be due to auxin overproduction
in plants (via bacteria originating lateral gene transfer promoted activation and/or
overexpression of auxin biosynthesis genes) (Cardarelli *et al.,* 1987, Liu *et al.,* 2015) or
caused by external auxin stimulus produced by the bacteria themselves. Overall, it can be
concluded that the bacteria affected meristem growth (cell division and differentiation),
vascular tissue density and subcellular composition (both intra and extracellular densities),
but the exact mechanism remains unsolved.
276
Superior properties of intact fresh tissues 3D maps
Previous efforts to study the amount and localization of Se in plant tissues have relied mostly
on three very different approaches: µXRF (Wang *et al.,* 2013, Hu *et al.,* 2014, Zhao *et al.,*
2014) or spectroscopy imaging assays assays (Krajcarová *et al.,* 2017), cell organelle
fractionation, and indirect gene expression assays (Gupta & Gupta*,* 2016) in response to
exogenously applied chemical compounds. In fluorescence imaging assays conducted at
room temperature, imaged samples were either rather large (whole roots, seedlings of
leaves), fixed in resin, or freeze dried to minimize sample movement during imaging. All
these approaches have their caveats: The spatial resolution in imaging of large samples will
not be sufficient to reveal subcellular differences or fine features. Fixing a sample in resin
includes dehydration steps, infusion with resins, and possibly sectioning – leading potentially
to alteration of chemical composition and anatomical deformation. Same issues apply to
freeze drying, in which all the water is often removed using *e.g.* ethanol series followed by
lyophilisation. As water content affects critically the conformation, density, mechanical
properties and chemical composition of biological tissues, every modification will inevitably
contort the obtained data compared to samples imaged *in vivo*. While electron microscope
imaging methods can provide direct information of chemical elements in high resolution, the
samples are sectioned to thin slices. The imaged samples are also often dried to immobilize
them, loosing information on water and soluble compounds. All these described procedures
leave plenty of room for improvement on sample integrity but also on volume of information
gained from such narrow "views" of 2D slices. Together, the previously used methods have
provided mostly either low resolution, indirect or spatially limited evidence of both the
physical and chemical properties of biological samples.
300
We overcame all these limitations by studying Se uptake and accumulation in fresh
Arabidopsis thaliana root tissues using state-of-the-art synchrotron X-ray radiation based
PXCT, XNH and XRNF tomographies at about 100 nm scale resolution. All imaging assays
were conducted in cryogenic conditions in order to preserve sample features as intact as
possible for reliable chemical element analysis. When samples are frozen, the movement
caused by radiation damage, *i.e.* volatilization and reactive oxygen species formation due
to high photon flux, is minimal allowing long measuring times. Long measuring times allow
retrieval of > 1000 projections from each sample, leading to the desired image resolution of
about 100 nm. As Se can be processed to volatile compounds, sample freezing also
prevented enzymatic activity and putative loss of Se signal from samples. As a bonus, cryo-
imaging allowed inspection of transient chemical signaling events in plant tissues,

preserving fresh samples features and local chemical element concentration peaks in
samples as if the samples were in their native state.
314
High-resolution imaging resolution reveals subcellular organelle features
The reconstructed high-resolution 3D electron density volumes from PXCT and XNH
showed highly similar results on sample anatomy. This is noteworthy, as the electron density
data was obtained using two different cryogenic X-ray imaging methods and the
measurements were performed at different facilities from different biological sample
batches. One difference was that unlike PXCT, XNH datasets appeared to highlight network
like subcellular compartments (like ER and vacuolic membranes) based on their properties
that enhance edges on compartments of different texture (e.g. fat vs. water). Due to the
choice of large field of view, the XNH results were also of slightly lower resolution than the
PXCT as evidenced by the subcellular features. The XNH datasets contained also some
artefacts at samples edges (seen as "ripple effects" at the epidermal cell edges, Figure 2,
panels D – F). These artefacts could be reduced by using higher incident X-ray energy.
327
The volumes allowed inspection of individual cells and subcellular compartments in fine
detail. For example, phloem sieve elements could be identified (Figure 2 G and Extended
Figure 6) based on the cell wall properties and developmental status of different cells in the
stele. In *Arabidopsis thaliana* root, phloem sieve element cells are first vascular cells to
differentiate at the transition zone (at the exact position where the in dataset were collected).
Because of this, phloem sieve element cells have denser cell walls than the adjacent phloem
companion cells, procambial or phloem pole pericycle cells (visible in Extended Figure 6,
cells with asterisks). Sieve element differentiation is accompanied by clearing of cellular
contents in a process called enucleation and phloem cell clearing (Furuta *et al.,* 2014, Blob
et al., 2018) (visible in Figure 2 G, cell type 5). The visualization of phloem cell clearing
process was originally reported using laborious high-resolution cryo-EM section assay that
relies on imaging 2D slices of electron microscopy samples (Furuta *et al.,* 2014,).
Remarkably, we were able to obtain information on phloem sieve element cell wall density
properties and visualize cellular clearing from 3D stacks of intact fresh samples (albeit not
at same ultra-high resolution as with 2D EM imaging). In addition to cell wall density
properties, very fine subcellular anatomical features were distinguishable within the 3D
datasets of soft plant tissues simply based on minor differences in their electron densities.
With the obtained resolution, it was possible to produce strikingly detailed images (like
Figure 2 H – L) and distinguish various subcellular organelles within imaged intact samples,
including mitochondria, vacuoles, and different nuclear compartments (i.e. nuclear
envelope, ribosomal protein enriched nucleolus, nucleoplasm, nuclear cavities, and
chromatin). To the authors' knowledge, similar high-quality absolute electron density 3D data
has not been reported from fresh multicellular plant samples, especially in such detail. **This
makes our PXCT and XNH results truly novel in two ways: 1) Subcellular features
previously only recorded with cryo-EM methods can now be determined in 3D from
intact fresh plant tissues. 2) Our datasets report actual sample electron densities
instead of mere anatomical features.**

Elemental information at subcellular organelle resolution

Our multi-modal 3D cryo-imaging approach yielded high-resolution electron density and elemental maps of intact fresh tissues at cell organelle resolution providing direct evidence of e.g. aggregate composition. Due to time constraints XRNF tomographic imaging was done using a larger pixel size of 200 nm and only a single or a few slices were imaged. To visualize and quantify distribution of Se and other micronutrients in root tissue and subcellular compartments, the higher resolution anatomical XNH electron density features were matched to individual chemical elements (mg/cm^3 values) obtained via XRNF measurements.

We observed differences in Se levels between samples and treatments, but overall Se accumulation profiles were similar between different tissues (epidermis, cortex, endodermis, and stele) (Graph 1). Se accumulation appeared to be enhanced in roots co-cultured with bacteria. Thus, based on our data, we could conclude that bacterial co-culturing does indeed promote Se accumulation to root tissues. According to previous reports plants accumulate Se in vacuoles (Gupta & Gupta, 2016). Using direct fluorescence measurements, we were able to quantify Se levels from various subcellular compartments (cytoplasm, cell walls and vacuoles). In both Se and Se+bacteria co-cultured roots highest values were measured from cytoplasm and cell walls, not from vacuoles. It should however be noted that due the cell wall masks used in segmentation analysis, some Se signal from cytoplasmic was likely included in the cell wall fraction, emphasizing subcellular cell wall measurements over other fractions. In addition, due to limited resolution, vacuoles were pooled as individual areas (including both membranes and vacuolic fluid). When analysis was done from individual hand selected points across the root, Se measurements looked slightly different: highest Se values came from cytoplasm, followed by aggregates in vacuolic membranes (lowest values were measured from vacuolic fluid). Nonetheless, we confirmed that Se does accumulate and aggregate in plants, however not exactly in similar manner as bacterial and mammalian cells.

In addition to differences in Se accumulation and subcellular localization, we observed changes in other micronutrient levels in response Se+bacteria supplementations, including high concentration Fe deposits in cell walls and Zn and Ca aggregates (also clearly visible in the videos of 3D XRNF volumes provided in online material). Interestingly, unlike initially expected, the observed high electron density aggregates seen in PXCT and XNH datasets did not match with XRNF Se maps but rather to Zn and Ca measurements. It seemed like the overall number and size of such dense aggregates was larger in bacteria co-cultured roots than in control or Se alone treated roots (Figure 2), indicating that bacteria affect root chemical composition in a complex manner. While bacteria have been shown to change the redox status of Se in the growing medium (Lusa *et al.,* 2017), possibly affecting uptake of Se, the methods we used in our research could not determine chemical speciation of the imaged elements. For such data chemical element speciation sensitive methods should be applied. Therefore, the questions about the possible influence of bacteria supplementation to the speciation of elements in the roots and the effect of chemical speciation to uptake of Se and Fe (and other elements) into tissues remains to be addressed.

Catching mechano-sensing "mid-flight"

One remarkable finding was that cryo-imaging allowed visualization of transient "moments frozen in time" preserved over extended imaging times. In the imaged samples, majority of Ca signal was seen evenly distributed at cell membranes/cell walls (Figure 4, Extended Figure 7). This is interesting, as Ca is known to act as a signalling component for mechanical stress, free oxygen radicals, and gravistimulus in plant cells, and Ca is locally released from cell walls and subcellular compartments rapidly upon such stimuli (Demidchik *et al.,* 2003, Monshause *at al.,* 2011, Kurusu *et al.,* 2013). Our measurements show high local Ca peaks in Se+bacteria supplemented roots in cells that were slightly damaged just prior to freezing (Extended Figure 7). In these cells, Ca levels were 2.5-fold higher than in the surrounding intact cell walls (Extended Figure 7, graph). While wounding and subsequent Ca burst was most likely induced by direct mechanical stimulus during sample preparation, we cannot exclude the possibility that the bacteria residing at the surface of the root might have contributed to cellular damage. Regardless of the cause, this example illustrates that non-invasive cryogenic imaging is especially well suited for preserving *in vivo* features of fresh tissues over extended imaging times (in our case up to 18h). In addition, it is worth noting that these same cells could be segmented from the voxel-wise Se vs. Fe concentration scatter plots as they form a clear outlier region distinct from other regions (Extended Figure 9). This indicates that utilization of multichannel data and advanced segmentation algorithms has potential to classify and reveal anatomical and physiological information which could be otherwise difficult to see. However, such considerations are beyond the scope of the current work.

Bioremediation and future of micronutrient accumulation assays

Using a multimodal cryogenic 3D and 2D imaging approach we were able to obtain comprehensive data of fresh *Arabidopsis thaliana* root samples in their native state with higher spatial resolution than what is currently reported in literature. Our observations on the chemical landscapes of individual cells and cellular organelles can promote understanding of metabolic processes involved in storing, processing and translocating Se and other micronutrients within the plant body. In addition, our data offers possible explanations to how soil bacteria affect root anatomy and growth. We suspect that the bacteria elicit phytohormone like growth regulators that lead to observed cellular division and tissue patterning defects. These abnormal cellular divisions then lead to super numerous root hair formation promoting micronutrient uptake (explaining the observed increase in Se and Fe accumulation). It is also possible that the bacteria promoted increase in micronutrient accumulation exceeds the toxicity limit for Se (and possibly Fe), leading to the observed growth arrest/tissue deformation and micronutrient hyperaccumulation. Regardless of the exact mechanisms which require more work until solved, the observed effects of bacterial supplementation open up an important promising possibility: *Pseudomonas* sp. strain T5-6-I soil bacteria could be utilized in both Se bio-remediation and Fe bio-fortification (likely even without Se supplementations), improving the nutritional values in agriculturally important crops grown on Fe poor areas**. Our findings demonstrate the great potential of soil-microbe co-culturing on bioremediation & biofortification efforts and illustrate how such interactions and their effects can be studied *in planta*.** This in turn will improve our

443 agricultural practices, especially in eroded areas where soil health, micro- and macronutrient
444 availability and plant uptake would benefit tremendously from the introduction of beneficial
445 soil microbes.
446
METHODS

MATERIAL

Surface sterilized (0.1 % Triton X in 95% ethanol) *Arabidopsis thaliana* wild type (Col-0
ecotype) seeds were sown on 1xMS-agar (Murashige and Skoog Basal Salt Mixture, Sigma
M5524) (Xi et al. 2016) plates (50 mL) containing 1% sucrose (w/v) and 0.7% (w/v) agar
(Sigma A1296).The seeds were kept at 4 °C in darkness for minimum of 2 days
(stratification) and then transferred to a growth chamber set at 23°C in 16/8 h light and dark
cycles (long day cycle, LD). Selenium and Bacterial suspensions of Pseudomonas sp. strain
T5-6-I (5·10^7 CFU of exponential growth phase bacteria) were added on one-week old
seedlings, by spreading in total 5 mL of the suspensions on the surface of the plant agar.
The plants were thereafter co-cultured with Se and bacteria for 7 days before imaging
(Extended figure 1). The plants were transported to synchrotron facilities in hand luggage
and kept under LD during sample preparation and imaging procedures. As we have reported
before (Lusa *et al.,* 2019) the treatments affect root phenotypes (Extended figure 1).
Therefore, the effect of each treatment was verified on site with light microscopy. An overall
visual inspection of plants used for PXCT and XNH verified that soil bacteria co-culturing
affects root anatomy and architecture (Extended Figure 2). While plants grown on normal
media showed normal primary and lateral root elongation rates and lateral root branching
pattern (Extended Figure 2 A, D), plants supplemented with Se showed slight reduction in
root growth rates (Extended Figure 2 B and E). However, the overall lateral root branching
pattern seemed unaltered in Se supplemented plants. In clear contrast, the plants
supplemented with Se+bacteria showed a reduction in root growth rate and abnormal lateral
root branching phenotype at a high frequency (not quantified) (Extended Figure 2 C, G and
H). Bacteria supplemented roots also exhibited a hairy root phenotype, as was reported with
kale (Lusa *et al.,* 2019) (Extended Figure 3, panels G - L). Root hair imaging was done at
the Light Microscopy unit at the Institute of Biotechnology, Helsinki, Finland. Imaged roots
were extracted from media and stained with propidium iodine (PI) before imaging with a
Leica SP5 inverted confocal microscope using a solid-state blue laser for GFP (480
nm/270mW).
476
In addition to increased root hair formation, bacteria co-cultured roots showed high rate of
lateral root meristem arrest (white root tips, Extended Figure 2 G) (not quantified). Such
arrested roots were not included in assays and imaging was only done with healthy looking
roots of similar age (Extended Figure 2 D - F) at around 2 d post emergence. Healthy looking
lateral root tips were mounted on cryogenic sample holders (OMNY-pins [Holler *et al.,* 2017]
and aluminium Huber-tubes at the Swiss Light Source (SLS) and at the European
Synchrotron Radiation Facility (ESRF), respectively, and snap frozen in liquid nitrogen.
484
485

IMAGING AND DATA RECONSTRUCTION

486
All synchrotron imaging assays (PXCT, XNH, and XRNF) were done at a position
corresponding to the transition zone (blue boxes in figure 1), which acts as a divider between
meristematic cells and elongating and differentiating cells. This zone is especially interesting
as cells of this zone act as sensors for various environmental stimuli, they are active in
cytoskeletal rearrangements and mediate growth responses leading to initiation of both root
hairs and organs (Baluška and Mancuso, 2013).
493
The cryo-PXCT measurements were performed at the cSAXS beamline at the SLS at the
Paul Scherrer Institut (PSI), Switzerland, using the OMNY instrument (Holler *et al.,* 2018).
Samples were glued to OMNY pins (Holler *et al.,* 2017) at room temperature and then frozen
in liquid nitrogen. The pins were transferred to the vacuum chamber and measured at a
sample stage temperature of 90 K. Cryogenic conditions can help maintain tissue and cell
shape, and reduce possible volatilization of Se. Samples were centered at about 100 μm
distance from the root tips (variation in root tilt affected distance). The heights of imaged
areas ranged between 15-25 μm, depending on the angle and/or width of the roots. A gold
Fresnel zone plate fabricated at the Laboratory for Micro- and Nanotechnology at the PSI
with a diameter of 250 μm and 50 nm outermost zone width was coherently illuminated at
6.2 keV photon energy. Spatial horizontal coherence was achieved by a horizontal upstream
slit and temporal coherence by a Si(111) double crystal monochromator. For the
measurement the samples were placed 1.6 mm after the focal spot of the Fresnel zone
plate, where the beam size had reached 6.5 μm diameter. The diffracted X-rays have been
recorded by an in-vacuum Eiger detector (Guizar-Sicairos *et al.,* 2014) placed at a distance
of 5.213 m after the sample.
510
Ptychographic projections were recorded scanning the sample in the beam following the
Fermat spiral trajectory (Huang *et al.,* 2014) with a step of 2 μm. The field of view and
number of projections was adjusted according to sample diameter and available
measurement time. Table 1 gives an overview of the different samples measured and the
parameters used. In order to keep imaging times reasonable (limiting to max 23-24 hours
per sample, corresponding to ~1200 projections), the resolution vs stack height was chosen
separately for each sample. The projections were reconstructed using between 1000 and
1600 iterations of the difference map algorithm (Thibault *et al.,* 2008) followed by 600
iterations of maximum likelihood refinement (Thibault & Guizar-Sicairos, 2012). For the
reconstructions, either 500 x 500 or 400 x 400 pixels of the detector were used, resulting in
a reconstructed pixel size of 27.8 or 34.7 nm, respectively. The projections were aligned
(Guizar-Sicairos *et al.,* 2011 and 2015) and combined to 3D volumes using a modified
filtered back projection (Guizar-Sicairos *et al.,* 2011) using a GPU-based matlab routine as
described in Ref. (Odstrcil *et al.,* manuscript in preparation). The 3D resolution was
estimated for each tomogram using Fourier shell correlation (FSC) between two
subtomograms with half the number of projections each and using the half-bit criterion (van
Heel & Schatz, 2005). An illustration of raw data and tissue identification from control root
samples is shown in Figure 1. The results of the FSC analysis are also given in table 1.
XNH and XRNF were performed under cryogenic conditions (LN2 temperature) under ultra-

high vacuum (Figure 1, B) at ESRF ID16A beamline. XNH volumes measured were centered
at 115μm distance from the root tips (full volumes shown in Figure 3). A brilliant beam (10^{11}
phot/s) of highly focused X-rays (30 (H) x 30 (V) nm^2) at 17 keV energy was produced by an
undulator source and a KB mirror optic (cf. Figure 1 B) (Morawe *et al.,* 2015). The focus was
used as a virtual point source for the XNH imaging by placing the sample some distance
downstream from the focus and recording a transmission image on the detector. These XNH
images show intricate interference patterns that carry information about the X-ray phase
shift happening in the sample. A tomographic scan was performed over 180-degree range
with 0.1-degree step size. At each rotation position the sample was randomly (but
controllably) displaced to reduce systematic artefacts from the incident beam and detector
non-uniformities. Such tomographic scan was performed at 4 different propagation
distances by varying the sample distance from the X-ray focus. A linear phase retrieval
algorithm (Paganin *et al.* 2002) extended to multiple propagation distances and followed by
iterative corrections was then applied to the set of 4 projection images at each rotation angle.
This produced 1800 quantitative phase projection images covering the 180-degree rotation,
which were subsequently used for tomographic reconstruction with PyHST software
package (Mirone *et al.,* 2014) to get a 3D volumetric image of the electron density in the
sample. The resulting voxel size of the images was 50 nm, which was chosen so that the
whole cross section of the sample would fit into the image, resulting in a total imaged volume
of about 100 ×100 ×100 μm^3.
550
The XRNF measurements were conducted by placing the sample in the X-ray focus, and
then raster scanning the sample to produce 2D maps and sinograms. The raster scanning
was performed with 200 nm spatial step size, 27 ms dwell time, and 0.72-degree step over
180 degrees rotation. Either 1 or 7 slices (spaced by 200 nm) were recorded in these scans
depending on the sample. The corresponding XRNF 2D slices and 3D stacks were located
at the center of the XNH field of view, at 115 μm distance from the root tips. The incident
energy of the X-rays was high enough to ionize all element species of interest in the sample.
A set of energy discriminating detectors placed at the sides of the sample were used to
record the fluorescence spectra coming from the sample. The summed spectra from these
detectors was then used to do peak fitting using the PyMCA software package (Solé *et al.,*
2007). The measurements were brought to quantitative units by using a previously
measured calibration sample. The end results of this peak fitting step were therefore images
of different elements in units of surface density (ng/mm^3). The fitted images were then
organized as sonograms, which were then reconstructed using the PyHST software
package with an iterative algorithm that minimizes the total variation in the object by using
Chambolle-Pock type method (Chambolle & Pock, 2010). This resulted in cross sectional
slice images of local concentration (gm/cm^3) for each of the analyzed elements. Due to the
high incident flux and small background signal, the XRNF images have a final detection limit
of about 1 ppm for each element.
570
DATA ANALYSIS, VISUALIZATION AND SEGMENTATION
For electron density analysis PXCT datasets were imported to and calibrated in Fiji using
experiment specific values for each dataset. These calibrated volumes were then used for

electron density measurements of various tissues and subcellular organelles from each
sample. In case of subcellular organelle measurements average values (min, max, median
and mean) were calculated from over 30 measuring individual areas around the sample and
presented as averages (table 2) and as bar charts including error bars (Extended Figure 4).
The sizes of measure areas ranged from tens of pixels (cell walls, vacuolic membranes) to
larger areas covering several thousand pixels per measured point (cytoplasmic and vacuolic
fluid), depending on the size of analyzed cellular organ.

XNH volumes and XRNF slices were imported, analyzed and processed in Fiji for electron
density visualizations and chemical element measurements. Chemical element
measurements done both at root, tissue and cell organelle level (for Fe and Se,) were done
in Fiji. Tissue averages and subcellular measurement averages are presented in Graphs 1
– 3 created using Microsoft Excel. Se levels were measured from each tissue type from in
all samples, from 20-30 areas per each cell type (different cell types are areas from which
measurements were taken are illustrated in Extended Figure 7). Subcellular Se
measurements were done from Se+bacteria sample with the most distinct anatomy for
reliable scoring of subcellular compartments. In this case, numerous small areas around the
root were selected (Extended Figure 8) and measured as pooled areas from epidermis,
cortex and endodermis. Resulting measurements for vacuole membranes, vacuolic fluid and
cytoplasm are presented in Graph 2. Fe levels were measured from 5+5 points around the
root per sample for cell walls and intracellular areas (Graph 3). All graphs and table 2 were
generated from measurements done in Fiji by exporting measured values to Microsoft Excel.

For visualizations individual images were exported in 16-bit tiff format and processed in
Corel PaintShop Pro X9 to create multilayered composite and panel images. Pseudo-
colored images of root cell types shown in Figure 1 were created from reconstructed PXCT
electron density image slices by hand using Corel PaintShop Pro X9. Each cell type was
colored onto a separate layer on top of the underlaying electron density image. Pseudo-
colored slice images (Figure 2, panels G and H, Extended Figure 3) consisting originally of
a single channel (grey) were produced with Fiji color palette function reflecting the
differences in intensity of the object. 3D visualizations of XNH datasets (of which slice
images are shown in Figure 3) were generated by exporting image stacks of calibrated
datasets from Fiji as tiff. and bmp. stacks and processed with CTvox from Bruker.

Concentrations of Se and Fe in various tissues and subcellular compartments in Graphs 1,
2 and 3, and online material Se_Fe statistics.xlsx were determined by tracing the anatomical
features from XNH slices using a combination of pixel and vector graphics programs (Corel
PaintShop Pro X9, Gimp2, Inkscape) and overlaying the traced image with the
corresponding XRNF slice to produce bitmap masks for voxel discrimination. The
distributions of concentrations in voxels over masked areas were calculated using custom
Python 3 scripts utilizing Numpy, Scipy, and Pandas packages, and the violin plots were
produced using Matplotlib and modified Seaborn packages.

For Se vs. Fe scatter plots, region of interest selections and the corresponding voxel visualizations (Extended Figure 9), the corresponding XNH and XRNF slices were aligned by forward projecting the reconstructed slices in x- and y-directions and finding the translational offset in x and y by cross-correlating the resulting 1D projections. The contribution of background signal was minimized by calculating masks over the reconstructed root by thresholding the data and calculating a convex hull around the remaining voxels. For scatter plots 2D histograms were evaluated from the voxelwise Se-Fe concentrations. To visualize the locations of voxels in the sample on the basis of scatter plots, regions of interest were defined as polygons in Se-Fe plots and XRNF voxels were masked on top of the anatomical XNH slice based on their inclusion in the polygons. All the computational and visualization were made using custom Python 3 scripts utilizing Numpy, Scipy, Matplotlib, Shapely, and Scikit-image packages. The analysis codes are available from the authors by request.

REFERENCES

1. Amselem J & Tepfer M. (1992). Molecular basis for novel root phenotypes induced by Agrobacterium rhizogenes A4 on cucumber. Plant Molecular Biology, 19 (3); 421–432.
2. Arrivault S, Senger T & Krämer U. (2006). The Arabidopsis metal tolerance protein AtMTP3 maintains metal homeostasis by mediating Zn exclusion from the shoot under Fe deficiency and Zn oversupply. The Plant Journal, doi.org/10.1111/j.1365-313X.2006.02746.x
3. Baluška F and Mancuso S. (2013). Root Apex Transition Zone As Oscillatory Zone. Frontiers in Plant Science. 2013; 4: 354. doi: 10.3389/fpls.2013.00354
4. Blob B, Heo J & Helariutta Y. (2018). Phloem differentiation: an integrative model for cell specification. Journal of Plant Research, 131(1): 31–36. doi: 10.1007/s10265-017-0999-0
5. Bodnar M, Konieczka P & Namiesnik J. (2017). The Properties, Functions, and Use of Selenium Compounds in Living Organisms. Journal of Environmental Science and Health Part C Environmental Carcinogenesis & Ecotoxicology Reviews. 2012;30(3):225-52. doi: 10.1080/10590501.2012.705164.
6. Boerjan W, Cervera M, Delarue M, Beeckman T, Dewitte W, Bellini C, Caboche M, Van Onckelen H, Van Montagu M, Inzé D. (1995). Superroot, a recessive mutation in Arabidopsis, confers auxin overproduction. The Plant Cell, 7: 1405-1419. doi.org/10.1105/tpc.7.9.1405
7. Cardarelli M, Mariotti D, Pomponi M, Spanò L, Capone I & Costantino P. (1987). Agrobacterium rhizogenes T-DNA genes capable of inducing hairy root phenotype Molecular Genetics and Genomics, 209(3):475-80.
8. Chen S, Deng J, Yuan Y, Flachenecker C, Mak R, Hornberger B, Jin Q, Shu D, Lai B, Maser J, Roehrig C, Paunesku T, Gleber S, Vine D, Finney L, VonOsinski J, Bolbat M, Spink I, Chen Z, Steele J, Trapp D, Irwin J, Feser M, Snyder E, Brister K, Jacobsen C, Woloschak G and Vogt S. (2014). The Bionanoprobe: hard X-ray fluorescence nanoprobe with cryogenic capabilities. JOURNAL OF SYNCHROTRON RADIATION ISSN: 1600-5775 21(19): 66-75. doi.org/10.1107/S1600577513029676

9. Cheng Y, Dai X & Zhao Y. (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. Genes Development. 2006 Jul 1; 20(13): 1790–1799.doi: 10.1101/gad.1415106

10. Chambolle A & Pock T. (2010). A first-order primal-dual algorithm for convex problems with applications to imaging. Journal of Mathematical Imaging and Vision. 40(1): 120–145

11. Cruz E, Becker S, Becker S & Sussulini A. (2018). Imaging of Selenium by Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA-ICP-MS) in 2-D Electrophoresis Gels and Biological Tissues. Methods in Molecular Biology. 1661:219-227. doi: 10.1007/978-1-4939-7258-6_16.

12. Cui J, Liu T, Li Y& Li F. (2018). Selenium reduces cadmium uptake into rice suspension cells by regulating the expression of lignin synthesis and cadmium-related genes. Science of the Total Environment, 644: 602–610.

13. da Silva J C, Pacureanu A, Yang Y, Bohic S, Morawe C, Barrett R, Cloetens P, Efficient concentration of high-energy x-rays for diffraction-limited imaging res-olution, Optica 4, 492-495, 2017.

14. Demidchik V, Shabala S, Coutts K, Tester M & Davies J. (2003). Free oxygen radicals regulate plasma membrane Ca^{2+}- and K^+-permeable channels in plant root cells. Journal of Cell Science, 1;116(1):81-8.

15. Desai M, Haigh M & Walkington H. (2018). Phytoremediation: Metal decontamination of soils after the sequential forestation of former opencast coal land. Science of the Total Environment, 15;656:670-680. doi: 10.1016/j.scitotenv.

16. Dierolf M, Menzel A, Thibault P, Schneider P, Kewish C, Wepf R, Bunk O & Pfeiffer F. (2010). Ptychographic X-ray computed tomography at the nanoscale. *Nature*, 467: 436–439.

17. Durán P, Viscardi S, Acuña J, Cornejo,Rosario Azcón P & de la Luz Mora M. (2018). Endophytic selenobacteria and arbuscular mycorrhizal fungus for Selenium biofortification and Gaeumannomyces graminis biocontrol. Journal of soil science and plant nutrition, 18(4) Temuco dic.

18. Esitken A. (1970). Use of Plant Growth Promoting Rhizobacteria in Horticultural Crops. In book: Bacteria in Agrobiology: Crop Ecosystems, Chapter 8. pages 189-235.

19. Furuta K, Yadav S, Lehesranta S, Belevich I, Miyashima S, Heo J, Vatén A, Lindgren O, De Rybel B, Van Isterdael G, Somervuo P, Lichtenberger R, Rocha R, Thitamadee S, Tähtiharju S, Auvinen P, Beeckman T, Jokitalo E & Helariutta Y. (2014). Arabidopsis NAC45/86 direct sieve element morphogenesis culminating in enucleation. Science, 345(6199): 933-937. DOI: 10.1126/science.1253736

20. Guizar-Sicairos M, Boon J, Mader K, Diaz A, Menzel A & Bunk O. (2015). Quantitative interior x-ray nanotomography by a hybrid imaging technique. Optica, 2(3): 259-266. https://doi.org/10.1364/OPTICA.2.000259

21. Guizar-Sicairos M, Diaz A, Holler M, Lucas M, Menzel A, Wepf R & Bunk O. (2011). Phase tomography from x-ray coherent diffractive imaging projections. Optics Express, 1(22): 21345-21357. https://doi.org/10.1364/OE.19.021345

22. Guizar-Sicairos M, Johnson I, Diaz A, Holler M, Karvinen P, Stadler H-C, Dinapoli R, Bunk O& Menzel A. (2014). High-throughput ptychography using Eiger: scanning X-ray nano-imaging of extended regions. Optics Express, 22(12): 14859-14870. https://doi.org/10.1364/OE.22.014859

23. Gupta M & Gupta S. (2016). An overview of Selenium Uptake, Metabolism and Toxicity in plants, Frontiers in Plant Science, 2016(7):2074. doi: 10.3389/fpls.2016.02074.

24. Helin J, Ikonen A & Hjerpe T. (2010) Review of element specific data for biosphere assessment BSA-2009. Working Report 2010-37, Posiva Oy.

25. Holler M, Raabe J, Wepf R, Shahmoradian SH, Diaz A, Sarafimov B, Lachat T, Walther H & Vitins M (2017). OMNY PIN — A versatile sample holder for tomographic measurements at room and cryogenic temperatures. Review of Scientific Instruments, 88(11):10.1063/1.4996092

26. Holler M, Raabe J, Diaz A, Guizar-Sicairos M, Wepf R, Odstrcil M, Shai F, Panneels V, Menzel A, Sarafimov B, Maag S, Wang X, Thominet V, Walther H, Lachat T, Vitins M & Bunk O. (2018). OMNY—A tOMography Nano crYo stage. Review of Scientific Instruments, 89, 043706. https://doi.org/10.1063/1.5020247

27. Hu P, Wang Y, Przybyłowicz W, Li Z, Barnabas A, Wu L, Luo Y & Mesjasz-Przybyłowicz J. (2015). Elemental distribution by cryo-micro-PIXE in the zinc and cadmium hyperaccumulator Sedum plumbizincicola grown naturally. Plant and Soil. 388(1–2) 267–282.

28. Huang X, Yan H, Harder R, Hwu Y, Robinson I & Chu Y. (2014). Optimization of overlap uniformness for ptychography. Optics Express, 22(10): 12634-12644. https://doi.org/10.1364/OE.22.012634

29. Jiang Y, El Mehdawi A, Tripti , Lima L, Stonehouse G, Fakra S, Hu Y, Qi H & Pilon-Smits E. (2018). Characterization of Selenium Accumulation, Localization and Speciation in Buckwheat–Implications for Biofortification. Frontiers in Plant Science, 2018; 9: 1583. doi: 10.3389/fpls.2018.01583

30. Khan S, Zada S, Ahmad S, Lv J & Fu P. (2019). Concurrent biomineralization of silver ions into Ag0 and AgxO by Leptolyngbya strain JSC-1 and the establishment of its axenic culture. Chemosphere, 215, Pages 693-702.

31. Krajcarová L, Novotný K, Kummerová M, Dubová J, Gloser V & Kaiser J. (2017). Mapping of the spatial distribution of silver nanoparticles in root tissues of Vicia faba by laser-induced breakdown spectroscopy (LIBS) Talanta .173: 28-35. doi.org/10.1016/j.talanta.

32. Kurusu T, Kuchitsu K, Nakano M, Nakayama Y & Iida H. (2013). Plant mechanosensing and Ca2+ transport. Trends in Plant Science, 18, ISSUE 4, P227-233, April 01, 2013. DOI:https://doi.org/10.1016/j.tplants.2012.12.002

33. Li H, McGrath S & Zhao F (2008). Selenium uptake, translocation and speciation in wheat supplied with selenate or selenite. New Phytology, 178: 92-102, 10.1111/j.1469-8137.2007.02343.x

34. Lincoln C, Britton J & Estelle M. (1990). Growth and Development of the axr1 Mutants of Arabidopsis. The Plant Cell, Vol. 2, 1071-1 080

35. Lintern M, Anand R, Ryan C & Paterson D. (2013). Natural gold particles in Eucalyptus leaves and their relevance to exploration for buried gold deposits. Nature Communications, 4: 2614

36. Liu S, Hu Q, Luo S, Li Q, Yang X, Wang X & Wang S. (2015) Expression of wild-type PtrIAA14.1, a poplar Aux/IAA gene causes morphological changes in Arabidopsis. Frontiers in Plant Science,02 June 2015. doi.org/10.3389/fpls.2015.00388

37. Longchamp M, Angeli N & Castrec-Rouelle M. (2016). Effects on the accumulation of calcium, magnesium, iron, manganese, copper and zinc of adding the two inorganic forms of selenium to solution cultures of Zea mays. Plant Physiology and Biochemistry. 98:128-37. doi: 10.1016/j.plaphy.2015.11.013.

38. Lusa M, Bomberg M, Aromaa H, Knuutinen & Lehto J. (2015). The microbial impact on the sorption behaviour of selenite in an acidic, nutrient-poor boreal bog. Journal of Environmental Radioactivity, 147: 85-96.

39. Lusa M, Help H, Honkanen AP, Parkkonen J, Kalasova D & Blomberg M. (2019). Bacterial effects on selenium uptake in plants. Environmental Research. Available online 13 August 2019, 108642. https://doi.org/10.1016/j.envres.2019.108642

40. Lusa M, Knuutinen J & Blomberg M. (2017). Uptake and reduction of Se(IV) in two heterotrophic aerobic Pseudomonads strains isolated from boreal bog environment. AIMS Microbiology, 4: 798-814.

41. Lusa M, Lehto J, Aromaa H, Knuutinen J & Bomberg M. (2016) The Uptake of Radioiodide by Paenibacillus sp., Pseudomonas sp., Burkholderia sp. and Rhodococcus sp. isolated from a Boreal Nutrient-poor Bog. Journal of Environmental Sciences. 44, 26-37.

42. Mirone A, Brun E, Gouillart E, Tafforeau P & Kieffer J. (2014) .The PyHST2 hybrid distributed code for high speed tomographic reconstruction with iterative reconstruction and a priori knowledge capabilities. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms, 324: 41-48.

43. Monshausen G, Miller N, Murphy S & Gilroy S. (2011). Dynamics of auxin-dependent Ca2+ and pH signaling in root growth revealed by integrating high-resolution imaging with automated computer vision-based analysis. Plant Journal. 65(2):309-18. doi: 10.1111/j.1365-313X.2010.04423.x.

44. Morawe Ch, Barrett R, Cloetens P, Lantelme B; Peffen J-Ch, Vivo A. (2015) Graded multilayers for figured Kirkpatrick-Baez mirrors on the new ESRF end station ID16A. Proc. SPIE, 9588, 958803.

45. Odstrcil M. et al, *manuscript in preparation*

46. Ojuederie O & Babalola O. (2017). Microbial and Plant-Assisted Bioremediation of Heavy Metal Polluted Environments. International Journal of Environmental Research and Public Health, 4;14(12). pii: E1504. doi: 10.3390/ijerph14121504.e 4

47. Ondrasek G, Rengel Z, Clode P, Kilburn M, Guagliardo P & Romic D. (2019). Zinc and cadmium mapping by NanoSIMS within the root apex after short-term exposure to metal contamination. Ecotoxicology and Environmental Safety. 171: 571-578. doi.org/10.1016/j.ecoenv.

48. Paganin D, Mayo S C, Gureyev T E, Miller P R, Wilkins S W, Simultaneous phase and amplitude extraction from a single defocused image of a homogeneous object, J. Microsc. 206, 33 (2002)

49. Perrot-Rechenmann C. (2010). Cellular Responses to Auxin: Division versus Expansion. Cold Spring Harbor Perspectives in Biology, 2(5): a001446. doi: 10.1101/cshperspect.a001446

50. Pirozzi N, Hoogenboom J & Giepmans B (2018). ColorEM: analytical electron microscopy for element-guided identification and imaging of the building blocks of life. Histochemistry and Cell Biology. 150,(5): 509–520.

51. Reynolds R & Pilon-Smits E. (2018). Plant selenium hyperaccumulation- Ecological effects and potential implications for selenium cycling and community structure. Biochimica et Biophysica Acta (BBA) - General Subjects. 1862(11): 2372-2382.

52. Rodrigues E, Gomes M, Duran N, Cassanji J, da Cruz T, Sant'Anna Neto A, Savassa S, de Almeida E & Carvalho H. (2018). Laboratory Microprobe X-Ray Fluorescence in Plant Science: Emerging Applications and Case Studies. Frontiers in Plant Science, 9: 1588. doi: 10.3389/fpls.2018.01588

53. Shahmoradian S, Tsai E, Diaz A, Guizar-Sicairos M, Raabe J, Spycher L, Britschgi M, Ruf A, Stahlberg H & Holler M. (2017). Three-Dimensional Imaging of Biological Tissue by Cryo X-Ray Ptychography. Scientific Reports, 24; 7(1):6291.

54. Singh P, Mohanta T & Sinha A. (2015). Unraveling the Intricate Nexus of Molecular Mechanisms Governing Rice Root Development: OsMPK3/6 and Auxin-Cytokinin Interplay. PLOS ONE, DOI:10.1371/journal.pone.0123620

55. Solé V, Papillon E, Cotte M, Walter Ph & Susini J. (2007). A multiplatform code for the analysis of energy-dispersive X-ray fluorescence spectra. Spectrochimica Acta Part B: Atomic Spectroscopy, 62(1): 63-68.

56. Staicu L, van Hullebusch E & Lens P. (2017). Wastewaters and Physical–Chemical Treatment Technologies. Bioremediation of Selenium Contaminated Wastewater pp 103-130 | Industrial Selenium Pollution:

57. Tangahu B, Abdullah S, Basri H, Idris M, Anuar N & Mukhlisin M. (2011). A Review on Heavy Metals (As, Pb, and Hg) Uptake by Plants through Phytoremediation. International Journal of Chemical Engineering. 2011; 939161. doi.org/10.1155/2011/939161

58. Thibault P, Dierolf M, Menzel A, Bunk O, David C & Pfeiffer F. (2008). High-Resolution Scanning X-ray Diffraction Microscopy. Science, 321(5887): 379-382. DOI: 10.1126/science.1158573

59. Thibault P & Guizar-Sicairos M. (2012). Maximum-likelihood refinement for coherent diffractive imaging. New Journal of Physics, Volume 14.

60. Tran T, Zhou F, Yang W, Wang M, Dinha Q, Wang D & Liang D. (2018). Detoxification of mercury in soil by selenite and related mechanisms. Ecotoxicology and Environmental Safety, 159, Pages 77-84

61. van Heel M & Schatz M. (2005). Fourier shell correlation threshold criteria. Journal of Structural Biology, 151 (3): 250-262. doi.org/10.1016/j.jsb.2005.05.009

62. Wang P, Menzies N , Lombi E, McKenna B, de Jonge M, Donner E, Blamey F, Ryan C, Paterson D, Howard D, James S & Kopittke P. (2013). Quantitative determination of metal and metalloid spatial distribution in hydrated and fresh roots of cowpea using

synchrotron-based X-ray fluorescence microscopy. Science of The Total Environment.
463–464: 131-139.
63. Weinhardt V, Chen JH, Ekman A, McDermott G, Le Gros MA and Larabell C. (2019).
Imaging cell morphology and physiology using X-rays, Biochemical Society Transactions.
BST20180036; DOI: 10.1042/BST20180036
64. Xi W, Gong X, Yang Q, Yu H & Liou YC. (2016). Pin1At regulates PIN1 polar localization
and root gravitropism. Nature Communications. 2016/01/21/online, DOI:
10.1038/ncomms10430
65. Ximénez-Embún P, Alonso I, Madrid-Albarrán Y, Cámara C. (2004). Establishment of
Selenium Uptake and Species Distribution in Lupine, Indian Mustard, and Sunflower
Plants. Journal of Agricultural and Food Chemistry, 52: 832−838.
66. Yu Y, Fu P, Huang Q, Zhang J & Li H. (2019). Accumulation, subcellular distribution, and
oxidative stress of cadmium in Brassica chinensis supplied with selenite and selenate at
different growth stages. Chemosphere, 216: 331-340
67. Zhao FJ, Moore K, Lombi E, Zhu YG. (2014). Imaging element distribution and speciation
in plant cells. Trends in Plant Science. 19(3): 183-192
68. Zhao Y, Christensen S, Fankhauser C, Cashman J, Cohen J, Weigel D & Chory J. (2001).
A Role for Flavin Monooxygenase–Like Enzymes in Auxin Biosynthesis. Science, 291;
306-309.
850

FIGURE LEGENDS

Figure 1: Plant material and visualization of the ID16A imaging protocol

A) Arabidopsis seedlings were grown in 3 different conditions: Normal 1% suc 1xMS media
(left, top), 1 w on normal media + 1 w on Se supplemented media (left, middle) and 1 w on
normal media + 1 w on Se supplemented media with bacterial co-culturing (left, bottom).
Roots of same age were collected few days after emergence (middle panels, note the
growing conditions affected root elongation rates) from ~2w old plants in vegetative growing
phase showing few rosette leaves (illustration on the right). Collected roots were mounted
in cryogenic imaging holders. The imaged areas are shown in small white boxes in middle
panes (middle-top, control root; middle-middle Se supplemented; middle-bottom,
Se+bacteria cultured root). B) illustration of the ID16A imaging protocol: 2D XNH
holotomographic projection images taken at different distance from X-ray focus were used
to retrieve phase map projection images. These 2D phase images were then reconstructed
to 3D volumes. XRNF measurements were conducted by placing the sample in the X-ray
focus, and then raster scanning the sample pixel by pixel to produce 2D maps and
sinograms. The XRNF 2D slices and 3D stacks (calculated from sinograms) were located at
the center of the XNH field of view, at 115 μm distance from the root tips.
868

Figure 2: Anatomical comparisons and details of PXCT and XNH imaged roots

Electron density cross section images from PXCT (A – C) and XNH (D - F) reconstructions.
A) control root, B) Se supplemented root, and C) Se+bacteria supplemented root. D) control
root, E) Se supplemented root, and F) Se+bacteria supplemented root. Overall the imaged
roots appear to be anatomically very similar within each treatment (A and D, control roots)
(B and E, Se supplemented roots) (C and F, Se + bacteria supplemented roots). Identities

of different cell layers are indicated with colored dots in the greyscale electron density images; epidermis (brown), cortex (yellow), endodermis (red) and pericycle (purple). An electron density signal intensity-based color image illustrates different cell types in a longitudinal cross section of control root: 1) epidermis, 2) cortex, 3) endodermis, 4) pericycle, 5) phloem sieve element (cleared from cytoplasmic organelles), x) procambium/ metaphloem cells. H - L) Close up images of heat-map colored control root (H), Se supplemented root (I) and Se+bacteria supplemented root (J, K and L). These close ups of epidermal and cortical cells nicely illustrate that cryogenic 3D PXCT imaging allows inspection of density features in very fine detail, even at the level of subcellular organelles. Marked in the image are various subcellular and nuclear compartments: nuclear envelopes (white arrows) and nucleoli (orange arrows). Nuclear cavities and chromatin are within the nuclear envelope. The nucleus is surrounded by mitochondria (dark blue arrows), which appear denser that the surrounding cytoplasmic fuid and vacuolic "fluid" within the expanding vacuolic compartments (lilac arrows). Red arrows point to dense aggregates. Yellow circle in panel K illustrates high density cytoplasmic membranes (ER), and red circle illustrates dense vacuolic membranes, All scale bars 10µm.

Figure 3: Meristem anatomical features and root hair densities
Meristem anatomy of control (A and D), Se (B and E) and Se + bacteria treated roots (C and F) differ from each other. Panels A, B and C represent longitudinal cross sections showing the vasculature, whereas panels D, E and F show ground tissue layers. Arabidopsis control root hairs in panel G, Se supplemented root in H, and Se+bacteria supplemented root in I were imaged using a confocal laser microscope. The root hair phenotypes are likely a combination resulting from selenium toxicity (with additive effects of other micronutrient accumulation,) and bacterial elicitor sensing and uptake (that mimic plant phytohormone, e.g. auxin signaling). Scale bar in panel A is 10µm and 50µm in panels G - I.

Figure 4: XNH electron density slice images and specific chemical element maps
This figure illustrates XNH slices of differently treated samples [1 (control), 2 & 3 (Se supplemented) and 4 & 5 (Se+bacteria supplemented)] and 2D chemical element maps of selected elements, Se, Fe, Cr and Ca at the exact position of XNH scans. Top row marked with XNH present physical electron density maps (XNH slices). Second row illustates XRNF signal of Se in samples. Third row shows signal of Fe, fourth row Cr and fourth row Ca signal in control roots. Composite overlay images of XRNF maps of Se, Fe, Cr and Ca are shown in the sixth panel. Composite images showning the physico-chemical maps are shown in bottom panel to illustrate the overall chemical composition differences between samples.

Figure 5: Closeups of Se and Fe in different treatments
First column on the left presents XNH imaged roots. Second column close ups of selenium and third column iron in root sections. In the fourth column Se and Fe signals are overlaid and presented in blue and red, respectively. The column in the right shows Se and Fe signals overlaid with XNH images for tissue visualization. Rows A – E represent imaged samples, control (A), Se supplemented (B and C) and Se+bacteria supplemented roots (D and E).

Graph 1: Selenium and iron contents in various root tissues of different treatments

Top: Tissues from which Se and Fe levels were measured: epidermis (red), cortex (green), endodermis (light blue), pericycle (yellow), stele (dark blue) and background (black). Pink areas represent damaged cells that were excluded from analysis. Middle: The solid horizontal lines within the violin plots of various tissues and adjacent cell walls represent medians, and the dashed lines mark 25 and 75 percentile tails. Peak values are indicated with numbers. The bottom and top values represent minimum and maximun signal intensities, respectively (anything below 0 should be considered a measurement artefact). Bottom: The Fe levels of control and Se supplemented roots are in similar range, with slightly higher values in control root. The Se+bacteria supplemented roots show considerably higher levels of Fe (notice the different scale on y-axes).

Graph 2: Selenium and iron contents measured from cells and walls from various tissues from different treatments

Se and Fe levels were measured from within all ground tissue layers, and from different cell wall positions (e.g. cell walls connecting only epidermis cells, and separately from cell walls connecting epidermis and cortex cells etc.). The masks representing measured areas are presented on the top: superficial cell walls (burgundy), epidermal cell contents (dark blue), epidermal cell walls (red), epidermocortical cell walls (brown), cortical cell contents (dark purple), cortical cell walls (dark green), corticoendodermal cell walls (bright green), endodermal cell contents (fuchsia), endodermal cell walls (turquoise) and endodermo-pericydal cell walls (light blue). Yellow areas are damaged cells, and white cell walls that were both excluded from measurements. Black areas, background. The solid horizontal lines within the violin plots of various tissues and adjacent cell walls represent medians, and the dashed lines mark 25 and 75 percentile tails. The bottom and top values represent minimum and maximum signal intensities, respectively (anything below 0 should be considered a measurement artefact). Medians are shown in solid lines and 25 and 75 percentile tails are separated with dashed lines. Peak values are indicated with numbers. Accumulation of Se appears quite even between intracellular and cell wall areas in Se and Se+bacteria supplemented roots. Opposite to Se, Fe appears to accumulate predominantly into cell walls, and in all samples, the superficial cell walls facing the growing medium showed highest values for all treatments. Fe accumulation was extremely pronounced in Se+bacteria supplemented root samples, where the median (solid lines) and highest values (peaks) were ~10X higher than in control root (notice the differentially scaled y-axes).

Graph 3: Selenium and iron contents in different subcellular organelles

Se and Fe levels were quantified at cell organelle level from all XRNF imaged samples. Masks used in quantification are shown on top row. Not all cells would be included into analysis due to suboptimal tissue resolution. Cell walls are shown in red, cytoplasmic areas in yellow and vacuolic compartments in blue. The solid horisontal lines within the violin plots of various tissues and adjacent cell walls represent medians, and the dashed lines mark 25 and 75 percentile tails. The bottom and top values represent minimum and maximun signal intensities, respectively (anything below 0 should be considered a measurement artefact). Based on these segmentation analyses, cell walls and cytoplasm contained on average

more Se than vacuolic compartments. Notice the different scale on y-axes.

Table 1: Overview of the cryo-PXCT measurement parameters
Overview of the cryo-PXCT measurement parameters. (*) The FSC estimation for id134 is unrealistically low due to the insufficient amount of projections in each half-tomogram, given the large diameter of this sample.

EXTENDED MATERIAL

Extended Figure 1: Electron densities of cryo- PXCT imaged roots
These 3 graphs represent the electron density measurements from control, Se supplemented and Se+bacteria supplemented roots imaged using cryo-PXCT. For each average value shown in the attached graphs, >30 points across the samples were measured and averaged. Minimum (light blue bars), maximum (orange bars), median (light yellow bars) and mean (dark yellow bars) represent the electron densities of cell walls, cytoplasm and organelles, vacuole fluid and membranes. Error bars are included.

Extended Figure 2: XRNF data for Se. Fe. Cr. Br and Ca
While most chemical elements showed little difference in signal levels between different treatments, Se, Fe, Cr and Br showed different accumulation levels in roots, of which Se and Fe were most noticeable. Ca is presented as an element that did not change between treatments. Zn measurements with clear aggregates are shown in the bottom panel.

Extended Figure 3: Example of Se measurements at organelle level in Se+bacteria supplanted root
Se levels measured at organelle level, here shown in Se+bacteria supplemented root #1, of which individual cell types and organelles were most distinguishable. Small yellow boxes in the top panels represent regions from which Se was measured in cortex cells. Left, cytoplasm and ER, brightest signal corresponds to highest Se levels, darkest signal to low Se levels. Middle panel shows vacuole membranes from which signals were measured, and right panel cell vacuolic fluid areas. Highest Se signal levels were measured in cytoplasm (and ER) in all analyzed cell types; epidermis (brown color in insert picture), cortex (yellow) and endodermis (pink).The bars on the graph do not show error bars as they represent averaged values from multiple measurement points (Extended Figure 8). In all measured tissues, Se levels are highest cytoplasm (including ER).

Extended Figure 4: Damaged root epidermal cells show increased Ca signal
The XNH 3D reconstruction of a Se+bacteria supplemented root sample #2 on the left shows rod like Pseudomonas bacteria visible on the root surface. The cells marked with red arrow are also visibly damaged, and correspondingly show higher Ca signal in fluorescent map (top panel on the right). The Ca signal on damaged areas was ~2,5-fold stronger than in intact cell walls, and is shown in table. 17 individual points/areas were measured for both damaged and intact root epidermis values (n=17 + 17).

Extended Figure 5: Root density plots for Se and Fe visualization of plotted areas
Voxel-wise concentrations of Fe vs. Se in XRNF reconstructions of the measured samples. Scatter plots A-E correspond to the control, Se 1, Se 2, Se+bac 1, and Se+bac2, respectively. Panel F presents the same data in a single figure together with 5 regions of interest (labelled 1-5). The XRNF voxels falling inside the marked rois are presented for each sample over the XNH data in the bottom part of the figure. Notably the outlier region 5 corresponds to the damaged cells of epidermis in Se+bacteria supplemented samples.

Extended Figure 6: A single slice electron density image of a Se+bacteria treated root imaged using 3D cryo-PXCT
Pseudo-color image of a Se+bacteria supplemented root imaged using PXCT. Vacuolic cell organelles can be distinguished by their yellow color, and their density is smaller than cytoplasm (orange), endoplasmatic reticulum (red), cell walls (red, pink and white) and vacuolic membranes (red membranes inside yellow areas). As was shown with fluorescence measurements Fe localizes primarily to root cell walls, especially in cell corners. These high density areas show white signal. Fluorescence data also showed that Se accumulates primarily inside the cells; in cytoplasm and ER (orange and red areas) and in vacuole membranes shown here in orange and red. The white aggregates marked with black hollow arrows are likely Zn or Ca aggregates. Orange organelles in the cytoplasm (marked with narrow arrows) are mitochondria and/or other small cell organelles. Most cells in different tissues exhibit similar anatomy and density features. What is noticeable, is that the resolution of cryogenic PXCT imaging at 80nm resolution allows detection and distinction of individual cellular organelles: mitochondria, plastids, nuclei, cytoplasmic fluid, ER, vacuole membranes. The cells at the center of the root are vascular cells, and the ones with thicker cell walls (marked with asterisks) are phloem sieve elements.

Extended Figure 7: Different root cell identities can be distiguished from PXCT reconstructions
Panels on the left show electron density maps of a PXCT imaged control root, cross section (top panel) and longitudinal section (bottom panel). Panels on the right side are artificially colored to illustrate different cell layers normally present in roots (cross section at the top, longitudinal section in the middle, color legends at the bottom). Outermost cell layer is epidermis (brown), below it is cortex (yellow), endodermis (rosa), pericycle (purple), proto xylem (blue), metaxylem (light blue), procambium (light green), phloem companion cells (pink) and phloem sieve elements (red). All scale bars are 10 µm.

Extended Figure 8: Overall root architectures of different treatments, growth arrested roots and abnormal lateral root anatomy of bacteria supplemented samples.
This figure show overall architecture of imaged roots. Roots grown on normal media for 2 weeks (A), look similar to roots grown on normal media for 1 w and supplemented with Se for 1 weeks (B) whereas roots grown with Se+bacteria for 1 w (C) grow differently than agar controls; bacteria supplemented roots show defects in primary root elongation and branching. D is a close up of a control lateral root, E is a Se supplemented root and F , G and H show Se+bacteria supplemented lateral roots of similar age as D and E. Root in panel

1051 F exhibits stunted growth and swelling, and the roots in panel G and H show arrest in growth
1052 (dying meristem is white and opaque) and an extraordinary branching phenotype. In control
1053 samples, roots never emerge side by side. Scale bar in H is 100µm.

1054

1055 **Table 2: Electron densities of cell compartments**
1056 Presented values of subcellular organelle density minimum, maximum, median and mean
1057 from different treatments are averages, each measured from n>30 areas around the sample.
1058 The size of the measured areas depended on the cell organelle size.

1059

ACKNOWLEDGEMENTS

Hanna Help and Henrik Mäkinen were supported by the Academy of Finland (Grant
1295696). Merja Lusa supported by the project "Biosorption and bioaccumulation of heavy
metals and radionuclides - use of environmental bacteria in the bioremediation of mining
waste- waters" from KAUTE foundation. Ari-Pekka Honkanen was funded by University of
Helsinki Doctoral Program in Materials Research and Nanosciences (MATRENA).

1066

AUTHOR CONTRIBUTIONS

H.H. designed the experiments, participated in material preparations, conducted all sample
preparations, participated in measurements, performed data analysis, and wrote the
manuscript. **M.L.** participated in experimental design, prepared the plant material,
participated in measurements and manuscript preparation. **A.-P.H.** participated in
measurements, cryo-PXCT data reconstruction, analysis of cryo-PXCT, -XNH and -XRNF
data, and manuscript preparation. **A.D.** and **M.H.** performed the cryo-PXCT measurements
and data reconstructions and participated in manuscript preparation. **H.M.** participated in
cryo-PXCT measurements and data reconstructions. **M.S.** and **P.C.** performed the cryo-XNH
and -XRNF measurements and data reconstructions. **S.H.** participated in experimental
design and manuscript preparation as well as in overall project supervision. **H.S.** participated
in experimental design, cryo-XNH and -XRNF measurements and data reconstructions, and
manuscript preparation.

1080

1081 ## COMPETING INTERESTS

1082 The authors declare that there is no conflict of interest regarding the publication of this article.
1083 The also authors declare that they have no known competing financial interests or personal
1084 relationships that could have appeared to influence the work reported in this paper.

1085

1086 ## MATERIALS & CORRESPONDENCE

1087 All correspondence regarding the manuscript and scientific content should be addressed to
1088 Dr Hanna Help. Requests for materials should be addressed to Dr Merja Lusa. Requests for
1089 segmentation code should be addressed to Ari-Pekka Honkanen. All data requests should
1090 be addressed to Dr Heikki Suhonen.

1091

1092

FIGURES

Figure 1: Plant material and visualization of the ID16A imaging protocol

1118 **Figure 2: Anatomical comparisons and details of PXCT and XNH imaged roots**

Figure 3: Meristem anatomical features and root hair densities

Figure 4: XNH electron density slice images and spesific chemical element mps

Figure 5: Closeups of Se and Fe in different treatments

Graph 1: Selenium and iron contents in various root tissues of different treatments

Graph 2: Selenium and iron contents measured from cells and walls of various tissues from different treatments

1241 **Graph 3: Selenium and iron contents in different subcellular organelles**

1242

1243

1244

1245

1246

1247

1248

1249

1250 **Table 1: Electron densities of cell compartments**

Sample ID	e- density			
	min	max	median	mean
control cell wall	0.36	0.42	**0.39**	**0.39**
control cytoplasm and organelles	0.28	0.37	0.32	0.32
control vacuole fluid	0.26	0.34	0.30	0.30
control vacuole membranes	0.33	0.42	0.38	0.38
Se cell wall	0.31	0.43	0.38	0.38
Se cytoplasm and organelles	0.26	0.38	0.32	0.32
Se vacuole fluid	0.25	0.35	0.30	0.30
Se vacuole membranes	0.28	0.40	**0.34**	**0.34**
Se+bact cell wall	0.33	0.47	**0.42**	**0.42**
Se+bact cytoplasm and organelles	0.28	0.40	0.33	0.33
Se+bact vacuole fluid	0.27	0.34	0.31	0.31
Se+bact vacuole membranes	0.32	0.45	**0.40**	**0.39**

1251 Presented values of subcellular organelle density minimum. maximum. median and mean from
1252 different treatments are averages. Each measured from n>30 areas around the sample. The size of
1253 the measured areas depended on the cell organelle size.

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264 **EXTENDED MATERIAL**

1265 **Extended Figure 1: Electron densities of cryo- PXCT imaged roots**

1277

1278

1279

1280

1281

1282

Extended Figure 2: XRNF data for Se. Fe. Cr. Br and Ca

Extended Figure 3: Example of Se measurements at organelle level in Se+bacteria supplemented root

Extended Figure 4: Damaged root epidermal cells show increased Ca signal

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

Extended Figure 5 – Root density plots for Se and Fe visualization of plotted areas

1337 **Extended Figure 6: A single slice electron density image of a Se+bacteria treated root**
1338 **imaged using 3D cryo-PXCT**

1339
1340
1341

Extended Figure 7: Different root cell identities can be distiguished from PXCT reconstructions

.

Extended Figure 8: Overall root architectures of different treatments. growth arrested roots and abnormal lateral root anatomy of bacteria supplemented samples.

1391

1393 **Table 2: Overview of the cryo-PXCT measurement parameters**

Sample ID	id131	id132	id133	id134	id135	id136
Name	control_s1	bakt_sel_1	sel_7	bakt_sel_2	control_2	sel_9
Field of view [μm^2]	100 x 20	85 x 22	85 x 22	110 x 12	70 x 24	130 x 20
Voxel size (nm)	27.8	34.7	34.7	34.7	34.7	34.7
# Projections	1200	1200	1200	600	600	950 (1000)
Dose [MGy]	2.5	15	15	6.5	7	
Measurement time (projection)	60 s	54 s	54 s	41 s	50 s	72 s
Measurement time (total)	22 h	21 h	21 h	10 h	10 h	
Sample diameter/ number of projections	70 nm	60 nm	70 nm	160 nm	90 nm	120 nm
FSC 3D	90 nm	80 nm	90 nm	NA*	124 nm	130 nm
Figures	2A, 2B, S1, S10	2C, S4		2D		

1394 Overview of the cryo-PXCT measurement parameters. (*) The FSC estimation for id134 is
1395 unrealistically low due to the insufficient amount of projections in each half-tomogram, given the large
1396 diameter of this sample.