TRAF6 controls excitatory spinogenesis and excitation-inhibition balance through binding neuroplastin

Running tittle: TRAF6 mediates cell-autonomously synapse formation

Sampath Kumar Vemula[1,*], Ayse Malci[1,*], Lennart Junge[1], Anne-Christin Lehmann[1], Ramya Rama[1], Johannes Hradsky[1], Ricardo A. Matute[2,3], Michael Naumann[4], Michael R. Kreutz[1,5], Constanze I. Seidenbecher[1,6], Eckart D. Gundelfinger[1,6], Rodrigo Herrera-Molina[1,3,§]

[1]Leibniz Institute for Neurobiology, Magdeburg, Germany

[2]Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California, United States

[3]Centro Integrativo de Biología y Química Aplicada, Universidad Bernardo OHiggins, Santiago, Chile

[4]Institute of Experimental Internal Medicine, Medical Faculty, Otto von Guericke University, Magdeburg, Germany

[5]Leibniz Group 'Dendritic Organelles and Synaptic Function', Center for Molecular Neurobiology (ZMNH), University Medical Center Hamburg-Eppendorf Hamburg, Germany.

[6]Center for Behavioral Brain Sciences and Medical Faculty, Otto von Guericke University, Magdeburg, Germany

* These authors contributed equally to this work

§ Address correspondence to: Rodrigo Herrera-Molina rherrera@lin-magdeburg.de, Laboratory of Neuronal and Synaptic Signals, Department of Neurochemistry and Molecular Biology, Leibniz Institute for Neurobiology, Magdeburg, Germany.

Summary

Cell-autonomous mechanisms of early synaptogenesis and their impact on the excitatory-inhibitory brain balance are poorly understood. By analyzing binding motifs in cytoplasmic regions of synaptogenic cell adhesion molecules, we identified a tumor necrosis factor receptor-associated factor 6 (TRAF6) binding motif in neuroplastin. Three-dimensional molecular modelling and biochemical approaches identified amino acids in neuroplastin binding the TRAF-C of TRAF6 with micromolar affinity. TRAF6 is required for spinogenesis and its association with neuroplastin fostered formation of new postsynapses in young hippocampal neurons. Also, TRAF6 is strictly necessary to restore failed spinogenesis in neuroplastin-deficient neurons via neuroplastin expression. These features are independent from neuroplastin's extracellular adhesive properties or its known interaction with plasma-membrane Ca^{2+} ATPases. Furthermore, TRAF6-mediated neuroplastin-dependent spinogenesis determinates the excitatory synapse density and in turn the balance of E-I synapses in mature neurons. These findings provide a highly specific cell-encoded mechanism for early synaptogenesis crucial for neuronal connectivity.

Keywords

Synapse formation, excitatory spinogenesis, TRAF6, E-I synapse balance, neuroplastin, dendritic protrusion, neuronal connectivity.

Introduction

Synapse formation is a highly coordinated cellular process, which sets up neuronal connectivity in the developing nervous system. Indeed, correct establishment of synapses is fundamental for the information flow in the healthy brain (McAllister 2007; Südhof 2008, 2017) and inaccuracies in synapse formation can underlie altered connectivity in neurological disorders including mental retardation, autism spectrum disorders, and schizophrenia (Südhof 2008, 2017; Zhang et al, 2009; Boda et al, 2010). The early appearance of synaptogenic structures in young dendrites, named dendritic protrusions or filopodia, does not seem to be triggered by neuronal activity (Verhage et al., 2000; Sando et al., 2017; Sigler et al. 2017), global intracellular calcium transients (Lohmann et al., 2005; Lohmann and Bonhoeffer 2008) or calcium-dependent signaling (Zhang and Murphy 2004). Rather, it might be controlled by cell-autonomous expression of synaptogenic molecules (Okawa et al., 2014; Jiang et al., 2017a; Südhof 2017). Currently, there is only limited knowledge on how such molecules instruct and organize the formation of synapses. Also, it has not been appreciated how early synaptogenic events contribute to future connectivity or plasticity of mature neurons (Yoshihara et al., 2009; Südhof 2017).

Recently, the tumor necrosis factor (TNF) receptor-associated factor 6 (TRAF6) has been shown critical for spine plasticity in mature neurons (Ma et al., 2017). Also, TRAF6 has been linked to neuronal survival and apoptosis by regulating p75 neurotrophin receptors (p75NTR) degradation and signaling (Lomaga et al., 2000; Yeiser et al., 2004 Geetha et al., 2005). Nevertheless, whether TRAF6 plays a role in synapse formation in young neurons is unknown. TRAF6 is a largely recognized mediator of inflammatory cell processes, differentiation, activation and tolerance of immune cells as well as morphology and migration of cancer cells (Lomaga et al., 1999; Kobayashi et al., 2001; Xie 2013). TRAF6 is a cytoplasmic E3 ligase and an adaptor protein with an N-terminal region formed by a RING domain and four zinc fingers and a C-terminal region that comprises a coiled coil domain and a TRAF-C domain. The TRAF-C domain is responsible for the binding of TRAF6 to a specific motif in cytoplasmic domains of transmembrane proteins (Chung et al., 2002; Yin et al., 2009). Upon activation, TRAF6 dimers undergo *homo*-oligomerization by lateral engagement of neighboring RING domains leading to a high-order assembly of a three-dimensional lattice-like structure (Yin et al., 2009; Ferrao et al., 2012; Wu 2013). This high-order TRAF6 structures are reported as plasma membrane-associated "fluorescent spots"

on the micrometer scale where hundreds of cell signaling intermediaries would nest (Ferrao et al., 2012; Wu 2013).

Neuroplastin is a type-1 transmembrane glycoprotein with a short intracellular tail (Langnaese et al., 1997; Beesley et al., 2014). The isoform neuroplastin 55 (Np55) displays two extracellular immunoglobulin-like (Ig-like) domains whereas neuroplastin 65 (Np65) has an additional N-terminal Ig-like domain with *trans*-homophilic adhesive capacity (Smalla et al., 2000; Owczarek et al., 2011; Herrera-Molina et al., 2014). In humans, neuroplastin has been associated with cortical thickness and cognitive capabilities in adolescents (Desrivieres et al., 2015), schizophrenia (Saito et al., 2007), Alzheimer's disease (Ilic et al., 2018). In adult mice, neuroplastin deficiency goes along with retrograde amnesia and impaired associative learning, cortical activity, and synaptic plasticity (Bhattacharya et al., 2017; Herrera-Molina et al. 2017). Neuroplastin has been shown to mediate the formation of a specific group of excitatory synapses in the hippocampus (Herrera-Molina et al., 2014; Amuti et al., 2016) and in the inner hear (Carrott et al., 2016; Zeng et al., 2016). In hippocampal neurons, inactivation of the *Nptn* gene leads to unbalanced excitatory/inhibitory (E/I) synapse ratio resulting from a decreased glutamatergic with normal GABAergic synapse density (Bhattacharya et al., 2017; Herrera-Molina et al., 2014). The absence of neuroplastin also results in malformed glutamatergic postsynapses partnering presynapses with impaired vesicle release activity in mature neurons (Herrera-Molina et al., 2014, Amuti et al., 2016; Carrott et al., 2016). However, it remains unknown, how neuroplastin mechanistically contributes to synapse formation and whether its synaptogenic function impacts the E/I balance in mature neuronal circuits.

Here, we uncover a hitherto unanticipated function for TRAF6 in synapse formation. We identified a TRAF6 binding motif in neuroplastin, but not in others synaptogenic cell adhesion molecules (CAMs). We show the molecular interface and properties of the TRAF6-neuroplastin binding and that; TRAF6 confers to neuroplastin a specific and cell-autonomous capacity to control spinogenesis. Furthermore, we demonstrate that neuroplastin-TRAF6-initiated spinogenesis is essential to establish the density of excitatory synapses what directly implicates it in the determination of the E/I synapse balance of mature neurons.

Results

Neuroplastins promote early spinogenesis in young hippocampal neurons

To assess the role of neuroplastin in early spinogenesis, we studied the formation of filopodia-like protrusions from dendrites, which can act as precursors of dendritic spines, i.e. postsynaptic compartments of glutamatergic synapses (Ziv and Smith, 1996; McClelland et al., 2010). By confocal microscopy we quantified the number of protrusions per 10 µm length expanding from MAP2-stained dendrites of GFP-filled pyramidal neurons (Figure 1A,B). This indicated that ablation of neuroplastin gene expression results in reduced density of dendritic protrusions in $Nptn^{-/-}$ compared to $Nptn^{+/+}$ hippocampal neurons at 9 DIV. The phenotype was rescued by transfection of mutant neurons at 6-7 DIV with recombinant neuroplastin isoforms Np55-GFP or Np65-GFP (Figure 1C). In parallel experiments with rat primary hippocampal neurons, we observed that the over-expression of either neuroplastin isoforms promotes dendritic protrusion density (Figure 1D,E). Rat neurons transfected with either Np55-GFP or Np65-GFP at 7 DIV displayed higher density of dendritic protrusion than control GPF-transfected neurons when evaluated at 8 DIV (Figure 1D,E) or 9 DIV (cf. Figure 1F). Surprisingly, later transfections of Np65- or Np55-GFP performed at 9 DIV were less effective to raise the protrusion density in rat neurons analyzed at 10 or 11 DIV (Table 1). Therefore, expression of either Np65 or Np55 is most effective to raise the density of dendritic protrusions in cultured neurons during the early period of spinogenesis, i.e. around 6 to 9 DIV.

To characterize the nature of neuroplastin-promoted dendritic protrusions, we evaluated the appearance of Shank2, a well-established early marker (Grabrucker et al., 2011; Sarowar and Grabrucker 2016) and key organizer (Roussignol et al., 2005) of functional postsynapses. While transfection of Np65-GFP at 7DIV increased dendritic protrusion density compared to transfection with control GFP, the relative abundance of Shank2-positive vs. Shank2-negative protrusions (Protrusion fraction) was not different between Np65-GFP-tranfected and GFP-transfected rat neurons at 9 DIV (Figure 1F,H). Although area and intensity of Shank2 clusters were not different between Np65-GFP- and GFP-expressing dendrites at 9 DIV (Figure 1I,J), the number of Shank2 clusters per dendritic protrusion was higher in Np65-GFP- vs. GFP-expressing dendrites at 9 DIV (Figure 1K). The data show that Np65-GFP-overexpressing neurons display an increased number of newly formed postsynapses defined as dendritic protrusions containing a higher number of Shank2 clusters

compared to GPF-filled control neurons (Figure 1L). These results are consistent with the idea that both neuroplastin isoforms share the potential to promote the formation of dendritic protrusion during a time period of spinogenesis. Accordingly, we looked for additional *cis*-acting mechanisms related to both neuroplastins to increase the density of spinogenic dendritic protrusions.

A specific TRAF6 binding motif is present in neuroplastin, but not in others synaptogenic CAMs

To decipher underlying signaling mechanisms of spinogenesis, we sought for potential intracellular binding partners of neuroplastins employing *in silico* approaches. Using the ELM database (http://elm.eu.org/) we identified a single TRAF6 binding motif in the cytoplasmic tail of all neuroplastins from human, rat, and mouse (Figure 2A) which fully matches the well-characterized TRAF6 binding motif (Ye at al., 2002; Sorrentino et al., 2008; Yin et al., 2009). Very surprisingly, the TRAF6 binding motif was not found amount a number of other known spinogenic type-1 CAMs namely N-Cadherin (Bozdagi et al, 2010), LRRTM (Linhoff et al, 2009), neuroligins (Varoqueaux et al, 2006), neurexins (Missler et al; 2003), SynCAM1 (Robbins et al, 2010), EphB2 (Henderson et al, 2001), PTPR0 (Jiang et al., 2017), and others (Figure S1). These results highlight both the specificity and novelty of the association of TRAF6 to neuroplastin to mediate spinogenesis. After further analysis, we noticed that the alternative mini-exon-encoded Asp-Asp-Glu-Pro (DDEP) of neuroplastin is close to its TRAF6 binding motif (Figure 2A). Based on crystallographic studies on the interaction of the TRAF6 TRAF-C domain with the TRANCE receptor (Ye et al., 2002), *in silico* modelling was applied to TRAF6 TRAF-C domain-neuroplastin interaction (Figure 2B; Figure S2A). A strikingly similar three-dimensional structure was predicted for the TRAF6 binding site of neuroplastin when compared to the TRANCE receptor TRAF6 binding motif (Figure 2B). In particular, the coordinates and stereo specificity of key amino acids (Figure 2B: P_{-2}= Pro, P_0= Glu, and P_3= Aromatic/Acidic) involved in docking of the TRANCE receptor to TRAF6 TRAF-C domain (TRAF-C) were conserved in the TRAF6 binding site of neuroplastin (Figure 2C).

We decided to test whether there is a direct interaction between TRAF6 and neuroplastins. First, we characterized the binding of purified neuroplastin intracellular peptide containing the TRAF6 binding motif to immobilized recombinant TRAF6 by surface plasmon resonance (Figure 2D,E; Figure S2B,C). Binding of neuroplastin

peptide to TRAF6 was dependent on peptide concentration, saturable, and displayed a 1:1 stoichiometry. We calculated a K_d value of 88 µM for the neuroplastin-TRAF6 interaction (Figure 2D,E), which is very similar to the K_d of 84 µM for the TRANCE receptor-TRAF6 interaction (Yin et al., 2009). To establish whether the TRAF6 motif in neuroplastin binds TRAF6 in living cells, we performed co-immunoprecipitation assays from HEK cells transfected with different GFP-tagged constructs of neuroplastins and flag-tagged TRAF6. Due to alternative splicing of the primary transcript both major neuroplastin isoforms can contain an alternative DDEP insert close to their TRAF6 binding site. To consider potential differences in binding, splicing variants with and without DDEP were tested. As shown in Figure 2F,G both splice variants of Np65-GFP co-precipitated flag-TRAF6 suggesting that the mini exon-encoded insertion is not critical for the interaction. Similarly, Np55 with and without DDEP insertion co-precipitated with TRAF6 (Figure S2D). In contrast, co-precipitation was largely abolished when GFP-tagged versions of Np65 either with deleted TRAF6 binding motif (Np65Δ-GFP) or with triple (Np65$_{PED}$-GFP) or single (Np65$_P$-GFP) amino acid substitutions in the binding site (Figure 2F-G) were used. Additionally, pull-down assays demonstrate that Np65-GFP isolated from HEK cell homogenates equally well binds equally well to purified recombinant GST-TRAF6 or to GST-TRAF6 C-domain (coiled coil and TRAF-C domains GST-TRAF6$_{cc-c}$) (Figure EV1B,C). Finally, neuroplastin also co-precipitates TRAF6 from total brain homogenates (Figure S2E). Based on these analyses, we conclude that the TRAF6 binding site in the cytoplasmic tail of neuroplastins is fully capable of binding the TRAF-C domain of TRAF6.

TRAF6 mediates the formation filopodial structures by neuroplastin

We have reported disorganization of polymerized actin in dendrites of $Nptn^{-/-}$ primary hippocampal neurons (Herrera-Molina et al., 2014). Coincidently, TRAF6 has been associated to regulation of actin polymerization (Armstrong et al., 2002; Wang et al., 2006; Yamashita et al., 2008). As a first test to understand the significance of the neuroplastin-TRAF6 interaction, we therefore performed experiments to explore if and how neuroplastins and TRAF6 interact to regulate actin-based filopodia formation in HEK cells. Over-expression of Np65 or Np55 in HEK cells was sufficient to induce a massive increase of filopodia number and length as compared to control cells transfected with either soluble or membrane-attached GFP (Figure 3A-C). Variants of Np55 or Np65 lacking the DDEP insert were as effective as the ones that carry the

insert to promote filopodial structures (Figure S3A-D). The capacity of neuroplastin to promote filopodia was eliminated by mutants abolishing TRAF6 binding (i.e. Np65Δ-GFP, Np65PED-GFP, Np65P-GFP) (Figures 3A-C). Furthermore, after decreasing protein levels of endogenous TRAF6 by ~80% using a specific siRNA (Figure S3E,F), neither expression of Np65-GFP nor of Np55-GFP did increase the number or length of filopodia in HEK cells (Figure 3A-C). Thus, Np55 and Np65 (±DDEP) are equally effective to promote the formation of filopodial structures and they seem to require endogenous TRAF6 and binding to their TRAF6 motifs to do so.

TRAF6 translocates from the cytoplasm to the membrane by recruitment to integral membrane proteins with TRAF6 binding domains (Yin et al., 2009; Wu 2013). Therefore, we tested whether neuroplastins *via* their C-terminal TRAF6 binding motif have the capacity to recruit endogenous TRAF6 to the plasma membrane. In HEK cells transfected with GPI-anchored GFP or with Np65Δ-GFP, TRAF6 immunoreactivity was primarily located in the cytoplasm (Figure 3D). In contrast, TRAF6 immunoreactivity was abundantly associated with the plasma membrane in cells expressing recombinant Np65-GFP (Figure 3D) or other variants of neuroplastin (Figure S3A). Analyses of co-distribution (Figure 3E) and co-localization (Figure 3F) confirmed that plasma membrane-associated TRAF6 co-localizes with Np65. Thus, neuroplastin can recruit TRAF6 to the plasma membrane and thereby regulate its subcellular localization. This is independent of the presence or absence of the DDEP insert.

Next, we asked whether the recruitment and binding of TRAF6 by neuroplastin mediate filopodia formation. To test this prediction, we co-expressed GFP-tagged TRAF6 (TRAF6-GFP) with Np55-RFP. Clearly, co-expression of TRAF6-GFP fostered the increase of filopodia number by Np55-GFP (Figure 3G-I). Very interestingly, endogenous TRAF6 and TRAF6-GFP co-localized with Np55-RFP in filopodia-associated microscopic spots (Figure 3G). Indeed, analyses of fluorescent intensity and distribution revealed high co-localization of TRAF6-GFP with Np55-RFP in single spots of filopodia (Figure 3J). The potential involvement of the N-terminal RING domain of TRAF6 was tested using TRAF6cc-c-GFP containing the coiled coil and TRAF-C domains and lacking the N-terminal domain (see Figure S2B,C). Despite being recruited to the plasma membrane and co-localized with Np55-RFP (Figure 3G,J), TRAF6cc-c-GFP blocked neuroplastin-induced filopodia formation (Figure 3G-I). Accordingly, the recruitment and binding of TRAF6cc-c by neuroplastin is insufficient to promote filopodial structures. Because RING domain is well-known to be responsible

for three-dimensional assembly of functional TRAF6 lattice-like structures (Yin et al., 2009; Ferrao et al., 2012; Wu 2013), we conclude that only the recruitment and binding of fully functional TRAF6 can mediate formation of filopodial structures by neuroplastin.

TRAF6 confers the spinogenetic capacity to neuroplastin

So far, our results document the capacity of TRAF6 to mediate neuroplastin-promoted filopodia formation in non-neuronal cells, i.e. HEK cells. Next, we tested the hypothesis that TRAF6 is essential for the spinogenic function of neuroplastin in hippocampal pyramidal neurons. First, we used $Nptn^{-/-}$ hippocampal neurons with significantly reduced number of spinogenic protrusions (Figure 1). The protrusion density in $Nptn^{-/-}$ dendrites expressing Np65-GFP was higher than in control $Nptn^{-/-}$ dendrites expressing GFP at 9 DIV (Figure 4A,B). Clearly, Np65Δ-GFP failed to rescue the dendrite protrusion density in $Nptn^{-/-}$ neurons (Figure 4A,B). In independent experiments, we found that Np65Δ-GFP-failed to promote the density of dendritic protrusions as Np65Δ-GFP-expressing rat neurons displayed a similar density of dendritic protrusions as GPF-expressing rat neurons at 8 DIV (Figure 4C,D). Additionally, we evaluated whether Np65Δ-GFP affects the number of protrusions and interferes with the normal enrichment of Shank2 in dendritic protrusions in rat neurons at 9 DIV. The density of dendritic protrusions and the distribution of Shank2-positive vs. Shank2-negative protrusions were similar between GFP- and Np65Δ-GFP-expressing rat neurons at 9DIV (Figure 4E-G). These data show that, in contrast to Np65-GFP, Np65Δ-GFP did neither rescue impaired spinogenesis in $Nptn^{-/-}$ neurons nor increase the number of dendritic protrusions in rat neurons. Thus, independently of the rodent model from which neurons were derived, Np65 requires its TRAF6 motif to regulate the density of spinogenic protrusions in hippocampal neurons.

Next, we evaluated whether endogenous TRAF6 is involved in neuroplastin-mediated spinogenesis in rat hippocampal neurons at 9 DIV. We used the small molecule inhibitor (SMI) 6860766, which binds the TRAF-C domain of TRAF6 and abrogates its capacity to interact with its binding sites (van den Berg et al., 2014; Chatzigeorgiou et al., 2015) (for simplicity hereafter called SMI TRAF6). Addition of SMI TRAF6 (2 µM) reduced the density of protrusions in GFP- and in Np65-GFP-expressing neurons compared to vehicle-treated (0.01% DMSO) GFP- and Np65-GFP-expressing neurons, respectively (Figure 4H,I). SMI TRAF6 application did not decrease significantly the distribution of Shank2-positive vs. Shank2-negative

protrusions in GFP-filled neurons (Figure 4J). However, treatment with SMI TRAF6 decreased the fraction of Shank2-positive protrusions in Np65-GFP-expressing neurons slightly but significantly (Figure 4J). SMI TRAF6 also decreased the area, but not the intensity of Shank2 clusters, and it reduced the number of Shank2 clusters per protrusion in Np65-GFP-expressing neurons to the level of controls (Figure 4K-M). Moreover, SMI TRAF6 treatment evidenced that the size of Shank2 clusters depends on TRAF6 (Figure 4K). We conclude that neuroplastin strictly requires both its TRAF6 binding site and TRAF6 to increase the density of dendritic protrusions in hippocampal neurons. Either deficiency of these pre-requisites abrogates the spinogenic capacity of neuroplastin (Figure 4N).

As additional evidence for functional interaction during early spinogenesis, we confirmed the co-localization/co-distribution of TRAF6 and neuroplastin in dendritic protrusions using high-resolution microscopy and image deconvolution procedures in young neurons but not in mature neurons (Figure S4). In addition, we evaluated the density of dendritic protrusions after decreasing levels of TRAF6 by siRNA knockdown (as characterized in Figure S3E,F) in rat neurons transfected with GFP-, Np65-GFP or Np65Δ-GFP at 9 DIV. When TRAF6 levels were knocked down by 60% or more, the dendritic protrusion density was reduced in GFP-, Np65-GFP-, and Np65Δ-GFP-expressing neurons (Figure S3C,D).

Very recently, it has been discovered that neuroplastin interacts with all four plasma membrane Ca^{2+} ATPases (PMCA1-4) stabilizing the surface expression of these pumps (Bhattacharya et al., 2017; Herrera-Molina et al., 2017; Korthals et al., 2017; Schmidt et al., 2017; Gong et al., 2018). Thus, we addressed the obvious question whether TRAF6 or neuroplastin require PMCA to promote dendritic protrusion density. First, we examined whether the TRAF6 binding motif of neuroplastin is necessary to maintain PMCA protein. Consistent with our previous report (Herrera-Molina et al., 2017), full-length Np65-GFP increased protein levels of PMCA2 compared to GFP when co-transfected in HEK cells (Figure S5A,B). Np65Δ-GFP was found as effective as Np65-GFP to increase PMCA2 levels in co-transfected HEK cells (Figure S5A,B). In rat hippocampal neurons at 9 DIV, confocal microscopy revealed that both Np65-GFP and Np65Δ-GFP co-localized with endogenous PMCA and increased PMCA protein levels (Figure S5C,D). Also, SMI TRAF6 treatment from 7-9 DIV did not modified total PMCA protein levels in young hippocampal neurons (data not shown). Therefore, either elimination of the TRAF6 binding motif of neuroplastin or

SMI TRAF6 treatment abrogates TRAF6-mediated spinogenetic function (Figure 4), but does not affect endogenous PMCA protein levels or the capacity of neuroplastin to promote the expression levels of the pump. Although PMCA inhibition with Caloxin 2a1 seemed to be slightly enlarge protrusions, the density of protrusions was not affected in GFP-filled (Figure S5E,F) nor in Np65-GFP-expressing neurons at 9 DIV (data not shown). Thus, PMCA is not critically required by TRAF6 to mediate neuroplastin-promoted spinogenesis.

TRAF6 mediates spinogenesis and contributes to the balance of number and activity of synapses

To our knowledge, TRAF6 has never being linked to spinogenesis or synapse formation. Based our observation that TRAF6 is essential mediating spinogenesis controlled by neuroplastin in young neurons, we wondered whether early TRAF6 inhibition results in a reduced number of excitatory synapses. To test this, rat hippocampal neurons were treated with SMI TRAF6 (2 µM) or with vehicle (0.01% DMSO) from 7 to 9 DIV and analyzed at 12 DIV (Figure 5A). The number of excitatory synapses (homer-positive spines matching synapsin-positive presynapses, Herrera-Molina et al., 2014) per 10 µm dendrite was significantly reduced in neurons treated with SMI TRAF6 from 7 to 9 DIV (Figure 5B) as compared to vehicle-treated neurons. In contrast, neurons treated with SMI TRAF6 from 10 to 12 DIV and analyzed at 12 DIV displayed similar number of synapses as vehicle-treated neurons (Figure 5B). These data indicate that TRAF6 is critical for the early spinogenesis of some ~25% of hippocampal excitatory synapses *in vitro*.

Next, we quantified the area and fluorescence intensity of the homer-positive and synapsin-positive puncta that were still present after the treatment with SMI TRAF6. The area of homer-positive puncta was reduced in rat neurons treated with SMI TRAF6 compared to control vehicle-treated rat neurons from 7 to 9 DIV. Neurons treated with SMI TRAF6 from 10 to 12 DIV displayed similar area of homer clusters compared to control vehicle-treated neurons (Figure 5C). The intensity of homer-positive puncta as well as the area and intensity of synapsin-positive puncta were not affected by SMI TRAF6 (Figure 5C-D). Thus, TRAF6 inhibition during spinogenesis from 7 to 9 DIV decreases the enrichment of the postsynaptic marker homer, but not of the presynaptic protein synapsin. This phenotype resulting from TRAF6 inhibition is

very similar to the one previously reported in neuroplastin-deficient neurons (Herrera-Molina et al., 2014).

TRAF6 is critical for early spinogenesis of Shank2-containing spines at 9 DIV (Figure 4 and Figure S3) and for the formation of excitatory synapses at 12 DIV (Figure 5). Thus, we tested whether the presynaptic activity is consequentially affected by TRAF6 inhibition in mature synapses. Not surprisingly, evaluation of the presynaptic uptake of synaptotagmin-1 antibody – reporting vesicle release and recycling driven by intrinsic network activity – showed a decreased activity in mature VGAT-negative excitatory presynapses after treatment with SMI TRAF6 from 7 to 9 DIV, but not from 10 to 12 DIV (Figure 5F). Therefore, in addition to the anatomical evidence (Figure 4, Figure S4, and Figure 5), this result shows that spinogenetic function of TRAF6 in young neurons is critically linked to a long-term establishment of a balanced E/ I synapse activity in mature neurons.

Interestingly, the activity of VGAT-positive inhibitory terminals was also reduced in at 16 DIV when neurons were treated with SMI TRAF6 from 7 to 9 DIV, but not from 10 to 12 DIV (Figure 5F). Similarly, VGAT-positive inhibitory terminals placed on the soma of neurons displayed decreased activity after the treatment with SMI TRAF6 from 7 to 9 DIV, but not from 10 to 12 DIV (not shown). To unravel the nature of these results, we calculated statistical parameters to estimate the area of vesicle release (mean area of puncta) and the activity level (mean intensity per pixel) for each presynapse type. From these data (Table 2), we concluded that decreased activity of inhibitory synapses results from an adaptation of the network activity in response to the earlier reduction of the density of spinogenic protrusions directly caused by disruption of the neuroplastin-TRAF6 interaction or by TRAF6 activity from 7 to 9 DIV (Figures 1, 4, Figure S4, and Figure 5). Consistent with this explanation, TRAF6 inhibition after early spinogenesis (10 to 12 DIV) did not affect the formation of protrusion nor resulted in lower number of excitatory synapses. Furthermore, neurons treated with TRAF6 inhibitor from 10 to 12 DIV kept unchanged and balanced excitatory/ inhibitory presynaptic activities.

Discussion

Our study addresses the question of how neurons set an adequate density and functional balance of synapses to communicate with each other. In particular, we demonstrate a novel role for TRAF6 – mainly know as a key mediator of inflammation and cell morphology and proliferation of cancer cells – in early glutamatergic synaptogenesis, establishment of the correct density of excitatory synapses as well as in proper balance of E/I synapse function through its binding to neuroplastin.

The discovery of TRAF6 as strictly necessary for early spinogenesis reveals a novel function of this factor. Our data demonstrate that TRAF6 is effective to mediate neuroplastin-dependent spinogenesis during a defined time window in neuronal development, i.e. when most dendritic protrusions are forming (6 to 9 DIV). After this time period, neither TRAF6 inhibition nor neuroplastin expression modified the density of protrusions or the number of excitatory synapses. Furthermore, we observed a regularly spaced and cell membrane-associated co-localization of TRAF6 with neuroplastin in dendritic protrusions of young neurons only. In contrast, TRAF6 did not co-localized with neuroplastin in mature neurons. This can be explained by a developmental shift of the binding of TRAF6 to neuroplastin in young neurons towards PSD-95, a much more prominent interaction partner, in mature neurons. Accordantly, PSD-95 levels are lower than neuroplastin levels in young neurons (Buckby et al., 2004) but highly abundant in all mature glutamatergic spines. Consistent with this proposition, it has been reported recently that TRAF6 efficiently binds to and regulates the stability of PSD-95 as a necessary step for the plasticity of hippocampal mature spines (Ma et al, 2017). Alternatively, neuroplastin could also change its predilection for interaction partners in young neurons as it becomes an essential subunit for the four paralogs in mature neurons (Bhattacharya et al., 2017; Herrera-Molina et al., 2017; Korthals et al., 2017; Schmidt et al., 2017; Gong et al., 2018).

We demonstrated here that Np55 and Np65 harbor a single intracellular TRAF6 binding site. The intracellular sequence RKRPDEVPD of neuroplastin fulfilled structural and three-dimensional criteria to be considered as a proper TRAF6 binding site (Ye et al., 2002; Sorrentino et al., 2008; Yin et al., 2009). The binding affinity between neuroplastin-derived peptide carrying the TRAF6 binding site with TRAF6 C-terminal domain was very close to the expected one according to solid crystallographic, structural and functional data demonstrating the specialization of the TRAF6 binding site for binding TRAF6 (Ye et al., 2002; Sorrentino et al., 2008; Yin et al., 2009). Co-

precipitation between the neuroplastin isoforms with TRAF6 was not affected by the presence of the alternative splicing DDEP proximal to the TRAF6 binding site. However, it was drastically reduced or eliminated by mutation or deletion of the TRAF6 binding site of neuroplastin. Endogenous TRAF6 and GFP-tagged TRAF6 were recruited into the regularly spaced cell membrane-associated puncta by neuroplastin only when an intact TRAF6 binding site was present. This is coherent with TRAF6 moving towards the cell membrane and forming micrometric and geometrically organized lattice-like supramolecular structures able to regulate the multimerization of bound transmembrane proteins and to host downstream cell signaling elements (Schultheiss et al., 2001; Yin et al., 2009; Ferrao et al., 2012; Wu et al., 2013). Elimination of the lattice-forming RING domain did not prevented TRAF6 recruitment by neuroplastin but abrogated the capacity of the transmembrane glycoprotein to promote the formation of filopodial structures. Thus, it is realistic to propose that upon TRAF6 binding and higher-order oligomerization of the factor, neuroplastin might become a part of such supramolecular complexes to initiate downstream events of cell signaling causing the formation of filopodial structures in HEK cells and spinogenic protrusion in neurons.

The selective and specific binding of TRAF6 places neuroplastin as novel candidate to organize prolific signaling mechanisms related to spinogenesis and/or to stabilization young spines. Indeed, preliminary experiments suggest that inhibition of p38 MAPK, ERK1/2 or PI3 kinase reduces the number of neuroplastin-promoted filopodia in HEK cells and dendritic protrusions in young hippocampal neurons (Vemula and Herrera-Molina, unpublished data). It has been reported that extracellular engagement of neuroplastin activates p38 MAPK (Empson et al., 2006), ERK1/2 and PI3 kinase (Owczarek et al., 2011). These signalling pathways could also be related to homophilic *trans*-synaptic engagement of Np65 to foster and or stabilize young protrusion. The literature recognized TRAF6 largely as a main upstream activator of transcriptional factor NFκB pathway (Darnay et al., 1999; Xie 2013). In young hippocampal neurons, NFκB activity is not regulated by neuronal activity but, it is necessary for the formation of excitatory synapses (Boersma et al., 2011). Also, the constitutively high NFκB activity in young neurons maintains glutamatergic spinogenesis contributing in turn to the future establishment of E/I synapse balance in mature neurons (Boersma et al., 2011; Dresselhaus et al., 2018). Preliminary experiments suggest that inhibition of NFκB nuclear translocation reduces the density

of neuroplastin-induced dendritic protrusions in young hippocampal neurons. Thus, it is possible that neuroplastin regulates NFκB activity in a TRAF6-dependent manner as a necessary step to promote the formation of filopodial structures in HEK cells and spinogenic protrusions in hippocampal neurons – a hypothesis that needs to be tested in future.

TRAF6 is strictly required by neuroplastin to promote spinogenesis in young hippocampal neurons. Expression of Np55 or Np65 equally rescued the reduced number of dendritic protrusions in $Nptn^{-/-}$ neurons and enhanced normal formation of spinogenic protrusions in rat neurons upon overexpression. This implies that TRAF6 is a spinogenic mechanism that do not essentially dependent of the Np65-specific *trans*-adhesive extracellular domain (Smalla et al., 2000). In mouse models, elimination of both Np55 and Np65 is required to reduce the density of hippocampal excitatory synapses (Herrera-Molina et al., 2014, Amuti et al., 2016; Bhattacharya et al., 2017) and ribbon synapses (Carrott et al., 2016). In contrast, elimination of Np65 only is not sufficient to alter the synapse density; however, it results in morphological alterations of hippocampal spines in Np65 KO mice (Amuti et al., 2016). These data, however, do not rule out the possibility of a later participation of the Np65-specific adhesive Ig-like domain in the *trans*-stabilization of postsynapses (Smalla et al., 2000; Herrera-Molina et al., 2014). This needs to be addressed by future experiments. The present finding that over-expression of Np65 fosters the formation Shank2-containing protrusions indicates that these additional and newly formed structures are spinogenic in transfected neurons under development. Additionally, this finding matches with previous observations, i.e. a tight correlation between the fast up-regulation of expression of neuroplastin and intracellular synaptic proteins during early synaptogenesis in cultured and in acutely isolated rat hippocampal slices (Buckby et al., 2004) as well as the stabilization and maturation of spines (Böckers et al., 1999; Sarowar and Grabrucker 2016).

TRAF6 mediated spinogenesis seems to be rather restricted to neuroplastin, because TRAF6 binding motif in not present in other spinogenic CAMs. Cell adhesion molecules have been proposed as key participants in synapse formation. Nonetheless, knocking out *trans*-synaptic CAMs, that are crucially involved in synapse maturation, transmission and/or plasticity, does not significantly affect the number of synapses formed *in vivo* (Henderson et al., 2001; Missler et al.; 2003; Varoqueaux et al., 2006; Chubykin et al., 2007; Linhoff et al., 2009; Bozdagi et al., 2010; Robbins et al., 2010).

Thus, it has been considered that still unexplored molecular machineries interplaying with CAMs may promote synapse formation (Yoshihara et al., 2009; Jiang et al., 2017; Jang et al., 2017; Südhof 2017). Independent of their adhesive properties some CAMs can form transmembrane complexes in *cis* that in turn can recruit intracellular proteins and activate signaling mechanisms underlying neuronal development (Cavallaro and Dejana, 2011; Jang et al., 2017). Our results shown that neuroplastin contributes to the formation of spinogenic dendritic protrusions cell-autonomously and, the binding of TRAF6 confers to neuroplastin unique mechanistic possibilities to potentially regulate cell signalling and gene expression related to control spinogenesis.

Figure Legends

Figure 1. Neuroplastin regulates spinogenesis.

A-**C**. Reduced number of dendritic protrusions in *Nptn*$^{-/-}$ compared to *Nptn*$^{+/+}$ mouse primary hippocampal neurons at 9 DIV. (**A**) *Nptn*$^{-/-}$ and *Nptn*$^{+/+}$ neurons transfected with GFP-encoding plasmids at 6-7 DIV using Lipofectamine. At 9 DIV, neurons were fixed and stained with anti-GFP antibodies followed by an Alexa 488-conjugated antibody to enhance their intrinsic fluorescence (green) and with anti-MAP2 antibodies followed by a proper secondary antibody to detect dendrites (magenta). Images were obtained using a confocal microscope. Scale bar=100 μm. (**B**) Protrusion density (number of dendritic protrusions per 10μm) of GFP-filled *Nptn*$^{-/-}$ and *Nptn*$^{+/+}$ neurons (circles) is expressed as mean ± SEM from three independent cultures. ***p<0.001 between genotypes using Student's t-test (*Nptn*$^{+/+}$ GFP=4.12 ± 0.18 N=33; *Nptn*$^{-/-}$ GFP=1.72 ± 0.19 N=36). (**C**) Protrusion density of GFP-, Np65-GFP- or Np55-GFP-expressing *Nptn*$^{-/-}$ neurons from two independent cultures. ***p<0.001 or **p<0.01 vs. *Nptn*$^{-/-}$ GFP

using Student's t-test ($Nptn^{-/-}$ GFP=1.92 ± 0.22 N=26; $Nptn^{-/-}$ Np65-GFP=3.67 ± 0.18 N=20; $Nptn^{-/-}$ Np55-GFP=3.77 ± 0.19 N=26).

D, **E**. Both neuroplastin isoforms increase dendritic protrusion density in rat neurons at 8 DIV. (**D**) Confocal images show rat neurons transfected with plasmids encoding GFP, Np65-GFP or Np55-GFP at 7 DIV. At 8 DIV, neurons were fixed and stained with anti-GFP antibodies followed by an Alexa 488-conjugated antibody (white). Scale bar=10 µm (**E**) Protrusion densities of 40-50 neurons per group (circles) from three-four independent cultures. ***p<0.001 or **p<0.01 vs. GFP transfected cells using Student's t-test (GFP=1.95 ± 0.19 N=39; Np65-GFP=3.23 ± 0.14 N=56; Np55-GFP=3.58 ± 0.16 N=38).

F-**H**. Overexpression of Np65-GFP increases the number of newly formed Shank2-containing spines. (**F**) Confocal images of dendritic segments of rat neurons transfected with GFP or Np65-GFP at 7 DIV. At 9 DIV, neurons were fixed and stained with primary antibodies against GFP (white) and Shank2 (red). Scale bar=10 µm (**G**) Protrusion density (GFP=3.151 ± 0.182 N=48; Np65-GFP=4.642 ± 0.145 N=54) and (**H**) Distribution of Shank2-positve and Shank2-negative protrusions were calculated as a fraction from 40-50 neurons per group from three independent experiments. Plots display mean ± SEM as indicated. **p<0.01 for Np65-GFP vs GFP using Student's t-test (Shank2(+): GFP=0.54 ± 0.07; Np65-GFP=0.60 ± 0.06).

I-**K**. (**I**) Size of puncta (area; GFP=0.10 ± 0.01 N=747; Np65-GFP=0.11 ± 0.01 N=738), (**J**) Fluorescence intensity (GFP=127.6 ± 2.1; Np65-GFP=131.5 ± 1.9) and (**K**) Number of Shank2 clusters/protrusion in neurons (GFP=1.46 ± 0.17 N=43; Np65-GFP=1.91 ± 0.18 N=49) of the experiments displayed in Figure 1F. *p<0.05 between Np65-GFP-expressing and GFP-expressing neurons using Student's t-test.

L. The upper sketch on the left illustrates dendritic protrusions enriched on Shank2 in control GFP-filled hippocampal neurons at 9 DIV. Np65-GFP-expressing neurons (lower sketch) display more spinogenic protrusions with Shank2 clusters.

A

Np65 intracellular tail:
Homo sapiens E$_{361}$K**R**K**R**P**DE**VPD<u>**DDE**PAG</u>$_{377}$
Rattus norvegicus E$_{360}$K**R**K**R**P**DE**VPD<u>**DDE**PAG</u>$_{376}$
Mus musculus E$_{360}$K**R**K**R**P**DE**VPD<u>**DDE**PAG</u>$_{376}$
TRAF6 site: **(Bs)**xx**P**x**E**xx**(Ar/Ac)**

D

E

K_d = 87,73 x10^{6}M
Np peptide (M^{-4})

B **C**

Neuroplastin: K R **P** D **E** V P **D**
TRANCE-R: Q M **P** T **E** D E **Y**
P$_{-2}$ P$_0$ P$_3$

F
Np65-GFP: E3$_{60}$K**R**K**R**PDEVPD<u>DDE</u>PAG......N-GFP
Np65$_{DDEP(-)}$-GFP: E$_{360}$K**R**K**R**PDEVPDAG.....N-GFP
Np65$_\Delta$-GFP: E$_{360}$K**R**KR.....N-GFP
Np65$_{PED}$-GFP: E$_{360}$K**R**KRADAVPAAG.....N-GFP
Np65$_P$-GFP: E$_{360}$K**R**KRADEVPDAG.....N-GFP

G

Figure 2. Neuroplastin binds TRAF6.

A. Potential TRAF6 binding site in the intracellular tail of neuroplastin 65 (and identical in neuroplastin 55) fits the canonical and specific motif recognized by TRAF6. The alternatively spliced DDEP sequence is underlined. Bs, Ac, and Ar stand for basic, acidic, and aromatic amino acids, respectively.

B, **C**. Neuroplastin-TRAF6 binding *in silico*. (**B**) Three-dimensional model of the TRAF6 binding site in the intracellular tail of neuroplastin (cyan) and key amino acids responsible for the binding to TRAF6 fit to the well-known TRAF6 binding motif present in the TRANCE receptor (green). (**C**) Docking of the TRAF6 C-domain with the TRAF6 binding site of neuroplastin. Similar to the binding of TRANCE receptor to TRAF6 documented by a crystallographic study (Yin et al., 2009), interaction of neuroplastin with TRAF6 would be mediated by the Proline (P) in the position P$_{-2}$ Glutamic acid (E) in P0, and Aspartic acid (D) in P$_3$.

D, **E**. Direct binding of the neuroplastin-derived intracellular peptide comprising the TRAF6 binding site to purified recombinant TRAF6. The binding curve (**D**) and the binding curve (**E**) for the neuroplastin-TRAF6 binding where obtained using surface plasmon resonance.

F, **G**. Neuroplastin-TRAF6 co-precipitation is drastically decreased by deletion or mutation of key amino acids in the TRAF6 binding site of neuroplastin. (**F**) Neuroplastin constructs included into the experiments. (**G**) HEK cells were co-transfected with constructs encoding either GFP, Np65-GFP or $Np65_{DDEP(-)}$-GFP and with TRAF6-flag or flag alone for 24 hours. Alternatively, cells were co-transfected with GFP, Np65-GFP, Np65Δ-GFP (TRAF6 binding site deficient construct), $Np65_{PED}$-GFP (containing a TRAF6 binding site with triple substitution to alanine) or $Np65_P$-GFP (with single substitution to alanine) and with TRAF6-flag or flag constructs for 24 hours. After homogenization, anti-GFP antibody-coupled beads were used to precipitate GFP-tagged complexes. We used anti-Flag or anti-GFP antibodies to detect the proteins as indicated.

Figure 3. Neuroplastin requires its TRAF6 binding site and TRAF6 to promote filopodia formation.

For these experiments, HEK cells were transfected with plasmids coding for either soluble GFP, membrane-attached GFP-GPI, DDEP insert-containing isoforms Np65-GFP or Np55-GFP, TRAF6 binding site-deficient Np65Δ-GFP, Np65$_{PED}$-GFP (containing triple substitution to alanine in the TRAF6 binding site) or Np65$_P$-GFP (with single substitution to alanine), full-length TRAF6-GFP or coiled coil-TRAF-C-GFP (TRAF6$_{cc-c}$-GFP). To knockdown endogenous TRAF6, cells were transfected with siRNA against TRAF6 or scrambled siRNA for 24 hours before the transfection with plasmids. After 24 hours, cells were fixed with methanol and immunostained with anti-GFP rabbit antibody overnight and with an Alexa-488 secondary antibody. Alternatively, cells were additionally stained with an anti-TRAF6 rabbit antibody followed by a Cy3-conjugated secondary antibody and DAPI.

A-**C**. Deletion or mutations of the TRAF6 binding site of neuroplastin or knockdown of endogenous TRAF6 decrease neuroplastin capacity to promote filopodia. (**A**) Scale bar=10 μm. (**B**) Filopodia number per micron of plasma membrane (GFP=0.15 ± 0.01 N=70; GFP-GPI=0.20 ± 0.01 N=59; Np55-GFP=0.51 ± 0.02 N=51; Np65-GFP=0.48 ± 0.01 N=126; Np65Δ-GFP=0.25 ± 0.01 N=34; Np65$_{PED}$-GFP=0.27 ± 0.02 N=34; Np65$_P$-GFP=0.33 ± 0.01 N=40; siTRAF6 GFP-GPI=0.29 ± 0.01 N=30; siTRAF6 Np55-GFP=0.17 ± 0.01 N=28; siTRAF6 Np65-GFP=0.18 ± 0.01 N=27) and (**C**) Filopodia length (GFP=5.07 ± 0.24; GFP GPI=8.41 ± 0.33; Np55-GFP=16.68 ± 0.76; Np65-

GFP=16.67 ± 0.48; Np65Δ-GFP=9.48 ± 0.57; Np65$_{PED}$-GFP= 10.79 ± 0.58; Np65$_{P}$-GFP=10.78 ± 0.64; siTRAF6 GFP-GPI= 1.73 ± 0.72; siTRAF6 Np55-GFP=10.18 ± 0.72; siTRAF6 Np65-GFP= 9.77 ± 0.85) were quantified using a semi-automatized Matlab-based algorithm. ***p<0.001 for the indicated condition vs. GFP and ##p<0.01 vs. Np65 using Student's t-test.

D-F. Full-length neuroplastin recruits cytosolic TRAF6 to the cell membrane. (**D**) Endogenous TRAF6 is recruited by and co-localizes with Np65-GFP, but not with Np65Δ-GFP nor with GFP-GPI at the plasma membrane. Scale bar=10 µm. The lower confocal pictures are single z plains and on them, a line scan served to quantify the fluorescence distribution of the GFP-tagged proteins and TRAF6 as shown in (**E**). (**F**) Co-localization index (Pearson's coefficient) is displayed for each of the condition as indicated.

G-J. Elimination of RING domain abrogates TRAF6 capacity to mediate neuroplastin-promoted filopodia formation. (**G**) The pictures are single z plains acquired by confocal microscopy. Arrow heads point to fluorescent spots formed by Np55-RFP co-localizing with TRAF6-GFP or with TRAF6$_{cc-c}$-GFP. (**H**) Filopodia number (TRAF6=0.53 ± 0.02 N=54; TRAF6-GFP=0.88 ± 0.04 N=69; TRAF6$_{cc-c}$-GFP=0.17 ± 0.02 N=25) and (**I**) Filopodia length (TRAF6=13.44 ± 0.35; TRAF6GFP=13.89 ± 0.37; TRAF6$_{cc-c}$-GFP= 4.3 ± 0.29) were obtained using a semi-automatized Matlab-based algorithm. **p<0.01 vs. TRAF6 using Student's t-test. (**J**) Line scan analysis of TRAF6-GFP- and Np55-RFP-associated fluorescent signals of single spots. Also, the corresponding co-localization index (Pearson's coefficient) is displayed for each plotting.

Figure 4. Neuroplastin mediates spinogenesis through TRAF6.

A, **B**. The TRAF6 binding site-deficient Np65Δ-GFP does not rescue spinogenesis in *Nptn⁻/⁻* neurons. (**A**) Confocal images of segments of dendrites of *Nptn⁺/⁺* and *Nptn⁻/⁻* neurons transfected with plasmids encoding GFP, Np65-GFP or Np65Δ-GFP at 7 DIV. At 9 DIV, these neurons were fixed and stained with anti-GFP antibodies followed by an Alexa 488-conjugated antibody (green). (**B**) Protrusion densities from two independent cultures were used to obtain the mean ± SEM as indicated (*Nptn⁺/⁺* GFP=4.12 ± 0.18 N=34; *Nptn⁻/⁻* GFP=1.72 ± 0.19 N=27; *Nptn⁻/⁻* Np65-GFP=3.67 ± 0.18 N=33; *Nptn⁻/⁻* Np65Δ-GFP= 1.79 ± 0.16 N=33). ***p<0.001 vs. GFP-filled wild type neurons and ###p<0.001 vs. GFP-filled *Nptn⁻/⁻* neurons using Student's t-test.

C, **D**. Np65Δ-GFP does not foster spinogenesis. (**C**) Dendritic segments of 8 DIV-old rat neurons expressing the indicated proteins upon transfection are shown. (**D**) Protrusion densities from three independent cultures are expressed as the mean ± SEM (GFP=1.72 ± 0.15 N=52, Np65-GFP=3.63 ± 0.11 N=43; Np65Δ-GFP= 1.64 ± 0.16 N=28). ***p<0.001 vs. GFP using Student's t-test.

E-**G**. Expression of Np65Δ-GFP does not promote spinogenesis in 9 DIV-old rat hippocampal neurons. (**E**) Dendritic segments of neurons expressing GFP or Np65Δ-GFP and stained with antibodies against GFP (white) and Shank2 (red clusters) were photographed using confocal microscopy. Scale bar=10 μm. (**F**) Quantification of the protrusion densities and (**G**) the distribution of Shank2-positve and Shank2-negative protrusions from 20-30 neurons per group from three independent cultures (Shank2(+): GFP=0.55 ± 0.06; Np65Δ-GFP=0.62 ± 0.04).

H-**J**. TRAF6 inhibition decreases spinogenesis. (**H**) 7 DIV-old rat neurons transfected with Np65-GFP and treated with the TRAF6 inhibitor SMI 6860766 (SMI TRAF6, 2 μm) for 48 hours were fixed and stained for GFP (white) and Shank2 (red clusters). Scale bar=10 μm. (**I**) Protrusion density (DMSO GFP=3.24 ± 0.118 N=47; SMI TRAF6 GFP=2.22 ± 0.23 N=16; DMSO Np65-GFP=4.59 ± 0.16 N=56; SMI TRAF6 Np65-GFP=3.34 ± 0.16 N=28) and (**J**) Distribution of Shank2-positve and Shank2-negative protrusions from transfected neurons per group from three independent cultures are displayed. *p<0.05 or ***p<0.001 vs. DMSO GFP and ###p<0.001 vs. SMI TRAF6 GFP using Student's t-test (Shank2(+): GFP DMSO=0.58 ± 0.08; SMI TRAF6=0.52 ± 0.19; Np65-GFP DMSO=0.55 ± 0.06; Np65-GFP SMI TRAF6=0.42 ± 0.09).

K-**M**. We calculated (**K**) the area (DMSO GFP=0.105 ± 0.004; DMSO Np65-GFP=0.137 ± 0.004; DMSO Np65Δ-GFP=0.106 ± 0.003; SMI TRAF6 GFP=0.093 ± 0.003; SMI TRAF6 Np65-GFP=0.094 ± 0.006; SMI TRAF6 Np65Δ-GFP=0.098 ± 0.005), (**L**) the fluorescence intensity (DMSO GFP= 134.6 ± 1.4; DMSO Np65-GFP= 139.5 ± 1.9; DMSO Np65Δ-GFP= 138.4 ± 2.1; SMI TRAF6 GFP= 133.0 ± 1.7; SMI TRAF6 Np65-GFP= 134.0 ± 1.8; SMI TRAF6 Np65Δ-GFP= 134.3 ± 1.6), and (**M**) the number of Shank2 clusters per protrusion (DMSO GFP=1.36 ± 0.17; DMSO Np65-GFP=1.92 ± 0.14; DMSO Np65Δ-GFP=1.35 ± 0.14; SMI TRAF6 GFP=1.39 ± 0.13; SMI TRAF6 Np65-GFP=1.28 ± 0.09; SMI TRAF6 Np65Δ-GFP=1.40 ± 0.10) of the experiments in Figure 4H-J. *p<0.05 between Np65-GFP-expressing and GFP-

expressing neurons using Student's t-test. #p<0.05 between the treatments for the same transfection.

N. Neuroplastin requires both its TRAF6 binding site and endogenous TRAF6 activity to promote spinogenic protrusion density. The illustration in the middle shows Np65-GFP-expressing neurons with increased density of Shank2-containing spinogenic protrusions. This phenotype is no longer observed when the TRAF6 binding site is deleted from the Np65 intracellular tail (Np65Δ-GFP, left). Incubation with SMI 6860766 (SMI TRAF6) decreases both the density of protrusions and fraction of protrusions with Shank2 clusters (right).

Figure 5. TRAF6 activity during spinogenesis is necessary for normal synapse density and balanced E/I ratio.

A-**D**. Treatment with SMI 6860766 (SMI TRAF6) from 7 to 9 DIV, but not from 10 to 12 DIV reduces the number of excitatory synapses. (**A**) Representative confocal images of dendritic segments stained with antibodies against synaptic markers (red, postsynaptic Homer; cyan, presynaptic Synapsin-1) at 12 DIV. As indicated, rat hippocampal neurons were previously treated with SMI TRAF6 for 48 hours between days 7 to 9 or 10 to 12. Scale bar=10 µm (**B**) Quantification of the number excitatory synapses per 10 µm of dendritic segment was obtained from SMI TRAF6-treated neurons at days 7 to 9 or 10 to 12 in two independent experiments (control=7.36 ± 0.58 N=19; 7-9=4.84 ± 0.35 N=19; 10-12=7.36 ± 0.45 N=9). **p<0.01 vs. control and ###p<0.001 vs. 10-12 using Student's t-test. (**C**) Quantification of the area (Homer: control=0.172 ± 0.004; 7-9=0.143 ± 0.004; 10-12=0.162 ± 0.006. Synapsin: control=0.236 ± 0.006; 7-9=0.232 ± 0.007; 10-12=0.248 ± 0.003) and (**D**) fluorescence intensity (Homer: control=148.58 ± 0.99; 7-9=147.13 ± 0.99; 10-12=145.58 ± 1.12. Synapsin: control=90.37 ± 0.79; 7-9=92.21 ± 0.84; 10-12=96.77 ± 2.01) of homer- and synapsin-positive puncta. For Homer, **p<0.01 vs. control and ##p<0.01 vs. 10-12 using Student's t-test.

E, **F**. Reduced presynaptic vesicle recycling in SMI 6860766 (SMI TRAF6)-treated 16 DIV-old neurons. (**E**) Dendritic segments of rat neurons, treated with SMI TRAF6 when indicated, tested for uptake of fluorescently labelled antibody against the luminal part of the presynaptic protein synaptotagmin at 16 DIV (see Material and Methods), fixed, stained with an anti-VGAT antibody to discriminate inhibitory presynapses from the excitatory ones. Photomicrographs with a confocal microscope are shown. The heat scale bar indicates uptake levels. Scale bar=10 µm. (**F**) Quantification of presynaptic activities from three independent experiments (Excitatory: control=55.97 ± 1.28 N=130; 7-9=49.58 ± 2.025 N=67; 10-12=58.73 ± 1.568 N=69; Inhibitory: control=66.00 ± 2.63 N=130; 7-9=54.99 ± 3.30 N=67, 10-12=67.11 ± 3.62 N=69). **$p<0.01$ vs. control and ##$p<0.01$ between synapse type using Student's t-test.

DIV						Condition: mean ± SE; N	P value
6	7	8	9	10	11		

rat neurons

6	7	8	9	10	11	Condition: mean ± SE; N	P value
▨	▨	▨				GFP: 2.169 ± 0.371; N=12 Np65-GFP: 3.654 ± 0.343; N=13 Np65$_{DDEP(-)}$-GFP: 3.716 ± 0.293; N=13 Np55-GFP: 3.628 ± 0.267; N=10 Np55$_{DDEP(-)}$-GFP: 3.352 ± 0.185; N=13	p<0.01 p<0.01 p<0.01
	Fig. 1 D, E					GFP: 1.951 ± 0.196; N=39 Np65-GFP: 3.233 ± 0.142; N=56 Np55-GFP: 3.586 ± 0.161; N=38	p<0.001 p<0.001
	Fig. 1 F, G					GFP: 3.151 ± 0.182; N=48 Np65-GFP: 4.642 ± 0.145; N=54	p<0.01
		▨	▨			GFP: 3.334 ± 0.189; N=26 Np65-GFP: 3.985 ± 0.229; N=24	p=0.03
			▨	▨		GFP: 4.225 ± 0.229; N=16 Np65-GFP: 4.551 ± 0.422; N=18	p>0.05
			▨	▨	▨	GFP: 4.991 ± 0.667; N=39 Np65-GFP: 5.081 ± 0.442; N=49 Np55-GFP: 5.122 ± 0.321; N=40	p>0.05

Nptn-/- neurons

6	7	8	9	10	11	Condition: mean ± SE; N	P value
	Fig. 1 A-C					GFP: 1.924 ± 0.222; N=26 Np65-GFP: 3.674 ± 0.182; N=20 Np55-GFP:3.771 ± 0.195; N=26	p<0.01 p<0.001

Table 1 (related to Fig 1). Time course of TRAF6 mediated neuroplastin-promoted spinogenesis. The table is explained in the main text.

Presynapse:	Condition	Area of vesicle release (area of puncta)	P value vs. control	Activity level (intensity per pixel)	P value vs. control
Glutamatergic	control	0.3924 ± 0.009 N=130		138.3 ± 0.56 N=130	
	SMI 7-9	0.3446 ± 0.012 N=67	p<0.001	139.8 ± 0.99 N=67	p>0.05
	SMI 10-12	0.4011 ± 0.009 N=69	p>0.05	140.3 ± 0.87 N=69	p<0.05
Gabaergic	control	0.5520 ± 0.012 N=130		114.7 ± 2.8 N=130	
	SMI 7-9	0.5535 ± 0.019 N=67	p>0.05	100.2 ± 5.062 N=67	p<0.01
	SMI 10-12	0.5574 ± 0.017 N=69	p>0.05	113.8 ± 5.0 N=69	p>0.05

Table 2 (related to Fig 5). Synaptotagmin-1 assay in inhibitory and excitatory presynapses. The table is explained in the main text.

Materials and Methods

Cells. Primary *Nptn*[-/-] neurons were derived from hippocampi of *Nptn*[-/-] mice and compared to primary *Nptn*[+/+] *neurons* derived from their proper control *Nptn*[+/+] mice (Herrera-Molina et al., 2014; Bhattacharya et al., 2017). Cultures of rat hippocampal neurons were obtained as described (Herrera-Molina et al., 2005; 2012). Human embryonic kidney (HEK) 293T cells were cultured as previously described (Herrera-Molina et al., 2017).

DNA constructs and transfections. GFP-tagged neuroplastin constructs have been described (Herrera-Molina et al., 2017). Neuroplastin mutants flanked by HindIII and BamH1 restriction sites were generated from Np65-GFP plasmid by PCR amplification using the following primers for Np65-GFP forward: 5'-TCA AGC TTG CCA CCA TGT CG-3' reverse: 5'- GGC GAT GGA TCC ATT TGT GTT TC-3'; Np65Δ-GFP reverse 5'-GGA TCC TGG CCT CTT CCT CTT CTC ATA C-3': Np65$_{PED}$-GFP forward 5'-GAG GAA GAG GGC AGA TGC GGT TCC TGC TG-3' reverse 5'-CAG CAG GAA CCG CAT CTG CCC TCT TCC TC-3'. The mouse N-terminal Flag-tagged TRAF6 (Flag-TRAF6) mammalian expression plasmid was purchased from Addgene (#21624, GenBank: BAA12705.1). N-terminally GST-tagged TRAF6 (GST-TRAF6) and RING domain deficient TRAF6 with coiled-coil domain and TRAF6-C domain (289-530aa; GST-TRAF6$_{cc-c}$) plasmids with BamH1 and EcoR1 restriction sites were generated by PCR amplification. GST-TRAF6 forward 5'-GAC AGG ATC CTC ATG AGT CTC TTA AAC-3' reverse 5- TAC GAA TTC CTA CAC CCC CGC ATC AGT A-3'; GST-TRAF6$_{cc-c}$ forward 5-GCG TCG GAT CCA TAT GGC CGC CTC T-3'; TRAF6-GFP forward 5'-GTG AAG CTTCTA ATG AGT CTC TTA AAC TGT GA -3' reverse 5'- ATA AGG ATC CCT ACA CCC CCG CAT C -3'; TRAF6$_{cc-c}$-GFP forward 5'- GTG AAG CTT CTA ATG GCC GCC TCT -3'. Scrambled siRNA (sc-37007) and TRAF6 siRNA (sc-36717) were purchased from Santa Cruz. HEK cells were transiently transfected with plasmid DNA constructs using Lipofectamine 2000 (Invitrogen/ThermoFisher) in optiMEM media (Gibco). Primary neuronal cultures were transfected at 7 days *in vitro* (DIV). Control scrambled siRNA or TRAF6 siRNA (30 nM) were transfected using siLentFect (Bio-Rad).

***In silico* modelling.** We performed local peptide docking based on interaction similarity and energy optimization as implemented in the GalaxyPepDock docking tool

(Lee et al., 2015). The protein–peptide complex structure of the hTRANCE-R peptide bound to the TRAF6 protein as provided by Ye et al., 2002 was used as input (PDB: 1LB5). The docking employs constraints of local regions of the TRAF6 surface based on the interaction template. The energy-based optimization algorithm of the docking tool allows efficient sampling of the backbone and side-chains in the conformational space thus dealing with the structural differences between the template and target complexes. Models were sorted according to protein structure similarity, interaction similarity, and estimated accuracy. The fraction of correctly predicted binding site residues and the template-target similarity was used in a linear model to estimate the prediction accuracy. The model using target-template interactions based on the QMPTEDEY motif of the hTRANCE-R template was selected (TM score: 0.991; Interaction similarity score 108.0; Estimated accuracy: 0.868).

Surface plasmon resonance. Protein-Protein interaction measurements were carried out on a BIACORE X100 (GE Healthcare Life Sciences). Sensorgrams were obtained as single cycle kinetics runs. Therefore increasing concentrations of neuroplastin peptide (2.5, 5, 100, 200, 400µM) or just running buffer (startup) were sequentially injected on GST-TRAF6 coated CM5 sensor chip (GE). Unspecific bindings were calculated by using a GST-coated sensor as reference response. Immobilization of these proteins was done using the amine coupling kit as we described in Reddy et al. 2014. All runs were performed in HBS-P buffer. Analysis of affinity was performed using the BIACORE X100 Evaluation Software 2.0.1 (Reddy et al. 2014).

GST pull-down assay. GST, GST-TRAF6 and GST-TRAF6$_{cc-c}$ were transformed into *Escherichia coli* BL21 (DE3) bacterial strain and induced by 0.5 mM of isopropyl-1-thio-b-D-galactopyranoside (IPTG) for 6 h at 25°C. The cells were lysed in resuspension buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl and protease inhibitor cocktail (Roche) with sonication on ice. The purifications of these proteins from transformed bacterial cell extract were performed according to manufacturer instructions (GST bulk kit, GE Healthcare Life Sciences). The purified soluble GST proteins were immobilized on glutathione sepharose 4B beads (GE Healthcare Life Sciences). The beads were washed with binding buffer at least four times, and the pull-down samples were subsequently subjected to immunoblot analyses. The 5 µg of fusion protein coupled beads (GST, GST-TRAF6 and GST-TRAF6$_{cc-c}$) were incubated

with lysate from HEK cells transfected with Np65-GFP for 1 h at 4°C in 500 µl RIPA lysis buffer. The beads were washed and eluted with pre-warmed SDS sample buffer. The eluted complexes were resolved by SDS-PAGE.

Immunoblotting. Proteins were separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) on 10% gels and transferred to a nitrocellulose membrane (Whatman). After blocking with 5% non-fat milk in Tris-buffered saline (TBS) containing 0.1% of Tween 20 for 1 h, the membranes were incubated with indicated antibodies overnight, washed with TBS three times, and then incubated with corresponding secondary antibody conjugated to horseradish peroxidase enzyme for 1 h. Immunodetection was performed with the following antibodies: anti-Flag mouse (Sigma, #F1804; 1:2,000) , anti-GFP rabbit (Abcam, #ab290; 1:2,500), anti-TRAF6 mouse (Santa Cruz, #sc-8709; 1:1,000) and anti-β-actin mouse (Sigma, #A5441; 1:1,000), horseradish peroxidase-conjugated anti-mouse (Dako, #P0447; 1:4,000) or anti-rabbit IgG (gamma-chain specific, Sigma, #A1949-1VL; 1:4,000) antibodies.

Immunocytochemistry. Hippocampal neurons were fixed with 4% PFA for 8 mins and then washed with a solution containing 10% horse serum, 0.1 mM glycine, and 0.1% Triton X-100 in Hanks' balanced salt solution four times for 5 min. Fixed samples were incubated with indicated primary antibodies for overnight at 4°C. To visualize dendritic protrusions, after transfection, pyramidal neurons were morphologically identified based on the side and shape of cell body as observed using anti-MAP2 guinea pig (Synaptic Systems, #188 004; 1:1,000) and anti-GFP mouse (Sigma Aldrich, #11814460001; 1:1,000) antibodies. Routinely, neuron identity was confirmed using an anti-Ctip2 rat (Abcam, #25B6; 1:250) (Herrera-Molina et al., 2014). Subsequently, samples were incubated with anti-guinea pig Cy5-, anti-mouse Alexa 488-conjugated secondary antibodies (1:1,000) generated in donkey (Jackson ImmunoResearch) for 1 hour at RT. Other primary antibodies used were: anti-Synapsin 1 rabbit (Synaptic Systems, #106 103; 1:500), anti-Shank2 guinea pig antibody (Synaptic Systems, #162 204; 1:1,000), anti-Homer1 mouse (Synaptic Systems, #160 011; 1:500), anti-RelA (p65) rabbit (Santa Cruz, #sc-372; 1:500); anti-MAP2 guinea pig (Synaptic Systems, #188 004; 1:1,000) primary antibodies for overnight at 4°C. Subsequently, samples were incubated with anti-rabbit 405-, anti-mouse Cy5-, anti-rat Alexa 488- and/ or anti-guinea pig Cy3-conjugated donkey secondary antibodies (1:1,000) for 1 h. Then

samples were washed and mounted with Mowiol. Quantification of synapse marker signals was performed as in detail described in Herrera-Molina et al., 2014.

Image acquisition and quantification of filopodia/ dendritic protrusions. Images were acquired using HCX APO 63/1.40 NA or 100/1.4NA objectives coupled to a TCS SP5 confocal microscope under sequential scanning mode with 4.0- to 6.0-fold digital magnification. Z-stacks (41.01 x 41.01 x 5 μm physical lengths) were digitalized in a 512 x 512 pixels format file. In HEK cells, filopodia number and length were quantified using a MATLAB-based algorithm, FiloDetect, with some modifications (Nilufar et al., 2013). The algorithm was run for every single image, and the image threshold was adjusted to avoid false filopodia detection and to quantify precise filopodia length and number. The filopodia number per μm was calculated from perimeter of the cell using ImageJ. In neurons, the dendritic protrusions were quantified manually using maximum intensity and Z-projection method of ImageJ software. The dendritic protrusions were considered between 0.25 μm and 20 μm length. Shank clusters were quantified from cropped images using original GFP fluorescent as reference to identify puncta of interest.

Co-immunoprecipitation assay. HEK cells overexpressing GFP-tagged constructs were washed in ice-cold PBS and lysed using radioimmunoprecipitation assay (RIPA) buffer contained 20 mM of Tris (pH 7.5), 100 mM of NaCl, 1 mM EDTA, 10% glycerin, 0.1% SDS, 1% Triton X-100, 1 mM AEBSF, 1 mM sodium orthovanadate, 1 mM sodium molybdate, 1 mM N-Ethylmalemide, 20 mM sodium fluoride, 20 mM glycerol-2-phosphate, 10 mM potassium hydrogen phosphate, 10 mM sodium pyrophosphate and protease inhibitor cocktail (Roche). Samples were incubated with GFP antibody-coupled magnetic beads (μMACS) at 4° C for 4 h. Immunoprecipitated complexes were eluted using μMACS GFP isolation kit (#130-091-125) according to manufacturer's instructions. Eluted complexes were subjected SDS-PAGE.

Synaptotagmin Uptake Assay. Presynaptic activity driven by endogenous network activity was monitored as described before (Herrera-Molina et al., 2014). Hippocampal neurons were washed once with pre-warmed Tyrodes buffer (119 mM NaCl, 2.5 mM KCl, 25 mM HEPES, pH 7.4, 30 mM glucose, 2 mM $MgCl_2$, 2 mM $CaCl_2$) and immediately incubated with an Oyster 550-labeled anti-synaptotagmin-1 rabbit antibody (Synaptic Systems, #105 103C3; 1:500) for 20 min at 37°C. After the antibody

uptake, neurons were washed, fixed, and stained with anti-VGAT guinea pig (Synaptic Systems, #131 004; 1:1,000) and anti-synaptophysin mouse (company, catalog number; 1:1,100) primary antibodies overnight at 4°C. Subsequently, samples were incubated with anti-rabbit Cy3-, anti-guinea pig Cy5- and anti-mouse Alexa 488-conjugated donkey secondary antibodies (1:1,000) for 1 hour. Z-stack images of soma and secondary/tertiary dendrites were acquired using an oil-immersion (HCX APO 63/1.40 NA) objective coupled to a TCS SP5 confocal microscope under sequential scanning mode with a 4.0-fold digital magnification, and digitalized in a 512 x 512 pixels format file (61.51 x 61.51 μm physical lengths). All parameters were rigorously maintained during the image acquisition. For quantification, z-stacks were projected using "sum slices" Z-projection method of Fiji software. We quantified the synaptotagmin-associated fluorescence co-localizing with 1-bit masks derived from VGAT-positive (inhibitory presynapses) or VGAT-negative synaptophysin-positive (excitatory presynapses) puncta using the "image calculator" in the Fiji software. During image processing the original settings of the synaptotagmin channel were carefully maintained as the original. One-bit masks were generated using the analyze particle in the Fiji software for a segmented image of each presynaptic marker (range of particle size $0.15 - 2.25$ μm^2 for inhibitory presynapses and $0.15 - 1.50$ μm^2 for excitatory presynapses).

Statistical analysis. The results are presented as mean $\pm$ SEM (standard error of the mean) accompanied by N number of cells in the figure legends. For statistical analysis, Prism 5 software (GraphPad) was used. Mean between two groups was compared with the Student's t-test. Means of two or more groups were analyzed by one-way ANOVA (analysis of variance) followed by Tukey's multiple comparison test for post hoc comparisons. Normalized data were compared using Mann-Whitney test.

Acknowledgements.

We thank to Dr. Dirk Montag for kindly providing *Nptn*[-/-] mice for breading and to Kathrin Pohlmann for her technical work. S.K.V. thanks to the GRK 1167. A.M. thanks the federal state Saxony-Anhalt and the European Structural and Investment Funds (ESF, 2014-2020), project number ZS/2016/08/80645. R.A.M. thanks the FONDECYT Grant No. 1181260. R.H-M., C.I.S., and E.D.G. thanks to SFB 854 and BMBF 01DN17002.

Author Contributions.

S.K.V. and A.M. conducted most of the experiments. L.J. conducted PMCA experiments. A-C.L. and R.R. made constructs. J.H. and M.K. conducted SPR experiments and data interpretation. R.A.M. conducted in silico modelling. M.N., C.I.S., E.D.G. concept and data interpretation. R.H.M. concept, conducted experiments, data interpretation, wrote manuscript draft. All authors contributed to the manuscript final version.

References

Amuti, S., Y. Tang, et al. (2016). "Neuroplastin 65 mediates cognitive functions via excitatory/inhibitory synapse imbalance and ERK signal pathway." Neurobiol Learn Mem **127**: 72-83.

Armstrong, A. P., M. E. Tometsko, et al. (2002). "A RANK/TRAF6-dependent signal transduction pathway is essential for osteoclast cytoskeletal organization and resorptive function." J Biol Chem **277**(46): 44347-44356.

Beesley, P. W., R. Herrera-Molina, et al. (2014). "The Neuroplastin adhesion molecules: key regulators of neuronal plasticity and synaptic function." J Neurochem **131**(3): 268-283.

Bhattacharya, S., R. Herrera-Molina, et al. (2017). "Genetically Induced Retrograde Amnesia of Associative Memories after Neuroplastin Ablation." Biol Psychiatry **81**(2): 124-135.

Boda, B., A. Dubos, et al. (2010). "Signaling mechanisms regulating synapse formation and function in mental retardation." Curr Opin Neurobiol **20**(4): 519-527.

Boeckers, T. M., M. R. Kreutz, et al. (1999). "Proline-rich synapse-associated protein-1/cortactin binding protein 1 (ProSAP1/CortBP1) is a PDZ-domain protein highly enriched in the postsynaptic density." J Neurosci **19**(15): 6506-6518.

Boersma, M. C., E. C. Dresselhaus, et al. (2011). "A requirement for nuclear factor-kappaB in developmental and plasticity-associated synaptogenesis." J Neurosci **31**(14): 5414-5425.

Bozdagi, O., X. B. Wang, et al. (2010). "Persistence of coordinated long-term potentiation and dendritic spine enlargement at mature hippocampal CA1 synapses requires N-cadherin." J Neurosci **30**(30): 9984-9989.

Buckby, L. E., R. Mummery, et al. (2004). "Comparison of neuroplastin and synaptic marker protein expression in acute and cultured organotypic hippocampal slices from rat." Brain Res Dev Brain Res **150**(1): 1-7.

Carrott, L., M. R. Bowl, et al. (2016). "Absence of Neuroplastin-65 Affects Synaptogenesis in Mouse Inner Hair Cells and Causes Profound Hearing Loss." J Neurosci **36**(1): 222-234.

Cavallaro, U. and E. Dejana (2011). "Adhesion molecule signaling: not always a sticky business." Nat Rev Mol Cell Biol **12**(3): 189-197.

Chatzigeorgiou, A., T. Seijkens, et al. (2014). "Blocking CD40-TRAF6 signaling is a therapeutic target in obesity-associated insulin resistance." Proc Natl Acad Sci U S A **111**(7): 2686-2691

Chubykin AA, Atasoy D, Etherton MR, Brose N, Kavalali ET, Gibson JR, Südhof TC (2007) Activity-dependent validation of excitatory versus inhibitory synapses by neuroligin-1 versus neuroligin-2. *Neuron* **54:** 919-931

Chung, J. Y., Y. C. Park, et al. (2002). "All TRAFs are not created equal: common and distinct molecular mechanisms of TRAF-mediated signal transduction." J Cell Sci **115**(Pt 4): 679-688.

Darnay, B. G., J. Ni, et al. (1999). "Activation of NF-kappaB by RANK requires tumor necrosis factor receptor-associated factor (TRAF) 6 and NF-kappaB-inducing kinase. Identification of a novel TRAF6 interaction motif." J Biol Chem **274**(12): 7724-7731.

Desrivieres, S., A. Lourdusamy, et al. (2015). "Single nucleotide polymorphism in the neuroplastin locus associates with cortical thickness and intellectual ability in adolescents." Mol Psychiatry **20**(2): 263-274.

Dresselhaus, E. C., M. C. H. Boersma, et al. (2018). "Targeting of NF-kappaB to Dendritic Spines Is Required for Synaptic Signaling and Spine Development." J Neurosci **38**(17): 4093-4103.

Empson, R. M., L. E. Buckby, et al. (2006). "The cell adhesion molecule neuroplastin-65 inhibits hippocampal long-term potentiation via a mitogen-activated protein kinase p38-dependent reduction in surface expression of GluR1-containing glutamate receptors." J Neurochem **99**(3): 850-860.

Ferrao, R., J. Li, et al. (2012). "Structural insights into the assembly of large oligomeric signalosomes in the Toll-like receptor-interleukin-1 receptor superfamily." Sci Signal **5**(226):

re3.

Grabrucker, A. M., M. J. Knight, et al. (2011). "Concerted action of zinc and ProSAP/Shank in synaptogenesis and synapse maturation." EMBO J **30**(3): 569-581.

Gong, D., X. Chi, et al. (2018). "Structure of the human plasma membrane Ca(2+)-ATPase 1 in complex with its obligatory subunit neuroplastin." Nat Commun **9**(1): 3623.

Henderson, J. T., J. Georgiou, et al. (2001). "The receptor tyrosine kinase EphB2 regulates NMDA-dependent synaptic function." Neuron **32**(6): 1041-1056.

Herrera-Molina, R., K. Mlinac-Jerkovic, et al. (2017). "Neuroplastin deletion in glutamatergic neurons impairs selective brain functions and calcium regulation: implication for cognitive deterioration." Sci Rep **7**(1): 7273.

Herrera-Molina, R., I. Sarto-Jackson, et al. (2014). "Structure of excitatory synapses and GABAA receptor localization at inhibitory synapses are regulated by neuroplastin-65." J Biol Chem **289**(13): 8973-8988.

Herrera-Molina, R., R. Frischknecht, et al. (2012). "Astrocytic alphaVbeta3 integrin inhibits neurite outgrowth and promotes retraction of neuronal processes by clustering Thy-1." PLoS One **7**(3): e34295.

Herrera-Molina, R. and R. von Bernhardi (2005). "Transforming growth factor-beta 1 produced by hippocampal cells modulates microglial reactivity in culture." Neurobiol Dis **19**(1-2): 229-236.

Ilic, K., K. Mlinac-Jerkovic, et al. (2019). "Hippocampal expression of cell-adhesion glycoprotein neuroplastin is altered in Alzheimer's disease." J Cell Mol Med **23**(2): 1602-1607.

Jang, S. Lee Hyejin Lee, et al. (2017). "Synaptic adhesion molecules and excitatory synaptic transmission." Current Opinion in Neurobiology 45: 45-50.

Jiang, W., M. Wei, et al. (2017). "Identification of Protein Tyrosine Phosphatase Receptor Type O (PTPRO) as a Synaptic Adhesion Molecule that Promotes Synapse Formation." J Neurosci **37**(41): 9828-9843.

Kip, S. N., N. W. Gray, et al. (2006). "Changes in the expression of plasma membrane calcium extrusion systems during the maturation of hippocampal neurons." Hippocampus **16**(1): 20-34.

Kobayashi, N., Y. Kadono, et al. (2001). "Segregation of TRAF6-mediated signaling pathways clarifies its role in osteoclastogenesis." EMBO J **20**(6): 1271-1280.

Korthals, M., K. Langnaese, et al. (2017). "A complex of Neuroplastin and Plasma Membrane Ca(2+) ATPase controls T cell activation." Sci Rep **7**(1): 8358.

Langnaese, K., P. W. Beesley, et al. (1997). "Synaptic membrane glycoproteins gp65 and gp55 are new members of the immunoglobulin superfamily." J Biol Chem **272**(2): 821-827.

Linhoff, M. W., J. Lauren, et al. (2009). "An unbiased expression screen for synaptogenic proteins identifies the LRRTM protein family as synaptic organizers." Neuron **61**(5): 734-749.

Lohmann, C. and T. Bonhoeffer (2008). "A role for local calcium signaling in rapid synaptic partner selection by dendritic filopodia." Neuron **59**(2): 253-260.

Lohmann, C., A. Finski, et al. (2005). "Local calcium transients regulate the spontaneous motility of dendritic filopodia." Nat Neurosci **8**(3): 305-312.

Lomaga, M. A., W. C. Yeh, et al. (1999). "TRAF6 deficiency results in osteopetrosis and defective interleukin-1, CD40, and LPS signaling." Genes Dev **13**(8): 1015-1024.

Ma, Q., H. Ruan, et al. (2017). "Proteasome-independent polyubiquitin linkage regulates synapse scaffolding, efficacy, and plasticity." Proc Natl Acad Sci U S A **114**(41): E8760-E8769.

McAllister, A. K. (2007). "Dynamic aspects of CNS synapse formation." Annu Rev Neurosci **30**: 425-450.

McClelland, A.C., M. Hruska, et al. (2010). "Trans-synaptic EphB2–ephrin–B3 interaction regulates excitatory synapse density by inhibition of postsynaptic MAPK signaling." Proc Natl Acad Sci U S A **107** (19) 8830-8835.

Missler, M., W. Zhang, et al. (2003). "Alpha-neurexins couple Ca2+ channels to synaptic vesicle exocytosis." Nature **423**(6943): 939-948.

Nilufar, S., A. A. Morrow, et al. (2013). "FiloDetect: automatic detection of filopodia from fluorescence microscopy images." BMC Syst Biol **7**: 66.

Okawa, H., L. Della Santina, et al. (2014). "Interplay of cell-autonomous and nonautonomous mechanisms tailors synaptic connectivity of converging axons in vivo." Neuron **82**(1): 125-137.

Owczarek, S., V. Soroka, et al. (2011). "Neuroplastin-65 and a mimetic peptide derived from its homophilic binding site modulate neuritogenesis and neuronal plasticity." J Neurochem **117**(6): 984-994.

Reddy, P. P., V. Raghuram, et al. (2014). "Molecular dynamics of the neuronal EF-hand Ca2+-sensor Caldendrin." PLoS One **9**(7): e103186.

Robbins, E. M., A. J. Krupp, et al. (2010). "SynCAM 1 adhesion dynamically regulates synapse number and impacts plasticity and learning." Neuron **68**(5): 894-906.

Roussignol, G., F. Ango, et al. (2005). "Shank expression is sufficient to induce functional dendritic spine synapses in aspiny neurons." J Neurosci **25**(14): 3560-3570.

Russo, S. J., M. B. Wilkinson, et al.. (2009). "Nuclear factor kappa B signaling regulates neuronal morphology and cocaine reward." J Neurosci 29, 3529-3537.

Ziv, N. and S.J. Smith (1996). "Evidence for a Role of Dendritic Filopodia in Synaptogenesis and Spine Formation." Neuron **17**(1):91-102.

Sarowar, T. and A. M. Grabrucker (2016). "Actin-Dependent Alterations of Dendritic Spine Morphology in Shankopathies." Neural Plast **2016**: 8051861.

Saito, A., Y. Fujikura-Ouchi, et al. (2007). "Association study of putative promoter polymorphisms in the neuroplastin gene and schizophrenia." Neurosci Lett **411**(3): 168-173.

Sakaguchi, M., M. Yamamoto, et al. (2016). "Identification of an S100A8 Receptor Neuroplastin-beta and its Heterodimer Formation with EMMPRIN." J Invest Dermatol **136**(11): 2240-2250.

Sando, R., E. Bushong, et al. (2017). "Assembly of Excitatory Synapses in the Absence of Glutamatergic Neurotransmission." Neuron **94**(2): 312-321 e313.

Schmeisser, M. J., B. Baumann, et al. (2012). "IkappaB kinase/nuclear factor kappaB-dependent insulin-like 76 growth factor 2 (Igf2) expression regulates synapse formation and spine maturation via Igf2 receptor signaling." J Neurosci 32, 5688-5703

Schmidt, N., A. Kollewe, et al. (2017). "Neuroplastin and Basigin Are Essential Auxiliary Subunits of Plasma Membrane Ca(2+)-ATPases and Key Regulators of Ca(2+) Clearance." Neuron **96**(4): 827-838 e829.

Schultheiss, U., S. Puschner, et al. (2001). "TRAF6 is a critical mediator of signal transduction by the viral oncogene latent membrane protein 1." EMBO J **20**(20): 5678-5691.

Sigler, A., W. C. Oh, et al. (2017). "Formation and Maintenance of Functional Spines in the Absence of Presynaptic Glutamate Release." Neuron **94**(2): 304-311 e304.

Smalla, K. H., H. Matthies, et al. (2000). "The synaptic glycoprotein neuroplastin is involved in long-term potentiation at hippocampal CA1 synapses." Proc Natl Acad Sci U S A **97**(8): 4327-4332.

Sorrentino, A., N. Thakur, et al. (2008). "The type I TGF-beta receptor engages TRAF6 to activate TAK1 in a receptor kinase-independent manner." Nat Cell Biol **10**(10): 1199-1207

Sudhof, T. C. (2008). "Neuroligins and neurexins link synaptic function to cognitive disease." Nature **455**(7215): 903-911.

Sudhof, T. C. (2017). "Molecular Neuroscience in the 21(st) Century: A Personal Perspective." Neuron **96**(3): 536-541.

Sylwestrak, E. L. and A. Ghosh (2012). "Elfn1 regulates target-specific release probability at CA1-interneuron synapses." Science **338**(6106): 536-540.

van den Berg, S. M., T. T. Seijkens, et al. (2015). "Blocking CD40-TRAF6 interactions by small-molecule inhibitor 6860766 ameliorates the complications of diet-induced obesity in mice." Int J Obes (Lond) **39**(5): 782-790.

Varoqueaux, F., G. Aramuni, et al. (2006). "Neuroligins determine synapse maturation and function." Neuron **51**(6): 741-754.

Verhage, M., A. S. Maia, et al. (2000). "Synaptic assembly of the brain in the absence of neurotransmitter secretion." <u>Science</u> **287**(5454): 864-869.

Wang, K. Z., N. Wara-Aswapati, et al. (2006). "TRAF6 activation of PI 3-kinase-dependent cytoskeletal changes is cooperative with Ras and is mediated by an interaction with cytoplasmic Src." <u>J Cell Sci</u> **119**(Pt 8): 1579-1591.

Williams, M. E., S. A. Wilke, et al. (2011). "Cadherin-9 regulates synapse-specific differentiation in the developing hippocampus." <u>Neuron</u> **71**(4): 640-655.

Wu, H. (2013). "Higher-order assemblies in a new paradigm of signal transduction." <u>Cell</u> **153**(2): 287-292.

Xie, P. (2013). "TRAF molecules in cell signaling and in human diseases." <u>J Mol Signal</u> **8**(1): 7.

Yamashita, M., K. Fatyol, et al. (2008). "TRAF6 mediates Smad-independent activation of JNK and p38 by TGF-beta." <u>Mol Cell</u> **31**(6): 918-924.

Ye, H., J. R. Arron, et al. (2002). "Distinct molecular mechanism for initiating TRAF6 signaling." <u>Nature</u> **418**(6896): 443-447.

Yin, Q., S. C. Lin, et al. (2009). "E2 interaction and dimerization in the crystal structure of TRAF6." <u>Nat Struct Mol Biol</u> **16**(6): 658-666.

Yoshihara, Y., M. De Roo, et al. (2009). "Dendritic spine formation and stabilization." <u>Curr Opin Neurobiol</u> **19**(2): 146-153.

Zeng, W. Z., N. Grillet, et al. (2016). "Neuroplastin Isoform Np55 Is Expressed in the Stereocilia of Outer Hair Cells and Required for Normal Outer Hair Cell Function." <u>J Neurosci</u> **36**(35): 9201-9216.

Zhang, S. and T. H. Murphy (2004). "Ca2+-independent spine dynamics in cultured hippocampal neurons." <u>Mol and Cell Neurosci</u> **25**(2):334-344.

Zhang, C., J. M. Milunsky, et al. (2009). "A neuroligin-4 missense mutation associated with autism impairs neuroligin-4 folding and endoplasmic reticulum export." <u>J Neurosci</u> **29**(35): 10843-10854.