

1 Genetic correlations across genetically
2 determined and developmentally plastic
3 alternative reproductive tactics
4

5 Jessica K. Abbott^{1,2}, Oscar Rios-Cardenas³ & Molly Morris⁴
6

7 1. Section for Evolutionary Ecology, Department of Biology, Lund University, 10 Sölvegatan 37, 22362
8 Lund, Sweden. Email: Jessica.abbott@biol.lu.se

9 2. Corresponding author

10 3. Red de Biología Evolutiva, Instituto de Ecología, A. C. Xalapa, Veracruz, Mexico

11 4. Department of Biological Sciences, Ohio University, Athens OH, USA
12

13 Running title: Inbreeding limits sexual antagonism

14 Keywords: Intralocus conflict, alternative reproductive tactics, genetic correlation, simulations,
15 plasticity
16

17

18

19 Number of words in main text: 4300
20

21

22 Number of figures: 4
23

24

25 Data archiving: Not applicable
26

27

24 Abstract

25 Alternative reproductive tactics occur when individuals of the same sex have a suite of
26 morphological and/or behavioural traits that allow them to pursue different reproductive strategies.
27 A common pattern is e.g. the existence of “courter” and “sneaker” tactics within males. We have
28 previously argued that alternative reproductive tactics should be subject to genetic conflict over the
29 phenotypic expression of traits, similar to sexual antagonism. In this process, which we called intra-
30 locus tactical conflict, genetically determined tactics experience conflicting selection on a shared
31 phenotypic trait, such as body size, but a positive genetic correlation between tactics in body size
32 prevents either tactic from reaching its optimum. Recently, other authors have attempted to extend
33 this idea to developmentally plastic alternative reproductive tactics, with mixed results. However, it
34 is not clear whether we should expect intra-locus tactical conflict in developmentally plastic tactics
35 or not. We have therefore run a series of simulation models investigating under what conditions we
36 should expect to see positive estimates of the inter-tactical genetic correlation, since a positive
37 genetic correlation is a prerequisite for the existence of intra-locus tactical conflict. We found that
38 for autosomal, X-linked, and Y-linked genetically-determined tactics, estimated inter-tactical genetic
39 correlations were generally high. However, for developmentally plastic tactics, the genetic
40 correlation depends on the properties of the switching threshold between tactics. If it is fixed, then
41 estimated genetic correlations are positive, but if there is genetic variation in the switch-point, then
42 any sign and magnitude of estimated genetic correlation is possible, even for highly heritable traits
43 where the true underlying correlation is perfect. This means that caution should be used when
44 investigating genetic constraints in plastic phenotypes.

45

46 Introduction

47 Alternative reproductive tactics (ARTs) are different ways of achieving reproductive success within a
48 sex, and often involve suites of behavioral, morphological and life history traits. ARTs are expected
49 to evolve when sexual selection is strong and multiple strategies are possible (reviewed in Shuster
50 and Wade 2003). Studies of ARTs across a range of taxa (reviewed in Oliveira et al. 2008) have led to
51 a better understanding of the maintenance of genetic variation given strong sexual selection, as well
52 the adaptive nature of that variation (i.e. the best phenotype for using one tactic is often not the
53 best phenotype for a different tactic). The discontinuous expression of at least one or more traits in
54 either males or females is often the first indication of an ART (e.g. Gross 1996; Brockmann 2001).
55 However, there is a growing appreciation for the role of intralocus tactical conflict (IATC) in
56 constraining the evolution of differences between the ARTs (tactical dimorphism), such that even
57 traits that are more or less continuous within a sex may have more than one underlying optimum
58 (Abbott et al. 2019).

59 Individuals from different ARTs will share many homologous traits, however if the optimal state for
60 these traits differ depending on the ART, this will lead to opposing selection (i.e. tactically disruptive
61 selection). When traits that are genetically correlated across the ARTs are not at their adaptive
62 optimum, tactically disruptive selection can generate intralocus tactical conflict (Morris et al. 2013;
63 Buzatto et al. 2015). Studies of intralocus tactical conflict have the potential to increase our
64 awareness of cases where expressed states are not necessarily optimal due to the evolutionary
65 constraints (e.g. slower growth rates, a lack of behavioural plasticity; Abbott et al. 2019). IATC also
66 has the potential to lead to a better understanding of the role of ecological variation across
67 populations in producing the patterns of divergence between ARTs, as well as provide us with a
68 better understanding of the relationship between ARTs and rapid speciation (Abbott et al. 2019).
69 The criteria for demonstrating intralocus tactical conflict include a positive genetic correlation
70 between the ARTs, detecting different optima for the trait across ARTs, and evidence that the ARTs
71 are not at their optima for the trait (Morris et al. 2013).

72 Inter-tactical genetic correlations measure the extent of similarity between the additive effects of
73 alleles when expressed in different tactics. The ideal method for determining inter-tactical genetic
74 correlations when ARTs are genetically fixed is a multigeneration half-sib breeding design, followed
75 by the statistical decomposition of the genetic variance into its many different components
76 (Falconer and Mackay 1996). Given the prevalence of genetic correlations across the sexes even
77 when sexual dimorphism has evolved (e.g. Harano et al. 2010; Poissant et al. 2010), it can be
78 assumed that these correlations are not temporary or transitional stages, highlighting the
79 importance of their estimation across ARTs as well as across the sexes. By current estimations, most

80 ARTs are developmentally plastic (Oliveira et al. 2008); however, this consensus may change as the
81 studies of the proximal mechanisms behind these ARTs increase.

82 West-Eberhard (1986) described how the loss of an alternative phenotype could play a role in
83 speciation through the release “from constraints of having to accommodate multiple alternatives”
84 (pg 1388) within a shared genome. And yet, the idea that developmental plasticity decouples the
85 development of the alternative morphs, allowing them to evolve independently, is prevalent in the
86 literature (reviewed in Tomkins and Hazel 2007). Therefore, a better understanding of the potential
87 for genetic correlations between ARTs that are both genetically fixed and developmentally plastic is
88 needed to determine the extent to which intralocus tactical conflict may be influencing the evolution
89 of ARTs. There are a few empirical studies that have examined genetic correlations across
90 developmentally plastic ARTs. Considering male traits across two species with tactically dimorphic
91 male ARTs that are environmentally influenced, Pike et al. (2017), detected very weak genetic
92 correlations in one species (earwigs, *Forficula auricularia*), and significant correlations in another
93 species (acarid mites, *Rhizoglyphus echinopus*). The genetic correlations in the acarid mite have been
94 further confirmed through artificial selection experiments (Buzatto et al. 2018).

95 Genetically-determined ARTs can be divided into two types that could potentially differ in their
96 propensity for genetic correlations. First, allelic variation at autosomal loci can influence
97 polymorphisms, as in a marine isopod (Shuster and Wade 1991). Second, genetic polymorphisms
98 may be correlated with the genes influencing sex, as in the swordtail fishes where male ARTs have
99 been linked to genetic variation on the Y-chromosome (Zimmerer and Kallman 1989; Lampert et al.
100 2010). Developmentally plastic ARTs, on the other hand, have been proposed to evolve via
101 environmental thresholds. In this model, the environment influences alternative phenotypes
102 through a genetically-determined threshold (Hazel et al. 1990; Hutchings and Myers 1994), which
103 may be able to respond rapidly to artificial selection (Emlen 1996). Studies of several species have
104 provided evidence for the environmental threshold model of developmentally plastic ARTs: Atlantic
105 salmon, where males can either mature sexually early in life in freshwater or more commonly only
106 after completing a migration at sea (Lepais et al. 2017); horned beetles, where some males develop
107 horns and some do not (Emlen 1996); and bulb mites (Buzatto et al. 2015). In the current study, we
108 use simulations to examine the potential for genetic correlations across ARTs that vary in their
109 underlying mechanisms, including an autosomal ART locus, an X- or Y-linked ART locus, a fixed
110 threshold for plastically determined tactics, and genetic variation in the switching threshold between
111 plastic tactics.

112 Intertactical additive genetic correlations, similar to r_{mf} between the sexes (Lande 1980), are
113 predictive of the potential for future independent evolution of the ARTs within a population. Here
114 we considered the genetic correlation of traits across ARTs that are not directly linked to the locus
115 (or loci) producing the ART. For example, in the case of the autosomal supergene producing
116 differences across male ARTs in ruffs (Küpper et al. 2015), the inversion has linked a suite of traits
117 involved in the ARTs. However, by considering the extent to which these correlations will be present
118 for traits not directly linked to the ARTs, we can determine the extent to which IATC can influence
119 and potentially constrain their evolution. Our results suggest that for autosomal, X-linked, and Y-
120 linked genetically-determined tactics, estimated genetic correlations are, as expected, generally
121 high. However, for developmentally plastic tactics, the estimated genetic correlation depends on the
122 properties of the switching threshold between tactics, such that genetic variation in the switching
123 threshold can lead to a range of estimates even when the true underlying genetic correlation is
124 perfect.

125 Simulation model

126 In the case of genetically-determined tactics, all simulations assume a single ART locus with two
127 alternative alleles producing different alternative reproductive tactics. If tactic is developmentally
128 plastic, then the phenotype determines the threshold for switching between tactics. Different
129 relationships between the phenotype and the threshold are considered (see below). The trait that
130 can experience tactical antagonism is a quantitative trait determined by many loci that are spread
131 throughout the genome (e.g. body size). This means that there is no effect of linkage between the
132 ART locus and the trait.

133 The simulations are designed in such a way that they resemble the simplest form of an animal
134 model:

$$135 \quad y_i = \mu + a_i + e_i$$

136 Where y_i is the phenotypic trait value of individual i , μ is the phenotypic mean in the population, a_i
137 is the breeding value, and e_i is the residual error. Parameters in the simulations are therefore the
138 number of families (fixed at 1000), the population mean (arbitrarily fixed at 50), the population
139 standard deviation (used in randomly generating a breeding value for each family; arbitrarily fixed at
140 5 unless otherwise stated), the within-family standard deviation (used in generating developmental
141 noise for each individual, and equal to the residual standard error), the magnitude of the trait
142 difference between the ARTs (arbitrarily fixed at 5 unless otherwise stated), and the offspring
143 number (arbitrarily fixed at 10 unless otherwise stated). Heritability measured as V_a/V_p is therefore
144 equal to the population SD/(population SD + within-family SD). For developmentally plastic tactics


145 there is an additional parameter for the switching threshold. The inter-tactical correlation was
146 estimated from within-family means, as in Buzatto et al. (2015). For Y-linked tactics, we also included
147 an analysis using MCMCglmm (see below). Datasets were simulated 1000 times each, in order to
148 obtain information about stochastic variation in correlation estimates for a given set of parameter
149 values.

150 We examine several different scenarios, including an autosomal ART locus, an X- or Y-linked ART
151 locus, a fixed threshold for switching tactics, and various ways of relating the switching threshold to
152 the breeding value. Note that in all cases the underlying genetic architecture of the quantitative trait
153 is a perfect genetic correlation between tactics, so what we investigate with our simulations is
154 variation in the empirical estimate of the genetic correlation, and how well it corresponds to the
155 true genetic architecture.

156 Autosomal ART locus

157 This is the simplest possible case, and was simply analysed to provide a baseline for comparison with
158 the other scenarios. We assume that at least some families are capable of producing a mix of
159 different ARTs, for example when two heterozygotes for a dominant ART locus mate with each other
160 and produce offspring of all possible genotypes. We varied the population standard deviation, the
161 within-family standard deviation, offspring number, and the mean trait difference between ARTs,
162 and investigated heritability and the magnitude of the inter-tactical genetic correlation.

163 We found that, as expected, estimated inter-tactical correlations were high for an autosomal ART
164 locus, and increased with increasing heritability (Figure 1). This is a logical result; if the variation
165 within a family is as large as (or larger than) the variation in breeding values across the entire
166 population, it will be more difficult to detect the inter-tactical correlation. Interestingly, we could
167 also show that the estimate of the inter-tactical genetic correlation was often higher than the
168 heritability of the trait (Figure 1, dashed line shows 1:1 relationship), which suggests that the
169 underlying genetic architecture is more important in determining the estimated genetic correlation
170 than the observed heritability. Increasing or decreasing the trait difference between the tactics does
171 not affect these conclusions, but assigning different levels of developmental noise to the two ARTs,
172 or reducing family size, both decrease the estimate of the correlation (Figure 1). This is also an
173 intuitive result since there will be more error in the estimate when the family size is low. The effects
174 of trait difference between tactics and family size is consistent across all other scenarios discussed
175 below (data not shown).



176

177 *Figure 1: Estimated inter-tactical genetic correlation for a quantitative trait dependent on its heritability, in the case of an*
178 *autosomal ART locus. Error bars denote 95% confidence intervals and points are jittered to avoid overlap. Estimates*
179 *increase with increasing heritability. Increasing the magnitude of the difference between ARTs (here by a factor of two)*
180 *does not affect the estimated genetic correlation. However, reducing the offspring number within each family (by 50%*
181 *shown here) and a higher variance in one ART ($SD\ ART1 = 2 * SD\ ART2$ shown here) both decrease the magnitude of the*
182 *estimated genetic correlation. Points falling above the dashed 1:1 line have a higher estimated genetic correlation than the*
183 *heritability of the trait.*


184 Sex-linked ART locus

185 We assume XY sex determination for simplicity, but the results of these simulations should be
186 equally applicable to species with ZW sex determination. In the case of X-linkage, the results are the
187 same as for an autosomal ART locus, as long as it is possible to produce families with a mix of ARTs.
188 For a male-limited X-linked polymorphism, heterozygote female carriers will produce mixed broods,
189 and for a dominant female-limited X-linked polymorphism, mixed broods can be produced either by
190 heterozygote mothers mating males carrying the recessive allele, or by homozygote mothers who

191 mate with males carrying the opposite allele. Given that we assume that the trait and the morph
192 locus are unlinked, results for an X-linked locus are therefore the same as for an autosomal locus
193 since an individual's morph assignment is independent of its breeding value for the trait. In practice,
194 estimates will probably often be lower for X-linked loci compared to autosomal loci because of the
195 difficulty in obtaining large numbers of offspring of each ART from the same family (Figure 1).

196 For a Y-linked locus, it will never be possible to produce full siblings of different ARTs, which means
197 that some sort of maternal half-sib design is necessary for estimating inter-tactical genetic
198 correlations. We therefore assumed a design where each mother is mated to two males, one from
199 each ART. In this case, the family means method of estimation will consistently underestimate the
200 genetic correlation due to the lower level of relatedness between siblings of different ARTs, so we
201 also analysed simulated Y-linked datasets using MCMCglmm (Hadfield 2010). MCMCglmm analyses
202 were run with a modified inverse Wishart prior, and assume that 50% of the variance in the trait is
203 genetic in origin (Mousseau and Roff 1987). The number of iterations was 11000 with a burn-in
204 period of 1000 and thinning interval of 50. This produced a reasonable autocorrelation in the trial
205 runs that were carried out. The effective sample size is rather small with this combination of
206 parameters, but was retained in the interest of saving time. Because MCMCglmm analyses are
207 considerably more time-consuming to run than calculating the correlation based on family means
208 (since each run involves a large number of permutations), the MCMCglmm analyses were carried out
209 on 50 simulated datasets for each parameter combination, instead of 1000, resulting in wider
210 confidence intervals for the MCMCglmm estimated compared to the family means estimates (Figure
211 2).

212 As expected, uncorrected inter-tactical correlations estimated from family means were about half
213 the magnitude for Y-linked tactics compared to autosomal ones, due to the half-sib breeding design.
214 Although accuracy of the estimates can be improved by multiplying the calculated correlation by two
215 to take reduced relatedness into account, precision will still be considerably lower (i.e. high
216 confidence intervals) for Y-linked tactics compared to autosomal tactics. The MCMCglmm analysis
217 was more effective in producing estimates for Y-linked tactics that were similar to those for an
218 autosomal locus, at least for lower heritability values (Figure 2). The decrease in estimates at high
219 heritabilities (>0.6) is likely a side effect of the prior which assumes a heritability of 0.5. This suggests
220 that unless information about the heritability of the quantitative trait is incorporated into the
221 analysis, intertactical genetic correlations may be inaccurately estimated even when using
222 MCMCglmm.



223


224 *Figure 2: Estimated inter-tactical genetic correlation for a quantitative trait dependent on its heritability, in the case of a Y-*
225 *linked ART locus. Baseline values for the autosomal case are included for the sake of comparison. Error bars denote 95%*
226 *confidence intervals and points are jittered to avoid overlap. Raw family means values are approximately half that of the*
227 *autosomal case, and confidence intervals are wider. MCMCglimm performs well when the prior accurately reflects the true*
228 *additive genetic variance (see main text for details). Points falling above the dashed 1:1 line have a higher estimated*
229 *genetic correlation than the heritability of the trait.*

230

231 Developmentally plastic ARTs

232 In the case of developmentally plastic ARTs, which tactic an individual becomes is determined by a
233 phenotypic switching threshold. In many species, this is determined by body size, which is usually
234 heritable, but also tends to reflect overall condition. Following Buzatto *et al.* (2015), we assumed
235 that the switching threshold is genetically fixed in the population and the same as the population
236 mean. We also assumed that the trait that determines the switching threshold is also the one for

237 which we wish to estimate the intertactical genetic correlation. In practice, this means that only
238 families with a breeding value relatively close to the population mean for the quantitative trait will
239 produce offspring that belong to both ARTs, unless environmental manipulations make it possible to
240 influence the developmental trajectory (we chose not to simulate this sort of manipulation for the
241 sake of simplicity). We chose this approach because it is likely to be the least favourable scenario for
242 detecting intertactical genetic correlations; if the trait of interest is not the



243

244 *Figure 3: Estimated inter-tactical genetic correlation for a quantitative trait dependent on its heritability, in the case of*
245 *developmentally plastic ARTs. Baseline values for the autosomal case are included for the sake of comparison. Error bars*
246 *denote 95% confidence intervals and points are jittered to avoid overlap. If the switching threshold is genetically fixed in the*
247 *population, then estimates are lower and more variable overall, but follow a similar pattern as in the autosomal case. If*
248 *there is genetic variation for the switching threshold, then estimates often include zero (indicated by the dashed line) even*
249 *though the true underlying genetic architecture is a perfect correlation.*

250

251 trait that determines the switching threshold, then it should be much easier to obtain mixed
252 families.


253 If we take body size as an example, then in this scenario, all males that are below the population
254 mean body size at the developmental decision point become the “small” ART (i.e. sneaker or minor
255 males) and all males that are above the population mean body size at the developmental decision
256 point become the “large” ART (i.e. courter or major males). We found that the estimated inter-
257 tactical genetic correlation is lower for developmentally plastic tactics compared to an autosomal
258 ART locus, but that the estimates are still always significant as long as the heritability of the
259 quantitative trait is moderately high (>0.3 ; Figure 3, dark blue triangles). The confidence interval for
260 the estimates is also higher, which is likely a result of decreased effective sample size since not all
261 families will produce mixed broods.

262 We predict that under intra-locus tactical conflict, it would be advantageous for developmentally
263 plastic ARTs to evolve the ability to adjust the switching threshold to match the value of the
264 quantitative trait that determines the developmental trajectory. For example, a family with genes
265 for becoming large will tend to produce particularly large “small” males, which may be inefficient at
266 obtaining sneaky matings. Similarly, a family with genes for becoming small will tend to produce
267 particularly small “large” males, which may likewise be inefficient at courting females. A family with
268 genes for low values of the switching threshold trait should therefore adjust the threshold upwards
269 if possible (Figure 4), and vice versa for families with high values of the trait.

270 There are multiple ways in which the switching threshold could be adjusted. We examine two main
271 types of adjustment, including or excluding the effect of developmental noise. It is not our aim here
272 to explore all possible means of adjusting the switching threshold – although there has been much
273 speculation that species can harbour genetic variation in switching threshold, demonstrating that
274 this is not trivial, since it requires raising similar genotypes across a range of environmental factors.
275 This means that data on what sort of switching adjustment patterns may exist is sparse (Taborsky
276 2017). We therefore chose to examine two simple types of adjustment scenarios to see if and how
277 threshold adjustment can alter the estimate of the inter-tactical genetic correlation.

278 If developmental noise is not taken into account when adjusting the switching threshold, then a
279 simple rule is to move the threshold upwards or downwards the same amount as the family
280 deviation from the population mean. For example, a family with a breeding value 0.5 standard
281 deviations above the mean would move its switching threshold to 0.5 standard deviations below the
282 mean, ensuring that only individuals with a trait value that is 1 population standard deviation below
283 the family mean will develop into the “small” tactic (Figure 4). We found that this type of adjustment

284 results in universally low and non-significant estimates of the inter-tactical genetic correlation, since
285 the adjustment cancels out the effect of underlying genetic differences in the quantitative trait
286 (Figure 3, light blue diamonds). Changing the magnitude of the adjustment (e.g. by multiplying the
287 family deviation from the population mean by a fixed constant) alters the slope of the relationship
288 between the estimate and V_a/V_p , but the general conclusions hold (Figure S1).



289
290 *Figure 4: In a family of small individuals, individuals above the switching threshold will develop as “large” (i.e.*
291 *major/courter, shaded area below the curve) males but be at the lower end of the size distribution for this ART (A). It would*
292 *therefore be potentially advantageous for males in this family to adjust their switching threshold upwards (B). Conversely,*
293 *in a family of large individuals, individuals below the switching threshold will develop as “small” (i.e. minor/sneaker, shaded*
294 *area below the curve) males but be at the higher end of the size distribution for this ART (C). It would therefore be*
295 *potentially advantageous for males in this family to adjust their switching threshold downwards (D). Figure partially*
296 *adapted from Buzatto et al. (2015).*

297

298 When developmental noise is not taken into account, as in the scenario above, this means that
299 families with extreme values of the focal trait will not produce mixed broods unless the
300 developmental noise parameter is very large. We therefore decided to explore what would happen if
301 the switching threshold was adjusted relative to both the family breeding value and the degree of
302 developmental noise. In this scenario, the threshold adjustment was scaled according to the
303 developmental noise, such that a family with a breeding value 0.5 standard deviations above the
304 population mean will move its threshold 0.5 within-family standard deviations (i.e. the
305 developmental noise parameter) below the population mean. This means that the magnitude of the
306 adjustment increases as the developmental noise increases, which we feel is a reasonable approach.
307 The greater the uncertainty in offspring phenotype, the more scope for adjustment is needed. We
308 found that this scenario caused estimates of the inter-tactical genetic correlation to fluctuate from
309 negative to positive, depending on the heritability of the trait (Figure 3, purple squares). Again,
310 changing the magnitude of the adjustment (e.g. by multiplying the threshold displacement from the
311 family mean by a fixed constant) moves the location of the inflection point of the relationship
312 between the estimate and V_a/V_p , but the general conclusions hold (Figure S2).

313 Conclusions

314 Genetic correlations across ARTs is an essential criterion for intralocus tactical conflict to be a
315 constraint on the evolution of ARTs. Our results suggest that such conflict is possible for
316 developmentally plastic tactics, which is consistent with the detection of genetic correlations in
317 some recent studies (Pike et al. 2017; Buzatto et al. 2018). In addition, we suggest that different
318 methods are needed to detect genetic correlations across developmentally plastic ARTs, and that
319 even for genetically influenced ARTs, the methods that can be applied will determine the extent to
320 which real genetic correlations can be detected.

321 Inter-tactical genetic correlations are potentially more difficult to measure in
322 developmentally plastic ARTs, unless experimental manipulations of conditions influencing the
323 switch are possible (e.g. rearing soapberry bugs under a wide range of food and conspecific density
324 conditions that influence a wing polyphenism, Fawcett et al. 2018; manipulated pheromones in
325 mites to produce high status scramblers and low status fighters, Michalczyk et al. 2018). Estimates of
326 genetic correlations were also lower for developmentally plastic tactics (at least using the family
327 means method), meaning that non-significant results might be likely even if there is a true
328 correlation. If there is an ability to adjust the switching threshold, all estimates become suspect.
329 Plausible adjustment mechanisms include thresholds that are regulated by transcriptional switches.
330 Gene-expression differences associated with polyphenic morphs have provided evidence for

331 transcriptional switches, e.g. differences in the expression of genes encoding insulin signaling
332 components, which alters the reaction norm for the influence of nutrition on a wing polyphenism in
333 the soapberry bug (Fawcett et al. 2018), and additional studies reviewed by Projecto-Garcia et al.
334 (2017). In the case of male-limited ARTs, the maternal focal trait value could also be a source of
335 information about the family breeding value. This means that phenotypic and maternal effects cues
336 could be combined to decide how to adjust the switching threshold (McNamara et al. 2016). Finally,
337 in cases where the ARTs are Y-linked, half-sib breeding designs are necessary (as mixed families are
338 not possible), and inter-tactical correlations estimated from family means will be more difficult to
339 detect compared to the genetic correlations across autosomal tactics. In this case, experimental
340 tests such as experimental evolution may be the only way to conclusively confirm or deny the
341 existence of an inter-tactical genetic correlation.

342 In summary, substantial inter-tactical genetic correlations and therefore IATC are clearly
343 possible for developmentally plastic tactics. However, the study of these correlations will require
344 further understanding of both the genetic and environmental influences on the underlying
345 thresholds.


346

347 References

348

- 349 Abbott, J. K., O. Rios-Cardenas, and M. R. Morris. 2019. Insights from intralocus tactical conflict:
350 adaptive states, interactions with ecology, and population divergence. *Oikos* In press.
- 351 Brockmann, H. J. 2001. The evolution of alternative strategies and tactics. *Advances in the Study of*
352 *Behavior* 30:1-51.
- 353 Buzatto, B. A., H. L. Clark, and J. L. Tomkins. 2018. Morph-specific artificial selection reveals a
354 constraint on the evolution of polyphenisms. *Proc. R. Soc. Lond. B Biol. Sci* 285:20180335.
- 355 Buzatto, B. A., J. S. Kotiaho, J. L. Tomkins, and L. W. Simmons. 2015. Intralocus tactical conflict:
356 genetic correlations between fighters and sneakers of the dung beetle *Onthophagus taurus*.
357 *J. Evol. Biol* 28:730-738.
- 358 Emlen, D. J. 1996. Artificial selection on horn length-body size allometry in the horned beetle
359 *Onthophagus acuminatus* (Coleoptera: Scarabaeidae). *Evolution* 50:1219-1230.
- 360 Falconer, D. S. and T. F. C. Mackay. 1996. Introduction to quantitative genetics. Pearson Education
361 Limited, Harlow.
- 362 Fawcett, M. M., M. C. Parks, A. E. Tibbetts, J. S. Swart, E. M. Richards, J. C. Vanegas, M. Censer, L.
363 Crowley, W. R. Simmons, W. S. Hou, and D. R. Angelini. 2018. Manipulation of insulin
364 signaling phenocopies evolution of a host-associated polyphenism. *Nature Communications*
365 9:1699.
- 366 Gross, M. R. 1996. Alternative reproductive strategies and tactics: diversity within sexes. *Trends Ecol.*
367 *Evol* 11:92-98.
- 368 Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear models: the MCMCglmm
369 R package. *Journal of Statistical Software* 33:1-22.
- 370 Harano, T., K. Okada, S. Nakayama, T. Miyatake, and D. J. Hosken. 2010. Intralocus sexual conflict
371 unresolved by sex-limited trait expression. *Current Biology* 20:2036-2039.

- 372 Hazel, W., R. Smock, and M. D. Johnson. 1990. A polygenic model for the evolution and maintenance
373 of conditional strategies. *Proc. R. Soc. Lond. B Biol. Sci* 242:181-187.
- 374 Hutchings, J. A. and R. A. Myers. 1994. The evolution of alternative mating strategies in variable
375 environments. *Evol. Ecol* 8:256-268.
- 376 Küpper, C., M. Stocks, J. E. Risse, N. dos Remedios, L. L. Farrell, S. B. McRae, T. C. Morgan, N.
377 Karlinova, P. Pinchuk, Y. I. Verkuil, A. S. Kitaysky, J. C. Wingfield, T. Piersma, K. Zeng, J. Slate,
378 M. Blaxter, D. B. Lank, and T. Burke. 2015. A supergene determines highly divergent male
379 reproductive morphs in the ruff. *Nat. Genet* 48:79-83.
- 380 Lampert, K. P., C. Schmidt, P. Fischer, J.-N. Volff, C. Hoffman, J. Muck, M. Lohse, M. J. Ryan, and M.
381 Scharfl. 2010. Determination of onset of sexual maturation and mating behavior by
382 melanocortin receptor 4 polymorphisms. *Current Biology* 20:1-6.
- 383 Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.
384 *Evolution* 34:292-305.
- 385 Lepais, O., A. Manicki, S. Glise, M. Buoro, and A. Bardonnnet. 2017. Genetic architecture of threshold
386 reaction norms for male alternative reproductive tactics in Atlantic salmon (*Salmo salar* L.).
387 *Scientific Reports* 7:43552.
- 388 McNamara, J. M., S. R. X. Dall, P. Hammerstein, and O. Leimar. 2016. Detection vs. selection:
389 integration of genetic, epigenetic and environmental cues in fluctuating environments.
390 *Ecology Letters* 19:1267-1276.
- 391 Michalczyk, Ł., M. Dudziak, J. Radwan, and J. L. Tomkins. 2018. Fitness consequences of threshold
392 trait expression subject to environmental cues. *Proc. R. Soc. Lond. B Biol. Sci* 285:20180783.
- 393 Morris, M. R., D. Goedert, J. K. Abbott, D. Robinson, and O. Rios-Cardenas. 2013. Intralocus tactical
394 conflict and the evolution of alternative reproductive tactics. *Advances in the Study of*
395 *Behavior* 45:447-478.
- 396 Mousseau, T. A. and D. A. Roff. 1987. Natural selection and the heritability of fitness components.
397 *Heredity* 59:181-197.
- 398 Oliveira, R. F., M. Taborsky, and H. J. Brockmann. 2008. Alternative reproductive tactics: an
399 integrative approach. Cambridge University Press, Cambridge, UK.
- 400 Pike, K. N., J. L. Tomkins, and B. A. Buzatto. 2017. Mixed evidence for the erosion of intertactical
401 genetic correlations through intralocus sexual conflict. *J. Evol. Biol* 30:1195-1204.
- 402 Poissant, J., A. J. Wilson, and D. W. Coltman. 2010. Sex-specific genetic variance and the evolution of
403 sexual dimorphism: a systematic review of cross-sex genetic correlations. *Evolution* 64:97-
404 107.
- 405 Projecto-Garcia, J., J. F. Biddle, and E. J. Ragsdale. 2017. Decoding the architecture and origins of
406 mechanisms for developmental polyphenism. *Current Opinion in Genetics and Development*
407 47:1-8.
- 408 Shuster, S. M. and M. J. Wade. 1991. Equal mating success among male reproductive strategies in a
409 marine isopod. *Nature* 350:606-610.
- 410 Shuster, S. M. and M. J. Wade. 2003. Mating systems and strategies. Princeton University Press,
411 Princeton NJ.
- 412 Taborsky, B. 2017. Developmental plasticity: preparing for life in a complex world. *Advances in the*
413 *Study of Behavior* 49:49-99.
- 414 Tomkins, J. L. and W. Hazel. 2007. The status of the conditional evolutionarily stable strategy. *Trends*
415 *Ecol. Evol* 22:522-528.
- 416 West-Eberhard, M. J. 1986. Alternative adaptations, speciation, and phylogeny (a review). *Proc. Nat.*
417 *Acad. Sci. USA* 83:1388-1392.
- 418 Zimmerer, E. J. and K. D. Kallman. 1989. Genetic basis for alternative reproductive tactics in the
419 pygmy swordtail, *Xiphophorus nigrensis*. *Evolution* 43:1298-1307.
- 420




421

422 *Figure S1: Estimated inter-tactical genetic correlation for a quantitative trait dependent on its heritability, in the case of*
423 *developmentally plastic ARTs where development noise is not taken into account. Baseline values for the fixed switching*
424 *threshold case are included for the sake of comparison. Error bars denote 95% confidence intervals and points are jittered*
425 *to avoid overlap. Multiplying the adjustment value by a fixed constant (here 20% lower or higher, i.e. 0.8 and 1.2) changes*
426 *the slope of the relationship but not the qualitative effect; estimated genetic correlations are still low and usually not*
427 *significantly different from zero.*

428

429



430

431 *Figure S2: Estimated inter-tactical genetic correlation for a quantitative trait dependent on its heritability, in the case of*
432 *developmentally plastic ARTs where development noise is taken into account. Baseline values for the fixed switching*
433 *threshold case are included for the sake of comparison. Error bars denote 95% confidence intervals and points are jittered*
434 *to avoid overlap. Multiplying the adjustment value by a fixed constant (here 20% lower or higher, i.e. 0.8 and 1.2) changes*
435 *the inflection point but not the qualitative effect; estimated genetic correlations are still low and point estimates vary from*
436 *negative to positive depending on the heritability of the trait.*

