When does a minor outbreak become a major epidemic? Linking the risk from

invading pathogens to practical definitions of a major epidemic

AUTHORS

R.N. Thompson[1,2,3,4,*], C.A. Gilligan[4], N.J. Cunniffe[4]

*Correspondence to: robin.thompson@chch.ox.ac.uk

AFFILIATIONS

[1]Mathematical Institute, University of Oxford, Oxford, UK

[2]Department of Zoology, University of Oxford, Oxford, UK

[3]Christ Church, University of Oxford, Oxford, UK

[4]Department of Plant Sciences, University of Cambridge, Cambridge, UK

ABSTRACT

Forecasting whether or not initial reports of disease will be followed by a major epidemic

is an important component of disease management. Most estimates of the probability of

a major epidemic involve assuming that infections occur according to a branching

process. Surprisingly, however, these calculations can be carried out without the factors

that differentiate a major epidemic from a minor outbreak ever being defined precisely.

We assess potential implications of this lack of explicitness by considering the differences

between three practically relevant possible definitions of a major epidemic. Specifically,

we consider a major epidemic as an outbreak in which: i) a large number of hosts are

infected simultaneously; ii) a large number of infections occur in total; and iii) disease

persists in the population for a long time. We calculate the probability of a major epidemic under each of these definitions, initially considering the commonly used stochastic Susceptible-Infected-Susceptible model as a tractable case study allowing analytical progress. We show how the probability of a major epidemic can either be similar to, or very different from, the branching process estimate, depending on which definition of a major epidemic is used. We also show, using other models, that this ambiguity continues to hold in different epidemiological settings. In particular, we consider two additional models: the stochastic Susceptible-Infected-Removed model, and a more complex host-vector model parameterised for Zika virus. Our work highlights how the precise definition of a major epidemic must be considered carefully when estimating the risk posed by a new outbreak. It also demonstrates that the precise definition of a major epidemic must be tailored to the epidemiology of the host-pathogen system under consideration.

KEYWORDS

mathematical modelling; infectious disease epidemiology; major epidemic; forecasting

SHORT TITLE

Assessing the practically relevant risk of a major epidemic

AUTHOR SUMMARY

When cases of disease are first reported in a new region or country, policy-makers must decide if a disease control programme should be rolled out, as well as how to deploy limited resources. Central to this assessment is understanding the risk of initial cases

generating a major epidemic as opposed to fading out as a minor outbreak in which only

few individuals ever become infected. We show how this risk can be evaluated with

practical relevance, by considering consequences for disease control. For example, a

"major epidemic" might be defined as an outbreak likely to require more treatments than

are stockpiled. Once a major epidemic is defined precisely, a practically relevant estimate

for the probability of a major epidemic can be inferred using stochastic epidemiological

models. Estimating the risk posed by invading pathogens in this way contrasts with using

standard out-of-the-box estimates for the probability of a major epidemic, which do not

require a major epidemic to be defined formally and are instead based on restrictive

simplifying assumptions. At the start of any outbreak, careful consideration of precisely

what constitutes a "major epidemic" in that particular setting is vital for accurate

quantification of risk.

1. INTRODUCTION

Infectious disease epidemics in populations of humans, animals and plants represent a

recurring risk worldwide [1–7]. An important question for policy-makers towards the start

of an outbreak is whether initial cases will lead on to a major epidemic, or whether the

pathogen will rapidly die out instead [8,9]. An important practical consequence is that, if

an outbreak is likely to simply fade out, then costly interventions such as vaccination

[10,11], culling/felling [12–18] and workplace or school closure [19] may well be

unnecessary [20].

There is a well-known estimate for the probability of a major epidemic when a pathogen

is newly arrived in a host population, which in its simplest form is given by

$$\text{Prob(major epidemic)} = 1 - \left(\frac{1}{R_0}\right)^{I(0)}, \qquad (1)$$

in which R_0 is the basic reproduction number of the pathogen and $I(0)$ is the number of

individuals that are currently infected. The estimate in equation (1) applies to a wide range

of models, including commonly-used models such as the Susceptible-Infected-

Susceptible (SIS) and Susceptible-Infected-Removed (SIR) models [21]. It can be derived

by assuming that infections occur according to a branching process (see Methods). In

models in which the infected class is sub-divided into different compartments, the value

of $I(0)$ must be interpreted as the total number of individuals infected at $t = 0$. For example,

for the Susceptible-Exposed-Infectious-Removed model, the exponent in equation (1)

would in fact become $E(0) + I(0)$ [9]. More sophisticated estimates that are based on the

branching process approximation can be derived for models including additional

epidemiological detail, such as more complex population structure [22–24] and/or

infectious periods that are not exponentially distributed [25,26].

The quantity in equation (1), and particularly the version in which $I(0) = 1$, is used

extensively in the epidemiological modelling literature [8,9,21,26–35]. It is also used in

real-time during emerging outbreaks. For example, it was used during the 2013-16

epidemic of Ebola virus disease in West Africa to estimate the chance that, if the virus

arrived in Nigeria, sustained transmission would follow in that country [30], and it has

been considered in the context of flare-ups in new locations for the ongoing Ebola

epidemic in the Democratic Republic of the Congo [26].

However, the probability of a major epidemic in equation (1) is derived without specifying

a precise definition of a major epidemic. Over many outbreaks under identical conditions,

if the population size is large and R_0 is much greater than one, then the distribution of

possible epidemic sizes is bimodal according to simple epidemic models such as the

stochastic SIR model (Fig 1d – see also [36–38]): in other words, the final size of any

single outbreak can be in one of two possible ranges. For example, in Fig 1d, most

outbreaks either lead to 1-20 hosts ever infected or to 60-95 hosts ever infected. The

estimate for the chance of a major epidemic in equation (1) corresponds approximately

to the proportion of outbreaks that have a final size in the higher of these ranges. This

provides a natural definition for "minor outbreaks" and "major epidemics", motivated

largely by the elegance of the mathematical modelling analysis. However, for practical

application it would often be more appropriate for the notion of a major epidemic to instead

be grounded in consequences for control of disease, depending on the specific system

and outbreak under consideration.

Here we consider three potential definitions of a major epidemic that might be practically

relevant in different outbreak scenarios. Specifically, these are:

Concurrent size. An outbreak in which the number of individuals simultaneously

infected exceeds the capacity for treatment.

Final size. An outbreak in which the number ever infected exceeds the total number

of available treatments.

Duration. An outbreak that is not contained quickly and therefore persists for an

unacceptably long period.

We investigate under which circumstances the probability of a major epidemic under each

of these definitions matches the branching process estimate from equation (1). In our

analyses, we consider three stochastic epidemiological models that are representative of

different host responses to infection and capture different routes of transmission:

specifically, the SIS model, the SIR model, and a host-vector model parameterised for

Zika virus transmission. For the SIS and SIR models, the branching process estimate

corresponds to equation (1), and in the case of Zika virus the branching process estimate

is an adapted version of equation (1) that accounts for host-vector-host transmission (see

Methods).

The definitions of a major epidemic above might each be applicable in different scenarios.

For example, it might be natural to assume that, if the number of individuals infected

simultaneously remains below the capacity for treatment, then the outbreak can be

controlled and is therefore minor. The threshold capacity might derive from the number

of available hospital beds [39] or the availability of care workers [40]. This motivates the

"Concurrent size" definition above.

However, such a definition cannot be applied ubiquitously. Policy-makers often have to

make decisions concerning how much treatment to stockpile. For example, in response

to growing awareness of the threat of an influenza pandemic, between 2006 and 2013

policy-makers in the UK stockpiled around 40 million units of antivirals at a cost of £424

million, leading to severe criticism when only 2.4 million units were needed; the majority

of which were used during the 2009 H1N1 influenza pandemic [41]. Another possible

definition of a major epidemic is therefore an outbreak in which the total number of

individuals ever infected and requiring treatment exceeds a critical value (the "Final size"

definition, above). This critical value might be set by the stock of available treatments for

use during the outbreak.

Finally, we consider a third definition of a major epidemic (the "Duration" definition). Under

this definition, a major epidemic is an outbreak that persists for an unacceptably long time.

An outbreak that fades out quickly may escape public attention. Even if an outbreak leads

to a significant number of hosts infected, if it ends relatively quickly then it might be

considered minor. For example, the first Ebola outbreak in the Democratic Republic of

the Congo in 2018 resulted in 53 cases, but was not considered a major epidemic due to

its fast containment [42], leading to commendation of the success of public health

measures. Consequently, an outbreak might only be classified as a major epidemic if it

persists for a threshold length of time.

We note that these definitions do not always coincide. In 1665-66, plague affected the

village of Eyam in the UK. The epidemic in the village was long-running, and a large

number of individuals were killed (267 out of 350 in the village), yet model fits suggest

that a maximum of only 30 people were ever infected simultaneously [43,44]. As a result,

this epidemic might have been classified as a major epidemic according to the "Final size"

and "Duration" definitions, yet not under the "Concurrent size" definition, depending on

the precise values of the thresholds set in each case. This highlights the need to define

a major epidemic carefully, since an individual outbreak may or may not qualify as a major

epidemic, depending on the definition used.

We will show that the probability of a major epidemic depends on precisely how a major

epidemic is defined. The probability of a major epidemic under practical definitions may

or may not match the branching process estimate. The definition to use should therefore

be considered carefully before the risk of a major epidemic is assessed at the beginning

of each future outbreak. Only once the notion of a major epidemic has been formally

defined – based on criteria of practical relevance – can this risk be properly assessed.

Figure 1. Schematic diagrams illustrating the population structures for the different models considered, and

an example distribution of final sizes for the stochastic SIR model. (a) The SIS model. (b) The SIR model.

(c) The model of Zika virus transmission. (d) Distribution of final sizes in the stochastic SIR model, with

population size $N = 100$, $R_0 = 1.8$, $I(0) = 1$ and the rest of the population susceptible initially. The y-axis has

been truncated – the height of the leftmost bar is 0.33.

183   **2. METHODS**

We conduct five main analyses. In the first analysis, we calculate the probability of a major

epidemic under the Concurrent size definition for the stochastic SIS model. We then

conduct two further analyses in which we calculate the probability of a major epidemic

under that definition for two other epidemiological models (the SIR model and Zika host-

vector model). We then focus on the stochastic SIS model, and perform two more

analyses in which we calculate the probability of a major epidemic under different

definitions of a major epidemic (the Final size and Duration definitions). In each of these

five analyses, we compare the probability of a major epidemic for the particular

model/major epidemic definition pairing considered to the commonly used branching

process approximation for the probability of a major epidemic which does not require a

major epidemic to be defined formally.

Here, we describe the epidemiological models that we use, calculation of the branching

process approximation to the probability of a major epidemic for each model, and how the

probability of a major epidemic can be calculated under the Concurrent size definition for

each of the models considered. We then explain how the probability of a major epidemic

under each of the practically relevant definitions of a major epidemic can be obtained for

the SIS model.

<u>Epidemiological models</u>

Susceptible-Infected-Susceptible (SIS) model

According to the SIS model, at any time each individual in the population is classified to

be either (*S*)usceptible to or (*I*)nfected by the disease. The deterministic SIS model is

given by

211
$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta IS + \mu I,$$

212
$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta IS - \mu I. \quad (2)$$

214   We use the analogous stochastic model in most of our analyses, in which the net rate at

215   which any epidemiological event occurs is $\beta IS + \mu I$. The probability that this next event is

216   an infection is $\frac{\beta IS}{\beta IS + \mu I}$ and the probability that the next event is a recovery is $\frac{\mu I}{\beta IS + \mu I}$.

218   In this model, the basic reproduction number is given by

219
$$R_0 = \frac{\beta N}{\mu}.$$

221   *Susceptible-Infected-Removed (SIR) model*

222   Under the SIR model, at any time each individual in the population is classified according

223   to whether they are (*S*)usceptible to infection, (*I*)nfected, or (*R*)emoved and no longer

224   spreading the pathogen or available for infection. The deterministic SIR model is given by

226
$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta IS,$$

227
$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta IS - \mu I.$$

228
$$\frac{\mathrm{d}R}{\mathrm{d}t} = \mu I. \quad (3)$$

230   In the analogous stochastic model, the net rate at which any epidemiological event occurs

231   is still $\beta IS + \mu I$, and the probability that the next event is an infection event is similarly

232   unchanged at $\frac{\beta IS}{\beta IS + \mu I}$. However, the other possible next event is a removal, which occurs

233   with probability $\frac{\mu I}{\beta IS + \mu I}$. The basic reproduction number is again

234
$$R_0 = \frac{\beta N}{\mu}.$$

Zika transmission model

We consider the transmission of Zika virus according to the host-vector model of

Kucharski *et al.* [45], which we chose so that we could demonstrate how the probability

of a major epidemic can be calculated in a relatively complex epidemiological setting. In

the model, the numbers of the *N* hosts that are (*S*)usceptible, (*E*)xposed, (*I*)nfectious and

(*R*)emoved are tracked, as well as the vectors that are (S^V)usceptible, (E^V)xposed and

(I^V)nfectious. The deterministic version of this model is given by

244
$$\frac{dS}{dt} = -\beta I^V S,$$

245
$$\frac{dE}{dt} = \beta I^V S - \alpha_H E,$$

246
$$\frac{dI}{dt} = \alpha_H E - \mu I,$$

247
$$\frac{dR}{dt} = \mu I,$$

248
$$\frac{dS^V}{dt} = \delta N^V - \beta_V S^V \frac{I}{N} - \delta S^V.$$

249
$$\frac{dE^V}{dt} = \beta_V S^V \frac{I}{N} - (\delta + \alpha_V) E^V,$$

250
$$\frac{dI^V}{dt} = \alpha_V E^V - \delta I^V. \qquad (4)$$

In the analogous stochastic model, the number of infected human hosts arising from a single infected human (accounting for human-vector-human transmission) in an otherwise entirely susceptible population of humans and vectors is given by

$$R_0^{HV} \times \rho^{E^V \to I^V} \times R_0^{VH} = \frac{\beta_V \alpha_V \beta N^V}{\mu(\delta + \alpha_V)\delta},$$

where $R_0^{HV} = \frac{1}{\mu}\frac{\beta_V N^V}{N}$ is the expected number of vectors infected (and going on to enter the exposed class) by a single infectious human, $\rho^{E^V \to I^V} = \frac{\alpha_V}{\delta + \alpha_V}$ is the proportion of exposed vectors that become infectious and $R_0^{VH} = \frac{1}{\delta}\beta N$ is the expected number of humans infected by a single infectious vector.

The basic reproduction number is given by

$$R_0 = \sqrt{\frac{\beta_V \alpha_V \beta N^V}{\mu(\delta + \alpha_V)\delta}},$$

where the square root accounts for the fact that it takes two generations for infected humans to generate new infections, since new infections require host-vector-host transmission [46,47]. We note that in some studies, e.g. [45], the square root is omitted from the definition of R_0. In contrast to the expression calculated by Kucharski *et al*. [45],

to facilitate simulation of the stochastic model we also explicitly track the total number of

vectors, N^V, rather than the density.

<u>Probability of a major epidemic (branching process estimate)</u>

Standard estimate (for stochastic SIS/SIR models)

The commonly used estimate for the probability of a major epidemic when a pathogen

first arrives in a host population [8,9,21,27,29–35] can be derived by assuming that

infections occur according to a branching process, making the assumptions that the

susceptible population is large and that infection lineages arising from different infected

hosts are independent. When a single infected host arrives in an otherwise susceptible

population, the branching process estimate for the probability of a major epidemic is given

by

285
$$\text{Prob(major epidemic)} \approx \begin{cases} 0 \ \text{ for } R_0 \leq 1, \\ 1 - \dfrac{1}{R_0} \ \text{ for } R_0 > 1. \end{cases}$$

This expression is derived in Text S1.

If instead there are $I(0)$ infected individuals initially rather than one, then for no major

epidemic to occur, it is necessary for each initial infection lineage to die out, leading to

the approximation given in equation (1) whenever $R_0 > 1$.

Extension to more complex models

Here we show how the branching process estimate for the probability of a major epidemic

can be extended to more complex systems by considering the stochastic version of the

model of Zika virus given by the system of equations (4).

We denote the probability of no major epidemic occurring starting from i exposed or

infectious human hosts, j exposed vectors and k infectious vectors by q_{ijk}. We must

consider exposed and infectious vectors separately to account for the possibility that

exposed vectors die before becoming infectious.

Starting from a single infectious host introduced into an entirely susceptible population of

hosts and vectors, and conditioning on the next event, gives

$$q_{100} = \frac{\mu}{\mu + \frac{\beta_V N^V}{N}} q_{000} + \frac{\frac{\beta_V N^V}{N}}{\mu + \frac{\beta_V N^V}{N}} q_{110}.$$

Similarly,

$$q_{010} = \frac{\delta}{\delta + \alpha_V} q_{000} + \frac{\alpha_V}{\delta + \alpha_V} q_{001},$$

$$q_{001} = \frac{\delta}{\delta + \beta N} q_{000} + \frac{\beta N}{\delta + \beta N} q_{101}.$$

We again assume that infection lineages are independent, permitting us to approximate

terms with two exposed or infectious individuals by non-linear terms involving single

exposed or infectious individuals, e.g. $q_{110} \approx q_{100} q_{010}$. Noting that $q_{000} = 1$, then the

three equations above can be solved to give expressions for q_{100}, q_{010} and q_{001}. In

particular, the probability of a major epidemic starting from a single infected host is then

$$1 - q_{100} = \begin{cases} 0 & \text{for } R_0 \leq 1, \\ \dfrac{(R_0)^2 - 1}{(R_0)^2 + R_0^{VH}} & \text{for } R_0 > 1. \end{cases} \quad (5)$$

In this expression, R_0^{VH} is the expected number of humans infected by a single infectious

vector in an otherwise entirely susceptible population of humans and vectors.

<u>Probability of a major epidemic ("Concurrent size" definition)</u>

As described in the introduction, we first define a major epidemic to be an outbreak in

which the number of simultaneously infected individuals ever exceeds a threshold value,

which we denote by *M*. The value of *M* of relevance in practical applications might be set

by the capacity for treatment.

Deterministic SIS and SIR models

In the deterministic SIS model, the maximum number of simultaneously infected

individuals over the course of the outbreak is given by

$$I_{\max} = \begin{cases} N\left(1 - \dfrac{1}{R_0}\right) & \text{for } I(0) \leq N\left(1 - \dfrac{1}{R_0}\right), \\ I(0) & \text{for } I(0) > N\left(1 - \dfrac{1}{R_0}\right), \end{cases}$$

as shown in Text S1.

In the deterministic SIR model, on the other hand,

337
$$I_{\max} = \begin{cases} I^* & \text{for } I(0) \leq I^*, \\ I(0) & \text{for } I(0) > I^*, \end{cases}$$

339 where $I^* = -\frac{N}{R_0} + \frac{N}{R_0} \ln\left(\frac{N}{S(0)R_0}\right) + S(0) + I(0)$, and again this is shown in Text S1.

341 The probability of a major epidemic according to the "Concurrent size" definition under

342 the deterministic SIS and SIR models is then

344
$$\text{Prob(major epidemic)} = \begin{cases} 1 & \text{for } M \leq I_{\max}, \\ 0 & \text{for } M > I_{\max}, \end{cases}$$

where $I_{\max}$ is given by the corresponding expressions above depending which model is

used.

Stochastic SIS model

Under the stochastic SIS model, however, calculating the probability of a major epidemic

is more challenging [48]. Nevertheless, we can calculate this value analytically, which is

advantageous since approximating this quantity using model simulations can be time

consuming given that outbreaks under the SIS model can persist for long periods.

We denote the probability of no major epidemic starting with I infected individuals as q_I,

and assume that the rest of the population is susceptible. Conditioning on whether the

first event is an infection or recovery event gives

359
$$q_I = \frac{\beta IS}{\beta IS + \mu I} q_{I+1} + \frac{\mu I}{\beta IS + \mu I} q_{I-1},$$

361 for $I = 1,2,\ldots,M$-1. We solve this tridiagonal system of equations with boundary conditions

362 $q_0 = 1$ and $q_M = 0$, since a major epidemic certainly does not occur if there are initially no

363 infected individuals, and certainly does if there are initially M infected individuals. Noting

364 that $S = N - I$, these equations can be put in the form

366
$$q_{I+1} - q_I = C_I(q_I - q_{I-1}),$$

368 where

369
$$C_I = \frac{\mu I}{\beta I(N-I)}.$$

371 Iterating this gives

372
$$q_{I+1} - q_I = A_I(q_1 - q_0), \qquad (6)$$

374 where

375
$$A_I = \prod_{j=1}^{I} C_j,$$

376
$$= \frac{\left(\frac{N}{R_0}\right)^I}{I! \binom{N-1}{I}}.$$

378 Adding equation (6) to itself for $I = 1,2,\ldots,M - 1$ and rearranging gives

380
$$q_1 = \frac{\sum_{I=1}^{M-1} A_I}{1 + \sum_{I=1}^{M-1} A_I}.$$

382 Instead adding equation (6) to itself for $I = 1,2,\ldots,k - 1$ gives

384
$$q_k = \frac{\sum_{I=k}^{M-1} A_I}{1 + \sum_{I=1}^{M-1} A_I}.$$

386 The probability of a major epidemic is then given by

388
$$\text{Prob(major epidemic)} = 1 - q_k = \begin{cases} 0 \text{ for } k = 0, \\ \dfrac{1}{1 + \sum_{I=1}^{M-1} A_I} \text{ for } k = 1, \\ \dfrac{1 + \sum_{I=1}^{k-1} A_I}{1 + \sum_{I=1}^{M-1} A_I} \text{ for } 1 < k < M, \\ 1 \text{ for } k \geq M. \end{cases} \quad (7)$$

391 *Stochastic SIR model*

392 Under the stochastic SIR model, the probability of a major epidemic according to the

393 "Concurrent size" definition starting from any state (I,R) is calculated using an iterative

394   approach. Denoting the probability of a major epidemic starting from state (I,R) by $p_{I,R}$,

395   then conditioning on what happens next gives

397
$$p_{I,R} = \frac{\beta I(N - I - R)}{\beta I(N - I - R) + \mu I} p_{I+1,R} + \frac{\mu I}{\beta I(N - I - R) + \mu I} p_{I-1,R}$$

This system can be solved with boundary conditions $p_{0,R} = 0$, $p_{I,N-M+1} = 0$ and $p_{M,R} = 1$.

To do this, the probability of a major epidemic is deduced for the following states (in

order): (I,R) = (M-1,N-M), (M-2,N-M), …, (1,N-M), (M-1,N-M-1), …,(1,N-M-1),… (M-

1,0),…,(1,0). For a schematic showing the order in which these probabilities are deduced,

see Fig S1.

Zika transmission model

The probability of a major epidemic under the "Concurrent size" definition is approximated

by simulating the model 10,000 times using the Gillespie direct method [49], and then

calculating the proportion of simulations in which the number of infected human hosts

exceeds M at any point during the simulation.

Probability of a major epidemic ("Final size" and "Duration" definitions)

We also consider the probability of a major epidemic according to the deterministic and

stochastic SIS model for the "Final size" and "Duration" definitions of a major epidemic.

Under the "Final size" definition, a major epidemic is assumed to be an outbreak in which

at least *F* infections occur over the course of the outbreak. Under the "Duration" definition,

a major epidemic is defined to be an outbreak that persists for at least *T* days.

In the deterministic SIS model, whenever $R_0 > 1$, the outbreak persists indefinitely with

an infinite number of infection events. As a result, any outbreak is a major epidemic under

the "Final size" and "Duration" definitions.

In contrast, in any simulation of the stochastic SIS model, the number of infected

individuals will always reach zero, even if this takes a long time. As a result, the probability

of a major epidemic under the "Final size" and "Duration" definitions is not simply one or

zero depending on the value of R_0. We approximate the probability of a major epidemic

under these definitions by simulating the model 10,000 times using the Gillespie direct

method [49], and recording the proportion of simulations in which there are at least *F*

infections or that have a duration of at least *T* days.

<div align="center">**3. RESULTS**</div>

To begin to understand outbreak dynamics under the SIS, SIR and Zika transmission

models, we first numerically solved the deterministic models given by the systems of

equations (2), (3) and (4) with $R_0 = 1.5$ in each case (Fig 2). In a deterministic setting, the

SIS model predicts the largest number of individuals simultaneously infected as well as

the most infections in total. Epidemics persisted forever (i.e. *I* remained larger than zero)

under all three models, although the number of infected hosts tended to zero under the

SIR and Zika transmission models.

However, our main focus is the probability of a major epidemic soon after the pathogen

enters the system. For any definition of "major epidemic", according to a deterministic

model the corresponding probability is either zero or one depending on the values of

model parameters. We therefore considered the more realistic stochastic models, in

which demographic stochasticity is included. In the following sections, first we calculate

the probability of a major epidemic for the stochastic SIS model under the Concurrent

size definition. We then consider different epidemiological models, as well as different

definitions of a major epidemic. In each case, the probability of a major epidemic for the

particular epidemiological model-definition of a major epidemic pair under consideration

is compared to the branching process approximation to the probability of a major epidemic

for that model. Results are shown in Figs 3-5, as well as summarised in Tables 1 and 2.

Figure 2. Numerical solutions of the deterministic SIS, SIR and Zika virus transmission models when the

basic reproduction number is $R_0 = 1.5$. (a) Number of infected individuals through time according to the

deterministic SIS model. (b) Cumulative number of infections through time according to the deterministic

SIS model. (c)-(d) Equivalent to a-b but for the deterministic SIR model. (e)-(f) Equivalent to a-b but for the

deterministic Zika virus transmission model. Parameters for deterministic SIS and SIR models: $N = 1{,}000$,

$\beta = 0.00015$ per day, $1/\mu = 10$ days. Parameters for deterministic Zika virus transmission model: $N =$

$1{,}000$, $N^V = 10{,}000$, $1/\alpha_V = 10.5$ days, $1/\alpha_H = 5.9$ days, $1/\mu = 5$ days, $1/\delta = 7.8$ days, $\beta = 0.0000615$ per

day, $\beta_V = 0.22$ per day.

<u>The probability of a major epidemic</u>

We calculated the probability of a major epidemic according to the stochastic SIS model

under the "Concurrent size" definition of a major epidemic – i.e. major epidemics defined

as outbreaks in which a threshold number of simultaneously infected individuals is

exceeded. In this case, as described in Methods, it is possible to calculate the probability

of a major epidemic analytically.

We show the probability of a major epidemic for a range of values of the major epidemic

threshold, M, in Fig 3a. For R_0 much larger than one, we found that the probability of a

major epidemic was approximated closely by the standard branching process estimate

for many values of the major epidemic threshold, M. When, however, R_0 was close to

unity, the standard estimate corresponded to a single choice of M (see e.g. blue lines in

Fig 3a, where the solid line is close to the corresponding dotted line in only one place, i.e.

for a single value of M). The parameter regime in which R_0 is close to one is important in

many real epidemiological systems since the aim of eradicative control strategies is

usually to reduce the reproduction number below one [50,51].

Figure 3. Probability of a major epidemic under the SIS model, where a major epidemic is defined as an

outbreak in which M simultaneously infected individuals is exceeded ("Concurrent size" definition). (a)

Dependence on R_0. Solid lines represent the true probability of a major epidemic (system of equations (7)),

dotted lines represent the branching process estimate (equation (1)), and dots show the maximum number

simultaneously infected in the analogous deterministic models (for values of M below this, the probability

of a major epidemic in the relevant deterministic model is 1). R_0 is varied by changing the value of β. (b)

Equivalent to a, but showing dependence on the population size, N. (c) Equivalent to a, but showing

dependence on the initial number of infected individuals, $I(0)$. (d). Single simulation of the stochastic SIS

model (blue), and numerical solution of deterministic SIS model (red dotted). The value of I in the stochastic

simulation will continue to fluctuate about the deterministic value until I reaches 0. Parameter values

(except where stated): N = 1,000, R_0 = 1.5, $I(0)$ = 1 and $R(0)$ = 0 and the remainder of the population

susceptible. In panel d, $\beta = 0.000015$ per day and $1/\mu = 10$ days.

In large host populations, the probability of a major epidemic as a function of M took the

form of a step function (Fig 3b). This suggested that, if the pathogen successfully invaded

the population, then the number of infected individuals would definitely reach a specific

maximum value which is determined by R_0. For example, for outbreaks with R_0 = 1.5, the

pathogen will invade the population with probability 0.33, and, if this occurs, then 67% of

the population will be infected simultaneously at some time during the epidemic.

The maximum value of I that will be reached in the stochastic SIS model in the large N

case is approximately

506
$$\max(I) = \max\left(2N\left(1 - \frac{1}{R_0}\right), N\right).$$

The expression $2N\left(1 - \frac{1}{R_0}\right)$ is twice the maximum value of I in the corresponding

deterministic model. This is because, if the pathogen invades in the stochastic SIS model,

then I will fluctuate approximately symmetrically around the deterministic equilibrium

value (Fig 3d). It is likely, then, that by the time the pathogen dies out by reaching I = 0,

the number of infected individuals will at some stage also have reached approximately

double the value it fluctuated around too.

We also note that, if the expression above for max(I) is reformulated to give the maximum

proportion of the population that is simultaneously infected, the resulting expression is

independent of the population size N. This can also be seen graphically – in Fig 3b, if the

pathogen successfully invaded the population then the maximum proportion of hosts that

were simultaneously infected was approximately independent of the size of the host

population, so long as N was sufficiently large.

<u>Different epidemiological models</u>

We considered the probability of a major epidemic ("Concurrent size" defintion) under the

SIR and Zika virus transmission models. For the stochastic SIR model, we used an

iterative method to calculate the probability of a major epidemic as described in Methods.

For the stochastic Zika virus transmission model, we simulated the model in a population

of N = 1,000 human hosts and N^V = 10,000 vectors using the Gillespie direct algorithm

[49], using parameter values from Kucharski $et\ al.$, 2016 [45] – see caption of Fig 4. The

value of R_0 was then varied in Fig 4b by altering the parameter β that governs the rate at

which infected vectors infect susceptible human hosts.

Figure 4. Probability of a major epidemic under the "Concurrent size" definition of a major epidemic, for the

SIR and Zika virus transmission models. (a) SIR model. Solid lines represent the true probability of a major

epidemic calculated using the iterative method described in Methods, dotted lines represent the branching

process estimate (equation (1)), and dots show the maximum number simultaneously infected in the

analogous deterministic models (for values of M below this, the probability of a major epidemic in the

relevant deterministic model is 1). R_0 is varied by changing the value of β. (b) Equivalent to a, but for the

Zika virus transmission model (where M refers to the number of simultaneously infected hosts). For the

Zika transmission model, the true probability of a major epidemic is calculated by simulation, and the

branching process estimate is given by equation (5). For both models, $N = 1{,}000$. Other parameters for the

Zika virus transmission model: $N^V = 10{,}000$, $1/\alpha_V = 10.5$ days, $1/\alpha_H = 5.9$ days, $1/\mu = 5$ days, $1/\delta = 7.8$

days, $\beta_V = 0.22$ per day. Initial conditions for both models comprise of a single infected host, with all other

individuals (for the Zika transmission model, hosts and vectors) susceptible.

Under the stochastic SIR and Zika models, for R_0 larger than and not close to one, the

maximum number of simultaneously infected individuals whenever the pathogen invaded

the host population was typically smaller than under the SIS model (*cf.* Fig 2).

Nonetheless, we found qualitatively similar behaviour in these cases – the probability of

a major epidemic approximated using a branching process corresponded to a wide range

of values of the major epidemic threshold when R_0 was high (Fig 4). However, even if that

is the case, the practically relevant value of the major epidemic threshold (e.g. the number

of treatments that have been stockpiled) may not give a probability of a major epidemic

that matches the branching process estimate. For example, if R_0 = 2 and 250 treatments

were to have been stockpiled, for the SIR model the probability of a major epidemic under

the "Final size" definition is 0 (solid grey line in Fig 4a), yet the branching process estimate

for the probability of a major epidemic is 0.5 (dotted grey line in Fig 4a).

<u>Alternative definitions of a major epidemic</u>

For the stochastic SIS model, we then calculated the probability of a major epidemic for

different definitions of a major epidemic – specifically, outbreaks in which there are at

least F infection events (the "Final size" definition – Fig 5a) or outbreaks that persist for

at least T days (the "Duration" definition – Fig 5b).

Figure 5. Probability of a major epidemic under the SIS model, for alternative definitions of a major epidemic.

(a) A major epidemic is defined as an outbreak in which more than F infections occur ("Final size" definition).

(b) A major epidemic is defined as an outbreak that persists for more than T days ("Duration" definition).

Solid lines represent the true probability of a major epidemic assessed via simulation of the stochastic model, and dotted lines represent the branching process estimate (equation (1)). Results of the deterministic model are not included in the figure, since under the deterministic SIS model epidemics persist indefinitely and generate an infinite number of infections whenever $R_0 > 1$. Parameter values: N = 1,000, $I(0)$ = 1 and $R(0)$ = 0 and the remainder of the population susceptible. In both panels, R_0 is varied by changing the value of β. In panel b, $1/\mu = 10$ days.

In the stochastic SIS model, if the pathogen invaded the host population then it tended to persist for long periods. Consequently, the branching process estimate corresponded to a very wide range of major epidemic thresholds under the "Final size" or "Duration" definitions (i.e. values of F or T) compared to under the "Concurrent size" definition. As a result, in these specific cases (i.e. when the stochastic SIS model was used and a major epidemic was defined according to the "Final size" or "Duration" definitions) it can be

588 concluded that the branching process approximation often leads to sensible estimates of

589 the risk posed by invading pathogens. Nonetheless, even in these cases, for very large

590 values of the major epidemic thresholds the probability of a major epidemic does not

591 match the branching process estimate, particularly when R_0 was larger than but close to

592 one (see e.g. red line in Fig 5b).

	SIS model	SIR model	Zika host-vector model
R_0 = 1.1	M in range [21,54] (Fig 3a)	M in range [18,21] (Fig 4a)	M in range [6,7] (Fig 4b)
R_0 = 1.6	M in range [8,681] (Fig 3a)	M in range [8,55] (Fig 4a)	M in range [3,47] (Fig 4b)
R_0 = 2	M in range [6,866] (Fig 3a)	M in range [7,117] (Fig 4a)	M in range [2,83] (Fig 4b)

594
Table 1. Effect of underlying epidemiology. For which values of the threshold (number of hosts
simultaneously infected, M) in the Concurrent size definition of a major epidemic is the branching process
approximation to the probability of a major epidemic accurate? Threshold values here are those for which
the branching process approximation is within 0.01 of the probability of a major epidemic. Results are
summarised for the SIS, SIR and Zika virus models, for the parameter values shown in the relevant figure
captions.
601
602
603

	Concurrent size	Final size	Duration
R_0 = 1.1	M in range [21,54] (Fig 3a)	F in range [138,878] (Fig 5a)	T in range [183,458] (Fig 5b)
R_0 = 1.6	M in range [8,681] (Fig 3a)	F greater than 18 (Fig 5a)	T greater than 51 (Fig 5b)
R_0 = 2	M in range [6,866] (Fig 3a)	F greater than 10 (Fig 5a)	T greater than 34 (Fig 5b)

604
Table 2. Effect of definition of a major epidemic. For which values of the thresholds in the more practically
relevant definitions of a major epidemic is the branching process approximation to the probability of a
major epidemic accurate? Threshold values here are those for which the branching process
approximation is within 0.01 of the probability of a major epidemic. Results are summarised for the SIS

model, for the parameter values shown in the relevant figure captions. Final size values were tested up to
a maximum of $F = 2,000$ (Fig 5a) and duration values were tested up to $T = 6,000$ (Fig 5b).

4. DISCUSSION

Determining whether or not an emerging outbreak is likely to develop into a major

epidemic is useful for planning intervention and containment strategies. When a pathogen

arrives in a new location, the probability of a major epidemic can be derived by assuming

that infections occur according to a branching process. For simple models such as the

stochastic SIS and SIR models, this leads to the probability of a major epidemic in

equation (1). It is also possible to calculate the probability of a major epidemic according

to the branching process approximation using models with other epidemiological

complexity, as we showed by considering the case of host-vector transmission (see

equations (4) and (5)).

However, the branching process estimate for the probability of a major epidemic is not

necessarily accurate when a definition of a major epidemic is used that addresses

practical aspects of disease control (Fig 3). The branching process estimate corresponds

to a range of choices of the epidemic thresholds in our definitions when R_0 is much greater

than one, or when the population size is extremely large (see e.g. different values of M in

Fig 3a,b). However, when R_0 is close to one and the population is not large, the standard

estimate can correspond to a single choice of the epidemic threshold (see e.g. blue and

red lines in Fig 3a and Tables 1 and 2). For specific outbreaks, even when both R_0 and

the population size are large, the branching process estimate may not be relevant – since

the range of choices of the major epidemic threshold that the standard estimate

corresponds to may or may not include the specific threshold of practical importance in

the outbreak, for example the number of hospital beds available. Consequently, using the

standard branching process estimate for the probability of a major epidemic could lead to

the risk of a major epidemic being incorrectly assessed, potentially including scenarios in

which a major epidemic develops when previously deemed unlikely. Our main conclusion

that the branching process estimate for the probability of a major epidemic may or may

not match the true probability of a major epidemic when a practically relevant definition is

used holds for a range of epidemiological systems (Fig 4) as well as different definitions

of a major epidemic that apply in alternative settings (Fig 5). We note that exactly how

well the branching process estimate reflects a range of threshold values depends on the

underlying model as well as the relevant quantity differentiating major epidemics from

minor outbreaks (*cf*. Figs 3a and 5).

We considered practical definitions of a major epidemic that were based on thresholds

such as the availability of treatment. A previous study has concluded that epidemiological

modellers should report the precise cutoff used to define a major epidemic in model

simulations [52]. We support this conclusion, and indeed some authors have done this –

for example, Keeling *et al*. [53] define a major epidemic to be an outbreak in which at

least one-third of the population becomes infected. However, we contend furthermore that

the precise threshold used should be motivated by the value that is most practically

relevant in the particular system under consideration.

Under the first definition of a major epidemic that we considered (the "Concurrent size"

definition), the probability of a major epidemic was assessed in the context of the capacity

for treatment by estimating whether or not a threshold number of simultaneously infected

individuals was likely to be exceeded. This definition may be practically relevant in a range

of scenarios. For example, real-time analysis of a diphtheria epidemic in Cox Bazar's in

Bangladesh involved assessing the number of hospital beds that were needed [39]. The

number of hospital beds that were already available might have provided a practically

relevant major epidemic threshold. Another example is citrus greening disease in Brazil,

for which a law was introduced that stated that a citrus grove must be destroyed if 28%

of trees in the grove were infected and symptomatic [54,55]. This fraction of trees could

correspond to a threshold in a representative model similar to the thresholds that we

considered here. Other examples for which interventions are introduced as soon as a

threshold in the number simultaneously infected is reached include the development of

the National Chlamydia Screening Programme in the United Kingdom in 2002 in response

to the large size of the infected population [56].

However, no single definition of a major epidemic will be relevant in all situations. We also

considered two other definitions of a major epidemic. In one of these (the "Final size"

definition), whether or not an outbreak was classified as a major epidemic referred to the

total number of infection events over the course of the outbreak, rather than the maximum

number simultaneously infected. This might correspond to the total number of individuals

ever requiring treatment, which may be an important threshold if treatments have been

stockpiled prior to the outbreak [41]. This definition might also be relevant if, for example,

a policy-maker has to choose how to deploy resources between two different epidemics.

If there are only sufficient resources to contain one outbreak, then it might be preferable

to choose the one that is likely to generate more infections. In other real-world scenarios,

other definitions might be appropriate. We also considered major epidemics defined as

outbreaks that persist for a threshold length of time (the "Duration" definition). Different

definitions of a major epidemic might appear contradictory – for example, treatment can

act to reduce the total number of infections yet increase the duration of the outbreak [57],

making a major epidemic less likely under the "Final size" definition of a major epidemic

but more likely under the "Duration" definition. In fact, integrating interventions into our

models to examine their effects on the probability of a major epidemic under different

practical definitions is a candidate for further study.

Our intention here was to use very simple models to demonstrate the principle that

different definitions of a major epidemic lead to different probabilities of a major epidemic

each of which might be practically relevant in different scenarios. Although simple models

are commonly used, accurate outbreak forecasts require a model carefully matched to

the epidemiology of the host-pathogen system, potentially including asymptomatic

transmission [9,58], spread between spatially distinct regions [29], or explicit modelling of

control interventions [12,20,59–61]. For certain definitions, it may be necessary to include

convalescent hosts in the model explicitly. For example, if convalescing individuals take

up resources, such as beds in treatment rooms or hospitals, and the definition of a major

epidemic is linked to the availability of resources (as in the case of the "Concurrent size"

definition), then these individuals should be modelled, potentially by including them in a

new compartment following the infectious class. More complex definitions of major

epidemics could also be used, for example requiring multiple criteria to be satisfied for an

outbreak to be classified as a major epidemic. In these more complicated scenarios,

analytic calculations of the probability of a major epidemic might not be possible. Model

simulations can then be used to assess the probability of a major epidemic, as we showed

using a host-vector model of Zika virus disease transmission (Fig 4b).

In summary, we have shown that how precisely to define a major epidemic should be

considered carefully in future studies. The definition of a major epidemic should be

designed to match the questions of interest in the particular setting being considered.

Standard approximations based on branching processes should not always be used as

the default. Only once a "major epidemic" has been defined precisely can the probability

of a major epidemic occurring be properly assessed. Providing an explicit demonstration

of the consequences of not considering a practically relevant definition in evaluating the

risk of an epidemic is the key contribution of this paper.

SUPPLEMENTARY MATERIAL

S1 Text.

S1 Figure. Schematic showing how the probability of a major epidemic under the

"Concurrent size" definition can be deduced for the stochastic SIR model. The probability

corresponding to each state (I,R) is deduced iteratively in the order 1,2,3... (red). The black arrows indicate which previous values are used to inform each new deduction.

COMPETING INTERESTS

We have no competing interests.

AUTHORS' CONTRIBUTIONS

All authors designed the study. RNT undertook the research and wrote the first draft of the manuscript. All authors revised the manuscript.

FUNDING

This research was funded by BBSRC via a G2O PhD studentship (RNT), and by Christ Church, Oxford, via a Junior Research Fellowship (RNT).

REFERENCES

1. Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife — threats to biodiversity and human health. Science (80-). 2000;287: 443–450.

2. Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P. Emerging infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers. Trends Ecol Evol. 2004;19: 535–544. doi:10.1016/j.tree.2004.07.021

3. Taylor LH, Latham SM, Woolhouse MEJ. Risk factors for human disease

emergence. Philos Trans R Soc B. 2001;356: 983–989.

doi:10.1098/rstb.2001.0888

4. Morens DM, Folkers GK, Fauci AS. The challenge of emergin and re-emerging

infectious diseases. Nature. 2004;430: 242–249.

5. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global

trends in emerging infectious diseases. Nature. 2008; 990–994.

doi:10.1038/nature06536

6. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, et al.

Emerging fungal threats to animal, plant and ecosystem health. Nature. 2012;484:

186–194. doi:10.1038/nature10947

7. Thompson RN, Brooks-Pollock E. Detection, forecasting and control of infectious

disease epidemics: modelling outbreaks in humans, animals and plants. Philos

Trans R Soc B. 2019;374: 20190038.

8. Craft ME, Beyer HL, Haydon DT. Estimating the probability of a major outbreak

from the timing of early cases: an indeterminate problem ? PLoS One. 2013;8:

e57878. doi:10.1371/journal.pone.0057878

9. Thompson RN, Gilligan CA, Cunniffe NJ. Detecting presymptomatic infection is

necessary to forecast major epidemics in the earliest stages of infectious disease

outbreaks. PLoS Comput Biol. 2016;12: e1004836.

doi:10.1371/journal.pcbi.1004836

10. Ramsay ME. Measles: the legacy of low vaccine coverage. Arch Dis Child.

2013;98: 752–754. doi:10.1136/archdischild-2013-304292

11. Rohani P, Earn DJD, Grenfell BT. Impact of immunisation on pertussis

771  transmission in England and Wales. Lancet. 2000;355: 285–286.

772 12. Ferguson NM, Donnelly CA, Anderson RM. The foot-and-mouth epidemic in Great

773  Britain: Pattern of spread and impact of interventions. Science (80-). 2001;292:

774  1155–1160.

775 13. Kao RR. The role of mathematical modelling in the control of the 2001 FMD

776  epidemic in the UK. Trends Microbiol. 2002;10: 279–286.

777 14. Keeling MJ. Models of foot-and-mouth disease. Proc R Soc B Biol Sci. 2005;272:

778  1195–1202.

779 15. Keeling MJ, Woolhouse MEJ, Shaw DJ, Matthews L, Chase-Topping M, Haydon

780  DT, et al. Dynamics of the 2001 UK foot and mouth epidemic: Stochastic

781  dispersal in a heterogeneous landscape. Science (80-). 2001;294: 813–818.

782 16. Pautasso M, Aas G, Queloz V, Holdenrieder O. European ash (Fraxinus

783  excelsior) dieback – A conservation biology challenge. Biol Conserv. Elsevier Ltd;

784  2013;158: 37–49. doi:10.1016/j.biocon.2012.08.026

785 17. Cunniffe NJ, Laranjeira FF, Neri FM, DeSimone RE, Gilligan CA. Cost-effective

786  control of plant disease when epidemiological knowledge is incomplete: Modelling

787  Bahia bark scaling of citrus. PLoS Comput Biol. 2014;10: e1003753.

788  doi:10.1371/journal.pcbi.1003753

789 18. Chan M-S, Jeger MJ. An analytical model of plant virus disease dynamics with

790  roguing and replanting. J Appl Ecol. 1994;31: 413–427.

791 19. Wu JT, Cowling BJ, Lau EHY, Ip DKM, Ho L-M, Tsang T, et al. School closure

792  and mitigation of pandemic (H1N1) 2009, Hong Kong. Emerg Infect Dis. 2010;16:

793  538. doi:10.3201/eid1603.091216

794   20.   Thompson RN, Gilligan CA, Cunniffe NJ. Control fast or control smart: When

795   should invading pathogens be controlled? PLoS Comput Biol. 2018;14.

796   doi:10.1371/journal.pcbi.1006014

797   21.   Keeling MJ, Rohani P. Modeling infectious diseases in humans and animals.

798   Princeton University Press; 2008.

799   22.   Lloyd AL, Zhang J, Root AM. Stochasticity and heterogeneity in host-vector

800   models. J R Soc Interface. 2007;4: 851–863. doi:10.1098/rsif.2007.1064

801   23.   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect

802   of individual variation on disease emergence. Nature. 2005;438: 355–359.

803   doi:10.1038/nature04153

804   24.   Nishiura H, Cook AR, Cowling BJ. Assortativity and the probability of epidemic

805   extinction: a case study of pandemic influenza A (H1N1-2009). Interdiscip

806   Perspect Infect Dis. 2011; 194507.

807   25.   Anderson D, Watson R. On the spread of a disease with gamma distributed latent

808   and infectious periods. Biometrika. 1980;67: 191–198.

809   26.   Thompson RN, Jalava K, Obolski U. Sustained transmission of Ebola in new

810   locations: more likely than previously thought. Lancet Infect Dis. 2019;

811   doi:10.1016/S1473- 3099(19)30360-3

812   27.   Swinton J. Extinction times and phase transitions for spatially structured closed

813   epidemics. Bull Math Biol. 1998;60: 215–230.

814   28.   Merler S, Ajelli M, Fumanelli L, Parlamento S, Pastore y Piontti A, Dean NE, et al.

815   Containing Ebola at the Source with Ring Vaccination. PLoS Negl Trop Dis.

816   2016;10: 1–11. doi:10.1371/journal.pntd.0005093

29.   Thompson RN, Thompson C, Pelerman O, Gupta S, Obolski U. Increased

frequency of travel in the presence of cross-immunity may act to decrease the

chance of a global pandemic. Philos Trans R Soc B. 2019;374: 20180274.

doi:10.1098/rstb.2018.0274

30.   Althaus CL, Low N, Musa EO, Shuaib F, Gsteiger. Ebola virus disease outbreak

in Nigeria: Transmission dynamics and rapid control. Epidemics. 2015;11: 80–84.

31.   Abdullah N, Kelly JT, Graham SC, Birch J, Gonçalves-Carneiro D, Mitchell T, et

al. Structure-guided identification of a nonhuman morbillivirus with zoonotic

potential. J Virol. 2018;92: e01248-18.

32.   Keeling MJ, Danon L. Mathematical modelling of infectious diseases. Br Med Bull.

2009;92: 33–42. doi:10.1093/bmb/ldp038

33.   Keeling MJ, Ross J V. On methods for studying stochastic disease dynamics. J R

Soc Inferface. 2008;5: 171–181. doi:10.1098/rsif.2007.1106

34.   Kessler DA, Shnerb NM. Solution of an infection model near threshold. Phys Rev

E. 2007;76: 010901. doi:10.1103/PhysRevE.76.010901

35.   Park AW, Gubbins S, Gilligan CA. Invasion and persistence of plant parasites in a

spatially structured host population. Oikos. 2001;94: 162–174.

36.   Kendall DG. Deterministic and stochastic epidemics in closed populations. Proc

3rd Berkeley Symp Math Stat Prob. 1956;4: 149–165.

37.   de Jong MCM. Mathematical modelling in veterinary epidemiology: why model

building is important. Prev Vet Med. 1995;25: 183–193.

38.   Nasell I. 'The threshold concept in stochastic epidemic and endemic models' in D.

Mollison, Epidemic models: their structure and relation to data. Cambridge

University Press; 1995.

39. Finger F, Funk S, White K, Siddiqui MR, Edmunds WJ, Kucharski AJ. Real-time

analysis of the diphtheria outbreak in forcibly displaced Myanmar nationals in

Bangladesh. BMC Med. 2019;17: 58. doi:10.1038/d41586-018-05810-w

40. Lamontagne F, Clément C, Fletcher T, Jacob ST, Fischer II WA, Fowler RA.

Doing today's work superbly well - treating Ebola with current tools. N Engl J Med.

2014;371: 1565–1566.

41. O'Dowd A. Government says it would stockpile Tamiflu again. Br Med J.

2014;349: g6386. doi:10.1136/bmj.g6386

42. Butler D. Africa's latest Ebola outbreak comes to swift end. Nature. 2018; 2018–

2020.

43. Brauer F. "Compartmental models in epidemiology" in F. Brauer, P. van den

Driessche, J. Wu - Mathematical Epidemiology. Springer; 2014. doi:10.1007/978-

3-540-78911-6

44. Raggett GF. A stochastic model of the Eyam plague. J Appl Stat. 1982;9: 212–

225. doi:10.1080/02664768200000021

45. Kucharski AJ, Funk S, Eggo RM, Mallet H-P, Edmunds WJ, Nilles EJ.

Transmission Dynamics of Zika Virus in Island Populations : A Modelling Analysis

of the 2013 – 14 French Polynesia Outbreak. PLoS Negl Trop Dis. 2016;10:

e0004726. doi:10.1371/journal.pntd.0004726

46. van den Driessche P, Watmough J. Further notes on the basic reproduction

number. In "Mathematical Epidemiology." Springer; 2008.

47. Heffernan JM, Smith RJ, Wahl LM. Perspectives on the basic reproductive ratio. J

R Soc Inferface. 2005;2: 281–293. doi:10.1098/rsif.2005.0042

48. Artalejo JR, Economou A, Lopez-Herrero MJ. The maximum number of infected

individuals in SIS epidemic models: Computational techniques and quasi-

stationary distributions. J Comput Appl Math. 2010;233: 2563–2574.

doi:10.1016/j.cam.2009.11.003

49. Gillespie DT. Exact stochastic simulation of coupled chemical reactions. J Phys

Chem. 1977;8: 2340–2361.

50. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to

estimate time-varying reproduction numbers during epidemics. Am J Epidemiol.

2013;178: 1505–1512. doi:10.1093/aje/kwt133

51. Thompson RN, Stockwin JE, Gaalen RD Van, Polonsky JA, Kamvar ZN, Demarsh

PA, et al. Improved inference of time-varying reproduction numbers during

infectious disease outbreaks. Epidemics. 2019;19: 100356.

doi:10.1016/j.epidem.2019.100356

52. Orbann C, Sattenspiel L, Miller E, Dimka J. Defining epidemics in computer

simulation models: How do definitions influence conclusions? Epidemics. Elsevier

B.V.; 2017;19: 24–32. doi:10.1016/j.epidem.2016.12.001

53. Keeling MJ. The implications of network structure for epidemic dynamics. Theor

Popul Biol. 2005;67: 1–8. doi:10.1016/j.tpb.2004.08.002

54. Craig AP, Cunniffe NJ, Parry M, Laranjeira FF, Gilligan CA. Grower and regulator

conflict in management of the citrus disease Huanglongbing in Brazil: A modelling

study. J Appl Ecol. 2017;55: 1956–1965. doi:10.1111/1365-2664.13122

55. Belasque Jr. J, Bassanezi RB, Yamamoto PT, Ayres AJ, Tachibana A, Violante

AR, et al. Letter to the editor: Lessons from Huanglongbing management in Sao

Paulo state, Brazil. J Plant Pathol. 2010;92: 285–302.

56. LaMontagne DS, Fenton KA, Randall S, Anderson S, Carter P. Establishing the

National Chlamydia Screening Programme in England: results from the first full

year of screening. Sex Transm Infect. 2004;80: 335–341.

57. Hollingsworth TD, Klinkenberg D, Heesterbeek H, Anderson RM. Mitigation

strategies for pandemic influenza A: Balancing conflicting policy objectives. PLoS

Comput Biol. 2011;7: e1001076. doi:10.1371/journal.pcbi.1001076

58. Hyatt-Twynam SR, Parnell S, Stutt ROJH, Gottwald TR, Gilligan CA, Cunniffe NJ.

Risk-based management of invading plant disease. New Phytol. 2017;214: 1317–

1329.

59. Thompson RN, Hart WS. Effect of confusing symptoms and infectiousness on

forecasting and control of Ebola outbreaks. Clin Infect Dis. 2018;67.

doi:10.1093/cid/ciy248

60. Thompson RN, Cobb RC, Gilligan CA, Cunniffe NJ. Management of invading

pathogens should be informed by epidemiology rather than administrative

boundaries. Ecol Modell. 2016;324. doi:10.1016/j.ecolmodel.2015.12.014

61. Cunniffe NJ, Cobb RC, Meentemeyer RK, Rizzo DM, Gilligan CA. Modeling when,

where, and how to manage a forest epidemic, motivated by sudden oak death in

California. Proc Natl Acad Sci USA. 2016;113: 5640–5645.

doi:10.1073/pnas.1602153113

907