When does a minor outbreak become a major epidemic? Linking the risk from

invading pathogens to practical definitions of a major epidemic

AUTHORS

R.N. Thompson[1,2,3,4,*], C.A. Gilligan[4], N.J. Cunniffe[4]

*Correspondence to: robin.thompson@chch.ox.ac.uk

AFFILIATIONS

[1]Mathematical Institute, University of Oxford, Oxford, UK

[2]Department of Zoology, University of Oxford, Oxford, UK

[3]Christ Church, University of Oxford, Oxford, UK

[4]Department of Plant Sciences, University of Cambridge, Cambridge, UK

ABSTRACT

Forecasting whether or not initial reports of disease will be followed by a major epidemic

is an important component of disease management, guiding optimal deployment of limited

resources for surveillance and control. For example, the probability that undetected cases

arriving in different countries would lead to a major epidemic was estimated during the

2014-16 Ebola epidemic in West Africa, and in the ongoing epidemic in DR Congo.

Standard epidemic risk estimates involve assuming that infections occur according to a

branching process. Surprisingly, however, these calculations are carried out without the

factors differentiating major epidemics from minor outbreaks being defined precisely. We

assess implications of this lack of explicitness by considering three practically relevant

potential definitions of a major epidemic; namely, an outbreak in which: i) a large number

of hosts are infected simultaneously; ii) a large number of infections occur in total; and iii)

disease remains in the population for a long period. We show that the major epidemic

probability under these definitions can be similar to, or different from, the standard

estimate. This holds in a range of systems, highlighting that careful consideration of what

constitutes a "major epidemic" in each outbreak is vital for accurate quantification of risk.

KEYWORDS

mathematical modelling; infectious disease epidemiology; major epidemic; forecasting

SHORT TITLE

Assessing the practically relevant risk of a major epidemic

38   **1.   INTRODUCTION**

Infectious disease epidemics in populations of humans, animals and plants represent a

recurring risk worldwide [1–7]. An important question for policy-makers towards the start

of an outbreak is whether initial cases will lead on to a major epidemic, or whether the

pathogen will rapidly die out instead [8,9]. An important practical consequence is that, if

an outbreak is likely to simply fade out, then costly interventions such as vaccination

[10,11], culling/felling/roguing of plants or agricultural animals [12–18] and workplace or

school closure [19] may well be unnecessary [20].

There is a well-known estimate for the probability of a major epidemic when a pathogen is newly arrived in a host population, which in its simplest form is given by

$$\text{Prob(major epidemic)} = 1 - \left(\frac{1}{R_0}\right)^{I(0)}, \qquad (1)$$

in which R_0 is the basic reproduction number of the pathogen and $I(0)$ is the number of individuals that are currently infected. The estimate in equation (1) applies to a wide range of models, including the commonly used Susceptible-Infected-Susceptible (SIS) and Susceptible-Infected-Removed (SIR) models [21]. It can be derived by assuming that infections occur according to a branching process (see Methods). In some models in which the infected class is sub-divided into different compartments, the value of $I(0)$ must instead be interpreted as the total number of individuals infected at the current time. For example, for the Susceptible-Exposed-Infectious-Removed model, the exponent in equation (1) would in fact become $E(0) + I(0)$ [9]. More sophisticated estimates that are based on similar branching process approximations can be derived for models including additional epidemiological detail, such as more complex population structure [22–24] and/or infectious periods that are not exponentially distributed [25,26].

The quantity in equation (1), and particularly the version in which $I(0) = 1$, is used extensively in the epidemiological modelling literature [8,9,21,26–35]. It is also increasingly used in real-time during emerging outbreaks. For example, it was used during the 2014-16 epidemic of Ebola virus disease in West Africa to estimate the chance

that, if the virus arrived in Nigeria, sustained transmission would follow in that country

[30]. It was also considered in the context of flare-ups in new locations for the ongoing

Ebola epidemic in the Democratic Republic of the Congo [26].

However, the probability of a major epidemic in equation (1) is derived without specifying

a precise definition of a major epidemic. Over many outbreaks under identical conditions,

if the population size is large and R_0 is much greater than one, then the distribution of

possible epidemic sizes is bimodal according to simple epidemic models such as the

stochastic SIR model (Fig 1d – see also [36–38]). In other words, the final size of any

single outbreak is almost always in one of two possible ranges. For example, in Fig 1d,

virtually all outbreaks either lead to 1-20 hosts ever infected or to 700-860 hosts ever

infected, where the precise ranges depend on the population size and the value of R_0.

The estimate for the probability of a major epidemic in equation (1) corresponds

approximately to the proportion of outbreaks that have a final size in the higher of these

ranges. This provides a natural definition for "minor outbreaks" and "major epidemics",

motivated largely by the elegance of the mathematical modelling analysis. However, for

practical application it would often be more appropriate for the notion of a major epidemic

to instead be grounded in consequences for control of disease, depending on the specific

system and outbreak under consideration.

Here we consider three potential definitions of a major epidemic that might be practically

relevant in different outbreak scenarios. Specifically, these are:

Concurrent size. An outbreak in which the number of individuals simultaneously

infected exceeds the capacity for treatment.

Total infections. An outbreak in which the total number of infections exceeds the

number of available treatments.

Duration. An outbreak that is not contained quickly and therefore persists for an

unacceptably long period.

We investigate under which circumstances the probability of a major epidemic according

to each of these definitions coincides with the branching process estimate. In our

analyses, as examples we consider three stochastic epidemiological models that are

representative of different host responses to infection and capture different routes of

transmission. Specifically, we consider the SIS model, the SIR model, and a host-vector

model parameterised for Zika virus transmission. For the SIS and SIR models, the

standard branching process estimate corresponds to equation (1), and in the case of Zika

virus the standard branching process estimate is an adapted version of equation (1) that

accounts for host-vector-host transmission (see Methods).

To motivate our analyses, we note that the three definitions of a major epidemic above

might each be applicable in different scenarios. For example, it might be natural to

assume that, if the number of individuals infected at any time always remains below the

capacity for treatment, then the outbreak is minor since medical care is available for all

sick individuals. The threshold capacity might derive from the number of available hospital

beds [39] or the availability of care workers [40]. This motivates the "Concurrent size"

definition above.

However, a definition based on numbers of simultaneously infected hosts cannot be

applied ubiquitously. Policy-makers often have to make decisions concerning how much

treatment to stockpile; if all cases must be treated, this corresponds to the total number

of infections over the course of the outbreak. For example, in response to growing

awareness of the threat of a potential influenza pandemic, between 2006 and 2013 policy-

makers in the UK stockpiled around 40 million units of antivirals at a cost of £424 million.

This led to severe criticism when only 2.4 million units were needed; the majority of which

were used during the 2009 H1N1 influenza pandemic [41]. Another possible definition of

a major epidemic is therefore an outbreak in which the total number of infections exceeds

a critical value (the "Total infections" definition, above). This critical value might be set by

the stock of available treatments for use during the outbreak.

Finally, we consider a third definition of a major epidemic (the "Duration" definition). Under

this definition, a major epidemic is an outbreak that persists for an unacceptably long time.

An outbreak that fades out quickly may escape public attention. Even if an outbreak leads

to a significant number of hosts infected, if it ends relatively quickly then it might be

considered minor. For example, the first Ebola outbreak in the Democratic Republic of

the Congo in 2018 resulted in 53 cases, but was not considered a major epidemic due to

its fast containment [42], leading to commendation of the success of public health

measures. Consequently, an outbreak might only be classified as a major epidemic if it

persists for a threshold length of time.

It is important to recognise that these definitions do not always coincide. In 1665-66,

plague affected the village of Eyam in the UK, which famously isolated itself via a self-

imposed quarantine [43,44]. The outbreak in the village was long-running, and a large

number of individuals were killed (most reports suggest 250-260 out of a total of 350 in

the village died, although there is some ambiguity particularly regarding the size of the at-

risk population [45]). However, model fits suggest that a maximum of only around 30

people were ever infected simultaneously [46–48]. As a result, this epidemic might have

been classified as a major epidemic according to the "Total infections" and "Duration"

definitions, yet not under the "Concurrent size" definition, depending on the precise values

of the thresholds set in each case. This highlights the need to define a major epidemic

carefully, since an individual outbreak may or may not qualify as a major epidemic,

depending on the definition used.

We will show that the probability of a major epidemic depends on precisely how a major

epidemic is defined. The probability of a major epidemic under practical definitions may

or may not match the branching process estimate. The definition to use should therefore

be considered carefully before the major epidemic risk is estimated at the beginning of an

emerging outbreak. Only once the notion of a major epidemic has been formally defined

– based on criteria of practical relevance – can this risk be properly assessed.

Figure 1. Schematic diagrams illustrating the population structures for the different models considered, and

an example distribution of final sizes for the stochastic SIR model. (a) The SIS model. (b) The SIR model.

(c) The model of Zika virus transmission. (d) Distribution of final sizes in the stochastic SIR model, with

population size $N = 1000$, $R_0 = 2$, $I(0) = 1$ and the rest of the population susceptible initially. The x-axis has

been split into bars of width 20 (so that, for example, the first bar corresponds to the probability that between

1 and 20 individuals are ever infected).

169   **2. METHODS**

We conduct five analyses. In the first three analyses, we calculate the probability of a

major epidemic under the "Concurrent size" definition for the stochastic SIS model, the

SIR model and the Zika host-vector model. Our final two analyses focus on the stochastic

SIS model. We calculate the probability of a major epidemic under other definitions of a

major epidemic (the "Total infections" and "Duration" definitions). In each of these five

analyses, we compare the probability of a major epidemic for the particular model-major

epidemic definition pairing considered to the commonly used branching process

approximation for the probability of a major epidemic which does not require a major

epidemic to be defined formally.

Here, we describe the epidemiological models that we use, calculation of the branching

process approximation to the probability of a major epidemic for each model, and how the

probability of a major epidemic can be calculated under the "Concurrent size" definition

for each of the models considered. We then explain how the probability of a major

epidemic under the other practically relevant definitions of a major epidemic can be

obtained for the SIS model, although our methodology generalises immediately to any

model for which a method of stochastic simulation is available.

Epidemiological models

Susceptible-Infected-Susceptible (SIS) model

According to the SIS model, at any time each individual in the population is classified to

be either (*S*)usceptible to or (*I*)nfected by the pathogen. The deterministic SIS model is

given by

196
$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta IS + \mu I,$$

197
$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta IS - \mu I, \quad (2)$$

where β represents the rate of infection between each susceptible-infected pair, and μ is

the rate at which each infected host recovers and becomes susceptible again. We use

the analogous stochastic model in most of our analyses, in which the net rate at which

any epidemiological event occurs is $\beta IS + \mu I$. At any time prior to the end of the outbreak,

the probability that this next event is an infection is $\frac{\beta IS}{\beta IS + \mu I}$ and the probability that the next

event is a recovery is $\frac{\mu I}{\beta IS + \mu I}$.

In this model, if the total population size is N, the basic reproduction number is given by

206
$$R_0 = \frac{\beta N}{\mu}.$$

208   *Susceptible-Infected-Removed (SIR) model*

209   Under the SIR model, at any time each individual in the population is classified according

210   to whether they are (*S*)usceptible to infection, (*I*)nfected, or (*R*)emoved and no longer

211   spreading the pathogen or available for infection. The deterministic SIR model is given by

213
$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta IS,$$

214
$$\frac{\mathrm{d}I}{\mathrm{d}t} = \beta IS - \mu I,$$

215
$$\frac{\mathrm{d}R}{\mathrm{d}t} = \mu I, \quad (3)$$

216 in which β is again a parameter governing the infection rate, and μ is the rate of

217 removal. In the analogous stochastic model, the net rate at which any epidemiological

218 event occurs is still $\beta IS + \mu I$, and the probability that the next event is an infection event

219 is similarly unchanged at $\frac{\beta IS}{\beta IS + \mu I}$. However, the other possible next event is a removal,

220 which occurs with probability $\frac{\mu I}{\beta IS + \mu I}$. The basic reproduction number is again

221
$$R_0 = \frac{\beta N}{\mu}.$$

Zika transmission model

We consider the transmission of Zika virus according to the host-vector model of

Kucharski *et al.* [49], which we chose to demonstrate how the probability of a major

epidemic can be calculated in a relatively complex epidemiological setting. In the model,

the numbers of the *N* hosts that are (*S*)usceptible, (*E*)xposed, (*I*)nfectious and (*R*)emoved

are tracked, as well as the number of the N^V vectors that are (S^V)usceptible, (E^V)xposed

and (I^V)nfectious. The deterministic version of this model is given by

231
$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta I^V S,$$

232
$$\frac{\mathrm{d}E}{\mathrm{d}t} = \beta I^V S - \alpha_H E,$$

233
$$\frac{\mathrm{d}I}{\mathrm{d}t} = \alpha_H E - \mu I,$$

234
$$\frac{\mathrm{d}R}{\mathrm{d}t} = \mu I,$$

235
$$\frac{\mathrm{d}S^V}{\mathrm{d}t} = \delta N^V - \beta_V S^V \frac{I}{N} - \delta S^V,$$

$$236 \qquad \frac{\mathrm{d}E^V}{\mathrm{d}t} = \beta_V S^V \frac{I}{N} - (\delta + \alpha_V)E^V,$$

$$237 \qquad \frac{\mathrm{d}I^V}{\mathrm{d}t} = \alpha_V E^V - \delta I^V. \qquad (4)$$

The parameters β and β_V govern the rates at which infectious vectors infect susceptible

hosts and infectious hosts infect susceptible vectors, respectively. The mean latent

periods of infections in hosts and vectors are given by $1/\alpha_H$ and $1/\alpha_V$. The parameter μ

is the rate of removal of infectious hosts, and δ describes the death rate of every vector.

In the analogous stochastic model, the number of infected human hosts arising from a

single infected human (accounting for human-vector-human transmission) in an

otherwise entirely susceptible population of humans and vectors is given by

$$246 \qquad R_0^{HV} \times \rho^{E^V \to I^V} \times R_0^{VH} = \frac{\beta_V \alpha_V \beta N^V}{\mu(\delta + \alpha_V)\delta},$$

248 where $R_0^{HV} = \frac{1}{\mu}\frac{\beta_V N^V}{N}$ is the expected number of vectors infected (and going on to enter

249 the exposed class) by a single infectious human, $\rho^{E^V \to I^V} = \frac{\alpha_V}{\delta + \alpha_V}$ is the proportion of

250 exposed vectors that become infectious and $R_0^{VH} = \frac{1}{\delta}\beta N$ is the expected number of

251 humans infected by a single infectious vector.

253 The basic reproduction number is given by

$$255 \qquad R_0 = \sqrt{\frac{\beta_V \alpha_V \beta N^V}{\mu(\delta + \alpha_V)\delta}},$$

where the square root accounts for the fact that it takes two generations for infected

humans to generate new infections, since new infections require host-vector-host

transmission [50,51]. We note that in some studies, e.g. [49], the square root is omitted

from the definition of R_0. In contrast to the expression calculated by Kucharski *et al*. [49],

to facilitate simulation of the stochastic model we also explicitly track the total number of

vectors, N^V, rather than the density.

<u>Probability of a major epidemic (branching process estimate)</u>

Standard estimate (stochastic SIS/SIR models)

The commonly used estimate for the probability of a major epidemic when a pathogen

first arrives in a host population [8,9,21,27,29–35] can be derived by assuming that

infections occur according to a branching process, making the assumptions that the

susceptible population is large and that infection lineages arising from different infected

hosts are independent. When a single infected host arrives in an otherwise susceptible

population, the branching process estimate for the probability of a major epidemic is given

by

$$\text{Prob(major epidemic)} \approx \begin{cases} 0 \ \text{for } R_0 \leq 1, \\ 1 - \dfrac{1}{R_0} \ \ \text{for } R_0 > 1. \end{cases}$$

This expression is derived in Text S1.

If instead there are $I(0)$ infected individuals initially rather than one, then for no major

epidemic to occur, it is necessary for each initial infection lineage to die out, leading to

the approximation given in equation (1) whenever $R_0 > 1$.

Standard estimate (more complex models)

Here we show how the branching process estimate for the probability of a major epidemic

can be derived for more complex systems by considering the stochastic version of the

model of Zika virus given by the system of equations (4). Invasion probabilities have been

estimated previously for vector-borne pathogens, for example by considering the

distributions of secondary infections from infected hosts/vectors and their associated

probability generating functions [22], however we use an alternative approach here that

is straightforward to understand and extend to a range of epidemiological settings.

We denote the probability of no major epidemic occurring starting from i exposed or

infectious human hosts, j exposed vectors and k infectious vectors by q_{ijk}. We must

consider exposed and infectious vectors separately to account for the possibility that

exposed vectors die before becoming infectious.

Starting from a single infectious host introduced into an entirely susceptible population of

hosts and vectors, and conditioning on the next event, gives

299
$$q_{100} = \frac{\mu}{\mu + \frac{\beta_V N^V}{N}} q_{000} + \frac{\frac{\beta_V N^V}{N}}{\mu + \frac{\beta_V N^V}{N}} q_{110},$$

300   because the next event can then either be recovery of the infectious host (with

301   probability $\frac{\mu}{\mu+\frac{\beta_V N^V}{N}}$) or infection of a susceptible vector (with probability $\frac{\frac{\beta_V N^V}{N}}{\mu+\frac{\beta_V N^V}{N}}$). Similarly,

302   starting from a single exposed or infectious vector gives

303
$$q_{010} = \frac{\delta}{\delta+\alpha_V} q_{000} + \frac{\alpha_V}{\delta+\alpha_V} q_{001},$$

304
$$q_{001} = \frac{\delta}{\delta+\beta N} q_{000} + \frac{\beta N}{\delta+\beta N} q_{101}.$$

305   We again assume that infection lineages are independent, permitting us to approximate

306   terms with two exposed or infectious individuals by non-linear terms involving single

307   exposed or infectious individuals, e.g. $q_{110} \approx q_{100}q_{010}$. Noting that $q_{000} = 1$, the three

308   equations above can be solved to give expressions for q_{100}, q_{010} and q_{001}. In particular,

309   the probability of a major epidemic starting from a single infected host is then

310
$$1 - q_{100} = \begin{cases} 0 & \text{for } R_0 \leq 1, \\ \dfrac{(R_0)^2 - 1}{(R_0)^2 + R_0^{VH}} & \text{for } R_0 > 1. \end{cases} \quad (5)$$

In this expression, R_0^{VH} is the expected number of humans infected by a single infectious

vector in an otherwise entirely susceptible population of humans and vectors.

<u>Probability of a major epidemic ("Concurrent size" definition)</u>

As described in the introduction, we first define a major epidemic to be an outbreak in

which the maximum number of simultaneously infected individuals is above a threshold

value, which we denote by *M*. The value of *M* of relevance in practical applications might

be set by the capacity for treatment.

Deterministic SIS and SIR models

In the deterministic SIS model, the maximum number of simultaneously infected

individuals over the course of the outbreak is given by

325
$$I_{\max} = \begin{cases} N\left(1 - \dfrac{1}{R_0}\right) \text{ for } I(0) \le N\left(1 - \dfrac{1}{R_0}\right), \\ I(0) \text{ for } I(0) > N\left(1 - \dfrac{1}{R_0}\right), \end{cases}$$

327   as shown in Text S1.

329   In the deterministic SIR model, on the other hand,

331
$$I_{\max} = \begin{cases} I^* \text{ for } I(0) \le I^*, \\ I(0) \text{ for } I(0) > I^*, \end{cases}$$

333   where $I^* = -\dfrac{N}{R_0} + \dfrac{N}{R_0}\ln\left(\dfrac{N}{S(0)R_0}\right) + S(0) + I(0)$, and again this is shown in Text S1.

335   The probability of a major epidemic according to the "Concurrent size" definition under

336   the deterministic SIS and SIR models is then

338
$$\text{Prob(major epidemic)} = \begin{cases} 1 \text{ for } M \le I_{\max}, \\ 0 \text{ for } M > I_{\max}, \end{cases}$$

where I_{max} is given by the corresponding expressions above depending which model is

used.

Stochastic SIS model

Under the stochastic SIS model, however, calculating the probability of a major epidemic

is more challenging [52]. Nevertheless, we can calculate this value analytically, which is

advantageous since approximating this quantity using model simulations can be time

consuming given that outbreaks under the SIS model can persist for indefinitely long

periods.

We denote the probability of no major epidemic starting with I infected individuals as q_I,

and assume that the rest of the population is susceptible. Conditioning on whether the

first event is an infection or recovery event gives

354
$$q_I = \frac{\beta IS}{\beta IS + \mu I} q_{I+1} + \frac{\mu I}{\beta IS + \mu I} q_{I-1},$$

356   for I = 1,2,…,M-1. We solve this tridiagonal system of equations with boundary conditions

357   q_0 = 1 and q_M = 0, since a major epidemic certainly does not occur if there are initially no

358   infected individuals, and certainly does if there are initially M infected individuals. Noting

359   that $S = N - I$, these equations can be put in the form

361
$$q_{I+1} - q_I = C_I(q_I - q_{I-1}),$$

363 where

$$C_I = \frac{\mu I}{\beta I (N - I)}.$$

366 Iterating this gives

$$q_{I+1} - q_I = A_I(q_1 - q_0), \qquad (6)$$

369 where

$$A_I = \prod_{j=1}^{I} C_j,$$

$$= \frac{\left(\frac{N}{R_0}\right)^I}{I! \binom{N-1}{I}}.$$

373 Adding equation (6) to itself for $I = 1,2,\ldots,M - 1$ and rearranging gives

$$q_1 = \frac{\sum_{I=1}^{M-1} A_I}{1 + \sum_{I=1}^{M-1} A_I}.$$

377 Instead adding equation (6) to itself for $I = 1,2,\ldots,k - 1$ gives

$$q_k = \frac{\sum_{I=k}^{M-1} A_I}{1 + \sum_{I=1}^{M-1} A_I}.$$

381 The probability of a major epidemic is then given by

$$\text{Prob(major epidemic)} = 1 - q_{I(0)} = \begin{cases} 0 \text{ for } I(0) = 0, \\ \dfrac{1}{1 + \sum_{I=1}^{M-1} A_I} \text{ for } I(0) = 1, \\ \dfrac{1 + \sum_{I=1}^{I(0)-1} A_I}{1 + \sum_{I=1}^{M-1} A_I} \text{ for } 1 < I(0) < M, \\ 1 \text{ for } I(0) \geq M. \end{cases} \quad (7)$$

Stochastic SIR model

Under the stochastic SIR model, the probability of a major epidemic according to the "Concurrent size" definition starting from any state (I,R) is calculated using an iterative approach. Denoting the probability of a major epidemic starting from state (I,R) by $p_{I,R}$, then conditioning on what happens next gives

$$p_{I,R} = \frac{\beta I(N - I - R)}{\beta I(N - I - R) + \mu I} p_{I+1,R} + \frac{\mu I}{\beta I(N - I - R) + \mu I} p_{I-1,R+1}.$$

This system can be solved with boundary conditions $p_{0,R} = 0$, $p_{I,N-M+1} = 0$ and $p_{M,R} = 1$. To do this, the probability of a major epidemic is deduced for the following states (in order): (I,R) = $(M\text{-}1,N\text{-}M)$, $(M\text{-}2,N\text{-}M)$, …, $(1,N\text{-}M)$, $(M\text{-}1,N\text{-}M\text{-}1)$, …,$(1,N\text{-}M\text{-}1)$,… $(M\text{-}1,0)$,…,$(1,0)$. For a schematic showing the order in which these probabilities are deduced, see Fig S1.

Zika transmission model

The probability of a major epidemic under the "Concurrent size" definition is approximated

by simulating the model 10,000 times using the Gillespie direct method [53], and then

calculating the proportion of simulations in which the number of infected human hosts

exceeds M at any point during the simulation.

<u>Probability of a major epidemic ("Total infections" and "Duration" definitions)</u>

We also consider the probability of a major epidemic according to the deterministic and

stochastic SIS model for the "Total infections" and "Duration" definitions of a major

epidemic. Under the "Total infections" definition, a major epidemic is assumed to be an

outbreak in which at least F infections occur over the course of the outbreak. Under the

"Duration" definition, a major epidemic is defined to be an outbreak that persists for at

least T days.

In the deterministic SIS model, whenever $R_0 > 1$, the outbreak persists indefinitely with

an infinite number of infection events. As a result, any outbreak is a major epidemic under

the "Total infections" and "Duration" definitions.

In contrast, in any simulation of the stochastic SIS model, the number of infected

individuals will always reach zero, even if this takes a long time. As a result, the probability

of a major epidemic under the "Total infections" and "Duration" definitions is not simply

one or zero depending on the value of R_0. We approximate the probability of a major

epidemic under these definitions by simulating the model 10,000 times using the Gillespie

direct method [53], and recording the proportion of simulations in which there are at least

F infections or that have a duration of at least *T* days.

427   **3.   RESULTS**

To begin to understand outbreak dynamics under the SIS, SIR and Zika transmission

models, we first numerically solved the deterministic models given by the systems of

equations (2), (3) and (4) with $R_0 = 1.5$ in each case (Fig 2). In a deterministic setting, the

SIS model predicts the largest number of individuals simultaneously infected as well as

the most infections in total. Epidemics persisted forever (i.e. *I* remained larger than zero)

under all three models, although the number of infected hosts tended to zero under the

SIR and Zika transmission models.

However, our main focus is the probability of a major epidemic soon after the pathogen

enters the system. For any definition of "major epidemic", according to a deterministic

model the corresponding probability is either zero or one depending on the values of

model parameters. We therefore considered the more realistic stochastic models, in

which demographic stochasticity is included. In the following sections, first we calculate

the probability of a major epidemic for the stochastic SIS model under the "Concurrent

size" definition. We then consider different epidemiological models, as well as different

definitions of a major epidemic. In each case, the probability of a major epidemic for the

particular epidemiological model-definition of a major epidemic pair under consideration

is compared to the branching process approximation to the probability of a major epidemic

for that model. Results are shown in Figs 3-5, as well as summarised in Tables 1 and 2.

Figure 2. Numerical solutions of the deterministic SIS, SIR and Zika virus transmission models when the

basic reproduction number is $R_0 = 1.5$. (a) Number of infected individuals through time according to the

deterministic SIS model. (b) Cumulative number of infections through time according to the deterministic

SIS model. (c)-(d) Equivalent to a-b but for the deterministic SIR model. (e)-(f) Equivalent to a-b but for the

deterministic Zika virus transmission model. Parameters for deterministic SIS and SIR models: $N = 1,000$,

$\beta = 0.00015$ per day, $1/\mu = 10$ days. Parameters for deterministic Zika virus transmission model: $N =$

$1,000$, $N^V = 10,000$, $1/\alpha_V = 10.5$ days, $1/\alpha_H = 5.9$ days, $1/\mu = 5$ days, $1/\delta = 7.8$ days, $\beta = 6.15 \times 10^{-5}$ per

day, $\beta_V = 0.22$ per day [49].

<u>The probability of a major epidemic</u>

We calculated the probability of a major epidemic according to the stochastic SIS model

under the "Concurrent size" definition of a major epidemic – i.e. an outbreak in which a

threshold number of simultaneously infected individuals is exceeded. In this case, as

described in Methods, it is possible to calculate the probability of a major epidemic

analytically.

We show the probability of a major epidemic for a range of values of the major epidemic

threshold, M, in Fig 3a. For R_0 much larger than one, we found that the probability of a

major epidemic was approximated closely by the standard branching process estimate

for many values of the major epidemic threshold, M. When, however, R_0 was close to

one, the standard estimate corresponded to a single choice of M (see e.g. blue lines in

Fig 3a, where the solid line is close to the corresponding dotted line in only one place, i.e.

for a single value of M). The parameter regime in which R_0 is close to one is important in

many real epidemiological systems since the aim of control strategies is usually to reduce

the reproduction number below one [54,55].

Figure 3. Probability of a major epidemic under the SIS model, where a major epidemic is defined as an outbreak in which M simultaneously infected individuals is exceeded ("Concurrent size" definition). (a) Dependence on R_0. Solid lines represent the true probability of a major epidemic (system of equations (7)), dotted lines represent the branching process estimate (equation (1)), and dots show the maximum number simultaneously infected in the analogous deterministic models (for values of M below this, the probability of a major epidemic in the relevant deterministic model is 1). R_0 is varied by changing the value of β. (b) Equivalent to a, but showing dependence on the population size, N. (c) Equivalent to a, but showing dependence on the initial number of infected individuals, $I(0)$. (d). Single simulation of the stochastic SIS

model (blue), and numerical solution of deterministic SIS model (red dotted). The value of I in the stochastic

simulation will continue to fluctuate about the deterministic value until I reaches 0. Parameter values

(except where stated): N = 1,000, R_0 = 1.5, $I(0)$ = 1 and $R(0)$ = 0 and the remainder of the population

susceptible initially. In panel d, $\beta = 0.000015$ per day and $1/\mu = 10$ days.

In large host populations, the probability of a major epidemic as a function of M took the

form of a step function (Fig 3b). This suggested that, if the pathogen successfully invaded

the population, then the number of infected individuals would definitely reach a specific

maximum value which is determined by R_0. For example, for outbreaks with R_0 = 1.5, the

pathogen will invade the population with probability 0.33, and, if this occurs, then around

two-thirds of the population will be infected simultaneously at some time during the

epidemic.

An approximation to the maximum value of I that will be reached in the stochastic SIS

model in the large N case is given by

503
$$\max(I) \approx \max\left(2N\left(1 - \frac{1}{R_0}\right), N\right).$$

The expression $2N\left(1 - \frac{1}{R_0}\right)$ is twice the maximum value of I in the corresponding

deterministic model. This approximation involves the assumption that, if the pathogen

invades in the stochastic SIS model, I will fluctuate approximately symmetrically around

the deterministic equilibrium value (Fig 3d). It is likely, then, that by the time the pathogen

dies out by reaching $I = 0$, the number of infected individuals will at some stage also have

reached approximately double the value it fluctuated around too. In Fig 3b, when $N =$

10,000, this approximation gives a maximum proportion of the population simultaneously

infected of 0.67, when the true value is 0.62.

We also note that, if the expression above for max(I) is reformulated to give the maximum

proportion of the population that is simultaneously infected, the resulting expression is

independent of the population size N. This can also be seen graphically – in Fig 3b, if the

pathogen successfully invaded the population then the maximum proportion of hosts that

were simultaneously infected was approximately independent of the size of the host

population, so long as N was sufficiently large.

<u>Different epidemiological models</u>

We considered the probability of a major epidemic ("Concurrent size" defintion) under the

SIR and Zika virus transmission models. For the stochastic SIR model, we used an

iterative method to calculate the probability of a major epidemic as described in Methods.

For the stochastic Zika virus transmission model, we simulated the model in a population

of $N = 1,000$ human hosts and $N^V = 10,000$ vectors using the Gillespie direct algorithm

[53], using parameter values from Kucharski *et al.*, 2016 [49] – see caption of Fig 4. The

value of R_0 was then varied in Fig 4b by altering the parameter β that governs the rate at

which infected vectors infect susceptible human hosts.

Figure 4. Probability of a major epidemic under the "Concurrent size" definition of a major epidemic, for the

SIR and Zika virus transmission models. (a) SIR model. Solid lines represent the true probability of a major

epidemic calculated using the iterative method described in Methods, dotted lines represent the branching

process estimate (equation (1)), and dots show the maximum number simultaneously infected in the

analogous deterministic models (for values of M below this, the probability of a major epidemic in the

relevant deterministic model is 1). R_0 is varied by changing the value of β. (b) Equivalent to a, but for the

Zika virus transmission model (where M refers to the number of simultaneously infected hosts). For the

Zika transmission model, the true probability of a major epidemic is calculated by simulation, and the

branching process estimate is given by equation (5). For both models, $N = 1,000$. Other parameters for the

Zika virus transmission model: $N^V = 10,000$, $1/\alpha_V = 10.5$ days, $1/\alpha_H = 5.9$ days, $1/\mu = 5$ days, $1/\delta = 7.8$

days, $\beta_V = 0.22$ per day [49]. Initial conditions for both models comprise of a single infected host, with all

other individuals (for the Zika transmission model, hosts and vectors) susceptible.

Under the stochastic SIR and Zika models, for R_0 larger than and not close to one, the

maximum number of simultaneously infected individuals whenever the pathogen invaded

the host population was typically smaller than under the SIS model (*cf.* Fig 2).

Nonetheless, we found qualitatively similar behaviour in these cases – the probability of

a major epidemic approximated using a branching process corresponded to a wide range

of values of the major epidemic threshold when R_0 was high (Fig 4). However, even if that

is the case, the practically relevant value of the major epidemic threshold (e.g. the number

of available hospital beds) may not give a probability of a major epidemic that matches

the branching process estimate. For example, if $R_0 = 2$ and 250 beds are available, for

the SIR model the probability of a major epidemic under the "Concurrent size" definition

is 0 (solid grey line in Fig 4a), yet the branching process estimate for the probability of a

major epidemic is 0.5 (dotted grey line in Fig 4a).

<u>Alternative definitions of a major epidemic</u>

For the stochastic SIS model, we then calculated the probability of a major epidemic for

different definitions of a major epidemic – specifically, outbreaks in which there are at

least F infection events (the "Total infections" definition – Fig 5a) or outbreaks that persist

for at least T days (the "Duration" definition – Fig 5b).

Figure 5. Probability of a major epidemic under the SIS model, for alternative definitions of a major epidemic.

(a) A major epidemic is defined as an outbreak in which at least F infections occur ("Total infections" definition). (b) A major epidemic is defined as an outbreak that persists for at least T days ("Duration" definition). Solid lines represent the true probability of a major epidemic assessed via simulation of the stochastic model, and dotted lines represent the branching process estimate (equation (1)). The x-axis is shown on a log-scale, for F between 1 and 2000 (panel a) and T between 1 and 6000 (panel b). The step function in panel a reflects the fact that the total number of infections can only take discrete values. Results of the deterministic model are not included in the figure, since under the deterministic SIS model epidemics persist indefinitely and generate an infinite number of infections whenever $R_0 > 1$. Parameter values: $N =$ 1,000, $I(0) = 1$ and $R(0) = 0$ and the remainder of the population susceptible initially. In both panels, R_0 is varied by changing the value of β. In panel b, $1/\mu = 10$ days.

In the stochastic SIS model, if the pathogen invaded the host population then it tended to persist for long periods. Consequently, the branching process estimate corresponded to a very wide range of major epidemic thresholds under the "Total infections" or "Duration" definitions (i.e. values of F or T) compared to under the "Concurrent size" definition. As a

result, in these specific cases (i.e. when the stochastic SIS model was used and a major

epidemic was defined according to the "Total infections" or "Duration" definitions) it can

be concluded that the branching process approximation often leads to sensible estimates

of the risk posed by invading pathogens. Nonetheless, even in these cases, for small or

very large values of the major epidemic thresholds the probability of a major epidemic

does not match the branching process estimate, particularly when R_0 was larger than but

close to one (see e.g. red line in Fig 5b).

	SIS model	SIR model	Zika host-vector model
$R_0 = 1.1$	$21 \leq M \leq 54$ (Fig 3a)	$18 \leq M \leq 21$ (Fig 4a)	$6 \leq M \leq 7$ (Fig 4b)
$R_0 = 1.6$	$8 \leq M \leq 681$ (Fig 3a)	$8 \leq M \leq 55$ (Fig 4a)	$3 \leq M \leq 47$ (Fig 4b)
$R_0 = 2$	$6 \leq M \leq 866$ (Fig 3a)	$7 \leq M \leq 117$ (Fig 4a)	$2 \leq M \leq 83$ (Fig 4b)

595
Table 1. Effect of underlying epidemiology. For which values of the threshold (number of hosts
simultaneously infected, M) in the "Concurrent size" definition of a major epidemic is the branching
process approximation to the probability of a major epidemic accurate? Range of values here are those
for which the branching process approximation is within 0.01 of the probability of a major epidemic.
Results are summarised for the SIS, SIR and Zika virus models, for the parameter values shown in the
relevant figure captions.
602
603
604

	Concurrent size	Total infections	Duration
$R_0 = 1.1$	$21 \leq M \leq 54$ (Fig 3a)	$138 \leq F \leq 878$ (Fig 5a)	$183 \leq T \leq 458$ (Fig 5b)
$R_0 = 1.6$	$8 \leq M \leq 681$ (Fig 3a)	$F \geq 18$ (Fig 5a)	$T \geq 51$ (Fig 5b)
$R_0 = 2$	$6 \leq M \leq 866$ (Fig 3a)	$F \geq 10$ (Fig 5a)	$T \geq 34$ (Fig 5b)

605
Table 2. Effect of definition of a major epidemic. For which values of the thresholds in the more practically
relevant definitions of a major epidemic is the branching process approximation to the probability of a
major epidemic accurate? Range of values here are those for which the branching process approximation
is within 0.01 of the probability of a major epidemic. Results are summarised for the SIS model, for the
parameter values shown in the relevant figure captions. "Total infections" threshold values were tested up
to a maximum of $F = 2,000$ (Fig 5a) and "Duration" threshold values were tested up to $T = 6,000$ (Fig 5b).
612

4. DISCUSSION

Determining the risk of an emerging outbreak developing into a major epidemic is vital for

planning whether or not intervention and/or containment strategies will be necessary.

When a pathogen arrives in a new location, the probability of a major epidemic can be

approximated by assuming that infections occur according to a branching process. For

simple models such as the stochastic SIS and SIR models, this leads to the probability of

a major epidemic in equation (1). It is also possible to estimate the probability of a major

epidemic according to the branching process approximation using models with more

complexity, as we showed by considering the case of host-vector transmission (see

equations (4) and (5)).

However, the branching process estimate for the probability of a major epidemic is not

necessarily accurate when definitions of a major epidemic are used that address practical

aspects of disease control (Fig 3). The branching process estimate corresponds to a

range of choices of the epidemic thresholds in our definitions when R_0 is much greater

than one, or when the population size is extremely large (see e.g. different values of M in

Fig 3a,b). However, when R_0 is close to one and the population is not large, the standard

estimate can correspond to a single choice of the epidemic threshold (see e.g. blue and

red lines in Fig 3a, and Tables 1 and 2). For specific outbreaks, even when both R_0 and

the population size are large, the branching process estimate may not be relevant – since

the range of choices of the major epidemic threshold that the standard estimate

corresponds to may or may not include the specific threshold of practical importance in

the outbreak, for example the number of hospital beds available. Consequently, using the

standard branching process estimate for the probability of a major epidemic could lead to

the risk of a major epidemic being incorrectly assessed, potentially including scenarios in

which a major epidemic develops when previously deemed unlikely. Our main conclusion

that the branching process estimate for the probability of a major epidemic may or may

not match the true probability of a major epidemic when a practically relevant definition is

used holds for a range of epidemiological systems (Fig 4) as well as different definitions

of a major epidemic that apply in alternative settings (Fig 5). We note that exactly how

well the branching process estimate reflects a range of threshold values depends on the

underlying model as well as the relevant quantity differentiating major epidemics from

minor outbreaks (*cf*. Figs 3a and 5).

We considered practical definitions of a major epidemic that were based on thresholds

such as the availability of treatment. A previous study defined major epidemics according

to a threshold in the percentage of the population ever infected, and concluded that

epidemiological modellers should report the precise cutoff used to define a major

epidemic in model simulations [56]. Their conclusion was based on the observation that

different thresholds in the percentage of hosts ever infected corresponded to wide

variations in the other outputs of model simulations including the number of dead hosts

or the time of the epidemic peak. We support this conclusion, and indeed some authors

have reported the definition of a major epidemic they used clearly – for example, Keeling

et al. [57] define a major epidemic to be an outbreak in which at least one-third of the

population becomes infected. However, we also emphasise that the precise type of

threshold, and the value used, should be chosen according to practical relevance in the

particular system under consideration, rather than simply an arbitrary threshold in the

number of individuals ever infected.

Under the first definition of a major epidemic that we considered (the "Concurrent size"

definition), the probability of a major epidemic was assessed in the context of the capacity

for treatment by estimating whether or not a threshold number of simultaneously infected

individuals was likely to be exceeded. This definition may be practically relevant in a range

of scenarios. For example, real-time analysis of a diphtheria epidemic in Cox Bazar's in

Bangladesh involved assessing the number of hospital beds that were needed [39]. The

number of beds required was approximated in that study by using a model to forecast

disease incidence, assuming that 15% of reported cases would require treatment as

inpatients with an average hospital stay of five days for each case. The number of hospital

beds that were already available might have provided a practically relevant major

epidemic threshold. Another example for which this type of threshold might apply is citrus

greening disease in Brazil, for which a law was introduced stating that a citrus grove must

be destroyed if 28% of trees in the grove were infected and symptomatic [58,59]. Major

epidemics could therefore be defined as outbreaks in which more than 28% of trees in a

grove are infected and symptomatic concurrently. Other examples for which interventions

were introduced as soon as a threshold in the number simultaneously infected was

reached include the development of the National Chlamydia Screening Programme in the

United Kingdom in 2002 in response to the large size of the infected population [60].

However, no single definition of a major epidemic will be relevant in all situations. To

illustrate the ambiguity to which this might lead, we also considered two other definitions

of a major epidemic. In one of these (the "Total infections" definition), whether or not an

outbreak was classified as a major epidemic referred to the total number of infection

events over the course of the outbreak, rather than the maximum number simultaneously

infected. This might correspond to the total number of treatments required, which may be

an important threshold if treatments have been stockpiled prior to the outbreak [41]. This

definition might also be relevant if, for example, a policy-maker has to choose how to

deploy resources between two different epidemics. If there are only sufficient resources

to contain one outbreak, and both epidemics are equally controllable, then it might be

preferable to choose the one that is likely to generate more infections. In other real-world

scenarios, alternative definitions might be appropriate. We also considered major

epidemics defined as outbreaks that persist for a threshold length of time (the "Duration"

definition). Different definitions of a major epidemic might appear contradictory – for

example, treatment can act to reduce the total number of infections yet increase the

duration of the outbreak [61], making a major epidemic less likely under the "Total

infections" definition of a major epidemic but more likely under the "Duration" definition.

Our intention here was to use very simple models to demonstrate the principle that

different definitions of a major epidemic lead to different probabilities of a major epidemic.

Although simple models are commonly used, accurate outbreak forecasts require a model

carefully matched to the epidemiology of the host-pathogen system, potentially including

asymptomatic transmission [9,62,63] or spread between spatially distinct regions [29,64].

For certain definitions, it may be necessary to include convalescent hosts in the model

explicitly. For example, if convalescent individuals require resources, such as beds in

treatment rooms or hospitals, and the definition of a major epidemic is linked to the

availability of resources (as in the case of the "Concurrent size" definition), then these

individuals should be modelled, potentially by including them in a new compartment

following the infectious class. More complex definitions of major epidemics could also be

used, for example requiring multiple criteria to be satisfied for an outbreak to be classified

as a major epidemic. In these more complicated scenarios, analytic calculations of the

probability of a major epidemic might not be possible. Model simulations can then be used

to assess the probability of a major epidemic, as we showed for a host-vector model of

Zika virus transmission (Fig 4b).

We note that practical use of the methods presented here at the start of an emerging

outbreak to assess the major epidemic risk would require the wide range of interventions

that are introduced in outbreak response settings to be integrated into the models

explicitly. One way in which control can be included is to consider the effective

reproduction number when the pathogen arrives in the system instead of the basic

reproduction number, since the effective reproduction number accounts for interventions

[26,54,55,65–67]. In that case, the results that we presented would be unchanged (except

that e.g. the lines in Fig 3a would correspond to different values of the effective

reproduction number). Temporal changes in interventions once an outbreak is underway

have been approximated in epidemiological models by assuming that the values of the

parameters governing transmission change during the outbreak, either by assuming that

transmissibility changes at a single timepoint [68,69] or continuously as the outbreak

progresses [70,71]. However, for detailed descriptions of control to be included in

estimates of the major epidemic risk, more complex interventions should be included in

model simulations. Since models are often used to test possible control strategies

[12,20,72–74], this is a simple extension of the results presented here.

In summary, we have shown that how precisely to define a major epidemic should be

considered carefully in future studies. The definition of a major epidemic should be

designed to match the questions of interest in the particular setting being considered.

Standard approximations based on branching processes should not always be used as

the default. Only once a "major epidemic" has been defined precisely can the probability

of a major epidemic occurring be properly assessed. Providing an explicit demonstration

of the consequences of not considering a practically relevant definition in evaluating the

risk of an epidemic is the key contribution of this paper.

SUPPLEMENTARY MATERIAL

Text S1.

Figure S1. Schematic showing how the probability of a major epidemic under the

"Concurrent size" definition can be deduced for the stochastic SIR model. The probability

corresponding to each state (I,R) is deduced iteratively in the order 1,2,3... (red). The

black arrows indicate which previous values are used to inform each new deduction.

COMPETING INTERESTS

We have no competing interests.

AUTHORS' CONTRIBUTIONS

All authors designed the study. RNT undertook the research and wrote the first draft of

the manuscript. All authors revised the manuscript.

FUNDING

This research was funded by BBSRC via a G2O PhD studentship (RNT), and by Christ

Church, Oxford, via a Junior Research Fellowship (RNT).

764   **REFERENCES**

1.   Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife —

threats to biodiversity and human health. Science (80-). 2000;287: 443–450.

2.   Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P.

Emerging infectious diseases of plants: pathogen pollution, climate change and

agrotechnology drivers. Trends Ecol Evol. 2004;19: 535–544.

doi:10.1016/j.tree.2004.07.021

3. Taylor LH, Latham SM, Woolhouse MEJ. Risk factors for human disease

emergence. Philos Trans R Soc B. 2001;356: 983–989.

doi:10.1098/rstb.2001.0888

4. Morens DM, Folkers GK, Fauci AS. The challenge of emergin and re-emerging

infectious diseases. Nature. 2004;430: 242–249.

5. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global

trends in emerging infectious diseases. Nature. 2008; 990–994.

doi:10.1038/nature06536

6. Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, et al.

Emerging fungal threats to animal, plant and ecosystem health. Nature. 2012;484:

186–194. doi:10.1038/nature10947

7. Thompson RN, Brooks-Pollock E. Detection, forecasting and control of infectious

disease epidemics: modelling outbreaks in humans, animals and plants. Philos

Trans R Soc B. 2019;374: 20190038.

8. Craft ME, Beyer HL, Haydon DT. Estimating the probability of a major outbreak

from the timing of early cases: an indeterminate problem ? PLoS One. 2013;8:

e57878. doi:10.1371/journal.pone.0057878

9. Thompson RN, Gilligan CA, Cunniffe NJ. Detecting presymptomatic infection is

necessary to forecast major epidemics in the earliest stages of infectious disease

outbreaks. PLoS Comput Biol. 2016;12: e1004836.

792   doi:10.1371/journal.pcbi.1004836

793 10. Ramsay ME. Measles: the legacy of low vaccine coverage. Arch Dis Child.

794   2013;98: 752–754. doi:10.1136/archdischild-2013-304292

795 11. Rohani P, Earn DJD, Grenfell BT. Impact of immunisation on pertussis

796   transmission in England and Wales. Lancet. 2000;355: 285–286.

797 12. Ferguson NM, Donnelly CA, Anderson RM. The foot-and-mouth epidemic in Great

798   Britain: Pattern of spread and impact of interventions. Science (80-). 2001;292:

799   1155–1160.

800 13. Kao RR. The role of mathematical modelling in the control of the 2001 FMD

801   epidemic in the UK. Trends Microbiol. 2002;10: 279–286.

802 14. Keeling MJ. Models of foot-and-mouth disease. Proc R Soc B Biol Sci. 2005;272:

803   1195–1202.

804 15. Keeling MJ, Woolhouse MEJ, Shaw DJ, Matthews L, Chase-Topping M, Haydon

805   DT, et al. Dynamics of the 2001 UK foot and mouth epidemic: Stochastic

806   dispersal in a heterogeneous landscape. Science (80-). 2001;294: 813–818.

807 16. Pautasso M, Aas G, Queloz V, Holdenrieder O. European ash (Fraxinus

808   excelsior) dieback – A conservation biology challenge. Biol Conserv. Elsevier Ltd;

809   2013;158: 37–49. doi:10.1016/j.biocon.2012.08.026

810 17. Cunniffe NJ, Laranjeira FF, Neri FM, DeSimone RE, Gilligan CA. Cost-effective

811   control of plant disease when epidemiological knowledge is incomplete: Modelling

812   Bahia bark scaling of citrus. PLoS Comput Biol. 2014;10: e1003753.

813   doi:10.1371/journal.pcbi.1003753

814 18. Chan M-S, Jeger MJ. An analytical model of plant virus disease dynamics with

roguing and replanting. J Appl Ecol. 1994;31: 413–427.

19.   Wu JT, Cowling BJ, Lau EHY, Ip DKM, Ho L-M, Tsang T, et al. School closure

and mitigation of pandemic (H1N1) 2009, Hong Kong. Emerg Infect Dis. 2010;16:

538. doi:10.3201/eid1603.091216

20.   Thompson RN, Gilligan CA, Cunniffe NJ. Control fast or control smart: When

should invading pathogens be controlled? PLoS Comput Biol. 2018;14.

doi:10.1371/journal.pcbi.1006014

21.   Keeling MJ, Rohani P. Modeling infectious diseases in humans and animals.

Princeton University Press; 2008.

22.   Lloyd AL, Zhang J, Root AM. Stochasticity and heterogeneity in host-vector

models. J R Soc Interface. 2007;4: 851–863. doi:10.1098/rsif.2007.1064

23.   Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect

of individual variation on disease emergence. Nature. 2005;438: 355–359.

doi:10.1038/nature04153

24.   Nishiura H, Cook AR, Cowling BJ. Assortativity and the probability of epidemic

extinction: a case study of pandemic influenza A (H1N1-2009). Interdiscip

Perspect Infect Dis. 2011; 194507.

25.   Anderson D, Watson R. On the spread of a disease with gamma distributed latent

and infectious periods. Biometrika. 1980;67: 191–198.

26.   Thompson RN, Jalava K, Obolski U. Sustained transmission of Ebola in new

locations: more likely than previously thought. Lancet Infect Dis. 2019;19: 1058–

1059. doi:10.1016/S1473- 3099(19)30360-3

27.   Swinton J. Extinction times and phase transitions for spatially structured closed

epidemics. Bull Math Biol. 1998;60: 215–230.

28.   Merler S, Ajelli M, Fumanelli L, Parlamento S, Pastore y Piontti A, Dean NE, et al.

Containing Ebola at the Source with Ring Vaccination. PLoS Negl Trop Dis.

2016;10: 1–11. doi:10.1371/journal.pntd.0005093

29.   Thompson RN, Thompson C, Pelerman O, Gupta S, Obolski U. Increased

frequency of travel in the presence of cross-immunity may act to decrease the

chance of a global pandemic. Philos Trans R Soc B. 2019;374: 20180274.

doi:10.1098/rstb.2018.0274

30.   Althaus CL, Low N, Musa EO, Shuaib F, Gsteiger. Ebola virus disease outbreak

in Nigeria: Transmission dynamics and rapid control. Epidemics. 2015;11: 80–84.

31.   Abdullah N, Kelly JT, Graham SC, Birch J, Gonçalves-Carneiro D, Mitchell T, et

al. Structure-guided identification of a nonhuman morbillivirus with zoonotic

potential. J Virol. 2018;92: e01248-18.

32.   Keeling MJ, Danon L. Mathematical modelling of infectious diseases. Br Med Bull.

2009;92: 33–42. doi:10.1093/bmb/ldp038

33.   Keeling MJ, Ross J V. On methods for studying stochastic disease dynamics. J R

Soc Inferface. 2008;5: 171–181. doi:10.1098/rsif.2007.1106

34.   Kessler DA, Shnerb NM. Solution of an infection model near threshold. Phys Rev

E. 2007;76: 010901. doi:10.1103/PhysRevE.76.010901

35.   Park AW, Gubbins S, Gilligan CA. Invasion and persistence of plant parasites in a

spatially structured host population. Oikos. 2001;94: 162–174.

36.   Kendall DG. Deterministic and stochastic epidemics in closed populations. Proc

3rd Berkeley Symp Math Stat Prob. 1956;4: 149–165.

37.   de Jong MCM. Mathematical modelling in veterinary epidemiology: why model

building is important. Prev Vet Med. 1995;25: 183–193.

38.   Nasell I. 'The threshold concept in stochastic epidemic and endemic models' in D.

Mollison, Epidemic models: their structure and relation to data. Cambridge

University Press; 1995.

39.   Finger F, Funk S, White K, Siddiqui MR, Edmunds WJ, Kucharski AJ. Real-time

analysis of the diphtheria outbreak in forcibly displaced Myanmar nationals in

Bangladesh. BMC Med. 2019;17: 58. doi:10.1038/d41586-018-05810-w

40.   Lamontagne F, Clément C, Fletcher T, Jacob ST, Fischer II WA, Fowler RA.

Doing today's work superbly well - treating Ebola with current tools. N Engl J Med.

2014;371: 1565–1566.

41.   O'Dowd A. Government says it would stockpile Tamiflu again. Br Med J.

2014;349: g6386. doi:10.1136/bmj.g6386

42.   Butler D. Africa's latest Ebola outbreak comes to swift end. Nature. 2018; 2018–

2020.

43.   Wood W. The History and Antiquities of Eyam. Whitefish, MT: Kessinger

Publishing; 1859.

44.   Scott S, Duncan CJ. Biology of plagues: evidence from historical populations.

Cambridge University Press; 2001.

45.   Whittles LK, Didelot X. Epidemiological analysis of the Eyam plague outbreak of

1665-1666. Proc R Soc B Biol Sci. 2016;283: 20160618.

46.   Brauer F. "Compartmental models in epidemiology" in F. Brauer, P. van den

Driessche, J. Wu - Mathematical Epidemiology. Springer; 2014. doi:10.1007/978-

3-540-78911-6

47.   Raggett GF. A stochastic model of the Eyam plague. J Appl Stat. 1982;9: 212–

225. doi:10.1080/02664768200000021

48.   Cullerne JP, French A, Poon D, Baxter A, Thompson RN. The pedagogical power

of context: extending the epidemiology of Eyam. Phys Educ. 2019;

49.   Kucharski AJ, Funk S, Eggo RM, Mallet H-P, Edmunds WJ, Nilles EJ.

Transmission Dynamics of Zika Virus in Island Populations : A Modelling Analysis

of the 2013 – 14 French Polynesia Outbreak. PLoS Negl Trop Dis. 2016;10:

e0004726. doi:10.1371/journal.pntd.0004726

50.   van den Driessche P, Watmough J. Further notes on the basic reproduction

number. In "Mathematical Epidemiology." Springer; 2008.

51.   Heffernan JM, Smith RJ, Wahl LM. Perspectives on the basic reproductive ratio. J

R Soc Inferface. 2005;2: 281–293. doi:10.1098/rsif.2005.0042

52.   Artalejo JR, Economou A, Lopez-Herrero MJ. The maximum number of infected

individuals in SIS epidemic models: Computational techniques and quasi-

stationary distributions. J Comput Appl Math. 2010;233: 2563–2574.

doi:10.1016/j.cam.2009.11.003

53.   Gillespie DT. Exact stochastic simulation of coupled chemical reactions. J Phys

Chem. 1977;8: 2340–2361.

54.   Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to

estimate time-varying reproduction numbers during epidemics. Am J Epidemiol.

2013;178: 1505–1512. doi:10.1093/aje/kwt133

55.   Thompson RN, Stockwin JE, Gaalen RD Van, Polonsky JA, Kamvar ZN, Demarsh

PA, et al. Improved inference of time-varying reproduction numbers during

infectious disease outbreaks. Epidemics. 2019;19: 100356.

doi:10.1016/j.epidem.2019.100356

56. Orbann C, Sattenspiel L, Miller E, Dimka J. Defining epidemics in computer

simulation models: How do definitions influence conclusions? Epidemics. Elsevier

B.V.; 2017;19: 24–32. doi:10.1016/j.epidem.2016.12.001

57. Keeling MJ. The implications of network structure for epidemic dynamics. Theor

Popul Biol. 2005;67: 1–8. doi:10.1016/j.tpb.2004.08.002

58. Craig AP, Cunniffe NJ, Parry M, Laranjeira FF, Gilligan CA. Grower and regulator

conflict in management of the citrus disease Huanglongbing in Brazil: A modelling

study. J Appl Ecol. 2017;55: 1956–1965. doi:10.1111/1365-2664.13122

59. Belasque Jr. J, Bassanezi RB, Yamamoto PT, Ayres AJ, Tachibana A, Violante

AR, et al. Letter to the editor: Lessons from Huanglongbing management in Sao

Paulo state, Brazil. J Plant Pathol. 2010;92: 285–302.

60. LaMontagne DS, Fenton KA, Randall S, Anderson S, Carter P. Establishing the

National Chlamydia Screening Programme in England: results from the first full

year of screening. Sex Transm Infect. 2004;80: 335–341.

61. Hollingsworth TD, Klinkenberg D, Heesterbeek H, Anderson RM. Mitigation

strategies for pandemic influenza A: Balancing conflicting policy objectives. PLoS

Comput Biol. 2011;7: e1001076. doi:10.1371/journal.pcbi.1001076

62. Hyatt-Twynam SR, Parnell S, Stutt ROJH, Gottwald TR, Gilligan CA, Cunniffe NJ.

Risk-based management of invading plant disease. New Phytol. 2017;214: 1317–

1329.

63. Thompson RN, Morgan OW, Jalava K. Rigorous surveillance is necessary for

high confidence in end-of-outbreak declarations for Ebola and other infectious

diseases. Philos Trans R Soc B. 2019; doi:10.1098/rstb.2018.0431

64. Kraemer MUG, Faria NR, Reiner RC, Golding N, Nikolay B, Stasse S, et al.

Spread of yellow fever virus outbreak in Angola and the Democratic Republic of

the Congo 2015–16: a modelling study. Lancet Infect Dis. 2017;17.

doi:10.1016/S1473-3099(16)30513-8

65. Nishiura H, Chowell G. The effective reproduction number as a prelude to

statistical estimation of time-dependent epidemic trends. Mathematical and

Statistical Estimation Approaches in Epidemiology. 2009. doi:10.1007/978-90-

481-2313-1_5

66. Delamater PL, Street EJ, Leslie TF, Yang YT, Jacobsen KH. Complexity of the

basic reproduction number (R0). Emerg Infect Dis. 2019;25.

doi:10.3201/eid2501.171901

67. Anderson RM. The concept of herd immunity and the design of community-based

immunization programmes. Vaccine. 1992;10: 928–935. doi:10.1016/0264-

410X(92)90327-G

68. Hart WS, Hochfilzer LFR, Cunniffe NJ, Lee H, Nishiura H, Thompson RN.

Accurate forecasts of the effectiveness of interventions against Ebola may require

models that account for variations in symptoms during infection. Epidemics. 2019;

69. Meakin SR, Tildesley MJ, Davis E, Keeling MJ. A metapopulation model for the

2018 Ebola outbreak in Equateur province in the Democratic Republic of the

Congo. bioRxiv. 2018; 1–30.

70.   Chowell G, Nishiura H. Characterizing the transmission dynamics and control of

954   Ebola Virus Disease. PLoS Biol. 2015;13: 1002057.

955   doi:10.1371/journal.pbio.1002057

956   71.   Chowell G, Hengartner NW, Castillo-Chavez C, Fenimore PW, Hyman JM. The

957   basic reproductive number of Ebola and the effects of public health measures:

958   The cases of Congo and Uganda. J Theor Biol. 2004;229: 119–126.

959   doi:10.1016/j.jtbi.2004.03.006

960   72.   Thompson RN, Hart WS. Effect of confusing symptoms and infectiousness on

961   forecasting and control of Ebola outbreaks. Clin Infect Dis. 2018;67.

962   doi:10.1093/cid/ciy248

963   73.   Thompson RN, Cobb RC, Gilligan CA, Cunniffe NJ. Management of invading

964   pathogens should be informed by epidemiology rather than administrative

965   boundaries. Ecol Modell. 2016;324. doi:10.1016/j.ecolmodel.2015.12.014

966   74.   Cunniffe NJ, Cobb RC, Meentemeyer RK, Rizzo DM, Gilligan CA. Modeling when,

967   where, and how to manage a forest epidemic, motivated by sudden oak death in

968   California. Proc Natl Acad Sci USA. 2016;113: 5640–5645.

969   doi:10.1073/pnas.1602153113

970