A cross-scale assessment of productivity-diversity relationships

Dylan Craven[1,2,3,4*], Masha T. van der Sande[3,4,5,6,7], Carsten Meyer[4,8,9], Katharina Gerstner[4,8], Joanne M. Bennett[4,10], Darren P. Giling[4,8,11], Jes Hines[4,8], Helen R. P. Phillips[4,8], Felix May[4,12], Katherine H. Bannar-Martin[4,13], Jonathan M. Chase[4,14] & Petr Keil[4,14]

[1] Centro de Modelación y Monitoreo de Ecosistemas, Facultad de Ciencias, Universidad Mayor, Santiago, Chile
[2] Biodiversity, Macroecology & Biogeography, Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Büsgenweg 1, 37077 Göttingen, Germany
[3] Department of Community Ecology, Helmholtz Centre for Environmental Research – UFZ, 06120 Halle (Saale), Germany
[4] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany
[5] Department of Biological Sciences, Florida Institute of Technology, Melbourne, FL, USA
[6] Institute for Biodiversity & Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands
[7] Forest Ecology and Forest Management Group, Wageningen University and Research, Wageningen, the Netherlands
[8] Institute of Biology, Leipzig University, Leipzig, Germany
[9] Institute of Geosciences and Geography, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany
[10] Institute of Biology, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany
[11] Institute of Ecology and Evolution, Friedrich Schiller University Jena, Jena, Germany
[12] Methodology Centre and Institute of Ecology, Leuphana University Lüneburg, Lüneburg, Germany
[13] Quantitative Assessment Methods Section, Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, BC, Canada
[14] Department of Computer Sciences, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

* Correspondence: dylan.craven@aya.yale.edu

Keywords: spatial grain, biomass, carbon sequestration, climate, machine learning, more individuals hypothesis, species-energy

Abstract

Biodiversity and ecosystem productivity vary across the globe and considerable effort has been made to describe their relationships. Biodiversity-ecosystem functioning research has traditionally focused on how experimentally controlled species richness affects net primary productivity (S→NPP) at small spatial grains. In contrast, the influence of productivity on richness (NPP→S) has been explored at many grains in naturally assembled communities. Mismatches in spatial scale between approaches have fostered debate about the strength and direction of biodiversity-productivity relationships. Here we examine the direction and strength of productivity's influence on diversity (NPP→S) and of diversity's influence on productivity (S→NPP), and how this varies across spatial grains using data from North American forests at grains from local (672 m^2) to coarse spatial units (median area = 35,677 km^2). We assess relationships using structural equation and Random Forest models, while accounting for variation in climate, environmental heterogeneity, management, and forest age. We show that relationships between S and NPP strengthen with spatial grain. Within each grain, S→NPP and NPP→S have similar magnitudes, meaning that processes underlying S→NPP and NPP→S either operate simultaneously, or that one of them is real and the other is an artifact. At all spatial grains, S was one of the weakest predictors of forest productivity, which was largely driven by biomass, temperature, and forest management and age. We conclude that spatial grain mediates relationships between biodiversity and productivity in real-world ecosystems and that results supporting predictions from each approach (NPP→S and S→NPP) serve as an impetus for future studies testing underlying mechanisms.

Significance statement

The relationships between diversity and productivity are central to efforts explaining global variation of biodiversity and rates of carbon sequestration. However, little is known about the relative importance of biodiversity as the driving force, or as the consequence of ecosystem-level productivity. Our analysis of a comprehensive database of North American forests reveals that biodiversity and productivity can be both cause and effect and that their relationship strengthens with spatial grain. Importantly, we show that environmental context is more important in determining biodiversity and productivity than either biodiversity or productivity alone. Productivity-diversity relationships emerge at multiple spatial grains, which should widen the focus of national and global policy and research to larger spatial grains.

84 Introduction

One of the most prominent questions in ecology is how to describe relationships between biodiversity and

ecosystem-level productivity (1–10). Two fields of research with different motives have tried to

understand causality between these variables (11). The first examines how biodiversity varies across

space as a result of different levels of productivity created by environmental variation (e.g., temperature,

precipitation), and has resulted in a voluminous literature on the shapes of the patterns and their potential

underlying causality (1, 2, 9, 10, 12–14). The second aims to quantify changes in vital ecosystem

functions such as productivity following anthropogenically-induced changes in diversity (5, 15–17). As a

result of the different perspectives on the direction of causality, there remains considerable debate and

confusion surrounding the relationship between diversity and productivity (18), which is exacerbated by

differing spatial grains at which studies are conducted (4, 19).

Recently, there has been growing interest in assessing biodiversity ecosystem functioning (BEF)

relationships in real-world, non-experimental ecosystems over large geographic extents, but likely due to

logistical constraints, relationships are typically measured at local spatial grains (20–22). Results suggest

that the positive effect of species richness on productivity and other ecosystem functions can be as, or

more, important than abiotic environmental drivers' effects on productivity, suggesting that diversity-

productivity relationships can be even stronger in real-world communities than in controlled experiments

(21). However, to fully understand the influence of diversity on productivity, and vice versa, it is critical

to recognize that traditional bivariate analyses may underestimate the strength of these relationships by

not accounting for the effects of spatial grain, as well as those of biomass, shading, macro-climate, and

management (11, 18, 23, 24).

The striking mismatch between the spatial grains of BEF experiments (cm^2 to m^2; 4), observational

studies of BEF (0.04 to 1.0 ha; 20, 25), and macroecological diversity-productivity correlations (m^2 to

thousands of km^2; 1, 10, 26, 27) further obscures comparisons between perspectives. However, there is a

diverse array of theoretical expectations for grain dependency of the effects of productivity on diversity

(NPP→S) and of diversity on productivity (S→NPP), which predict effects to either strengthen or weaken

as the spatial grain increases (Table 1). For example, spatial turnover of species that are functionally

equivalent within the regional grain can offset low species richness at local grains, resulting in a

strengthening of S→NPP with increasing spatial grain. The effects of NPP→S are also hypothesized to

increase with spatial grain, because higher NPP is associated with greater heterogeneity at larger spatial

grains, which enhances coexistence of more species at the regional grain. Moreover, other components of

a community, such as biomass, can mediate relationships between productivity and diversity via their

effects on competitive dominance (18). These theoretical expectations have been supported by

observational data for the effects of productivity on diversity (10, 28, 29). In the case of BEF relationships

(i.e. S→NPP), there is also empirical and theoretical support for grain dependence, which comes from a

restricted range of small spatial grains (30–32).

Here, we aim to address the dual nature by which productivity influences diversity (NPP→S) and

diversity influences on productivity (S→NPP) across spatial grains by combining structural equation

models (SEM) and Random Forest models (RFs) to explicitly account for the bidirectionality of NPP→S

and S→NPP. Using SEM, we propose and test hypothesis-based models (Fig. S2) that estimate the

direction and strength of NPP→S and S→NPP. Next, we use RFs, an assumption-free machine learning

approach (33), to quantify the relative importance of predictors of species richness and productivity. We

examine both hypothesized directions of the relationship, along with a number of important covariates

that influence both diversity and productivity, such as biomass, precipitation, temperature, and forest age,

using a comprehensive observational dataset of North American forests at fine (area = 672 m^2; n =

46,211 plots), medium (median area = 1,386 km^2; n = 1,956 spatial units), and coarse spatial grains

(median area = 35,677 km^2; 98 spatial units). We specifically ask whether the influence of productivity

(and its environmental drivers) on diversity (NPP→S) played a predominant role, or whether the

influence of diversity on productivity (S→NPP) was predominant, and how these relationships manifest

across grains in real-world ecosystems.

Table. 1 Overview of hypotheses predicting grain dependence of relationships between net primary productivity (NPP) and species richness (S).

No.	Direction	Mechanism of grain dependence	Weakens or strengthens towards coarse grain?	Reference
I	NPP→S and S→NPP	Spatially asynchronous demographic stochasticity impacts small populations (or small grains) and averages out over large grains.	Both NPP→S and S→NPP strengthen towards coarse grains	(34)
II	NPP→S	At larger grains, higher NPP is associated with increased heterogeneity and/or dissimilarity of local patches, allowing for greater regional coexistence.	NPP→S strengthens towards coarse grains	(28, 35, 36)
III	NPP→S	A statistical interaction between NPP and grain in their effect on S emerges as a consequence of increasing occupancy with NPP.	NPP→S weakens towards coarse grains	(37)
IV	NPP→S	At very large grains (thousands of km^2 and larger), high productivity increases occupancy and population size, thus increasing the probability of reproductive isolation and speciation	NPP→S strengthens towards coarse grains	(38)
V	S→NPP	Stochastic sampling effects dominate at small grains, resource partitioning at larger grains ('spatial insurance'), and their relative magnitude determines the grain dependency.	Both strengthening or weakening possible	(39, 40)
VI	S→NPP	Functionally redundant species at the regional grain can compensate for low richness at local grains.	S→NPP strengthens towards coarse grains	(41)
VII	S→NPP	With incomplete compositional turn-over, proportional changes in larger-grain richness are always less than proportional changes in smaller-grain richness such that the explanatory power of richness on changes in functioning decreases with spatial scale.	S → NPP strengthens strengthens towards coarse grains until species richness saturates	(42)

Fig.1. Maps of species richness (S), net primary productivity (NPP) [gC/m^2/year], and biomass [Mg/ha] of forests at three spatial grains across the contiguous USA. The values in all plots are on log$_{10}$ scale.

Results

Spatial patterns in productivity (NPP) and richness (S) emerged at coarser spatial grains, with higher S and NPP usually observed in the eastern USA than in the western USA (Fig. 1). Biomass, a time-integrated measure of NPP that also influences diversity, also exhibited similar patterns (Fig. 1). Bivariate relationships among S and NPP exhibited scale dependence, with their slopes increasing with spatial grain (Fig. 2).

Fig. 2. Bivariate relationships between observed species richness (S) and productivity (NPP) of forests at three spatial grains across the contiguous USA, on linear (A, C) and log_{10} (B, D) scales. Panels A and B show NPP as a response to S, panels C and D show NPP as a predictor. Solid lines are least-squares linear regressions fitted at each grain, shaded areas are standard errors. Analyses were performed using stratified random samples of 1000, 500 and 98 spatial units at the fine, intermediate and coarse spatial grains, respectively.

Structural Equation Models (SEM). We examined relationships between species richness and net primary productivity (NPP) across spatial grains using two SEMs for each spatial grain: the first (S→NPP) testing the direct effect of S on NPP and the indirect effect of NPP on S (via biomass), and the second (NPP→S) testing both the direct and indirect effects of NPP on S (Fig. 3). In both SEMs, environmental variables (e.g., mean annual precipitation (MAP), mean annual temperature (MAT),

temperature seasonality, and elevation range), size of the species pool, forest age, and management were

used to explain variation in S, biomass, and NPP. At the intermediate and coarse grains, we also included

area (of each spatial unit) to account for variation in species richness due to the effects of area (see

Methods).

Fig. 3. Structural equation models (SEM) testing the influence of diversity (S) on productivity (NPP)('S
→ NPP'; A, B, C) and that of NPP on S ('NPP → S'; D, E, F), once controlling for environmental
variables (e.g., mean annual precipitation, mean annual temperature, temperature seasonality, and
elevation range), size of the species pool, forest age, and management, in forests across the contiguous
USA at three spatial grains. Both models fit the data well at all spatial grains (P-value of the Chi-square
test > 0.1; Table S1). Boxes represent measured variables and arrows represent relationships among
variables. Solid blue and red arrows represent significant (P< 0.05) positive and negative standardized
path coefficients, respectively, and their width is scaled by the corresponding standardized path
coefficient. Solid and dashed gray arrows represent non-significant (P>0.05) positive and negative
standardized path coefficients, respectively. AGE is forest age, MANAGED is forest management,
ANN.PREC is mean annual precipitation, ANN.TEMP is mean annual temperature, TEMP.SEAS is
temperature seasonality, ELEV.RANGE is elevation range, S.POOL is the regional species pool, and
AREA is area. S, BIOMASS, NPP, and AREA were natural log transformed prior to analysis.

Fig. 4. Direct effects of diversity on productivity (S → NPP) and productivity on diversity (NPP → S)
estimated with structural equation models (SEM) in forests across the contiguous USA at three spatial
grains. Points are standardized path coefficients and solid lines are 95% confidence intervals.

Both models fit the data well for all spatial grains (P-value of the Chi-square test > 0.1; Table S1). At

each spatial grain, both SEMs had similar R^2 values averaged over S, biomass and NPP, indicating a

similar fit of the model to the data. R^2 values for both SEMs increased with spatial grain, from 0.34 at the

fine grain, to 0.56 at the intermediate grain and 0.85 at the coarse grain. Generally, the strength of effects

of S → NPP and NPP → S were similar within each spatial grain, but both increased in strength with

increasing spatial grain (Fig. 3 & 4). At the fine spatial grain, we found a weak direct effect of S → NPP

(Fig. 3A) and NPP → S (Fig. 3D), but an indirect effect of NPP on S via biomass (standardized path

coefficient of indirect effect = 0.23; Fig. 3D). At the intermediate spatial grain, we found a similarly

strong direct effect of S on NPP (standardized path coefficient of direct effect = 0.11, Fig. 3B and 4) as

NPP on S (standardized path coefficient of direct effect = 0.24; Fig. 3E and 4) and weak indirect effects

of NPP on S (standardized path coefficient of indirect effect = 0.04; Fig. 3E). Similarly at the coarse

spatial grain, we found strong direct effects of S on NPP (0.42, Fig. 3C and 4) and of NPP on S (0.47, Fig.

3F and 4) and weak indirect effects of NPP on S (standardized path coefficient of indirect effect = 0.08;

Fig. 3F).

Overall, the SEMs show that the productivity-diversity relationship increases in strength with spatial

grain, and both relationships (S→NPP and NPP→S) explain similar amounts of variation, albeit with

some differences in the direct and indirect effects. At fine spatial grains, our SEMs show greater support

for a strong indirect effect of NPP on S via biomass, but do not support the inverse effect of S on NPP.

Towards coarser spatial grains, our SEMs do not conclusively show stronger support for one direction of

causality over the other. These patterns were robust to the direction of paths between S and biomass (Fig.

S3A, B, and C; Table S1) and between biomass and NPP (Fig. S3D, E, and F; Table S1).

Random Forest models (RFs). To assess the relative importance of each predictor of species richness

and NPP, and to provide an assumption-free alternative to the SEMs, we fitted two random forest models

for each of the three spatial grains: one with NPP and the other with S as response variables. We found

that NPP was an important predictor of S at the fine and intermediate spatial grains (Fig. 5A), with

unimodal and linear effects respectively (Fig. 5), but was less important relative to other predictors at the

coarse spatial grain. For S, we found that species pool, MAT, MAP, and forest age were the best

predictors at all spatial grains (Fig. 5). In line with the SEM analyses, the overall explained variation of S

increased towards coarse spatial grains, from 0.38 at the fine grain to 0.51 at the intermediate and 0.85 at

coarse grains. For NPP, we found that species richness was one of the weakest predictors relative to other

predictors at all spatial grains (Fig. 5B), with management, forest age, MAP, and especially biomass

being the most important predictors (Fig. 5B). The overall explained variation of NPP also increased

towards coarse spatial grains, from 0.62 at the fine spatial, to 0.82 at the intermediate spatial grain and

0.83 at the coarse spatial grain.

222
223
Fig. 5. Relative variable importance from random forest models explaining NPP and species richness (S)
and NPP at three spatial grains, which is the mean decrease in squared error caused by each of the
variables, rescaled such that it sums up to the total pseudo R^2. The curves in the insets show shapes of the
partial response of ln NPP or ln S (after accounting for all of the covariates); all of the insets have the
same span of the y-axis. For complete responses with axis scales and tick-marks see Fig. S5.

Discussion

The first important result is the similar magnitude of the S→NPP (18) and NPP→S (10, 26, 43)

relationships at all grains. This reflects, in part, that both productivity and species richness have many

environmental and geographical drivers in common (44), which complicates distinguishing correlation

from causation, even when using SEMs (45, 46). There are two possible interpretations of this result: (i) it

may indicate that diversity's causal effects on productivity and productivity's causal effects on diversity

operate simultaneously, which was suggested by (18), but never demonstrated on observational data from

large spatial grains. Alternatively (ii), if only one direction of the diversity-productivity relationship is

real and causal, it may be possible to fit another model assuming the opposite direction because of

multicollinearity in the data or non-identifiability of the causal direction (47). Without large-grain

experiments that manipulate diversity in ways that mimic biodiversity change (i.e. species gains and

losses) in real-world ecosystems (11, 48–50), we see little hope for resolving this with contemporary data

and approaches.

Our second important result is that both S→NPP and NPP→S strengthen from the fine to the intermediate

grain, and in the case of the SEM both relationships continue strengthening towards the coarsest grain.

While grain-dependent shifts are often expected (Table 1), this had not been shown previously with

empirical data for S→NPP using spatial grains coarser than several hectares (25, 31, 32). If the S→NPP

direction is the real causal one, then our results from SEM and RF analyses support several theoretical

expectations (Table 1) and give further impetus to efforts quantifying biodiversity effects in naturally

assembled ecosystems at broad spatial scales (51). If the NPP→S direction is the real causal one, then our

results are in line with (27, 35), but are in contrast with (29, 37), particularly when upscaling from the fine

grain to intermediate grain, where both the SEM and RF analyses give congruent results. Intriguingly, a

third possibility is that both NPP→S and S→NPP are real and that they operate simultaneously, as

suggested by our SEM results. In this case, we are unaware of any theory that considers how this

reciprocal relationship would be expected to change with increasing spatial grain. The one caveat

applicable to interpreting any direction of diversity-productivity relationships is that of demographic

stochasticity (mechanism I in Table1), which may weaken both NPP→S and S→NPP, or their synergistic

interplay, at fine spatial grains. In our study, the strong local effect of demographic stochasticity appears

plausible given the small area of the forest plots (0.067 ha) and small population sizes (12.24 ± 0.02 trees

per plot; range = 1- 157 trees per plot) therein. This would suggest that temporal changes in local scale

biodiversity (52, 53) may have under-appreciated effects on ecosystem function (54).

The third key result is that other predictors, such as temperature and biomass, were particularly influential

in all our analyses. That is, the grain dependence of the relationship between S and NPP was coupled with

a clear increase in the effect of annual temperature (but not precipitation) on both S and NPP towards

coarse grains, which supports the notion that either temperature-dependent diversification (55, 56) or

ecological limits (43) shape diversity at these spatial grains. The consistently weak effect of precipitation

is expected since we focus on forests, which only grow above certain precipitation thresholds (57).

Second, we found a positive, indirect effect of NPP on species richness via forest biomass at the fine

spatial grain, which supports multiple hypotheses (Table 1) such as the view that higher ecosystem

productivity enhances species diversity by enabling larger numbers of individuals per species to persist

due to lower extinction rates (35, 37, 58), particularly at fine grains where stochastic extinctions occur

(34). The clear importance of temperature, biomass, and other predictors such as forest age, seasonality,

or species pool (Figs. 3, 4) highlights that even when the NPP→S relationship holds across grains, other

drivers are considerably more important in predicting both (e.g., 59). Hence, integrating the

environmental context surrounding ecological communities into modeling diversity-productivity

relationships is a necessary step towards making robust predictions of either biodiversity or ecosystem

functioning at any spatial grain.

Our results reveal that mechanisms associated with one direction of diversity-productivity relationships

may provide insight to observed patterns of either direction, despite being initially formulated at a

different spatial grain. For example, the positive, indirect effect of NPP on species richness via forest

biomass at the fine spatial grain provides support for the more individuals hypothesis (37), although it is

typically tested at regional to continental spatial scales. Increasingly, macroecological mechanisms such

as speciation gradients (60) and water-energy variables are being examined in small-grain experimental

grasslands to explore their role in mediating niche-based processes (61) and biodiversity effects (62),

respectively. Similarly, efforts to upscale biodiversity effects on productivity - developed initially to

identify local scale mechanisms such as complementarity and selection effects (63, 64) - may emerge as a

crucial mechanism in determining spatial variation in ecosystem functioning at large spatial scales. Rather

than uniquely focusing on the direction and strength of S-NPP once accounting for other factors, our

results show that mechanisms associated with S→NPP and NPP→S likely underpin the context

dependency of diversity-productivity relationships across spatial grains (Table 1). These recent

developments in BEF research and macroecology suggest that conceptual integration between these two

disciplines is just beginning (65), yet further efforts to bridge disciplinary gaps are essential to deepen

current understanding of mechanisms that underpin the shifts in diversity-productivity relationships

across spatial scales.

To conclude, we show that the relationship between diversity and productivity strengthens toward coarse

grains. This result is in line with expectations from both BEF theory, and some (but not all) expectations

from macroecological studies on NPP→S, and highlights the potential of demographic stochasticity to

distort diversity-productivity relationships at fine grains. Moreover, we find similar support for both

directions of diversity-productivity relationships across spatial grains, revealing that biodiversity and

productivity can be both cause and effect. Future research on this relationship needs to move from fine-

grain experiments and observational studies to coarse grains in order to fully understand and predict the

impacts of anthropogenic biodiversity change on ecosystem function.

Methods

Data

Geographic extent and grain. We conducted analyses across the contiguous USA at three spatial grains

(Fig. 1): (1) fine grain (46,211 plots, 672 m^2 or 0.000672 km^2 each), (2) intermediate grain (1956 units,

median 1,386 km^2) created by aggregating US counties to larger units based on the forested area within

them (see 'spatial aggregation algorithm' below), and (3) coarse grain (95 units, median 35,677 km^2)

created by further aggregating the intermediate grain units. We restricted our analyses to forested areas to

make comparisons within and among spatial grains in similar ecosystems. For the intermediate and coarse

grains, we defined an area as forested if it fell into a 1 km^2 pixel with non-zero forest biomass following

(66).

Spatial aggregation algorithm. To minimize variation in species diversity (defined below) due to

differences in area between spatial units within the intermediate and coarse spatial grains, we created an

algorithm that minimized variation of forested area at each spatial grain. The algorithm worked in an

iterative process as follows: (1) Calculate variance of area across all counties. (2) Randomly select a focal

county with a probability proportional to 1/sqrt(forest area+1), which will more likely select small rather

than large counties. This was because small counties can be merged to approach the grain of the large

counties, thus decreasing variation in area. (3) Select a random adjacent county and merge it with the

focal county. (4) Updated the variance of area across all spatial units in the USA and compare it to the

original variance. If the new variance is lower than the original variance, the algorithm accepts the

merged unit and returns to step one. If the variance does not decrease, the algorithm repeats step 3 until it

decreases, with the maximum number of attempts of 1,000. If the variance still does not decrease even

after 1,000 attempts, the algorithm rejects the merge, and returned to step one. The algorithm started with

3,107 US counties, and we first terminated it when 1,956 merged spatial units were created (intermediate

grain dataset) and then after it reached 98 merged spatial units (coarse grain dataset) (Fig. 1). Although

the algorithm substantially reduced variation in area within both spatial grains (Fig. S9), it did not

eliminate the variation entirely, and thus we still used area as a covariate in the statistical analyses at the

intermediate and coarse spatial grains.

Species richness (S). For all spatial grains, we estimated diversity as species richness (S) because it is the

most commonly used and best understood metric of biodiversity, although other measures of diversity

may be better predictors of net primary productivity (67–69). We extracted S at the fine spatial grain from

the Forest Inventory and Analysis National Program (FIA) database v. 1.7.0 (70). We restricted our

analysis to plots on forested land that were sampled using the national FIA design (plot design code 1)

(71). All plots were surveyed between 1998 and 2016 and contain four circular 168 m^2 sub-plots with a

total area of 0.067 ha in which all individuals larger than 12.7 cm diameter at 1.3 m were recorded and

identified to species level. In total, our final dataset included 344 woody species and 93,771 plots. To

estimate S at the intermediate and coarse spatial grains, we used data on the extant native species of trees

for the contiguous USA provided in The Biota of North America Program's (BONAP) North American

Plant Atlas (72).

Taxonomic harmonization of species names. We cleaned scientific names from the FIA data and the

BONAP data and harmonized them to accepted species based on The Plant List (73) and the Taxonomic

Name Resolution Service (TNRS)(74), following the protocol described in (75). We included hybrid

forms but excluded any names that could not be resolved to the species level.

Filtering of species occurrences. We restricted our analyses to woody species occurring in forest. To this

end, we initially filtered the BONAP data to species classified as 'trees' in BONAP's taxonomic query

database (72). We further filtered out 70 county-level occurrences of 5 non-woody species from the

BONAP dataset. Species woodiness was inferred from woodiness data (76) and species lists of trees,

shrubs and subshrubs (77), except for 37 species without such data for which we instead inferred

woodiness from online searches or assumed resemblance among congeneric species. We also filtered out

8 FIA plot-level species occurrences and 1595 BONAP county-level species occurrences that we deemed

unlikely to be forest occurrences, as inferred from independent species occurrences within forested pixels

recorded in FIA plots and Global Biodiversity Information Facility (GBIF) point-occurrence records

(Downloaded via https://www.gbif.org/ on 26th September 2016; doi:10.15468/dl.mka2y5;

Supplementary Note). To make species richness data internally consistent across the different spatial

grains, we added a further 6,593 quality-vetted county-level forest occurrences of woody species from

FIA plot records to the 282,991 occurrences in the taxonomically harmonized BONAP dataset.

Net primary productivity (NPP). For a large subset of plots in the FIA database that have been

measured at least twice (n = 46,211, on average plots remeasured every 5.8 years), we calculated tree

productivity. This was measured as the change in tree C over time due to growth ($gC\ m^{-2}\ y^{-1}$), and is the

sum of aboveground C increment of living trees between two measurements and conservatively excludes

recruits and dead trees (67). Tree-level carbon was estimated by multiplying tree-level biomass (see

below) by 0.48, but recognize that gymnosperms may have higher carbon content than that of

angiosperms (78). For plots with more than two inventories, tree productivity was calculated for each

period and then averaged.

At intermediate and coarse spatial grains, we calculated NPP using the MODIS-derived MOD17 A3

product (79, 80), which gives annual values of NPP as $gC\ m^{-2}\ yr^{-1}$ in 30 arc-sec pixels (roughly 1 km^2

around the equator). Here, NPP is defined as the annual sum of daily net photosynthesis minus the cost of

growth and maintenance of living cells in permanent woody tissue. We averaged the annual values from

2000 to 2015 for each pixel, and then averaged these across the intermediate and coarse grains.

Biomass. At the fine scale, we estimated tree-level biomass using generalized biomass equations

developed for North American tree species (81). For each plot we calculated aboveground biomass (Mg

ha^{-1}) as the sum of individual biomass of living trees per hectare.

For the intermediate and coarse spatial grains we derived biomass values using a map of aboveground

forest biomass of the USA, which is derived by modeling FIA plot biomass as a function of geospatial

predictor variables (66). This data layer had a grain of 250 x 250 m^2, therefore, the average within each of

the intermediate- and coarse-grain spatial units was taken.

Forest age and management. For each plot in the fine-scale dataset, we extracted forest age and

management history from the FIA data set. Forest age is estimated using dendrochronological records

(71). Management regime was a binary variable that indicated whether any forest management activity,

e.g. harvest, thinning, tree planting, had been observed in any inventory or not.

At the intermediate and coarse grain, forest age was calculated as the average forest age from NASA

NACP 1 km^2 resolution layer (82). Management regime at the intermediate and coarse grains was

calculated as the proportion of managed FIA plots within all FIA plots that were within each spatial unit.

Climatic variables. For all grains, we used WorldClim (83) bioclimatic variables at 30 sec resolution.

Many of the WorldClim variables are strongly collinear with one another, or with other variables in the

analysis (Table 1). Thus, only three variables that captured different aspects of the climate were selected;

mean annual temperature (BIO1), mean precipitation (BIO12), and temperature seasonality (BIO4). At

the fine scale, for each FIA plot we extracted the values of the 30 sec pixel in which the plot was found.

For the intermediate and coarse grains, we averaged the values across all pixels within each spatial unit.

Elevation range. We used elevation range as a proxy for topographic and habitat heterogeneity, a

variable that has been shown to be a good predictor of species richness (84). The USGS SRTM1 dataset

(85) with 1 sec (approx. 30 x 30 m^2) resolution was used for all spatial grains. At the fine-scale, we

calculated a 250 m diameter buffer around each FIA plot and calculated the elevation range using all 1 sec

SRTM pixels within the buffer. At the intermediate and coarse scale, elevation range was calculated as

the difference between the minimum and maximum elevation points within each spatial unit.

Species pools. We calculated regional species pools for each spatial grain as probabilistic dispersal pools

(86). For each intermediate-grain spatial unit and each species in our data set, we first estimated the

species' probability of being part of the unit's species pool as the joint probabilities that dispersal might

happen between that unit and any of the species' intermediate-grain occurrences within the contiguous

US. Due to insufficient data on species' dispersal abilities, we assumed that dispersal probability between

focal units and species' occurrences would decay with great-circle distance between the respective

regions' centroids. We explored five alternative exponential distance-decay functions, with scaling

coefficients that determined the probability of a species occurring in neighboring units would disperse to

the focal unit of 0.975, 0.95, 0.90, 0.80, and 0.60. We chose the function whose resulting species pool

estimate most strongly correlated with species richness. Finally, we calculated species pools for each

spatial unit as the sum of all species' individual probabilities of dispersal from any of their respective

occurrences. For each coarse-grain unit, we summed the species' joint probabilities of dispersal between

any of their intermediate-grain occurrences and any of the intermediate-grain units nested within the

coarse unit. For fine-grain units, we assumed that their species pools would equal those of the

intermediate-grain spatial units in which they were nested.

All of the variables used in our analyses are listed and summarized in Table 2.

417   **Table 2.** Overview of all variables used in the SEM and RFs.

Variable	Abbreviation	Units	The grain at which data are used	Source, reference, or link	Layer resolution, if applicable
Species richness	S	# of species	fine	FIA	
			intermediate, coarse	BONAP	
Net Primary Productivity	NPP	gC/m^2/year	fine	FIA	
			intermediate, coarse	MODIS (79)	1 km^2
Biomass	BIOMASS	Mg/ha	fine	FIA	
			intermediate, coarse	(66)	1 km^2
Forest age	AGE	years	fine	FIA	
		years	intermediate, coarse	NACP (81)	1 km^2
Management regime	MGMT	managed or not	fine	FIA	
		proportion of managed FIA plots	intermediate, coarse	FIA	
Mean annual precipitation	MAP	average mm/y	all grains	WorldClim	1 km^2
Mean annual temperature	MAT	average □/y	all grains	WorldClim	1 km^2
Temperature seasonality	TEMP.SEAS	standard deviation of monthly □ * 100	all grains	WorldClim	1 km^2
Elevation range (altitudinal span)	ELEV.RANGE	m	all grains	SRTM v 2.1	30 arc-sec
Size of species pool	S.POOL	# of species	all grains	FIA, GBIF, BONAP	

418
Stratified random sampling. Large areas of the contiguous US are environmentally homogeneous, while

other parts are environmentally unique and small. We employed stratified random sampling (87) for the

fine and intermediate spatial grains in order to (1) enhance environmental representativeness of the data,

(2) prevent excessive statistical leverage of the large number of data points from homogeneous areas and

(3) reduce spatial pseudoreplication (autocorrelation) by increasing the geographic distance between data

points. We first identified 11 strata at the fine and intermediate grains respectively, using multivariate

regression trees with S, NPP and biomass as response variables and all covariates as predictors (Fig. 1).

We then took a random and proportionally sized sample of spatial units from each strata (fine grain, N =

1,000; intermediate grain, N = 500). We did not use stratified random sampling at the coarse spatial grain

because the number of spatial units was small (N = 98) and spatial autocorrelation was low. The spatial

locations of the stratified samples are in Fig. S1. All of the analyses presented here, as well as our main

conclusions, are based on these stratified sub-samples of the data.

Data transformation and standardization. Prior to analysis, species richness, biomass, NPP, and area

were natural-log transformed.

Data Analyses

Two complementary statistical approaches were used. First, we fitted structural equation models (SEMs),

which allow the assessment of indirect effects including feedback loops, address causality, and take into

account potential collinearity among covariates (45, 46). The paths in our candidate SEMs were based on

previous evidence of causal links between S, biomass, and NPP (Figure S2; 18). Second, to better

understand the relative importance of each variable in explaining variation in the response variables

within models, we fitted Random Forest models (RFs) (33). The results from SEMs provide insight into

differences among models (i.e. between the two causal pathways per spatial grain, and among spatial

grains), while results from RFs provide additional insights into the relative importance of different

predictors variables within models.

Structural Equation Modelling (SEM). To test the relative importance of S on NPP ('S→NPP') and

NPP on S ('NPP→S') we fitted two SEMs per spatial grain. For each SEM, we started with a 'saturated'

model, which included the relationships between S, NPP, and biomass, and relationships of all additional

covariates on S, NPP, and biomass (except for area at the fine spatial grain) (Figure S2). An effect of area

on NPP and biomass was included because in many models this proved necessary to obtain an accepted

model. The S→NPP model evaluated how S directly affects NPP and how NPP indirectly affects S via

biomass and, therefore, included a feedback loop. The NPP→S model tested the direct effect of NPP on

richness and, unlike the S→NPP model, did not include a feedback loop. This way, we tested the direct

effect of S on NPP (S→NPP model), the direct effect of NPP on S (NPP→S), and the indirect effect of

NPP on S (included in both models).

To test the sensitivity of the relative importance of S on NPP to the direction of the relationships between

S and biomass ('S → Biomass') or biomass and NPP ('Biomass → NPP', we fitted two additional SEMs

per spatial grain as described above. The S → Biomass model evaluated how S directly affects NPP and

biomass and how S indirectly affects biomass via NPP. The Biomass → NPP model tested the direct

effect of S on NPP and the direct effect of biomass on S. Neither Biomass → NPP or S → Biomass

models includes an indirect effect of NPP on S, as was the case for the S → NPP or NPP → models.

Model fit can only be tested on unsaturated models, i.e. those that have at least one missing path.

Therefore, we removed the path with the lowest standardized path coefficient from the model. As SEMs

had equal number of paths, we could compare model fit across all models within each spatial grain using

their unadjusted R^2 values. After excluding the additional paths, path coefficients of S, NPP, and biomass

remained qualitatively the same, and model fit to the data were still accepted (Chi-square test; P>0.05).

This indicates that the models are identifiable and their results are robust. Therefore, we did not further

reduce the model, and models maintained the same number of paths within each scale.

To assess the differences among scales in the relationships between S, NPP and biomass for each model,

we compared the standardized regression coefficients using their 95% confidence intervals. All SEMs

were fitted using the 'sem' function of the 'lavaan' package in R (88).

Random Forest models (RFs). To assess the relative importance of each variable in predicting the

response variables within models, we used Random Forest models (RFs) (33, 89, 90). We used the

'randomForest' function in the R package 'randomForest', with all RF models produced using the default

settings: 500 trees, one third of predictors sampled in each tree, sampling with replacement of the entire

dataset, and terminal node size of 5.

At each of the three spatial grains we fitted two RFs, one with S as a response variable and the other with

NPP as a response variable. All predictors that were used in the SEMs were used in the RF models

(including biomass). To quantify the relative importance of each predictor, we calculated the mean

decrease of squared error across all 500 trees using the function 'importance'. The importances were then

scaled between 0 and 1, with 1 being the most important predictor. Using the function 'partialPlot', we

extracted the partial responses of S and NPP to visualize the relationship between the two variables after

accounting for all other covariates.

Reproducibility. All data on species richness, biomass, NPP, covariates, and R code used for the data

processing and analyses are available on Figshare (DOI: 10.6084/m9.figshare.5948155) under a CC-BY

license.

Acknowledgments

All authors recognize support from the German Centre for Integrative Biodiversity Research (iDiv) Halle-

Jena-Leipzig (DFG-FZT 118). MvdS is supported by the Rubicon research programme with project

number 019.171LW.023, which is financed by the Netherlands Organisation for Scientific Research

(NWO). CM acknowledges funding from the Volkswagen Foundation through a Freigeist Fellowship. We

thank John Kartesz and Misako Nishino for generously providing access to BONAP data. We thank

David Currie and Antonin Machac for initial discussions and Christian Wirth, Katie Barry, Nico

Eisenhauer, Stan Harpole, Miguel Mahecha, and the CAFE discussion group for their suggestions to

improve analyses.

Author contributions

494 Conceived project and designed research: DC, MvdS, KBM, JMB, JC, KG, DG, JH, FM, CM, PK.

495 Compiled data: DC, KG, CM, FM, PK. Analyzed data: MvdS, PK. Wrote the paper: All authors.

References

1. Adler PB, et al. (2011) Productivity Is a Poor Predictor of Plant Species Richness. *Science*
333(6050):1750–1753.
2. Currie DJ (1991) Energy and large-scale patterns of animal-and plant-species richness. *The*
American Naturalist 137(1):27–49.
3. Balvanera P, et al. (2006) Quantifying the evidence for biodiversity effects on ecosystem
functioning and services. *Ecology Letters* 9:1146–1156.
4. Cardinale BJ, et al. (2011) The functional role of producer diversity in ecosystems. *American*
Journal of Botany 98(3):1–21.
5. Cardinale BJ, et al. (2012) Biodiversity loss and its impact on humanity. *Nature* 486:59–67.
6. Hooper DU, et al. (2012) A global synthesis reveals biodiversity loss as a major driver of ecosystem
change. *Nature* 486(7401):105–108.
7. Naeem S, Duffy JE, Zavaleta E (2012) The Functions of Biological Diversity in an Age of
Extinction. *Science* 336(6087):1401–1406.
8. Tilman D, Isbell F, Cowles JM (2014) Biodiversity and Ecosystem Functioning. *Annu Rev Ecol*
Evol Syst 45(1):471–493.
9. Rosenzweig ML (1995) *Species diversity in space and time* (Cambridge University Press).
10. Mittelbach GG, et al. (2001) What Is the Observed Relationship Between Species Richness and
Productivity? *Ecology* 82(9):2381–2396.
11. Loreau M, et al. (2001) Biodiversity and Ecosystem Functioning: Current Knowledge and Future
Challenges. *Science* 294(5543):804–808.
12. Connell JH, Orias E (1964) The Ecological Regulation of Species Diversity. *The American*
Naturalist 98(903):399–414.
13. Waide RB, et al. (1999) The Relationship Between Productivity and Species Richness. *Annual*
Review of Ecology and Systematics 30(1):257–300.
14. Fraser LH, et al. (2015) Worldwide evidence of a unimodal relationship between productivity and
plant species richness. *Science* 349(6245):302–305.
15. Schulze E, Mooney H (1993) *Biodiversity and Ecosystem Function,* (Springer, Berlin).
16. Tilman D (1999) The ecological consequences of changes in biodiversity: a search for general
principles. *Ecology* 80(5):1455–1474.
17. Isbell F, et al. (2017) Linking the influence and dependence of people on biodiversity across scales.
Nature 546(7656):65–72.
18. Grace JB, et al. (2016) Integrative modelling reveals mechanisms linking productivity and plant
species richness. *Nature* 529(7586):390–393.
19. Whittaker RJ (2010) Meta-analyses and mega-mistakes: calling time on meta-analysis of the species
richness–productivity relationship. *Ecology* 91(9):2522–2533.
20. Liang J, et al. (2016) Positive biodiversity-productivity relationship predominant in global forests.
Science 354(6309):aaf8957.
21. Duffy JE, Godwin CM, Cardinale BJ (2017) Biodiversity effects in the wild are common and as
strong as key drivers of productivity. *Nature* 549(7671):261–264.
22. van der Plas F (2019) Biodiversity and ecosystem functioning in naturally assembled communities.
Biological Reviews 0(0). doi:10.1111/brv.12499.

23. Oberle B, Grace JB, Chase JM (2009) Beneath the veil: plant growth form influences the strength of
species richness–productivity relationships in forests. *Global Ecology and Biogeography*
18(4):416–425.
24. Cardinale BJ, Bennett DM, Nelson CE, Gross K (2009) Does productivity drive diversity or vice
versa? A test of the multivariate productivity–diversity hypothesis in streams. *Ecology* 90(5):1227–
1241.
25. Chisholm RA, et al. (2013) Scale-dependent relationships between tree species richness and
ecosystem function in forests. *Journal of Ecology* 101(5):1214–1224.
26. Hawkins BA, et al. (2003) Energy, water, and broad□scale geographic patterns of species richness.
Ecology v. 84(12):3105–3117.
27. Field R, et al. (2009) Spatial species-richness gradients across scales: a meta-analysis. *Journal of
Biogeography* 36(1):132–147.
28. Chase JM, Leibold MA (2002) Spatial scale dictates the productivity–biodiversity relationship.
Nature 416(6879):427–430.
29. Belmaker J, Jetz W (2011) Cross-scale variation in species richness–environment associations.
Global Ecology and Biogeography 20(3):464–474.
30. Chalcraft DR (2013) Changes in ecological stability across realistic biodiversity gradients depend
on spatial scale. *Global Ecology and Biogeography* 22(1):19–28.
31. Luo W, Liang J, Gatti RC, Zhao X, Zhang C Parameterization of biodiversity–productivity
relationship and its scale dependency using georeferenced tree-level data. *Journal of Ecology* 0(ja).
doi:10.1111/1365-2745.13129.
32. Hao M, Zhang C, Zhao X, Gadow K von (2018) Functional and phylogenetic diversity determine
woody productivity in a temperate forest. *Ecology and Evolution* 8(5):2395–2406.
33. Hastie T, Tibshirani R, Friedman JH (2009) *The elements of statistical learning: data mining,
inference, and prediction* (New York, NY: Springer). 2nd Ed.
34. Lande R, Engen S, Saether B-E (2003) *Stochastic population dynamics in ecology and conservation*
(Oxford University Press).
35. Wright DH, Currie DJ, Maurer BA (1993) Energy supply and patterns of species richness on local
and regional scales. *Species diversity in ecological communities: historical and geographical
perspectives*:66–74.
36. Abrams PA (1988) Resource Productivity-Consumer Species Diversity: Simple Models of
Competition in Spatially Heterogeneous Environments. *Ecology* 69(5):1418–1433.
37. Storch D, Evans KL, Gaston KJ (2005) The species–area–energy relationship. *Ecology Letters*
8(5):487–492.
38. Jetz W, Fine PVA (2012) Global Gradients in Vertebrate Diversity Predicted by Historical Area-
Productivity Dynamics and Contemporary Environment. *PLOS Biology* 10(3):e1001292.
39. Loreau M, Mouquet N, Gonzalez A (2003) Biodiversity as spatial insurance in heterogeneous
landscapes. *PNAS* 100(22):12765–12770.
40. Cardinale BJ, Ives AR, Inchausti P (2004) Effects of species diversity on the primary productivity
of ecosystems: extending our spatial and temporal scales of inference. *Oikos* 104(3):437–450.
41. Srivastava DS, Vellend M (2005) Biodiversity-Ecosystem Function Research: Is It Relevant to
Conservation? *Annual Review of Ecology, Evolution, and Systematics* 36(1):267–294.
42. Thompson PL, Isbell F, Loreau M, O'Connor MI, Gonzalez A (2018) The strength of the
biodiversity–ecosystem function relationship depends on spatial scale. *Proc Biol Sci* 285(1880).
doi:10.1098/rspb.2018.0038.
43. Šímová I, et al. (2011) Global species–energy relationship in forest plots: role of abundance,
temperature and species climatic tolerances. *Global Ecology and Biogeography* 20(6):842–856.
44. Lavers C, Field R (2006) A resource-based conceptual model of plant diversity that reassesses
causality in the productivity–diversity relationship. *Global Ecology and Biogeography* 15(3):213–
224.
45. Shipley B (2016) *Cause and Correlation in Biology: A User's Guide to Path Analysis, Structural*

Equations and Causal Inference with R (Cambridge University Press). 2nd Ed.

46. Grace JB, Anderson TM, Olff H, Scheiner SM (2010) On the specification of structural equation
models for ecological systems. *Ecological Monographs* 80(1):67–87.

47. Petersen ML, van der Laan MJ (2014) Causal models and learning from data: integrating causal
modeling and statistical estimation. *Epidemiology* 25(3):418–426.

48. Wardle DA (2016) Do experiments exploring plant diversity–ecosystem functioning relationships
inform how biodiversity loss impacts natural ecosystems? *Journal of Vegetation Science*:n/a–n/a.

49. Hillebrand H, et al. (2018) Biodiversity change is uncoupled from species richness trends:
Consequences for conservation and monitoring. *J Appl Ecol* 55(1):169–184.

50. Manning P, et al. (2019) Transferring biodiversity-ecosystem function research to the management
of 'real-world' ecosystems. *Advances in Ecological Research* (Academic Press).
doi:10.1016/bs.aecr.2019.06.009.

51. Isbell F, et al. (2018) Quantifying effects of biodiversity on ecosystem functioning across times and
places. *Ecology Letters* 21(6):763–778.

52. Dornelas M, et al. (2014) Assemblage Time Series Reveal Biodiversity Change but Not Systematic
Loss. *Science* 344(6181):296–299.

53. Magurran AE, et al. (2018) Divergent biodiversity change within ecosystems. *PNAS*:201712594.

54. Bannar☐Martin KH, et al. (2018) Integrating community assembly and biodiversity to better
understand ecosystem function: the Community Assembly and the Functioning of Ecosystems
(CAFE) approach. *Ecology Letters* 21(2):167–180.

55. Rohde K (1992) Latitudinal Gradients in Species Diversity: The Search for the Primary Cause.
Oikos 65(3):514–527.

56. Allen AP, Brown JH, Gillooly JF (2002) Global Biodiversity, Biochemical Kinetics, and the
Energetic-Equivalence Rule. *Science* 297(5586):1545–1548.

57. Whittaker R (1975) *Communities and Ecosystems* (MacMillan, New York, USA). 2nd Ed.

58. Evans KL, Greenwood JJD, Gaston KJ (2005) Dissecting the species–energy relationship.
Proceedings of the Royal Society B: Biological Sciences 272(1577):2155–2163.

59. Ratcliffe S, et al. (2017) Biodiversity and ecosystem functioning relations in European forests
depend on environmental context. *Ecol Lett* 20(11):1414–1426.

60. Schluter D, Pennell MW (2017) Speciation gradients and the distribution of biodiversity. *Nature*
546(7656):48–55.

61. Zuppinger-Dingley D, et al. (2014) Selection for niche differentiation in plant communities
increases biodiversity effects. *Nature* 515(7525):108–111.

62. Wagg C, et al. (2017) Plant diversity maintains long-term ecosystem productivity under frequent
drought by increasing short-term variation. *Ecology* 98(11):2952–2961.

63. Loreau M, Hector A (2001) Partitioning selection and complementarity in biodiversity experiments.
Nature 412(6842):72–76.

64. Turnbull LA, Isbell F, Purves DW, Loreau M, Hector A (2016) Understanding the value of plant
diversity for ecosystem functioning through niche theory. *Proceedings of the Royal Society of
London B: Biological Sciences* 283(1844). doi:10.1098/rspb.2016.0536.

65. Craven D, et al. (2019) Evolution of interdisciplinarity in biodiversity science. *Ecology and
Evolution* 9(12):6744–6755.

66. Blackard JA, et al. (2008) Mapping U.S. forest biomass using nationwide forest inventory data and
moderate resolution information. *Remote Sensing of Environment* 112(4):1658–1677.

67. Paquette A, Messier C (2011) The effect of biodiversity on tree productivity: from temperate to
boreal forests. *Global Ecology and Biogeography* 20(1):170–180.

68. Venail P, et al. (2015) Species richness, but not phylogenetic diversity, influences community
biomass production and temporal stability in a re-examination of 16 grassland biodiversity studies.
Functional Ecology 29(5):615–626.

69. Cadotte MW (2015) Phylogenetic diversity and productivity: gauging interpretations from
experiments that do not manipulate phylogenetic diversity. *Functional Ecology* 29(12):1603–1606.

70. USDA Forest Service (2017) *Forest Inventory and Analysis National Program (FIA) database v*
170. Available at: http://fia.fs.fed.us/tools-data/default.asp.
71. Burrill EA, et al. (2018) *The Forest Inventory and Analysis Database: database description and*
user guide version 8.0 for Phase 2 (U.S. Department of Agriculture, Forest Service) Available at:
http://www.fia.fs.fed.us/library/database-documentation.
72. Kartesz JT (2015) The Biota of North America Program (BONAP). *Taxonomic Data Center*.
Available at: (http://www.bonap.net/tdc.
73. The Plant List (2013) Version 1.1 URL: http://www.theplantlist.org/ [accessed 1 December 2017].
Available at: http://www.theplantlist.org/ [Accessed January 1, 2017].
74. The Taxonomic Name Resolution Service iPlant Collaborative. Version 4.0. Available at:
http://tnrs.iplantcollaborative.org.
75. Meyer C, Weigelt P, Kreft H (2016) Multidimensional biases, gaps and uncertainties in global plant
occurrence information. Ecology Letters 19(8):992–1006
76. Zanne A, et al. (2013) *Data from: Three keys to the radiation of angiosperms into freezing*
environments (Dryad Digital Repository) doi:10.5061/dryad.63q27.2.
77. USDA NRCS (2018) The PLANTS Database (http://plants.usda.gov). Available at:
http://plants.usda.gov.
78. Thomas SC, Martin AR (2012) Carbon Content of Tree Tissues: A Synthesis. *Forests* 3(2):332–
352.
79. Zhao M, Heinsch FA, Nemani RR, Running SW (2005) Improvements of the MODIS terrestrial
gross and net primary production global data set. *Remote Sensing of Environment* 95(2):164–176.
80. Zhao M, Running SW (2010) Drought-induced reduction in global terrestrial net primary production
from 2000 through 2009. *Science* 329(5994):940–943.
81. Chojnacky DC, Heath LS, Jenkins JC (2013) Updated generalized biomass equations for North
American tree species. *Forestry* 87(1):129–151.
82. Pan Y, et al. (2012) *NACP Forest Age Maps at 1-km Resolution for Canada (2004) and the U.S.A.*
(2006). Data set. (ORNL DAAC, Oak Ridge, Tennessee, USA).
83. Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated
climate surfaces for global land areas. *International Journal of Climatology* 25(15):1965–1978.
84. Stein A, Gerstner K, Kreft H (2014) Environmental heterogeneity as a universal driver of species
richness across taxa, biomes and spatial scales. *Ecol Lett* 17(7):866–880.
85. USGS (2009) SRTM v2.1 Topography. Available at:
http://dds.cr.usgs.gov/srtm/version21/Documentation/SRTMTopo.pdf.
86. Karger DN, et al. (2016) Delineating probabilistic species pools in ecology and biogeography.
Global Ecology and Biogeography 25(4):489–501.
87. Cochran WG (1977) *Sampling techniques* (John Wiley and Sons, New York, USA). 3rd Ed.
88. Rosseel Y (2012) lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical*
Software 48(2):1–36.
89. Breiman L (2001) Random forests. *Machine learning* 45(1):5–32.
90. Liaw A, Wiener M (2002) Classification and regression by randomForest. *R news* 2(3):18–22.