The conserved regulatory basis of mRNA contributions to the early *Drosophila* embryo differs between the maternal and zygotic genomes

Charles S. Omura* and Susan E. Lott*

Department of Evolution and Ecology, University of California, Davis CA 95616

*Corresponding Authors, e-mail: csomura@ucdavis.edu (CSO), selott@ucdavis.edu (SEL)

Abstract

The gene products that drive early development are critical for setting up developmental trajectories in all animals. The earliest stages of development are fueled by maternally provided mRNAs until the zygote can take over transcription of its own genome. In early development, both maternally deposited and zygotically transcribed gene products have been well characterized in model systems. Previously, we demonstrated that across the genus *Drosophila*, maternal and zygotic mRNAs are largely conserved but also showed a surprising amount of change across species, with more differences evolving at the zygotic stage than the maternal stage. In this study, we use comparative methods to elucidate the regulatory mechanisms underlying maternal deposition and zygotic transcription across species. Through motif analysis, we discovered considerable conservation of regulatory mechanisms associated with maternal transcription, as compared to zygotic transcription. We also found that the regulatory mechanisms active in the two genomes, maternal versus zygotic, are quite different. For maternally deposited genes, we uncovered many signals that are consistent with transcriptional regulation through control at the level of chromatin through factors enriched in the ovary, rather than precisely controlled gene-specific factors. For genes expressed only by the zygotic genome, we found evidence for previously identified regulators such as Zelda and GAGA-factor, with multiple analyses pointing toward gene-specific regulation. The observed mechanisms of regulation are consistent with what is known about regulation in these two genomes: during oogenesis, the maternal genome is optimized to quickly produce a large volume of transcripts to provide to the oocyte; after zygotic genome activation, mechanisms are employed to activate transcription of specific genes in a spatiotemporally precise manner. Thus the genetic architecture of the maternal and zygotic genomes and the specific requirements for the transcripts present at each stage of embryogenesis determine the regulatory mechanisms responsible for transcripts present at these stages.

Introduction

Development is a sequential process, where each step builds on the one before it. The earliest stages of embryonic development are therefore critical, as processes such as cleavage cycles and the beginnings of axial patterning become the basis for all subsequent developmental processes. Regulation of these important tasks is controlled by mRNAs and proteins, and perhaps unsurprisingly then, mRNA levels in *Drosophila* are found to be precisely controlled during early embryogenesis [1,2]. This precise control of transcript levels is especially remarkable, given that the transcripts at early stages of development come from two different genomes. The first set of transcripts are those deposited into the egg by the mother, while the second set are transcribed from the zygotic genome [3–5]. However, the regulatory mechanisms responsible for this precise control are not yet fully understood.

During oogenesis, the oocyte itself is mostly transcriptionally silent [6]. Instead, support cells called nurse cells synthesize RNA, proteins, and organelles which are transported into the oocyte [7]. These maternally produced mRNAs are responsible for many of the critical events of early embryogenesis, such as the rapid cleavage cycles, the establishment of body axis, and the coordination of the handoff of control to the zygotic genome. This handoff of developmental control from mother to zygote, known as the maternal to zygotic transition (MZT), is complex from a regulatory standpoint. Critical housekeeping genes retain a steady transcript level, despite changing the genome of origin. New transcripts must be synthesized from the newly activated zygotic genome, and maternal transcripts must be degraded, in a highly regulated and time-specific manner [8]. This transition is well studied in model systems such as *Drosophila melanogaster*, where maternal mRNA degradation regulators such as *smaug* (*smg*) [8] and regulators critical to the activation of the zygotic genome such as *zelda* (*zld*) [9,10] have been identified. When the transition of developmental control between the two genomes is complete, the zygotic genome must be poised to carry out the rest of development in a precise manner. One process that

exemplifies the precision required at the handoff to the zygotic genome is segmentation in *Drosophila*.

This process begins with broad maternal gradients which control transcription of early zygotic gap

genes, and later pair-rule genes, at precise locations within the embryo at specific developmental times

[11,12].

Regulation of transcripts in development has been the subject of considerable study in *D.melanogaster*.

Much of this study has been focused around the process of the MZT or other important events in early

development, such as patterning along the anterior-posterior or dorsal-ventral axes. For example, a

number of regulators of maternal transcript degradation at or prior to the MZT have been identified

[13–16]. Zygotic transcription activation has also been the subject of considerable study, and has

implicated critical transcription factors such as *zelda* and *grainy head* [4]. How transcripts are

transported into eggs has been the subject of some study [7,17,18], as has how those maternal

transcripts are regulated post-transcriptionally [3,19–23]. Post-transcriptional mRNA regulation is

especially crucial at the maternal stage as new transcripts cannot be produced after the completion of

oogenesis. However, how transcript production is regulated in the nurse cells is largely unknown. As

transcript pools at both the maternal and zygotic stages are highly conserved over evolutionary time

[24], we employed a comparative approach to investigate gene regulation at these stages.

In this study, we uncover regulatory elements that are associated with transcription in the early

Drosophila embryo, from both maternally deposited and zygotically transcribed genes. We use motif

analysis to compare regulation of maternal versus zygotic transcription, and also investigate how

regulation at these two stages is different across *Drosophila* species. To this end, we used a previously

generated RNAseq dataset from Atallah and Lott, 2018, which sampled embryos from a developmental

stage where all transcripts are maternal (stage 2[25,26]) and a stage after zygotic genome activation

(end of stage 5 [23,24], across 14 species, representing ~50 million years of divergence time. Here, we

used the transcript abundance data from 11 of these species (due to limitations in genome annotation

quality, see Methods), representing the same span in divergence time, to examine putative regulatory

regions of maternally deposited or zygotically transcribed genes. Through comparisons of these

sequences and associated gene transcription levels, we identified a number of sequence motifs as being

enriched in either maternally deposited or early zygotically expressed genes. We found a high similarity

between motifs across all species, suggesting a high level of conservation for regulation of transcription

within each genome (maternal and zygotic). At the stage controlled by maternal transcripts, we found a

high number of motifs that bind to proteins annotated with insulator function or that have previously

been associated with boundaries between topologically associating domains (TADs). Our findings

suggest that maternal transcription is largely controlled through regulation of chromatin state, and not

through gene-specific mechanisms. Many transcription factors predicted to bind the identified motifs

were found to be enriched in ovaries [27]. After zygotic genome activation (stage 5), we find many of

the motifs known to be associated with early zygotic transcription, such as the binding site for the

pioneer transcription factor, Zelda, reinforcing many previously identified aspects of transcriptional

regulation at this stage. We also find a larger number of motifs with less significant enrichment at this

stage, with evidence that points to these motifs regulating a smaller subset of genes. This study provides

evidence for global control of maternal transcription at the level of chromatin, while zygotic

transcription is regulated in a more gene-specific manner. This is especially striking considering that the

maternal transcript pool is more highly conserved than that of the early zygote [24].

Results

Discovered maternal-associated motifs are bind architectural proteins; discovered

zygotic-associated motifs bind to known zygotic regulators

To examine the regulatory basis of maternal and zygotic transcription, we surveyed the genomes of 11

Drosophila species for regulatory elements. These species represent the evolutionary divergence of the

Drosophila genus, encompassing divergence times from 250,000 to 50 million years [28]. The RNAseq

datasets produced from Atallah and Lott (2018)[24] were used. These data sampled two developmental

stages, one where all transcripts present are maternally derived (stage 2, Bownes' stages [25,26]) and

the other after zygotic genome activation (the end of stage 5, or the end of blastoderm stage). The

transcript abundance data was used to classify each gene as being on or off at both stage 2 and stage 5

for each species (see Methods). For each gene, we extracted sequences at likely locations for proximal

regulatory elements (see Methods). To accommodate the varying annotation quality of the various

species, this search encompassed introns, exons, and a 2kb region upstream of the gene.

To identify motifs associated with maternally deposited genes, we employed HOMER [29]. For most

species, a characteristic pattern emerged where the most enriched motifs were present in the upstream

region of the maternally deposited genes, with less enriched motifs appearing in exons (Fig S1). Some

motifs, possibly representing repressor binding sites, were enriched in the upstream and intron region

of genes that were not maternally deposited as compared to the genes that were maternally deposited

(Fig S1).

Analyzing regulatory elements at the post-zygotic genome activation stage (stage 5) presents a

challenge, as it is difficult to distinguish newly transcribed zygotic mRNAs from residual maternally

deposited mRNAs. At this stage, roughly half of the transcripts present are maternal transcripts that

have not yet been degraded [8,30–32]. Therefore, to interrogate regulatory elements associated with

zygotic transcription, we restricted our search to genes that do not have transcripts present at stage 2

but do have transcripts present by stage 5. Because of these stricter requirements for zygotically

transcribed genes, there were far fewer genes in the dataset, resulting in less statistical power.

However, without these assumptions, we risk failing to identify signals associated specifically with

zygotic transcription amongst the signal of maternal transcription.

To determine which proteins are likely to bind to maternal or zygotic motifs, we used Tomtom [33] to

evaluate the similarity of the discovered motifs to several motif databases (Table 1) for *D. melanogaster*.

For motifs associated with maternally deposited transcripts, we found a striking similarity between the

motifs generated by Ramírez et al. in 2018[34], who discovered a number of motifs associated with

topologically associated domain (TAD) boundaries in *Drosophila* S2 cells. TAD boundaries are genomic

regions where the chromatin on one side of the boundary interacts substantially less than expected with

the chromatin on the other side. We hypothesize that the regulatory mechanisms responsible for

generating TAD boundaries in the Ramírez et al. (2018) paper are also responsible for maternal

deposition, and that maternal deposition is regulated by the establishment of TAD boundaries. This set

of motifs that are found to be associated with TAD boundaries in our study are also similar to the set

that were found to be associated with housekeeping core promoters by Zabidi et al. in 2015[35]. This is

consistent with existing data showing that functions of maternally deposited genes are enriched for

genes with housekeeping activities [36].

The motifs associated with maternally deposited genes are predicted to bind several insulators or

architectural proteins. An insulator is a regulatory element that suppresses the interactions of other

regulatory elements with genes, or prevents the spread of chromatin state. An architectural protein is a

protein that organizes and regulates chromatin structure. The most prominent motif by q-value binds to

DNA replication-related element factor (DREF), a known architectural protein and the "master key-like

factor for cell proliferation" [37]. It is required for normal progression through the cell cycle. It is known

to occur in the promoters of many cell proliferation genes and to interact with chromatin remodeling

proteins. Interestingly, DREF binding site overlaps with the binding site for BEAF-32, another well-

researched protein that acts as an insulator [38,39] that often appears between head-to-head genes

(genes with adjacent promoters that get transcribed in opposite directions). Another identified motif is

predicted to bind ZIPIC, which is known to bind and recruit CP190, an insulator. A previous study

provides evidence for the co-localization of ZIPC and BEAF-32[40], which likely work together with

CP190 to perform insulator functions. Thus of the most enriched motifs in maternal genes (DREF, BEAF-

32, ZIPC), many have previously identified roles as insulators or in other ways regulating chromatin

state.

Another maternal motif identified is predicted to bind M1BP (motif-1 binding protein), which causes

RNA polymerase II (Pol II) to pause on the gene [41]. Pol II pausing is critical to early zygotic

expression[42,43] but its function in producing the maternal transcriptome is unknown. Several

functions have been suggested for this Pol II pausing behavior, including maximizing transcription speed

once certain conditions are met, synchronizing with RNA processing machinery, reacting to other

developmental or environmental signals, keeping chromatin accessible, and acting as an insulator. Given

that M1BP is both maternally deposited at high levels and has increased expression in the early embryo,

it is possible that M1BP has multiple functions at different time points. During oogenesis, pausing to

wait for external signals or RNA processing machinery seems counterproductive to maximizing

transcription in the ovary, but the other function of maintaining a state of open chromatin and

solidifying TAD boundaries may be very important. In contrast, at stage 5 it may be much more

important to maximize expression in response to certain signals.

In searching for motifs associated with zygotic expression, we recovered motifs for well-known

regulators of the zygotic genome (Table 1). We only identified a small number of highly enriched motifs

at this stage, and thus were able to predict a much smaller number of predicted factors binding to these

motifs, including Trl (or GAGA factor) and Zelda. Trl is a known early zygotic activator and chromatin

remodeler [44–46] and Zelda is known as a "master key regulator" to early developmental genes [9,47]

and appears to be a pioneer transcription factor that establishes the initial chromatin landscape of the

zygotic genome [5] . In addition to these high-quality motifs, we found a large number of motifs with

lower quality scores (Table S1). These motifs may regulate spatio-temporal specific genes that we

observe in the early embryo, and thus have a lower enrichment score due to our whole-embryo

approach being ill-equipped to finding such specific patterns.

167   Similar motifs appear in different species

To quantify the conservation of the discovered motifs across the 11 species in our study, we used

Tomtom [33] to measure the similarity between the sets of motifs discovered in different species. For a

motif to be considered conserved between two species, we required that it be discovered by HOMER in

both species and for Tomtom to report a statistically significant alignment score (see Methods). At the

maternal stage, we found that high quality (q-value < 1e-100 by HOMER, see Methods) motifs tended to

be well-conserved (Fig 1A) with a large percentage of the total discovered motif content shared across

species. We observed that sister species *D. pseudoobscura* and *D. persimilis* are unique in that they have

the highest number of motifs that are either species-specific or are only shared with each other, and

have the fewest number of motifs shared with the rest of the species. This is especially noteworthy

considering that this lineage is roughly in the middle of the distribution of divergence times from most

of the other species, and thus many more distantly related species comparisons have a higher degree of

motif conservation than do any comparisons with these two species. This is consistent with previous

results [24] that this lineage has a disproportionately high number of changes in transcript abundance

for its phylogenetic position, and suggests that these large number of changes in transcript abundance

may be due to the large scale changes in regulation in these species observed here . When comparing

the rest of the species, we found a relatively higher number of conserved motifs shared between pairs

of species within the *Drosophila melanogaster* species group (*D. melanogaster*, *D. simulans*, *D. sechellia*,

D. yakuba, *D. erecta*, *D. ananassae*), and a slightly reduced number of conserved motifs between the *D.*

melanogaster group species and the more distantly related species (*D. willistoni*, *D. mojavensis*, *D. virilis*)

(Fig 1B). At stage 5, we do not observe a high percentage of conserved motifs between species, rather

we observe many motifs that are significantly enriched in just one or two species. We also observe little

phylogenetic signal in the data, with the only detectable pattern being that the species with the longest

divergence time from the rest of the species, *D. virilis* and *D.mojavensis,* have slightly fewer shared

motifs (Fig 1 C,D). If the unique motifs at either stage indeed represent newly evolved regulatory

mechanisms, we expect that these motifs to be rare or to have a smaller frequency difference between

transcribed and non-transcribed genes. Either of these effects would raise the false discovery rate as

reported by HOMER, which makes the number of species-specific zygotic motifs identified all the more

remarkable.

Motif conservation by gene

While these results show that some motifs are important to regulation in the genomes of multiple

species, do not speak to whether orthologous genes in different species tend to contain similar motifs.

To investigate whether regulation was conserved at the level of individual genes, we compared the

motif content of each *D. melanogaster* gene (see Methods) to the motif content of each of its orthologs

from other species. We counted motifs as conserved between two species if the motif appeared in both

orthologs. For both stage 2 and stage 5, we categorized motifs based on the percent of orthologs for

which the motif was conserved (Fig 1E). Motifs have different levels of gene-specific conservation

between stages, with maternal stage motifs appearing to have lower conservation across orthologues

than zygotic stage motifs, where a larger proportion of orthologues possess the same motif. It is possible

that the genes that are missing motifs compared to their orthologues are regulated by different motifs,

or that the same motifs that are in radically different positions in different species. As many different

maternal motifs appear to be regulating transcription at the level of chromatin state, these motifs may

be able to function interchangeably, thus leading to more motif turnover at this stage but with higher

conservation of transcription overall [24], as compared to the zygotic stage.

Motif position

While similar binding motifs identified in multiple species implies that regulatory proteins with similar binding domains are acting in these species, we can also verify the similarity in the regulatory machinery by the relative positions of the binding sites relative to the genes they are regulating. To investigate whether the discovered motifs had the same positional relationship with the transcription start site (TSS) across all species, we generated position frequency data for each motif. For each gene, we examined each position starting from 2kb upstream of the TSS to the 3' end of the gene body, and whether there was a motif at that position. Many of the most prominent motifs shared a similar distribution pattern, characterized by a strong peak at -100bp, and sometimes a secondary peak at -340bp (Fig S2). To quantify this similarity, we performed an Anderson-Darling test on each motif for each pair of species, which indicated that 65% (stage 2) and 91% (stage 5) of motif distributions are identical between species (percent of motifs for which $p < .05$). This suggests conservation of the relationship between binding to these motifs and initiation of transcription. The higher conservation of motif position in stage 5, which has fewer conserved motifs between species than stage 2, may be consistent with this stage having more gene-specific regulation, as discussed further below.

Motif Strandedness

While some studies focus on finding motifs with a particular orientation relative to their proximal genes [48], there is some evidence that motifs do not behave in a strand-specific manner[49]. To evaluate the importance of the strandness of the discovered motifs, we generated a regression to predict expression level that differentiated between forward and reverse versions of each motif (see Methods). This regression indicated a significant difference between the forward and reverse versions of many motifs. For example, we found the E-box motif affects the log-odds of maternal deposition by .192 in the forward orientation but only .115 in the reverse orientation (t-test, $p < .001$). Almost all motifs had the

same sign coefficients, but with different magnitudes, indicating that while motifs had the same effect

regardless of orientation, their efficiency could be increased if the orientation was optimal.

While the strandedness of motifs may play a small role in their overall effect, we want to know if

strandedness makes a qualitative difference to our motifs effects on transcript level, and if we can use

motif strand to improve our model. To determine this, we ran HOMER exclusively on the same strand

that the gene appeared on, rather than the default mode of scanning both strands. This resulted in the

same set of motifs being discovered. This is consistent with the regression results that show that the

two strands of the motifs generally have the same sign, indicating an effect in the same direction on

transcript level. To evaluate the overall effectiveness of this strategy, we constructed another regression

using only the data from the same-strand motifs. This regression performed less well than the

regression using motifs from both strands (AIC = 7915.8 for the unstranded regression, AIC = 8612.5 for

the stranded regression for a representative species *D. ananassae*). Overall, this suggests that motif

binding elements need not bind in a strand specific manner to induce their effects, though the optimal

orientation provides measurable increase in their effect on transcription. This result is the same at both

stage 2 and stage 5.

GO analysis

While we have identified a set of motifs that together seem to be responsible for early embryonic RNA

content, we next asked if these motifs are likely to be regulating genes with specific types of functions.

To this end, we performed gene ontology (GO) analysis on groups of genes, based on their motif

content. To simplify this analysis, we chose to focus on the top 8 motifs as reported by HOMER, and for

each of the 8 motifs, we compared their enrichment relative the enrichGO function in the clusterProfiler

R package [50,51]. 8 motifs were chosen to include representatives from both known and unknown

motifs. When comparing stages, we observe no overlap between GO terms. (Fig 3A), which is consistent

with our expectations that the genes that are activated in the zygote have different functionality to

those transcripts that are maternally deposited. Many of the stage 2 motifs show a similar pattern in the

GO categories they are associated with, with the strongest associations belonging to the DREF motif,

which is strongly associated with most identified categories (Fig 3B). This could be an indication that

there is a high degree of homogeneity in terms of the types of genes these motifs may regulate. In

contrast, the stage 5 motifs show more variety in the GO terms of genes they are associated with (Fig

3C), which could be indicative of more specific regulation at this stage.

Predicted maternal motif binding proteins are enriched in the ovary

Next, we investigated whether the potential motif binding proteins we identified were plausible

regulators of maternal deposition. It is unclear whether the motifs we identified as enriched in

maternally deposited genes are associated specifically with maternal deposition, given that chromatin

regulators are important at all stages in all tissues. To investigate, we used modENCODE [52] transcript

abundance data to compare the mRNA transcript levels for proteins predicted to bind our discovered

motifs, and found increased expression in ovaries (Fig 4A) as compared to other tissues sampled. This

pattern exists, though to a lesser extent, in the FlyAtlas 2 dataset [53] , which is a tissue-specific

database of transcript levels that utilizes RNA-seq data rather than microarray analysis. The discrepancy

between the two datasets could be due to the differences in gene expression measurement method or

in experimental methods. While it has been demonstrated that mRNA levels do not necessarily mirror

protein levels [54], the enrichment of mRNA in ovaries compared to other tissues is reasonable evidence

that these proteins are important in ovaries.

To investigate whether these proteins are acting to affect transcription in the ovaries specifically, we

examined the expression profiles of RNA in various tissue types (referenced in Fig 3B) from existing RNA

quantification datasets [52,55]. For each instance of a motif of interest, we extracted the transcript level

from within a 20kb window surrounding the motif and measured the normalized relative transcript level

for each position (an example of this is shown in Fig 3B). While the relative normalized transcript level

changes in each of the measured tissues, the effect is strongest in ovaries, indicating that the presence

of one of these binding sites is associated with a higher increase in transcript levels in the ovary

compared to other tissues.

As the motifs associated with maternal transcription also act to some degree in other tissues, we next

wanted to ask whether the motifs were more enriched in maternally deposited genes than in genes

expressed in other tissues. To determine whether regulation in different tissue types were associated

with different motifs, we ran HOMER in the same manner as with the maternal stage data to discover

enriched motifs (see Methods) in transcripts present in other tissues, as identified from ModENCODE

data [56]. We found that most other tissue types were also enriched in the same motifs discovered in

transcripts present in stage 2 embryos. However, examining the frequency of motifs in specific genes

revealed that the majority of those motifs were from genes that were shared between those tissue

types and stage 2 embryos. When we exclude genes that are expressed in stage 2 embryos, HOMER fails

to identify the original set of motifs as enriched in male larval gonads, male reproductive tract, adult

heads and adult midgut. Furthermore, HOMER detects the motifs at a lesser rate in larval ovaries, larval

CNS, and intestinal tract. Despite being identified in fewer tissue types and at a lesser rate in other

tissue types as compared to the stage 2 expression levels, the observation that these motifs may also

have important functions in other tissue types is consistent with the literature. For example, DREF is

known to be important for cell proliferation and chromatin regulation, and is active in many other

tissues [57,58]. These motifs are likely associated with many housekeeping genes that are vital to a

variety of tissue types.

Maternally deposited genes are physically clustered on the genome

In addition to motifs, we observed several other effects that were related to early embryonic RNA content. Given that many of our discovered motifs bind architectural proteins, we hypothesize many effects may be linked to the physical location of genes on the chromosome. We examined the positional distribution of transcribed genes in various tissue types. As previous papers utilizing the Hi-C method have shown correlation with active topologically associated domains (TADs) and gene expression [59,60], we predicted that any tissue type where regulation is dominated by architectural proteins to transcribe a set of genes physically clustered on the chromosome. To compare the physical gene clustering of transcription at the maternal stage with that of other tissue types, we acquired several RNAseq datasets from NCBI/GEO [61] and performed a Wald–Wolfowitz runs test [62] on each tissue of the previously described tissue types. While all tissues examined showed a strong preference for groupings of transcribed genes, ovaries were the most grouped (Fig 4B). This result was robust to changes in the threshold of what is considered to be expressed (see Methods). This pattern of physical co-expressed gene clustering on the chromosome is consistent with our model of regulation via architectural proteins.

While these results speak to the pattern of clustering of expression for maternal genes in terms of adjacent genes being on or off, they do not account for the distance between genes. To answer the question of whether this clustering phenomenon is dependent on distance, we examined the distance to adjacent genes. We observed a trend whereby proximity to an active promoter increases the odds of maternal deposition (Fig 4C). This effect was slightly affected by the strandedness of the two genes whereby genes that have an opposite orientation are more likely to have different expression. This is consistent with observations from previous studies [34] that consecutive genes on the same strand were more likely to show co-expression, while consecutive genes on opposite strands were more likely to have different expression.

Many previous studies have observed that zygotic genes tend to be short in length [24,30,63,64]. In

addition to affecting transcription speed, shorter gene lengths result in a smaller distance between

transcriptional units along the chromosome, especially when considering which strand the gene is on. To

explore gene length in maternal genes and the relationship between gene length and the position on

the chromosome, we measured the maternal deposition rates with respect to gene length. We observed

a trend that in most species, shorter genes are less likely to be maternally deposited. There are

differences in the length of maternal genes across species, and this trend could be partly due to the bias

for more highly annotated genomes to be enriched in shorter genes (Fig 4D). Additionally, chromatin

context seems to heavily influence this effect: when the adjacent genes are off, gene length is much

more important (Fig 4C) and very short genes are very likely to be off. This could be because shorter

genes are more likely to be influenced by the regulatory machinery of a nearby gene. Alternatively,

longer genes might be long enough to physically isolate themselves more effectively and establish their

own unique regulatory environment.

Given that a number of motifs found in this study are bound by proteins annotated as insulators, and

the motifs are similar to those that are associated with TAD boundaries, we asked whether the dataset

shows properties consistent with these motifs acting as insulators. To determine whether these motif

binding sites were associated with proteins acting as insulators, in the sense that they were blocking the

action of a nearby enhancer or the spread of chromatin state, we assessed the transcription of nearby

genes relative to the transcription of a gene with these identified motifs. For each regulatory region, the

gene nearest to that regulatory region was examined, as well as two genes downstream and two

upstream. The frequency of motifs was measured based on the transcript abundance pattern of these

five genes. Many of the top motifs including Dref, M1BP, Zipic, and E-box, occur more frequently in the

center of maternally deposited gene clusters, rather than on the edge of clusters. (t-test p-values 7e-3,

2e-6,3e-10, and 1e-4 respectively). This is consistent with previous results that suggest architectural

proteins are prevalent in the centers of TADs as well as the edges[34].

Stage-specific genes are isolated on the genome

Given that maternally deposited genes are physically clustered together in the genome, we wanted to

examine if this pattern held with the set of genes that were stage-specific. To determine if consecutively

expressed cluster size is related to stage-specificity of transcript representation, we examined maternal-

only (transcripts present at stage 2 and entirely degraded by stage 5) and zygotic-only genes (transcripts

present at stage 5, not present at stage 2; for both stage-specific categories, see Methods for further

definitions) and their frequencies in clusters of different sizes. We determined that for most species,

both maternal-only and zygotic-only genes are more likely to be in smaller (1-3 consecutive maternally

deposited) groups than in larger groups (more than 3 consecutive genes) (Fig 5, A and B). For zygotic-

only genes, this could be an indication that control at the zygotic level is more specific, affecting single

genes rather than larger clusters. Meanwhile, maternal-only genes being in small groups could more

likely to have more specific regulation than that of all maternal genes as a whole

GC-content of upstream regions

In *Drosophila*, transcription start sites are frequently associated with a spike in GC content [65]. To

explore this phenomenon with respect to the two developmental stages we examined, we evaluated the

average GC content of upstream regions for genes in stage 2 and stage 5. When comparing the GC-

content of putative cis-regulatory sequences in maternally versus non-maternally deposited genes, we

observed an increase in GC-content upstream of the TSS (Fig S3), as well as a dip in GC content ~200bp

upstream of these genes. This modulation in GC-content has been previously observed in *Drosophila* as

well as humans [65]. To determine whether this modulation of GC-content was predictive of maternal

deposition, we constructed four generalized linear models using the GC-content, the motif data, and

both the motif data and GC-content as data sources (see Methods). Adding the GC-content to the model

that already included motif data improved the model (AIC: 185589 without GC content AIC:183079 with

GC content), hence increased GC content upstream of TSS is somewhat predictive of maternal

deposition, even when accounting for motif presence in this region.

Discussion

Maternally deposited gene products are responsible for the first stages of embryonic development in all

animals [66]. It is therefore critical that the required kind and amount of mRNAs and proteins are

deposited into the unfertilized egg. Later in development, the zygotic genome becomes transcriptionally

active and takes over control of development from maternal mRNAs. Failure in maternal mRNA

deposition, zygotic genome activation, or the transfer of developmental control between the two

genomes can lead to lethality [1,9,13], thus the gene products regulating early development are critical

to organismal survival.

Previous research has shown that the maternal and zygotic mRNA expression profiles of different

species of *Drosophila* are generally conserved, but with some noticeable differences [24]. To identify the

regulatory basis of trascription at these stages, we leveraged a large comparative dataset to identify the

transcription factor binding motifs found in the *cis*-regulatory sequences of these genes. We identified

transcription factor binding motifs that are associated with mRNA expression across species for

maternally deposited transcripts and zygotically expressed transcripts. We also investigated the effects

of other regulatory mechanisms such as chromatin state on maternal and zygotic expression of mRNAs,

as well as the association of transcript levels at these two stages of embryogenesis with gene length,

strandedness, and GC content.

Maternal Regulation

We found that motifs associated with putative *cis*-regulatory regions of maternally deposited genes

are predominantly annotated as insulator binding sites. An insulator is a type of regulatory element

that can block the interactions of *cis*-regulatory elements with promoters or prevent the spread of

chromatin state. Insulators are known to be important in creating and maintaining the gene

expression patterns, ubiquitous in *Drosophila*, and potentially a key factor for *Drosophila* to maintain

such a high gene density [40]. Here, we find that the process of maternal deposition may rely heavily

on insulators to express a large percentage of the genome. Because the roles and mechanisms of

factors annotated as insulators are not well understood, using the term "Architectural Protein"

instead of insulator binding protein may be more appropriate [67]. Recently, these proteins have

been studied using genome-wide chromatin organization methods, such as Hi-C, which detects

regions of interacting chromatin known as Topologically Associated Domains (TADs) and identifies

boundaries between them. Histone marks appear to be enriched in certain TADs but stop abruptly at

TAD boundaries, supporting the idea that certain TADs are entirely transcriptionally silenced while

others are expressed [34]. Furthermore, ChIP-seq has demonstrated that TAD boundaries in other

tissues are enriched in architectural protein binding sites [34] , including several those that we

identified in this study.

To understand the connection between architectural proteins and maternal deposition, we need to

examine where these transcripts are produced to understand the cellular context. In the ovary, nurse

cells are responsible for the transcription of maternally deposited genes, and there is a considerable

body of literature devoted to nurse cell biology. Much study has been directed towards elucidating

how nurse cells transport their products into the oocyte and how post translational control

mechanisms fine-tune protein levels of maternal transcripts [3,7,18–23,68]. However, despite this

wealth of knowledge, the regulatory mechanisms by which the nurse cells specify which genes to

transcribe are largely unknown. One unusual feature of nurse cells is that they are highly polyploid

[69,70]. One of the major benefits of this could be an across-the-board increase in transcription rates

necessary to provision the embryo with all necessary transcripts. These transcripts represent a large

proportion of the genome, with estimates ranging from 50-75%, depending on experimental

conditions [3], and necessitate large amount of transcription overall in a short period of time. We

extract >100ng total RNA from an embryo; this is an astonishingly large amount of RNA to be present

in what is essentially at the time of fertilization a single, albeit a highly specialized, cell. One point of

comparison is Abruzzi et al. 2015, [71] who extracted 2-5pg RNA per *Drosophila* neuron. A

transcriptional environment that is optimized to quickly transcribe huge numbers of genes might be

more amenable to control via chromatin state.

Given the amount of overlap between the motifs enriched in the *cis*-regulatory regions of maternally

deposited genes and the motifs associated with TAD boundaries, it is possible that these same

architectural proteins are functioning to define which genes are maternally transcribed and then

deposited into the embryo. We found that the maternally deposited genes are highly clustered on

the genome, which is indicative of control via architectural proteins. Additionally, we uncovered that

proximity to nearby expressed genes is highly correlated with expression. We also identified a

pattern whereby the relative strandedness of adjacent genes is indicative of whether they will be

maternally deposited, which is a pattern that has been previously observed with insulators [34]. Each

of these results is consistent with known behavior of architectural proteins, suggesting that

expression at stage 2 is controlled locally on the chromosome by activating TADs rather than specific

genes.

As architectural proteins are important in determining genome organization and regulating transcription

to some degree in all tissues and stages, we investigated whether the regulatory patterns we observed

for maternal genes were ovary-specific or shared across all stages and tissues. Many of the motif binding

elements discovered in this analysis appear to be enriched in ovaries, although these proteins have

important functions in other tissues as well. Some of the proteins predicted to bind our motifs have

been noted for being enriched in the regulation of housekeeping genes, and as maternally deposited

genes themselves are enriched in housekeeping genes, this result is perhaps unsurprising. A number of

studies have suggested that in addition to the common architectural proteins shared across conditions

and developmental stages, there may exist tissue-specific architectural proteins that integrate into the

canonical protein complex to produce tissue-specific TAD patterns [72–74]. Perhaps this is the case with

the ovary, and further study will reveal whether there are ovary-specific factors that may interact with

the common architectural proteins whose binding sites we find enriched here. For example, the authors

of Mataz et al. 2012 [75] suggest that Shep may be a tissue-specific factor interacting with architectural

proteins in the central nervous system. The enrichment Shep in the central nervous system is even less

extreme than the enrichment we observe of CP190 (a known interaction partner of ZIPC, one of our

maternal expression associated motifs) in ovaries, suggesting that CP190 could also qualify as tissue-

specific. Alternatively, the polyploid nature of nurse cells and the extensive and rapid transcription that

occurs in these cells may instead provide an extreme enrichment of the common architectural proteins,

without the need for stage or tissue specific architectural proteins.

Our results show that regulation in ovaries is accomplished primarily through architectural proteins that

establish general regions of open chromatin. This process can turn on a large percentage of the genome,

without the need to maintain specific motifs within specific genes. However, this leaves us with the

question of how the stage 2 mRNA content is so highly conserved across species overall [24], as

regulation at the chromatin level would appear less precise than gene-specific regulation. Perhaps

regulatory control primarily at the level of chromatin provides redundancy to maintain transcription

despite the gain or loss of individual binding sites. Alternatively, there could be other levels of regulatory

control that we are unable to detect, with the signal from chromatin-level control being so strong during

this time. The high level of conservation of maternal transcripts is also remarkable given the importance

of post-transcriptional regulators at this stage [3,19,23,76], as it is not clear if conservation at the

transcript level is necessary for conservation at the protein level.

Zygotic Regulation

Our examination of motifs that are associated with zygotic mRNA expression revealed several previously

discovered motifs, including those that bind Zelda and GAGA factor (Trl). Additionally, several motifs are

likely binding sites for other well-characterized developmental proteins (Table S1) which are sometimes

highly localized in the embryo. If transcripts are produced in a spatially localized manner, they are

necessarily not expressed in the entire embryo, and thus their signal may be more difficult to detect in

our data from whole embryos. Overall, we observe few motifs at stage 5 that are conserved across

species, in comparison to motifs for maternally deposited genes. However, the motifs that we do find at

stage 5 tend to higher conservation within specific genes than the motifs we discover at stage 2. This

highlights that it may be more important for specific genes to have precise signals after ZGA.

Additionally, in our zygotic analysis, we focused only transcripts that are present at stage 5 and do not

have a maternal component, as many maternally deposited transcripts are still present at stage 5

(roughly half of maternal transcripts are still present at this stage [8,30–32]). Because many maternal

transcripts are still present, analysis of the total stage 5 transcriptome would largely recapitulate the

stage 2 results, especially as stage 5 transcripts are much more likely to be expressed in specific spatio-

temporal patterns, which to our whole-embryo analysis would appear as low or noisy signal. Our

decision to remove transcripts with maternal deposition highlights the signals that are unique to stage 5,

but comes at the cost of an overall reduction in the number of genes available for analysis, resulting in

higher false discovery rates for all motifs.

Conclusions

In this study, we examined regulatory elements associated with maternal transcripts present at stage 2 of embryogenesis and zygotic transcripts present at stage 5 across species of *Drosophila*. At both stages, we found regulatory motifs that are conserved throughout the ~50 million years of divergence represented by these species*,* which speaks to a conservation of regulatory mechanisms across the genus. In general, the high degree of conservation in regulatory elements at the maternal stage and the zygotic stage, while different from one another, speaks to the critical nature of the complement of transcripts present to direct early embryogenesis. The differing patterns observed in the *obscura* group species (*D. pseudoobscura* and *D. persimilis*), and the regulatory basis of changes in transcript representation between species are the subject of ongoing study. At the maternal stage, we found many regulators that appear to be defining general regions of the genome to be transcribed via chromatin regulation through architectural proteins and likely at the level of TADs. Given the exceptionally high level of conservation of maternal transcript deposition, the relatively non-specific mechanism of maternal gene regulation appears contradictory. In contrast, we found zygotic regulatory elements to be considerably more gene-specific. The different patterns of regulation for transcripts present at these two stages of embryogenesis is consistent with the specific transcriptional contexts of these two genomes, with the non-specific mechanism active in highly transcriptionally active polyploid nurse cells in oogenesis in the mother, and the gene-specific mechanism acting in the zygote where transcription is often localized in time and space.

Methods

Data Acquisition

RNA-seq data utilized for this study was generated previously [24], and is available at NCBI/GEO at accession number GSE112858. This dataset contains RNA-Seq data from single embryos. Embryos were collected either at stage 2, representing a time point before zygotic genome activation, and at the end of stage 5, representing a time point after widespread zygotic genome activation. Embryos were collected from 14 species, however we only used the data from 11 (*D. simulans, D. sechellia*, *D. melanogaster*, *D. yakuba*, *D. erecta*, *D. ananassae*, *D. persimilis*, *D. willistoni*, *D. mojavensis*, *D. virilis*) due to annotation deficiencies in the remaining 3. GTF files and references genomes from previously sequenced species [28] were downloaded from Flybase [77].

To determine whether a gene would be labeled as 'off' or as 'on', the overall distribution of FPKMs was analyzed. For all species, for both stage 2 and stage 5, a bimodal distribution appeared, with one peak at 0 and another at approximately $e^{3.5}$. The commonly used cutoff of FPKM=1 [78,79] was chosen as it falls between these two distributions.

To determine which genes were orthologues, we used the FlyBase orthology table "gene_orthologs_fb_2014_06_fixed.tsv".

Sequence Selection

Preliminary tests were performed to determine which regions were most likely to have regulatory elements. For each gene, several regions were extracted: 10kb upstream, 5kb upstream, 2kb upstream, 1kb upstream, 500bp upstream, 5' UTR, total introns, total exons, and 3' UTR. For each region, boundaries were obtained from the appropriate GTF and sequences were extracted using BioPython (Version 1.73, (Cock et al., 2009)). The 2kb upstream region showed the highest quality motifs (Fig S1),

530 and thus were used for matching motifs in external databases, measuring motif overlap between

531 species, analysing motif position distributions, and GO analysis. For these analysis (featured in figures 1

532 through 3 double check), UTRs were ignored as not every species had annotated UTRs.

Motif Discovery

We used HOMER [29] to discover motifs in test sets using the background sets as control FASTA files,

test and background sets are defined below. Deviations from the default settings include the use of the -

fasta flag to specify a custom background file. For stage 2 queries, the test FASTA files included genes

that had a FPKM >= 1 at stage 2 while the control FASTA files included genes that had an FPKM < 1. For

the stage 5 queries, the test FASTA files contained genes where the stage 5 FPKM >= 1 and the stage 2

FPKM < 1, while the control FASTA files included genes whose stage 5 FPKM < 1 and stage 2 FPKM < 1.

Additionally, we used the -p flag to utilize our computational resources more efficiently. We used -

norevopp flag in the case of strand-specific searches. Motif quality was evaluated based on the HOMER-

outputted q-values.

To validate the HOMER output files we used MEME[33] v4.12.0 and RSAT [80]. MEME was run using-

mod zoops -nmotifs 2 -minw 8 -maxw 12 -revcomp. The RSAT analysis uses the purge-sequences tool,

followed by oligo-analysis using the following parameters: -lth occ_sig 0 -uth rank 5000 -return

occ,proba,rank -2str -noov -quick_if_possible -seqtype dna -l 8, followed by pattern-assembly using the

following parameters: -v 1 -subst 1 -toppat 5000 -2str, followed by matrix-from-patterns using the

following parameters: -v 1 -logo -min_weight 5 -flanks 2 -max_asmb_nb 10 -uth Pval 0.00025 -bginput -

markov 0 -o purged_result.

Zygotic-Only Motifs

551 To examine the regulatory elements associated with zygotic transcripts, we defined genes as being

552 zygotic only if they were off at stage 2 (FPKM <1) and on at stage 5 (FPKM >1). It is necessary to impose

553   such a restriction, as a large percentage (approximately 95%) of genes that are maternally deposited are

554   still expressed at stage 5, and analysis of their regulatory mechanisms would confound the signal of

555   stage 5 only genes.

Motif Sharing

To determine weather motifs were shared between species, the HOMER-formatted motifs were

converted to meme-formatted motifs using chem2meme from the MEME Suit [33]. Tomtom, also from

the MEME Suit, was then used to find matching motifs, using default parameters. For a motif to be

considered shared with another species, the Tomtom output threshold of α = .05 was used. this

technique was used to calculate the similarity of motifs found in different species, as well as to evaluate

the similarity of different motif discovery strategies using MEME, RSAT, or HOMER with alternative

parameters.

To refine the results of shared motifs, we applied an additional quality cutoff. For stage 2, motifs were

first filtered for a q-value of less than 1e-100, and for stage 5, motifs were first filtered for a q-value of

1e-10. The difference in the cutoffs used at the two different stages was due to the differences in the

overall distribution of q-values for these stages due to a reduced number of zygotic-only genes (see

zygotic-only motifs above).

Because sharing was calculated on a by-species basis, it is possible that one species has a motif that

meets the criteria for being shared among all other species while other species' version of that same

motif failing to meet the criteria. This can occur, for example, when a motif is an intermediary version of

two motifs that fall just outside the cutoff.

To find proteins that bind to the discovered motifs, we used Tomtom to query JASPAR and Combined

Drosophila Databases using the default parameters[81].

Motif Position and Count

Motif position was determined by using the scanMotifGenomeWide tool to in the HOMER package.

Queries were performed by scanning the discovered motifs against the fasta files for each gene. The 5'

boundary of the motif was used as the motif position. For the motif counts per gene used in many

downstream analyses analysing motif position distributions, GO analysis, GC content analysis, and motif

strand analysis. We used this output and counted the occurrence of a given motif in the target region.

To quantify positional distribution similarity, we used the stats.anderson_ksamp function from the scipy

library V1.2.1[82]. Distributions were considered to be different at α = .05 after Bonferroni correction.

Transcript Enrichment by Tissue

Expression data for various adult tissues was downloaded from modENCODE[56]. To compare

enrichment for transcripts with different magnitudes of abundance, we applied an additional

normalization. For each transcript, transcript levels in FPKMs were divided by a scaling factor equal to

the average of the expression levels in ovaries. This normalization preserves the relative abundances

within each transcript, but allows for visualization of transcript levels with dramatically different overall

expression levels.

Expression by Position

591   *D. melanogaster* expression data by position was downloaded from modENCODE [56] for several tissue

592   types. Positions for each motif was determined as previously described in the Motif Position and Count

593   section above. For each instance of the motif of interest, we determined expression values in area from

594   -10kb to +10kb. Transcript abundance in FPKMs were then normalized by the average FPKM reported on

595   the track.

GO Analysis

GO Analysis

We used the R package clusterProfiler 3.10.1 [50] and the org.Dm.eg.db 3.7.0 [83] dictionary to perform

gene ontology (GO) analysis. For the stage 2 comparison, we generated a test set of the melanogaster

gene names for every gene in our dataset that was maternally deposited in at least any 7 of our species,

and performed an enrichment analysis using enrichGO's default parameters using a background set of

all *D. melanogaster* genes. For the stage 5 comparison, we generated a test set of the *D. melanogaster*

gene names for which at least two orthologues in our dataset showed zygotic-only expression (see

Zygotic-only motifs section above for definition). This threshold approximates the percent of the

genome that we observed to be zygotic only. We then performed an enrichment analysis using

enrichGO's default parameters using a background set of *D. melanogaster* genes that are not maternally

deposited in at least two species. This analysis therefore specifically examines the zygotically activated

genes in the context of genes that are "off" at stage 2 (FPKM<1 at this stage). For our analysis of stage 2

motifs, we generated a test set for each motif consisting of genes that contained that motif in at least

two species and were maternally deposited (FPKM > 1) in at least two species. We then performed an

enrichment analysis using enrichGO's default parameters using a background set of all *D. melanogaster*

genes. For our analysis of stage 5 motifs, we generated a test set for each motif using genes that were

represented by transcripts >1 FPKM at stage 5 in at least two species and had the motif of interest in at

least two species. We then performed an enrichment analysis using enrichGO's default parameters

using a background set of *D. melanogaster* genes that were represented by transcripts >1 FPKM at stage

5. To visualize our results, we employed the dotplot method for enrichGO objects, also from the

clusterProfiler package. For each motif, the top 3 GO terms were identified and added to the y-axis

labels. Whenever any GO category from another motif was identified as statistically significant (α = .05),

that GO category was shaded appropriately.

Model Fitting

Logistic regression was performed using the "glm" function in R, using the logit link function. As inputs,

we used the list of motifs generated from HOMER and their counts as described in the "Motif Position

and Count" section above. To avoid redundant motifs in our model, only motifs of size 10 were

considered. To evaluate the strand-specificity of motifs, we compared two generalized linear models

using the formulas indicated in S1_Model_Generation.pdf. To identify the most important motifs, the R

function stepAIC from the MASS library 7.3-51.4[84] was used to find generate an ordered list of motifs.

The base model used contained no additional features (chromatin state, etc). StepAIC was run 8 steps to

generate a short list of motifs for evaluation.

Analysis of physical clustering of co-expressed genes

To evaluate the effect of gene cluster size on expression, we iterated through each species for both

stage 2 and stage 5 and assigned sizes of co-expressed gene clusters on the chromosome, based on how

many adjacent genes were coexpressed, resulting in cluster size frequencies for each genome. Errors

were calculated using 95% confidence interval for a two-tailed binomial distribution.

To compare the clustering of different datasets with varying percents of "on" genes, we employed the

Wald–Wolfowitz runs test.

Tissue-specific RNA Levels

636 modENCODE tissue profiles [52] were downloaded from flybase.org . Flyatlas2 tissue profiles were

637 downloaded from http://flyatlas.gla.ac.uk/FlyAtlas2/ [53].

Gene length

639 To determine gene length, we examined the relevant line of the appropriate .GFF file and took the

640 difference between the end and the start positions.

Distance between genes

To determine the distance between genes, we look at the appropriate .GFF file and took the difference of positions between adjacent genes from transcription start site (TSS) to TSS.

Maternal deposition rates as compared to gene length, distance, and orientation

Genes were binned by category and by either distance or length. For the top plot, 150 bins of 70bp width were used. For the bottom plot, 60 bins of 70bp width were used and bins with fewer than 6 genes were disregarded. confidence intervals were calculated using the binomial distribution with α = .05 after Bonferroni.

GC content

GC content levels associated with each gene were evaluated by calculating the number of GC nucleotides within a sliding window of size 50bp for each of 1950 window positions to cover the upstream 2kb of each gene. To evaluate the first bin of each gene, the region from -1bp to -50bp was extracted, and the number of G and C nucleotides was counted. The result was divided by 50 to get the %GC for this window. To calculate the GC content for the next bin, this process was repeated on the region from -2bp to -51bp. Each bin had its GC content evaluated this way until the final bin of -451bp to -500bp. To evaluate how closely a particular upstream region resembled a maternally deposited-like distribution or a non maternally deposited-like distribution for the purposes of modeling, we calculated the average GC content for each position of maternally deposited, and not maternally deposited genes. Then for each gene, we measured the correlation between the GC content and that of both category averages. We used the difference in these correlations as a metric to evaluate similarity in GC content for each gene.

Acknowledgements

We would like to thank Joel Atallah for his work on the original dataset, Gizem Kalay, Anna Feitzinger, and Emily Cartwright for comments on the manuscript, and all members of the Lott Lab for feedback.

References

1. Driever W, Nüsslein-Volhard C. The bicoid protein determines position in the Drosophila embryo in a concentration-dependent manner. Cell. 1988;54: 95–104.

2. Lécuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, Cerovina T, et al. Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. Cell. 2007;131: 174–187.

3. Vastenhouw NL, Cao WX, Lipshitz HD. The maternal-to-zygotic transition revisited. Development. 2019;146. doi:10.1242/dev.161471

4. Schulz KN, Harrison MM. Mechanisms regulating zygotic genome activation. Nat Rev Genet. 2019;20: 221–234.

5. Ventos-Alfonso A, Ylla G, Belles X. Zelda and the maternal-to-zygotic transition in cockroaches. FEBS J. 2019; doi:10.1111/febs.14856

6. Navarro-Costa P, McCarthy A, Prudêncio P, Greer C, Guilgur LG, Becker JD, et al. Early programming of the oocyte epigenome temporally controls late prophase I transcription and chromatin remodelling. Nat Commun. 2016;7: 12331.

7. Mische S, Li M, Serr M, Hays TS. Direct observation of regulated ribonucleoprotein transport

across the nurse cell/oocyte boundary. Mol Biol Cell. 2007;18: 2254–2263.

8. Tadros W, Westwood JT, Lipshitz HD. The mother-to-child transition. Dev Cell. 2007;12: 847–

849.

9. Liang H-L, Nien C-Y, Liu H-Y, Metzstein MM, Kirov N, Rushlow C. The zinc-finger protein Zelda is a

key activator of the early zygotic genome in Drosophila. Nature. 2008;456: 400–403.

10.   Harrison MM, Li X-Y, Kaplan T, Botchan MR, Eisen MB. Zelda binding in the early

Drosophila melanogaster embryo marks regions subsequently activated at the maternal-to-zygotic

transition. PLoS Genet. 2011;7: e1002266.

11.   Akam M. The molecular basis for metameric pattern in the Drosophila embryo.

Development. 1987;101: 1–22.

12.   Ingham PW. The molecular genetics of embryonic pattern formation in Drosophila.

Nature. 1988;335: 25–34.

13.   Tadros W, Goldman AL, Babak T, Menzies F, Vardy L, Orr-Weaver T, et al. SMAUG is a

major regulator of maternal mRNA destabilization in Drosophila and its translation is activated by

the PAN GU kinase. Dev Cell. 2007;12: 143–155.

14.   Benoit B, He CH, Zhang F, Votruba SM, Tadros W, Westwood JT, et al. An essential role

for the RNA-binding protein Smaug during the Drosophila maternal-to-zygotic transition.

Development. 2009;136: 923–932.

15.   Laver JD, Li X, Ray D, Cook KB, Hahn NA, Nabeel-Shah S, et al. Brain tumor is a sequence-

specific RNA-binding protein that directs maternal mRNA clearance during the Drosophila

maternal-to-zygotic transition. Genome Biol. 2015;16: 94.

16.   Bushati N, Stark A, Brennecke J, Cohen SM. Temporal reciprocity of miRNAs and their

targets during the maternal-to-zygotic transition in Drosophila. Curr Biol. 2008;18: 501–506.

17.   Becalska AN, Gavis ER. Lighting up mRNA localization in Drosophila oogenesis.

Development. 2009;136: 2493–2503.

18.   Clark A, Meignin C, Davis I. A Dynein-dependent shortcut rapidly delivers axis

determination transcripts into the Drosophila oocyte. Development. 2007;134: 1955–1965.

19.   Barckmann B, Simonelig M. Control of maternal mRNA stability in germ cells and early

embryos. Biochim Biophys Acta. 2013;1829: 714–724.

20.   Cui J, Sackton KL, Horner VL, Kumar KE, Wolfner MF. Wispy, the Drosophila homolog of

GLD-2, is required during oogenesis and egg activation. Genetics. 2008;178: 2017–2029.

21.   Benoit P, Papin C, Kwak JE, Wickens M, Simonelig M. PAP- and GLD-2-type poly(A)

polymerases are required sequentially in cytoplasmic polyadenylation and oogenesis in Drosophila.

Development. 2008;135: 1969–1979.

22.   Sallés FJ, Lieberfarb ME, Wreden C, Gergen JP, Strickland S. Coordinate initiation of

Drosophila development by regulated polyadenylation of maternal messenger RNAs. Science.

1994;266: 1996–1999.

23.   Temme C, Simonelig M, Wahle E. Deadenylation of mRNA by the CCR4-NOT complex in

Drosophila: molecular and developmental aspects. Front Genet. 2014;5: 143.

24.   Atallah J, Lott SE. Evolution of maternal and zygotic mRNA complements in the early

Drosophila embryo. PLoS Genet. 2018;14: e1007838.

25.   Bownes M. A photographic study of development in the living embryo of Drosophila

melanogaster. J Embryol Exp Morphol. 1975;33: 789–801.

26.   Campos-Ortega JA, Hartenstein V. The Embryonic Development of Drosophila

melanogaster. Springer, Berlin, Heidelberg; 1985.

27.   Nègre N, Brown CD, Ma L, Bristow CA, Miller SW, Wagner U, et al. A cis-regulatory map

of the Drosophila genome. Nature. 2011;471: 527–531.

28.   Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, et al. Evolution of

genes and genomes on the Drosophila phylogeny. Nature. 2007;450: 203–218.

29.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations of

lineage-determining transcription factors prime cis-regulatory elements required for macrophage

and B cell identities. Mol Cell. 2010;38: 576–589.

30.   De Renzis S, Elemento O, Tavazoie S, Wieschaus EF. Unmasking activation of the zygotic

genome using chromosomal deletions in the Drosophila embryo. PLoS Biol. 2007;5: e117.

31.   Thomsen S, Anders S, Janga SC, Huber W, Alonso CR. Genome-wide analysis of mRNA

decay patterns during early Drosophila development. Genome Biol. 2010;11: R93.

32.   Lott SE, Villalta JE, Zhou Q, Bachtrog D, Eisen MB. Sex-specific embryonic gene

expression in species with newly evolved sex chromosomes. PLoS Genet. 2014;10: e1004159.

33.   Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME SUITE: tools

for motif discovery and searching. Nucleic Acids Res. 2009;37: W202–8.

34.   Ramírez F, Bhardwaj V, Arrigoni L, Lam KC, Grüning BA, Villaveces J, et al. High-

resolution TADs reveal DNA sequences underlying genome organization in flies. Nat Commun.

2018;9: 189.

35.   Zabidi MA, Arnold CD, Schernhuber K, Pagani M, Rath M, Frank O, et al. Enhancer-core-

promoter specificity separates developmental and housekeeping gene regulation. Nature.

2015;518: 556–559.

36.   Liu MM, Davey JW, Jackson DJ, Blaxter ML, Davison A. A conserved set of maternal

genes? Insights from a molluscan transcriptome. Int J Dev Biol. 2014;58: 501–511.

37.   Matsukage A, Hirose F, Yoo M-A, Yamaguchi M. The DRE/DREF transcriptional regulatory

system: a master key for cell proliferation. Biochim Biophys Acta. 2008;1779: 81–89.

38.   Yang J, Ramos E, Corces VG. The BEAF-32 insulator coordinates genome organization

and function during the evolution of Drosophila species. Genome Res. 2012;22: 2199–2207.

39.   Nègre N, Brown CD, Shah PK, Kheradpour P, Morrison CA, Henikoff JG, et al. A

comprehensive map of insulator elements for the Drosophila genome. PLoS Genet. 2010;6:

e1000814.

40.   Maksimenko O, Bartkuhn M, Stakhov V, Herold M, Zolotarev N, Jox T, et al. Two new

insulator proteins, Pita and ZIPIC, target CP190 to chromatin. Genome Res. 2015;25: 89–99.

41.   Li J, Gilmour DS. Distinct mechanisms of transcriptional pausing orchestrated by GAGA

factor and M1BP, a novel transcription factor. EMBO J. 2013;32: 1829–1841.

42.   Chen K, Johnston J, Shao W, Meier S, Staber C, Zeitlinger J. A global change in RNA

polymerase II pausing during the Drosophila midblastula transition. Elife. 2013;2: e00861.

43.   Levine M. Paused RNA polymerase II as a developmental checkpoint. Cell. 2011;145:

502–511.

44.   Benyajati C, Mueller L, Xu N, Pappano M, Gao J, Mosammaparast M, et al. Multiple

isoforms of GAGA factor, a critical component of chromatin structure. Nucleic Acids Res. 1997;25:

3345–3353.

45.   Tsai S-Y, Chang Y-L, Swamy KBS, Chiang R-L, Huang D-H. GAGA factor, a positive

regulator of global gene expression, modulates transcriptional pausing and organization of

upstream nucleosomes. Epigenetics Chromatin. 2016;9: 32.

46.   Granok H, Leibovitch BA, Shaffer CD, Elgin SC. Chromatin. Ga-ga over GAGA factor. Curr

Biol. 1995;5: 238–241.

47.   Harrison MM, Botchan MR, Cline TW. Grainyhead and Zelda compete for binding to the

promoters of the earliest-expressed Drosophila genes. Dev Biol. 2010;345: 248–255.

48.   Ohler U, Liao G-C, Niemann H, Rubin GM. Computational analysis of core promoters in

the Drosophila genome. Genome Biol. 2002;3: RESEARCH0087.

49.   Lis M, Walther D. The orientation of transcription factor binding site motifs in gene

promoter regions: does it matter? BMC Genomics. 2016;17: 185.

50.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological

themes among gene clusters. OMICS. 2012;16: 284–287.

51.   R Core Team. R: A Language and Environment for Statistical Computing [Internet].

Vienna, Austria: R Foundation for Statistical Computing; 2014. Available: http://www.R-project.org/

52.   Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. Diversity and

dynamics of the Drosophila transcriptome. Nature. 2014;512: 393–399.

53.   Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. FlyAtlas 2: a new version of the

Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data. Nucleic

Acids Res. 2018;46: D809–D815.

54.   Vogel C, Marcotte EM. Insights into the regulation of protein abundance from proteomic

and transcriptomic analyses. Nat Rev Genet. 2012;13: 227–232.

55.   Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, et al. The

developmental transcriptome of Drosophila melanogaster. Nature. 2011;471: 473–479.

56.   modENCODE Consortium, Roy S, Ernst J, Kharchenko PV, Kheradpour P, Negre N, et al.

Identification of functional elements and regulatory circuits by Drosophila modENCODE. Science.

2010;330: 1787–1797.

57.   Bauke A-C, Sasse S, Matzat T, Klämbt C. A transcriptional network controlling glial

development in the Drosophila visual system. Development. 2015;142: 2184–2193.

58.   Gurudatta BV, Yang J, Van Bortle K, Donlin-Asp PG, Corces VG. Dynamic changes in the

genomic localization of DNA replication-related element binding factor during the cell cycle. Cell

Cycle. 2013;12: 1605–1615.

59.   Ulianov SV, Khrameeva EE, Gavrilov AA, Flyamer IM, Kos P, Mikhaleva EA, et al. Active

chromatin and transcription play a key role in chromosome partitioning into topologically

associating domains. Genome Res. 2016;26: 70–84.

60.   Hou C, Li L, Qin ZS, Corces VG. Gene density, transcription, and insulators contribute to

the partition of the Drosophila genome into physical domains. Mol Cell. 2012;48: 471–484.

61.   Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI GEO:

archive for functional genomics data sets--update. Nucleic Acids Res. 2013;41: D991–5.

62.   Bradley JV. Distribution-free statistical tests. Prentice-Hall; 1968.

63.   Artieri CG, Fraser HB. Transcript length mediates developmental timing of gene

expression across Drosophila. Mol Biol Evol. 2014;31: 2879–2889.

64.   Heyn P, Kircher M, Dahl A, Kelso J, Tomancak P, Kalinka AT, et al. The earliest

transcribed zygotic genes are short, newly evolved, and different across species. Cell Rep. 2014;6:

285–292.

65.   Zhang L, Kasif S, Cantor CR, Broude NE. GC/AT-content spikes as genomic punctuation

marks. Proc Natl Acad Sci U S A. 2004;101: 16855–16860.

66.   Tadros W, Lipshitz HD. The maternal-to-zygotic transition: a play in two acts.

Development. 2009;136: 3033–3042.

67.   Van Bortle K, Nichols MH, Li L, Ong C-T, Takenaka N, Qin ZS, et al. Insulator function and

topological domain border strength scale with architectural protein occupancy. Genome Biol.

2014;15: R82.

68.   Jambor H, Surendranath V, Kalinka AT, Mejstrik P, Saalfeld S, Tomancak P. Systematic

imaging reveals features and changing localization of mRNAs in Drosophila development. Elife.

2015;4. doi:10.7554/eLife.05003

69.   Dej KJ, Spradling AC. The endocycle controls nurse cell polytene chromosome structure

during Drosophila oogenesis. Development. 1999;126: 293–303.

70.   Zhimulev IF, Belyaeva ES, Semeshin VF, Koryakov DE, Demakov SA, Demakova OV, et al.

Polytene Chromosomes: 70 Years of Genetic Research. International Review of Cytology. Academic

Press; 2004. pp. 203–275.

71.   Abruzzi K, Chen X, Nagoshi E, Zadina A, Rosbash M. Chapter Seventeen - RNA-seq

Profiling of Small Numbers of Drosophila Neurons. In: Sehgal A, editor. Methods in Enzymology.

Academic Press; 2015. pp. 369–386.

72.   Liang J, Lacroix L, Gamot A, Cuddapah S, Queille S, Lhoumaud P, et al. Chromatin

immunoprecipitation indirect peaks highlight long-range interactions of insulator proteins and Pol II

pausing. Mol Cell. 2014;53: 672–681.

73.   Phillips-Cremins JE, Corces VG. Chromatin insulators: linking genome organization to

cellular function. Mol Cell. 2013;50: 461–474.

74.   Matzat LH, Lei EP. Surviving an identity crisis: a revised view of chromatin insulators in

the genomics era. Biochim Biophys Acta. 2014;1839: 203–214.

75.   Matzat LH, Dale RK, Moshkovich N, Lei EP. Tissue-specific regulation of chromatin

insulator function. PLoS Genet. 2012;8: e1003069.

76.   Vardy L, Orr-Weaver TL. Regulating translation of maternal messages: multiple

repression mechanisms. Trends Cell Biol. 2007;17: 547–554.

77.   Gramates LS, Marygold SJ, Santos GD, Urbano J-M, Antonazzo G, Matthews BB, et al.

FlyBase at 25: looking to the future. Nucleic Acids Res. 2017;45: D663–D671.

78.   Brooks MJ, Rajasimha HK, Roger JE, Swaroop A. Next-generation sequencing facilitates

quantitative analysis of wild-type and Nrl(-/-) retinal transcriptomes. Mol Vis. 2011;17: 3034–3054.

79.   Tao T, Zhao L, Lv Y, Chen J, Hu Y, Zhang T, et al. Transcriptome sequencing and

differential gene expression analysis of delayed gland morphogenesis in Gossypium australe during

seed germination. PLoS One. 2013;8: e75323.

80.   Medina-Rivera A, Defrance M, Sand O, Herrmann C, Castro-Mondragon JA, Delerce J, et

al. RSAT 2015: Regulatory Sequence Analysis Tools. Nucleic Acids Res. 2015;43: W50–6.

81.   Khan A, Fornes O, Stigliani A, Gheorghe M, Castro-Mondragon JA, van der Lee R, et al.

JASPAR 2018: update of the open-access database of transcription factor binding profiles and its

web framework. Nucleic Acids Res. 2018;46: D260–D266.

82.   Jones E, Oliphant T, Peterson P, Others. SciPy: Open source scientific tools for Python

[Internet]. Available: http://www.scipy.org/

83.   Carlson M. org.Dm.eg.db: Genome wide annotation for Fly. 2018.

84.   Venables WN, Ripley BD. Modern Applied Statistics with S [Internet]. Fourth. New York:

Springer; 2002. Available: http://www.stats.ox.ac.uk/pub/MASS4

Figure Captions

Table 1: A summary of the top ranked motifs. HOMER was used to find motifs enriched in the 2kb

windows upstream of maternally deposited genes (stage 2) and zygotically transcribed genes (stage 5).

Sequence logo shows the consensus motif where the probability of each base is proportional to its

representative character. P-value is given by HOMER. %target represents the percent of either

maternally deposited or zygotically expressed genes that contain at least one instance of the motif.

%background indicates the percent of all genes that contain this motif. Best match indicates protein

with a previously identified binding site that mostly closely matches the discovered motif (see Methods).

Fig 1: Motifs associated with maternal deposition are largely shared across species, zygotic motifs are

likely to be species-specific. For each analysis represented in A-D, motif enrichment was determined for

each group of genes at each stage (all maternally deposited genes at stage 2; or zygotic genes at stage 5)

separately in each species, then lists of enriched motifs at each stage were compared across species. For

stage 2 motifs, we required motifs to have a -log qvalue > 100, while for stage 5 motifs we required

motifs to have a -log qvalue > 10 (see Methods). (A, C) Percent of motif content in the upstream region

that is found to be shared between species at stage 2 and stage 5, respectively. The number of species

that share each motif is indicated by the color of the bar. Note that in stage 2, a large majority of motifs

are shared in all (11 species) or almost all (9 or 10 species), with the exception of *D. pseudoobscura* and

D. persimilis, sister species that share common motifs between themselves but are different from the

rest of the species. Zygotic motifs identified at stage 5 are much more likely to be species specific or

shared by only a couple of species. (B, D) Number of motifs shared between each pair of species at stage

2 and stage 5, respectively. Comparisons of one species to itself indicate the total number of motifs that

fit quality criteria discovered in that species. Comparing the number of shared motifs between pairs of

species, there is some signal of the phylogeny in stage 2 (B), with *D. melanogaster* subgroup species

sharing more motifs in common with one another than they do with the more distantly related species,

and *D. pseudoobscura* and *D. persimilis* with the highest number of motifs in common but the most

differences from the remaining species. For stage 5 (D), apparent patterns include both the number of

species-specific motifs (diagonal) and less apparent phylogenetic structure. (E) Conservation of top

motifs in orthologous genes across species. Y-axis indicates all of the instances of the motif of interest

within the upstream region. Coloration represents how many species' orthologues also contain that

motif. In general, top motifs at the zygotic stage (stage 5) are more likely to be conserved in orthologous

genes at this stage. This sets up a contrast with parts A-D, where maternal deposition is broadly

associated with a shared set of motifs across species, but part E shows that orthologous maternal genes

are less likely to share a specific motif.

Fig 2: Top GO terms show that motifs regulate broader set of genes at the maternal stage, and a more

specific set of developmentally associated genes at the zygotic stage. (A) GO terms associated with each

stage. Note that the set of identified GO categories does not overlap between stages. (B) GO terms

associated with top motifs in stage 2, where a majority of motifs are associated with similar broad GO

categories (C) GO terms associated with top motifs in stage 5, some motifs are associated with the same

categories, some appear to be more specialized, with identified categories showing more specificity

than categories associated with stage 2

Fig 3: Identified maternal regulators are ovary enriched, as is their effect on transcription (A) RNA levels

of putative binding proteins by tissue type. Transcript abundances within each gene have been

normalized such that the average abundance in ovaries is equal to 1. While identified maternal

regulators have regulatory functions in multiple tissue types, they are highly enriched in ovaries

compared to other tissues. (B) Average normalized expression levels versus proximity to motif by tissue

type. Normalization was performed by dividing each expression value by the average expression from

9.9-10kb away. While binding sites for identified maternal regulators are present in multiple tissues, the

effect on gene expression is stronger in ovaries compared to other tissues.

Fig 4: Chromatin stage and maternal deposition. For the analyses in A and B, genes were categorized as

either expressed or not expressed (see Methods) and adjacent expressed genes were considered to be

clustered, with a cluster size equal to the number of constituent genes. (A) Physical clustering of

maternally deposited genes along the chromosome, in a representative species (*D. simulans*). The

shaded blue region represents the observed frequency of co-expressed maternal gene clusters of

various sizes. The red region represents the 95% CI constructed with 10,000 bootstrap iterations.

Maternal genes are co-expressed in clusters along the chromosome more often than expected, given

the percent of the genome that is transcribed at this stage. (B) Physical clustering of co-expressed genes

on chromosomes in various tissue types. In order to compensate for differing proportions of the

genome that are expressed in each tissue type, physical clustering was measured by performing a Wald-

Wolfowitz runs test and taking the z-score (see Methods). Maternally expressed genes, represented by

stage 2 embryos, show the highest proportion of physical clustering of co-expressed genes, though

other tissues such as intestinal stem cells and larval CNS also have highly physically clustered co-

expressed genes. (C) Gene length by number of adjacent maternally expressed genes, "open" indicating

both adjacent genes are expressed, "border" indicating that one is expressed, and "closed" indicating

that neither are expressed. Genes that with more expressed neighbors are more likely to be maternally

deposited, regardless of length. Genes without expressed neighbors are less likely to be maternally

deposited, with the odds increasing as length increases. (D) Odds of maternal deposition versus distance

to the nearest upstream gene by upstream expression and strand. Distance is measured by from

transcription start site (TSS) to TSS. When the upstream gene is maternally deposited, odds of maternal

deposition are high, but decrease with distance regardless of strand. When the upstream gene is not

maternally deposited, odds of maternal deposition are low and have a strand-dependent relationship

with distance.

Fig 5: Stage-specific genes are more likely to be different from their chromatin neighborhood. *D.*

simulans was chosen as a representative species. (A) Cluster size distribution of maternal only genes

(green bars) compared with the expected frequencies based on the overall cluster size frequencies

observed at stage two (blue region). The expected frequencies are based on the distribution in Fig 4A

multiplied by a scale factor equal to the proportion of maternally deposited genes that are maternal

only, with the shaded region representing a 95% confidence interval. (B) Cluster size distribution for

zygotic only genes (green bars) compared with the expected frequencies based on the overall cluster

size frequencies observed at stage 5 (blue region) in a manner similar to Fig 5A. the shaded region

represents a 95% confidence interval. For both stages, stage-specific genes are more likely to be the

single gene (or one of a small number of genes) that are expressed where their neighboring genes are

not, representing small numbers of "on" genes in an "off" chromatin environment.

Supporting Information Figure Captions

S1 Table: A summary of the top ranked zygotic motifs. Motifs were selected if had enrichment if they

were enriched in the combined upstream regions of all species with a q-value < 1e-50 and a Tomtom

match to any motif in an existing database with q < .1. If there were more than one, the best two

matches to motifs in existing databases were reported in the Best Match column. Some motifs are

plausible binding sites for known embryonic regulators.

S1 Fig: Distribution of motif qualities by location in a representative species in each stage. *D. ananassae*

was selected as a representative species. Motif qualities are given by the negative natural logarithm of

the q-value outputted by HOMER. High quality motifs enriched for stage 2 (A) are most likely to be

found in the 2kb upstream of a gene. Motifs for stage 5 are generally less high quality by this metric, and

while the highest quality tend to also be enriched 2kb upstream, some are enriched in 2kb upstream

regions of non-expressed genes or enriched in exons.

S2 Fig: Representative positional distributions of motifs. Distributions for both maternally deposited

genes ("on") and non- maternally deposited genes ("off") are shown. (A) The positional distribution of

the DREF motif, which follows the same pattern as M1BP, Zipic, Ohler-6, and E-box, and many motifs

without identified factors that bind them. These motifs are found upstream of maternally deposited

genes (red), with a higher frequency closer to the transcription start site. They are not found with any

frequency in non-maternally deposited genes (blue). (B,C) Positional distribution patterns of some rare,

undocumented motifs. In both, we see that the motif is more enriched in maternally deposited genes

than in non-maternally deposited genes, but that the enrichment difference is less than those motifs

represented by (A) above. In (B), this motif is most highly enriched upstream, less enriched around the

transcription start site (TSS), and more highly enriched again downstream of the TSS (though less so

than upstream). In (C), we see the highest enrichment downstream of the TSS, with a dip in enrichment

around the TSS, and less enrichment upstream of the TSS than downstream

S3 Fig: GC content of the region upstream of the TSS. GC content for each gene in a sliding window with

50bp width is summed for each gene in the category. (A) Maternally deposited genes. (B) Non-

maternally deposited genes. (C) Zygotic only genes. Note the high number of genes with higher GC

content immediately upstream of maternally deposited genes, and the lower GC content upstream of

this GC-enriched region.

	Logo	-log p-value	%Target	%Background	Best Match	Description
	TATCGATAGT	2091	35.49%	12.32%	DREF	• "Master key-like factor for cell proliferation" (Akio Matsukage et al. 2008) • Shares binding site with BEAF-32
	TATCGATAGT	2091	35.49%	12.32%	BEAF-32	• Insulator (Yang, Ramos, and Corces 2012; Nègre et al. 2010) • Shares binding site with DREF
Stage 2	CAGTGTGACCGT	1591	27.85%	9.42%	M1BP	• Causes PolII to pause on the gene gene (Li and Gilmour 2013)
	TTTGCAACACTG	934	27.35%	12.73%	ZIPIC	• Recruits insulator CP190 (Maksimenko et al. 2015).
	CGACGGTATTTT	843	40.25%	23.98%	Ohler-6	• Commonly found between TAD boundaries (Ramirez et al 2018)
	CAAACAGCTGTT	692	20.13%	9.05%	E-box	• Regulates gene expression
	GTCTACCTGCTC	163	18.45%	8.77%	Zld	• "master regulator of genome activation"
Stage 5	CGCTCGCTCT	84	35.74%	26.05%	Trl	• Required for embryogenesis • Known to regulate developmental genes
	CGTCTCTCTC	56	48.41%	39.9%	Trl	• Required for embryogenesis • Known to regulate developmental genes

Table 1

Fig 1

A

B

C

Fig 2

Fig 3

Fig 4

A

B

Fig 5

Supplementary

Location	Motif	Best Match	Tomtom q-value
Upstream	AHTATCGATA	Beaf-32	0.00809
Upstream	AHTATCGATA	Dref	0.03381
Upstream	BGYGYGYGCG	sr	0.01908
Upstream	BGYGYGYGCG	klu	0.04051
Upstream	CGCTCDCNCK	Trl	0.00634
Upstream	CGGCGCAGCG	Mad	0.08778
Upstream	CGTCTCTCTC	GAF	0.01615
Upstream	CGTCTCTCTC	Trl	0.01615
Upstream	GAGAGAGAGA	GAF	0.00001
Upstream	GAGAGAGAGA	Trl	0.00001
Upstream	KCGCTCGCTC	Trl	0.00059
Upstream	KCTACCTGYB	Zld	0.00002
Upstream	NNNNTACCTG	Zld	0.09762
Upstream	NNNNTACCTG	vvl	0.09762
Exon	TACATACATA	Top2	0.00277

S1 Table

A

B

S1 Fig

A

B

C

S2 Fig

A

B

C

S3 Fig