Cervicovaginal tissue residence imprints a distinct differentiation program upon memory

CD8 T cells

Veronica Davé[1,2], E. Fabian Cardozo-Ojeda[1], Florian Mair[1], Jami Erickson[1], Amanda S.

Woodward-Davis[1], Amanda Koehne[3], Andrew Soerens[1], Julie Czartoski[1], Candice Teague[1],

Nicole Potchen[1,2], Susanne Oberle[4], Dietmar Zehn[4,†], Joshua T. Schiffer[1,5,6], Jennifer M.

Lund[1,2,*], Martin Prlic[1,2,7*]

[1] Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle,

WA 98109

[2] Graduate Program in Pathobiology, Department of Global Health, University of Washington,

Seattle, WA 98195

[3] Comparative Pathology, Fred Hutchinson Cancer Research Center, Seattle, WA 98109

[4] Division of Immunology and Allergy, Department of Medicine, Lausanne University Hospital,

1011 Lausanne, Switzerland

[5] Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109

[6] Department of Medicine, University of Washington, Seattle, WA 98195

[7] Department of Immunology, University of Washington, Seattle, WA 98195

[*]Co-corresponding authors (jlund@fredhutch.org, mprlic@fredhutch.org)

[†]Present address: Division of Animal Physiology and Immunology, School of Life Sciences

Weihenstephan, Technical University of Munich, 85354 Freising, Germany

Running title: Cervicovaginal CD8 T cell memory

Summary (~40 words)

The tissue-resident memory (T_{RM}) CD8 T cell compartment in human and mouse cervicovaginal

tissue (CVT) is remarkably similar. The CVT T_{RM} compartment is maintained autonomously and

does not reach phenoypical or numerical equilibrium. The numerical decline leads to impaired

viral control in a secondary challenge.

Abstract (160 words)

Tissue-resident memory CD8 T cells (CD8 T_{RM}) are critical for maintaining barrier immunity.

CD8 T_{RM} have been mainly studied in the skin and gut with recent studies suggesting that the

signals that control tissue-residence and phenotype are highly tissue-dependent. We examined

the T cell compartment in healthy human cervicovaginal tissue (CVT) and found that most CD8

T cells were granzyme B^+ and TCF-1^-. To address if this phenotype is driven by CVT tissue-

residence, we used a mouse model to control for environmental factors. Using localized and

systemic infection models, we found that CD8 T_{RM} in the mouse CVT gradually acquired a

granzyme B^+, TCF-1^- phenotype as seen in human CVT. In contrast to CD8 T_{RM} in the gut,

these CD8 T_{RM} were not stably maintained regardless of the initial infection route, which led to

reductions in local immunity. Our data show that residence in the CVT is sufficient to

progressively shape the size and function of its CD8 T_{RM} compartment.

Introduction

After infection or immunization, naive CD8 T cells differentiate into several major populations of memory T cells with distinct functions and trafficking patterns. Tissue-resident memory CD8 T cells (CD8 T_{RM}) are a major subset defined by the fact that they do not recirculate via blood and lymph, instead remaining in the tissue sites where they were initially seeded by effector T cells during the primary response (Gebhardt et al., 2009; Masopust et al., 2010; Masopust and Soerens, 2019; Szabo et al., 2019; Wakim et al., 2008). Because of their privileged location, CD8 T_{RM} are uniquely poised to respond to subsequent encounters with their cognate antigen by directly killing infected cells (Casey et al., 2012; Steinbach et al., 2016), proliferating to boost the local CD8 T cell pool (Beura et al., 2018; Park et al., 2018), and activating and recruiting additional immune cells from both resident and circulating populations (Ariotti et al., 2014; Schenkel et al., 2013).

Many factors influence the formation of T_{RM}, including the route of priming, exposure to antigen and inflammation, and the type of tissue in which residency is established. For example, the presence of local antigen boosts the size and function of T_{RM} populations within the matched tissue site (Davies et al., 2017; Gebhardt et al., 2009; Khan et al., 2016; Schenkel et al., 2013). Similarly, local inflammation enhances the number and function of T_{RM} in an antigen-independent manner (Mackay et al., 2012; Shin and Iwasaki, 2012) and cytokine cues such as TGF-β, IL-15, and CCL5 appear to directly regulate T_{RM} formation and maintenance (Bergsbaken et al., 2017; Iijima and Iwasaki, 2014; Mackay et al., 2015; Zhang and Bevan, 2013).

Despite the importance of local exposure to antigen and inflammation in T_{RM} development, strong evidence also exists that T_{RM} can broadly seed distal tissues after both systemic and local immunization or infection. Systemic immunization and infection models are routinely used in mice to generate T_{RM} populations across many body sites (Masopust et al.,

2001; Milner et al., 2017; Pope et al., 2001; Reinhardt et al., 2001; Steinert et al., 2015).

Likewise, localized infections such as herpes simplex virus (HSV) and vaccinia virus seed CD8

T_{RM} at distal tissue sites, albeit to a much lower extent than at the primary site of infection

(Davies et al., 2017; Jiang et al., 2012; Khan et al., 2016; Masopust et al., 2004). Together, these

studies suggest that T_{RM} can develop within diverse tissue sites across a range of levels of

inflammation and antigen availability, and it is currently unclear how the combination of these

variables may ultimately tune the characteristics of the resulting T_{RM} population.

The tissue of residence itself also impacts the phenotype of CD8 T_{RM} populations. CD8

T_{RM} have been extensively studied in skin of both humans and mice, and it is now well-known

that skin-resident CD8 T cells express the putative residency markers CD69 and integrin α_E

(CD103), respond to local HSV infection, and rely on TGF-β and IL-15 signals for their

development (Gebhardt et al., 2009; Mackay et al., 2013; Watanabe et al., 2015; Zhu et al.,

2013). Many reports suggest that CD8 T_{RM} in other organs may differ in their phenotype or

developmental requirements. For example, it is well-documented that CD69 and CD103 are not

necessarily expressed by T_{RM} in the uterus or pancreas (Steinert et al., 2015) but may be

required by T_{RM} in other organs such as the salivary gland, lung, or kidney (Hofmann and

Pircher, 2011; Takamura et al., 2016; Walsh et al., 2019). Likewise, some organs such as the

small intestine maintain T_{RM} that constitutively express granzyme B in the absence of antigen or

re-challenge (Casey et al., 2012; Kim et al., 1997), a phenotype which has not been observed as

robustly in other tissue sites, including the lung and uterus (Beura et al., 2019; Piet et al., 2011).

It remains an open question whether CD8 T_{RM} phenotype is inextricably linked to tissue of

residence, or whether alterations to the priming method or the inflammatory milieu within a

tissue could elicit features such as constitutive granzyme B expression in tissue sites where

they are not normally observed.

Mucosal barrier tissues contain large populations of CD8 T_{RM}, and pathogens that infect

mucosal barrier surfaces represent a significant global health burden. Specifically, the study of

CD8 T cell memory within the cervicovaginal tissue (CVT) has broad importance in

understanding the pathogenesis and vaccinology of sexually-transmitted infections. Compared

to memory CD8 T cells in the upper female reproductive tract and other mucosal and lymphoid

tissues, the phenotypic and functional characteristics of memory CD8 T cells in the CVT remain

relatively understudied. We recently reported that the CD8 T cell compartment in human CVT

includes a subset of CD8 T cells that robustly express granzyme B (Pattacini et al., 2019). It is

unclear to what extent the distinctive features we observed in human CVT may be driven by

recurring local infections, exposure to inflammatory cues, or a result of signals intrinsically

associated with the cervicovaginal microenvironment.

Here, we report that CD8 T cells isolated from healthy human CVT lacked expression of

the memory-associated transcription factor TCF-1 and resembled effector or terminally-

differentiated cells rather than a self-renewing memory T cell population. To determine if this

characteristic was driven by residence in the tissue itself or was a consequence of local

exposure to antigenic insults or inflammation in human CVT, we utilized multiple immunization

strategies to assess memory CD8 T cell differentiation and maintenance in the mouse CVT. We

found that memory CD8 T cells in the mouse CVT at later memory time points closely

resembled the CD8 T cells found in human CVT regardless of the initial immunization route and

early phenotypic differences. Memory CD8 T cell numbers where stably maintained in the

periphery and gut, but gradually declined in the CVT over five months post-immunization, which

was associated with a delay in viral control upon HSV vaginal challenge. We conclude that

residence in the CVT is sufficient to alter the canonical differentiation and maintenance program

of memory CD8 T cells.

Results

Healthy human cervicovaginal tissue contains CD8 T cells that express granzyme B and lack TCF-1

In order to study the phenotypic profile of CD8 T cells within human CVT from healthy women without any genital infections, we isolated lymphocytes from human vaginal biopsies via enzymatic digestion (**Fig. S1A**) and observed that the majority of memory T cells in this tissue site expressed the residency markers CD69 and CD103 (**Fig. 1A-B**). A large proportion of these CD8 T cells expressed granzyme B *ex vivo* (**Fig. 1C**) and the majority of granzyme B$^+$ cells co-expressed CD103 (**Fig. 1D**), suggesting that they were potentially resident within the tissue rather than transient inflammatory cells. We observed that the memory CD8 T cell compartment in human CVT lacked expression of TCF-1 (**Fig. 1E**), a transcription factor associated with self-renewal potential in CD8 T cells (Zhou et al., 2010). These data indicated that human memory CD8 T cells within the CVT may possess functional and proliferative characteristics distinct from such cells in other tissue locations.

Vaginal immunization in mice results in a CD8 T cell compartment that resembles that of human CVT

We next wanted to address if this human CVT CD8 T cell phenotype was attributable to ongoing basal levels of antigenic insult or inflammation in the tissue, or might represent more generalized characteristics of CD8 T cells in CVT. To stringently control for external environmental factors, we sought to generate a comparable population of memory CD8 T cells in mouse CVT. We adoptively transferred naive gBT-I CD8 T cells specific to the SSIEFARL epitope from HSV and primed these cells by vaginal infection with non-lethal thymidine-kinase deficient HSV-2 (HSV-2 TK-) (**Fig. 2A**). One month following infection, HSV gB tetramer$^+$ CD8 T cells in the CVT expressed CD69, CD103, and granzyme B in similar proportions to those observed in human CVT (**Fig. 2B-C**). In addition, we found that these CD8 T cells largely lacked

expression of TCF-1, especially compared to gB tetramer⁺ cells in the vaginal-draining lymph

nodes (dLN) (**Fig. 2D**). Given that these mice had received a vaginal infection, we wondered if

this granzyme B⁺ TCF-1⁻ phenotype might be a result of ongoing inflammatory responses in the

CVT. Indeed, analysis of hematoxylin and eosin-stained sections of CVT from infected mice

revealed that clusters of inflammatory cells remained in the CVT lamina propria as long as 22d

after infection (**Fig. 2E**), suggesting that inflammation was not resolved by this time point. Thus,

our data suggested that prolonged local tissue inflammation could be driving this distinct

memory CD8 T cell phenotype within the CVT.

Systemic immunization with LM-gB elicits stable HSV-specific CD8 T cell memory in secondary

lymphoid organs and the small intestine, but not the CVT

To determine if prolonged tissue inflammation drives this CVT memory T cell phenotype,

we used a systemic immunization approach to prime gBT-I cells in the absence of significant

vaginal inflammation. We generated a recombinant strain of *Listeria monocytogenes* that

expresses the HSV-derived SSIEFARL peptide (LM-gB) and immunized mice intravenously

(**Fig. 3A**). We confirmed that this immunization strategy did not result in detectable levels of

vaginal inflammation compared to uninfected mice, and caused significantly less inflammation

than observed in mice vaginally immunized with HSV-2 TK- (**Fig. 3B**). In addition, immunized

mice mounted a robust systemic HSV-specific CD8 T cell response, with gB-specific T cells

making up an average of 36.4% of CD8 T cells in the blood by 1wk after immunization (**Fig. 3C**).

At 1mo after immunization, we could identify these cells in multiple body sites including the CVT

and vaginal-draining lymph nodes (**Fig. S1B** and **Fig. 3D**). Again, CD103⁺ gB-specific CD8 T

cells in the CVT lacked expression of TCF-1, replicating the phenotype observed in both human

vaginal tissue and CVT of vaginally-immunized mice (**Fig. 3E**), despite the lack of tissue

inflammation.

Given this evidence that memory CD8 T cells in CVT lack significant expression of TCF-

1 regardless of immunization route, we next asked whether this memory compartment was

deficient in self-renewal and would undergo gradual decay. While the frequency of memory gB-

specific CD8 T cells remained relatively stable in the blood following contraction (**Fig. 3F**), and

the number of gB-specific CD8 T cells in the spleen, CVT-draining lymph nodes, and small

intestine lamina propria (SI LP) beyond 1mo after immunization was stable, this population

underwent a threefold loss in number within the CVT, with some mice having very few gB-

specific CD8 T cells remaining in the CVT by 5 months post-immunization (**Fig. 3G**).

Interestingly, gB-specific T cells in the SI LP also lacked expression of TCF-1 (**Fig. S2A**),

suggesting that memory T cells induced by the same immunization may have different

requirements for self-renewal.

Loss of memory CD8 T cells in the CVT occurs mainly among the CD69$^-$ CD103$^-$ subset

Given that gB-specific CD8 T cells in the CVT continued to decline in number after T cell

contraction had concluded in other tissues, we next examined the expression of canonical

markers of tissue residency and effector function. To reliably determine what proportion of the

gB-specific CD8 T cells in each organ were located within the tissue, we performed

intravascular (i.v.) antibody labeling (Galkina et al., 2005) (**Fig. S1C**). We found that an average

of 85.6% of the gB-specific CD8 T cells in the CVT were protected from i.v. labeling by 1mo

after immunization, similar to the fraction within the small intestine lamina propria (**Fig. 4A**). The

proportion of CVT gB-specific CD8 T cells that were protected from i.v. labeling remained stable

across time points, suggesting that the smaller population of T cells remaining at late time points

continued to be composed of cells that were mostly located within the tissue parenchyma (**Fig.

4A**).

We next assessed whether gB-specific CD8 T cells expressed CD69 and CD103. We

found that these markers were mostly non-expressed on gB-specific CD8 T cells at the earliest

memory time point. However, the average frequency of CD103$^+$ cells within the gB-specific CD8

T cell subset increased from 20.6% to 54.4% between 1mo and 5mo after LM-gB immunization

(**Fig. 4B-D**). While the i.v.-label$^+$ fraction of gB-specific CD8 T cells in the CVT remained almost

completely CD103$^-$ for the entire duration of the experiment, the shift from a predominantly

CD103$^-$ population to a predominantly CD103$^+$ population was especially clear within the i.v.-

label$^-$ fraction of CVT gB-specific CD8 T cells (**Fig. 4C-D**) and was mainly attributable to a

relative loss of CD103$^-$ cells from the CVT over time (**Fig. 4E**). Conversely, the equivalent i.v.

label$^-$ fraction of gB-specific CD8 T cells in the SI LP, did not undergo a population-level shift

towards becoming CD103$^+$ (**Fig. 4C**). Finally, the total CD69$^+$ CD103$^+$ population in the CVT

was stably maintained over time (**Fig. 4E**), which could be due to conversion of other subsets to

a CD69$^+$ CD103$^+$ phenotype or the ability to self-renew at a rate sufficient for stable

maintenance.

CVT memory CD8 T cells induced by systemic immunization gradually acquire expression of

granzyme B

We next wanted to determine if systemic immunization resulted in constitutive granzyme

B expression by CD8 T cells in the CVT, as we observed after vaginal immunization (**Fig. 2**) and

as described for tissue-resident CD8 T cells in the SI LP, which express granzyme B in the

absence of antigen re-exposure (Casey et al., 2012; Masopust et al., 2006). Few gB-specific

CD8 T cells in the CVT expressed granzyme B at 1mo after LM-gB immunization, but the

granzyme B$^+$ population substantially increased in both number and frequency at later time

points (**Fig. 5A-B**). In contrast, a subset of SI LP gB-specific CD8 T cells expressed granzyme B

by 1mo after immunization, and expression remained stable thereafter (**Fig. 5A**). As expected,

granzyme B expression was rarely observed among gB-specific CD8 T cells in the spleen or

dLN, regardless of the memory timepoint (**Fig. 5A**).

We next examined the phenotypic identity of these granzyme B-expressing cells. By

3mo post-immunization, granzyme B$^+$ gB-specific CD8 T cells in the CVT were almost

exclusively CD103$^+$ (**Fig. 5C-E**), resembling our results in human samples (**Fig. 1D**) and

suggesting that these two dynamically-expressed markers were demarcating the same cellular

population. By contrast, the granzyme B$^+$ population in the SI LP comprised both CD103$^+$ and

CD103$^-$ cells (**Fig. 5D-E**). These data suggest that immunization-induced gB-specific memory

CD8 T cells differentiate asynchronously in the CVT relative to the gut mucosa or peripheral

lymphoid organs.

Numeric and phenotypic changes of HSV-specific memory T cells in the CVT lead to delayed

protection against viral challenge.

Given the decline in the number of gB-specific CD8 T cells in the CVT combined with

their shift towards CD103 and granzyme B expression, we next evaluated how these changes

affected the immunoprotective response to vaginal HSV-2 challenge. Because LM-gB does not

prime HSV-specific CD4 T or B cells responses, this immunization system allowed us to

specifically assess the protective effect of CD8 T cells in the absence of other antigen-specific

responses. We presumed that the most rapid antiviral responses would likely be mediated by

tissue-resident gB-specific CD8 T cells, while circulating gB-specific CD8 T cells would

contribute to later responses. We challenged immunized mice at either 1mo ("Early Memory") or

4mo ("Late Memory") after immunization (**Fig. 6A**). A similar proportion of mice in each group

survived lethal challenge (**Fig. 6B**), demonstrating that gB-specific memory CD8 T cells were

sufficient to confer protection against HSV-2 lethality in a subset of immunized mice. However,

mice challenged earlier after immunization began to show evidence of faster viral clearance

starting as early as 3.5d after challenge (**Fig. 6C**) and underwent greater T cell expansion in the

first 2.5d after challenge (**Fig. 6D**). We hypothesized that this was due to quantitative and

qualitative differences between the early and late CD8 T cell compartment in the CVT.

To better understand the relationship between CVT gB-specific CD8 T cells and viral

control, we built upon our previously described approaches to model the acute T cell response

to HSV-2 infection (Schiffer et al., 2010; Schiffer et al., 2013) and used the mathematical model

in **Equation 1** to characterize the different gB-specific T cell response kinetics between the

early and late memory groups. In this model, susceptible cells in the CVT are infected by free

HSV-2, allowing them to produce new virus or be killed by HSV-specific CD8 T cells (**Fig. 6E**).

We performed 100 model-fit rounds to all observations simultaneously for each group for

experimentally-determined CVT gB-specific CD8 T cell counts and viral load and overlaid these

predictions onto our experimental data (**Fig. 6D)**. In naive mice, gB-specific CD8 T cells were

absent from or rare in the CVT at 0-3.5d after challenge but were abundant by 6.5d after

challenge (**Fig. 6D**, top left), corresponding with the entry of effector T cells into the CVT

following primary infection. In mice challenged at either early or late time points, gB-specific

CD8 T cells increased dramatically in number between 0d and 2.5d after HSV-2 infection, and

HSV viral titer fell by 6.5 post-challenge (**Fig. 6D**). The best model fits predicted that naive mice

had the highest basic reproductive number, meaning that they had the highest number of new

cellular infections predicted to derive from a single HSV-infected cell in a susceptible pool (**Fig.**

6F). In addition, the greater areas under fitted gB tet$^+$ T cell curves (shown in **Fig. 6D**, top row)

were associated with earlier viral clearance (**Fig. 6G**). These data suggest that the numeric and

phenotypic changes we observed within the CVT memory T cell compartment led to a decrease

in the efficacy of the anti-viral response in the first days after challenge.

Progressive differentiation by the memory CD8 T cell compartment in the CVT does not require

new input of cells from circulation

To determine if the CVT-specific changes in the memory T cell compartment occurred in

a tissue-autonomous manner, we depleted Thy1.1$^+$ gBT-I cells from the blood and lymphoid

tissues of LM-gB immunized mice 1mo after immunization (**Fig. 7A)**, as previously described

([Schenkel et al., 2013](#)). This method depleted splenic and circulating gBT-I cells but left the CVT

gBT-I compartment intact (**Fig. 7B-C**) and did not have a major impact on protection against

vaginal challenge with HSV-2 (**Fig. S3A-B**). By 3mo after depletion, we observed that a similar

level of decay had occurred among the CVT gB-specific CD8 T cell population to that seen in

undepleted mice between 1mo and 5mo after immunization (**Figs. 3G and 7C**). Similar to our

previous results, we observed that this decay occurred mostly among the CD103$^-$ subset of T

cells (**Fig. 7D-E**) and that the persisting cells had upregulated granzyme B by 3mo after

immunization (**Fig. 7F**). Low TCF-1 expression remained stable from 1mo to 3mo after

immunization but was reduced by gBT-I depletion, suggesting that the small TCF-1$^+$ subset in

the CVT of immunized mice was partially but not completely attributable to recirculating gBT-I

cells (**Fig. 7G**). These data indicate that the granzyme B$^+$ subset of gB-specific cells stems from

granzyme B$^-$ cells that were present in the tissue 1mo after immunization.

The memory CD8 T cell compartment in the CVT undergoes numeric decay regardless of

priming infection and immunization route

289   Finally, we wished to determine whether site-matched immunization would result in

improved maintenance of the CVT memory CD8 T cell compartment. We compared the

maintenance of HSV-specific CD8 T cells in the CVT of mice immunized vaginally with HSV-2

TK- or intravenously with LM-gB (**Fig. 8A**). We observed that vaginal immunization with HSV-2

TK- did not result in a larger memory population of HSV-specific CD8 T cells in the CVT than

systemic immunization with LM-gB (**Fig. 8B**). In addition, the gB-specific population induced by

vaginal immunization was relatively stably maintained in the spleen and SI LP, but underwent

decay in the CVT between 1mo and 3mo after immunization (**Fig. 8B**). Despite this decay in the

CD8 T cell compartment, HSV-2 TK- immunized mice remained completely protected against

lethal vaginal challenge with wildtype HSV-2 (**Fig. S3**), presumably due to CD4 T cells providing

substantial protection in this model ([Milligan et al., 1998](#)).

Discussion

Here, we report that CVT microenvironment plays a dynamic role in shaping the

numeric, phenotypic, and functional characteristics of the tissue memory CD8 T cells, which

thereby impacts their immunoprotective potential. The cervicovaginal mucosa is distinct from

other parts of the female reproductive tract and other mucosal tissues in terms of epithelium

type and physiology. Specifically, the vagina and ectocervix are composed of type II mucosa

and the endocervix and uterus are composed of type I mucosa, with the cervical transformation

zone marking the area where the simple columnar epithelial cells of the endocervix meet the

stratified squamous epithelial cells of the ectocervix and vagina. In contrast, the gut and the

lung, other mucosal tissues commonly studied in the case of CD8 T cell memory, are type I

mucosal surfaces (Iwasaki, 2010). In addition to these differences in the structure of the

epithelium, there are differences in the complexity and makeup of the microbiota at different

mucosal tissue sites (Anahtar et al., 2018), which may also contribute to tissue-specific

environmental differences that could affect memory CD8 T cell differentiation and function.

Vaginal-resident CD8 T cells in healthy women did not express TCF-1 (**Fig. 1**), which could

either simply indicate recent activation of these T cells (recent infection) or a memory population

with an impaired ability to self-renew. Importantly, given that vaginal CD8 T cells from multiple

donors uniformly lacked TCF-1 expression, it suggested that a tissue-driven phenotype was

more likely than a recent infection in a cohort of healthy donors with no current genital

infections. Using a mouse model, we similarly found that both vaginal (**Fig. 2**) and systemic

(**Fig. 3**) immunization resulted in a TCF-1^{low} phenotype among CD8 T cells in the mouse CVT.

These data demonstrate that the mouse and human memory CD8 T cell phenotype are

remarkably well conserved in the CVT.

Over the course of 5 months following immunization, antigen-specific memory CD8 T

cells in the CVT numerically declined and gradually acquired expression of CD103 and

granzyme B. Conversely, HSV-specific CD8 T cells that were circulating (spleen, LN) or resident

in the small intestine were numerically and phenotypically stable over the same time course.

This decline in CVT T_{RM} memory occurred at a similar rate even when the circulating gBT-I T

cells were selectively depleted, suggesting that peripheral T cells do not reseed the CVT under

homeostatic conditions (**Fig. 7**). In addition, the CVT CD8 T cell compartment upregulated

granzyme B in the absence of new input from the periphery, supporting the occurrence of

tissue-intrinsic memory differentiation. Thus, our data show that memory CD8 T cell

maintenance and differentiation in the CVT is ultimately directly and uniquely regulated by the

tissue environment. A possible explanation for these differences between the vagina and other

mucosal tissues such as the gut is that the CVT tissue microenvironment is lacking signals that

are important for resident-memory formation and maintenance (Liu et al., 2018). It is noteworthy

that we did not observe strong TCF-1 expression in the gut T_{RM} cells either, but these cells were

stably maintained over time. This could either suggest that TCF-1 is not required for self-

renewal of T_{RM}, or the existence of other compensatory mechanisms to maintain the T_{RM}

compartment in certain mucosal tissues. TCF-1 activity is thought to be regulated by β-catenin

and the Wnt signaling pathway. However, β-catenin does not regulate the phenotype of memory

T cells (Driessens et al., 2010) and β-catenin-deficient T cells mount normal recall responses

(Prlic and Bevan, 2011). Since β-catenin is sufficient, but not necessary to activate TCF-1

(Tiemessen et al., 2014), a β-catenin-independent pathway may control TCF-1 activity in tertiary

tissues with distinct cell fate outcomes compared to β-catenin-dependent activation. A recent

study provides some genetic evidence that TCF-1 negatively regulates CD103 expression in the

lung tissue (Wu et al., 2020), indicating the need to further study and dissect the role of TCF-1

in controlling differentiation and maintenance of T_{RM} cells.

Understanding the relationship between T_{RM} in mucosal tissues and the periphery is

highly relevant, because memory T cell characteristics and frequencies in the blood are used as

a benchmark in clinical studies to assess immune responses to vaccines and to establish

correlates of protection (Koup et al., 2011; Lewinsohn et al., 2017; Stephenson et al., 2016;

Streeck, 2016). Our data indicate that there is a disconnect between memory CD8 T cell

frequency, phenotype, and stability in the periphery and the CVT. The numerical and

phenotypical changes of the memory CD8 T cell compartment one month vs. 5 months post-

immunization ultimately influenced the rapidity of the immune response to vaginal viral

challenge (**Fig. 6**). This may be of particular relevance to the success of a vaccine-induced

memory T cell response within the CVT, as the host-pathogen events occurring within this early

window likely result in the ultimate success or failure of the protective response (Haase, 2010).

Barrier immunity in the CVT remains poorly understood, but is of particular interest,

because sexually-transmitted infections (STI) have a significant impact on global health. We

currently lack effective vaccines to prevent the majority of these infections, including HIV, HSV,

and bacterial STIs. Our data demonstrate that following systemic immunization, animals were

partially protected against HSV-2 lethality by gB-specific memory CD8 T cells (**Fig. 6**).

Importantly, this protective effect was observed despite the lack of HSV-specific CD4 T or B cell

memory, elements of the immune response that are often considered crucial for protection

against HSV-2 (Dudley et al., 2000; Milligan et al., 1998; Morrison et al., 1998; Petro et al.,

2016). In line with this observation, results from participants with recurrent genital HSV-2

infection indeed suggest that CD8 T cell activity within lesions is associated with viral control

(Peng et al., 2012; Schiffer et al., 2010; Zhu et al., 2007; Zhu et al., 2013). It is possible that a

new focus on vaccine strategies that elicit memory CD8 T cells within the genital mucosa, the

site of first exposure to STIs, will increase vaccine efficacy. However, successful design of these

approaches, not just in regards to adjuvant and immunization route, but also boosting intervals

to optimize protection, relies on understanding how memory CD8 T cells are maintained within

these genital tract mucosal tissues.

In summary, we demonstrate that the CVT tissue environment controls memory CD8 T

cell differentiation and maintenance, which occurs in a tissue autonomous manner and differs

across mucosal tissue compartments. The CD8 T_{RM} compartment in the CVT declines over time

with a concomitant decrease in the ability to mediate rapid antigen-specific protection.

Methods

Mice

C57BL/6J and B6.PL-*Thy1ᵃ*/CyJ mice were purchased from The Jackson Laboratory and maintained in specific pathogen-free conditions at the Fred Hutchinson Cancer Research Center (FHCRC). All mice used in experiments were female and were 8–12 weeks of age. Mice that rejected transferred Thy1.1⁺ gBT-I cells or had an undetectable HSV-2 titer in vaginal wash at 2.5d post-challenge were excluded from experiments. Experiments were approved by the FHCRC Institutional Animal Care and Use Committee.

Study population

Women recruited for this study (n=13) consented to vaginal biopsy and blood sampling. Eligibility criteria included aged > 18 to < 45, assigned female sex at birth, in good general health, normal PAP smear within the past 3-5 years, not menopausal, Hepatitis C negative and no report of active genital tract irritation or infection. Additionally, participants were screened to be negative for HIV, HSV-2, bacterial vaginosis, chlamydia, gonorrhea, and trichomonas vaginalis. Subjects were not using any type of steroid or medication for treatment of autoimmune disorders. Samples were not collected during menses. Informed written consent was obtained from all participants. The study and procedures were approved by the FHCRC Institutional Review Board.

Adoptive CD8 T cell transfers

CD44^{low} CD8 T cells were purified via negative selection from spleen and lymph nodes using mouse CD8a⁺ T Cell Isolation kits (Miltenyi) as previously described (Prlic et al., 2001). 5 × 10⁴ cells were injected into recipient B6 mice via i.v. injection into the tail vein.

Herpes simplex virus type 2 infections and quantification

Mice were injected subcutaneously with 2 mg of medroxyprogesterone acetate injectable

suspension (Greenstone). Seven days later, mice were infected intravaginally with 2×10^4 pfu

wildtype HSV-2 (strain 186 syn+) or 1.88×10^5 PFU of thymidine kinase-negative HSV-2 186

kpn to the vaginal canal in a 10 ul volume (Parr et al., 1994). After infection with wildtype HSV-2,

mice were monitored daily and clinical disease progression was scored as follows: 0, No sign; 1,

slight genital erythema; 2, moderate genital erythema and edema; 3, significant genital

inflammation with visible lesion; 4, hind leg paralysis or other severe condition requiring

euthanasia; 5, moribund or dead. Mice were euthanized if they were moribund or showed signs

of severe disease, including hind leg weakness, hind leg paralysis, or hunched posture. Viral

titer was determined by plaque assay on Vero cells (ATCC) or by PCR as previously described

(Jerome et al., 2002).

Listeria monocytogenes immunization

A strain of *Listeria monocytogenes* (LM) was generated to recombinantly express and

secrete ovalbumin containing the HSV-2 glycoprotein B-derived peptide SSIEFARL according to

previously described methods (Lauer et al., 2002; Zehn et al., 2009). Stocks were diluted to $2 \times$

10^4 CFU per ml in sterile PBS before immunization. Naive B6 mice received 4,000 CFU i.v. via

tail vein injection. Infectious dose was confirmed by plating on brain-heart infusion media plates

and enumerating colonies.

Labeling of T cells in the vasculature

To discriminate CD8 T cells in the vasculature from those located within tissues, we

utilized an intravascular labeling technique as previously described (Galkina et al., 2005).

Briefly, mice were injected i.v. with 3 ug anti-CD8b antibody conjugated to either FITC or APC

(Clone: H35-17.2, eBioscience) in PBS via the tail vein 2-5 min before euthanasia by CO_2

inhalation.

Depletion of Thy1.1-expressing T cells

Mice were injected intraperitoneally with 500 ng anti-Thy1.1 antibody (Clone 19E12, Bio

X Cell) to deplete Thy1.1-expressing cells from the circulation and peripheral lymphoid tissues.

Tissue collection and processing

Mouse

For pathologic assessment, murine cervicovaginal tracts (consisting of the vagina,

ectocervix, and endocervix) were fixed in 10% formalin, embedded in paraffin, sectioned, and

stained with hematoxylin and eosin.

To prepare single cell suspensions, cervicovaginal tracts were minced with scissors and

then incubated in 2 ml DMEM medium supplemented with 2% FBS, 2 mg/ml collagenase D

(Roche), and 1.5 mg/ml DNase I (Roche) for 30 minutes at 37C with gentle rocking. Single cell

suspensions were prepared from small intestines by removing the Peyer's patches and

mesentery, incubated in HBSS with 2% FBS, 5 mM EDTA, and 1 mM dithiothreitol for 15 min at

37C with gentle rocking, followed by collagenase treatment as outlined for the CVT.

Human

Blood was collected into acid citrate dextrose tubes. Tissues were transported from the clinic in

ice cold PBS and immediately processed upon arrival. Biopsy samples were trimmed to $2mm^2$

pieces and digested with collagenase II (700 units/ml, Sigma-Aldrich) and DNAse I (1unit/ml,

Sigma-Aldrich) for two subsequent 30 minute-digestions at 37°C as previously described

(McKinnon et al., 2014). Single cell suspensions were then washed and filtered before use.

Pathologic assessment of mouse vaginal tissue

Hematoxylin and eosin-stained sections of mouse CVT were scored based on signs of

inflammation within the vaginal lamina propria as follows: 1, Rare neutrophils and other

leukocytes present in the lamina propria; 2, Small clusters of neutrophils and other leukocytes

present in the lamina propria; 3, Multifocal clusters of leukocytes beginning to coalesce and/or

form follicles within the lamina propria; 4, Leukocytes arranged in sheets within the lamina

propria. Scoring was performed by a veterinary pathologist who was blinded to group identities.

Cell Staining for Flow Cytometry and Sorting

Cells were incubated in LIVE/DEAD fixable amine-reactive viability dye (Invitrogen),

blocked for Fc binding (Clone 2.4G2 for mice or TruStain (BioLegend) for human), and then

stained with tetramers and antibodies.

The following fluorochrome-conjugated antibodies were used to stain mouse cells for

flow cytometry: anti-CD8 (Clone 53-6.7, PerCP-eFluor710), anti-CD4 (Clone GK1.5, APC-

eFluor780), anti-CD45.1 (Clone A20, Brilliant Ultraviolet 395), anti-CD45.2 (Clone 104, Brilliant

Ultraviolet 737), anti-Thy1.1 (Clone HIS51, FITC or SuperBright645), anti-CD44 (Clone IM7,

APC or AlexaFluor700), anti-CD69 (Clone H1.2F3, PE-Dazzle594 or PE-Cy7), anti-CD103

(Clone M290, Brilliant Violet 786), anti-granzyme B (Clone GB11, Pacific Blue), and anti-TCF-1

(Clone C63D9, PE). Anti-CD8, anti-CD4, and anti-Thy1.1 were purchased from eBioscience.

Anti-CD45.1, anti-CD45.2, anti-CD44, and anti-CD103 were purchased from BD. Anti-CD69 and

anti-granzyme B antibodies were purchased from Biolegend. Anti-TCF-1 antibody was

purchased from Cell Signaling Technologies.

The following fluorochrome-conjugated antibodies were used to stain human cells for

flow cytometry: anti-CD3 (Clone UCHT1, Brilliant Ultraviolet 661), anti-CD45 (Clone HI30,

Brilliant Ultraviolet 805), anti-CD8 (Clone RPA-T8, Brilliant Ultraviolet 496), anti-CD4 (Clone

RPA-T4, Brilliant Violet 605), anti-CD45RA (Clone HI100, APC-H7), anti-CCR7 (Clone G043H7,

AlexaFluor488), anti-granzyme B (Clone GB11, AlexaFluor700), anti-CD103 (Clone Ber-ACT8,

Brilliant Violet 750), and anti-TCF-1 (Clone C63D9, PE). Anti-CD3, anti-CD45, anti-CD4, anti-

CD8, anti-CD45RA, anti-CD103, and anti-granzyme B were purchased from BD. Anti-CCR7

antibody was purchased from Biolegend. Anti-TCF-1 antibody was purchased from Cell

Signaling Technologies.

Intracellular staining was conducted after fixation and membrane permeabilization using

Cytofix/Cytoperm Fixation/Permeabilization Kits (BD). If TCF-1 was included in the staining

panel, intracellular staining was conducted after nuclear permeabilization using

Foxp3/Transcription Factor Staining Kits (eBioscience) according to the manufacturer's protocol.

Samples were acquired on a FACSymphony instrument (BD) or sorted on a FACSAria (BD). To

estimate cell counts, 2×10^5 AccuCheck Counting Beads (ThermoFisher) were added to each

sample immediately before acquisition.

Statistical Analysis

Sample means between two unpaired groups were compared by *t*-test. Sample means

between three or more unpaired groups were compared by ordinary one-way ANOVA with

Tukey's multiple comparisons test. Sample means between three paired groups were compared

by repeated measures ANOVA with Greenhouse-Geisser correction. Paired mean fluorescence

intensity or relative frequency data were analyzed via paired *t*-test. All cell count and viral titer

data were $\log_{10}$-transformed before graphing and analysis. Error bars represent standard

deviations. Exact *P*-values are given for all results where $P < 0.05$. Analysis was performed

using Prism 7 (GraphPad Software).

HSV-2 and CD8 T cell Dynamics Modeling

To understand the virus and gB-specific CD8 T cell dynamic differences between the

Naive, Early Memory, and Late Memory groups, we used an acute viral infection model that

included a cytotoxic effect of the CD8 T cell compartment. In this model, susceptible cells (*S*)

are infected at rate *βVS* by free HSV-2 virus (*V*). Productively HSV-2-infected cells (*I*) have a

clearance that is intrinsic or mediated by a humoral response with rate *δ$_I$I* or a clearance

mediated by HSV-specific CD8 T cells (*E*) with killing rate *kEI*. Free virus is produced at a rate

pI and cleared at rate *cV*. HSV-specific CD8 T cells are cleared at rate *δ$_E$E* and proliferate in the

presence of infected cells at a maximum rate *ωEI*. We also assumed the proliferation of HSV-

specific CD8 T cells is constrained when infected and effector cells grow over saturation levels

I$_{50}$ and *E$_{50}$*, respectively. Under these assumptions the model has the form (see schematics in

Fig. 6D):

519
$$\frac{dS}{dt} = -\beta VS$$
$$\frac{dI}{dt} = \beta VS - \delta_I I - kEI$$
$$\frac{dV}{dt} = pI - cV \qquad \textbf{(Equation 1)}$$
$$\frac{dE}{dt} = \frac{\omega EI}{\left(1+\frac{I}{I_{50}}\right)\left(1+\frac{E}{E_{50}}\right)} - \delta_E E.$$

The basic reproductive ratio, R_0, which represents the number of secondary infections produced

by an HSV-infected cell when introduced into a susceptible population, was calculated as $R_0 =$

$\frac{\beta p T_0}{c(\delta_I + kE_0)}$. When $R_0 < 1$, viremia is controlled.

We performed 100 rounds of model fits to observations from all individual animals for

each group with different initial parameter guesses in the optimization algorithm. We assumed

that all outcomes from the 100 fit rounds were plausible predictions of the data and represented

the variability in the infection dynamics in each group. For each round, we re-parameterized the

model in function of R_0, and used a nonlinear least-squares approach using the differential

evolution and the L-BFGS-B algorithms in R (R Development Core Team) to fit the model and

to estimate parameters: V_0, pT_0, E_0, R_0, δ, c, k, ω, δ_E, I_{50} and E_{50}. Since mice were inoculated

with ~2.3×10^6 HSV genomes, we set that number as the upper limit for V_0. We also constrained

the value estimate E_0 to the maximum and minimum HSV-specific CD8 T cell observations at

the moment of virus challenge. Finally, we ensured that $R_0 > 1$ in all fitting rounds. In all

simulations $t = 0$ represented the time of the HSV-2 vaginal challenge. From the best fits, we

computed the predicted time of viral clearance as the time when viral load crossed the detection

limit for the PCR assay of 4615.4 genomes per vaginal wash. We also computed the predicted

area under the curve of HSV-specific CD8 T cells using best fits of variable E from challenge

until the time of viral clearance.

We repeated the fits by exploring different possibilities for constraining HSV-specific

CD8 T cell growth. Specifically, we explored a model were CD8 growth is constrained only by a

saturation level in HSV-infected cells, i.e. $\frac{dE}{dt} = \frac{\omega EI}{\left(1+\frac{I}{I_{50}}\right)} - \delta_E E$, or only by a saturation for the

number of effector cells, i.e. $\frac{dE}{dt} = \frac{\omega EI}{\left(1+\frac{E}{E_{50}}\right)} - \delta_E E$. To determine the best and most parsimonious

model, we computed the Akaike Information Criteria, $AIC = n \ln \overline{SSE} + 2m$, where m is the

number of parameters estimated, n the number of data points in each group and $\overline{SSE}$ is the

average sum of squares error of all the 100 model-fit rounds in each group. We assumed a

model had similar support from the data if the difference between its AIC and the best model

(lowest) AIC was less than two. We selected and showed results from the model with the lowest

AIC.

Abbreviations

AIC, Akaike information criteria

AUC, area under curve

CVT, cervicovaginal tissue

dLN, CVT-draining lymph nodes

gB, HSV glycoprotein B

HSV-2 TK-, thymidine kinase-deficient herpes simplex virus type 2

LM-gB, *Listeria monocytogenes* strain expressing the SSIEFARL peptide from HSV glycoprotein

B

LN, lymph node

SI, small intestine

SI LP, small intestine lamina propria

STI, sexually-transmitted infection

Acknowledgements

We thank Kristin Weakly for excellent technical assistance and Sean M. Hughes for assistance

with statistical analyses. This work was supported by NIH grants R01 AI123323 (MP), R01

AI121129 to (JTS, JML and MP), R01 AI131914 (JML), R01 AI141435 (JML), T32 AI07140 and

the Doug and Maggie Walker Fellowship (ASWD), and T32 AI007509 and F31 AI140514 (VD).

FM is an ISAC scholar.

References

Anahtar, M.N., D.B. Gootenberg, C.M. Mitchell, and D.S. Kwon. 2018. Cervicovaginal

Microbiota and Reproductive Health: The Virtue of Simplicity. *Cell Host Microbe* 23:159-

168.

Ariotti, S., M.A. Hogenbirk, F.E. Dijkgraaf, L.L. Visser, M.E. Hoekstra, J.Y. Song, H. Jacobs, J.B.

Haanen, and T.N. Schumacher. 2014. T cell memory. Skin-resident memory CD8(+) T

cells trigger a state of tissue-wide pathogen alert. *Science* 346:101-105.

Bergsbaken, T., M.J. Bevan, and P.J. Fink. 2017. Local Inflammatory Cues Regulate

Differentiation and Persistence of CD8(+) Tissue-Resident Memory T Cells. *Cell Rep*

19:114-124.

Beura, L.K., N.J. Fares-Frederickson, E.M. Steinert, M.C. Scott, E.A. Thompson, K.A. Fraser,

J.M. Schenkel, V. Vezys, and D. Masopust. 2019. CD4(+) resident memory T cells

dominate immunosurveillance and orchestrate local recall responses. *J Exp Med*

216:1214-1229.

Beura, L.K., J.S. Mitchell, E.A. Thompson, J.M. Schenkel, J. Mohammed, S. Wijeyesinghe, R.

Fonseca, B.J. Burbach, H.D. Hickman, V. Vezys, B.T. Fife, and D. Masopust. 2018.

Intravital mucosal imaging of CD8(+) resident memory T cells shows tissue-autonomous

recall responses that amplify secondary memory. *Nature immunology* 19:173-182.

Casey, K.A., K.A. Fraser, J.M. Schenkel, A. Moran, M.C. Abt, L.K. Beura, P.J. Lucas, D. Artis,

E.J. Wherry, K. Hogquist, V. Vezys, and D. Masopust. 2012. Antigen-independent

differentiation and maintenance of effector-like resident memory T cells in tissues.

Journal of immunology (Baltimore, Md. : 1950) 188:4866-4875.

Davies, B., J.E. Prier, C.M. Jones, T. Gebhardt, F.R. Carbone, and L.K. Mackay. 2017. Cutting

Edge: Tissue-Resident Memory T Cells Generated by Multiple Immunizations or

Localized Deposition Provide Enhanced Immunity. *Journal of immunology (Baltimore,*

Md. : 1950) 198:2233-2237.

Driessens, G., Y. Zheng, and T.F. Gajewski. 2010. Beta-catenin does not regulate memory T
cell phenotype. *Nat Med* 16:513-514; author reply 514-515.

Dudley, K.L., N. Bourne, and G.N. Milligan. 2000. Immune protection against HSV-2 in B-cell-
deficient mice. *Virology* 270:454-463.

Galkina, E., J. Thatte, V. Dabak, M.B. Williams, K. Ley, and T.J. Braciale. 2005. Preferential
migration of effector CD8+ T cells into the interstitium of the normal lung. *J Clin Invest*
115:3473-3483.

Gebhardt, T., L.M. Wakim, L. Eidsmo, P.C. Reading, W.R. Heath, and F.R. Carbone. 2009.
Memory T cells in nonlymphoid tissue that provide enhanced local immunity during
infection with herpes simplex virus. *Nature immunology* 10:524-530.

Haase, A.T. 2010. Targeting early infection to prevent HIV-1 mucosal transmission. *Nature*
464:217-223.

Hofmann, M., and H. Pircher. 2011. E-cadherin promotes accumulation of a unique memory
CD8 T-cell population in murine salivary glands. *Proceedings of the National Academy*
of Sciences of the United States of America 108:16741-16746.

Iijima, N., and A. Iwasaki. 2014. T cell memory. A local macrophage chemokine network
sustains protective tissue-resident memory CD4 T cells. *Science* 346:93-98.

Iwasaki, A. 2010. Antiviral immune responses in the genital tract: clues for vaccines. *Nat Rev*
Immunol 10:699-711.

Jerome, K.R., M.L. Huang, A. Wald, S. Selke, and L. Corey. 2002. Quantitative stability of DNA
after extended storage of clinical specimens as determined by real-time PCR. *J Clin*
Microbiol 40:2609-2611.

Jiang, X., R.A. Clark, L. Liu, A.J. Wagers, R.C. Fuhlbrigge, and T.S. Kupper. 2012. Skin
infection generates non-migratory memory CD8+ T(RM) cells providing global skin
immunity. *Nature* 483:227-231.

Khan, T.N., J.L. Mooster, A.M. Kilgore, J.F. Osborn, and J.C. Nolz. 2016. Local antigen in

nonlymphoid tissue promotes resident memory CD8+ T cell formation during viral

infection. *J Exp Med* 213:951-966.

Kim, S.K., D.S. Reed, W.R. Heath, F. Carbone, and L. Lefrancois. 1997. Activation and

migration of CD8 T cells in the intestinal mucosa. *Journal of immunology (Baltimore, Md.*

: 1950) 159:4295-4306.

Koup, R.A., B.S. Graham, and D.C. Douek. 2011. The quest for a T cell-based immune

correlate of protection against HIV: a story of trials and errors. *Nat Rev Immunol* 11:65-

70.

Lauer, P., M.Y. Chow, M.J. Loessner, D.A. Portnoy, and R. Calendar. 2002. Construction,

characterization, and use of two Listeria monocytogenes site-specific phage integration

vectors. *Journal of bacteriology* 184:4177-4186.

Lewinsohn, D.A., D.M. Lewinsohn, and T.J. Scriba. 2017. Polyfunctional CD4(+) T Cells As

Targets for Tuberculosis Vaccination. *Front Immunol* 8:1262.

Liu, Y., C. Ma, and N. Zhang. 2018. Tissue-Specific Control of Tissue-Resident Memory T Cells.

Crit Rev Immunol 38:79-103.

Mackay, L.K., A. Rahimpour, J.Z. Ma, N. Collins, A.T. Stock, M.L. Hafon, J. Vega-Ramos, P.

Lauzurica, S.N. Mueller, T. Stefanovic, D.C. Tscharke, W.R. Heath, M. Inouye, F.R.

Carbone, and T. Gebhardt. 2013. The developmental pathway for CD103(+)CD8+

tissue-resident memory T cells of skin. *Nature immunology* 14:1294-1301.

Mackay, L.K., A.T. Stock, J.Z. Ma, C.M. Jones, S.J. Kent, S.N. Mueller, W.R. Heath, F.R.

Carbone, and T. Gebhardt. 2012. Long-lived epithelial immunity by tissue-resident

memory T (TRM) cells in the absence of persisting local antigen presentation.

Proceedings of the National Academy of Sciences of the United States of America

109:7037-7042.

Mackay, L.K., E. Wynne-Jones, D. Freestone, D.G. Pellicci, L.A. Mielke, D.M. Newman, A.

Braun, F. Masson, A. Kallies, G.T. Belz, and F.R. Carbone. 2015. T-box Transcription

Factors Combine with the Cytokines TGF-beta and IL-15 to Control Tissue-Resident

Memory T Cell Fate. *Immunity* 43:1101-1111.

Masopust, D., D. Choo, V. Vezys, E.J. Wherry, J. Duraiswamy, R. Akondy, J. Wang, K.A.

Casey, D.L. Barber, K.S. Kawamura, K.A. Fraser, R.J. Webby, V. Brinkmann, E.C.

Butcher, K.A. Newell, and R. Ahmed. 2010. Dynamic T cell migration program provides

resident memory within intestinal epithelium. *J Exp Med* 207:553-564.

Masopust, D., and A.G. Soerens. 2019. Tissue-Resident T Cells and Other Resident

Leukocytes. *Annu Rev Immunol* 37:521-546.

Masopust, D., V. Vezys, A.L. Marzo, and L. Lefrancois. 2001. Preferential localization of effector

memory cells in nonlymphoid tissue. *Science* 291:2413-2417.

Masopust, D., V. Vezys, E.J. Usherwood, L.S. Cauley, S. Olson, A.L. Marzo, R.L. Ward, D.L.

Woodland, and L. Lefrancois. 2004. Activated primary and memory CD8 T cells migrate

to nonlymphoid tissues regardless of site of activation or tissue of origin. *Journal of*

immunology (Baltimore, Md. : 1950) 172:4875-4882.

Masopust, D., V. Vezys, E.J. Wherry, D.L. Barber, and R. Ahmed. 2006. Cutting edge: gut

microenvironment promotes differentiation of a unique memory CD8 T cell population.

Journal of immunology (Baltimore, Md. : 1950) 176:2079-2083.

McKinnon, L.R., S.M. Hughes, S.C. De Rosa, J.A. Martinson, J. Plants, K.E. Brady, P.P. Gumbi,

D.J. Adams, L. Vojtech, C.G. Galloway, M. Fialkow, G. Lentz, D. Gao, Z. Shu, B.

Nyanga, P. Izulla, J. Kimani, S. Kimwaki, A. Bere, Z. Moodie, A.L. Landay, J.A.

Passmore, R. Kaul, R.M. Novak, M.J. McElrath, and F. Hladik. 2014. Optimizing viable

leukocyte sampling from the female genital tract for clinical trials: an international multi-

site study. *PLoS One* 9:e85675.

Milligan, G.N., D.I. Bernstein, and N. Bourne. 1998. T lymphocytes are required for protection of

the vaginal mucosae and sensory ganglia of immune mice against reinfection with

herpes simplex virus type 2. *Journal of immunology (Baltimore, Md. : 1950)* 160:6093-

6100.

Milner, J.J., C. Toma, B. Yu, K. Zhang, K. Omilusik, A.T. Phan, D. Wang, A.J. Getzler, T.

Nguyen, S. Crotty, W. Wang, M.E. Pipkin, and A.W. Goldrath. 2017. Runx3 programs

CD8(+) T cell residency in non-lymphoid tissues and tumours. *Nature* 552:253-257.

Morrison, L.A., X.J. Da Costa, and D.M. Knipe. 1998. Influence of mucosal and parenteral

immunization with a replication-defective mutant of HSV-2 on immune responses and

protection from genital challenge. *Virology* 243:178-187.

Park, S.L., A. Zaid, J.L. Hor, S.N. Christo, J.E. Prier, B. Davies, Y.O. Alexandre, J.L. Gregory,

T.A. Russell, T. Gebhardt, F.R. Carbone, D.C. Tscharke, W.R. Heath, S.N. Mueller, and

L.K. Mackay. 2018. Local proliferation maintains a stable pool of tissue-resident memory

T cells after antiviral recall responses. *Nature immunology* 19:183-191.

Parr, M.B., L. Kepple, M.R. McDermott, M.D. Drew, J.J. Bozzola, and E.L. Parr. 1994. A mouse

model for studies of mucosal immunity to vaginal infection by herpes simplex virus type

2. *Laboratory investigation; a journal of technical methods and pathology* 70:369-380.

Pattacini, L., A. Woodward Davis, J. Czartoski, F. Mair, S. Presnell, S.M. Hughes, O. Hyrien,

G.M. Lentz, A.C. Kirby, M.F. Fialkow, F. Hladik, M. Prlic, and J.M. Lund. 2019. A pro-

inflammatory CD8+ T-cell subset patrols the cervicovaginal tract. *Mucosal immunology*

Peng, T., J. Zhu, K. Phasouk, D.M. Koelle, A. Wald, and L. Corey. 2012. An effector phenotype

of CD8+ T cells at the junction epithelium during clinical quiescence of herpes simplex

virus 2 infection. *J Virol* 86:10587-10596.

Petro, C.D., B. Weinrick, N. Khajoueinejad, C. Burn, R. Sellers, W.R. Jacobs, Jr., and B.C.

Herold. 2016. HSV-2 DeltagD elicits FcgammaR-effector antibodies that protect against

clinical isolates. *JCI insight* 1:

Piet, B., G.J. de Bree, B.S. Smids-Dierdorp, C.M. van der Loos, E.B. Remmerswaal, J.H. von

der Thusen, J.M. van Haarst, J.P. Eerenberg, A. ten Brinke, W. van der Bij, W. Timens,

R.A. van Lier, and R.E. Jonkers. 2011. CD8(+) T cells with an intraepithelial phenotype

upregulate cytotoxic function upon influenza infection in human lung. *J Clin Invest*

121:2254-2263.

Pope, C., S.K. Kim, A. Marzo, D. Masopust, K. Williams, J. Jiang, H. Shen, and L. Lefrancois.

2001. Organ-specific regulation of the CD8 T cell response to Listeria monocytogenes

infection. *Journal of immunology (Baltimore, Md. : 1950)* 166:3402-3409.

Prlic, M., and M.J. Bevan. 2011. Cutting edge: beta-catenin is dispensable for T cell effector

differentiation, memory formation, and recall responses. *Journal of immunology*

(Baltimore, Md. : 1950) 187:1542-1546.

Prlic, M., B.R. Blazar, A. Khoruts, T. Zell, and S.C. Jameson. 2001. Homeostatic expansion

occurs independently of costimulatory signals. *Journal of immunology (Baltimore, Md. :*

1950) 167:5664-5668.

Reinhardt, R.L., A. Khoruts, R. Merica, T. Zell, and M.K. Jenkins. 2001. Visualizing the

generation of memory CD4 T cells in the whole body. *Nature* 410:101-105.

Schenkel, J.M., K.A. Fraser, V. Vezys, and D. Masopust. 2013. Sensing and alarm function of

resident memory CD8(+) T cells. *Nature immunology* 14:509-513.

Schiffer, J.T., L. Abu-Raddad, K.E. Mark, J. Zhu, S. Selke, D.M. Koelle, A. Wald, and L. Corey.

2010. Mucosal host immune response predicts the severity and duration of herpes

simplex virus-2 genital tract shedding episodes. *Proceedings of the National Academy of*

Sciences of the United States of America 107:18973-18978.

Schiffer, J.T., D. Swan, R. Al Sallaq, A. Magaret, C. Johnston, K.E. Mark, S. Selke, N.

Ocbamichael, S. Kuntz, J. Zhu, B. Robinson, M.L. Huang, K.R. Jerome, A. Wald, and L.

Corey. 2013. Rapid localized spread and immunologic containment define Herpes

simplex virus-2 reactivation in the human genital tract. *Elife* 2:e00288.

Shin, H., and A. Iwasaki. 2012. A vaccine strategy that protects against genital herpes by

establishing local memory T cells. *Nature* 491:463-467.

Steinbach, K., I. Vincenti, M. Kreutzfeldt, N. Page, A. Muschaweckh, I. Wagner, I. Drexler, D.

Pinschewer, T. Korn, and D. Merkler. 2016. Brain-resident memory T cells represent an

autonomous cytotoxic barrier to viral infection. *J Exp Med* 213:1571-1587.

Steinert, E.M., J.M. Schenkel, K.A. Fraser, L.K. Beura, L.S. Manlove, B.Z. Igyarto, P.J.

Southern, and D. Masopust. 2015. Quantifying Memory CD8 T Cells Reveals

Regionalization of Immunosurveillance. *Cell* 161:737-749.

Stephenson, K.E., H.T. D'Couto, and D.H. Barouch. 2016. New concepts in HIV-1 vaccine

development. *Curr Opin Immunol* 41:39-46.

Streeck, H. 2016. Designing optimal HIV-vaccine T-cell responses. *Curr Opin HIV AIDS* 11:593-

600.

Szabo, P.A., M. Miron, and D.L. Farber. 2019. Location, location, location: Tissue resident

memory T cells in mice and humans. *Sci Immunol* 4:

Takamura, S., H. Yagi, Y. Hakata, C. Motozono, S.R. McMaster, T. Masumoto, M. Fujisawa, T.

Chikaishi, J. Komeda, J. Itoh, M. Umemura, A. Kyusai, M. Tomura, T. Nakayama, D.L.

Woodland, J.E. Kohlmeier, and M. Miyazawa. 2016. Specific niches for lung-resident

memory CD8+ T cells at the site of tissue regeneration enable CD69-independent

maintenance. *J Exp Med* 213:3057-3073.

Tiemessen, M.M., M.R. Baert, L. Kok, M.C. van Eggermond, P.J. van den Elsen, R. Arens, and

F.J. Staal. 2014. T Cell factor 1 represses CD8+ effector T cell formation and function.

Journal of immunology (Baltimore, Md. : 1950) 193:5480-5487.

Wakim, L.M., J. Waithman, N. van Rooijen, W.R. Heath, and F.R. Carbone. 2008. Dendritic cell-

induced memory T cell activation in nonlymphoid tissues. *Science* 319:198-202.

Walsh, D.A., H. Borges da Silva, L.K. Beura, C. Peng, S.E. Hamilton, D. Masopust, and S.C.

Jameson. 2019. The Functional Requirement for CD69 in Establishment of Resident

Memory CD8(+) T Cells Varies with Tissue Location. *Journal of immunology (Baltimore,*

Md. : 1950) 203:946-955.

Watanabe, R., A. Gehad, C. Yang, L.L. Scott, J.E. Teague, C. Schlapbach, C.P. Elco, V.

Huang, T.R. Matos, T.S. Kupper, and R.A. Clark. 2015. Human skin is protected by four

functionally and phenotypically discrete populations of resident and recirculating memory

T cells. *Sci Transl Med* 7:279ra239.

Wu, J., A. Madi, A. Mieg, A. Hotz-Wagenblatt, N. Weisshaar, S. Ma, K. Mohr, T. Schlimbach, M.

Hering, H. Borgers, and G. Cui. 2020. T Cell Factor 1 Suppresses CD103+ Lung Tissue-

Resident Memory T Cell Development. *Cell Rep* 31:107484.

Zehn, D., S.Y. Lee, and M.J. Bevan. 2009. Complete but curtailed T-cell response to very low-

affinity antigen. *Nature* 458:211-214.

Zhang, N., and M.J. Bevan. 2013. Transforming growth factor-beta signaling controls the

formation and maintenance of gut-resident memory T cells by regulating migration and

retention. *Immunity* 39:687-696.

Zhou, X., S. Yu, D.M. Zhao, J.T. Harty, V.P. Badovinac, and H.H. Xue. 2010. Differentiation and

persistence of memory CD8(+) T cells depend on T cell factor 1. *Immunity* 33:229-240.

Zhu, J., D.M. Koelle, J. Cao, J. Vazquez, M.L. Huang, F. Hladik, A. Wald, and L. Corey. 2007.

Virus-specific CD8+ T cells accumulate near sensory nerve endings in genital skin

during subclinical HSV-2 reactivation. *J Exp Med* 204:595-603.

Zhu, J., T. Peng, C. Johnston, K. Phasouk, A.S. Kask, A. Klock, L. Jin, K. Diem, D.M. Koelle, A.

Wald, H. Robins, and L. Corey. 2013. Immune surveillance by CD8alphaalpha+ skin-

resident T cells in human herpes virus infection. *Nature* 497:494-497.

Figure Legends

Figure 1. Human cervicovaginal-resident CD8 T cells lack expression of TCF-1. (A) Gating

strategy to identify memory CD8 T cells in blood and CVT from healthy STI and HSV-2-negative

women. **(B)** Relative abundance of CD69 and CD103 subsets in blood and CVT. **(C)**

Representative flow plot and abundance of granzyme B$^+$ cells within the memory CD8

population in blood and CVT. **(D)** Representative flow plot and abundance of CD103 and

granzyme B co-expression in blood and CVT. **(E)** Representative TCF-1 staining and TCF-1$^+$

frequency from CD103$^+$ subset of CD8 T cells from the CVT compared to memory CD8 T cells

in matched blood. Flow plots from A and C-E are from one representative participant. Data from

B-E represent pooled results from study participants (n=13). Each dot in C-E represents an

individual participant. Error bars represent mean $\pm$ SD. *P*-value in E was calculated via paired *t*-

test. Exact *P*-values are given for all comparisons.

Figure 2. Mouse cervicovaginal-resident CD8 T cells induced by vaginal immunization

mirror the phenotype observed in human samples and lack TCF-1. (A) Experiment

schematic to induce CVT-resident CD8 T cells in mice via vaginal infection with 1.88×10^5 PFU

of thymidine kinase-negative HSV-2 186 kpn. **(B)** HSV-specific cells were identified in CVT via

staining with an H-2Kb SSIEFARL tetramer conjugated to APC. Left, representative flow plots

gated on CD8 T cells. Right, relative abundance of CD69 and CD103 subsets within gB

tetramer$^+$ CD8 T cells in CVT. **(C)** Representative flow plot and abundance of granzyme B$^+$ cells

within gB tetramer$^+$ CD8 T cells in CVT and CVT-draining lymph nodes (dLN). **(D)**

Representative TCF-1 staining and TCF-1$^+$ frequency within gB tetramer$^+$ CD8 T cells in CVT

and dLN. **(E)** Representative images of hematoxylin and eosin-stained sections of CVT at 10X

objective and average lamina propria inflammation scores. Arrow indicates coalescing clusters

of inflammatory cells with follicular organization. Arrowhead indicates clusters of inflammatory

cells within the mucosal epithelium. Flow plots and images from B-E are from one

representative mouse. Data from B-D represent pooled results from 2 independent experiments.

Each dot in E represents inflammation score averaged from 1-3 mice per time point. Error bars

represent mean $\pm$ SD. *P*-value in C was calculated via paired *t*-test. Exact *P*-values are given for

all comparisons.

Figure 3. The cervicovaginal CD8 T cell memory compartment of LM-gB-immunized mice

is poorly maintained after immunization. (A) Schematic of experiment to assess phenotype

and maintenance of HSV-specific T cell compartment of mice immunized with 4000 CFU LM-gB

i.v. **(B)** Representative image of hematoxylin and eosin-stained sections of CVT at 10X

objective and average lamina propria inflammation scores. **(C)** Representative flow plot gated

on CD8 T cells from blood collected 1wk after LM-gB immunization. **(D)** gB tetramer$^+$ CD8 T cell

abundance in spleen, dLN, and CVT 1mo after immunization. **(E)** Representative TCF-1 staining

and quantification of TCF-1 expression within CD103$^+$ and CD103$^-$ subsets of gB tetramer$^+$ CD8

T cells in CVT and dLN. **(F-G)** Relative abundance or number of HSV-specific CD8 T cells in

blood, spleen, dLN, small intestine lamina propria (SI LP), and CVT 1mo and 5mo after LM-gB

immunization. Images and flow plots from B, C, and E are from one representative mouse. Data

in D-G are pooled from 2-5 independent experiments. Each dot in B represents inflammation

score averaged from 2-3 mice per time point. Each dot in F represents the mean value of 20

mice from a representative experiment. Each dot in D, E, and G represents an individual mouse.

Error bars represent mean $\pm$ SD. *P*-value in B was calculated via *t*-test. *P*-values in E were

calculated with repeated measures ANOVA with Greenhouse-Geisser correction. *P*-values in G

were calculated via *t*-test using $\log_{10}$-transformed values. Exact *P*-values are given for all values

< 0.05.

Figure 4. Most HSV-specific CD8 T cells remaining in the CVT by five months after

immunization express CD69 and CD103. **(A)** Percentage of i.v. label⁻ HSV-specific CD8 T

cells in different lymphoid and non-lymphoid tissues at 1mo, 3mo, and 5mo after immunization

based on gating shown in Fig. S1C. **(B)** Representative staining showing CD69 and CD103

expression on HSV-specific CD8 T cells in the CVT 1mo and 5mo after immunization. **(C)**

Stacked bar plots showing relative abundance of each of four subsets of HSV-specific CD8 T

cells based on CD69 and CD103 expression in the i.v. label⁺ and i.v. label⁻ fractions of the CVT

and SI LP. **(D)** CD103 expression on i.v. label⁻ gB tetramer⁺ CD8 T cells in the CVT 1mo, 3mo,

and 5mo after immunization. **(E)** Total i.v. label⁻ count of each of four subsets of HSV-specific

CD8 T cells based on CD69 and CD103 expression 1mo, 3mo, and 5mo after immunization.

Data in A, C, D, and E are pooled from at least 2 independent experiments per time point. Data

in B are from one representative experiment. Each dot in A, D, and E represents an individual

mouse. Error bars represent mean $\pm$ SD. *P*-values in D and E were calculated via ordinary one-

way ANOVA with Tukey's post-hoc test using $\log_{10}$-transformed values for numbers and

untransformed values for percentages. Exact *P*-values are given for all values <0.05.

Figure 5. After systemic immunization, cervicovaginal-resident CD8 T cells progressively

upregulate granzyme B. **(A)** Frequency of granzyme B⁺ cells among HSV-specific CD8 T cells

in spleen, dLN, CVT, and SI LP at 1mo, 3mo, and 5mo post-immunization. **(B)** Number of

granzyme B-expressing cells among HSV-specific CD8 T cells at time points after LM-gB

immunization. **(C)** Concatenated flow files from 10 mice showing CD103 and granzyme B co-

expression among the HSV-specific CD8b i.v. label⁻ populations in the CVT and dLN 5mo after

LM-gB immunization. **(D-E)** Frequency of CD69 and CD103 expression within granzyme B⁺

HSV-specific CD8 T cells in the CVT and SI LP 1mo, 3mo, and 5mo after immunization. Data in

A-E are pooled from at least 2 independent experiments. Each dot or triangle in A, B, and D

represents an individual mouse. Error bars represent mean $\pm$ SD. *P*-values in A and B were

calculated via ordinary one-way ANOVA with Tukey's post-hoc test using log_{10}-transformed

values for numbers and untransformed values for percentages. Exact *P*-values are given for all

values <0.05.

Figure 6. LM-gB immunized mice are protected against severe disease after lethal

intravaginal challenge with HSV-2, but the efficacy of the immune response wanes over

time. (A) Experiment schematic to compare protective efficacy of LM-gB immunization 1mo and

4mo after administration. **(B)** Survival after vaginal HSV-2 challenge of naive and LM-gB

immunized mice. **(C)** Vaginal washes were collected after HSV-2 challenge and HSV-2 titer was

determined by plaque assay. Pink dots represent mice that survived challenge, as shown in B.

(D) Number of HSV-specific CD8 T cells (top) and HSV genomes (bottom) in the CVT with

modeling estimates overlaid in blue. **(E)** Schematic of mathematical model relating CVT-

infiltrating CD8 T cell immune response to HSV-2 viral expansion kinetics. *S* and *I* represent the

number of susceptible and infected cells per CVT, respectively; *E*, the number of gB tet$^+$ CD8 T

cells per CVT; and *V* the number of HSV genomes per vaginal wash. **(F)** Model estimates of

basic reproductive ratios (number of new cells infected by one infected cell when introduced into

a pool of susceptible cells) for each group of mice. **(G)** Scatter plot of areas under curves (AUC)

from D and corresponding predicted day of viral clearance. Dotted line represents a fitted line

from quadratic regression. Data in B, C, and D are pooled from at least 2 independent

experiments (n=8-14 per group in B). Each dot, square, or triangle in B, C, and D represents

data from an individual mouse. Each dot, square, or triangle in F and G represents one model

estimate. Error bars in C represent mean ± SD. *P*-values in C were calculated via unpaired *t*-

test using log_{10}-transformed values. *P*-values in F were calculated via pairwise-corrected Mann-

Whitney's test using Bonferroni's correction. *P*- and r_s-value in G were calculated via

Spearman's rank correlation. Exact *P*-values are given for all values <0.05.

Figure 7. Differentiation and decay of the CVT-resident CD8 T cell compartment is tissue-

intrinsic (A) Schematic of experiment to test effect of depleting circulating gBT-I cells. **(B)**

Example flow staining to identify gBT-I population within CD8 T cells from spleen and CVT of

depleted or undepleted mice 1mo after immunization. **(C)** Number of gBT-I cells recovered from

spleen or CVT of mice at indicated time points after immunization. **(D-E)** Relative abundance of

CD69 and CD103 subsets within i.v. label$^-$ gBT-I cells in CVT at indicated time points. **(F)**

Frequency of granzyme B expression among gBT-I cells. **(G)** Frequency of TCF-1 expression

among i.v. label$^-$ gBT-I cells. Data in C-G are pooled from 2 independent experiments. Flow

plots in B are derived from two representative mice. Each dot in C, E, F, and G represents an

individual mouse. Error bars represent mean ± SD. *P*-values in C, E, and F were calculated via

unpaired *t*-test comparing 1mo to 3mo within each depletion group. *P*-values in G were

calculated via ordinary one-way ANOVA with Tukey's post-hoc test. Exact *P*-values are given

for all values <0.05.

Figure 8. Decay of the CVT-resident CD8 T cell compartment occurs regardless of

priming infection and route. (A) Schematic of experiment to assess effect of vaginal

immunization on longevity of memory CD8 T cells in CVT. **(B)** Number of gB tetramer$^+$ cells

recovered from spleen, SI LP, or CVT of mice at indicated time points after immunization. Each

dot in B represents an individual mouse. Error bars represent mean ± SD. *P*-values in B were

calculated via unpaired *t*-test comparing 1mo to 3mo within each immunization group. Exact *P*-

values are given for all values <0.05.

Figure 1

A   Human Vaginal Biopsy

B   % of Memory CD8
   - CD69- CD103-
   - CD69+ CD103-
   - CD69- CD103+
   - CD69+ CD103+

C   Granzyme B+ 41.5
   GzmB+, % of Memory CD8

D   CD103+, % of GzmB+

E   ○ CVT, Memory CD8
   ○ Blood, Memory CD8
   P = 0.0008
   TCF-1+, % of Memory CD8

Figure 2

A

HSV-2 TK- i.vag.
gBT-I cells   Assess CD8 T cells

-7d 0d   22d   1mo

2 mg progesterone s.c.

Assess CVT pathology

B

Mouse 1mo CVT

HSV gB tet APC / CD8 PerCP-eF710: 17.0

CD103 BV786 / CD69 PE-Dz594: 2.88, 53.8, 17.3, 26.0

% of gB tet+ — CVT

☐ CD69- CD103-
▨ CD69+ CD103-
▨ CD69- CD103+
▨ CD69+ CD103+

C

GzmB Pacific Blue / CD8 PerCP-eF710: 42.3

P < 0.0001

GzmB+, % of gB tet+ — CVT, dLN

D

GzmB Pacific Blue / TCF-1 PE: 39.4, 0.96, 43.3, 16.3

P < 0.0001

TCF-1+, % of gB tet+ — CVT, dLN

E

Uninfected   22d, HSV TK- i.vag.

Lamina propria inflammation score — Day Post-Immunization

HSV TK- i.vag.

Figure 3

A

LM-gB i.v.
gBT-I cells

Assess gB-specific CD8⁺ T cells in
lymphoid and mucosal tissues

0d   22d   1mo   5mo

Assess CVT pathology

B

22d, LM-gB i.v.

Lamina propria inflammation score

HSV TK- i.vag.
LM-gB i.v.

P = 0.013

Day Post-Immunization

C

1wk Blood

HSV gB tet PE

36.7

CD44 APC-eF780

D

1mo

# of gB tet+ cells

CVT   Spleen   dLN

E

○ CVT, gB tet+ i.v. label-
○ dLN, gB tet+ i.v. label-

1mo CVT, i.v. label-

CD103 BV786

0.2   1.64
22.3   0.97

42   34.7
11.78   96.4

TCF-1 PE

TCF-1+, % of gB tet+

P < 0.0001
P = 0.0002
P = 0.0017

CVT CD103+   CVT CD103-   dLN

F

Blood

gB tet+, % of CD8 T cells

1wk   2wk   1mo   3mo   5mo

Post-Immunization Timepoint

G

# of gB tet+ cells

Spleen
n.s.
1mo   5mo

dLN
P = 0.0064
1mo   5mo

SI LP
n.s.
1mo   5mo

CVT
P < 0.0001
1mo   5mo

Post-Immunization Timepoint

Figure 4

A Spleen dLN CVT SI LP

B 1mo CVT, gB tet+   5mo CVT, gB tet+

C CVT, i.v. label-   CVT, i.v. label+   SI LP, i.v. label-

☐ CD69- CD103-
▨ CD69+ CD103-
▨ CD69- CD103+
▨ CD69+ CD103+

D CVT, i.v. label-

E CD69- CD103-   CD69+ CD103-   CD69- CD103+   CD69+ CD103+

Figure 5

A

Spleen   dLN

GzmB+, % of gB tet+

1mo 3mo 5mo   1mo 3mo 5mo

CVT   SI LP

P = 0.009
P = 0.0063
n.s.

1mo 3mo 5mo   1mo 3mo 5mo

B

CVT

# of GzmB+ gB tet+

P < 0.0004

1mo 3mo 5mo

C

○ CVT, gB tet⁺ i.v. label⁻
○ dLN, gB tet⁺ i.v. label⁻

CD103 BV786

7.31   0
33.8   31.2

34.4   0.53
92.2   0.51

GzmB Pacific Blue

D

CD103+, % of gB tet+ GzmB+

CVT

1mo 3mo 5mo

SI LP

1mo 3mo 5mo

E

% of gB tet+ GzmB+

CVT, Granzyme B+

1mo 3mo 5mo

SI LP, Granzyme B+

1mo 3mo 5mo

☐ CD69- CD103-
▨ CD69+ CD103-
▨ CD69- CD103+
▨ CD69+ CD103+

Figure 6

Figure 7

Figure 8

A

B

Spleen

SI LP

CVT