Running title: Sex Lability and Dimorphism

Sex Lability and Dimorphism in Diecious Palmer Amaranth (*Amaranthus palmeri*)

Mohsen B. Mesgaran[1, †], Maor Matzrafi[2, 3, †] and Sara Ohadi[4]

[1]Assistant Professor (ORCID 0000-0002-0841-6188), Department of Plant Sciences, University

of California, Davis, One Shields Avenue, Davis, CA 95616

[2]Former Postdoctoral Research Associate (ORCID 0000-0002-4867-0850), Department of Plant

Sciences, University of California, Davis, One Shields Avenue, Davis, CA 95616

[3]Current address: Department of Plant Pathology and Weed Research, Agricultural Research

Organization, Newe Ya'ar Research Center, Israel.

[4]Postdoctrol Research Associate, Department of Plant Sciences, University of California, Davis,

One Shields Avenue, Davis, CA 95616

Author for correspondence: Correspondence: Mohsen B. Mesgaran, Assistant Professor,

Department of Plant Sciences, University of California, Davis, One Shields Avenue, Davis, CA

95616

(E-mail: mbmesgaran@ucdavis.edu)

[†]These authors have contributed equally to this work

Abstract

Dioecious weeds (separate sexes) may benefit from a maximized outcrossing and optimal sex-

specific resource allocation but there are costs associated with the evolution of this breeding

system which can be exploited for long-term management of dioecious weeds. That is, seed

production in dioecious species is contingent upon the co-occurrence and co-flowering of the

two genders and can be further complicated by biases in sex ratio. We therefore explored the sex

ratio and dimorphism in secondary sex characters in three populations of Palmer amaranth

(*Amaranthus palmeri*) from California, Kansas and Texas and tested if water stress can change

the sex expression and/or the synchrony of flowering (anthesis) between male and female plants.

Sex ratio (proportion of males) was balanced and did not deviate from 1:1 in all experiments and

populations (California, Kansas, Texas) when plants received normal watering. Male and female

plants of *A. palmeri* did not differ in timing of emergence, plant height and relative growth rate.

While the initiation of flowering (emergence of inflorescence) occurred earlier in males than

females, females preceded males in timing of anthesis. Water stress delayed anthesis in males to

a larger extent than females giving rise to an anthesis mismatch as large as seven days between

the two sexes. Water stress induced a female sex expression in Kansas population giving rise to a

female to male ratio of 1.78 which significantly differed from the equal 1:1 sex ratio. Our data

provide the first evidence of sex lability and environment sex determination (ESD) in *A. palmeri,*

suggesting manipulation of sex expression and phenological asynchrony as novel tools for

ecological management of dioecious weeds.

Key words: Sex ratio, environmental sex determination (ESD), secondary sex characters,

flowering overlap, gender specialization, environmental stress

Introduction

Hermaphroditism represents the most common mode of sexual reproduction in weeds whilst dioecy (separate male and female individuals) is rare and have been observed in about 5-7% of weed species (Sutherland 2004). Whereas a selfing breeding system is often associated with a weedy character (Baker 1955; Razanajatovo et al. 2016), it might come as a surprise that some dioecious weeds such as palmer amaranth (*Amaranthus palmeri* S. Watson) and common waterhemp (*A. tuberculatus* (Moq.) J. D. Sauer) have been reported amongst the most the troublesome weeds in US (Van Wychen 2017). Contrary to hermaphroditism (particularly selfing), dioecy maximizes outcrossing and thereby reduces the likelihood of inbreeding depression (Thomson and Barrett 1981,1990; Charlesworth 1999). Dioecy also has the benefit of optimizing the allocation of resources between male and female functions by minimizing the competition between these two functions through sexual specialization, i.e. the so-call "division of labor" theory (Maynard Smith 1978; Charnov 1982).

Despite these advantages, a dioecious breeding system is subject to several demographic handicaps. First, the two sexes must co-occur in the same habitat and, unlike hermaphrodite weeds, dioecious plants cannot propagate from a single individual. Second, the two sexes need to flower at the same time. The smaller the degree of temporal overlapping between male and female flowering, the lower the chance of successful fertilization and hence seed production. Third, the fecundity of a dioecious population can be affected by the sex ratio (the proportion of males) and is likely to decline if the population becomes skewed towards one sex: a male-biased population may not achieve its maximum seed production because of the ovule limitation, whilst in a female-biased population, limited pollen availability may reduce the seed output. Therefore, it seems that there exists an optimal sex ratio at which the population-level seed production is

maximized, but beyond that optimal point, the seed output is likely to decline due to pollen or

ovule limitations (Mulcahy 1967).

We believe that novel weed management strategies can be developed by better understanding of

the above "cracks" in the evolution of a dioicous mating system. Two potential strategies

envisioned involve shifting the sex ratio towards one sex (sub-optimal sex ratio strategy) and

reducing the flowering overlap between the two sexes (phenological isolation strategy). The

proposed strategies rely on the assumptions that the sex is labile–hence sex ratio can be

changed–and that we can manipulate the flowering phenology of male and female individuals

differentially. For long sex was posited to be a fixed trait of the plant throughout its life cycle

and genetic sex determination (GSD), where genes/chromosomes specify the gender of an

individual, was considered as the sole mechanism by which the sex of an individual (dioecious)

plant is determined (Fuller and Ritchie 1967). Evidence, however, has mounted over the past half

century showing that sex can be labile and the genetic control of sex can be overridden by the

environment (reviewed in Freeman et al. 1980; Korpelainen 1998; Vega-Frutis et al. 2014):

Environmental sex determination (ESD) has been found in many plants where the expression of

sex has changed in response to the environmental factors such as soil moisture (Freeman and

Vitale 1985), nutrients (Dorken and Barrett 2004), light intensity (Zimmerman 1991),

photoperiod (Thompson 1956), plant density (Onyekwelu and Harper 1979), and exogenous

application of plant hormones (Chailakhyan 1979). Environmental stresses such as water stress

(drought) are more likely to induce maleness, while conditions conducive to growth e.g. high soil

nutrition favor a female sexuality (Freeman et al., 1980), perhaps because reproduction is more

costly to females than males (Delph, 1999; Meagher, 1988).

In many dioicous species, male and female individuals have shown to differ in secondary sex

characters (i.e. traits other than the sex itself) related to morphology, physiology, life history and

phenology (reviewed in Geber et al. 1999; Juvany and Munné-Bosc 2015; Lloyd and Webb

1977). Differences in progression of phenological events (e.g. timing of germination) can result

in temporal segregation of males and females, ultimately reducing the chance of successful

cross-pollination in a dioecious population. In fact, most dioecious species seems to be

protandrous i.e. males flower earlier than females (Forero-Montaña and Zimmerman 2010;

Forest 2014). As female's investment in reproduction is greater than males, it has been

hypothesized that females postpone flowering (reproduction) for longer period of resource

accumulation (Lloyd and Webb 1977; Purrington and Schmitt 1998). This hypothesis has direct

implications for manipulative separation of flowering timing between the two sexes. That is, if

requirement for resource accumulation delays flowering in females, then environmental stresses

can further isolate the females, exacerbating the flowering mismatch between the two sexes.

In this study, we attempted to test the viability of exploiting the demographic handicap

associated with a unisexual breeding system (i.e. shift in sex ratio and phenological isolation) for

novel management of diecious weeds. We used palmer amaranth (*Amaranthus palmeri* S.

Watson) as our diecious model system. The species has been ranked as the worst weed in US

corn fields in a recent survey (Van Wychen 2017) and evolved resistance to six classes of

herbicides with 56 herbicide-resistant biotypes found in the US (Heap 2018). Most of these

biotypes were found to be resistant to the widely used herbicide, glyphosate (Roundup®),

requiring the development of non-chemical control methods. The objectives of this study were

threefold: 1) determine if the male and female individuals of *A. palmeri* differ in secondary sex

characters related to growth and development, 2) examine sex lability (shift in sex ratio) in *A.*

palmeri in response to environmental stress (drought), and 3) test if the degree of flowering

synchrony between *A. palmeri* male and female plants can be affected by the environmental

stress (drought). As a proof of concept study, we used water stress for the sex change experiment

as this environmental factor has been found to affect sex expression in several plant species

(Freeman et al., 1980; Korpelainen, 1998).

Materials and Methods

Seeds of *A. palmeri* from California (CA) and Kansas (KS) were cordially provided by Dr. Anil

Shrestha (California State University, Fresno, California) and Dr. Dallas E. Peterson (Kansas

State University, Manhattan, Kansas), respectively. Seeds of a third population were collected

from mature plants (about 30 plants) from a set-aside field in College Station, Texas (TX), in the

summer of 2017. Seeds were stored at 4ºC before use. Two sets of experiments were conducted

in the greenhouse facility (Orchard Parks) of University of California, Davis, during the spring

and summer of 2018.

Experiment 1. Natural sex ratio and dimorphism in secondary sex characters

The aim of this experiment was to examine whether *A. palmeri* has undergone divergent

evolution in secondary sex traits related to growth and development. We therefore compared

male and female plants of *A. palmeri* for potential differences in emergence pattern, timing of

flowering and anthesis, plant height, and relative growth rate (RGR). By using three populations

(CA, KS, and TX) and large number of plants (~250), *Experiment 1* also served as a platform for

a robust estimate of baseline sex ratio in *A. palmeri*, which was the major trait to be examined in

the second experiment (*Experiment 2*, see below). In July 2018, we sow 200 seeds from each

population, into 5 by 5 by 5 cm plastic pots (one seed per pot) filled with a soil mix (1:1:1:3

sand:compost:peat:dolomite). We recorded the emergence time for each individual seed and then

transplanted the young plants (two-four leaf stage) into larger plastic pots (2.37 L) filled with a

soil mix (1:1 sand/peat) plus a controlled-release fertilizer (15–9–12, 150 g 75 L^{-1}; Scotts

Osmocote PLUS, Mississauga, ON). Pots were randomly placed in a net house under natural

California-summer conditions and watered twice a day through an automated drip irrigation.

Plant height was measured three times (11 July, 23 July, and 24 September) during the season.

We inspected plants daily and an individual was recorded to be at the flowering stage when the

tip of its inflorescence was visible either on the terminal or lateral buds. The onset of anthesis for

a flowering plant was then determined when the stigma (in pistillate flowers) or dehisced anther

(in staminate flower) become visible. At this stage, i.e. anthesis, we also recorded the sex of

plants.

Experiment 2. Sex expression and flowering synchrony in response to water stress

This experiment aimed at evaluating the effect of water stress on the sex ratio and flowering

synchrony between the male and female plants of *A. palmeri*. Because of space limitation we

only used two populations (CA and KS) in this experiment. The experiment was conducted twice

(April-July and July-September in 2018) in a completely randomized design with 50 replicates

(plants) per treatment (water stress *vs.* well-watered control). About 10 seeds were sown in each

round plastic pot (2.37 L), filled with a soil mix (1:1 sand/peat) plus a controlled-release

fertilizer (15 – 9 – 12, 150 g 75 L^{-1}; Scotts Osmocote PLUS, Mississauga, ON). Pots were placed

in a greenhouse set at a temperature of 32/22ºC (day/night) with a day length of 16 hours

provided through supplementary lighting between 5:00 to 9:00 am and 5:00 to 9:00 pm. Seedling

were thinned haphazardly several times to achieve one plant per pot by their four-leaf stage.

When plants reached six to seven true-leaf stage, 50 random plants from each population were

randomly assigned to well-watered or water deficit treatments. Well-watered plants received

water from four emitters inserted into the potting medium to deliver 65 mL of water min^{-1} for

two mins and twice per day (7 am and 2:00 pm). Water deficit treatment was achieved by

removing three out of four drip emitters from designated water stress treatment pots, reducing

the amount of water by approximately 75%. Additional irrigation (~100 mL twice per week) was

added to water deficit plants when severe visual stress symptoms were observed. Plants were

sexed by examining flowers for the presence of anther (in males) or stigma (in females). For

each plant we also measured time to flowering (inflorescences) and time to anthesis as described

in *Experiment 1*.

Statistical analysis

In Experiment 1, a binomial test (two-tailed) was used to test if the observed sex ratio in each

population statistically deviates from the expected 1:1 sex ratio (i.e. 50% male). Binomial test

was conducted by `binom.test()` function in `stats` package of R (R Core Team, 2018). Using a

t-test, male and female plants were compared for the secondary sex characters including time to

flowering (inflorescence), time to anthesis, plant height, and early stage RGR. Function `t.test()`

of 'stats' package (R Core Team, 2018) was used for the t-test analysis. The early stage RGR

was calculated based on the height gain between the first and second height measurements using

the following formula (Radosevich et al. 2007):

$$RGR = \frac{\log(H_2) - \log(H_1)}{t_2 - t_1} \qquad [1]$$

176   where H_1 and H_2 are plant heights measured at time t_1 (11 July) and t_2 (23 July), respectively (log

177   stands for natural logarithm). To explore difference in seedling emergence pattern between the

178   sexes, we fitted the following two-parameter logistic equation to cumulative emergence data of

179   *A. palmeri* using the `drm()` function of `drc` package (Ritz et al. 2016):

$$E(t) = \frac{1}{1 + \left(b(\log(t) - \log(t_{50}))\right)} \qquad [2]$$

where *E(t)* is the cumulative emergence over time, *t*, t_{50} is time to 50% emergence while *b*

denotes the steepness of emergence curve around t_{50}.

In *Experiment 2*, for each water treatment level we tested the deviation of sex ratio from 1:1 by

using a binomial test (two-tailed) as described above. Phenology data (time to inflorescence and

time to anthesis) were subject to analysis of variance, ANOVA, using the `lm()` and `Anova()`

functions of `stats` (R Core Team, 2018) and `car` (Fox et al. 2019) packages in R, respectively.

Inspection of residuals showed no predictable pattern when plotted against fitted values

suggesting that the ANOVA's assumption of homogeneity of variance has been met. The degree

of flowering/anthesis overlap between the two sexes was visually inspected by fitting a kernel

density smoother to time data (i.e. time to flowering and time to anthesis) using the

`geom_density_ridges()` function of `ggplot2` R package (Wickham et al. 2019).

Results and Discussion

Experiment 1. Natural sex ratio and dimorphism in secondary sex characters

Sex Ratio. The recorded sex ratio was balanced in all three tested populations and did not

deviate from the expected 1:1 male/female ratio as indicated by a binomial test (Figure 1).

Although California population had slightly higher proportion of males (57%) than females, the

departure from 50% sex ratio was not significant (P = 0.306) (Figure 1).

An equal sex ratio was found in *A. palmeri* plants grown under greenhouse conditions with no

water or nutrient limitations (Giacomini et al., 2014). A balanced sex ratio of 55% (22 males vs.

18 females) also was reported by Lemen (1980); however, he does not provide any information

about method of measuring this parameter e.g. whether sex has been determined in plants grown

from the seed or sex ratio has been measured under field conditions. Keeley et al. (1987) also

reported a male:female ratio of 47%:53%—which shows a balanced sex ratio—for a field grown

A. palmeri. However, the source of seed for this test population was collected from one single

plant suggesting that their reported sex ratio could best represent the sex ratio of that individual

and not the population. Of 300 pot-grown *A. palmeri* plants—for a pollen mediate gene flow

study—172 (~60%) and 162 (%54) plants became male in the first (2005) and second year

(2006) of experiment (Sosnoskie et al. 2012). The sex ratio has clearly (and statistically) been

male-based in the first year according to a binomial test (*P-vale* = 0.0009), though the sex ratio

data were not analyzed in the original article.

Seedling Emergence. Total emergence for California, Kansas, and Texas populations were 31%,

55%, and 35% (n=200 for each individual population), respectively. Logistic model (Eq.2) fitted

the emergence data adequately as evidenced in Figure 2 and small root mean square errors

(RMSE ≤ 4%) in all populations. Female and male plants of *A. palmeri* exhibited almost

identical emergence pattern and in none of the tested populations the emergence curves differed

between the sexes. Populations, however, differed in emergence rate (rapidity) with Kansas

being the fastest emerging population (t_{50} = 2.5±0.10 days) while California was the slowest (t_{50}

= 4.3±0.12 days) irrespective of the gender (see inset in Figure 2).

Time to Flowering and Anthesis. In all three populations, the initiation of flowering (i.e. tip of

inflorescence becoming visible) occurred earlier in male than female plants of *A. palmeri* (Figure

3). The difference in timing of flowering was particularly profound in Texas population where

males, on average, flowered ~9 days before females. Variability in flowering time seemed to be

similar between males and females as illustrated by the violin plots. The only exception was

observed with California population where the flowering bulge occurred close to the earliest

flowering times (i.e. at the bottom of the violin) as opposed to females that had their bulge

above the median time of flowering. Note that a bulge on a violin plot represents the value (here

flowering time) with the highest probability of occurrence. The male's earlier flowering may

suggest a protandrous flowering habit in *A. plameri*; however, when we investigated the timing

of anthesis, which might be regard as the effective flowering event, we arrived at an opposite

conclusion.

In both, California and Texas populations, anthesis happened four days earlier in females than

males (Figure 3). For Texas population, although the median time to anthesis was the same for

male and females, the most probable anthesis time (i.e. the bulge) was at an earlier time in

females compared to males. While male plants of *A. palmeri* began to flower earlier than female

plants, it took them a much longer time to mature as shown in Figure 4. For females, regardless

of the population, the length of time from the appearance of inflorescence to anthesis (visible

stigma) was no more than 4-5 days compared with a 11-day time period needed for males to

develop mature anthers following the initiation of flowering. This observation suggests that *A.*

palmeri should be classified as a protogynous species as its pistillate flowers mature earlier than

the staminate ones (Figure 4). Korres et al. (2017) also showed that female plants of *A. palmeri*

flowered 6 to 8 d earlier than male plants, a flowering pattern that is very similar to our result from

Kansan and California population. However, Korres et al. (2017) do not explicitly define "flowering"

stage and we are not sure if our "anthesis" represents the same reproductive stage as with the

"flowering" stage in Korres et al. (2017). A synchronous anthesis between male and female plants

observed with Texas population resembles the results of Giacomini et al. (2014) who found no

differences in days to first flowering (anthesis) between male and female plants of *A. palmeri*.

Plant Height and Relative Growth Rate (RGR). We observed no sex-specific differences in

final plant height for the three tested populations (Figure 5). The only significant sexual

dimorphism in growth was related to the height-based RGR in Texas population. At the early

stage of growth, males of this population grew tall at a rate that was ~13% faster than females.

However, the male's faster initial growth did not translate into significant larger plants as males

attained a shorter stature (125.3 ± 3.78 cm) than females (133.3 ± 4.69 cm) at the end of growing

season. In the study of Webster and Grey (2015) plant height did not differ between *A. palmeri*

sexes either grown alone or in competition with cotton, but sex differential growth was detected

in other traits e.g. females had wider canopy and produced more than double the biomass of

males. Several other studies have shown that female *A. palmeri* plants tend to be taller and grow

faster than males (e.g. Keeley et al., 1987; Korres et al., 2017). Whereas *A. palmeri* populations

originating from ten contrasting cropping systems differed in many morphological and growth

traits, no sexual dimorphism was reported in these populations (Bravo et al. 2017).

Experiment 2. Sex expression and flowering synchrony in response to water stress

Sex Ratio. While sex ratio remained balanced across water treatments in California population,

exposure to drought produced a heavily female-biased sex ratio (~1.78 times more female than

males) in Kansas population. Sex ratio significantly deviated from 50% (P-value = 0.009, n = 99)

in water stress treatment whilst it was balanced (P-value = 0.547, n = 99) when *A. palmeri* plants

received normal irrigation (Figure 6). Our data provide the first evidence of sex lability and

environment sex determination (ESD) in *A. palmeri:* water stress induced a female sex

expression in Kansas population.

To the best of our knowledge, our study provides the first case of a feminizing effect for water

stress: almost all previous work has reported a masculinizing effect for this stressor. Since

reproduction is more costly to females than males (Delph 1999; Meagher 1988), sex allocation

theory predicts that the environmental conditions conducive to plant growth should favor a

female sex expression while resource-poor environments should induce maleness (Charnov

1982). Empirical data support the prediction of sex allocation theory. Reviews by Freeman et al.

(1980) and Korpelainen (1998) show that water stress has induced maleness in all 11 species

(five dioecious and six monoecious) that have been tested for potential sex change in response to

draught. Limited field surveys have also recorded overrepresentation of males in resource-poor

habitats whereas females were more abundant in benign habitats (Policansky, 1981; Lovett

Doust and Cavers 1982). The observed overrepresentation of males under stressful environments

may be due to a sex differential mortality rather than being the result of an actual sex change.

Most studies on sex ratio, however, lack the capacity to disentangle these two processes i.e. sex

lability *vs.* sex-differential survival. By randomly thinning seedlings (with no bias towards

selection of vigorous seedlings), our study was able to rule out differential mortality as the

potential driver of shift in sex ratio. In our experiments, only two plants (one per water treatment

in Kansas population) died over the course of experiment.

Time to Flowering and Anthesis. Flowering phenology was recorded for plants from California

and Kansas populations grown under different irrigation regimes. Similar to *Experiment 1*, the

initiation of flowering happened earlier in males than females with water stress having no effects

on the flowering sequence between the two sexes (Figure 7). However, water stress slightly

delayed flowering, almost to the same extent, in both male and female plants giving rise to a

more temporally staggered flowering pattern as compared with the control. Consistent with

Experiment 1, *A. palmeri* female plants preceded males in timing of anthesis in the water stress

experiment (*Experiment 2*; Figure 8). Sex-differential responses to water stress were observed

with the timing of anthesis: water stress distorted the frequency distribution of anthesis timing to

a larger extent in males than females (Figure 8). Time to anthesis nearly followed a unimodal

distribution in both sexes when plants received normal irrigation but became multimodal and

"wavy" when plants were exposed to water stress. Anthesis was much synchronized between

male and female plants of *A. palmeri* under normal watering but the two sexes became more

separate (i.e. less overlapping in anthesis) with water stress. In both populations, the difference

between males and females in median time to anthesis (arrows in Figure 8) was only two days in

control but extended to seven days in water-stressed plants. The increased asynchrony between

the two sexes was mainly driven by a differential response of the sexes to water stress. While

water stress delayed anthesis in both sexes, the delay in males was much longer than females

(Figure 8). In California population, for example, water stress postponed anthesis (median time)

by seven days in male plants compared with a 2-day delay in females.

Decreasing the level of irradiance (i.e. greater stress) also postponed *A. palmeri* flowering in the

study of Korres et al. (2017). Plants experiencing stressful environments often accelerate their

transition to reproductive stage; perhaps as a potential adaptive strategy to produce seed before

they die due to stress (Wada and Takeno 2010). However, depending on the intensity and

duration of stress and plant development stage, stress may produce an opposite response (i.e.

delay the flowering) by slowing the metabolism in plant (Cho et al. 2017). The delay in

flowering of *A. palmeri* observed in Korres et al. (2017) and our study might have resulted from

a slowed metabolism. However, contrary to our initial hypothesis, the timing of

flowering/anthesis in females was less sensitive to stress than males. The prevalence of protandry

in diecious species has been attributed to the greater cost of reproduction incurred by females

(Lloyd and Webb 1977; Forest, 2014). As females need to produce seeds, they may undergo an

extended period of resources accumulation to ensure sufficient assimilates will be available for

allocation to seeds (Lloyd and Webb 1977; Purrington and Schmitt 1998). As a result, stress was

proposed to exert a stronger effect on the flowering phenology of females than that of males; a

hypothesis that was not supported by our data.

Implications for weed management

Our study showed that sex can be labile in *A. palmeri* and change in response to external

environmental conditions (here water stress). Sex lability can be exploited as new tool for long-

term management of *A. palmeri* (and other diecious weeds): by shifting the sex ratio towards the

extremes we might be able to reduce seed production and hence suppress the population growth.

A variety of external factors can be used to shift sex towards the desirable direction (e.g. plant

hormones, nutrient and water management, light intensity, and herbicides) but currently we have

no information about the effects of these factors on the sex expression of *A. palmeri*. Increasing

the number of males might offer a more viable and safer management option as males do not

contribute to seed production. However, if the population is severely pollen limited, increase in

the share of females may also reduce the seed production capacity of population. A quantitative

understanding of the relation between thee sex ratio and seed output is needed to determine the

proper sex conversion strategy that can potentially disrupt seed production.

Our study clearly demonstrated that water stress can result in temporal separation of the two

sexes in *A. palmeri*, offering a potential means for ecological management of this weed and

perhaps other diecious weeds. Asynchrony in timing of anthesis can reduce the chance of cross

pollination and hence reduce seed production in unisexual weeds such as *A. palmeri* and

common waterhemp (*A. tuberculatus* (Moq.) J. D. Sauer). While the direct (negative) effect of

water limitation on seed production is well understood, our study suggests a novel mechanism

through which water stress (and perhaps other stressors) can reduce seed production indirectly by

exacerbating the flowering/anthesis mismatch between the sexes in a diecious weed. Future work

should attempt to disentangle the direct effects of environmental stressors form their indirect

effects on seed production, a subject which was out of the scope of the current study.

Acknowledgments

This research was funded by UC Davis New Research Initiatives and Interdisciplinary Research

Grant. The authors would also like to thank Hannah Clifton and William Werner for greenhouse

maintenance.

References

Baker HG (1955) Self☐compatibility and establishment after '"long☐distance" dispersal.

Evolution 9:347-349.

Bravo W, Leon RG, Ferrell JA, Mulvaney MJ, Wood CW (2017) Differentiation of life-history

traits among Palmer amaranth populations (*Amaranthus palmeri*) and its relation to cropping

systems and glyphosate sensitivity. Weed Sci 65: 339-349.

Chailakhyan MK (1979) Genetic and hormonal regulation of growth, flowering, and sex

expression in plants. Am J Bot 66: 717–736.

Charlesworth D (1999) Theories of the evolution of dioecy. In Gender and sexual dimorphism in

flowering plants (pp. 33-60). Springer, Berlin, Heidelberg.

Charnov EL (1982) The Theory of Sex Allocation. Princeton University Press, Princeton, NJ.

Cho LH, Yoon J, An G (2017) The control of flowering time by environmental factors. Plant J

90:708-719.

Delph LF (1999) Sexual dimorphism in life history. *In* Gender and Sexual Dimorphism in

Flowering Plants (Geber, M. A., Dawson, T. E. & Delph, L. F. eds.), pp. 149–173, Springer-

Verlag, New York.

Dorken ME, Barrett SC (2004) Sex determination and the evolution of dioecy from monoecy in

Sagittaria latifolia (Alismataceae). Proc R Soc Lond [Biol] 271: 213-219.

Forero-Montaña J, Zimmerman JK (2010) Sexual dimorphism in the timing of flowering in two

dioecious trees in a subtropical wet forest, Puerto Rico. Caribbean J Sci 46: 88-95.

Forrest JR (2014) Plant size, sexual selection, and the evolution of protandry in dioecious plants.

Am Nat 184:338-351.

Fox J, Weisberg S, Adler D, Bates D, Baud-Bovy G, Ellison S, Firth D, Friendly M, Gorjanc G,

Graves S, Heiberger R (2019) Package 'car': Companion to applied regression. https://CRAN.R-

project.org/package=car

Freeman DC, Harper KT, Charnov EL (1980) Sex change in plants: old and new observations

and new hypotheses. Oecol 47:222-232.

Freeman DC, Vitale JJ (1985) The influence of environment on the sex ratio and fitness of

spinach. Botanical Gaz 146:137-142.

Fuller HJ, Ritchie DD (1967) General Botany, Barnes and Noble. Inc., New York. Geber MA

Dawson TE (1999) Gender and sexual dimorphism in flowering plants. Springer Science &

Business Media.

Giacomini D, Westra P, Ward SM (2014). Impact of genetic background in fitness cost studies:

an example from glyphosate-resistant Palmer amaranth. Weed Sci 62:29-37.

Heap I (2018) The International Survey of Herbicide Resistant Weeds. Online. Internet

Available at: www.weedscience.com (accessed Tuesday, June 17, 2019).

https://CRAN.R-project.org/package=car

Juvany M, Munné-Bosch S (2015) Sex-related differences in stress tolerance in dioecious

plants: a critical appraisal in a physiological context. J Exp Bot 66:6083-6092.

Keeley PE, Carter CH, Thullen RJ (1987) Influence of planting date on growth of Palmer

amaranth (*Amaranthus palmeri*). Weed Sci 35:199-204.

Khryanin VN (2002). Role of phytohormones in sex differentiation in plants. Russian J Plant

Physiol 49:545-551.

Korpelainen H (1998) Labile sex expression in plants. Biological Rev 73:157-180.

Korres NE, Norsworthy JK (2017) Palmer amaranth (*Amaranthus palmeri*) demographic and

biological characteristics in wide-row soybean Weed Sci 65:491-503.

Korres NE, Norsworthy JK, FitzSimons T, Roberts TL, Oosterhuis DM (2017) Differential

response of Palmer amaranth (*Amaranthus palmeri*) gender to abiotic stress. Weed Sci 65:213-

227.

Lemen C (1980) Allocation of reproductive effort to the male and female strategies in wind-

pollinated plants. Oecol 45:156-159.

Lloyd D G, Webb CJ (1977) Secondary sex characters in plants. Bot Rev 43:177-216.

Lovett Doust J, Cavers PB (1982) Sex and gender dynamics in jack-in-the-pulpit, *Arisaetna*

triphyllum (Araceae) Ecology 63:797-808.

Maynard Smith J (1978). The evolution of sex (No. 574.1 S5). Cambridge: Cambridge

University Press.

Meagher TR (1988). Sex determination in plants. *In* Plant reproductive ecology: patterns and

strategies. (Lovett Doust, J. & Lovett Doust, L. eds), pp. 125-138. Oxford University Press. New

York.

Morgan PW, Drew MC (1997) Ethylene and plant responses to stress. Physiol Plantarum,

100:620-630.

Mulcahy DL (1967) Optimal sex ratio in *Silene alba*. Heredity 22:411-423.

Onyekwelu SS, Harper JL (1979) Sex ratio and niche differentiation in spinach (*Spinacia*

oleracea L.). Nature 282:609-611.

Policansky D (1981) Sex choice and the size advantage model in jack-in-the-pulpit (*Arisaema*

triphyllum). PNAS 78:1306-1308.

Pospíšilová J, Synková H, Rulcová J (2000) Cytokinins and water stress. Biologia Plantarum

43:321-328.

Purrington CB, Schmitt J (1998) Consequences of sexually dimorphic timing of emergence and

flowering in *Silene latifolia*. J Ecol 86:397-404.

R Core Team (2018) R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Radosevich SR, Holt JS, Ghersa CM (2007) Ecology of weeds and invasive plants: relationship

to agriculture and natural resource management. John Wiley & Sons.

Razanajatovo M, Maurel N, Dawson W, Essl F, Kreft H, Pergl J, Pyšek P, Weigelt P, Winter M,

Van Kleunen M (2016). Plants capable of selfing are more likely to become naturalized. Nat

Comm 7:13313

Ritz C, Strebig JC, Ritz MC (2016) Package 'drc'. Analysis of dose-response curves.

http://www.bioassay.dk

Sosnoskie LM, Webster TM, Kichler JM, MacRae, AW, Grey TL, Culpepper AS (2012) Pollen-

mediated dispersal of glyphosate-resistance in Palmer amaranth under field conditions Weed Sci

60:366-373.

Sutherland S (2004) What makes a weed a weed: life history traits of native and exotic plants in

the USA. Oecol 141:24-39.

Thompson AE (1956) Methods of producing first-generation hybrid seed in spinach. Cornell

Univ Agr Exp Sta Mem 336:3-48.

Thompson FL, Eckert CG (2004) Trade☐offs between sexual and clonal reproduction in an

aquatic plant: experimental manipulations vs. phenotypic correlations. J Evol Biol 17:581-592.

Thomson J D, Barrett SC (1981) Selection for outcrossing, sexual selection, and the evolution of

dioecy in plants. Am Nat 118:443-449.

Thomson J D, Brunet J (1990) Hypotheses for the evolution of dioecy in seed plants. Trends

Ecol Evol 5:11-16.

Van Wychen L (2017) Survey of the most common and troublesome weeds in grass crops,

pasture and turf in the United States and Canada. Weed Science Society of America National

Weed Survey Dataset. Available: http://wssa.net/wp-content/uploads/2017-Weed-Survey_Grass-

crops.xlsx.

Vega☐Frutis R, Macías☐Ordóñez R, Guevara R, Fromhage L (2014) Sex change in plants and

animals: a unified perspective. J Evol Biol 27:667-675.

Wada KC, Takeno K (2010) Stress-induced flowering. Plant Signal Behav 5:944-947.

Webster TM, Grey TL (2015) Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*)

morphology, growth, and seed production in Georgia. Weed Sci 63:264-272.

Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, Woo K, Yutani H (2019)

Package 'ggplot2': Create elegant data visualisations using the grammar of graphics,

http://ggplot2.tidyverse.org

Zimmerman JK (1991) Ecological correlates of labile sex expression in the orchid *Catasetum*

viridiflavum. Ecology 72:597-608.

Figure 1. Observed sex ratios in three *A. palmeri* populations grown in pots during the summer

of 2018 in California, Davis. Also reported are the P-values (shown as P) for testing the

deviation of sex ratio (% male) from 50% based on a binomial test. Total number of plants

examined for sex ratio estimation was 61, 110, and 76 for California, Kansas, and Texas.

Abbreviations: F: Female and M: Male.

Figure 2. Comparison of cumulative seedling emergence between female (F) and male (M)

plants in three populations *A. palmeri* originating. Inset tables show parameter estimates for the

logistic model (equation 2). Values in the parentheses indicate stand errors. Note that emergence

data were normalized relative to the final number of seedlings counted for each sex within the

population (see text for definition of model's parameters).

Figure 3. Days to first flowering (inflorescence visible; top row) and anthesis (bottom row) in

female (F) and male (M) plants of *A. palmeri* from three populations (California, Kansas, and

Texas). The width of violin plot represents the relative frequency of the event of interest (i.e.

flowering or anthesis time). Horizontal line within the boxplot shows the median, the box

includes the interquartile range, IQR, (i.e. 25^{th} to 75^{th} percentile range), whiskers extend 1.5

times the IQR, and filled circles represent extreme values which fall within $> 1.5 \times$IQR and $<$

$3 \times$IQR.

Figure 4. Length of time (day) from flowering (inflorescence visible) to anthesis in female (F)

and male (M) plants of *A. palmeri* from three populations (California, Kansas, and Texas). The

width of violin plot shows the relative frequency of event (inflorescence to anthesis time). The

horizontal line within the boxplot shows the median, the box includes the interquartile range,

IQR, (i.e. 25th to 75th percentile range), whiskers extend 1.5 times the IQR, and filled circles

represent extreme values which fall within $> 1.5 \times$IQR and $< 3 \times$IQR.

Figure 5. Compression of final plant height (top row) and relative growth rate (bottom row)

between female (F) and male (M) plants in three populations of *A. palmeri*. Vertical lines on bars

indicate 95% confidence interval.

Figure 6. Observed sex ratios in two *A. palmeri* populations grown under well-watered (control)

and water stress conditions. Also reported are the P-values (shown as P) for testing the deviation

of sex ratio (% male) from 50% based on a binomial test. Total number of plants (sum of two

experiments) examined for sex ratio estimation was ~100 plants per water treatment.

Abbreviations: F: Female and M: Male.

Figure 7. Frequency distribution of time to flowering (emergence of inflorescence) in female (F)

and male (M) plants of two *A. palmeri* populations grown under normal watering (control) or

water stress conditions. Vertical lines show the location of median time to flowering while

symbols indicate observed data. The differences in median flowering time between the two

sexes are shown with horizontal arrows.

Figure 8. Frequency distribution of time to anthesis in female (F) and male (M) plants of two *A.*

palmeri populations grown under normal watering (control) or water stress conditions. Vertical

lines show the location of median time to flowering while symbols indicate observed data. The

differences in median flowering time between the two sexes are shown with horizontal arrows.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.