Classification: BIOLOGICAL SCIENCES, Microbiology

Controlling septum thickness by a large protein ring

Michaela Wenzel[1,§,*], Ilkay N. Celik Gulsoy[1*], Yongqiang Gao[1¤], Joost Willemse[2], Mariska G. M. van Rosmalen[3], Gijs J. L. Wuite[3], Wouter H. Roos[4], Leendert W. Hamoen[1#]

[1]Bacterial Cell Biology, Swammerdam Institute for Life Sciences, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

[2]Molecular Biotechnology, Institute of Biology, Leiden University, 2333 BE, Leiden, The Netherlands

[3]Department of Physics and Astronomy and Laser Lab, Free University of Amsterdam, 1081 HV Amsterdam, The Netherlands

[4]Molecular Biophysics, Zernike Institute, University of Groningen, 9747 AG Groningen, The Netherlands

[*]These authors contributed equally to this work.

[#]Corresponding author: Leendert Hamoen, l.w.hamoen@uva.nl

[§]Current address: Division of Chemical Biology, Department of Biology and Biological Engineering, Chalmers University of Technology, 41296 Gothenburg, Sweden

[¤]Current address: Department of Microbiology and Immunobiology, Harvard Medical School, Boston MA 02115, USA

Keywords: cell division, Bacillus subtilis, FtsZ, SepF

Abstract

Gram-positive bacteria divide by forming a thick cross wall. How the thickness of this septal wall is controlled is unknown. In this type of bacteria, the key cell division protein FtsZ is anchored to the cell membrane by two proteins, FtsA and SepF. We have isolated SepF homologues from different bacterial species and found that they all polymerize into large protein rings with diameters varying from 19 to 41 nm. Importantly, these values correlated well with the thickness of their septa. To test whether ring diameter determines septal thickness, we tried to construct different SepF chimeras with the purpose to manipulate the diameter of the SepF protein ring. This was indeed possible and confirmed that the conserved core domain of SepF determines ring diameter. Importantly, when SepF chimeras with a smaller diameter were expressed in the bacterial host *Bacillus subtilis*, the thickness of its septa also became smaller. These results strongly support a model in which septal thickness is controlled by curved molecular clamps formed by SepF polymers attached to the leading edge of nascent septa. This also implies that the intrinsic shape of a protein polymer can function as a mould to shape the cell wall.

Significance Statement

Many bacteria form a thick cell wall and divide by forming a cross wall. How they control the thickness of their cell wall and cross wall is unknown. In this study we show that in these bacteria the cell division protein SepF forms very large protein rings with diameters that correspond to the diameter of their cross walls. Importantly, when we reduced the diameter of SepF rings in the bacterial host *Bacillus subtilis* the cross wall also became thinner. These results provide strong evidence that a large protein ring can function as a mould to control the thickness of the cell wall that divides these bacterial cells.

Introduction

The hallmark of Gram-positive bacteria is their thick cell wall composed of multiple layers of peptidoglycan. They divide by synthesizing a crosswall in between the newly formed daughter cells, and the thickness of the nascent division septum approaches that of the lateral cell wall. How Gram-positive bacteria regulate the thickness of their division septum is not known.

Bacterial cell division is accomplished by a complex multi-protein machinery called the divisome. Assembly of the divisome begins with polymerization of the tubulin homologue FtsZ at midcell into a ring-like configuration, the so called Z-ring (1). This structure forms a scaffold for the late cell division proteins that are responsible for synthesis of the dividing septal wall (2). Several cell division proteins support the formation of the Z-ring, and a key step is the anchoring of FtsZ polymers to the cell membrane. This is achieved by the conserved peripheral membrane proteins FtsA and SepF. Both proteins directly interact with FtsZ and use an amphipathic α-helix to bind the cell membrane (3, 4). FtsA can be found in both Gram-positive and Gram-negative bacteria, whereas SepF is widely conserved in Gram-positive and Cyanobacteria, but has no known homologue in Gram-negatives (5, 6). Other Z-ring proteins are the conserved protein ZapA, which interlinks FtsZ polymers (7), and the bitopic transmembrane proteins EzrA (Gram-positives) and ZipA (Gram-negatives) (8, 9). Once the Z-ring is assembled, the late cell division proteins arrive. These conserved transmembrane proteins form a complex comprising the peptidoglycan glycosyltransferase FtsW (10), the transpeptidase Pbp2B (FtsI in Gram-negatives) (11, 12), and the heterotrimeric complex composed of FtsL, DivIC and DivIB (FtsL, FtsB and FtsQ in Gram-negatives, respectively) (13, 14). It is assumed that the latter three proteins

regulate the assembly of late cell division proteins, although it is not yet clear how

the late proteins are recruited to the Z-ring in Gram-positive bacteria.

Some bacteria, such as the Gram-positive model organism *Bacillus subtilis*,

contain both FtsA and SepF, and this organism needs only one of them for Z-ring

formation. However, the absence of SepF results in highly deformed septa, which

is not the case when FtsA is absent (6). This indicates that SepF must have an

additional function related to septum formation. A curious property of purified *B.*

subtilis SepF is that it forms large ring structures with an inner diameter of 40 nm

(15). Based on the SepF crystal structure, these rings must encompass at least 80

to 100 SepF molecules (15). *In vitro*, these protein rings are able to bundle FtsZ

polymers into very long microtubule-like structures with SepF rings stacked

perpendicularly to the FtsZ polymers (16). However, such microtubular structures

have never been observed in bacteria, and later studies showed that the

membrane-binding amphipathic α-helix of SepF is likely located inside the ring,

which seems to rule out ring formation *in vivo* (15).

Interestingly, the inner diameter of SepF rings is about the same size as the

thickness of the septal wall (43 nm). We wondered whether this relationship is

relevant, and if so, whether SepF rings might actually control the thickness of the

septum. To examine this, we purified SepF from different Gram-positive bacteria

and found that all these proteins bind to lipid membranes and form large protein

rings, albeit with different diameters. Importantly, also in these organisms there

was a correlation between SepF ring diameter and septum thickness. To confirm

that the SepF ring diameter determines septal width, we expressed SepF chimeras

with smaller diameters in *B. subtilis*. Indeed, this reduced the thickness of septa.

These results provide strong evidence that Gram-positive bacteria regulate the

thickness of their septal wall by the strong curvature of SepF polymers at the

leading edge of nascent septa. This also implies that the intrinsic form of a protein

polymer can function as a mold that can shape a cell wall.

Results

SepF rings and tubules

Purified *B. subtilis* SepF forms large protein rings when observed with transmission electron microscopy (TEM) using negative staining with uranyl acetate (15). To confirm these findings with an independent method, we examined whether the protein can form rings under more physiological conditions using atomic force microscopy (AFM). It appeared that under these conditions purified *B. subtilis* SepF also forms rings with diameters of 41 nm (Fig. 1). Rings were 2 to 4 nm high (Fig. 1C, D), which corresponds to a single or double molecule stacking, since the crystal packing has shown that SepF polymers can be either 1.5 or 2.6 nm wide, depending on the orientation of SepF dimers (15).

Previously, SepF was purified as a maltose-binding protein (MBP) fusion followed by proteolytic cleavage and removal of MBP using ion-exchange chromatography (15, 16). We noticed that after cleavage, SepF precipitates, possibly due to the presence of calcium in the digestion buffer. After resuspension in calcium-free buffer, SepF formed stacks and tubules with diameters corresponding to that of SepF rings when observed by TEM (Fig. 1E). After ion-exchange chromatography only rings are found (Fig. 1F). These findings, together with the AFM data, show that the circular polymerization of *B. subtilis* SepF is a robust feature that may also occur *in vivo*.

SepF from other species

To determine whether ring formation is a conserved feature of SepF, we purified the protein from different organisms. *Bacillus cereus* is an important food-spoiling bacterium (17, 18) and the causative agent of rainforest anthrax (19–21) and was

chosen as a close relative of *B. subtilis*. We further selected *Clostridium perfringens*, *Mycobacterium tuberculosis*, and *Streptococcus pneumoniae,* all of which are important human pathogens. *S. pneumoniae* differs from the rest since it forms cocci instead of rods, and *M. tuberculosis* is one of the bacterial species that lack an FtsA homologue. An amino acid sequence alignment of the different SepF homologues is shown in Fig. 2A. The central domain of the protein, which contains the FtsZ binding site (15), is particularly conserved. All proteins were successfully purified as MBP fusions. *C. perfringens, M. tuberculosis,* and *S. pneumoniae* SepF precipitated after proteolytic cleavage from MBP and were collected by centrifugation, whereas *B. cereus* SepF remained largely soluble and was isolated by ion-exchange chromatography.

Membrane binding

A key property of *B. subtilis* SepF is its capacity to bind to the cell membrane, which is achieved by an N-terminal amphipathic α-helix (amino acids 1-13) (15). This N-terminal amphipathic helix is reasonably conserved in the different SepF variants (Fig. 2A, see Fig. S1 for helical wheel depictions), although the helices differ in their hydrophobicity and hydrophobic moment (Fig. S2). To confirm that the different SepF molecules are able to bind to lipid membranes, the purified proteins were mixed with liposomes. This caused a strong aggregation of liposomes, and deformed them into small vesicles, when observed with high-resolution structured illumination microscopy (SIM). One example, *S. pneumoniae* SepF, is presented in Fig. 2B, C (see Fig. S3 for all variants). Such liposome deformation is a typical characteristic of membrane-interacting amphipathic α-helices (22), and a *B. subtilis* SepF variant without the N-terminal α-helix does not possess this property (15). These results indicate that the membrane binding

activity of SepF is conserved.

Ring formation

Another characteristic of *B. subtilis* SepF is its tendency to polymerize into a curved

structure. If this feature is important for the activity of SepF, it should not be

restricted to the *B. subtilis* protein. To check this, the purified SepF homologues

were spotted on TEM grids and negatively stained with uranyl acetate. Indeed, as

shown in Fig. 2D, all SepF variants formed large rings of a remarkably constant

size. However, the average ring diameters varied between species, from ~20 up

to ~40 nm (Fig. 2F).

Correlation with septum thickness

The different ring diameters provided a first proof of the hypothesis that SepF ring

diameter determines septum thickness. To confirm this we performed TEM of the

different bacterial species and measured the thickness of nascent cell division

septa (Fig. 2E, G). Because of its high pathogenicity we used published data for

M. tuberculosis (23). As shown in Fig. 2I, there is a clear correlation between inner

ring diameters and the thickness of septa in the different species. Only *B. cereus*

did not follow this trend, with SepF rings that are clearly smaller than the width of

its septa (Fig. 2F-H).

Core domain defines ring diameter

Proteolytic trimming has shown that the conserved C-terminal domain of *B. subtilis*

SepF, spanning amino acids 57 to 151, is sufficient for the formation of rings (15).

Therefore, we assumed that this core domain determines ring diameter. To test

this, we replaced the conserved core domain of *B. subtilis* SepF with those of the

SepF proteins from the other species, maintaining the first 67 and last 13 amino

acids of *B. subtilis* SepF. These chimeras were then expressed in *E. coli*, purified,

and visualized by TEM using negative staining. All SepF chimeras were able to form

rings, and importantly, the ring diameters of these chimeras corresponded very

well to the diameters of the original SepF variants (Fig. 3A-D).

Functional chimeras

If it is true that the diameter of SepF rings determines the thickness of septa, then

expressing a SepF variant with a smaller diameter in *B. subtilis* should result in

thinner septa. It is unlikely that SepF from other species function in *B. subtilis*,

since the N- and C-termini of SepF, which are much less conserved, are crucial for

its activity (15, 16). Therefore, the chimeras provided a unique opportunity to test

this. To determine whether these chimeras are active in *B. subtilis*, they were

expressed in a Δ*sepF* strain. *B. subtilis* is one of the few species that can grow

without SepF, although this leads to strongly deformed septa (6). Since these

deformed septa are lined by the cytoplasmic membrane, this phenotype can be

easily observed using high-resolution SIM microscopy by fluorescently labelling the

cell membrane (Fig. 4A). Only the *S. pneumoniae* chimera failed to rescue the

Δ*sepF* phenotype, while the *B. cereus, C. perfringens,* and *M. tuberculosis*

chimeras were all able to restore normal septum formation in the Δ*sepF*

background (Fig. 4A). To check that the membrane deformations observed by SIM

are indeed indicative of deformed cell wall septa, we confirmed these findings by

TEM (Fig. 4B).

The inner ring diameter of the *C. perfringens* SepF chimera (24 nm) is

considerably smaller than *B. subtilis* SepF (42 nm). When this chimera was

expressed in the Δ*sepF* strain, the septum thickness decreased substantially from

43 to 28 nm (Fig. 4C, D). A similar decrease in septum thickness was observed,

when the *M. tuberculosis* SepF chimera (35 nm diameter) was expressed in the

Δ*sepF* background (Fig. 4C-E). These results strongly suggests that the SepF ring

diameter control septum thickness.

Finally, the diameter of *B. cereus* SepF is approximately 10 nm smaller than

B. subtilis SepF (Fig. 2F), and so is the diameter of the *B. cereus* SepF chimera

(Fig. 3B). However, *B. cereus* has an average septum thickness comparable to that

of *B. subtilis* (Fig. 2G). If the core domain of SepF determines septum thickness,

it is likely that expression of the *B. cereus* chimera in *B. subtilis* Δ*sepF* results in

septa with a thickness comparable to that of wild type *B. subtilis* cells. This was

indeed the case (Fig. 4C-E).

Discussion

In this study we have shown that the membrane binding and ring forming activities are conserved in SepF homologues, that the correlation between ring diameter and septum thickness can be found in different Gram-positive bacteria, and that the septum thickness of *B. subtilis* cells can be reduced by expressing SepF mutants with a smaller diameter. Together these findings strongly suggest that the diameter of curved SepF polymers define septum thickness. The positioning of SepF polymers at the division site cannot be directly observed using high-resolution cryo-electron microscopy due to the electron density of the bacterial cytoplasm and cell wall. However, our findings provide support for a SepF clamp model (Fig. 5). Since the membrane-binding amphipathic α-helix of SepF is located inside the rings, it is likely that SepF polymers do not form rings, but instead form arcs, wrapping the leading edge of nascent septa, on top of which FtsZ polymers bind and align. Since these arcs control the freedom of movement of FtsZ polymers, including the peptidoglycan synthetic apparatus that is linked to them, the diameter of SepF rings will determine the thickness of the septal wall. Many SepF arcs must line the leading edge of the nascent septum to maintain an even thickness. This could be facilitated by the propensity of SepF rings to form stacks (Fig. 1F).

Previous structural work has indicated that SepF rings are built of tightly packed dimers. These dimers are formed by two parallel oriented monomers (15), as a consequence all SepF dimers have the same orientation in a ring. However, FtsZ polymers treadmill in both directions in the Z-ring (26), and it is likely that they bind SepF rings in both polymer orientations. SepF binds to the C-terminal domain of FtsZ (27), which is connected to the main protein domain by a long

flexible linker (28). This flexibility might enable FtsZ polymers with opposite

orientations to interact with the same SepF ring.

B. cereus SepF behaved slightly differently and showed an average inner

ring diameter of 30 nm, which is smaller than its average septum thickness of 43

nm. Possibly, *B. cereus* SepF polymers are more flexible in a cellular environment,

enabling a wider diameter, or there might be other factors involved in controlling

septum diameter in this organism in addition to the physical confinement of SepF

polymers. In fact, *B. subtilis* can still make septa, albeit irregular in shape, when

SepF is absent, suggesting that other factors provide some control of FtsZ

polymers. FtsA, which also forms polymers, might be such factor.

Gram-negative bacteria contain a cell wall that consists of a single-layer of

peptidoglycan. The lipoproteins LpoA and LpoB, which reach through the

peptidoglycan layer to contact and regulate the cell membrane-anchored

peptidoglycan-synthetic enzymes, are thought to regulate peptidoglycan thickness

in these species (24). Possibly, this is why Gram-negative bacteria do not need a

SepF-like protein for septum thickness control. It should be mentioned that *E. coli*

FtsA has been shown to form ring structures *in vitro* as well (25). However, the

inner diameter of these rings is ~15 nm, which is considerably smaller than the

width of septa.

Regulation of the shape of septal walls by SepF through its intrinsic

curvature is reminiscent of the activity of MreB, which controls the rod shape of

many bacteria (29). MreB forms polymers of defined curvature that orientate and

move transversely to the long axis of cells, along the direction of greatest

membrane curvature (30, 31), thereby guiding the cell wall synthetic machinery

in a way that enforces and maintains a rod shape (32, 33). However, unlike SepF,

the intrinsic curvature of MreB filaments, which has been estimated to have a

diameter of roughly 200 nm (30), does not correlate with the diameter of the cell,

which is much larger (30). Thus, the control of septum thickness by a molecular

clamp, such as a SepF arc, is a novel concept of how protein polymers can control

the shape of growing cell walls.

Material and Methods

Strain construction

All strains used in this study are listed in Table S1, plasmids in Table S2, and primers in Table S3. Plasmids for purification of SepF variants were constructed by PCR amplification of *sepF* from the respective organisms, followed by restriction cloning into the pMalC2 plasmid (34), using the *Xba*I and *Sma*I or *Eco*RI restriction sites. pMalC2-based plasmids for purification of chimera proteins and pAPNC-213-kan-based plasmids (35) for integration of *sepF* variants into the *aprE* locus in the *B. subtilis* genome were constructed by Gibson assembly (36). pMalC2-derived purification plasmids were transformed into calcium-competent *E. coli* BL21 (DE3). pAPNC-213-kan-derived integration plasmids were transformed into *B. subtilis* 168 using a standard starvation protocol (37). Deletions of the *sepF* and *ftsA* genes were introduced by transforming the resulting *B. subtilis* strains with chromosomal DNA isolated from YK204 (*sepF::spc*) (38), BFA2863 (*sepF::ery*) (6), and TB94 (*ftsA::cat*), respectively. TB94 was created by transformation of *B. subtilis* 168 with chromosomal DNA of strain MD133 *ftsA::cat aprE::kan Pspac-ftsA* (38) followed by selection on chloramphenicol resistance but kanamycin sensitivity.

Protein purification

E. coli BL21 (DE3) strains carrying pMalC2 plasmids for purification of SepF variants were grown overnight in LB containing 100 µg/ml ampicillin at 37 °C under continuous shaking. Cultures were diluted 1:100 and allowed to grow until an OD_{600} of 0.4 in the presence of ampicillin. Expression of maltose-binding protein (MBP)-tagged SepF proteins was induced by addition of 0.5 mM IPTG. Cells were harvested by centrifugation after 4 h of induction and subsequently washed in

phosphate-buffered saline supplemented with 1 mM phenylmethylsulfonyl fluoride to prevent protein degradation. Cell pellets were flash-frozen in liquid nitrogen and stored at -80 °C until further use. Cell pellets were resuspended in buffer AF (50 mM Tris-HCL, pH 7.4, 200 mM KCl, 5 mM EDTA, 0.5 mM DTT) supplemented with one Complete mini protease inhibitor tablet (Roche) and disrupted by French Press. Cell debris was removed by ultracentrifugation at 31,000 x g, and the resulting supernatant was filtered through 0.2 μm filter membranes prior to loading onto a Tricorn 10/20 column (GE Healthcare) packed with 2 ml amylose resin (New England Biolabs) equilibrated with buffer AF. After loading, the column was washed with 5 column volumes buffer AF, followed by buffer BF (50 mM Tris-HCl, pH 7.4) until the baseline was stable. MBP-tagged SepF variants were eluted with buffer BF containing 10 mM maltose. Appropriate fractions were pooled and digested with factor Xa protease (New England Biolabs) in the presence of 2 mM $CaCl_2$ at 4 °C overnight. *B. cereus* SepF was soluble while the others precipitated after cleavage (*B. subtilis, S. pneumoniae, C. perfringens, M. tuberculosis,* all chimera proteins) due to presence of calcium. Insoluble proteins were separated from soluble MBP and factor Xa by centrifugation at 10,000 x g. Pellets containing pure SepF were dissolved in buffer BF without $CaCl_2$, flash-frozen in liquid nitrogen and stored at -80 °C until further use. SepF variants that were soluble after factor Xa cleavage were separated from MBP and Factor Xa protease by ion exchange chromatography. To this end, digested samples were loaded onto a 1 ml HiTrap Q column (GE Healthcare) equilibrated with buffer BF. The column was washed with buffer BF until the baseline was stable, followed by washing with 7.5% buffer CF (50 mM Tris-HCl, pH 7.4, 1 M KCl) resulting in elution of MBP, and washing with 17.5% buffer CF resulting in elution of factor Xa. Pure SepF was eluted with 50% buffer CF, flash-frozen in liquid nitrogen, and stored at -80 °C.

Atomic force microscopy

Purified *B. subtilis* SepF was diluted 1:10 in buffer BF and added onto a freshly cleaved highly oriented pyrolytic graphite (HOPG) surface (Agar Scientific) and allowed to settle for 20 min at room temperature prior to imaging in buffer solution (39). Images were taken with an atomic force microscope from Nanotec Electronica (Tres Cantos) operated in jumping mode at room temperature. Olympus OMCL-RC800PSA rectangular, silicon-nitride cantilevers with a nominal tip radius of 15 nm and a nominal spring constant of 0.05 N/m were used. Images were processed and analyzed with WSxM software (40).

Fluorescence light microscopy of liposomes

Liposomes were prepared from *E. coli* polar lipid extract (Avanti Polar Lipids) as described previously (3). Liposomes were extruded through 0.2 µm filters. Samples were stained with 1 µg/ml Nile red, spotted on 1.2% agarose films, covered with poly-dopamine-coated cover slips (41), and immediately imaged with a Nikon Eclipse Ti equipped with a CFI Plan Apochromat DM 100x oil objective, an Intensilight HG 130 W lamp, a C11440-22CU Hamamatsu ORCA camera, and NIS elements software. Images were analyzed using ImageJ (National Institutes of Health).

Structured illumination (SIM) microscopy

Liposomes for SIM were prepared from *E. coli* polar lipid extract (Avanti Polar Lipids) as described previously (3). Liposomes were extruded through 0.8 µm filters to obtain large enough vesicles. 0.25 mg/ml of the respective SepF variants was mixed with 2 mg/ml liposomes in SepF binding buffer. Samples were stained

with 0.5 µg/ml mitotracker green, spotted on 1.2% agarose films, covered with

poly-dopamine-coated cover slips (41), and immediately imaged with a Nikon

Eclipse Ti N-SIM E microscope setup equipped with a CFI SR Apochromat TIRF

100x oil objective (NA1.49), a LU-N3-SIM laser unit, an Orca-Flash 4.0 sCMOS

camera (Hamamatsu Photonics K.K.), and NIS elements Ar software. SIM

microscopy of bacteria was performed by staining cells with 0.5 µg/ml mitotracker

green for 1 min, spotted on a thin film of 1.2% agarose (21).

Transmission electron microscopy of proteins

Protein samples were spotted on glow-discharged 200 mesh formvar/carbon-

coated copper grids (Agar Scientific) and incubated for 1 min at room temperature.

Excess liquid was removed with paper tissue and samples were negatively stained

by adding 100 µl 2% uranyl acetate drop by drop. Excess staining solution was

removed with paper tissue and samples were allowed to air dry. Samples were

examined with a JEOL1010 at 60 kV.

Growth conditions for microscopy

B. cereus, S. pneumoniae, and *C. perfringens* were grown on tryptic soy agar

plates containing 5 % sheep blood (BioMerieux). After three days of aerobic (*B.*

cereus and *S. pneumoniae*) or anaerobic incubation (*C. perfringens*) at 37 °C,

colonies were transferred to fresh plates and incubated for another 48 h prior to

suspension in phosphate-buffered saline and preparation for electron microscopy.

B. subtilis strains were grown in Luria-Bertani (LB) broth at 37 °C under steady

agitation. The medium was supplemented with 0.1 mM IPTG to induce expression

of SepF variants, where appropriate. It is important not to use more IPTG since

SepF overproduction causes membrane deformations that obscure septa (42).

Overnight cultures were grown with appropriate antibiotic concentrations (100

μg/ml spectinomycin, 10 μg/ml chloramphenicol, 5 μg/ml kanamycin, 1 μg/ml

erythromycin), where necessary. Cells were grown until exponential phase (OD_{600}

= 0.4) prior to microscopy.

Transmission electron microscopy of bacteria

Electron microscopy of bacteria was performed according to van Wezel *at al.* (43)

(*B. cereus*, *S. pneumoniae*, *C. perfringens)* or to a novel method that uses

immobilization of bacterial cells in one plane on an agarose layer prior to fixation

and embedding (44) (*B. subtilis*). Samples were examined with a JEOL 1010 at an

electron voltage of 60 kV.

Acknowledgements

We gratefully acknowledge Zehui Zhang and Daniel Antwi-berko for help with sample preparation, Wiep Klaas Smits for help with growing strains, Marien P. Dekker and Jan R. T. van Weering for support with TEM, and Gaurav Dugar for critically reading the manuscript. Electron microscopy was performed at the VU/VUMC EM facility, supported by the Netherlands Organization for Scientific Research (NWO, middelgroot 91111009). This work was financially supported by the Netherlands Organization for Scientific Research (NWO, STW-Vici 12128 to LWH and NWO-Vidi to WHR). MW was supported by a postdoc stipend from the Amsterdam Infection and Immunity Institute.

Conflict of Interest

The authors declare that this work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author contributions

MW planned and performed experiments and wrote the paper. INCG constructed strains and performed experiments. YG constructed strains. JW and MGM performed experiments. GJLW and WHR provided tools and planned experiments. LWH designed the study and wrote the paper.

References

References

1. J. Lutkenhaus, S. Pichoff, S. Du, Bacterial cytokinesis: From Z ring to

divisome. *Cytoskeleton* **69**, 778–790 (2012).

2. P. Gamba, J. W. Veening, N. J. Saunders, L. W. Hamoen, R. A. Daniel, Two-

step assembly dynamics of the Bacillus subtilis divisome. *J. Bacteriol.* **191**,

4186–4194 (2009).

3. H. Strahl, L. W. Hamoen, Membrane potential is important for bacterial cell

division. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 12281–12286 (2010).

4. S. Pichoff, J. Lutkenhaus, Tethering the Z ring to the membrane through a

conserved membrane targeting sequence in FtsA. *Mol. Microbiol.* **55**, 1722–

1734 (2005).

5. J. Errington, R. A. Daniel, D.-J. Scheffers, Cytokinesis in bacteria. *Microbiol.*

Mol. Biol. Rev. **67**, 52–65, table of contents (2003).

6. L. W. Hamoen, J.-C. Meile, W. De Jong, P. Noirot, J. Errington, SepF, a

novel FtsZ-interacting protein required for a late step in cell division. *Mol.*

Microbiol. **59**, 989–999 (2006).

7. F. J. Gueiros-Filho, R. Losick, A widely conserved bacterial cell division

protein that promotes assembly of the tubulin-like protein FtsZ. *Genes*

Dev. **16**, 2544–2556 (2002).

8. P. A. Levin, I. G. Kurtser, A. D. Grossman, Identification and

characterization of a negative regulator of FtsZ ring formation in Bacillus

subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 9642–9647 (1999).

9. C. A. Hale, P. A. de Boer, Direct binding of FtsZ to ZipA, an essential

component of the septal ring structure that mediates cell division in E. coli.

Cell **88**, 175–185 (1997).

10.   A. Taguchi, *et al.*, FtsW is a peptidoglycan polymerase that is functional

only in complex with its cognate penicillin-binding protein. *Nat. Microbiol.*

4, 587–594 (2019).

11.   R. A. Daniel, E. J. Harry, J. Errington, Role of penicillin-binding protein PBP

2B in assembly and functioning of the division machinery of Bacillus

subtilis. *Mol. Microbiol.* **35**, 299–311 (2000).

12.   L. Wang, M. K. Khattar, W. D. Donachie, J. Lutkenhaus, FtsI and FtsW are

localized to the septum in Escherichia coli. *J. Bacteriol.* **180**, 2810–2816

(1998).

13.   R. A. Daniel, M.-F. Noirot-Gros, P. Noirot, J. Errington, Multiple interactions

between the transmembrane division proteins of Bacillus subtilis and the

role of FtsL instability in divisome assembly. *J. Bacteriol.* **188**, 7396–7404

(2006).

14.   N. Buddelmeijer, J. Beckwith, A complex of the Escherichia coli cell division

proteins FtsL, FtsB and FtsQ forms independently of its localization to the

septal region. *Mol. Microbiol.* **52**, 1315–1327 (2004).

15.   R. Duman, *et al.*, Structural and genetic analyses reveal the protein SepF

as a new membrane anchor for the Z ring. *Proc. Natl. Acad. Sci. U. S. A.*

110, E4601-10 (2013).

16.   M. E. Gündoğdu, *et al.*, Large ring polymers align FtsZ polymers for normal

septum formation. *EMBO J.* **30**, 617–626 (2011).

17.   A. Kotiranta, K. Lounatmaa, M. Haapasalo, Epidemiology and pathogenesis

of Bacillus cereus infections. *Microbes Infect.* **2**, 189–198 (2000).

18.   E. J. Bottone, Bacillus cereus, a volatile human pathogen. *Clin. Microbiol.*

Rev. **23**, 382–398 (2010).

19.   K. S. Antonation, *et al.*, Bacillus cereus Biovar Anthracis Causing Anthrax in

Sub-Saharan Africa—Chromosomal Monophyly and Broad Geographic

Distribution. *PLoS Negl. Trop. Dis.* **10**, e0004923 (2016).

20. A. R. Hoffmaster, *et al.*, Identification of anthrax toxin genes in a Bacillus

cereus associated with an illness resembling inhalation anthrax. *Proc. Natl.*

Acad. Sci. **101**, 8449–8454 (2004).

21. C. Hoffmann, *et al.*, Persistent anthrax as a major driver of wildlife

mortality in a tropical rainforest. *Nature* **548**, 82 (2017).

22. W. A. Prinz, J. E. Hinshaw, Membrane-bending proteins. *Crit. Rev.*

Biochem. Mol. Biol. **44**, 278–291 (2009).

23. A. A. Velayati, P. Farnia, *Altas of Mycobacterium tuberculosis*, 1st Editio

(Academic Press, 2016).

24. A. Typas, *et al.*, Regulation of peptidoglycan synthesis by outer-membrane

proteins. *Cell* **143**, 1097–1109 (2010).

25. M. Krupka, *et al.*, Escherichia coli FtsA forms lipid-bound minirings that

antagonize lateral interactions between FtsZ protofilaments. *Nat. Commun.*

8, 15957 (2017).

26. A. W. Bisson-Filho, *et al.*, Treadmilling by FtsZ filaments drives

peptidoglycan synthesis and bacterial cell division. *Science* **355**, 739–743

(2017).

27. E. Krol, *et al.*, Bacillus subtilis SepF binds to the C-terminus of FtsZ. *PLoS*

One **7**, e43293 (2012).

28. P. J. Buske, A. Mittal, R. V Pappu, P. A. Levin, An intrinsically disordered

linker plays a critical role in bacterial cell division. *Semin. Cell Dev. Biol.*

37, 3–10 (2015).

29. L. J. Jones, R. Carballido-Lopez, J. Errington, Control of cell shape in

bacteria: helical, actin-like filaments in Bacillus subtilis. *Cell* **104**, 913–922

(2001).

30.   S. Hussain, *et al.*, MreB filaments align along greatest principal membrane curvature to orient cell wall synthesis. *Elife* **7**, e32471 (2018).

31.   T. S. Ursell, *et al.*, Rod-like bacterial shape is maintained by feedback between cell curvature and cytoskeletal localization. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1025-34 (2014).

32.   J. Dominguez-Escobar, *et al.*, Processive movement of MreB-associated cell wall biosynthetic complexes in bacteria. *Science* **333**, 225–228 (2011).

33.   E. C. Garner, *et al.*, Coupled, circumferential motions of the cell wall synthesis machinery and MreB filaments in B. subtilis. *Science* **333**, 222–225 (2011).

34.   P. Riggs, "Expression and Purification of Maltose-Binding Protein Fusions" in *Current Protocols in Molecular Biology*, (John Wiley & Sons, Inc., 2001) https://doi.org/10.1002/0471142727.mb1606s28.

35.   M. Yoshimura, T. Oshima, N. Ogasawara, Involvement of the YneS/YgiH and PlsX proteins in phospholipid biosynthesis in both Bacillus subtilis and Escherichia coli. *BMC Microbiol.* **7**, 69 (2007).

36.   D. G. Gibson, *et al.*, Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Meth* **6**, 343–345 (2009).

37.   P. M. Hauser, D. Karamata, A rapid and simple method for Bacillus subtilis transformation on solid media. *Microbiology* **140**, 1613–1617 (1994).

38.   S. Ishikawa, Y. Kawai, K. Hiramatsu, M. Kuwano, N. Ogasawara, A new FtsZ-interacting protein, YlmF, complements the activity of FtsA during progression of cell division in Bacillus subtilis. *Mol. Microbiol.* **60**, 1364–1380 (2006).

39.   K. Heinze, *et al.*, Protein Nanocontainers from Nonviral Origin: Testing the

Mechanics of Artificial and Natural Protein Cages by AFM. *J. Phys. Chem. B* **120**, 5945–5952 (2016).

40. I. Horcas, *et al.*, WSXM: A software for scanning probe microscopy and a tool for nanotechnology. *Rev. Sci. Instrum.* **78**, 13705 (2007).

41. J. D. te Winkel, D. A. Gray, K. H. Seistrup, L. W. Hamoen, H. Strahl, Analysis of antimicrobial-triggered membrane depolarisation using voltage sensitive dyes. *Front. Cell Dev. Biol.* **4**, 29 (2016).

42. Y. Gao, M. Wenzel, M. J. Jonker, L. W. Hamoen, Free SepF interferes with recruitment of late cell division proteins. *Sci. Rep.* **7**, 16928 (2017).

43. G. P. van Wezel, *et al.*, ssgA is essential for sporulation of Streptomyces coelicolor A3(2) and affects hyphal development by stimulating septum formation. *J. Bacteriol.* **182**, 5653–5662 (2000).

44. D. Saeloh, *et al.*, The novel antibiotic rhodomyrtone traps membrane proteins in vesicles with increased fluidity. *PLoS Pathog.* **14**, e1006876 (2018).

Figure legends

Fig. 1 SepF forms rings and tubules. **(A)** Plane AFM images of three different SepF rings imaged in buffer solution. **(B)** 3D projections of the same images. **(C)** Height measurements of the individual SepF rings shown in A and B. **(D)** Height measurements of SepF rings derived from AFM data. **(E)** TEM image of SepF stack and tubules after cleavage from MBP. **(F)** TEM image of SepF rings after ion-exchange chromatography. Scale bars 100 nm.

Fig. 2 Membrane binding, ring formation and septum thickness. **(A)** Sequence alignment (Clustal Omega) of SepF homologues used in this study. Dark and light grey marks identical and similar amino acids, respectively. Arrows mark the beginning and end of the highly conserved core domain. **(B)** Fluorescence light microscopy image showing the aggregation of small liposomes (~200 nm) by purified *S. pneumoniae* SepF. Liposomes were stained with Nile red. **(C)** SIM image showing deformation of large liposomes (800 nm) by purified *S. pneumoniae* SepF. Liposomes were stained with mitotracker green. 0.25 µg/ml SepF was mixed with 2 mg/ml liposomes. Scale bars 1 µm. See Fig. S3 for the results with other SepF variants. **(D)** TEM pictures of purified SepF from the different species. Scale bar 50 nm. **(E)** Quantification of the inner SepF ring diameter. Ring diameter was consistently measured as the inner diameter of the SepF ring at its widest point. Black bars show mean with standard error of the mean. A minimum of 45 rings was measured. **(F)** TEM images of nascent cell division septa of the respective species. **(G)** Quantification of the thickness of the nascent septa of the respective organisms. Black bars show mean with standard error of the mean. A minimum of 50 septa was measured. **(H)** Comparison of inner ring diameters with nascent

septum thickness. Average +/- standard deviations is indicated. *Data for *M. tuberculosis* were taken from the literature (23). **(I)** Correlation between inner SepF ring diameter and septum thickness (R^2=0.74).

Fig. 3 The conserved core domain of SepF determines ring diameter. **(A)** TEM pictures of purified SepF chimera proteins. **(B)** Quantification of ring diameter. Black bars represent mean with standard error of the mean. A minimum of 50 rings was measured. **(C)** Comparison of the inner ring diameters of the original and chimera proteins. Average +/- standard deviation is given. **(D)** Correlation between original and chimera ring diameter (R^2=0.71).

Fig. 4 SepF ring diameter determines septum thickness. **(A)** SIM images of *B. subtilis* Δ*sepF* cells expressing different SepF chimeras. The chimeras were expressed from the IPTG-inducible *Pspac* promotor. Cells were grown in the presence of 100 μM IPTG until mid-log phase prior to membrane staining and microscopy. Lack of SepF results in severely deformed septa, which shows as highly fluorescent membrane patches due to membrane invaginations and double membranes. Scale bar 2 μm. **(B)** Representative TEM images of septa of Δ*sepF* strains expressing the different SepF chimeras. Scale bar 1 μm. **(C)** Quantification of the septum thickness of the different strains. Black bars represent mean with standard error of the mean. p values for *C. perfringens* and *M. tuberculosis* chimeras compared to *B. subtilis* are <0.0001. **(D)** Comparison and **(E)** correlation of the inner ring diameter of SepF chimeras and nascent septum thickness of *B. subtilis* Δ*sepF* expressing the chimeras (R^2=0.57).

Fig. 5 Clamp model illustrating how SepF polymers can control septal thickness.

SepF forms large rings with an average inner ring diameter (d) that corresponds

to the thickness of septa (t). Crossections of a nascent septum (grey) is depicted.

FtsZ polymers (Z) bind to the outside of SepF rings, in a perpendicular fasion. The

transmembrane peptidoglycan synthesis machinery (P) is associated with FtsZ

polymers. Of note, it is not yet know how these transmembrane proteins are linked

to FtsZ polymers in Gram-positive bacteria, and this association might be indirect.

Fig. 1.

Fig. 1 SepF forms rings and tubules. **(A)** Plane AFM images of three different SepF rings imaged in buffer solution. **(B)** 3D projections of the same images. **(C)** Height measurements of the individual SepF rings shown in A and B. **(D)** Height measurements of SepF rings derived from AFM data. **(E)** TEM image of SepF stack and tubules after cleavage from MBP. **(F)** TEM image of SepF rings after ion-exchange chromatography. Scale bars 100 nm.

Fig. 2.

A

B

control

S. pneumoniae

C

control

S. pneumoniae

D

| *B. subtilis* | *B. cereus* | *S. pneumoniae* | *C. perfringens* | *M. tuberculosis* |

E

B. subtilis

B. cereus

S. pneumoniae

C. perfringens

F

G

I

H

	B. subtilis	*B. cereus*	*S. pneumoniae*	*C. perfringens*	*M. tuberculosis*
inner ring diameter (nm)	41 +/-3	30 +/-3	34 +/-4	28 +/-7	19 +/-7
nascent septum thickness (nm)	43 +/-8	43 +/-12	37 +/-11	33+/-7	21 nm*

Fig. 2 Membrane binding, ring formation and septum thickness. **(A)** Sequence alignment (Clustal Omega) of SepF homologues used in this study. Dark and light grey marks identical and similar amino acids, respectively. Arrows mark the beginning and end of the highly conserved core domain. **(B)** Fluorescence light microscopy image showing the aggregation of small liposomes (~200 nm) by purified *S. pneumoniae* SepF. Liposomes were stained with Nile red. **(C)** SIM image showing deformation of large liposomes (800 nm) by purified *S. pneumoniae* SepF. Liposomes were stained with mitotracker green. 0.25 µg/ml SepF was mixed with

2 mg/ml liposomes. Scale bars 1 µm. See Fig. S3 for the results with other SepF
variants. **(D)** TEM pictures of purified SepF from the different species. Scale bar
50 nm. **(E)** Quantification of the inner SepF ring diameter. Ring diameter was
consistently measured as the inner diameter of the SepF ring at its widest point.
Black bars show mean with standard error of the mean. A minimum of 45 rings
was measured. **(F)** TEM images of nascent cell division septa of the respective
species. **(G)** Quantification of the thickness of the nascent septa of the respective
organisms. Black bars show mean with standard error of the mean. A minimum of
50 septa was measured. **(H)** Comparison of inner ring diameters with nascent
septum thickness. Average +/- standard deviations is indicated. *Data for *M.*
tuberculosis were taken from the literature (23). **(I)** Correlation between inner
SepF ring diameter and septum thickness (R^2=0.74).

Fig. 3.

Fig. 3 The conserved core domain of SepF determines ring diameter. **(A)** TEM pictures of purified SepF chimera proteins. **(B)** Quantification of ring diameter. Black bars represent mean with standard error of the mean. A minimum of 50 rings was measured. **(C)** Comparison of the inner ring diameters of the original and chimera proteins. Average +/- standard deviation is given. **(D)** Correlation between original and chimera ring diameter (R^2=0.71).

Fig. 4.

Fig. 4 SepF ring diameter determines septum thickness. **(A)** SIM images of *B. subtilis ΔsepF* cells expressing different SepF chimeras. The chimeras were expressed from the IPTG-inducible *Pspac* promotor. Cells were grown in the presence of 100 µM IPTG until mid-log phase prior to membrane staining and microscopy. Lack of SepF results in severely deformed septa, which shows as highly fluorescent membrane patches due to membrane invaginations and double membranes. Scale bar 2 µm. **(B)** Representative TEM images of septa of *ΔsepF* strains expressing the different SepF chimeras. Scale bar 1 µm. **(C)** Quantification of the septum thickness of the different strains. Black bars represent mean with standard error of the mean. p values for *C. perfringens* and *M. tuberculosis* chimeras compared to *B. subtilis* are <0.0001. **(D)** Comparison and **(E)** correlation of the inner ring diameter of SepF chimeras and nascent septum thickness of *B. subtilis ΔsepF* expressing the chimeras (R^2=0.57).

Fig. 5.

Fig. 5 Clamp model illustrating how SepF polymers can control septal thickness. SepF forms large rings with an average inner ring diameter (d) that corresponds to the thickness of septa (t). Crossections of a nascent septum (grey) is depicted. FtsZ polymers (Z) bind to the outside of SepF rings, in a perpendicular fasion. The transmembrane peptidoglycan synthesis machinery (P) is associated with FtsZ polymers. Of note, it is not yet know how these transmembrane proteins are linked to FtsZ polymers in Gram-positive bacteria, and this association might be indirect.

Supplementary Information

Controlling septum thickness by a large protein ring

Michaela Wenzel, Ilkay N. Celik Gulsoy, Yongqiang Gao, Joost Willemse, Mariska G. M. van Rosmalen, Gijs J. L. Wuite, Wouter H. Roos, Leendert W. Hamoen

- **Table S1** Strains used in this study

- **Table S2** Plasmids used in this study

- **Table S3** Primers used in this study

- **Fig. S1** Comparison of the amphipathic helices of SepF variants

- **Fig. S2** Hydrophobic moment plot of the putative N-terminal α-helices of SepF variants

- **Fig. S3** Liposome aggregation and deformation by SepF variants

- **References**

Table S1 Strains used in this study.

strains	genotype	reference
B. subtilis 168	*trpC2*	(1)
B. subtilis TB92	*trpC2 sepF::spc* (YK204 in 168CA)	(2)
B. subtilis TB94	*trpC2 ftsA::cat* (MD133 in 168CA)	(2)
B. subtilis LH28	*trpC2 ezrA::cat*	(3)
B. subtilis BFA2863	*trpC2 sepF::ery*	(4)
B. subtilis GYQ457	*trpC2 aprE::kan Pspac-sepF(subtilis-cereus)*	this study
B. subtilis GYQ475	*trpC2 aprE::kan Pspac-sepF(subtilis-pneumoniae)*	this study
B. subtilis GYQ544	*trpC2 aprE::kan Pspac-sepF(subtilis-perfringens)*	this study
B. subtilis GYQ474	*trpC2 aprE::kan Pspac-sepF(subtilis-tuberculosis)*	this study
B. subtilis MW18	*trpC2 aprE::kan Pspac-sepF(WT) sepF::spc*	this study
B. subtilis GYQ458	*trpC2 aprE::kan Pspac-sepF(subtilis-cereus)*	this study
B. subtilis MW89	*trpC2 aprE::kan Pspac-sepF(subtilis-pneumoniae)*	this study
B. subtilis MW92	*trpC2 aprE::kan Pspac-sepF(subtilis-perfringens)*	this study
B. subtilis MW86	*trpC2 aprE::kan Pspac-sepF(subtilis-tuberculosis)*	this study
E. coli BL21(DE03) pNC1	*pMalC2-sepF(coelicolor3)*	this study
E. coli BL21(DE03) pNC2	*pMalC2-sepF(cereus)*	this study
E. coli BL21(DE03) pNC4	*pMalC2-sepF(tuberculosis)*	this study
E. coli BL21(DE03) pNC5	*pMalC2sepF(coelicolor2)*	this study
E. coli BL21(DE03) pNC6	*pMalC2-sepF(coelicolor1)*	this study
E. coli BL21(DE03) pNC7	*pMalC2-sepF(perfringens)*	this study
E. coli BL21(DE03) pNC9	*pMalC2-sepF(pneumoniae)*	this study
E. coli BL21(DE03) pNC12	*pMalC2-sepF(subtilis)*	(5)
E. coli BL21(DE03) pYQ62	*pMalC2-sepF(subtilis-cereus)*	this study
E. coli BL21(DE03) pYQ94	*pMalC2-sepF(subtilis-tuberculosis)*	this study
E. coli BL21(DE03) pYQ95	*pMalC2-sepF(subtilis-perfringens)*	this study
E. coli BL21(DE03) pYQ96	*pMalC2-sepF(subtilis-pneumoniae)*	this study
B. cereus ATCC14579		(6)
S. pneumoniae ATCC6305		(7, 8)
C. perfringens ATCC13124		(9)

Table S2 Plasmids used in this study.

plasmid	genotype	reference
pMW8	*pAPNC-213-kan-Pspac-sepF(WT)*	this study
pYQ90	*pAPNC-213-kan-Pspac-sepF(subtilis-cereus)*	this study
pYQ97	*pAPNC-213-kan-Pspac-sepF(subtilis-tuberculosis)*	this study
pYQ98	*pAPNC-213-kan-Pspac-sepF(subtilis-perfringens)*	this study
pYQ99	*pAPNC-213-kan-Pspac-sepF(subtilis-pneumoniae)*	this study
pNC1	*pMalC2-sepF(coelicolor3)*	this study
pNC2	*pMalC2-sepF(cereus)*	this study
pNC4	*pMalC2-sepF(tuberculosis)*	this study
pNC5	*pMalC2sepF(coelicolor2)*	this study
pNC6	*pMalC2-sepF(coelicolor1)*	this study
pNC7	*pMalC2-sepF(perfringens)*	this study
pNC9	*pMalC2-sepF(pneumoniae)*	this study
pNC12	*pMalC2-sepF(subtilis)*	(5)
pYQ62	*pMalC2-sepF(subtilis-cereus)*	this study
pYQ94	*pMalC2-sepF(subtilis-tuberculosis)*	this study
pYQ95	*pMalC2-sepF(subtilis-perfringens)*	this study
pYQ96	*pMalC2-sepF(subtilis-pneumoniae)*	this study

Table S3 Primers used in this study.

primer	description	sequence	used for
MW135	pAPNC-213-kan-GA-for	GGCGTTAGCCCAAGCGC	pMW8
MW139	pAPNC-213-kan-GA-rev	ACACCCCCTGTTTCATTTCCCTAGCAGGTCAATTGTGAGCGC	pMW8
MW140	*sepF*-WT-GA-for	GGAAATGAAACAGGGGGTGTACAGCAATGAGTATGAAAAATAAACTGAAAAACTTTTTCTCAATGGAAGATGAAG	pMW8
MW141	*sepF*-WT-GA-rev	GCGCTTGGGCTAACGCCTTACCACCTCTGATGTTCGTCTTCAGATATG	pMW8
eg122	*sepF-cereus*-for	GACGAATTCATGAGTTGGTCAAAAG	pNC2
eg123	*sepF-cereus*-rev	GACTCTAGATTACCACCTCTTTAT	pNC2
inc15	*sepF-pneumoniae*-for	GTAGGAGCCCGGGATGTCTTTAAAAGATAG	pNC9
inc16	*sepF-pneumoniae*-rev	GCTAGACTCTAGATTATCGTACTCTATTTC	pNC9
inc9	*sepF-perfringens*-for	GATTAGCCCGGGATGTGTATGTCAAAAG	pNC7
inc10	*sepF-perfringens*-rev	GACTATCTAGATTATTTTGAAGCCCAGTTG	pNC7
inc7	*sepF-tuberculosis*-for	GGTTACGGAATTCGTGAATAGTCACTGTAG	pNC4
inc8	*sepF-tuberculosis*-rev	GGCGACCGTCTAGACTATTGGTAGGCGTAG	pNC4
inc17	*sepF-coelicolor* 1-for	GTGAGAGGAGGAATTCATGGGATCGGTAC	pNC6
inc18	*sepF-coelicolor* 1-rev	TGTGCGGCTCTAGATCAGCTCTGGTTGAAG	pNC6
inc19	*sepF-coelicolor* 2-for	GAGGACTCCCGGGATGGCCGGCGCGATG	pNC5
inc20	*sepF-coelicolor* 2-rev	ACCGGTAGTCTAGATCAGCTCTGGTTGAAG	pNC5
eg139	*sepF-coelicolor* 3-for	GACGAATTCGTGAAATCGGGGGAGC	pNC1
eg140	*sepF-coelicolor* 3-rev	GACTCTAGATCACACTCCCGGCAC	pNC1
YQ276	*sepF-cereus*-for	AATCCTCTAAAGTGGTGTTGTTAGAACCACGCACATATTCGGA	pYQ62, 90
YQ277	*sepF-cereus*-rev	CGCACCAACAATATCTACATTTTC	pYQ62, 90
YQ278	pMalC2-for	ATGTAGATATTGTTGGTGCGATTTCTGAGCTCATATCTGAAGAC	pYQ62, 90
YQ279	pMalC2-rev	CAACACCACTTTAGAGGATTTCTG	pYQ62, 90
YQ349	pMalC2-rev	TTTCTGAACACTTTGCAAGCTCAC	pYQ94-99
YQ350	*sepF-tuberculosis*-for	GAGCTCATATCTGAAGACGAACATC	pYQ94, 97
YQ351	pMalC2-for	GCTTGCAAAGTGTTCAGAAACTCTCGAAGATCACCACGCT	pYQ94-99
YQ352	*sepF-tuberculosis*-rev	TCGTCTTCAGATATGAGCTCGCGCTCCTCGGGGGACACATC	pYQ94, 97
YQ353	*sepF-perfringens*-	GCTTGCAAAGTGTTCAGAAACAGCTAAATTA	pYQ95, 98

		ATGATAACTAAACCA	
YQ354	*sepF-perfringens-* rev	TCGTCTTCAGATATGAGCTCCTTTAATTCATTG CTTACTTCAAC	pYQ95, 98
YQ355	*sepF-pneumoniae-* for	GCTTGCAAAGTGTTCAGAAAAAGGTCATTATA GATGTTCGTTATC	pYQ96, 99
YQ356	*sepF-pneumoniae-* rev	TCGTCTTCAGATATGAGCTCGATATCTTCAAC ATTTACAATAACATTC	pYQ96, 99

Fig. S1 Comparison of the amphipathic helices of SepF variants. Helical wheel projections (RZ lab) show the first 13 amino acids of the different proteins. Hydrophilic residues are represented as circles, hydrophobic residues as diamonds, negatively charged as triangles, and positively charged as pentagons. Hydrophobicity is color coded with green indicating the highest and yellow the lowest (zero) hydrophobicity. Hydrophilic residues are depicted in red with the amount of red decreasing to orange proportionally to the hydrophilicity. Potentially charged residues are depicted in light blue.

Fig. S2 Hydrophobic moment plot of the putative N-terminal α-helices of SepF variants. High H values indicate strong hydrophobicity. The hydrophobic moment (µH) is a measure for the amphipathicity of an α-helix (high values indicating strong amphipathicity). The plot of H and µH indicates the probability of an α-helix to reside in the globular or surface-exposed domains of a protein and can be used to predict transmembrane or amphipathic membrane-binding helices. Black dots indicate known globular (*Bacillus thermoproteolyticus* thermolysin, aa 67-88), transmembrane (*B. subtilis* EzrA, aa 1-21) and amphipathic (*B. subtilis* FtsA, aa 424-440) helices for comparison. N-terminal α-helices of SepF homologues are depicted in red.

Fig. S3 Liposome aggregation and deformation by SepF variants. **(A)** Fluorescence light microscopy images of small liposomes (200 nm) stained with Nile red. **(B)** SIM images of large liposomes (800 nm) stained with mitotracker green. 0.25 µg/ml of the different SepF proteins (species names indicated) were mixed with 2 mg/ml liposomes prepared from *E. coli* polar lipid extract. Scale bars 1 µm.

References

1. Anagnostopoulos C, Spizizen J (1960) Requirements for transformation in Bacillus subtilis. *J Bacteriol* 81(5):741–746.
2. Ishikawa S, Kawai Y, Hiramatsu K, Kuwano M, Ogasawara N (2006) A new FtsZ-interacting protein, YlmF, complements the activity of FtsA during progression of cell division in Bacillus subtilis. *Mol Microbiol* 60(6):1364–1380.
3. Gamba P, Rietkotter E, Daniel RA, Hamoen LW (2015) Tetracycline hypersensitivity of an ezrA mutant links GalE and TseB (YpmB) to cell division. *Front Microbiol* 6:346.
4. Hamoen LW, Meile J-C, De Jong W, Noirot P, Errington J (2006) SepF, a novel FtsZ-interacting protein required for a late step in cell division. *Mol Microbiol* 59(3):989–999.
5. Duman R, et al. (2013) Structural and genetic analyses reveal the protein SepF as a new membrane anchor for the Z ring. *Proc Natl Acad Sci U S A* 110(48):E4601-10.
6. Frankland GC, Frankland PF (1887) XI. Studies on some new micro-organisms obtained from air. *Philos Trans R Soc London* 178:257–287.
7. Public Health Weekly Reports for APRIL 14, 1944 (1944) *Public Heal reports (Washington, DC 1896)* 59(15):485–512.
8. Public Health Weekly Reports for APRIL 7, 1944 (1944) *Public Heal reports (Washington, DC 1896)* 59(14):449–484.
9. Eastoe JE, Long JE (1959) The effect of nisin on the growth of cels and spores of Clostridium welchii in gelatine. *J Appl Bacteriol* 22(1):1–7.