Spatial transcriptional signatures define margin morphogenesis along the proximal-distal and medio-lateral axes in the complex leaf of tomato (*Solanum lycopersicum*)

Ciera C. Martinez[1,2,3], Siyu Li[3], Margaret R. Woodhouse[3], Keiko Sugimoto[4], and Neelima R. Sinha[3]*

[1]Department of Molecular and Cellular Biology, University of California at Berkeley, Berkeley, CA 94709, USA

[2]Berkeley Institute of Data Science, University of California at Berkeley, Berkeley, CA 94709, USA

[3]Department of Plant Biology, University of California at Davis, Davis, CA 95616, USA

[4]RIKEN Center for Sustainable Resource Science, Tsurumi, Yokohama, 15 230-0045 Japan

*Author for correspondence nrsinha@ucdavis.edu

Short title: Transcriptional signatures during leaf development

One-sentence summary: Rigorous structural characterization, laser capture microdissection, and transcriptomic sequencing reveal how gene expression patterns regulate early morphogenesis of the compound tomato leaf.

ABSTRACT

Leaf morphogenesis involves cell division, expansion, and differentiation in the

developing leaf, cells at different positions along the medio-lateral and proximal-distal

leaf axes divide, expand, and differentiate at different rates. The gene expression

changes that control cell fate along these axes remain elusive due to difficulties in

precisely isolating tissues. Here, we combined rigorous early leaf characterization, laser

capture microdissection, and transcriptomic sequencing to ask how gene expression

patterns regulate early leaf morphogenesis in wild-type tomato (*Solanum lycopersicum*)

and the leaf morphogenesis mutant *trifoliate*. We observed transcriptional regulation of

cell differentiation along the proximal-distal axis and identified molecular signatures

delineating the classically defined marginal meristem/blastozone region during early leaf

development. We describe the importance of endoreduplication during leaf

development, when and where leaf cells first achieve photosynthetic competency, and

the regulation of auxin transport and signaling along the leaf axes. Knockout mutants of

BLADE-ON-PETIOLE2 exhibited ectopic shoot apical meristem formation on leaves,

highlighting the role of this gene in regulating margin tissue identity. We mapped gene

expression signatures in specific leaf domains and evaluated the role of each domain in

conferring indeterminacy and permitting blade outgrowth. Finally, we generated a global

gene expression atlas of the early developing compound leaf.

INTRODUCTION

A major theme in plant development is the reiteration of patterning events, which are

influenced by the identity and relative arrangement of neighboring plant parts. The

phytomer concept describes reiterated units of the leaf, stem, and axillary bud that

make up the aboveground shoot (Sussex and Kerk, 2001). Molecular analyses

comparing development in various plant species suggest that the reiteration of

developmental patterning in plants is defined by the recruitment of a common molecular

toolbox and the dizzying array of leaf architecture found in plants results from variations

on a common genetic regulatory program (Tsukaya, 2014; Bendahmane and Theres,

2011; Blein et al., 2008).

The shoot apical meristem (SAM), which is located at the growing tip of the shoot, is a

dome-like structure containing reservoirs of continually self-renewing stem cells and is

characterized by spatially defined zones. The peripheral zone of the SAM gives rise to

most lateral organs, including leaves. Like the SAM, the angiosperm leaf has been

historically defined in terms of zones and spatial cell organization. Leaf development

begins with periclinal cell divisions on the periphery of the SAM and continues as cells

proceed through the specific steps of development beginning with cell division, followed

by cell expansion and cell specialization. In many instances, this specialization involves

endoreduplication. The timing of these stages varies depending on the cell position on

the leaf primordium.

Leaf morphogenesis and patterning occur along three main axes: the abaxial-adaxial,

proximal-distal, and medio-lateral axes. Many studies have focused on the importance

of the abaxial-adaxial boundary in establishing leaf polarity (Eshed et al., 2001; Moon

and Hake, 2011; Kidner and Timmermans, 2007), but less is known about the proximal-

distal and medio-lateral axes of the leaf. During the development of most eudicot

leaves, cells differentiate more rapidly in the distal (top) region than in the proximal

(base) region. Along the medio-lateral axis, the differentiation at the margin of a leaf is

decelerated relative to the more medial regions (midvein, rachis, petiole). Thus,

historically, the leaf margin is of particular interest because it maintains cellular

pluripotency longer than the other regions and has even been described as a

meristematic region termed the marginal meristem (Poethig and Sussex, 1985b; Avery,

1933) or marginal blastozone (Hagemann and Gleissberg, 1996).

Although the developmental fate, homology, and even the name of the margin region of

a leaf have been debated for roughly 100 years, there is general agreement that the

process of cell differentiation in this region largely determines final leaf shape (Ori et al.,

2007; Efroni et al., 2008; Scarpella and Helariutta, 2010). The regulation and

modulation of the margin identity of a leaf are responsible for blade expansion,

serrations, lobing, vascular patterning, and new organ initiation, as in the case of leaflet

initiation in compound leaves (Scarpella et al., 2010; Bilsborough et al., 2011).

The genetic regulation and coordination of leaf morphogenesis involve distinct changes

in gene expression, as revealed by leaf transcriptomic studies in spatially defined

regions across the proximal-distal axes of the simple-leaved plant *Arabidopsis thaliana*

(*A. thaliana*) (Beemster et al., 2005; Andriankaja et al., 2012; Efroni et al., 2008). These

studies revealed the role of endoreduplication (DNA replication without cell division) in

the acquisition of leaf morphogenic potential (Beemster et al., 2005; Andriankaja et al.,

2012; Efroni et al., 2008). The transcriptional mapping of gene expression changes in *A.*

thaliana (Beemster et al., 2005; Efroni et al., 2008; Andriankaja et al., 2012), *Solanum*

lycopersicum (tomato) (Ichihashi et al., 2014), and *Zea mays* (maize) (Li et al., 2010)

shed light on how patterning by cellular differentiation along the proximal distal axis is

established. However, this information has not yet been precisely mapped at the

transcriptome level with sufficient spatial resolution to define margin and

midvein/rachis/petiole transcriptional identity.

Interestingly, the tomato mutant *trifoliate* (*tf-2*) loses morphogenetic competence during

early leaf development and only produces three leaflets: a terminal leaflet and two

lateral leaflets subtended by a long petiole (Robinson and Rick, 1954; Naz et al., 2013).

The *tf-2* phenotype is caused by a nucleotide deletion resulting in a frameshift in the

translated amino acid sequence of an R2R3 MYB transcription factor gene

(Solyc05g007870) (Naz et al., 2013). Histological and scanning electron microscopy

(SEM) analyses of the *tf-2* mutant revealed that the marginal blastozone region is

narrower and has fewer cells, a three-fold increase in epidermal cell size, and faster cell

differentiation than the wild type (Naz et al., 2013). While the application of auxin to the

margins of wild-type *S. lycopersicum* leaf primordia causes leaflet initiation (Koenig et
al., 2009; Naz et al., 2013), in *tf-2,* the margin is unable to generate leaflets in response
to exogenous auxin , indicating that it lacks organogenic competency during early
development (Naz et al., 2013). Understanding why this mutant is incapable of initiating
more than two lateral leaflets, while wild-type leaves continue to generate an average of
ten leaflets at maturity (Naz et al., 2013), could help reveal the mechanisms regulating
margin maintenance and identity during complex leaf development.
129
Here, we used the complex tomato leaf as a model system to study the transcriptional
mechanisms directing spatial cell differentiation processes during a key developmental
stage in a young leaf, including the establishment of margin identity, proximal-distal
patterning, and leaflet initiation. Since a leaf primordium develops at varying rates in a
spatially defined manner, different developmental stages can be observed at the same
time in a single leaf (Hagemann and Gleissberg, 1996; Ori et al., 2007). We
anatomically characterized the earliest developmental stages in tomato and identified
leaf age P4 as the stage at which the medio-lateral and proximal-distal axes are first
identifiable while also containing multiple stages of leaflet organogenesis. We also
characterized the role of endoreduplication in tomato leaf morphogenesis. To map the
spatial transcriptional regulation of the P4 leaf using Laser Capture Microdissection
(LCM), we isolated six highly specific tissues previously unattainable during early
tomato leaf development and performed RNA-seq analysis to identify gene expression
changes that accompany the establishment of spatial cell differentiation patterning
during leaf organogenesis. We also included *tf-2* in our analysis, as *tf-2* lines have early
loss of morphogenetic potential in the leaf margin, thus helping us uncover a cluster of
genes whose expression differs only in regions that define organogenetic capacity in the
margin at the P4 stage. We further validated our results through molecular visualization,
providing the first evidence for when and where photosynthesis begins in a leaf. We
also used CRISPR knockout lines to explore the role of BLADE-ON-PETIOLE2 (BOP2)
(Solyc10g079460) in margin development. Our approach allowed us to predict multiple
gene expression differences that help explain the molecular identity of the classical

described, but never transcriptionally defined, marginal meristem/blastozone region and

built the first global transcriptome atlas of an early developing compound leaf.

RESULTS

Characterization of the P4 stage of tomato leaf development

The goal of this work was to characterize gene expression changes that occur during

tomato leaf morphogenesis. To define the scope of this our work, we focused on the

medio-lateral axis in an attempt to identify how the marginal blastozone maintains the

potential for leaflet organogenesis and the regulation of cell fate identity. We chose to

use the leaf stage primordium 4 (P4), the fourth oldest leaf emerging from the apical

meristem (**Figure 1A and B**). P4 provides a comprehensive snapshot of tomato leaflet

development, as it contains three distinct stages of leaflet development. During this

stage, the most distal region (destined to become the terminal leaflet) is undergoing

early blade expansion, while the most proximal region is undergoing lateral leaflet

initiation, and central to these positions is the recently initiated lateral leaflets. All three

regions can be anatomically defined, allowing the boundaries along both the medio-

lateral and proximal-distal axes to be clearly delineated. With our scope defined, we

began our analysis with a systematic survey of tissue differentiation patterns of the P4

leaf using a combination of scanning electron microscopy (SEM) and histological

approaches to establish the cellular context for detailed tissue-specific gene expression

analysis.

We defined three distinct regions of the P4 leaf along the proximal-distal axis, which are

referred to as the top, middle, and base hereafter (**Figure 1C-G**). These three regions

can further be divided into two distinct tissues types that define the medio-lateral axis:

the margin and the midrib/midvein/rachis, hereafter termed the rachis for brevity (**Figure**

1C,G, and F). The most distal region, the top, will ultimately become the terminal leaflet

of the mature leaf (**Figure 1C**). In P4 leaves, the top margin region has already begun

to develop lamina tissue (blade) and has not yet developed any tertiary vasculature, but

the future midvein in the top contains vascular cells including xylem and phloem (**Figure**

1C). The middle margin tissue has initiated the first lateral leaflets (henceforth called

LL1), the first leaflets to form from the marginal blastozone, and the rachis tissue

displays clear vascular bundles and more than four layers of cortex cells (**Figure 1F**).

The most proximal area is the base, where rachis tissue has established vascular

bundles (**Figure 1G**). Cells in the margins of all three regions along the proximal-distal

axis are small and non-vacuolated and have likely undergone little elongation, a

characteristic of marginal blastozone tissue (Hagemann and Gleissberg, 1996) (**Figure**

1C, F, and G). Tomato leaflets initiate in pairs proximal to previous leaflet initiation sites,

and therefore, the next leaflets to arise, Lateral Leaflets 2 (LL2), will occur at the base

margin region of a P4 leaf (**Figure 1D**).

To further delineate margin identity, we also characterized *tf-2*, a tomato mutant unable

to initiate leaflets past LL1, to compare its margin identity and marginal organogenesis

capacity with the wild type (**Figure 1E and J**). The developmental fate of the *tf-2* mutant

diverges from that of the wild type at P4, as the margin is unable to form leaflets after

LL1. Therefore, comparing *tf-2* and wild type allowed us to explore two leaves of

comparable developmental age but with different organogenic potential, i.e., different

abilities to form leaflets. The anatomical characterization of wild type and *tf-2* revealed

precise cell types present across a P4 leaf, serving as a proxy for defining cell

differentiation.

Cell division and endoreduplication in the P4 leaf

Previous transcriptomic studies tracing proximal-distal cell division patterning and

cellular processing indicated that changes in gene expression are responsible for the

regulation of cell division, cell elongation, and endoreduplication during differentiation in

developing *A. thaliana* leaves (Beemster et al., 2005; Efroni et al., 2008; Andriankaja et

al., 2012; Donnelly et al., 1999). Endoreduplication is thought to be a defining

component of *A. thaliana* leaf morphogenesis (Beemster et al., 2005; Gutierrez, 2005),

with ploidy levels varying from 2C to 32C (Melaragno et al., 1993; Beemster et al.,

2005). Endoreduplication occurs at the onset of leaf differentiation, and elongation

occurs after cell proliferation, when cell ploidy levels increase due to successive rounds

of DNA replication, often resulting in increased cell size (Kondorosi et al., 2000;

Sugimoto-Shirasu and Roberts, 2003; De Veylder et al., 2011). While endoreduplication

occurs at rates (256C to 512C) during tomato fruit development (Bergervoet et al.,

1996; Joubès et al., 2000; Cheniclet et al., 2005; Bourdon et al., 2010), where and to

what extent endoreduplication occurs during tomato leaf development are currently

unknown. Since no reports were available on endoreduplication during tomato leaf

development, we carefully characterized cell division and endoreduplication processes

at the P4 stage to identify similarities and differences between early leaf development in

tomato and *Arabidopsis*.

To observe where cell division is occurring throughout the P4 leaf, we used 5-ethynyl-

29-deoxy-uridine (EdU), which is incorporated during the S phase of the cell cycle and

serves as a proxy to map cell division locations. Along the mediolateral axis, in wild type

and to a lesser extent in *tf-2,* EdU fluorescence was more prominent in the margin

compared to rachis tissue (**Figure 2E-F**), indicating that the margin tissue is actively

undergoing cell division, as expected for marginal blastozone tissue. At the base margin

region of wild type, where Lateral Leaflet 2 (LL2) will arise, EdU was incorporated in a

cluster (**Figure 2E**), clearly demonstrating early cell division processes during LL2

initiation. Therefore, during early P4 development, LL2 initiation has already begun,

although this is not always obvious based on external views of the leaf (**Figure 1B and**

D). The *tf-2* mutant did not show clustering of EdU fluorescence in the base margin

(**Figure 2E and F**), revealing that the cell divisions needed for LL2 initiation have not

occurred. In conclusion, cell division across the mediolateral axis in wild type and *tf-2*

reflects similar processes that occur in *A. thaliana* (Donnelly et al., 1999), where cells

are actively dividing in the margin. The cell divisions needed for LL2 initiation at P4 have

already begun in the wild type but not in *tf-2*. Therefore, the mechanism that restricts

LL2 initiation in *tf-2* is likely in place at the P4 stage of development.

We used flow cytometry to measure DNA contents in tissues from the terminal leaflets

of leaves across several developmental ages. We grew all plants at Day 0 and sampled

the oldest leaf on the plant at each timepoint. Due to the limited availability of tissue

from the youngest leaf, we performed flow cytometry of whole terminal leaflet tissue

beginning at 8 days old (P6 stage leaf). We detected a combination of 2C and 4C nuclei

at all stages examined (**Figure 2G**). The 4C nuclei were likely G2 nuclei observed

following DNA replication and did not reflect the endoreduplication process, although a

few 8C nuclei were present at 30 and 60 days, perhaps representing cell-type-specific

endocycling (**Figure 2G**). In *A. thaliana* plants, there is a difference in ploidy level

between tip and base cells (Skirycz et al., 2011), but this was not the case in our

analysis (**Figure 2H**). We conclude that endoreduplication is not as pronounced in

tomato as in *Arabidopsis* and is likely not a vital aspect of tomato leaf morphogenesis,

illustrating the diversity of cellular processes in leaf morphogenetic strategies between

species.

Laser capture of six regions of the P4 tomato leaf

Since the P4 leaf is representative of two key developmental processes that define leaf

development, i.e., margin vs. rachis specification and leaflet initiation and

morphogenesis, we analyzed the P4 stage more carefully. We took advantage of our

comprehensive anatomical characterizations to generate a map delineating the medio-

lateral axis and leaflet organogenesis. We employed Laser Capture Microdissection

(LCM) following explicit rules for tissue collection (**S1 Figure and Movie 1**) on P4

leaves of both wild type and *tf-2* lines to capture gene expression differences that might

explain the morphogenetic differences in the margins of *tf-2* plants. Specifically, we

sectioned tomato apices transversely to isolate the same six sub-regions in both wild

type and *tf-2*, including the (1) top margin blastozone region (top margin), (2) top rachis,

(3) middle margin, (4) middle rachis, (5) base margin, and (6) base rachis (**Figure 1C-**

G, Movie 1). We attempted to collect enough tissue for seven replicates per sample, but

due to the fragility of RNA at such a small tissue size, a few replicates did not pass

quality control and were lost during various steps in the pipeline, resulting in a total of 3–

6 biological replicates per region. We collected tissue from 6–8 apices per biological

replicate to obtain a minimum of 2 ng of RNA per replicate. The number of cuts needed

to achieve minimum RNA levels varied depending on sample and tissue density, and

the total tissue area collected also varied among samples (**S2 Figure B and D**)**.** The

isolated mRNA from the collected tissues was further amplified and prepared for

Illumina sequencing (see Methods). Each replicate resulted in an average of 4.9 million

sequencing reads (**S2 Figure A and C**). To assess the overall similarity between

samples, we visualized gene expression using Multidimensional Scaling (MDS) for each

of the six subregions per genotype. Tissue types in each genotype generally clustered

together in multidimensional space (**S2 Figure A**) and we observed an additional

separation of margin and rachis tissue regions (**S2 Figure C**).

Differential gene expression between wild-type margin and rachis tissue along

the proximal-distal axis reveals signatures of morphogenetic states during early

leaf development

Identifying genes that are differentially regulated in margin versus rachis tissue in each

region would shed light on gene expression patterning along the medio-lateral axis. To

explore the differences between margin and rachis tissue in the three regions along the

proximal-distal axis, we performed pairwise differential gene expression analysis of wild-

type samples comparing the margin and rachis in each region (top, middle, base)

separately (**Dataset S1**) using edgeR (Robinson et al., 2010). We then performed Gene

Ontology (GO) enrichment analysis of the significantly up-regulated genes in these

samples (BH-adjusted p value < 0.05) (**Dataset S2**). More upregulated genes in the

margin region, which has historically been considered to proceed at a slower rate

through the morphogenetic stages, were enriched in GO terms associated with cell

processes that occur during early morphogenesis compared to rachis tissue. For

example, when we looked for differentially expressed genes between the margin and

rachis tissue in the top region, we identified 603 genes that were up-regulated in the

margin (**Figure S3 A**). These genes were enriched in GO terms that likely reflect cell

division processes, including chromatin and DNA processing (**Figure 3A, Dataset S2**).

Conversely, up-regulated genes in the rachis were enriched in GO terms reflecting the

cell specialization stage of morphogenesis, including transport, photosynthesis, sugar

biosynthesis, and carbohydrate metabolism (**Figure 3A-C, Dataset S2**).

A comparison of genes expressed in the margin and rachis in the most proximal region

(the base) revealed 1722 genes that were up-regulated in the rachis, which were

enriched for GO terms related to cell division, chromatin assembly, and DNA

processing. By contrast, only 94 differentially expressed genes were down-regulated

genes in the margin region vs. rachis at the base (**S3 Figure A**). These downregulated

genes were enriched for GO terms related to transcription factor activity and auxin influx

(**Figure 3A, Dataset S2**). The types of genes that were differentially expressed between

the margin and rachis also appeared to reflect that stage of morphogenesis of each

region and perhaps the distal-to-proximal wave of differentiation (**Figure 3B and C**).

The up-regulated genes in the top and middle margin regions were enriched in GO

terms describing active RNA, DNA, and chromatin processing, whereas those in the

base were enriched in GO terms similar to those in the rachis. The active processing of

RNA, DNA, and chromatin are key gene expression signatures of cell division and

expansion. The base region of the P4 leaf is still in these middle stages of

morphogenesis and just beginning to start secondary cell wall biosynthesis and to

become specialized for sucrose transport activity (**Figure 3A, Dataset S2**).

Taken together, the enriched GO terms of the differentially expressed genes describe

different stages of morphogenesis, pointing to two trajectories of development along the

leaf: development along the proximal-distal axis and the medio-lateral axis (**Figure 3B**

and C). Cells that have achieved specialized photosynthetic functions, leaf

development, and sugar transport define the final morphogenetic stages. Margin

regions undergoing active cell division are defined by chromatin assembly and DNA

processing (replication, integration, and recombination), which are required for proper

cell cycle progression, while the most highly meristematic tissue in the margin region at

the base is defined by only transcriptional activity and transcription factor and DNA

binding (**Figure 3A-C**). Thus, the P4 tomato leaf represents a complex mixture of

developmentally distinct regions that cannot be defined solely along the proximal-distal

or medio-lateral axes.

Modeling gene expression differences across the medio-lateral axis predicts that

photosynthetic activity first occurs in the rachis

Differential gene expression analysis in each region along the proximal-distal axis

revealed specific genes and enriched GO terms that are unique to the top, middle, or

base of the leaf. Next, we tried to identify gene activity that defines rachis and margin

identity across the entire P4 leaf primordium regardless of position on the longitudinal

axis. To address this issue, we performed differential gene expression analysis across

the margin and rachis tissue and adjusted for variability between the proximal-distal axis

by employing an additive linear model using the top, middle, and base identities as a

blocking factor in our experimental design using EdgeR (Robinson et al., 2010). In the

wild type, across the entire proximal-distal axis, 1,089 genes were significantly up-

regulated in the rachis and 188 genes were significantly up-regulated in the margin

(**Figure 4A, Dataset S3**). GO enrichment analysis of these upregulated genes revealed

24 enriched GO terms in the rachis (**Dataset S4**). These terms were categorized into

eight major categories: Sugar biosynthesis and transport, Metabolism (carbohydrate

and glucose), Transmembrane transport, Leaf development, Photosynthesis/light

harvesting, Catalytic activity, Cell wall organization, and Response to light (**Figure 4B**).

These categories reflect the activities of genes that were up-regulated in the rachis

compared to the margin across the entire proximal-distal axis. These results suggest

that the rachis region of a P4 leaf has many specialized tissue types and may already

be physiologically active.

Verifying photosynthetic gene expression patterns

Of the gene expression patterns described above, the most prominent pattern revealed

by both pairwise and modeled differential expression analyses was the persistent

presence of genes associated with GO terms related to photosynthetic processes; these

genes were up-regulated in the rachis compared to margin tissues (**Figure 3 and 4**).

While the up-regulation of genes involved in cell wall development, leaf development,

and transport might be expected in the rachis, a region of the leaf that acts as a

connective corridor to the rest of the plant, we were surprised to find up-regulation of so

many genes defined by GO terms related to photosynthesis. As noted in the pairwise

differential gene expression analysis described above, the most abundant enriched GO

terms for up-regulated genes in the rachis were related to sugar biosynthesis and

photosynthesis, indicating that the rachis region likely has functioning photosynthetic

machinery before it is acquired by the P4 margin, which is destined to become the

blade, the primary photosynthetic tissue of the leaf. Since little is known about when

photosynthesis first begins in a developing leaf, and no previous studies have described

photosynthesis specifically in the rachis, we attempted to verify the notion that the

rachis is a photosynthetic force during early leaf development.

To verify the photosynthetic signature repeatedly found to be up-regulated in rachis

compared to margin tissue, we searched for photosynthetic genes in our dataset that

showed significant differential expression between the rachis and margin in each

longitudinal region. We identified three *Light Harvesting Chlorophyll A-B binding* (*CAB*)

genes (Solyc03g005760 [*SlCAB1*], Solyc03g005760 [*SlCAB2*], and Solyc03g005760

[*SlCAB3*]) with significantly up-regulated expression in the rachis regions compared to

the margin (**Figure 4C**, **Data S1**). CAB proteins balance excitation energy between

Photosystem I and II during photosynthesis (Liu and Shen, 2004) and are important

components of photosynthesis.

In an attempt to verify the gene expression differences identified in our experimental

set-up and visualize when and where photosynthetic activity begins in a leaf

primordium, we constructed a transgenic line expressing a representative *CAB* gene

promoter attached to the β-glucuronidase (GUS) reporter (*pCAB1::CAB1::GUS*) (Mitra

et al., 2009; Tindamanyire et al., 2013). In the expanded leaflets of P8 leaves,

pCAB::CAB1::GUS expression was nearly ubiquitous across the entire blade (**Figure

4D**). At this age, the leaf had the anatomy of a fully functional photosynthetic organ. As

predicted from our gene expression analysis, in younger leaf primordia, we found a

clear *pCAB::CAB1::GUS* signal localized predominantly in the rachis region along the

proximal-distal axis in P4-P7 leaves (**Figure 4E and F**). The *pCAB::CAB1::GUS* signal

spread to the distal tips of newly established leaflets and lobes after P4 and continued

to spread to the margin tissue as development proceeded until the entire leaf showed

expression (**Figure 4D-G**).

Since pCAB::CAB1::GUS is predominantly expressed in the rachis region during early

development, we suggest that the rachis is the first region in a developing leaf to

function photosynthetically, as predicted in our RNA-seq analysis. Previous studies

have suggested that chloroplast retrograde signaling and sugar trigger leaf

differentiation (Andriankaja et al., 2012; Lastdrager et al., 2014). In the light of these

studies, the enrichment of photosynthetic genes in the rachis provides the first line of

evidence that during a very early developmental stage (P4), the rachis region does not

simply function as a conduit for nutrients and water transport, but it also functions in

photosynthesis and sugar production.

Self Organizing Maps (SOM) identify specific groups of genes that share similar

expression patterns

To refine our results and identify groups of genes sharing similar co-expression patterns

that may be too complex to define by differential expression analysis alone, we used

Self Organizing Mapping to cluster genes based on gene expression patterns across

the six tissue groups. SOM (Tamayo et al., 1999) begins by randomly assigning a gene

to a cluster. Genes are subsequently assigned to clusters based on similar expression

patterns via a reiterative process informed by previous cluster assignments. This

clustering method allows genes to be grouped based on specific expression patterns

shared across different tissues, allowing genes to be classified into smaller groups than

those generated by differential expression analysis alone. SOM analysis also allowed

us to survey the most prominent types of gene expression patterns found in our data.

To focus on the most variable genes across tissues, we selected the top 25% of genes

with the highest coefficients of variation, resulting in a dataset of 6,582 unique genes

(**Dataset S5**). We used principal component analysis to visualize groups of genes and
found that the first four principal components explained 31.9%, 26.2%, 19.0%, and
13.5% of the amount of variation in the dataset, respectively (**S5 Figure A**). Looking at
the expression of these genes in the PC space revealed distinct clusters of genes with
related expression patterns (**Figure 5A**).
434
To identify the most common gene expression patterns that describe the data, SOM
analysis was initially limited to six clusters (**Dataset S6**). One of these clusters, Cluster
4 (with 1090 genes), defines a clear separation of margin and rachis tissues, which
again reinforces our finding that the expression levels of many genes differ depending
on where the sampled tissue is localized along the medio-lateral axis (margin vs.
rachis). This cluster is enriched in genes defined by Carbohydrate metabolic processes,
hydrolase activity, protein dimerization, membrane, transporter activity, and
photosynthesis and light harvesting (**S5 Figure, Dataset** S7). These findings mirror the
results obtained by differential gene expression analysis and reflect the overall
abundance and diversity of genes up-regulated in the rachis, comprising the largest
signal in our dataset. These findings likely reflect the specialization that occurs in
tissues as the rachis develops an identity distinct from the margin.
447
448   **Auxin transport and regulation as a defining feature of margin identity**
449
To refine our analysis of gene expression patterns to genes that direct margin identity,
we generated a larger clustering map. We used this approach to obtain a smaller subset
of genes than could be obtained by differential gene expression analysis or SOM
clustering using a smaller number of clusters. We were especially interested in
identifying specific types of gene expression patterns that defined the medio-lateral axis;
in this case, we looked for groups of genes that were preferentially up or down-
regulated in the margin compared to the rachis. We specified 36 clusters in a 6x6
hexagonal topology, forcing interactions between multiple tissue types (**Figure 6A, S6**
Figure). We surveyed the gene expression patterns of each of the 36 clusters (**Dataset**
S8) and identified Clusters 10 (n =108) and 11 (n=112), which describe a group of

genes that were up-regulated in the margin and down-regulated in the rachis in stage P4 wild-type plants (**Figure 6B - C**). While over half of these genes (57.2%; 126/220) have no known function, many of the remaining genes are known to be involved in leaf margin identity (**Table 1**). Interestingly, Clusters 10 and 11 also contained genes related to auxin transport, biosynthesis, and regulation (*YUC4*, *PIN1*, *AUX2-11*) and genes (*ARGONAUTE7*/Solyc01g010970) known to interact with Auxin Response Factors (ARFs) (Yifhar et al. 2012).

Guided by the gene expression data in the wild type, and the results that auxin might play a role leaflet initiation in the base margin region, we wanted to see if auxin transport differences in *tf-2* could explain the striking feature of loss of meristematic potential in the basal margin of this mutant. To look specifically at the differences in auxin transport between *tf-2* and wild type and to verify the differences in *SlPIN1* gene expression found between the wild type and *tf-2,* we crossed a fluorescently labeled *pPIN1::PIN1::GFP* line (*PIN1::GFP*) (Benková et al., 2003; Koenig et al., 2009) with *tf-2* to visualize differences in PIN1 localization and expression in P4 leaves. In the wild type, PIN1::GFP was present along the entire margin region of a P4 leaf, with the strongest signal present at the site of the newly established LL1 (**Figure 7A and B**). In *tf-2,* there was an overall decrease in fluorescent signal along the margin of a P4 leaf. Also, *tf-2* had a noticeable decrease in PIN1::GFP fluorescent signal in the base margin region (**Figure 7C and D**). In addition, we visualized auxin presence using the auxin-inducible promoter DR5::Venus (Bayer et al., 2009). As observed previously (Shani et al., 2010; Martinez et al., 2016), in the wild type, DR5::Venus was expressed at the site of leaflet initiation as a sharp wedge-shaped focus region (**Figure 7E-F**). By contrast, in *tf-2*, there was a DR5::Venus focus region, but it was diffuse and located in the upper layers of the margin (**Figure 7G-H**). These results support the hypothesis that while *tf-2* is capable of forming auxin foci, it is incapable of maintaining proper auxin foci and canalization processes, as evidenced by the reduced *PIN1* expression in the basal margin region of the *tf-2* P4 leaf. The transcriptomic results and auxin visualization experiments suggest that misregulated auxin transport and biosynthesis, and

490 specifically *SlPIN1* misregulation, are important contributors to the *tf-2* phenotype and

491 that these processes are vital regulators of margin organogenesis.

493 **Differences in gene expression patterns between the wild type and *tf-2* help**

494 **define the loss of basal meristematic potential in *tf-2* leaves**

496 We included *tf-2* in this study because it has the intriguing phenotype of being unable to

497 form new leaflets after the first two LL1 leaflets. At the P4 stage, *tf-2* has already lost the

498 organogenetic ability to initiate new leaflets. As revealed by our auxin transport

499 visualization analysis, *tf-2* appears to receive a leaflet initiation signal, as it is capable of

500 forming auxin foci (**Figure 7H**), but the tissue is unable to initiate leaflet organs. We

501 examined our data to determine whether gene expression differences could explain the

502 loss of meristematic competency in *tf-2*. We performed differential gene expression

503 analysis of only *tf-2* reads. There were a lot fewer differentially expressed genes

504 between the margin and rachis in the top, middle, and base regions of this mutant

505 compared to the wild type (**S3 Figure A, Dataset S1**). However, *tf-2* followed similar

506 gene expression trends to the wild type when margin and rachis identity were

507 compared. The margin was more enriched in genes related to cell division and cell

508 expansion, while the rachis was enriched in genes related to specialization, including

509 water transport, metabolic processes, photosynthesis, and leaf development; however,

510 these differences were mostly apparent in the base region of the *tf-2* mutant (**S3 Figure**

511 **B**). The main difference between wild type and the *tf-2* mutant was a reduction in up-

512 regulated differentially expressed genes in the rachis region compared to the margin in

513 top, middle, and base regions (**S3 Figure A**). It should be noted that while wild type and

514 *tf-2* were similar morphologically at the P4 stage, the *tf-2* mutant appeared to be further

515 along in the morphogenesis process in all regions (top, middle, and base), a feature

516 described by Naz and coworkers (2013). This overall difference in the two genotypes

517 should be taken into account at the morphological level, and, as evidenced by our

518 transcriptional analysis, at the molecular level. In the margin of *tf-2*, we examined the

519 differentially expressed genes between the rachis and margin and found many genes

520 related to leaf development.

Taking into account the overall differences between these two genotypes, we were

interested in understanding why *tf-2* is unable to initiate lateral leaflets beyond LL1.

Could transcriptional differences explain the loss of morphogenic capacity in *tf-2*? To

address this issue, we combined both genotypes and used a generalized linear model

(glmQLFTest in edgeR) in which we defined each genotype as a group and therefore

could compare the top, middle, and base regions of the two genotypes. When we

compared the base margin region between *tf-2* and the wild type (**Figure 1A**), only 23

genes were differentially expressed: all were downregulated in the wild type compared

to *tf-2* (**Table 3**). We focused on the 12 genes that were functionally annotated and

noticed that *SlBOP2* was significantly up-regulated in the margin of *tf-2* compared to the

wild type (**Figure 8B**).

We explored the role of *SlBOP2* in regulating margin and rachis tissue identity by

phenotyping CRISPR/Cas9 gene edited loss-of-function *SlBOP2* mutants (*CR-slbop2)*

(Xu et al., 2016) with a focus on leaf phenotypes. Surprisingly, in *CR-slbop2* plants, we

observed ectopic meristems on mature leaves when plants were approximately two

months old. The ectopic meristems occurred along the adaxial rachis of mature leaves

at the bases of primary leaflets (**Figure 8C, D, E**). These ectopic SAM structures did not

persist as the leave aged, appearing to undergo tissue death approximately three

weeks after appearing on the rachis (S6 A) and only rarely did they generate complex

leaf-like organs (**S6 C Figure**). Loss of function of *SlBOP2* also resulted in increased

leaf complexity (**S6 B Figure**), as reported previously in these mutants (Xu et al., 2016)

and in *SlBOP2* knockdown lines (Ichihashi et al., 2014). Since TF is a known

transcription factor, we checked for TF binding site motifs in the 3KB upstream region of

BOP2 and found one TF binding site (**Figure 8F**). Taken together, these findings

indicate that *SlBOP2* functions in determining margin meristematic identity along the

rachis of the leaf and the possibility that *SlBOP2* functions via the direct binding of TF to

its upstream regulatory region – although more validation is needed to verify this

interaction.

DISCUSSION

Unique genetic signatures define leaf development along the proximal-distal and medio-lateral axes

The overall goal of this work was to use gene expression signatures to gain a better understanding of the processes that regulate morphogenesis along the medio-lateral axis in an early developing compound leaf. Based on anatomical analysis, we chose six unique regions in the P4 leaf (**Figure 1C-F**). We analyzed differential gene expression between margin and rachis tissue in the top, middle, and base regions and identified signature patterns of gene regulation along the proximal-distal (tip-base) axis (**Figure 3**) that help define leaf morphogenesis in the early tomato leaf primordium.

In addition to a basipetal wave of differentiation along the proximal-distal axis, the leaf differentiates from the midrib/rachis out into the margins at each region on the proximo-distal axis. These two regions, the margin and rachis, have distinct developmental trajectories: the rachis matures early and the marginal blastozone retains some meristematic potential and gives rise to the leaf blade region, as well as leaflets in compound leaves. Separating the rachis from the marginal blastozone region at three different points along the proximal-distal axis allowed us to determine whether development proceeds uniformly along the proximal-distal axis or if the leaf has a mosaic of developmental states in each segment along the proximal-distal axis. The further along in morphogenesis a region was, the more diverse the GO categories of genes that were upregulated in the region, likely because the last stage of leaf morphogenesis, cell specialization, had occurred. After summarizing the enriched GO terms in each of the three regions along the proximal-distal axis, patterns of developmentally distinct processes were identified in the rachis regions compared to other tissues (**Figure 3**). The margin regions, classically defined as the marginal blastozone or marginal meristem, retain the potential to divide and differentiate and also exhibit a basipetal gradient of gene expression changes of differentiation from the tip to the base of the leaf. Thus, this analysis suggests that defining leaf development or

capturing gene expression in the entire primordium, or even in regions along the

proximal-distal axis, does not provide an accurate picture of developmental patterns in a

leaf. Further dissecting these events at cellular resolution should help define these

patterns even more accurately.

Photosynthetic capability in the rachis as a regulator of medio-lateral

differentiation

To further define rachis and margin identity, we fitted an additive model that adjusts

differential gene expression comparisons based on baseline differences that occur

between the margin and rachis. We then performed differential gene expression

analysis to reveal gene expression trends that define margin and rachis tissue

regardless of the position on the proximal-distal axis. The most prevalent, though

unexpected, gene expression signature we observed was the enrichment of genes

associated with photosynthesis in the rachis, which we found by both differential

expression analysis (**Figure 3 and 7**) and cluster analysis (**Figure 5**). Since little is

known about when photosynthetic capacity is acquired during early leaf morphogenesis,

we further verified photosynthesis activity using a CAB::GUS reporter (**Figure 4C-G**).

This analysis suggested that photosynthetic activity is acquired as early as P4 and is

not uniformly distributed along the proximal-distal and medio-lateral axes. When viewed

in the context of cell differentiation processes along each axis, it is not surprising that

specialized functions are first acquired in regions that mature earliest, although the

function of photosynthesis has been traditionally assigned to the blade. What are the

developmental consequences of sugar biosynthesis in the rachis during early leaf

organogenesis? Could the rachis be the source of morphogenic signaling towards the

more immature base along the proximal-distal axis and along the medio-lateral axis to

the margin? Multiple studies in *A. thaliana* identified thousands of genes that respond to

changes in sugar levels by modifying transcript abundance (Price et al., 2004; Bläsing

et al., 2005; Osuna et al., 2007; Usadel et al., 2008). These studies suggest that the

main photosynthetic product, sugar, functions as a signal for plant development and

growth.

In the P4 primordium under study, while the rachis has acquired specialized functions, the margin is actively dividing, a process that relies on cell cycle progression. The cyclin genes *CYCD2* and *CYCD3*, encoding critical regulators of the cell cycle, are up-regulated in response to sugar (Riou-Khamlichi et al., 2000). Interestingly, sucrose has also been shown to influence auxin levels (Lilley et al., 2012; Sairanen et al., 2012), transport, and signal transduction (Stokes et al., 2013), and metabolism (Ljung, 2013). Moreover, sugar accumulation is spatiotemporally regulated in meristematic tissue in both the shoot and root apical meristem (Francis and Halford, 2006). Is the development of photosynthetic capacity in the rachis a cause or a consequence of its early differentiation? Do the acquisition of photosynthetic capability and the production of sugars represent a global mechanism for signaling quiescent regions to progress into the cell division phase? More work exploring photosynthesis, sugar transport, hormone regulation, and gene expression should help uncover a possible role for the rachis in regulating morphogenetic processes during early leaf organogenesis.

The presence of auxin as a defining feature of organogenic potential in margin tissue

PIN1-directed auxin transport is an important regulator of leaf development (Reinhardt et al., 2003; Heisler et al., 2005; Scarpella and Helariutta, 2010; Kawamura et al., 2010; Hay and Tsiantis, 2006; Scarpella et al., 2006) and leaflet initiation (Koenig et al., 2009). A common mechanism unites PIN1-directed development during leaf organogenesis across the systems studied: PIN1 directs auxin along the epidermal layer to sites of convergence on the meristem and transports the auxin subepidermally into the internal layers (Scarpella et al., 2010). *PIN1* can be split into two highly supported sister clades: *PIN1* and *Sister of PIN1* (*SoPIN1*) (O'Connor et al., 2014; Bennett et al., 2014; Abraham Juárez et al., 2015). The *SoPIN1* and *PIN1* clades might have disparate but complementary functions in auxin transport during organ initiation, where SoPIN1 mainly functions in epidermal auxin flux to establish organ initiation sites and PIN1 functions in the transport of auxin inward (O'Connor et al., 2014; Abraham Juárez et al.,

2015; Martinez et al., 2016). Tomato has one gene in the *PIN1* clade (*SlPIN1*) and two

genes in the *SoPIN1* clade (*SlSoPIN1a* [Solyc10g078370] and *SlSoPIN1b*

[Solyc10g080880]) (Pattison and Catalá, 2012; Nishio et al., 2010; Martinez et al.,

2016). The current findings suggest that in *tf-2, SlPIN1* is down-regulated at the region

of leaflet initiation compared to the wild type. Using *PIN1::GFP* as a reporter, we

observed a lack of fluorescence in the base marginal blastozone region of *tf-2* (**Figure**

7C-G). Using DR5:VENUS as a reporter, we observed the diffuse localization of auxin in

the base margin region of *tf-2* apices. Interestingly, even with external auxin application,

tf-2 is not capable of leaflet initiation (Naz et al., 2013), suggesting that the ability to

direct auxin inwards using PIN1, and not auxin accumulation itself, may be

compromised in this mutant. It remains to be seen whether a common theme for organ

formation will emerge in organisms in which the *PIN1* clade has diverged into two

groups. Based on our findings about the marginal blastozone region in *tf-2*, we suggest

that in addition to the creation of auxin foci, the drainage of auxin into internal leaf layers

might also be required for leaflet initiation. Analysis of higher-order mutants in the larger

PIN1 clade should help resolve this issue.

Organogenic potential of the margin and homology of the leaf margin and the

SAM

In the current study, we were especially interested in obtaining a genetic understanding

of the loss of organogenetic potential in the base margin of *tf-2*. We specifically looked

for the transcriptional differences that explain the loss of organogenetic potential in *tf-2*

compared with the wild type, specifically in the margin base region. This led us to a

small list of 23 differentially expressed genes including *SlBOP2* (**Figure 8, Table 2**).

Characterization of the *CR-bop2* line (**Figure 8**) revealed ectopic SAM production along

the adaxial rachis at the bases of primary leaflets of the complex leaf. This finding

supports the notion that the suppression of meristematic identity by BOP family

members is important during leaf morphogenesis. BOP1 was first introduced as a

suppressor of lamina differentiation on the petioles of simple *Arabidopsis* leaves (Ha et

al., 2003, 2004) that limits meristematic cell activity, as the *bop1* mutant displays ectopic

meristematic cells beyond the boundary between the base of the blade and petiole (Ha

et al., 2003, 2004). Further work using *SlBOP* knockdown and knockout tomato lines

demonstrated that SlBOP2 suppresses organogenetic potential (S6 Figure; (Xu et al.,

2016; Ichihashi et al., 2014), as lines with reduced or absent *SlBOP* function showed

increased leaflet organ initiation/leaf complexity. BOPs interact with transcription factors

to regulate floral identity, including the interaction of BOP with PERIANTHIA (PAN) in

Arabidopsis (Hepworth and Pautot, 2015) and the interaction of TERMINATING

FLOWER (TMF) with *SlBOPs* to repress meristematic maturation in tomato flowers

(MacAlister et al., 2012; Xu et al., 2016). We further hypothesize that *SlBOP2* and a

transcription factor interact to regulate organogenic potential in complex leaves.

Specifically, perhaps the TF transcription factor binds to the upstream regulatory region

of *SlBOP2* (**Figure 8F**). We suggest that both *TF and SlBOP2* function in suppressing

meristematic properties of the margin during an early developmental window that

gradually closes with leaf maturation, an idea consistent with the view of the marginal

blastozone described by Hagemann (1970).

While our understanding of the recruitment of genetic regulators in a spatiotemporal

context continues to increase, one of the more exciting questions still remains: Is the

marginal meristem evolutionarily derived from the SAM (Floyd and Bowman, 2010)?

Ectopic adventitious SAMs have been shown to occur on leaves of functional knockouts

of *CUP-SHAPED-COTYLEDONS2* (*CUC2*) and *CUC3* (Blein et al., 2008; Aichinger et

al., 2012; Hibara et al., 2003) and of Arabidopsis lines overexpressing homeobox genes

KNOTTED-1 (*KN1)* and *Kn1-like* (*KNAT1*) (Chuck et al., 1996; Sinha and Hake, 1994)

in a region analogous to the base of an emerging leaflet, suggesting developmental

analogy and possibly homology to axillary meristems. Axillary meristems form on the

adaxial surface at the boundary zone between the leaf and SAM, where BOP2 has

already been shown to play a regulatory role in this process in tomato (Izhaki et al.,

2018), barley (*Hordeum vulgare*) (Tavakol et al., 2015; Dong et al., 2017), and maize

(Dong et al., 2017). The ectopic meristem phenotype of *CR-slbop2* on the margins of

complex tomato leaves suggests that signals might be recruited in the margin that are

similar to those present in leaf initiation sites during axillary meristem formation. Our

findings add further evidence that the margin is analogous, and possibly homologous at

the process level, to the SAM. The leaf margin likely evolved via the genetic recruitment

of similar regulatory factors, including BOP, reinforcing the importance of the reiteration

of genetic mechanisms to establish distinct spatial identities in neighboring domains

during plant development.

Our current understanding of the leaf margin is based on foundational work that defined

the margin by explicitly tracking developmental landmarks (Avery, 1933; Poethig and

Sussex, 1985a; Dolan and Poethig, 1998; Wolf et al., 1986). Early literature defined the

leaf primordium as broadly meristematic during early development, with this

meristematic potential becoming restricted and gradually lost as the leaf develops

(Foster, 1936; Hagemann and Gleissberg, 1996; Sachs, 1969). Although such studies

provide a roadmap for describing growth patterns in the margin, a major challenge is to

understand how these patterns are specified at the genetic level (Whitewoods and

Coen, 2017; Coen et al., 2017) and how this fits with our interpretation of the

recruitment of regulatory mechanisms suppressing the morphogenetic potential of the

margin during the evolution of leaves in seed plants. Plant development is reliant on

reiterative patterning, and leaf development is no exception. Our findings suggest that

we can describe leaf development as the reiteration and modulation of similar

evolutionarily derived genetic programs that act to suppress the morphogenetic and

organogenetic potential of meristematic regions in order to achieve final leaf form.

Follow-up studies in additional species are needed to understand these evolutionarily

conserved mechanisms and how they have been modulated to sculpt the diversity of

leaf forms observed in nature.

METHODS

Plant growth and tissue embedding

Seeds of tomato mutant *tf-2* (LA0512) and wild-type line Condine Red (LA0533) were

obtained from the Tomato Genetics Resource Center (TGRC). The seeds were

sterilized with 50% bleach for two minutes and rinsed 10 times with distilled water. The

seeds were placed on moist paper towels in Phytotrays (Sigma-Aldrich), incubated in

the dark for two days, and allowed to germinate in a growth chamber for four days

before being transferred to soil for 8 days of growth; the seedlings were grown for a total

of 14 days. Generation of the transgenic DR5::Venus (cv. M82) line was described in

(Shani et al., 2010) and the *AtpPIN1::PIN1::GFP* (cv. Moneymaker*)* line was described

in (Bayer et al., 2009)*.*

CR-bop-2 RNAi lines were received from the Lippman Lab at Cold Spring Harbor. CR-

bop-2 RNAi and wild type plant (M82) were germinated and grown in growth chambers

for two weeks following methods above. Plants were further grown in a growth chamber

for one month. After one month, plants were transferred to a greenhouse and grown

under normal light conditions. The plants were observed, and any abnormalities were

noted, dated, and photographed throughout the lifetime of each plant. The images are

taken from mature leaves from plants that were 63 days old. Leaf complexity counts (S6

Figure) were taken at age 45 days old.

Plants were collected in the afternoon, vacuum infiltrated for one hour with ice-cold 3:1

(100% EtOH: 100% acetic acid) fixative, and fixed overnight at 4°C. The samples were

washed three times in 75% EtOH, passed through an EtOH series on a shaker at room

temperature for one hour per step (75%, 85%, 95%, 100%, 100%, 100%), and

incubated in 100% EtOH overnight at 4°C. All ethanol solutions were made using 2X

autoclaved diethylpyrocarbonate (DEPC) treated water. The tissue was passed through

a xylene/EtOH series for two hours per step (25%, 50%, 75%, 100%, 100%) on a

shaker at room temperature. The tissue was incubated overnight at room temperature in

100% xylene with 20-40 paraffin chips (Paraplast x-tra, Thermo Fisher Scientific),

followed by incubation at 42°C until the paraffin dissolved. The paraffin:xylene solution

was subsequently removed and replaced with 100% paraffin and the sample was

incubated for 3 days at 55°C, with the solution changed twice daily. The tissue was then

embedded using tools and surfaces that had been washed with RNAseZap (Thermo

Fisher Scientific) and DEPC. The embedded blocks were transversely sectioned at 5 to

7 µm thickness using a Leica RM2125RT rotary microtome (Leica Microsystems) on

RNase-free polyethylene naphthalate PEN membrane slides (Leica). The slides were

dried at room temperature and deparaffinized with 100% xylene.

EdU Visualization

Cell division was visualized by observing fluorescent signals derived from an EdU

incorporation assay in which EdU is incorporated into cells during the S-phase

(Kotogány et al., 2010). The EdU assay was performed as previously described

(Nakayama et al., 2014; Ichihashi et al., 2011) with some modifications using a Click-iT

® EdU Alexa Fluor® Imaging kit (Invitrogen). Fourteen-day-old seedlings were

dissected under a microscope. After removing older leaves, P4 leaf epidermis was

nicked using an insect mounting needle to increase infiltration in subsequent steps. The

plant apex was incubated in water containing 10 µM EdU for two hours. The samples

were washed in 1x phosphate-buffered saline solution (PBS, pH 7.4) and fixed in FAA

under vacuum infiltration for 3 h. Subsequently, the samples were fixed in 3.7%

formaldehyde in PBS (pH 7.4) for 30 min and washed three times in PBS with shaking.

Alexa Fluor coupling to EdU was performed in the dark following the manufacturer's

instructions. Photographs were taken under a Zeiss LSM 710 Confocal Microscope with

excitation wavelengths set at 488 and 420 nm.

Flow Cytometry and GUS Staining

Ploidy levels were measured using a PA-I ploidy analyzer (Partec) as described

previously (Sugimoto-Shirasu et al., 2002). We grew up 50 plants and sampled the

oldest two leaves on the plant at each timepoint. To identify the leaves by counting from

youngest leaves, requires apical meristem destruction, therefore we choose to use the

count Leaf age from oldest leaf - Leaf 1 corresponds to the oldest leaf, Leaf 2 is second

oldest and so on. For the first two timepoints (Day 8 and 18) we sampled L1 or L2 and

for the later timepoints, Day 30 and above, we sampled the oldest intact leaf, which

because of age was often damaged, and therefore the leaves sampled ranged from

Leaf 2 – Leaf 5. All leaves sampled beyond Day 30 had reached maturity. Fresh tissue

was extracted from whole leaves at the youngest leaf age (Day 8), while older stage

tissue was extracted from both the top and bottom sections of the leaf (Day 18 – 90).

The tissue was chopped with a razor blade. Cystain extraction buffer (Partec) was used

to release nuclei. The solution was filtered through a CellTrics filter (Partec) and stained

with Cystain fluorescent buffer (Partec). At least 4000 isolated nuclei isolated were used

for each ploidy measurement. Flow cytometry experiments were repeated at least three

times using independent biological replicates.

Histochemical localization of GUS activity was performed as previously described (Kang

and Dengler, 2002). Representative images were chosen from >15 samples stained in

three independent experiments.

Laser Capture Microdissection and RNA processing

Each tissue type was independently captured through serial sections using a Leica

LMD6000 Laser Microdissection System (Leica Microsystems). Each biological replicate

contained tissue collected from 5-8 apices. S1 Figure and Movie 1 show how tissue

regions were identified and dissected. Tissue was collected in lysis buffer from an

RNAqueous®-Micro Total RNA Isolation Kit (Ambion) and immediately stored at -80 °C.

RNA extraction was performed using an RNAqueous®-Micro Total RNA Isolation Kit

(Ambion) following the manufacturer's instructions. The RNA was amplified using the

WT-Ovation™ Pico RNA Amplification System (ver. 1.0, NuGEN Technologies Inc.).

The RNA was purified using RNAClean ® magnetic beads (Agencourt) and processed

within one month of fixation to ensure RNA quality.

RNA-seq libraries were created as described by Kumar and coworkers (Kumar et al.,

2012), starting with the second-strand synthesis step, with the following modifications:

For second-strand synthesis, 10 µL of cDNA (>250 ng) was combined with 0.5 µl of

random primers and 0.5 µl of dNTP. The sample was incubated at 80ºC for 2 min,

followed by 60ºC for 10 sec, 50ºC for 10 sec, 40ºC for 10 sec, 30ºC for 10 sec, and 4ºC

for at least 2-5 min. After adding 5 µl of 10x DNA pol buffer, 31 µL water, and 2.5 µL

DNA Pol I on ice, the sample was incubated at 16ºC for 2.5 h. The process was

continued following the published (Kumar et al., 2012) protocol starting with step 2.3:

Bead purification of double-stranded DNA. The libraries were quality checked and

quantified using a Bioanalyzer 2100 (Agilent) on RNA 6000 Pico Kit (Agilent) chips at

the UC Davis Genome Center. The libraries were sequenced in three lanes using the

HiSeq2000 Illumina Sequencer at the Vincent J Coates Genomics Sequencing

Laboratory at UC Berkeley.

Read Processing, Differential Expression, and Gene Ontology (GO) Enrichment

analysis

Quality filtering, N removal, and adaptor trimming were performed on data from each of

the three Illumina sequencing lanes separately. We first performed N removal using

read_N_remover.py. Sequences below a quality (phred) score of 20 were removed

without reducing the read size to below 35 bp. To remove adapter contamination, we

used adapterEffectRemover.py, setting the minimum read length to 41. To assess the

quality of the reads after pre-processing, we ran FASTQC (available at

http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/) before and after pre-processing.

To filter out reads from chloroplast or mitochondrial sequences, all libraries were

mapped to the S.lycopersicum_AFYB01.1 mitochondrial sequence from NCBI and the

NC_007898.3 chloroplast sequence from NCBI using STAR 2.4.0 (Dobin et al., 2013).

Reads that did not map to either organelle were mapped to the ITAG3.10 *Solanum*

lycopersicum genome using STAR 2.4.0, where non-genic sequences were masked

using the inverse coordinates of the ITAG3.10 gene model gff file. Bedtools (Quinlan,

2014) coverageBed was then used to count mapped reads, using a bed file generated

from ITAG3.10 gene models. We built an online visualization tool for the community to

manually explore the reads generated across the six tissue types in both wild type and

tf-2: bit.ly/2kkxsFQ (Winter et al., 2007; Patel et al., 2012).

Read processing and differential expression analysis were performed using the R

package edgeR (Robinson et al., 2010). Pairwise differential gene expression in each

region along the proximal-distal axis was calculated in each proximal-distal region (top,

middle, base) in separate analyses. Differential gene expression was determined using

`exactTest()`, multiple testing correction was performed using the Benjamini–Hochberg

procedure, and significance of differential expression was determined using a cutoff of

FDR < 0.05. To estimate differential expression of genes across the entire marginal

blastozone and rachis regions, we used an additive linear model where the proximal-

distal axis was assigned as a blocking factor, which adjusts for differences between the

margin and rachis in the top, middle, and base: model.matrix(~Region + Tissue). For

both pairwise and modeling analysis of differential expression, counts per million were

calculated from raw reads, and genes with < 5 reads in 2 or more reps were removed.

We estimated common negative binomial dispersion and normalized counts across all

samples using the trimmed mean of M-value (TMM) method (Robinson and Oshlack,

2010). Normalized Read counts, as calculated by Counts Per Million (cpm), are

available in **Dataset S9**. GO enrichment analysis was performed using the R libraries

GO.seq and GO.db. The GO terms were summarized using REVIGO

(http://revigo.irb.hr/). The full code used for these analyses is available at

http://github/iamciera/lcmProject (DOI provided upon publication).

SOM clustering

To explore the genes whose expression levels were the most variable across tissues,

we identified the top 25% genes based on coefficient of variation and ratio of standard

deviation compared to mean from our count data. To remove differences in counts

between samples, the magnitude of gene expression data was scaled between 2 and -2

in wild type and *tf-2* separately using the `scale()` function (Team, 2014). Hexagonal

layout was used for all SOM clustering (Kohonen). For basic SOM analysis, the SOM()

function was used for each genotype separately, while superSOMs were performed

using superSOM() in the Kohonen R package (Wehrens and Buydens, 2007). Training

for both methods was performed in 100 iterations in which the adaptive learning rate

decreased from 0.05 to 0.01. Codebook vectors and distance plots of cluster
assignments were generated using the visualization functions in the Kohonen R
package (Wehrens and Buydens, 2007) and ggplot2 (Wickham, 2009). To ensure that
the major variances in gene expression patterns were defined by SOM clustering and to
verify consistency in clustering, cluster assignments were projects onto the PC space.
All scripts used in clustering are available at http://github/iamciera/lcmProject (DOI
provided upon publication).
899
900
901 **AUTHOR CONTRIBUTIONS**
902
903 Conceived and designed the experiments: CM, NS. Performed molecular experiments
904 and plant characterizations: CM. Plant maintenance and phenotyping of *CR-bop2* lines:
905 SL. Contributed plant lines, protocols, and/or reagents: NS, KS. Performed
906 computational analysis: CM. Performed read mapping: MW. Analyzed the data: CM,
907 NS, KS. Wrote the paper: CM, NS. Edited the paper: CM, NS, KS.
908
909 **ACKNOWLEDGEMENTS**
910

The UC Davis Tomato Genetics Resource Center provided tomato germplasm,
AtpPIN1: PIN1: GFP and the DR5: VENUSx6 lines are gifts from Cris Kuhlemeier
(University of Bern) and Naomi Ori (Hebrew University, Israel), respectively. We thank
Zachary Lippman (Cold Spring Harbor Laboratory) for the *CR-slbop2* lines. We would
also like to acknowledge Eddi Esteban, Asher Pasha, and Nicholas J. Provart for
building the EFP browser. C.C.M. was supported by a National Science Foundation
Graduate Research Fellowship (DGE-1148897), Katherine Esau Summer Fellowship,
Walter R. and Roselinde H. Russell Fellowship, and Elsie Taylor Stocking Fellowship.
Part of the work was supported by NSF PGRP grants IOS–0820854 (to N.R. S., Julin
Maloof and Jie Peng), and S.L was partially supported by IOS 1856749 (to Julia Bailey-
Serres, Siobhan Brady, Roger Deal, Uta Paszcowsky, and N. R. S.). C.C.M. was also
supported by a collaboration between the National Science Foundation and the Japan

Society for the Promotion of Science with a Graduate Research Opportunities
Worldwide (GROW) award. This material utilized resources supported by the National
Science Foundation under Award Numbers DBI-0735191, DBI-1265383, and DBI-
1743442. URL: www.cyverse.org. We thank Daniel Chitwood (Michigan State
University) who provided computational training and insightful discussion which greatly
assisted the research and Kristina Zumstein (UC Davis) for her work and organization of
several experiments. We would also like to thank Siobhan Brady (UC Davis) and
Andrew Groover (USDA) for their helpful comments and John Harada (UC Davis) and
Julie Pelletier (UC Davis) for use and training on the laser capture microdissection
microscope.

Supplemental Data

S1_Dataset_allsig_DE_seperately.csv - Results of differential gene expression
analysis between the margin and rachis in the top, middle, and base regions of both
genotypes (WT and *tf-2*)*.*
S2_Dataset_sig_go_terms.csv - GO terms describing differentially expressed genes
between the margin and rachis in the top, middle, and base regions of both genotypes
(WT and *tf-2*)*.*
S3_Dataset_wt_modelled_DE.txt - Results of differential expression analysis across
the margin and rachis tissues performed with only wild-type reads and adjusted for
variability between the proximal-distal axis. An additive linear model was employed
using the top, middle, and base identities as a blocking factor in our experimental
design.
S5_Dataset_top25_coefficent_of_variation.csv - Genes with the most variable
expression. The top 25% of genes were selected based on coefficient of variation,
resulting in a dataset of 6,582 unique genes.
S6_Dataset_wt_SOM_small_cluster_assignments.csv - SOM cluster assignments
for wild type using a codemap vector of 6 showing the top six gene expression clusters.
S7_wt_SOM_small_sigGOterms.csv - GO terms derived from Dataset S6.

S8_wt_SOM_large_cluster_assignments.csv - SOM cluster analysis using a

codemap vector of 36 in wild type.

S9_Dataset_normalizedReadCount_cpm.csv - Normalized Read counts calculated

as counts per million (cpm).

S10_Dataset_tf2_modelled_DE.txt - Results of differential expression analysis across

the margin and rachis tissue performed with only *tf-2* reads and adjusted for variability

between the proximal-distal axis. An additive linear model was employed using the top,

middle, and base identities as a blocking factor in our experimental design.

REFERENCES

Abraham Juárez, M.J., Hernández Cárdenas, R., Santoyo Villa, J.N., O'Connor, D.,

Sluis, A., Hake, S., Ordaz-Ortiz, J., Terry, L., and Simpson, J. (2015).

Functionally different PIN proteins control auxin flux during bulbil development in

Agave tequilana. J. Exp. Bot. **66**: 3893–3905.

Aichinger, E., Kornet, N., Friedrich, T., and Laux, T. (2012). Plant stem cell niches.

Annu. Rev. Plant Biol. **63**: 615–636.

Andriankaja, M., Dhondt, S., De Bodt, S., Vanhaeren, H., Coppens, F., De Milde, L.,

Mühlenbock, P., Skirycz, A., Gonzalez, N., Beemster, G.T.S., and Inzé, D.

(2012). Exit from proliferation during leaf development in Arabidopsis thaliana: a not-

so-gradual process. Dev. Cell **22**: 64–78.

Avery, G.S. (1933). Structure and Development of the Tobacco Leaf. Am. J. Bot. **20**:

565–592.

Bayer, E.M., Smith, R.S., Mandel, T., Nakayama, N., Sauer, M., Prusinkiewicz, P.,

and Kuhlemeier, C. (2009). Integration of transport-based models for phyllotaxis

and midvein formation. Genes Dev. **23**: 373–384.

Beemster, G.T.S., De Veylder, L., Vercruysse, S., West, G., Rombaut, D., Van

Hummelen, P., Galichet, A., Gruissem, W., Inzé, D., and Vuylsteke, M. (2005).

Genome-wide analysis of gene expression profiles associated with cell cycle

transitions in growing organs of Arabidopsis. Plant Physiol. **138**: 734–743.

Bendahmane, A. and Theres, K. (2011). Shoot Branching and Leaf Dissection in
Tomato Are Regulated by Homologous Gene Modules. Plant Cell: 1–16.

Benková, E., Michniewicz, M., Sauer, M., Teichmann, T., Seifertová, D., Jürgens,
G., and Friml, J. (2003). Local, efflux-dependent auxin gradients as a common
module for plant organ formation. Cell **115**: 591–602.

Bennett, T., Brockington, S.F., Rothfels, C., Graham, S., Stevenson, D., Kutchan,
T., Rolf, M., Thomas, P., Wong, G.K.-S., Leyser, O., and Others (2014).
Paralagous radiations of PIN proteins with multiple origins of non-canonical PIN
structure. Mol. Biol. Evol.: msu147.

Bergervoet, J.H.W., Berhoeven, H.A., Gilissen, L.J.W., and Bino, R.J. (1996). High
amounts of nuclear DNA in tomato (Lycopersicon esculentum Mill.) pericarp. Plant
Sci. **116**: 141–145.

Bilsborough, G.D., Runions, A., Barkoulas, M., Jenkins, H.W., Hasson, A., Galinha,
C., Laufs, P., Hay, A., Prusinkiewicz, P., and Tsiantis, M. (2011). Model for the
regulation of Arabidopsis thaliana leaf margin development. Proc. Natl. Acad. Sci. U.
S. A. **108**: 3424–3429.

Bläsing, O.E., Gibon, Y., Günther, M., Höhne, M., Morcuende, R., Osuna, D.,
Thimm, O., Usadel, B., Scheible, W.-R., and Stitt, M. (2005). Sugars and
circadian regulation make major contributions to the global regulation of diurnal
gene expression in Arabidopsis. Plant Cell **17**: 3257–3281.

Blein, T., Pulido, A., Vialette-Guiraud, A., Nikovics, K., Morin, H., Hay, A.,
Johansen, I.E., Tsiantis, M., and Laufs, P. (2008). A conserved molecular
framework for compound leaf development. Science **322**: 1835–1839.

Bourdon, M., Frangne, N., Mathieu-Rivet, E., Nafati, M., Cheniclet, C., Renaudin, J.-
P., and Chevalier, C. (2010). Endoreduplication and Growth of Fleshy Fruits. In
Progress in Botany 71, U. Lüttge, W. Beyschlag, B. Büdel, and D. Francis, eds,

Progress in Botany. (Springer Berlin Heidelberg), pp. 101–132.

Cheniclet, C., Rong, W.Y., Causse, M., Frangne, N., Bolling, L., Carde, J.-P., and
Renaudin, J.-P. (2005). Cell expansion and endoreduplication show a large genetic
variability in pericarp and contribute strongly to tomato fruit growth. Plant Physiol.
139: 1984–1994.

Chuck, G., Lincoln, C., and Hake, S. (1996). KNAT1 induces lobed leaves with ectopic
meristems when overexpressed in Arabidopsis. Plant Cell **8**: 1277–1289.

Coen, E., Kennaway, R., and Whitewoods, C. (2017). On genes and form.
Development **144**: 4203–4213.

De Veylder, L., Larkin, J.C., and Schnittger, A. (2011). Molecular control and function
of endoreplication in development and physiology. Trends Plant Sci. **16**: 624–634.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq
aligner. Bioinformatics **29**: 15–21.

Dolan, L. and Poethig, R. (1998). Clonal analysis of leaf development in cotton. Am. J.
Bot. **85**: 315.

Dong, Z., Li, W., Unger-Wallace, E., Yang, J., Vollbrecht, E., and Chuck, G. (2017).
Ideal crop plant architecture is mediated by tassels replace upper ears1, a BTB/POZ
ankyrin repeat gene directly targeted by TEOSINTE BRANCHED1. Proc. Natl.
Acad. Sci. U. S. A. **114**: E8656–E8664.

Donnelly, P.M., Bonetta, D., Tsukaya, H., Dengler, R.E., and Dengler, N.G. (1999).
Cell cycling and cell enlargement in developing leaves of Arabidopsis. Dev. Biol.
215: 407–419.

Efroni, I., Blum, E., Goldshmidt, A., and Eshed, Y. (2008). A protracted and dynamic
maturation schedule underlies Arabidopsis leaf development. Plant Cell **20**: 2293–
2306.

Eshed, Y., Baum, S.F., Perea, J.V., and Bowman, J.L. (2001). Establishment of
polarity in lateral organs of plants. Curr. Biol. **11**: 1251–1260.

Floyd, S.K. and Bowman, J.L. (2010). Gene expression patterns in seed plant shoot
meristems and leaves: homoplasy or homology? J. Plant Res. **123**: 43–55.

Foster, A.S. (1936). Leaf differentiation in angiosperms. Bot. Rev. **2**: 349–372.

Francis, D. and Halford, N.G. (2006). Nutrient sensing in plant meristems. Plant Mol.
Biol. **60**: 981–993.

Gutierrez, C. (2005). Coupling cell proliferation and development in plants. Nat. Cell
Biol. **7**: 535–541.

Ha, C.M., Jun, J.H., Nam, H.G., and Fletcher, J.C. (2004). BLADE-ON-PETIOLE1
encodes a BTB/POZ domain protein required for leaf morphogenesis in Arabidopsis
thaliana. Plant Cell Physiol. **45**: 1361–1370.

Ha, C.M., Kim, G.-T., Kim, B.C., Jun, J.H., Soh, M.S., Ueno, Y., Machida, Y.,
Tsukaya, H., and Nam, H.G. (2003). The BLADE-ON-PETIOLE 1 gene controls
leaf pattern formation through the modulation of meristematic activity in Arabidopsis.
Development **130**: 161–172.

Hagemann, W. and Gleissberg, S. (1996). Organogenetic capacity of leaves: The
significance of marginal blastozones in angiosperms. Plant Syst. Evol. **199**: 121–
152.

Hay, A. and Tsiantis, M. (2006). The genetic basis for differences in leaf form between
Arabidopsis thaliana and its wild relative Cardamine hirsuta. Nat. Genet. **38**: 942–
947.

Heisler, M.G., Ohno, C., Das, P., Sieber, P., Reddy, G.V., Long, J. a., and
Meyerowitz, E.M. (2005). Patterns of auxin transport and gene expression during
primordium development revealed by live imaging of the Arabidopsis inflorescence
meristem. Curr. Biol. **15**: 1899–1911.

Hepworth, S.R. and Pautot, V.A. (2015). Beyond the Divide: Boundaries for Patterning
and Stem Cell Regulation in Plants. Front. Plant Sci. **6**: 1052.

Hibara, K.-I., Takada, S., and Tasaka, M. (2003). CUC1 gene activates the expression
of SAM-related genes to induce adventitious shoot formation. Plant J. **36**: 687–696.

Ichihashi, Y., Aguilar-Martínez, J.A., Farhi, M., Chitwood, D.H., Kumar, R., Millon,
L.V., Peng, J., Maloof, J.N., and Sinha, N.R. (2014). Evolutionary developmental
transcriptomics reveals a gene network module regulating interspecific diversity in
plant leaf shape. Proc. Natl. Acad. Sci. U. S. A. **111**: E2616–21.

Ichihashi, Y., Kawade, K., Usami, T., Horiguchi, G., Takahashi, T., and Tsukaya, H.
(2011). Key proliferative activity in the junction between the leaf blade and leaf
petiole of Arabidopsis. Plant Physiol. **157**: 1151–1162.

Izhaki, A., Alvarez, J.P., Cinnamon, Y., Genin, O., Liberman-Aloni, R., and Eyal, Y.
(2018). The Tomato BLADE ON PETIOLE and TERMINATING FLOWER Regulate
Leaf Axil Patterning Along the Proximal-Distal Axes. Frontiers in Plant Science **9**.

Joubès, J., Chevalier, C., Dudits, D., Heberle-Bors, E., Inzé, D., Umeda, M., and
Renaudin, J.-P. (2000). CDK-related protein kinases in plants. In The Plant Cell
Cycle, D. Inzé, ed (Springer Netherlands), pp. 63–76.

Kang, J. and Dengler, N. (2002). Cell cycling frequency and expression of the
homeobox gene ATHB-8 during leaf vein development in Arabidopsis. Planta **216**:
212–219.

Kawamura, E., Horiguchi, G., and Tsukaya, H. (2010). Mechanisms of leaf tooth
formation in Arabidopsis. Plant J. **62**: 429–441.

Kidner, C.A. and Timmermans, M.C.P. (2007). Mixing and matching pathways in leaf
polarity. Curr. Opin. Plant Biol. **10**: 13–20.

Koenig, D., Bayer, E., Kang, J., Kuhlemeier, C., and Sinha, N. (2009). Auxin patterns
Solanum lycopersicum leaf morphogenesis. Development **136**: 2997–3006.

Kohonen, T. Self-organized formation of topologically correct feature maps. Biol.
Cybern. **43**: 59–69.

Kondorosi, E., Roudier, F., and Gendreau, E. (2000). Plant cell-size control: growing
by ploidy? Curr. Opin. Plant Biol. **3**: 488–492.

Kotogány, E., Dudits, D., Horváth, G.V., and Ayaydin, F. (2010). A rapid and robust
assay for detection of S-phase cell cycle progression in plant cells and tissues by
using ethynyl deoxyuridine. Plant Methods **6**: 5.

Kumar, R., Ichihashi, Y., Kimura, S., Chitwood, D.H., Headland, L.R., Peng, J.,
Maloof, J.N., and Sinha, N.R. (2012). A High-Throughput Method for Illumina RNA-
Seq Library Preparation. Front. Plant Sci. **3**: 202.

Lastdrager, J., Hanson, J., and Smeekens, S. (2014). Sugar signals and the control
of plant growth and development. J. Exp. Bot. **65**: 799–807.

Lilley, J.L.S., Gee, C.W., Sairanen, I., Ljung, K., and Nemhauser, J.L. (2012). An
endogenous carbon-sensing pathway triggers increased auxin flux and hypocotyl
elongation. Plant Physiol. **160**: 2261–2270.

Li, P. et al. (2010). The developmental dynamics of the maize leaf transcriptome. Nat.
Genet. **42**: 1060–1067.

Liu, X.-D. and Shen, Y.-G. (2004). NaCl-induced phosphorylation of light harvesting
chlorophyll a/b proteins in thylakoid membranes from the halotolerant green alga,
Dunaliella salina. FEBS Lett. **569**: 337–340.

Ljung, K. (2013). Auxin metabolism and homeostasis during plant development.
Development **140**: 943–950.

MacAlister, C.A., Park, S.J., Jiang, K., Marcel, F., Bendahmane, A., Izkovich, Y.,
Eshed, Y., and Lippman, Z.B. (2012). Synchronization of the flowering transition by
the tomato TERMINATING FLOWER gene. Nat. Genet. **44**: 1393–1398.

Martinez, C.C., Koenig, D., Chitwood, D.H., and Sinha, N.R. (2016). A sister of PIN1

1112 gene in tomato (Solanum lycopersicum) defines leaf and flower organ initiation

1113 patterns by maintaining epidermal auxin flux. Dev. Biol. **419**: 85–98.

1114 **Melaragno, J.E., Mehrotra, B., and Coleman, A.W.** (1993). Relationship between

1115 Endopolyploidy and Cell Size in Epidermal Tissue of Arabidopsis. Plant Cell **5**:

1116 1661–1668.

1117 **Mitra, A., Han, J., Zhang, Z.J., and Mitra, A.** (2009). The intergenic region of

1118 Arabidopsis thaliana cab1 and cab2 divergent genes functions as a bidirectional

1119 promoter. Planta **229**: 1015–1022.

1120 **Moon, J. and Hake, S.** (2011). How a leaf gets its shape. Curr. Opin. Plant Biol. **14**:

1121 24–30.

1122 **Nakayama, H., Nakayama, N., Seiki, S., Kojima, M., Sakakibara, H., Sinha, N., and**

1123 **Kimura, S.** (2014). Regulation of the KNOX-GA gene module induces heterophyllic

1124 alteration in North American lake cress. Plant Cell **26**: 4733–4748.

1125 **Naz, A.A., Raman, S., Martinez, C.C., Sinha, N.R., Schmitz, G., and Theres, K.**

1126 (2013). Trifoliate encodes an MYB transcription factor that modulates leaf and shoot

1127 architecture in tomato. Proc. Natl. Acad. Sci. U. S. A. **110**: 2401–2406.

1128 **Nishio, S., Moriguchi, R., Ikeda, H., Takahashi, H., Takahashi, H., Fujii, N.,**

1129 **Guilfoyle, T.J., Kanahama, K., and Kanayama, Y.** (2010). Expression analysis of

1130 the auxin efflux carrier family in tomato fruit development. Planta **232**: 755–764.

1131 **O'Connor, D.L., Runions, A., Sluis, A., Bragg, J., Vogel, J.P., Prusinkiewicz, P.,**

1132 **and Hake, S.** (2014). A Division in PIN-Mediated Auxin Patterning during Organ

1133 Initiation in Grasses. PLoS Comput. Biol. **10**: e1003447.

1134 **Ori, N. et al.** (2007). Regulation of LANCEOLATE by miR319 is required for compound-

1135 leaf development in tomato. Nat. Genet. **39**: 787–791.

1136 **Osuna, D., Usadel, B., Morcuende, R., Gibon, Y., Bläsing, O.E., Höhne, M., Günter,**

1137 **M., Kamlage, B., Trethewey, R., Scheible, W.-R., and Stitt, M.** (2007). Temporal

1138 responses of transcripts, enzyme activities and metabolites after adding sucrose to

carbon-deprived Arabidopsis seedlings. Plant J. **49**: 463–491.

Patel, R.V., Nahal, H.K., Breit, R., and Provart, N.J. (2012). BAR expressolog

identification: expression profile similarity ranking of homologous genes in plant

species. Plant J. **71**: 1038–1050.

Pattison, R.J. and Catalá, C. (2012). Evaluating auxin distribution in tomato (Solanum

lycopersicum) through an analysis of the PIN and AUX/LAX gene families. Plant J.

70: 585–598.

Poethig, R.S. and Sussex, I.M. (1985a). The cellular parameters of leaf development

in tobacco: a clonal analysis. Planta **165**: 170–184.

Poethig, R.S. and Sussex, I.M. (1985b). The developmental morphology and growth

dynamics of the tobacco leaf. Planta **165**: 158–169.

Price, J., Laxmi, A., St Martin, S.K., and Jang, J.-C. (2004). Global transcription

profiling reveals multiple sugar signal transduction mechanisms in Arabidopsis.

Plant Cell **16**: 2128–2150.

Quinlan, A.R. (2014). BEDTools: the Swiss-army tool for genome feature analysis.

Curr. Protoc. Bioinformatics **47**: 11–12.

Reinhardt, D., Pesce, E.-R., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M.,

Traas, J., Friml, J., and Kuhlemeier, C. (2003). Regulation of phyllotaxis by polar

auxin transport. Nature **426**: 255–260.

Riou-Khamlichi, C., Menges, M., Healy, J.M., and Murray, J.A. (2000). Sugar control

of the plant cell cycle: differential regulation of Arabidopsis D-type cyclin gene

expression. Mol. Cell. Biol. **20**: 4513–4521.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics **26**: 139–140.

Robinson, M.D. and Oshlack, A. (2010). A scaling normalization method for

differential expression analysis of RNA-seq data. Genome Biol. **11**: R25.

Robinson, R. and Rick, C.M. (1954). New Tomato Seedling Characters and Their
Linkage Relationships.

Sachs, T. (1969). Regeneration experiments on the determination of the form of leaves.
Israel J Bot.

Sairanen, I., Novák, O., Pěnčík, A., Ikeda, Y., Jones, B., Sandberg, G., and Ljung,
K. (2012). Soluble carbohydrates regulate auxin biosynthesis via PIF proteins in
Arabidopsis. Plant Cell **24**: 4907–4916.

Scarpella, E., Barkoulas, M., and Tsiantis, M. (2010). Control of leaf and vein
development by auxin. Cold Spring Harb. Perspect. Biol. **2**: a001511.

Scarpella, E. and Helariutta, Y. (2010). Vascular pattern formation in plants. Curr. Top.
Dev. Biol. **91**: 221–265.

Scarpella, E., Marcos, D., Friml, J., and Berleth, T. (2006). Control of leaf vascular
patterning by polar auxin transport. Genes Dev. **20**: 1015–1027.

Shani, E., Ben-Gera, H., Shleizer-Burko, S., Burko, Y., Weiss, D., and Ori, N. (2010).
Cytokinin regulates compound leaf development in tomato. Plant Cell **22**: 3206–
3217.

Sinha, N. and Hake, S. (1994). The Knotted leaf blade is a mosaic of blade, sheath,
and auricle identities. Dev. Genet. **15**: 401–414.

Skirycz, A., Claeys, H., De Bodt, S., Oikawa, A., Shinoda, S., Andriankaja, M.,
Maleux, K., Eloy, N.B., Coppens, F., Yoo, S.-D., and Others (2011). Pause-and-
stop: the effects of osmotic stress on cell proliferation during early leaf development
in Arabidopsis and a role for ethylene signaling in cell cycle arrest. Plant Cell **23**:
1876–1888.

Stokes, M.E., Chattopadhyay, A., Wilkins, O., Nambara, E., and Campbell, M.M.
(2013). Interplay between sucrose and folate modulates auxin signaling in

Arabidopsis. Plant Physiol. **162**: 1552–1565.

Sugimoto-Shirasu, K. and Roberts, K. (2003). "Big it up": endoreduplication and cell-
size control in plants. Curr. Opin. Plant Biol. **6**: 544–553.

Sugimoto-Shirasu, K., Stacey, N.J., Corsar, J., Roberts, K., and McCann, M.C.
(2002). DNA topoisomerase VI is essential for endoreduplication in Arabidopsis.
Curr. Biol. **12**: 1782–1786.

Sussex, I.M. and Kerk, N.M. (2001). The evolution of plant architecture. Curr. Opin.
Plant Biol. **4**: 33–37.

Tamayo, P., Slonim, D., Mesirov, J., Zhu, Q., Kitareewan, S., Dmitrovsky, E.,
Lander, E.S., and Golub, T.R. (1999). Interpreting patterns of gene expression with
self-organizing maps: methods and application to hematopoietic differentiation.
Proc. Natl. Acad. Sci. U. S. A. **96**: 2907–2912.

Tavakol, E. et al. (2015). The barley Uniculme4 gene encodes a BLADE-ON-PETIOLE-
like protein that controls tillering and leaf patterning. Plant Physiol. **168**: 164–174.

Team, R.C. (2014). A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna.

Tindamanyire, J.M., Townsley, B., Kiggundu, A., Tushemereirwe, W., and Sinha,
N. (2013). Building a bi-directional promoter binary vector from the intergenic region
of Arabidopsis thaliana cab1 and cab2 divergent genes useful for plant
transformation. Afr. J. Biotechnol. **12**: 1203–1208.

Tsukaya, H. (2014). Comparative leaf development in angiosperms. Curr. Opin. Plant
Biol. **17**: 103–109.

Usadel, B., Bläsing, O.E., Gibon, Y., Retzlaff, K., Höhne, M., Günther, M., and Stitt,
M. (2008). Global transcript levels respond to small changes of the carbon status
during progressive exhaustion of carbohydrates in Arabidopsis rosettes. Plant
Physiol. **146**: 1834–1861.

Wehrens, R. and Buydens, L. (2007). Self- and Super-organizing Maps in R: The
kohonen Package. J. Stat. Softw. **21**: 1–19.

Whitewoods, C.D. and Coen, E. (2017). Growth and Development of Three-
Dimensional Plant Form. Curr. Biol. **27**: R910–R918.

Wickham, H. (2009). ggplot2: elegant graphics for data analysis (Springer Science &
Business Media).

Wind, J., Smeekens, S., and Hanson, J. (2010). Sucrose: metabolite and signaling
molecule. Phytochemistry **71**: 1610–1614.

Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G.V., and Provart, N.J.
(2007). An "Electronic Fluorescent Pictograph" Browser for Exploring and Analyzing
Large-Scale Biological Data Sets. PLoS ONE **2**: e718.

Wolf, S.D., Silk, W.K., and Plant, R.E. (1986). QUANTITATIVE PATTERNS OF LEAF
EXPANSION: COMPARISON OF NORMAL AND MALFORMED LEAF GROWTH
IN VITIS VINIFERA CV. RUBY RED. Am. J. Bot. **73**: 832–846.

Xu, C., Park, S.J., Van Eck, J., and Lippman, Z.B. (2016). Control of inflorescence
architecture in tomato by BTB/POZ transcriptional regulators. Genes Dev. **30**: 2048–
2061.

Yifhar, T., Pekker, I., Peled, D., Friedlander, G., Pistunov, A., Sabban, M.,
Wachsman, G., Alvarez, J.P., Amsellem, Z., and Eshed, Y. (2012). Failure of the
tomato trans-acting short interfering RNA program to regulate AUXIN RESPONSE
FACTOR3 and ARF4 underlies the wiry leaf syndrome. Plant Cell **24**: 3575–3589.

1238

1239

Figure 1 - Experimental set-up for sampling *S. lycopersicum* P4 leaves
(A) Transverse section from a wild-type apex showing leaf primordia P1-P5 in relation to the Shoot Apical Meristem (SAM). (B) Image of a wild-type apex during P4 leaf development. Images of transverse sections from the (C) top, (F) middle, and (G) base regions of a wild-type P4 leaf. Colors highlight the separation of the margin (lighter colors) and rachis (darker colors) along the top (purple), middle (brown), and base (green). Schematic diagram of a P4 leaf illustrating the six regions identified in (D) wild type and (E) *tf-2*. (H) Schematic diagram showing how the margin (gray) and rachis (blue) of a leaf were defined in this study. Images of leaves from wild type (I) and *tf-2* (J). Scale bars (A-E) = 100 µm and (I) and (J) = 5 mm.

Figure 2 – Characterization of the cell cycle using EdU fluorescence and flow cytometry

(A-F) Representative confocal images using EdU fluorescence to detect sites of cell division at the shoot apex of (A, C, E) wild type and (B, D, F) *tf-2* plants. Images in (C-D) show the top region of a P4 leaf, while (E-F) show the middle and base regions of a P4 leaf. Arrow in (E) points to clustering of EdU fluorescence suggesting sites of active cell division needed for leaflet initiation. (G) Bar graph showing DNA content peaks based on flow cytometry of leaf tissue. Tissue was collected from the oldest leaf of the plant at 8-90 days after germination. (H) Bar graph displaying the DNA content peaks from flow cytometry comparing leaf tissue sampled from the base (black) and tip (gray). Error bars show the standard deviation across at least three replicates at each time point. Scale bars = 100 μm.

A

B

C

Figure 3 - Pairwise differential gene expression between the rachis and margin in each region along the proximal-distal axis in a wild-type P4 leaf
(A) Graph summarizing representative enriched GO terms describing the significantly up-regulated genes in each region (top, mid, and base) from differential gene expression analyses performed on wild-type plants. Point size represents Benjamini-Hochberg (BH) corrected P-values in margin (gray) and rachis (blue) tissue. (B) Schematic diagram summarizing enriched GO terms of differentially up-regulated genes in each region of the P4 leaf. Colors highlight the separation of the margin (lighter colors) and rachis (darker colors) along the top (purple), middle (brown), and base (green). (C) Schematic diagram showing GO categories that help define each morphogenetic state along the leaf.

Figure 4 - Differential gene expression between the margin and rachis in a wild-type P4 leaf and chlorophyll A-B binding gene activity in the rachis compared to margin tissue during early leaf development

(A) Results of differential gene expression analysis in the wild type showing average log Counts Per Million (LogCPM) over log Fold Change (logFC) values. The number of significant differentially regulated genes (red) between margin and rachis tissue is shown. (B) Summary of GO terms describing up-regulated genes in each tissue, showing that rachis (blue) tissue is predominantly described by GO terms related to cell specialization compared to margin tissue (gray). (C) Normalized read count for chlorophyll A-B (CAB) binding genes in tomato (*SlCAB*). Colors highlight the separation of the margin (lighter colors) and rachis (darker colors) along the top (purple), middle (brown), and base (green). (D-G) pCAB::GUS expression showing photosynthetic activity during leaf development in tomato. pCAB::GUS is localized to the rachis of P4-P6 leaflets, illustrating differential regulation of *CAB* expression along the medio-lateral axis during early leaf development. pCAB::GUS is nearly ubiquitous in (G) P8 terminal leaflet. * p-value < 0.005 for significantly up-regulated genes in rachis tissue compared to the margin based on modeled differential expression analysis. Scale bars (D-E, G) = 1 mm, (F) = 100 μm.

Figure 5 - Top clusters identified by SOM analysis define each tissue region based on up-regulated genes
(A) Plotting of wild-type gene expression observed in the top 25% of genes based on coefficient of variation in the Principal Component (PC) space. (B) Projection of SOM cluster 4 onto the PC space explains one of the main clusters in the PC space (C) Codebook vector of a 2x3 SOM analysis showing the top six clusters (D) Gene expression patterns of Cluster 4 across the six tissue types. (C) and (D) Colors highlight the margin (lighter colors) and rachis (darker colors) along the top (purple), middle (brown), and base (green).

Figure 6 - Large SOM map describes a small gene cluster that defines margin identity
(A) Heatmap representing the gene expression patterns of the 36 gene clusters. The red box highlights clusters 10 and 11 (B) Heatmap of clusters 10 and 11, with genes that are up-regulated in the margin and down-regulated in the rachis. (C) Boxplot showing the gene expression patterns of cluster 10 and 11. See S5 Figure for full heatmap of all 36 clusters.

Figure 7 - Auxin visualization during leaflet initiation in wild type and *tf-2*
(A)-(D) Microscope images of apices from (A) and (B) wild type and (C) and (D) *tf-2*. (B) and (D) Fluorescence signals of PIN1::GFP (green) and chlorophyll autofluorescence (red) * marks the base marginal blastozone region. (B) shows clear PIN1::GFP signal in wild type along the entire margin of the P4 leaf, while in (D), *tf-2* has lost signal in the base marginal blastozone region. (E)-(H) DR5::Venus signal (green) observed by confocal microscopy. (E) and (F) wild type plant apices. (G) and (H) *tf-2* plants apices (F) and (H) close up on the site of leaflet initiation of the base margin region of P4 leaves. Scale bars = 100 μm.

Figure 8 - Differential gene expression analysis in base margin tissue between wild type and *tf-2* reveals BOP2 as a regulator that suppresses meristematic identity

(A) Schematic illustrating the regions (base margin) compared between wild type and *tf-2* using modeled differential gene expression analysis (B) Bar graph illustrating the expression patterns of *SlBOP2* across all six tissue types between wild type and *tf-2*, showing how *SlBOP2* is upregulated in the base margin only in *tf-2*. Colors highlight the separation of the margin (lighter colors) and rachis (darker colors) along the top (purple), middle (brown), and base (green). (C) - (E) *SlBOP2* CRISPR knockout line (*CR-slbop2*), which displays ectopic shoot apical meristems along the rachis of complex leaves. (F) *SlBOP2* genomic region. Black line shows the *TF-2* binding site 3kb upstream of *SlBOP2*. Scale bars (C) = 10 mm, (D) = 2 mm, (E) = 0.2 mm