1   **Flexible cobamide metabolism in *Clostridioides (Clostridium) difficile* 630 Δ*erm***
2
3   Amanda N. Shelton[1], Xun Lyu[2], Michiko E. Taga[1#]
4
5   [1]Department of Plant & Microbial Biology and [2]Department of Molecular & Cell Biology, University of California, Berkeley, Berkeley, CA 94720
6
7
8   #Correspondence to taga@berkeley.edu
9
Running title: B_{12} metabolism in *C. difficile*

Abstract

Clostridioides (Clostridium) difficile is an opportunistic pathogen known for its ability to

colonize the human gut under conditions of dysbiosis. Several aspects of its carbon and amino

acid metabolism have been investigated, but its cobamide (vitamin B_{12} and related cofactors)

metabolism remains largely unexplored. *C. difficile* has seven predicted cobamide-dependent

metabolisms encoded in its genome in addition to a nearly complete cobamide biosynthesis

pathway and a cobamide uptake system. To address the importance of cobamides to *C. difficile*,

we studied *C. difficile* 630 Δ*erm* and mutant derivatives under cobamide-dependent conditions *in*

vitro. Our results show that *C. difficile* can use a surprisingly diverse array of cobamides for

methionine and deoxyribonucleotide synthesis, and can use alternative metabolites or enzymes,

respectively, to bypass these cobamide-dependent processes. *C. difficile* 630 Δ*erm* produces the

cobamide pseudocobalamin when provided the early precursor 5-aminolevulinc acid or the late

intermediate cobinamide, and produces other cobamides if provided an alternative lower ligand.

The ability of *C. difficile* 630 Δ*erm* to take up cobamides and Cbi at micromolar or lower

concentrations requires the transporter BtuFCD. Genomic analysis revealed genetic variations in

in the *btuFCD* locus of different *C. difficile* strains, which may result in differences in the ability

to take up cobamides and Cbi. These results together demonstrate that, like other aspects of its

physiology, cobamide metabolism in *C. difficile* is versatile.

Importance

The ability of the opportunistic pathogen *Clostridioides difficile* to cause disease is closely linked

to its propensity to adapt to conditions created by dysbiosis of the human gut microbiota. The

cobamide (vitamin B_{12}) metabolism of *C. difficile* has been underexplored, though it has seven

metabolic pathways that are predicted to require cobamide-dependent enzymes. Here, we show

that *C. difficile* cobamide metabolism is versatile, as it can use a surprisingly wide variety of

cobamides and has alternative functions that can bypass some of its cobamide requirements.

Furthermore, *C. difficile* does not synthesize cobamides *de novo*, but produces them when given

cobamide precursors. Better understanding of *C. difficile* cobamide metabolism may lead to new

strategies to treat and prevent *C. difficile*-associated disease.

Introduction

The human gut microbiota is a complex community composed of hundreds to thousands of species of bacteria, archaea, and eukaryotic microbes (1). Members of this community compete for nutrients such as carbon sources, but also release metabolites that benefit other members. The exchange of B vitamins, particularly vitamin B_{12}, is thought to be prevalent in many environments because most bacteria lack the ability to synthesize some of the cofactors they require for enzyme catalysis (2–6), and instead must acquire them from other organisms (7). Such nutrient cross-feeding interactions can influence bacterial metabolism in ways that can affect not only the microbiota, but also host health (8, 9).

Clostridioides (Clostridium) difficile is a human intestinal pathogen that is among the most common causes of nosocomial infections, with nearly 300,000 healthcare-associated cases per year in the United States (10). *C. difficile* colonization of the gut is correlated with dysbiosis of the gut microbiota (11). Its abilities to germinate from spores, proliferate in the gut, and cause disease are impacted both positively and negatively by ecological and metabolic factors (12–14). The global alteration of the gut metabolome following antibiotic treatment is correlated with increased susceptibility to *C. difficile* infection, and recent work has linked changes in the relative abundance of specific metabolites to changes in the microbiome using model systems (11, 15–17). For example, succinate availability increases after disturbance of the microbiota, allowing *C. difficile* expansion in a mouse model (18). Additionally, specific commensal bacteria have been shown to produce compounds that stimulate *C. difficile* metabolism. In a bi-association, *Bacteroides thetaiotaomicron* can break down host mucin and produce sialic acid, which can be used by *C. difficile* for expansion in the gut (19). *C. difficile* can also induce other

members of the microbiota to produce indole, which is thought to create a more favorable

environment for the pathogen by inhibiting competing microbes (20).

Some interactions with microbiota members have also been shown to be inhibitory to *C.*

difficile. Co-culturing with certain *Bifidobacterium* spp. on particular carbon sources reduces *C.*

difficile toxin production relative to monoculture (21). While primary bile acids produced by the

host promote *C. difficile* spore germination, *Clostridium scindens* and other 7α-dehydroxylating

Clostridia transform these compounds into secondary bile acids, which are inhibitory to *C.*

difficile (22, 23). The latter example illustrates that compounds in the same class can have

different effects on the disease state. Given the complexity of metabolic interactions in the

mammalian gut, many additional microbial metabolites likely influence the ability of *C. difficile*

to colonize and persist in the gut.

One class of metabolites that has not been explored for its ability to affect *C. difficile*

growth and virulence is cobamides, the vitamin B_{12} (also called cobalamin) family of cofactors.

Cobamides are used in diverse microbial metabolisms including methionine synthesis,

deoxyribonucleotide synthesis, acetogenesis, and some carbon catabolism pathways. These

reactions are facilitated by fission of the Co-C bond to the cobamide upper ligand, which can be

a 5'-deoxyadenosyl group for radical reactions, a methyl group for methyltransferase reactions,

or a cyano group in the inactive vitamin form (24) (labeled as "R" in Fig. 1A). Over 80% of all

sequenced bacteria (25–27) and 80% of sequenced human gut bacteria (2, 28, 29) have one or

more cobamide-dependent enzymes, suggesting that cobamides are widely used cofactors across

microbial ecosystems. Strikingly, fewer than 40% of bacterial species are predicted to produce

cobamides *de novo* (2, 25–28), and therefore over half of bacteria that use cobamides must

acquire them from their environment. Cobamides vary in the structure of the lower ligand (Fig.

1A, B), and organisms studied to date are selective in which cobamides they can use (28, 30–37).

Seven cobamides in addition to the cobamide precursor cobinamide (Cbi, Fig. 1C) have been

detected in the human gut (38). In an environment with plentiful, diverse cobamides and

cobamide precursors, a microbial species that requires a particular cobamide can either import

that cobamide, synthesize it *de novo,* chemically remodel available cobamides to the preferred

structure, or alter its need for the cobamide by using alternative pathways (8, 39).

The seven predicted cobamide-dependent enzymes encoded in the *C. difficile* genome are

involved in methionine synthesis, nucleotide metabolism, and carbon metabolism (Fig. 2). When

grown with amino acids and glucose as carbon and energy sources *in vitro*, *C. difficile* does not

require cobalamin supplementation (40). However, in model infection systems, cobamide-

dependent metabolism may be important for virulence and growth. For example, access to

ethanolamine catabolism may be important in modulating virulence, as deletion of EutA, the

reactivating factor required for activity of the cobamide-dependent ethanolamine ammonia lyase

(EutBC), in *C. difficile* strain 630 Δ*erm* reduces the mean time to morbidity in a hamster model

(41). Additionally, metabolic models and transcriptomics (42, 43) suggest that the cobamide-

dependent Wood-Ljungdahl carbon fixation pathway is an important electron sink, and an

experimental study suggests that it may be used for autotrophic growth by some *C. difficile*

strains (44).

The observation that *C. difficile* can grow without added cobamides *in vitro* (40) suggests

that it may not require cobamides under those conditions, or that it can biosynthesize cobamides.

However, all sequenced strains of *C. difficile* are missing *hemA* and *hemL*, the first two enzymes

in the cobamide biosynthesis pathway required for production of the precursor 5-aminolevulinic

acid (ALA) (45) (Fig. 1C). Therefore, *C. difficile* is predicted to be able to produce a cobamide

only when ALA is available, as has been observed in three other bacteria (25) (Fig. 2). In order

to use cobamide-dependent metabolisms, we predict that *C. difficile* requires cobamides or

precursors such as ALA from the gut. While ALA is an intermediate made in all tetrapyrrole-

producing organisms, including the host, cobamides are only produced by some bacteria and

archaea (46).

To address the importance of cobamides for *C. difficile* metabolism and to understand

how *C. difficile* acquires cobamides, we examined *C. difficile* 630 Δ*erm* and mutant derivatives

in vitro under cobamide-dependent conditions. We found that the bacterium can use a

surprisingly diverse array of cobamides for methionine and deoxyribonucleotide synthesis, and

can use alternative nutrient sources or enzymes to fulfill its metabolic needs. In addition to

importing and using a variety of cobamides, when provided with ALA or the late intermediate

Cbi, *C. difficile* 630 Δ*erm* can produce the cobamide pseudocobalamin, and can produce other

cobamides if provided an alternative lower ligand. Together, these results show that *C. difficile* is

versatile in its cobamide metabolism.

Results

***C. difficile* requires methionine or a cobamide for growth**

To investigate cobamide-dependent metabolism in the model *C. difficile* strain 630 Δ*erm*,

we sought to culture the organism in conditions that require specific cobamide-dependent

enzymes. The *C. difficile* genome encodes the cobalamin-dependent methionine synthase MetH,

but does not contain the cobalamin-independent alternative enzyme MetE. The absence of a

complete cobamide biosynthesis pathway suggests that *C. difficile* requires either methionine or

a cobamide in its growth medium. Previously, methionine was classified as a "growth-

enhancing," but not essential, amino acid in a medium containing cyanocobalamin (vitamin B_{12})

for seven of eight strains tested (40, 47). To test whether *C. difficile* can use cobamides for

methionine synthesis and to identify the specific cobamides that support its MetH-dependent

growth, we cultured *C. difficile* in a defined medium lacking methionine with a range of

concentrations of cyanocobalamin, Cbi, and eight other cyanylated cobamides that we purified.

C. difficile was unable to grow in this medium without cobamide or methionine addition (Fig.

3A), suggesting that, as predicted, it cannot produce cobamides *de novo* to support the activity of

MetH. Remarkably, unlike other bacteria that have been reported to use a limited number of

cobamides for methionine synthase activity (28, 48, 49), all of the cobamides and Cbi were able

to confer high growth yields to *C. difficile* at concentrations as low as 1 nM (Fig. 3A).

Methionine addition also supported growth, though higher concentrations were required than for

cobamides (Fig. 3B). We also observed robust growth with addition of ALA (Fig. 3C).

***C. difficile* growth with ribonucleotide reductase NrdJ requires a more restricted set of**

cobamides

C. difficile genomes encode homologs of the cobalamin-dependent (class II)

ribonucleotide reductase (RNR) (*nrdJ,* CDIF630erm_RS07280), as well as two cobalamin-

independent RNRs: an oxygen-dependent (class I) RNR (encoded by *nrdE*,

CDIF630erm_RS16325, and *nrdF*, CDIF630erm_RS16320) and an oxygen-sensitive (class III)

RNR (*nrdD*, CDIF630erm_RS00990 and *nrdG*, CDIF630erm_RS00995). In principle, any of

these three isozymes could be used for deoxyribonucleotide synthesis from ribonucleotides,

although under anaerobic conditions only the class II and class III RNRs are expected to

function. Cobamide addition is not required for anaerobic growth of the parent strain *C. difficile*

630 Δ*erm* Δ*pyrE* in a casamino acid medium (CDDM) with glucose, and adding cobamides or

cobamide precursors did not affect growth yield (Supplemental Fig. 1), suggesting that the class

III RNR, NrdDG, is functional under these conditions. To test whether the class II RNR, NrdJ, is

functional, we deleted the *nrdD* and *nrdG* genes while providing exogenous cobalamin, using the

allelic exchange system in a Δ*pyrE* background (50). This strain could grow only with cobalamin

addition, suggesting that NrdJ is functional and NrdEF is not under these growth conditions (Fig.

4A). To determine which cobamides it requires, the Δ*nrdDG* strain was grown with the same

cobamides and precursors as in Fig. 3. In contrast to MetH, NrdJ is more selective in the

cobamides it can use (Fig. 4A), as expected based on studies with other class II RNRs (33, 36,

51, 52). There was little growth with [Cre]Cba, [Phe]Cba, and [5-OHBza]Cba (Fig. 4A).

Addition of ALA also supported NrdJ-dependent growth (Fig. 4B).

***C. difficile* produces pseudocobalamin from the precursor ALA via the *cbi* genes**

The observation that *C. difficile* could grow under cobamide-dependent conditions with

ALA or Cbi (Fig. 3A, C, Fig. 4) suggests that it can produce a cobamide from these precursors

using the cobamide biosynthetic genes encoded in its genome (25). To test this prediction, the

corrinoid fraction, which includes cobamides and late cobamide precursors including Cbi, was

extracted from the cell pellets of *C. difficile* 630 Δ*erm* grown with either ALA or Cbi. Consistent

with our predictions, HPLC analysis of the extracted corrinoids showed that *C. difficile* produced

a cobamide only when ALA or Cbi was added (Fig. 5A). Additionally, as predicted, corrinoid

analysis of a strain lacking the corrin ring biosynthesis genes *cbiKLJHGFTEDC* demonstrated

that these genes are necessary for cobamide synthesis from ALA, but not Cbi (Fig. 5A). Because

C. difficile lacks all known genes for biosynthesis of benzimidazoles and attachment of phenolic

lower ligands, it is predicted to be incapable of producing benzimidazolyl or phenolyl

cobamides, but may produce a purinyl cobamide (49, 53–59). Indeed, the major cobamide

present in *C. difficile* corrinoid extracts co-eluted with the purinyl cobamide pseudocobalamin

(Fig. 5A). The UV-Vis spectrum of the major cobamide was consistent with a pseudocobalamin

standard (Supplemental Fig. 2C). Mass spectrometry analysis verified that the major cobamide

extracted from cultures grown with ALA is pseudocobalamin (Supplemental Fig. 2A, B).

C. difficile can perform guided biosynthesis but does not remodel cobamides

Some bacteria can perform guided biosynthesis, a process in which an exogenously

provided, non-native lower ligand base is incorporated into a cobamide (32, 36, 48, 60, 61). To

test if *C. difficile* is capable of guided biosynthesis to produce cobamides other than its native

pseudocobalamin, either DMB (the lower ligand of cobalamin, Fig. 1A) or a related compound,

benzimidazole (Bza, Fig. 1B) was added to cultures containing either ALA or Cbi. Analysis of

corrinoid extracts showed that *C. difficile* could attach either of these exogenous lower ligands to

form cobalamin and [Bza]Cba, respectively, with both precursors (Fig. 5B). A small amount of

pseudocobalamin was also recovered in cultures containing Cbi with Bza (Fig. 5B).

Some bacteria and archaea are able to remodel cobamides by removing the lower ligand

and nucleotide loop with the amidohydrolase enzyme CbiZ and rebuilding the cobamide with a

different lower ligand (31, 62–64). We were unable to identify a *cbiZ* homolog in the *C. difficile*

genome, and accordingly, we did not observe evidence of remodeling; when cobalamin, [2-

MeAde]Cba, or [Cre]Cba was provided to *C. difficile*, the same cobamides were recovered from

the cells (Fig. 6A).

C. difficile requires *btuFCD* for efficient uptake of cobamides and Cbi

The presence of cobamides in the cellular fraction of cultures grown with either Cbi or a

cobamide at nanomolar concentrations (Fig. 5, 6A) suggested that *C. difficile* takes up Cbi and

cobamides via an active transporter. We identified a candidate cobalamin uptake operon

(*btuFCD*) downstream of a sequence annotated as a cobalamin riboswitch, suggesting that these

genes function in corrinoid import (27, 28, 65–70). No corrinoids could be detected in the

cellular fraction of the Δ*btuFCD* mutant grown with 10 nM Cbi or cobalamin (Fig. 6A). In

contrast, ALA uptake is apparently unaffected in the Δ*btuFCD* mutant, as pseudocobalamin can

be recovered from the cellular fraction when ALA is provided (Fig. 6A). Furthermore, the

Δ*btuFCD* mutant grew poorly in methionine-free medium even when Cbi or cobalamin was

added at concentrations 10^3 to 10^4-fold higher than required for growth of the parental strain

(Fig. 6B). The ability of methionine or ALA to support growth remained unaffected by the

Δ*btuFCD* mutation (Fig. 6B). Interestingly, genomic analysis identified strains of *C. difficile* that

contain a *tlpB* transposon insertion in *btuC*, likely rendering the BtuFCD transporter

nonfunctional (Supplemental Fig. 3) (71). Of the genomes analyzed, the *tlpB* insertion in this

locus appears to be restricted to strains in the PCR-ribotype 027 (RT027) clade, including the

hypervirulent strain R20291, based on a multi-locus sequence typing (MLST) tree of *C. difficile*

strains (Supplemental Fig. 3, red labels). This observation suggests that unlike strain 630 Δ*erm*

examined in this study, members of the RT027 clade may be unable to take up cobamides and

Cbi efficiently.

Discussion

The potential of *C. difficile* to cause disease is closely linked to its ability to fill

ecological niches made available by gut microbiota dysbiosis (13), using a suite of metabolic

pathways to make use of newly available nutrient sources. *C. difficile* has an unusually high

number of cobamide-dependent metabolisms encoded in its genome (25), but their functions

have been underexplored. Here, we show that *C. difficile* is able to use many cobamides and

cobamide precursors in two of its seven cobamide-dependent pathways. The promiscuous use of

cobamides and the ability to bypass these cobamide-dependent metabolisms highlights the

metabolic flexibility of *C. difficile.*

The cobalamin-dependent methionine synthase, MetH, is the most abundant cobamide-

dependent enzyme in bacterial genomes (25) and is found in numerous organisms in all three

domains of life, including humans (24). Compared to the majority of other MetH homologs that

have been studied, our results indicate that the *C. difficile* MetH homolog is unusually

promiscuous in its cobamide selectivity. For example, several eukaryotic algae grew robustly

under MetH-dependent conditions with cobalamin, but did not grow with pseudocobalamin at

the same concentrations (33). The human gut commensal bacterium *Bacteroides*

thetaiotaomicron could use benzimidazolyl and purinyl cobamides for MetH-dependent growth,

but could not use phenolyl cobamides (28). An example of MetH selectivity *in vitro* was in

Spirulina platensis, where the purified enzyme bound its native cobamide, pseudocobalamin,

with a higher affinity than cobalamin (72). An exception to this observed selectivity is another

gut pathogen, *Salmonella enterica*, which can use its native cobamide, pseudocobalamin, in

addition to cobalamin, [Phe]Cba, and [Cre]Cba for MetH-dependent growth, although other

cobamides were not tested (48, 49). The versatility of *C. difficile*'s cobamide use is notable given

the diversity of cobamides that have been detected in the gut (38).

In contrast to MetH, our growth experiments indicate that the selectivity of *C. difficile*'s

NrdJ is more similar to that of other organisms that rely on NrdJ for growth. For example,

Sinorhizobium meliloti was unable to grow with [Cre]Cba and grew poorly with

pseudocobalamin relative to its native cobamide, cobalamin (36); *Lactobacillus leichmannii*

could only use benzimidazolyl or purinyl cobamides (51); and *Euglena gracilis* grew well with

cobalamin and [Bza]Cba and poorly with pseudocobalamin, [5-OHBza]Cba, and [Cre]Cba (33,

52). Unlike MetH, the NrdJ enzyme requires cobamides that can adopt the "base-on"

configuration in which the lower ligand base is coordinated to the cobalt ion throughout the

catalytic cycle (24). Phenolyl cobamides are unable to adopt the base-on configuration, so their

inability to support growth in the NrdJ-dependent condition was expected. *C. difficile* 630 Δ*erm*

also contains an active class III cobamide-independent RNR, NrdDG, which may be an

important strategy to maintain deoxyribonucleotide synthesis when cobamides are scarce.

However, in other species, under certain conditions the class II RNR provides an advantage over

other RNR classes, such as during oxidative stress (73), although the conditions where NrdJ

would provide an advantage for *C. difficile* have yet to be uncovered.

Seven different cobamides and the precursor Cbi have been detected in human feces (38).

In stool samples of individuals not taking cobalamin supplements, the average total corrinoid

present is approximately 1300 ng per gram feces, roughly equivalent to 1 μM (38). Cbi is found

at tens of ng per gram (38). Growth experiments under MetH and NrdJ-dependent conditions

showed that *C. difficile* 630 Δ*erm* reached maximum growth yield with as little as 1 nM

cobamide or Cbi (Fig 3, 4). Based on the absence of corrinoids in the cellular fraction of a 630

Δ*erm* Δ*pyrE* Δ*btuFCD* strain (Fig. 6), we infer that strains with an insertion in *btuC*

(Supplemental Fig. 3)*,* including the hypervirulent R20291 and CD196 strains (71), would

require cobamides or Cbi at extracellular concentrations higher than 100 μM if relying on

cobamide-dependent enzymes. This suggests that these strains may not be able to use the

cobamides or Cbi present in the gut.

Our results show that not only is *C. difficile* able to use multiple cobamides to support its

metabolism, but it can also use the early precursor ALA to produce pseudocobalamin. The ability

to use ALA to produce a cobamide, and thus not strictly rely on cobamide or Cbi uptake, could

be important to strains with a transposon insertion in the *btuC* gene (70, 74). ALA concentrations

in the human gut have not been reported. However, we speculate that, similar to cobamides and

Cbi, ALA and possibly other early cobamide precursors could be provided by other members of

the microbiota. Alternatively, ALA could be provided by the host either through the diet or via

biosynthesis of heme, which also uses ALA as a precursor. Other commensal gut microbiota

have been reported to be able to salvage ALA (25), suggesting that ALA could be available in

the gut.

C. difficile is also able to incorporate non-native lower ligands to form benzimidazolyl

cobamides (guided biosynthesis). Free benzimidazole bases have been found in animal

gastrointestinal tracts such as rumen fluid and termite guts (75), but benzimidazole levels in the

human gut have not been measured. The cobamides used by *C. difficile* could therefore also vary

with the presence of different benzimidazole-producing organisms in the microbiota. Our results

show that pseudocobalamin and most benzimidazolyl cobamides support growth of *C. difficile*

equally for the two metabolisms we investigated in this study, but the cobamide preferences of

the other five cobamide-dependent metabolisms have not been investigated.

We have identified cobamides and precursors that *C. difficile* can use *in vitro,* but which

cobamides or cobamide precursors it predominantly uses in the gut remains to be discovered.

Evidence from transcriptomics is ambiguous with respect to expression of genes encoding

cobamide-dependent enzymes or cobamide biosynthesis during infection, likely due to

differences in study design (15, 43, 76–78). Since both diet and the microbiota can contribute to

the cobamide profile in the gut (38, 79, 80), the availability of cobamides may vary significantly

across infection systems and affect the expression and use of cobamide biosynthesis and

cobamide-dependent pathways by *C. difficile*. In one study, *hemB*, which encodes the enzyme

that converts ALA to the next intermediate, porphobilinogen, was among the most highly

expressed genes in *C. difficile* strain VPI 104363 in a mouse model (43), suggesting that *C.*

difficile produces cobamides from ALA in the gut. How the cobamide content in the gut

environment changes during *C. difficile* infection is unknown, but since much of the cobamide

content in the lower gastrointestinal tract is produced by resident gut microbes (79, 80), it is

possible that cobamide abundances change during dysbiosis. Further *in vivo* studies are needed to

determine the extent to which cobamide metabolism is important to *C. difficile* associated

disease.

Materials and Methods

Bacterial strains and growth conditions

C. difficile 630 Δ*erm*, an erythromycin-sensitive derivative of the isolate 630 (81), and *C.*

difficile 630 Δ*erm* Δ*pyrE*, a derivative of 630 Δ*erm* with a uracil auxotrophy (50), were streaked

from frozen stocks onto BHIS agar (82) before being transferred to *Clostridium difficile* defined

medium (CDDM) containing casamino acids (83) and 8 g/L glucose. Agar plates and 96-well

plates containing liquid cultures were incubated at 37ºC in an anaerobic chamber (Coy Labs)

containing 10% H_2, 10% CO_2, and 80% N_2. For *C. difficile* 630 Δ*erm* Δ*pyrE* and derived strains,

5 µg/ml uracil was included in all defined media. For corrinoid extractions and NrdJ phenotype

experiments, strains were cultured in CDDM plus 8 g/L glucose. For MetH phenotype

experiments, CDDMK medium plus 8 g/L glucose without methionine was used. CDDMK

contains the same salts, trace metals, and vitamins as CDDM, but the casamino acids, tryptophan

and cysteine are replaced with the individual amino acids as follows: 100 mg/L histidine, 100

mg/L tryptophan, 100 mg/L glycine, 100 mg/L tyrosine, 200 mg/L arginine, 200 mg/L

phenylalanine, 200 mg/L threonine, 200 mg/L alanine, 300 mg/L lysine, 300 mg/L serine, 300

mg/L valine, 300 mg/L isoleucine, 300 mg/L aspartic acid, 400 mg/L leucine, 500 mg/L

cysteine, 600 mg/L proline, 900 mg/L glutamic acid (40). All liquid defined media were

prepared by boiling under 80% N_2/20% CO_2 gas. After the pH stabilized between 6.8 and 7.2,

the medium was dispensed into stoppered tubes and autoclaved. Filter-sterilized glucose and

vitamins were added after autoclaving. Cultures in stoppered tubes were incubated at 37ºC.

For MetH phenotype assays, *C. difficile* 630 Δ*erm* was grown in CDDM, then washed

twice in CDDMK without methionine prior to inoculation in CDDMK at an optical density

(O.D.$_{600}$) of 0.01 in a 96-well plate. For NrdJ phenotype assays, *C. difficile* 630 Δ*erm* Δ*pyrE*

Δ*nrdDG* was grown in CDDM with 5 µg/ml uracil and 10 nM cyanocobalamin, and washed

three times in CDDM medium without cyanocobalamin prior to inoculation in CDDM at an

O.D.$_{600}$ of 0.01 in a 96 well plate. O.D.$_{600}$ was measured on a BioTek Synergy 2 plate reader after

23 to 24 hours of growth.

ALA, Cbi and cyanocobalamin were purchased from Sigma Aldrich. Other cobamides

were purified from bacterial cultures as described in Men *et al.* (84)

Strain and plasmid construction

The allelic coupled exchange (ACE) system of Ng *et al.* was used for construction of *C.*

difficile mutant strains (50). Briefly, 500-1000 bp sequences flanking the target gene(s) (arms of

homology) in the *C. difficile* 630 Δ*erm* genome (CP016318,

https://www.ncbi.nlm.nih.gov/nuccore/CP016318.1/) were amplified by PCR (Supplemental

Table 1) and then were cloned into pMTL-YN3 (Chain Biotech) by Gibson assembly (85) in *E.*

coli XL1-Blue. Plasmid inserts were sequenced by Sanger sequencing before transformation of

the plasmid into *E. coli* CA434 (Chain Biotech). Conjugation of *E. coli* CA434 and *C. difficile*

630 Δ*erm* Δ*pyrE* was performed as described (86), except that *C. difficile* and *E. coli* were each

cultured for 5-8 hours prior to pelleting *E. coli* and mixing with the *C. difficile* recipient. After 16

hours growth on BHIS agar, the mixed cells were resuspended in 1 ml PBS, and 100 µl of the

suspension was plated on each of 5-7 plates of BHIS agar with added 10 µg/ml thiamphenicol,

250 µg/ml D-cycloserine, and 16 µg/ml cefoxitin. Colonies were purified at least twice by

streaking onto BHIS with 15 µg/ml thiamphenicol, 250 µg/ml D-cycloserine, and 16 µg/ml

cefoxitin, before counterselection on CDDM agar supplemented with 2 mg/L 5-fluoroorotic acid

(5-FOA) and 5 µg/ml uracil. The resulting colonies were purified by streaking at least twice on

the counterselection medium prior to screening by colony PCR for the deletion and the presence

of the *C. difficile* toxin gene *tcdB* (86). For the deletion of *nrdDG,* 10 nM cobalamin was added

to all media during the ACE procedure.

Corrinoid extraction and analysis

C. difficile was grown in 50 ml CDDM plus 8 g/L glucose under 80% N_2/20% CO_2

headspace for 16-22 hours at 37ºC prior to corrinoid extraction. Two cultures were combined for

each condition for a total volume of 100 ml for each extraction. Corrinoid extractions were

performed as described (31), except that cell pellets were autoclaved for 35 minutes and cooled

prior to addition of methanol and potassium cyanide. Two or more biological replicates were

performed for each condition.

High-performance liquid chromatography (HPLC) analysis was performed with an

Agilent Series 1200 system (Agilent Technologies, Santa Clara, CA) equipped with a diode array

detector with detection wavelengths set at 360 and 525 nm. For Fig. 5B and 6A samples were

injected onto an Agilent Zorbax SB-Aq column (5 μm, 4.6 × 150 mm) at 30°C, with 1 mL/min

flow rate. Compounds in the samples were separated with a gradient of 25 to 34% acidified

methanol in acidified water (containing 0.1% formic acid) over 11 min, followed by a 34 to 50%

gradient over 2 min, and 50 to 75% over 9 min. For Fig. 5A, samples were injected onto an

Agilent Eclipse Plus C18 column (5 μm, 9.6 × 250 mm) at 30 ºC, with 2 mL/min flow rate.

Compounds in the samples were separated with a gradient of 10 to 42% acidified methanol in

acidified water over 20 min. The amount of standards that were injected were as follows: Cbi (**1**),

200 pmol; pseudocobalamin (**2**), 225 pmol; cobalamin (**3**), 50 pmol; [Bza]Cba (**4**), 114 pmol; [2-

MeAde]Cba (**5**), 114 pmol; [Cre]Cba (**6**), 151 pmol. 5% to 20% by volume *C. difficile* samples

were injected.

Supplemental material

Supplemental methods, figures, and tables are provided.

Acknowledgements

This work was supported by grants NIH DP2AI117984 and R01GM114535 to M.E.T.

We thank Aimee Shen for her extensive advice and protocols for working with *C. difficile*, Craig Ellermeier for advice on working with *C. difficile,* and John Taylor, Gary Andersen, Christian Sieber, and Eric Dubinsky for advice on creating the phylogenetic tree. *C. difficile* 630 Δ*erm* and 630 Δ*erm* Δ*pyrE* were provided to us by Andrew Hryckowian with permission from Nigel Minton. We thank Olga Sokolovkaya and Kenny Mok for help with HPLC analysis and Dr. Rita Nichiporuk at the QB3/Chemistry Mass Spectrometry Facility at UC Berkeley for performing the mass specrometry analysis. We thank Luis Valentin-Alvarado for helpful discussions about *C. difficle* metabolism and genetic tools. We also thank Ellen Simms, Olga Sokolovskaya, Sebastian Gude, Gordon Pherribo, and Kenny Mok for critical reading of this manuscript.

Author Contributions

A.N.S. performed growth experiments, corrinoid extractions, and phylogenetic analysis. X.L. and A.N.S. created the mutant strains. A.N.S. and M.E.T. wrote the manuscript.

References

1. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. 2012. Diversity, stability and resilience of the human gut microbiota. Nature 489:220–230.
2. Magnúsdóttir S, Ravcheev D, De Crécy-Lagard V, Thiele I. 2015. Systematic genome assessment of B-vitamin biosynthesis suggests cooperation among gut microbes. Front Genet 6:148.
3. D'Souza G, Waschina S, Pande S, Bohl K, Kaleta C, Kost C. 2014. Less is more: Selective advantages can explain the prevalent loss of biosynthetic genes in bacteria. Evolution (N Y) 68:2559–2570.
4. Zengler K, Zaramela LS. 2018. The social network of microorganisms - How auxotrophies shape complex communities. Nat Rev Microbiol 16:383–390.
5. Garcia SL, Buck M, McMahon KD, Grossart HP, Eiler A, Warnecke F. 2015. Auxotrophy and intrapopulation complementary in the interactome of a cultivated freshwater model

community. Mol Ecol 24:4449–4459.

6.   Hubalek V, Buck M, Tan B, Foght J, Wendeberg A, Berry D, Bertilsson S, Eiler A. 2017.
Vitamin and Amino Acid Auxotrophy in Anaerobic Consortia Operating under
Methanogenic Conditions. mSystems 2:e00038-17.

7.   Belzer C, Chia LW, Aalvink S, Chamlagain B, Piironen V, Knol J, de Vos WM. 2017.
Microbial Metabolic Networks at the Mucus Layer Lead to Diet-Independent Butyrate and
Vitamin B 12 Production by Intestinal Symbionts. MBio 8:1–14.

8.   Seth EC, Taga ME. 2014. Nutrient cross-feeding in the microbial world. Front Microbiol
5:350.

9.   Lee WJ, Hase K. 2014. Gut microbiota-generated metabolites in animal health and
disease. Nat Chem Biol 10:416–424.

10.   Lessa FC, Mu Y, Bamberg WM, Beldavs ZG, Dumyati GK, Dunn JR, Farley MM,
Holzbauer SM, Meek JI, Phipps EC, Wilson LE, Winston LG, Cohen JA, Limbago BM,
Fridkin SK, Gerding DN, McDonald LC. 2015. Burden of Clostridium difficile infection
in the United States. N Engl J Med 372:825–34.

11.   Theriot CM, Koenigsknecht MJ, Carlson PE, Hatton GE, Nelson AM, Li B, Huffnagle
GB, Z Li J, Young VB. 2014. Antibiotic-induced shifts in the mouse gut microbiome and
metabolome increase susceptibility to Clostridium difficile infection. Nat Commun
5:3114.

12.   Theriot CM, Young VB. 2014. Microbial and metabolic interactions between the
gastrointestinal tract and Clostridium difficile infection. Gut Microbes 5:86–95.

13.   Hryckowian AJ, Pruss KM, Sonnenburg JL. 2017. The emerging metabolic view of
Clostridium difficile pathogenesis. Curr Opin Microbiol 35:42–47.

14.   Libertucci J, Young VB. 2019. The role of the microbiota in infectious diseases. Nat
Microbiol 4:35–45.

15.   Jenior ML, Leslie JL, Young VB, Schloss PD. 2017. *Clostridium difficile* Colonizes
Alternative Nutrient Niches during Infection across Distinct Murine Gut Microbiomes.
mSystems 2:e00063-17.

16.   Jenior ML, Leslie JL, Young VB, Schloss PD. 2018. Clostridium difficile Alters the
Structure and Metabolism of Distinct Cecal Microbiomes during Initial Infection To
Promote Sustained Colonization. mSphere 3:1–15.

17.   Carlucci C, Jones CS, Oliphant K, Yen S, Daigneault M, Carriero C, Robinson A, Petrof
EO, Weese JS, Allen-Vercoe E. 2019. Effects of defined gut microbial ecosystem
components on virulence determinants of Clostridioides difficile. Sci Rep 9:1–11.

18.   Ferreyra JA, Wu KJ, Hryckowian AJ, Bouley DM, Weimer BC, Sonnenburg JL. 2014.
Gut microbiota-produced succinate promotes C. difficile infection after antibiotic
treatment or motility disturbance. Cell Host Microbe 16:770–777.

19.   Ng KM, Ferreyra J a, Higginbottom SK, Lynch JB, Kashyap PC, Gopinath S, Naidu N,
Choudhury B, Weimer BC, Monack DM, Sonnenburg JL. 2013. Microbiota-liberated host
sugars facilitate post-antibiotic expansion of enteric pathogens. Nature 502:96–9.

20.   Darkoh C, Plants-Paris K, Bishoff D, DuPont HL. 2019. Clostridium difficile Modulates
the Gut Microbiota by Inducing the Production of Indole, an Interkingdom Signaling and
Antimicrobial Molecule. mSystems 4:e00346-18.

21.   Valdés-Varela L, Hernández-Barranco AM, Ruas-Madiedo P, Gueimonde M. 2016. Effect
of Bifidobacterium upon Clostridium difficile growth and toxicity when co-cultured in
different prebiotic substrates. Front Microbiol 7:1–9.

22.   Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H,
Kinnebrew M, Viale A, Littmann E, van den Brink MRM, Jenq RR, Taur Y, Sander C,
Cross JR, Toussaint NC, Xavier JB, Pamer EG. 2014. Precision microbiome reconstitution
restores bile acid mediated resistance to Clostridium difficile. Nature 517:205–208.

23.   Kang JD, Myers CJ, Harris SC, Kakiyama G, Lee IK, Yun BS, Matsuzaki K, Furukawa
M, Min HK, Bajaj JS, Zhou H, Hylemon PB. 2019. Bile Acid 7α-Dehydroxylating Gut
Bacteria Secrete Antibiotics that Inhibit Clostridium difficile: Role of Secondary Bile
Acids. Cell Chem Biol 26:27-34.e4.

24.   Banerjee R, Ragsdale SW. 2003. The Many Faces of Vitamin B 12 : Catalysis by
Cobalamin-Dependent Enzymes. Annu Rev Biochem 72:209–247.

25.   Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, Taga ME. 2019. Uneven
distribution of cobamide biosynthesis and dependence in bacteria predicted by
comparative genomics. ISME J 13:789–804.

26.   Rodionov DA, Vitreschak AG, Mironov AA, Gelfand MS. 2003. Comparative Genomics
of the Vitamin B12 Metabolism and Regulation in Prokaryotes. J Biol Chem 278:41148–
41159.

27.   Zhang Y, Rodionov DA, Gelfand MS, Gladyshev VN. 2009. Comparative genomic
analyses of nickel, cobalt and vitamin B12 utilization. BMC Genomics 10:78.

28.   Degnan PH, Barry NA, Mok KC, Taga ME, Goodman AL. 2014. Human gut microbes
use multiple transporters to distinguish vitamin B 12 analogs and compete in the gut. Cell
Host Microbe 15:47–57.

29.   Rodionov DA, Arzamasov AA, Khoroshkin MS, Iablokov SN, Leyn SA, Peterson SN,
Novichkov PS, Osterman AL. 2019. Micronutrient Requirements and Sharing Capabilities
of the Human Gut Microbiome. Front Microbiol 10:1–22.

30.   Yan J, Ritalahti KM, Wagner DD, Löffler FE. 2012. Unexpected specificity of
interspecies cobamide transfer from Geobacter spp. to organohalide-respiring
Dehalococcoides mccartyi strains. Appl Environ Microbiol 78:6630–6636.

31.   Yi S, Seth E, Men Y, Stabler SP, Allen RH, Alvarez-Cohen L, Taga ME. 2012. Versatility
in corrinoid salvaging and remodeling pathways supports corrinoid-dependent metabolism
in Dehalococcoides mccartyi. Appl Environ Microbiol 78:7745–7752.

32.   Mok KC, Taga ME. 2013. Growth inhibition of Sporomusa ovata by incorporation of
benzimidazole bases into cobamides. J Bacteriol 195:1902–11.

33.   Helliwell KE, Lawrence AD, Holzer A, Kudahl UJ, Sasso S, Kräutler B, Scanlan DJ,
Warren MJ, Smith AG. 2016. Cyanobacteria and Eukaryotic Algae Use Different
Chemical Variants of Vitamin B12. Curr Biol 26:999–1008.

34.   Yan J, Şimşir B, Farmer AT, Bi M, Yang Y, Campagna SR, Löffler FE. 2016. The
corrinoid cofactor of reductive dehalogenases affects dechlorination rates and extents in
organohalide-respiring Dehalococcoides mccartyi. ISME J 10:1092–1101.

35.   Keller S, Kunze C, Bommer M, Paetz C, Menezes RC, Svatoš A, Dobbek H, Schubert T.
2018. Selective Utilization of Benzimidazolyl-Norcobamides as Cofactors by the
Tetrachloroethene Reductive Dehalogenase of Sulfurospirillum multivorans. J Bacteriol
200:e00584-17.

36.   Crofts TS, Seth EC, Hazra AB, Taga ME. 2013. Cobamide structure depends on both
lower ligand availability and CobT substrate specificity. Chem Biol 20:1265–1274.

37.   Sokolovskaya OM, Mok KC, Duk Park J, Tran JLA, Quanstrom KA, Taga ME. 2019.
Cofactor selectivity in methylmalonyl-CoA mutase, a model cobamide-dependent

505   enzyme. bioRxiv 637140.

506   38.   Allen RH, Stabler SP. 2008. Identification and quantitation of cobalamin and cobalamin
507   analogues in human feces. Am J Clin Nutr 87:1324–1335.

508   39.   Abreu NA, Taga ME. 2016. Decoding molecular interactions in microbial communities.
509   FEMS Microbiol Rev 40:648–663.

510   40.   Karasawa T, Ikoma S, Yamakawa K, Nakamura S. 1995. A defined growth medium for
511   Clostridium difficile. Microbiology 141:371–375.

512   41.   Nawrocki KL, Wetzel D, Jones JB, Woods EC, McBride SM. 2018. Ethanolamine is a
513   valuable nutrient source that impacts *Clostridium difficile* pathogenesis. Environ
514   Microbiol 20:1419–1435.

515   42.   Dannheim H, Will SE, Schomburg D, Neumann-Schaal M. 2017. Clostridioides difficile
516   630Δerm in silico and in vivo – quantitative growth and extensive polysaccharide
517   secretion. FEBS Open Bio 7:602–615.

518   43.   Fletcher JR, Erwin S, Lanzas C, Theriot CM. 2018. Shifts in the gut metabolome and
519   Clostridium difficile transcriptome throughout colonization and infection in a mouse
520   model. mSphere 3:e00089-18.

521   44.   Köpke M, Straub M, Dürre P. 2013. Clostridium difficile Is an Autotrophic Bacterial
522   Pathogen. PLoS One 8:1–7.

523   45.   Avissar YJ, Beale SI. 1989. Identification of the enzymatic basis for Δ-aminolevulinic
524   acid auxotrophy in a hemA mutant of Escherichia coli. J Bacteriol 171:2919–2924.

525   46.   Roth JR, Lawrence JG, Bobik TA. 1996. Cobalamin (Coenzyme B12): Synthesis and
526   Biological Significance. Annu Rev Microbiol 50:137–181.

527   47.   Haslam SC, Ketley JM, Mitchell TJ, Stephen J, Burdon DW, Candy DC. 1986. Growth of
528   Clostridium difficile and production of toxins A and B in complex and defined media. J
529   Med Microbiol 21:293–297.

530   48.   Anderson PJ, Lango J, Carkeet C, Britten A, Kräutler B, Hammock BD, Roth JR. 2008.
531   One pathway can incorporate either adenine or dimethylbenzimidazole as an α-axial
532   ligand of B12 cofactors in Salmonella enterica. J Bacteriol 190:1160–71.

533   49.   Chan CH, Escalante-Semerena JC. 2011. ArsAB, a novel enzyme from Sporomusa ovata
534   activates phenolic bases for adenosylcobamide biosynthesis. Mol Microbiol 81:952–967.

535   50.   Ng YK, Ehsaan M, Philip S, Collery MM, Janoir C, Collignon A, Cartman ST, Minton
536   NP. 2013. Expanding the Repertoire of Gene Tools for Precise Manipulation of the
537   Clostridium difficile Genome: Allelic Exchange Using pyrE Alleles. PLoS One 8.

538   51.   Barker HA. 1967. Biochemical Functions of Corrinoid Compounds: The sixth Hopkins
539   Memorial Lecture. Biochem J 105:1–15.

540   52.   Watanabe F, Nakano Y, Stupperich E. 1992. Different corrinoid specificities for cell
541   growth and the cobalamin uptake system in Euglena gracilis z. J Gen Microbiol
542   138:1807–1813.

543   53.   Campbell GRO, Taga ME, Mistry K, Lloret J, Anderson PJ, Roth JR, Walker GC. 2006.
544   Sinorhizobium meliloti bluB is necessary for production of 5,6-dimethylbenzimidazole,
545   the lower ligand of B12. Proc Natl Acad Sci 103:4634–4639.

546   54.   Taga ME, Larsen NA, Howard-Jones AR, Walsh CT, Walker GC. 2007. BluB
547   cannibalizes flavin to form the lower ligand of vitamin B12. Nature 446:449–453.

548   55.   Gray MJ, Escalante-Semerena JC. 2007. Single-enzyme conversion of FMNH2 to 5, 6-
549   dimethylbenzimidazole the lower ligand of B12. Proc Natl Acad Sci 104:2921–2926.

56.   Hazra AB, Tran JLA, Crofts TS, Taga ME. 2013. Analysis of substrate specificity in

CobT homologs reveals widespread preference for DMB, the lower axial ligand of
vitamin B12. Chem Biol 20:1275–1285.

57. Crofts TS, Hazra AB, Tran JLA, Sokolovskaya OM, Osadchiy V, Ad O, Bauer S, Taga
ME. 2014. Regiospecific Formation of Cobamide Isomers Is Directed by CobT.
Biochemistry 53:7805–7815.

58. Hazra AB, Han AW, Mehta AP, Mok KC, Osadchiy V, Begley TP, Taga ME. 2015.
Anaerobic biosynthesis of the lower ligand of vitamin B12. Proc Natl Acad Sci
112:10792–7.

59. Yan J, Bi M, Bourdon AK, Farmer AT, Wang PH, Molenda O, Quaile AT, Jiang N, Yang
Y, Yin Y, Şimşir B, Campagna SR, Edwards EA, Löffler FE. 2018. Purinyl-cobamide is a
native prosthetic group of reductive dehalogenases. Nat Chem Biol 14:8–14.

60. Keller S, Ruetz M, Kunze C, Kräutler B, Diekert G, Schubert T. 2013. Exogenous 5,6-
dimethylbenzimidazole caused production of a non-functional tetrachloroethene reductive
dehalogenase in Sulfurospirillum multivorans. Environ Microbiol 16:3361–3369.

61. Yan J, Im J, Yang Y, Löffler FE. 2013. Guided cobalamin biosynthesis supports
dehalococcoides mccartyi reductive dechlorination activity. Philos Trans R Soc B Biol Sci
368:20120320.

62. Woodson JD, Escalante-Semerena JC. 2004. CbiZ, an amidohydrolase enzyme required
for salvaging the coenzyme B12 precursor cobinamide in archaea. Proc Natl Acad Sci
101:3591–3596.

63. Gray MJ, Escalante-Semerena JC. 2009. The cobinamide amidohydrolase (cobyric acid-
forming) CbiZ enzyme: A critical activity of the cobamide remodelling system of
Rhodobacter sphaeroides. Mol Microbiol 74:1198–1210.

64. Gray MJ, Escalante-Semerena JC. 2009. In vivo analysis of cobinamide salvaging in
Rhodobacter sphaeroides strain 2.4.1. J Bacteriol 191:3842–51.

65. Borths EL, Locher KP, Lee AT, Rees DC. 2002. The structure of Escherichia coli BtuF
and binding to its cognate ATP binding cassette transporter. Proc Natl Acad Sci
99:16642–16647.

66. Mireku SA, Ruetz M, Zhou T, Korkhov VM, Kräutler B, Locher KP. 2017.
Conformational Change of a Tryptophan Residue in BtuF Facilitates Binding and
Transport of Cobinamide by the Vitamin B12 Transporter BtuCD-F. Sci Rep 7:41575.

67. Vitreschak AG, Rodionov DA, Mironov AA, Gelfand MS. 2003. Regulation of the
vitamin B12 metabolism and transport in bacteria by a conserved RNA structural element.
Rna 9:1084–1097.

68. Nahvi A, Sudarsan N, Ebert MS, Zou X, Brown KL, Breaker RR. 2002. Genetic control
by a metabolite binding mRNA. Chem Biol 9:1043–1049.

69. Nahvi A, Barrick JE, Breaker RR. 2004. Coenzyme B12 riboswitches are widespread
genetic control elements in prokaryotes. Nucleic Acids Res 32:143–150.

70. Perez AA, Rodionov DA, Bryant DA. 2016. Identification and regulation of genes for
cobalamin transport in the cyanobacterium Synechococcus sp. strain PCC 7002. J
Bacteriol 198:2753–2761.

71. Stabler RA, Rose G, Gerding DN, Lawley TD, Quail MA, Wren BW, Gibert M, Dawson
L, Parkhill J, Sebaihia M, Dougan G, Valiente E, Corton C, Martin M, Popoff MR, He M.
2009. Comparative genome and phenotypic analysis of Clostridium difficile 027 strains
provides insight into the evolution of a hypervirulent bacterium. Genome Biol 10:R102.

72. Tanioka Y, Miyamoto E, Yabuta Y, Ohnishi K, Fujita T, Yamaji R, Misono H, Shigeoka

S, Nakano Y, Inui H, Watanabe F. 2010. Methyladeninylcobamide functions as the
cofactor of methionine synthase in a Cyanobacterium, Spirulina platensis NIES-39. FEBS
Lett 584:3223–3226.

73. Taga ME, Walker GC. 2010. Sinorhizobium meliloti Requires a Cobalamin-Dependent
Ribonucleotide Reductase for Symbiosis With Its Plant Host. MPMI 23:1643–1654.

74. Cadieux N, Bradbeer C, Reeger-schneider E, Mohanty AK, Wiener MC, Kadner RJ. 2002.
Identification of the Periplasmic Cobalamin-Binding Protein BtuF of Escherichia coli. J
Bacteriol 184:706–717.

75. Crofts TS, Men Y, Alvarez-Cohen L, Taga ME. 2014. A bioassay for the detection of
benzimidazoles reveals their presence in a range of environmental samples. Front
Microbiol 5:592.

76. Janoir C, Denève C, Bouttier S, Barbut F, Hoys S, Caleechum L, Chapetón-Montes D,
Pereira FC, Henriques AO, Collignon A, Monot M, Dupuy B. 2013. Adaptive strategies
and pathogenesis of clostridium difficile from In vivo transcriptomics. Infect Immun
81:3757–3769.

77. Kansau I, Barketi-Klai A, Monot M, Hoys S, Dupuy B, Janoir C, Collignon A. 2016.
Deciphering adaptation strategies of the epidemic clostridium difficile 027 strain during
infection through in vivo transcriptional analysis. PLoS One 11:1–13.

78. Scaria J, Janvilisri T, Fubini S, Gleed RD, McDonough SP, Chang YF. 2011. Clostridium
difficile transcriptome analysis using pig ligated loop model reveals modulation of
pathways not modulated in vitro. J Infect Dis 203:1613–1620.

79. Herbert V. 1988. Vitamin B12: plant sources, requirements, and assay. Am J Clin Nutr
48:852–858.

80. Degnan PH, Taga ME, Goodman AL. 2014. Vitamin B12 as a modulator of gut microbial
ecology. Cell Metab 20:769–778.

81. O'Connor JR, Lyras D, Farrow KA, Adams V, Powell DR, Hinds J, Cheung JK, Rood JI.
2006. Construction and analysis of chromosomal Clostridium difficile mutants. Mol
Microbiol 61:1335–1351.

82. Sorg JA, Dineen SS. 2009. Laboratory maintenance of Clostridium difficile. Curr Protoc
Microbiol 1–10.

83. Cartman ST, Minton NP. 2010. A mariner -Based Transposon System for In Vivo
Random Mutagenesis of Clostridium difficile. Appl Environ Microbiol 76:1103–1109.

84. Men Y, Seth EC, Yi S, Allen RH, Taga ME, Alvarez-Cohen L. 2014. Sustainable growth
of Dehalococcoides mccartyi 195 by corrinoid salvaging and remodeling in defined
lactate-fermenting consortia. Appl Environ Microbiol 80:2133–2141.

85. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods
6:343–345.

86. Bouillaut L, McBride SM, Sorg JA. 2011. Genetic manipulation of Clostridium difficile.
Curr Protoc Microbiol 1–17.

87. Williams DR, Young DI, Young M. 1990. Conjugative plasmid transfer from Escherichia
coli to Clostridium acetobutylicum. J Gen Microbiol 136:819–826.

639
640
641

642   **Figures**

643
644
645   **Figure 1. Structures of cobamides and cobamide precursors. A.** Structure of cobalamin (B_{12}). The
646   corrin ring, nucleotide loop, and lower ligand are labeled. **B.** Lower ligands of cobamides analyzed in this
647   study, with the three structural classes labeled. The lower ligand name, abbreviation for the cobamide
648   containing the lower ligand, and alternative names of the cobamide (when applicable) are indicated. **C.**
649   Cobamide precursors used in this study. R, upper ligand (-CN, -OH, -CH$_3$ or 5'-deoxyadenosyl).
650

651
652
Figure 2. Predicted cobamide metabolism in *C. difficile* 630 Δ*erm*. The cobamide biosynthesis
pathway is shown with a purple background, homologs of cobalamin-dependent enzymes in purple text,
cobalamin-independent isozymes in green text, and the transporter BtuFCD as a black rectangle.
Abbreviations: Cba, cobamide; Cbi, cobinamide; ALA, 5-aminolevulinic acid; rSAM, radical *S*-
adenosylmethionine; NDPs, ribonucleoside diphosphates; NTPs, ribonucleoside triphosphates; dNDPs,
deoxyribonucleoside diphosphates; dNTPs, deoxyribonucleoside triphosphates. Enzymes: MetH,
cobalamin-dependent methionine synthase; NrdEF, cobalamin-independent, aerobic (oxygen-requiring,
class I) ribonucleotide reductase (RNR); NrdDG, cobalamin-independent, anaerobic (oxygen-sensitive,
class III) RNR; NrdJ, cobalamin-dependent (class II) RNR; QueG, epoxyqueuosine reductase; EutBC,
ethanolamine ammonia lyase; CFeSP, corrinoid iron-sulfur protein; OraSE, D-ornithine 4,5-aminomutase.
663
664

665
666 **Figure 3. *C. difficile* can use a broad range of cobamides for MetH-dependent growth.** The O.D.$_{600}$ of
667 *C. difficile* 630 Δ*erm* cultures grown to saturation (22.5 hours) in CDDMK medium plus glucose without
668 methionine with the addition of **A.** cobamides or Cbi, **B.** methionine, and **C.** ALA is shown. The mean
669 and standard deviation of four biological replicates are shown in the bars and error bars, respectively.
670
671

Compound added

Figure 4. *C. difficile* is selective in which cobamides it can use for NrdJ-dependent growth. The O.D.$_{600}$ of *C. difficile* 630 Δ*erm* Δ*pyrE* Δ*nrdDG* cultures grown to saturation (22.5 hours) in CDDM with added uracil and glucose is shown for **A.** cobamides and Cbi, and **B.** ALA added. The mean and standard deviation of three biological replicates are shown in the bars and error bars, respectively.

678
Figure 5. HPLC analysis of corrinoid extracts from *C. difficile* cultures. A. HPLC analysis of
corrinoid extracts from cell pellets of *C. difficile* 630 Δ*erm* (WT) and 630 Δ*erm* Δ*pyrE*
Δ*cbiKLJHGFTEDC* grown to saturation in CDDM with glucose with either 100 nM ALA or 10 nM
dicyanocobinamide (Cbi) added. An asterisk (*) indicates the corrinoid peak validated by mass
spectrometry (Supplemental Fig. 2). **B**. HPLC analysis of corrinoid extracts of *C. difficile* 630 Δ*erm*
grown with either 100 nM ALA or 10 nM Cbi and 100 nM lower ligand bases DMB or Bza. An Agilent
Eclipse Plus C18 column and an Agilent Zorbax SB-Aq column were used to separate corrinoid
extractions in panels A and B, respectively. Cbi (1), pseudocobalamin (2), cobalamin (3), and [Bza]Cba
(4) are shown as standards.
688

689
Figure 6. *C. difficile ΔbtuFCD* mutant is impaired in cobamide and Cbi uptake. A. HPLC analysis of
corrinoid extracts from cell pellets of *C. difficile* 630 Δ*erm* (WT) and *C. difficile* 630 Δ*erm* Δ*pyrE*
Δ*btuFCD* grown with 10 nM cobamides or 100 nM ALA. Cbi (1), pseudocobalamin (2), cobalamin (3),
[2-MeAde]Cba (5), [Cre]Cba (6) are shown as standards. **B.** Growth of *C. difficile* 630 Δ*erm* Δ*pyrE*
Δ*btuFCD* in MetH-dependent conditions. The O.D.$_{600}$ of saturated cultures (23.5 hours) in CDDMK
without methionine plus glucose and uracil is plotted as a function of the amount of compound added.
Bars and error bars are the mean and standard deviation of three biological replicates.

704 **Table 1: Bacterial strains and plasmids**

Strain or plasmid	Description	Source
Strains		
Escherichia coli XL1-Blue		QB3 MacroLab
Escherichia coli CA434	hsd20(rB-, mB-, recA13, rpsL20, leu, proA2, with IncPb conjugative plasmid R702	Chain Biotech(87)
Clostridioides difficile		
630 Δ*erm*	Erythromycin sensitive strain	(81)
630 Δ*erm* Δ*pyrE*	Strain CRG1496	(50)
630 Δ*erm* Δ*pyrE* Δ*btuFCD*		This study
630 Δ*erm* Δ*pyrE* Δ*cbiKLJHGFTEDC*		This study
630 Δ*erm* Δ*pyrE* Δ*nrdDG*		This study
Plasmids		
R702	Conjugation helper plasmid	(87)
pMTL-YN3	Allelic exchange vector	(50)
pXL001	pMTL-YN3 containing *btuFCD* deletion construct	This study
pXL002	pMTL-YN3 containing *cbiKLJHGFTEDC* deletion construct	This study
pXL003	pMTL-YN3 containing *nrdDG* deletion construct	This study